

# Mass Spectra of $D$ , $D_s$ Mesons using Dirac formalism with martin-like confinement potential

Manan Shah<sup>1,\*</sup> Bhavin Patel<sup>2,†</sup> and P C Vinodkumar<sup>1‡</sup>

<sup>1</sup>*Department of Physics, Sardar Patel University,  
Vallabh Vidyanagar-388120, INDIA and*

<sup>2</sup>*P. D. Patel Institute of Applied Sciences, CHARUSAT, Changa-388421, INDIA*

## Introduction

Remarkable progress at the experimental side, with various high energy machines such as BaBar, BELLE, B-factories, Tevatron, ARGUS collaborations, CLEO, CDF, DØ etc., for the study of hadrons has opened up new challenges in the theoretical understanding of light-heavy flavour hadrons. The existing results on excited heavy-light mesons are partially inconclusive, and even contradictory in several cases. The predictions of masses of heavy-light system for low-lying  $1S$  and  $1P_J$  states of  $D$  and  $D_s$  mesons were known from experiment [1] and few from the theory [2–4]. Here we study the mass spectra of  $D$  and  $D_s$  mesons in a relativistic framework.

## Theoretical Framework

The bound constituent quark and antiquark inside the meson are in definite energy states having no definite momenta. However one can find out the momentum distribution amplitude for the constituent quark and antiquark inside the meson immediately before their annihilation to a lepton pair. Though the colour confinement of quarks are understood in terms of multigluon exchanging at the non-perturbative regime of the hadronic size, it is not feasible to compute theoretically from the QCD first principles. Thus one assumes

various confinement mechanism to study the hadronic properties. In the present study, we assume that the constituent quarks in a meson core is independently confined by an average Martin-like potential of the form [5]

$$V(r) = \frac{1}{2}(1 + \gamma_0)(\lambda r^{0.1} + V_0) \quad (1)$$

To a first approximation, the confining part of the interaction is believed to provide the zeroth-order quark dynamics inside the meson core through the quark Lagrangian density

$$\mathcal{L}_q^0(x) = \bar{\psi}_q(x) \left[ \frac{i}{2} \gamma^\mu \overleftrightarrow{\partial}_\mu - V(r) - m_q \right] \psi_q(x). \quad (2)$$

TABLE I: S-wave  $D$  ( $c\bar{q}$ ,  $q = d, u$ ) spectrum (in MeV).

nL	State	Present	Experiment		[6]	[7]
			Meson	Mass [1]		
1S	$1^3S_1$	2013.3	$D^*(2010)$	$2010.28 \pm 0.13$	-	2010
	$1^1S_0$	1874.0		$1869.62 \pm 0.15$	-	1871
2S	$2^3S_1$	2581.0	$D^*(2600)$	2608.7 [8]	2639	2632
	$2^1S_0$	2501.7	$D(2550)$	2539.4 [8]	2567	2581
3S	$3^3S_1$	3088.9		-	3125	3096
	$3^1S_0$	3031.5		-	3065	3062
4S	$4^3S_1$	3567.8		-	-	3482
	$4^1S_0$	3521.6		-	-	3452
1P	$1^3P_2$	2455.1	$D_2^*(2460)$	$2462.6 \pm 0.7$	-	2460
	$1^3P_1$	2348.0		-	-	2469
	$1^3P_0$	2276.6	$D_0^*(2400)$	$2318 \pm 29$	-	2406
	$1^1P_1$	2317.3	$D_1(2420)$	$2421.3 \pm 0.6$	-	2426
2P	$2^3P_2$	2907.0		-	2965	3012
	$2^3P_1$	2834.4		-	2960	3021
	$2^3P_0$	2786.0		-	2880	2919
	$2^1P_1$	2812.3		-	2940	2932

\*Electronic address: mnshah09@gmail.com

†Electronic address: azadpatel2003@yahoo.co.in

‡Electronic address: p.c.vinodkumar@gmail.com

The normalized quark wave functions  $\psi(\vec{r})$  obtained from Eqn (2) satisfies the Dirac equation given by

$$[\gamma^0 E_q - \vec{\gamma} \cdot \vec{P} - m_q - V(r)]\psi_q(\vec{r}) = 0. \quad (3)$$

The two component solution of Dirac equation can be written as

$$\psi_{nlj}(r) = \begin{pmatrix} \psi_A \\ \psi_B \end{pmatrix} \quad (4)$$

where the positive and negative energy solutions are written as

$$\psi_A^{(+)}(\vec{r}) = N_{nlj} \left( \frac{ig(r)}{(\sigma \cdot \hat{r}) f(r)} \right) \mathcal{Y}_{ljm}(\hat{r}) \quad (5)$$

$$\psi_B^{(-)}(\vec{r}) = N_{nlj} \left( \frac{i(\sigma \cdot \hat{r}) f(r)}{g(r)} \right) (-1)^{j+m_j-l} \mathcal{Y}_{ljm}(\hat{r}) \quad (6)$$

and  $N_{nlj}$  is the overall normalization constant. The radial solutions  $f(r)$  and  $g(r)$  is obtained numerically to yield the energy eigen values. The parameters are fixed to get the ground state masses of  $D$  and  $D_s$  mesons. The meson radial wave function for  $q\bar{q}$  combination

is constructed with the respective quark and anti-quark wave functions given by Eqn. (5) and 6. The quark mass parameters  $m_c$ ,  $m_{u,d}$  and  $m_s$  are taken as 1.27 GeV, 0.37 GeV and 0.4 GeV respectively.

## Results and Discussion

The predicted S-wave masses of  $D$  and  $D_s$  mesons are in very good agreement with experimental [1] results as given in Table I and II respectively. The predicted results of P-wave  $D$  meson states,  $1^3P_2$  (2455.1 MeV) and  $1^3P_0$  (2276.6 MeV) are in good agreement with experimental results of  $2462.6 \pm 0.7$  MeV and  $2318 \pm 29$  MeV [1] respectively. The predicted results of P-wave  $D_s$  meson states  $1^3P_2$  (2572.3 MeV),  $1^3P_1$  (2433.7 MeV) and  $1^3P_0$  (2341.3 MeV) are also good found in good agreement with experimental results  $2571.9 \pm 0.8$  MeV,  $2459.6 \pm 0.6$  MeV and  $2317.8 \pm 0.6$  MeV [1] respectively.

## Acknowledgments

The work is part of Major research project NO. F. 40-457/2011(SR) funded by UGC. One of the author (BP) acknowledges the support through Fast Track project funded by DST (SR/FTP/PS-52/2011).

## References

- [1] J. Beringer et al. (Particle Data Group), Phys. Rev. D 86, (2012) 010001.
- [2] S Godfrey and R Kokoski, Phys. Rev. D 43, (1991) 1679.
- [3] E J Eichen, C T Hill and C Quigg, Phys. Rev. Lett. 71, (1993) 4116.
- [4] D Ebert, R N Faustov and V O Galkin, Phys. Rev. D 57, (1998) 014027.
- [5] N. Barik, B. K. Dash, and M. Das, Phys. Rev. D **31**, 1652 (1985).
- [6] A. M. Badalian, B. L. G. Bakker, Phys. Rev. D 84, (2011) 034006.

TABLE II: S-wave  $D_s$  ( $c\bar{s}$ ) spectrum (in MeV).

nL	State	Present	Experiment		[6]	[7]
			Meson	Mass [1]		
1S	$1^3S_1$	2112.3	$D_s^*$	$2112.3 \pm 0.5$	-	2111
	$1^1S_0$	1970.6	$D_s$	$1968.49 \pm 0.32$	-	1969
2S	$2^3S_1$	2684.4	$D_{s1}(2710)$	$2710_{-7}^{+12}$	2728	2731
	$2^1S_0$	2603.9		2638 [9]	2656	2688
3S	$3^3S_1$	3195.1		-	3200	3242
	$3^1S_0$	3136.9		-	3140	3219
4S	$4^3S_1$	3676.0		-	-	3669
	$4^1S_0$	3629.3		-	-	3652
1P	$1^3P_2$	2572.3	$D_{s2}(2573)$	$2571.9 \pm 0.8$	-	2571
	$1^3P_1$	2433.7	$D_{s1}(2460)$	$2459.6 \pm 0.6$	-	2574
	$1^3P_0$	2341.3	$D_{s0}^*(2317)$	$2317.8 \pm 0.6$	-	2509
	$1^1P_1$	2420.4	$D_{s1}(2536)$	$2535.12 \pm 0.13$	-	2536
2P	$2^3P_2$	3023.2		-	3045	3142
	$2^3P_1$	2927.7		-	3020	3154
	$2^3P_0$	2864.1		-	2970	3054
	$2^1P_1$	2918.4		$3044_{-9}^{+30}$	3040	3067

- [7] D. Ebert, R. N. Faustov and V. O. Galkin, Eur. Phys. J. C 66 (2010) 197.
- [8] P. Del A. Sanchez et al. (BABAR Collaboration), Phys. Rev. D 82, (2010) 111101.
- [9] A.V. Evdokimov et al. (SELEX Collaboration), Phys. Rev. Lett. 93 (2004) 242001.