

ON THE STRUCTURE OF INVOLUTIONS AND SYMMETRIC SPACES OF QUASI DIHEDRAL GROUP

ZAHID RAZA, IMRAN AND BIJAN DAVVAZ

ABSTRACT. Let $G = QD_{8k}$ be the quasi-dihedral group of order $8n$ and θ be an automorphism of QD_{8k} of finite order. The fixed-point set H of θ is defined as $H_\theta = G^\theta = \{x \in G \mid \theta(x) = x\}$ and generalized symmetric space Q of θ is given by $Q_\theta = \{g \in G \mid g = x\theta(x)^{-1} \text{ for some } x \in G\}$.

The characteristics of the sets H and Q have been calculated. It is shown that for any H and Q , $H.Q \neq QD_{8k}$. The H -orbits on Q are obtained under different conditions. Moreover, the formula to find the order of v -th root of unity in $\mathbb{Z}_{2k}$ for QD_{8k} has been calculated. The criteria to find the number of equivalence classes denoted by C_{4k} of the involution automorphism has also been constructed. Finally, the set of twisted involutions $R = R_\theta = \{x \in G \mid \theta(x) = x^{-1}\}$ has been explored.

1. INTRODUCTION

Serious work on groups generated by reflections began in the 19th century. The finite subgroups of $O(3)$ generated by isometric reflections on the 2-sphere (or equivalently, by orthogonal linear reflections on $\mathbb{R}^3$) had been determined by Möbius in 1852. In the second half of the 19th century, work also began on finite reflection groups on S^n for $n > 2$ (or equivalently, finite linear reflection groups on $\mathbb{R}^{n+1}$). By the late 19's other groups generated by isometric reflections across the edges of polygons (with 3 or more edges) in the hyperbolic plane had been studied by Klein and Poincare. The group of symmetries of such a root system was a finite group generated by reflections had been shown by Weyl in 1925. This intimate connection with the classification of semi-simple Lie groups cemented reflection groups into a central place in mathematics. The two lines of research were united in (Coxeter, 1930). He classified discrete groups generated by reflections on the n -dimensional sphere or Euclidean space. The notion of an abstract reflection group introduced by Jacques Tits which he called a coxeter group. He considered pairs (W, S) with W a group and S a set of involutions which generate W so that the group has a presentation of the following form: the set of generators is S and the set of relations

2010 *Mathematics Subject Classification.* 20F28 .

Key words and phrases. Automorphisms, involutions, fixed-point set, symmetric spaces, quasi-dihedral group.

is $\{(st)^{m(s,t)}\}$, where $m(s, t)$ denotes the order of st and the relations range over all unordered pairs $s, t \in S$ with $m(s, t) \neq \infty$. The pair (W, S) is a coxeter system and W is a coxeter group. This was demonstrated in (Bourbaki, 1968; Bjorner and Francesco, 2005).

The early work by Helminck on symmetric spaces concerned some algebraic results about their structure (Helminck, 1988). Classification of the real symmetric spaces and their fine structure had been included in (Helminck, 1994). By considering an open subgroup H of the fixed point group G^θ of an involution θ of G , the result was proved that “For k a field of characteristic not two, let G be a connected semi-simple algebraic k -group and θ an involution of G . Then θ is defined over k if and only if H^θ is defined over k ” (Helminck and Wang, 1993). They also proved that “if G is a connected semi-simple algebraic group, θ_i involution of G and $H_i = G^{\theta_i}$, ($i = 1, 2$). Let $H_1^{\theta_1}$ and $H_2^{\theta_2}$ be the identity components of H_1 and H_2 respectively. If $H_1^{\theta_1} = H_2^{\theta_2}$ then $\theta_1 = \theta_2$.”

A mini-workshop on generalizations of symmetric spaces was organized by Ralf Kohl at.el in 2012. Mathematicians in this area brought together to present their current research projects or trigger new collaboration via comparison, analogy, transfer, generalization and unification of method in this workshop.

The structure of automorphism group of dihedral group has been found by Cunningham at.el (2012). They also characterized the automorphism group with $\theta^2 = \text{id}$ and calculated H and Q for each automorphism for the family of finite dihedral groups. They also proved that the set of involutions is partitioned in such a way that there are at most two distinct equivalence classes of involutions in each of the partition. In involution case, Q is always a subgroup of G . Moreover, the infinite dihedral group is also discussed.

Let $G = D_{C_n}$ be the dicyclic group of order $4n$. Let ϕ be an automorphism of G of order k . Bishop at.el (2013) describe ϕ and the generalized symmetric space Q of G associated with ϕ . When ϕ is an involution, they describe its fixed point group H along with the H -orbits and G -orbits of Q corresponding to the action of ϕ -twisted conjugation.

In this paper, we extent the study to quasi-dihedral group QD_{8k} .

2. PRELIMINARIES

The quasi-dihedral group is denoted by QD_{8k} , where $QD_{8k} = \langle a, b : a^{4k} = b^2 = 1, ba = a^{2k-1}b \rangle$. In reset of the whole paper, we denote it by QD_{2m} , where $m = 4k$ and $n = \frac{m}{2} = 2k$. That is $QD_{2m} = \langle a, b : a^m = b^2 = 1, ba = a^{n-1}b \rangle$. It is clear from the presentation that $a^i b^j \in QD_{2m}$ is in normal form for some integer $0 \leq i < m$ and $j \in \{0, 1\}$. In the whole study, $\mathbb{Z}_n$ denotes the additive group of integers modulo n and U_m denotes the multiplicative group of units of $\mathbb{Z}_n$ and

$r \in U_m$ if and only if $\gcd(r, m) = 1$. The v -th root of unity in $\mathbb{Z}_n$ which can be described by $R_m^v = \{r \in U_m \mid r^v = 1 \pmod{n}\}$.

Let G be a group, then an element a is a conjugate with b in G if and only if there exists an element x such that $a = xbx^{-1}$. For $a \in G$, the function $\phi_a(x) = axa^{-1} \forall x \in G$ is called an inner automorphism of G . Let θ_1, θ_2 be two automorphisms, then $\theta_1 \sim \theta_2$ if and only if they are conjugate to each other. If there exists another automorphism σ such that $\sigma\theta_1\sigma^{-1} = \theta_2$.

Definition 3. For any $c \in \mathbb{Z}_n$, we define $ZDiv(c) = \{y \in \mathbb{Z}_n \mid cy \equiv 0 \pmod{n}\}$ and easy to see that $|ZDiv(c)| = \gcd(c, n)$

Notation 4. Let us fixed the following notations, which will be used in the whole paper.

- For fixed $r \in R_m^v$. The number of equivalence classes with fixed leading co-efficient r is denoted by $N_r = |\{\bar{\theta} \mid \theta = rx + s \in Aut_v(D_{2m})\}|$
- Let us consider the subset of automorphism denoted by $Aut_v(QD_{2m})$ and is defined as $Aut_v(QD_{2m}) = \{\theta \in Aut(QD_{2m}) \mid \theta^v = \text{id}\}$.

The following lemma is easy to prove.

Lemma 5. If QD_{2m} be the quasi-dihedral group, then

- (1) $ba^i = a^{i(n-1)}b$ for all $i = 0, 1, \dots, (m-1)$,
- (2) $|a^{2i}b| = 2$ and $|a^{2i+1}b| = 4$ for all $i = 0, 1, \dots, (n-1)$.

Lemma 6. The automorphism group of QD_{2m} is isomorphic to the group of affine linear transformations of $\mathbb{Z}_n$. In particular, we have

$$Aut(QD_{2m}) \cong Aff(\mathbb{Z}_n) = \{rx + s : \mathbb{Z}_n \rightarrow \mathbb{Z}_n \mid r \in U_m, s \in \mathbb{Z}_n\}$$

and the action of $rx + s$ on elements of QD_{2m} in normal form is given by

$$(6.1) \quad (rx + s)(a^j b^t) = a^{rj + [(t-1)n - (t-2)]s} b^t$$

Proof. Let θ be an automorphism, then $\theta(a) = a^r$ and $\theta(b) = (a^s b)$, where $\gcd(r, m) = 1$, $s = 0, 2, 4, \dots, (m-2)$. Consider arbitrary element $a^j b^t \in QD_{2m}$, and θ being homomorphism,

$$(6.2) \quad \theta(a^j b^t) = \theta(a^j) \theta(b^t),$$

then $\theta(a^j) = a^{rj}$ and $\theta(b^t) = (a^s b)(a^s b)(a^s b) \dots (a^s b) = a^s (ba^s)(ba^s)b \dots (a^s b)$.

By Lemma 5, we have

$$\begin{aligned} \theta(b^t) &= a^s [a^{s(n-1)} b] (ba^s) (ba^s) b \dots (a^s b) = a^{s+s(n-1)} (b^2 a^s) (ba^s) b \dots (a^s b) \\ \theta(b^t) &= a^{sn} (b^2 a^s) (ba^s) b \dots (a^s b) = a^{sn} (a^{s(n-1)} b^2) (ba^s) b \dots (a^s b) \\ \theta(b^t) &= a^{sn+sn-s} b^3 (ba^s) b \dots (a^s b) = a^{2sn-s} b^3 (ba^s) b \dots (a^s b) \text{ after } t\text{-steps, we have} \\ \theta(b^t) &= a^{(t-1)ns - (t-2)s} b^t. \text{ Put values in equation 6.2, we get} \\ \theta(a^j b^t) &= a^{rj} \cdot a^{(t-1)ns - (t-2)s} b^t = a^{rj + (t-1)ns - (t-2)s} b^t = a^{rj + [(t-1)n - (t-2)]s} b^t. \text{ Since } b^2 = \end{aligned}$$

1, so $t \in \{0, 1\}$. If $t = 0$, then $\theta(a^j) = a^{rj+(2-n)s} \implies rx + (2-n)s \in \text{Aff}(\mathbb{Z}_n)$. If $t = 1$, then $\theta(a^j b) = a^{rj+s} b \implies rx + s \in \text{Aff}(\mathbb{Z}_n)$.

Conversely, For $f \in \text{Aff}(\mathbb{Z}_n)$, that is $f = rx + s$ where $r \in U_m$, $s \in \mathbb{Z}_n$ and it is given as a group action $f(a^j b^t) = a^{rj+[(t-1)n-(t-2)]s} b^t$ for all $a^j b^t \in QD_{2m}$.

Now, we will show that f is an automorphism. For this let us define $\theta : QD_{2m} \longrightarrow QD_{2m}$ by $\theta(a^j b^t) = f(a^j b^t) = a^{rj+[(t-1)n-(t-2)]s} b^t$. We will show that θ is a bijective homomorphism. For this we will show that θ is one to one and onto.

one to one: Suppose that $\theta(a^{j_1} b^{t_1}) = \theta(a^{j_2} b^{t_2})$, then $a^{rj_1+[(t_1-1)n-(t_1-2)]s} b^{t_1} = a^{rj_2+[(t_2-1)n-(t_2-2)]s} b^{t_2} \implies a^{rj_1+[(t_1-1)n-(t_1-2)]s-rj_2-[(t_2-1)n-(t_2-2)]s} b^{t_1-t_2} = e$.

Case 1 :

$rj_1 + [(t_1 - 1)n - (t_1 - 2)]s - rj_2 - [(t_2 - 1)n - (t_2 - 2)]s = 0$ and $t_1 - t_2 = 2$, since, $t \in \{0, 1\}$ therefore $t_1 - t_2 \neq 2$ not possible.

Case 2 :

$rj_1 + [(t_1 - 1)n - (t_1 - 2)]s - rj_2 - [(t_2 - 1)n - (t_2 - 2)]s = m$ and $t_1 - t_2 = 0 \implies r(j_1 - j_2) + ns(t_1 - t_2) - s(t_1 - t_2) = m \implies r(j_1 - j_2) = m \implies r \mid m$ but $\gcd(r, m) = 1$, not possible. Now $r(j_1 - j_2) = m$ will be only possible when $j_1 - j_2 = 0$, $r \neq 0$ $j_1 = j_2 \implies a^{j_1} = a^{j_2} \implies a^{j_1} b^{t_1} = a^{j_2} b^{t_2} \implies \theta$ is one to one.

onto: It is cleared from the definition of the group action that for every $a^{rj+[(t-1)n-(t-2)]s} b^t$ there exists $a^j b^t \in QD_{2m}$ such that $\theta(a^j b^t) = a^{rj+[(t-1)n-(t-2)]s} b^t \implies \theta$ is onto. Hence $\text{Aut}(QD_{2m}) \cong \text{Aff}(\mathbb{Z}_n)$. $\square$

7. STRUCTURE OF AUTOMORPHISMS OF QD_{2m}

Proposition 7.1. For any integer $v \geq 1$ and $\theta = rx + s$ be an automorphism of QD_{2m} then $\theta^v = \text{id}$ if and only if $r \in R_m^v$ and $s \in Z\text{Div}(r^{v-1} + r^{v-2} + \dots + r + 1)$.

Proof. We will use the mathematical induction on v to show first the following statement: $\theta^v(x) = r^v x + (r^{v-1} + r^{v-2} + \dots + r + 1)s$.

For $v = 2$, 3 $\theta^2(x) = \theta[\theta(x)] = \theta(rx + s) = r(rx + s) + s = r^2 x + rs + s = r^2 x + (r + 1)s$. Similarly, $\theta^3(x) = r^3 x + r^2 s + rs + s = r^3 x + (r^2 + r + 1)s$.

Suppose that the statement is true for $v = i$ that is

$\theta^i(x) = r^i x + (r^{i-1} + r^{i-2} + \dots + r + 1)s$. We will show that the statement is true for $v = i + 1$. Take $\theta^{i+1}(x) = \theta[\theta^i(x)] = \theta[r^i x + (r^{i-1} + r^{i-2} + \dots + r + 1)s]$ $\theta^{i+1}(x) = r[r^i x + (r^{i-1} + r^{i-2} + \dots + r + 1)s] + s = r^{i+1} x + (r^i + r^{i-1} + \dots + r + 1)s$. Hence, the statement is true for all positive values of v . We have identified the identity automorphism with $x \in \text{Aff}(\mathbb{Z}_n)$, thus $r^v \equiv 1 \pmod{n}$ and $(r^{v-1} + r^{v-2} + \dots + r + 1)s \equiv 0 \pmod{n} \Leftrightarrow r \in R_m^v$ and $s \in Z\text{Div}(r^{v-1} + r^{v-2} + \dots + r + 1)$ as required. $\square$

Example 7.2. Let $G = QD_{56}$ and $v = 3$. In this case $R_{28}^3 = \{1, 9, 11, 15, 23, 25\}$, take $r = 9$, then $Z\text{Div}(9^2 + 9 + 1) = Z\text{Div}(91) = \{0, 2, 4, 6, 8, 10, 12\}$. Now, consider

$\theta_1 = 9x+6 \in \text{Aut}(QD_{56})$, then after simple calculation we get $\theta_1^3 = x = \text{id}$. Similarly, for $\theta_2 = 9x+12 \in \text{Aut}(QD_{56})$, we have $\theta_2^3 = x = \text{id}$.

Proposition 7.3. For any m and $v \geq 1$, we have

$$|\text{Aut}_v(QD_{2m})| = \sum_{r \in R_m^v} \gcd(r^{v-1} + r^{v-2} + \cdots + r + 1, n)$$

Proof. This follows from above Proposition: for any $r \in R_m^v$ there are

$$|Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1)| = \gcd(r^{v-1} + r^{v-2} + \cdots + r + 1, n)$$

element s such that $(r^{v-1} + r^{v-2} + \cdots + r + 1)s \equiv 0 \pmod{n}$ and every automorphism $\theta \in \text{Aut}_v(QD_{2m})$ must be of this form. $\square$

Proposition 7.4. Let $\theta_1 = r_1x + s_1$ and $\theta_2 = r_2x + s_2$ be two different automorphism of QD_{2m} . Then $\theta_1 \sim \theta_2$ if and only if $r_1 = r_2$ and $fs_1 - s_2 \in \langle r_1 - 1 \rangle \leq \mathbb{Z}_n$ for some $f \in U_m$.

Proof. Given that $\theta_1 = r_1x + s_1$ and $\theta_2 = r_2x + s_2 \in \text{Aut}(QD_{2m})$. Suppose that $\sigma = fx + g$ be any arbitrary automorphism of QD_{2m} and define $\sigma^{-1} = f^{-1}x - f^{-1}g$. We will check that $\sigma \cdot \sigma^{-1} = x$. For this take $\sigma \cdot \sigma^{-1} = \sigma(\sigma^{-1}(x)) = \sigma(f^{-1}x - f^{-1}g) = f(f^{-1}x - f^{-1}g) + g = x - g + g = x$. Now, we calculate

$$(7.1) \quad \sigma\theta_1\sigma^{-1} = \sigma \cdot (\theta_1 \cdot \sigma^{-1})$$

Take $\theta_1 \cdot \sigma^{-1} = \theta_1[\sigma^{-1}(x)] = \theta_1[f^{-1}x - f^{-1}g] = r_1(f^{-1}x - f^{-1}g) + s_1 = r_1f^{-1}x - r_1f^{-1}g + s_1$ put in 7.1, we get $\sigma\theta_1\sigma^{-1} = \sigma(r_1f^{-1}x - r_1f^{-1}g + s_1) = f(r_1f^{-1}x - r_1f^{-1}g + s_1) + g = r_1x - r_1g + fs_1 + g = r_1x + fs_1 - g(r_1 - 1)$. Now, by definition $\sigma\theta_1\sigma^{-1} = \theta_2$ if and only if $r_1 = r_2$ and $fs_1 - g(r_1 - 1) = s_2 \Rightarrow fs_1 - s_2 = g(r_1 - 1) \Rightarrow fs_1 - s_2 \in \langle r_1 - 1 \rangle$. $\square$

Example 7.5. Let $G = QD_{64}$ and consider $\theta_1 = 7x + 3$, $\theta_2 = 7x + 14$, $\theta_3 = 7x + 11$ three automorphisms of QD_{64} . Then it is easy to check that $\theta_1 \not\sim \theta_2$ because there does not exist such $f \in U_{32}$ for which $3f - 14 \in \langle 6 \rangle$. And $\theta_1 \sim \theta_3$ because there exist some $f \in U_{32}$ for which $3f - 14 \in \langle 6 \rangle$.

Proposition 7.6. Suppose m is fixed and let $v \geq 1$.

- (1) For any $r \in R_m^v$; $\langle r - 1 \rangle \leq Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1)$.
- (2) For any $r \in R_m^v$; U_m acts on the cosets $\frac{Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1)}{\langle r - 1 \rangle}$.
- (3) The set $\text{Aut}_v(QD_{2m})$ is partitioned into equivalence classes indexed by pair (r, A) , where $r \in R_m^v$ and A is an orbit of U_m on $\frac{Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1)}{\langle r - 1 \rangle}$.

Proof. (1) We know that $(r - 1) \cdot (r^{v-1} + r^{v-2} + \cdots + r + 1) = r^v - 1 \equiv 0 \pmod{n} \Rightarrow (r - 1) \cdot (r^{v-1} + r^{v-2} + \cdots + r + 1) \equiv 0 \pmod{n} \Rightarrow (r - 1) \in Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1) \Rightarrow \langle r - 1 \rangle \leq Z\text{Div}(r^{v-1} + r^{v-2} + \cdots + r + 1)$.

- (2) We simply recognize that $U_m = \text{Aut}(\mathbb{Z}_n)$ acts on $\mathbb{Z}_n$ by multiplication and since every subgroup is cyclic, U_m must stabilize the subgroups of $\mathbb{Z}_n$.
- (3) It is simply Proposition 7.4 in terms of the U_m -action on $\frac{Z\text{Div}(r^{v-1}+r^{v-2}+\dots+r+1)}{\langle r-1 \rangle}$.
-

Example 7.7. Let $G = QD_{40}$, and $v = 2$. Then $U_{20} = \{1, 3, 7, 9, 11, 13, 17, 19\}$ and $\mathbb{Z}_{10} = \{0, 1, 2, \dots, 9\}$.

- (1) Since $v = 2$, so $R_{20}^2 = \{1, 9, 11, 19\}$. Take $r = 9$ then $\langle 9 - 1 \rangle = \langle 8 \rangle = \{0, 2, 4, 6, 8\}$ and $Z\text{Div}(10) = \{0, 1, 2, \dots, 9\}$. So, $\{0, 2, 4, 6, 8\} \subseteq \{0, 1, 2, \dots, 9\}$
- (2) For U_{20} action on the cosets $\frac{Z\text{Div}(r+1)}{\langle r-1 \rangle}$. Take $r = 9$ then $\frac{Z\text{Div}(r+1)}{\langle r-1 \rangle} = \frac{Z\text{Div}(10)}{\langle 8 \rangle} = \frac{E=\{0,1,2,\dots,9\}}{F=\{0,2,4,6,8\}} = x + F \ \forall \ x \in E$ provide two sets F and $F_1 = \{1, 3, 5, 7, 9\}$. Now, the action of U_{20} on the cosets $\frac{Z\text{Div}(10)}{\langle 8 \rangle}$ has two sets. $\frac{Z\text{Div}(10)}{\langle 8 \rangle} = y.F = F \ \forall \ y \in U_{20}$ and $\frac{Z\text{Div}(10)}{\langle 8 \rangle} = y.F_1 = F_1 \ \forall \ y \in U_{20}$
- (3) The action of U_{20} on $\frac{Z\text{Div}(10)}{\langle 8 \rangle} = F, F_1$. Then the number of equivalence classes of $\text{Aut}(QD_{40})$ indexed by the pair (r, A) where $r \in R_{20}^2$ and A is an orbit of U_{20} on $\frac{Z\text{Div}(10)}{\langle 8 \rangle}$ is 8.

Theorem 7.8. If m is a fixed positive integer and $r \in R_m^v$, then the number of orbits of U_m on $\frac{Z\text{Div}(r^{v-1}+r^{v-2}+\dots+r+1)}{\langle r-1 \rangle}$ is equal to the number of divisors of $\frac{\text{gcd}(r-1, n) \cdot \text{gcd}(r^{v-1}+r^{v-2}+\dots+r+1, n)}{n}$.

Proof. The U_m -orbit on $\mathbb{Z}_n$ are indexed by the subgroups of $\mathbb{Z}_n$. Thus the U_m -orbit on $\frac{Z\text{Div}(r^{v-1}+r^{v-2}+\dots+r+1)}{\langle r-1 \rangle}$ are indexed by subgroups $L \leq \mathbb{Z}_n$ such that

$$\langle r-1 \rangle \leq L \leq Z\text{Div}(r^{v-1} + r^{v-2} + \dots + r + 1).$$

It is well known that the subgroup lattice of $\mathbb{Z}_n$ is isomorphic to the divisor lattice of n . The subgroup $\langle r-1 \rangle$ corresponds to the divisor $\text{gcd}(r-1, n)$ and the subgroup $Z\text{Div}(r^{v-1} + r^{v-2} + \dots + r + 1)$ correspond to the divisor $\frac{n}{\text{gcd}(r^{v-1}+r^{v-2}+\dots+r+1, n)}$, and the subgroups between these groups correspond to the divisor of n between $\text{gcd}(r-1, n)$ and $\frac{n}{\text{gcd}(r^{v-1}+r^{v-2}+\dots+r+1, n)}$ in the divisor lattice. Finally, it is known that this sub-lattice of the divisor of n is isomorphic to the divisor lattice of

$$\frac{\text{gcd}(r-1, n)}{\frac{n}{\text{gcd}(r^{v-1}+r^{v-2}+\dots+r+1, n)}} = \frac{\text{gcd}(r-1, n) \cdot \text{gcd}(r^{v-1} + r^{v-2} + \dots + r + 1, n)}{n}.$$

□

8. FIXED SSTS AND SYMMETRIC SPACES OF AUTOMORPHISMS

In this section if θ is an automorphism the fixed-point set H_θ and the symmetric space Q_θ will be discussed, where $H_\theta = \{x \in G \mid \theta(x) = x\}$ and $Q_\theta = \{g \in G \mid g =$

$x\theta(x)^{-1}$ for some $x \in G$. When θ is understood to be fixed the subscript from our notation has been dropped.

The following theorem characterizes the sets H and Q in the case of quasi-dihedral groups.

Theorem 8.1. If $\theta = rx + s$ is an automorphism of $G = QD_{2m}$ of finite order, then $H = \{a^j \mid j(r+2s-1) \equiv 0 \pmod{n}\} \cup \{a^{j'}b \mid j'(r+2s-1) \equiv -s \pmod{n}\}$ and $Q = \{a^j \mid j \in \langle r+2s-1 \rangle \cup [l(1-r)-s] \pmod{n}\}$ where, j' and l are even numbers.

Proof. Since, θ is an automorphism and $\theta(a^jb^t) = a^{rj+[(t-1)n-(t-2)]s}b^t$. First we will find set the H . Let us consider $a^j \in G$, then by definition of H , we have $\theta(a^j) = a^j$, $\theta(a^j) = a^{[r+(2-n)s]j} = a^j, \Rightarrow n \mid j(r+2s-1)$.

Now consider $a^{j'}b \in G$, then $\theta(a^{j'}b) = a^{j'}b$ so, $a^{j'}b = a^{[r+(2-n)s]j'}a^sb$, thus $n \mid (r+2s-1)j' + s$. Hence, $H = \{a^j \mid j(r+2s-1) \equiv 0 \pmod{n}\} \cup \{a^{j'}b \mid j'(r+2s-1) \equiv -s \pmod{n}\}$, where j' is even number.

Now we will find the set Q . Let $a^i \in G$ be an arbitrary element and let a^i be a fixed element of G , where $j \neq i$, then by definition of Q , $a^j = a^i\theta(a^i)^{-1} = a^i\theta(a^{-i}) = a^i.a^{-[r+(2-n)s]i} = a^{i[1-r-(2-n)s]} \Rightarrow j \equiv i[1-(r+2s)] \pmod{n} \Rightarrow j \in \langle r+2s-1 \rangle$.

Now consider $a^lb \in G$, where l is even number, then $a^j = (a^lb)\theta(a^lb)^{-1} = (a^lb)\theta(a^lb) = (a^lb)(a^{rl+s}b) = a^l(ba^{rl+s})b = a^la^{(rl+s)(n-1)}bb = a^{l+(rl+s)(n-1)} \Rightarrow j = l + (rl+s)(n-1) \Rightarrow j \equiv l[1+r(n-1)] + s(n-1) \Rightarrow j \equiv l(1-r) - s \pmod{n}$.

So, $Q = \{a^j \mid j \in \langle r+2s-1 \rangle \cup [l(1-r)-s] \pmod{n}\}$, where l is an even numbers. $\square$

Example 8.2. Let us consider $G = QD_{64}$ and $\theta = 3x + 12 \in \text{Aut}(D_{64})$, then $H = \{a^j \mid 10j \equiv 0 \pmod{16}\} \cup \{a^{j'}b \mid 10j' \equiv -4 \pmod{16}\} = \{1, a^8, a^2b, a^{10}b\}$ and $Q = \{a^j \mid j \in \langle 10 \rangle \cup (14j' + 4) \pmod{16}\} = \{1, a^2, a^4, \dots, a^{14}\}$

Using the descriptions of H and Q , we obtain further results as well.

Corollary 8.3. If $\theta = rx + s$ is an automorphism of QD_{2m} of finite order, where $m = 4k$, $n = 2k$ and k is even, then H will be cyclic or non-cyclic according to the following conditions

- (1) Let $\gcd(r+2s-1, n) = d_1$. If $s = 2w + 1$ is odd and $d_1 \nmid w$ then H will be cyclic that is $H \cong Z\text{Div}(r+2s-1)$ and $|H| = d_1$
- (2) Let $\gcd(r+2s-1, k) = d_2$. If $s = 2w$ is even and $d_2 \mid w$ then we have two cases
 - a): If w is odd, then H will be cyclic $|H| = d_1$ and $H \cong Z\text{Div}(r+2s-1)$.
 - b): If w is even
 - i): If $d_2 \nmid w$ then H will be cyclic $|H| = d_1$ and $H \cong Z\text{Div}(r+2s-1)$.
 - ii): If $d_2 \mid w$ then H will be non-cyclic. We have two cases

- I-):** if $4 \nmid k$ then $|H| = \frac{3}{2}d_1$ $H = \{\langle a^{\frac{k}{2}} \rangle, a^{2l}b, a^{2q}b \mid l \neq q \text{ for some } l, q \in \{1, 2, \dots, k\}\}$. It is not a subgroup of G .
II-): if $4 \mid k$ then $|H| = 2d_1$ and $H \cong ZDiv(r + 2s - 1) \times Z_2$.

Example 8.4. Let $G = QD_{48}$, then $\gcd(r, m) = 1 \Rightarrow r \in U_{24} = \{1, 5, 7, 11, 13, 17, 19, 23\}$ and $s \in \mathbb{Z}_{12} = \{0, 1, 2, \dots, 11\}$.

- (1) Consider $\theta_1 = 7x + 11 \in Aut(QD_{48})$ then $r = 7, s = 2(5) + 1 = 11 \Rightarrow r + 2s - 1 = 4$ and $d_1 = \gcd(28, 12) = 4 \Rightarrow 4 \nmid 5$ then $H_1 = \{1, a^3, a^6, a^9\}$ is cyclic. and $ZDiv4 = \{0, 3, 6, 9\} \Rightarrow H_1 \cong ZDiv4$ and $O(H_1) = d_1 = 4$
- (2) **a):** Take $\theta_2 = 11x + 10 \in Aut(QD_{48})$ then $r = 11, s = 2(5) = 10 \Rightarrow r + 2s - 1 = 6$ and $d_1 = \gcd(6, 12) = 6, d_2 = \gcd(6, 6) = 6$ then $H_2 = \{1, a^2, a^4, \dots, a^{10}\}$ is cyclic and $ZDiv6 = \{0, 2, 4, 6, 8, 10\} \Rightarrow H_2 \cong ZDiv6$ and $O(H_2) = d_1 = 6$
- b):** **i):** For $\theta_3 = 11x + 4 \in Aut(QD_{48})$ then $r = 11, s = 2(2) = 4 \Rightarrow r + 2s - 1 = 6$ and $d_1 = \gcd(6, 12) = 6, d_2 = \gcd(6, 6) = 6 \Rightarrow 6 \nmid -2$ then $H_3 = \{1, a^2, a^4, \dots, a^{10}\}$ is cyclic and $ZDiv6 = \{0, 2, 4, 6, 8, 10\} \Rightarrow H_3 \cong ZDiv6$ and $O(H_3) = d_1 = 6$
- ii):** **I-):** For $\theta_4 = 5x + 8 \in Aut(QD_{48})$ then $r = 5, s = 2(4) = 8 \Rightarrow r + 2s - 1 = 8$ and $d_1 = \gcd(8, 12) = 4, d_2 = \gcd(8, 6) = 2 \Rightarrow 2 \mid -4$ then $H_4 = \{1, a^3, a^6, a^9, a^{2l}b, a^{8l}b\}$ is non-cyclic and $O(H_4) = \frac{3}{2}.d_1 = 6$
- II-):** Take $\theta_5 = 7x + 4 \in Aut(QD_{48})$ then $r = 7, s = 2(2) = 4 \Rightarrow r + 2s - 1 = 2$ and $d_1 = \gcd(2, 12) = 2, d_2 = \gcd(2, 6) = 2 \Rightarrow 2 \mid -2$ then $H_5 = \{1, a^6, a^4b, a^{10}b\}$ is non-cyclic and $O(H_5) = 2.d_1 = 4$

Corollary 8.5. If $m = 4k, n = 2k$ and k is odd, then H will be cyclic or non-cyclic according to the following conditions

- (1) Let $\gcd(r + 2s - 1, n) = d_1$. If $s = 2w + 1$ is odd then H will be cyclic $|H| = d_1$ and $H \cong ZDiv(r + 2s - 1)$.
- (2) Let $\gcd(r + 2s - 1, n) = d_1$. If $s = 2w$ is even then
 - a):** If $d_1 = n$ then H will be cyclic $|H| = d_1 = n$ and $H \cong ZDiv(r + 2s - 1)$.
 - b):** If $d_1 = 2$ then H will be non-cyclic $|H| = \frac{3}{2}d_1 = \frac{3}{2}(2) = 3$ and $H = \{1, a^k, a^{2l}b\}$ for some $l \in \{1, 2, \dots, k\}$ is not a subgroup of G .

Example 8.6. Let $G = QD_{40}$ where $m = 20, n = 10, k = 5$. Since $\gcd(r, m) = 1 \Rightarrow r \in U_{20} = \{1, 3, 7, 9, 11, 13, 17, 19\}$ and $s \in \mathbb{Z}_{10} = \{0, 1, 2, \dots, 9\}$.

- (1) Consider $\theta_1 = 3x + 5 \in Aut(QD_{40})$ then $r = 3, s = 5 \Rightarrow r + 2s - 1 = 2$ and $\gcd(2, 10) = 2$ then $H_1 = \{1, a^5\}$ is cyclic and $ZDiv2 = \{0, 5\} \Rightarrow H_1 \cong ZDiv2$ and $|H_1| = 2$

- (2) **a):** For $\theta_2 = 7x + 2 \in \text{Aut}(QD_{40})$ then $r = 7, s = 2 \Rightarrow r + 2s - 1 = 0$ and $\gcd(0, 10) = 10$ then $H_2 = \{1, a^2, a^3, \dots, a^9\}$ is cyclic and $Z\text{Div}0 = \{0, 1, 2, \dots, 9\} \Rightarrow H_2 \cong Z\text{Div}0$ and $|H_2| = 10$
- b):** Take $\theta_3 = 7x + 4 \in \text{Aut}(QD_{40})$ then $r = 7, s = 4 \Rightarrow r + 2s - 1 = 4$ and $\gcd(4, 10) = 2$ then $H_3 = \{1, a^5, a^4b\}$ is non-cyclic and $|H_3| = \frac{3}{2} \cdot 2 = 3$

Corollary 8.7. If $\theta = rx + s \in \text{Aut}_v(QD_{2m})$ be of finite order, where $m = 4k, n = 2k$ and k is even number, then Q will be cyclic or non-cyclic according to the following conditions

- (1) If $s = 2w + 1$ is odd, then Q will be non-cyclic.
- (2) Let $\gcd(r + 2s - 1, n) = d_1$. If $s = 2w$ is even, then
- a):** If $d_1 \mid (1 - r)j' - s$ and
- i):** if $r + 2s - 1 \not\equiv k \pmod{n}$ then Q will be cyclic.
- ii):** If $r + 2s - 1 \equiv k \pmod{n}$ we have two cases
- if $1 - r \equiv k \pmod{n}$ then Q will be cyclic.
 - If $1 - r \not\equiv k \pmod{n}$ then Q will be non-cyclic.
- b):** If $d_1 \nmid (1 - r)j' - s$ and
- i):** if $-s \not\equiv k \pmod{n}$ then Q will be non-cyclic.
- ii):** If $-s \equiv k \pmod{n}$ then Q will be cyclic.

Example 8.8. Let $G = QD_{48}$ where $m = 24, n = 12, k = 6$. Since $\gcd(r, m) = 1 \Rightarrow r \in U_{24} = \{1, 5, 7, 11, 13, 17, 19, 23\}$ and $s \in \mathbb{Z}_{12} = \{0, 1, 2, \dots, 11\}$.

- (1) Consider $\theta_1 = 7x + 11 \in \text{Aut}(QD_{48})$ then $r = 7, s = 11 \Rightarrow r + 2s - 1 = 4$ then $Q_1 = \{1, a, a^4, a^8\}$ is non-cyclic.
- (2) **a):** **i):** For $\theta_2 = 5x + 8 \in \text{Aut}(QD_{48})$ then $r = 5, s = 8 \Rightarrow r + 2s - 1 = 8 \not\equiv k$ and $d_1 = \gcd(8, 12) = 4, (1 - r)j' - s = -4j' - 8$ at $j' = 2$ yeilds $4 \mid -16$ then $Q_2 = \{1, a^4, a^8\}$ is cyclic.
- ii):** Take $\theta_3 = 7x + 6 \in \text{Aut}(QD_{48})$ then $r = 7, s = 6 \Rightarrow r + 2s - 1 = 6 \equiv k$ and $d_1 = \gcd(6, 12) = 6, (1 - r)j' - s = -6j' - 6$ at $j' = 2$ yeilds $6 \mid -18$ and $1 - r = -6 \equiv 6 \equiv k$ then $Q_3 = \{1, a^6\}$ is cyclic. And for $\theta_4 = 11x + 4 \in \text{Aut}(QD_{48})$ then $r = 11, s = 4 \Rightarrow r + 2s - 1 = 6 \equiv k$ and $d_1 = \gcd(6, 12) = 6, (1 - r)j' - s = -10j' - 4$ at $j' = 2$ yeilds $6 \nmid -24$ and $1 - r = -10 \equiv 2 \not\equiv k$ then $Q_4 = \{1, a^4, a^6, a^8\}$ is non-cyclic.
- b):** **i):** Consider $\theta_5 = x + 10 \in \text{Aut}(QD_{48})$ then $r = 1, s = 10 \Rightarrow r + 2s - 1 = 8$ and $d_1 = \gcd(8, 12) = 4, (1 - r)j' - s = 0j' - 10 \equiv 2 \Rightarrow 4 \nmid 2$ and $-s = -10 \equiv 2 \not\equiv k$ then $Q_5 = \{1, a^2, a^4, a^8\}$ is non-cyclic.
- ii):** For $\theta_6 = 5x + 6 \in \text{Aut}(QD_{48})$ then $r = 5, s = 6 \Rightarrow r + 2s - 1 = 4$ and $d_1 = \gcd(4, 12) = 4, (1 - r)j' - s = -4j' - 6$ for any j' ,

$d_1 \nmid -4j' - 6$ and $-s = -6 \equiv 6 \equiv k$ then $Q_6 = \{1, a^2, a^4, \dots, a^{10}\}$ is cyclic.

Corollary 8.9. If $m = 4k$, $n = 2k$ and k is odd number, then Q will be cyclic or non-cyclic according to the following conditions

- (1) If $s = 2w$ is even then Q will be cyclic.
- (2) If $s = 2w + 1$ is odd
 - a): Q will be non-cyclic if either $r + 2s - 1 \equiv 0 \pmod{n}$ or $1 - r \equiv 0 \pmod{n}$.
 - b): Q will be cyclic
 - i): if $1 - r \equiv 0$ and $s \equiv k \pmod{n}$.
 - ii): if $r + 2s - 1 \not\equiv 0$ and $1 - r \not\equiv 0 \pmod{n}$.

Example 8.10. Let $G = QD_{40}$ where $m = 20$, $n = 10$, $k = 5$. Since $\gcd(r, m) = 1 \Rightarrow r \in U_{20} = \{1, 3, 7, 9, 11, 13, 17, 19\}$ and $s \in \mathbb{Z}_{10} = \{0, 1, 2, \dots, 9\}$.

- (1) For $\theta_1 = 7x + 6 \in \text{Aut}(QD_{40})$ then $r = 7$, $s = 6 \Rightarrow r + 2s - 1 = 8$ then $Q_1 = \{1, a^2, a^4, a^6, a^8\}$ is cyclic.
- (2) a): Take $\theta_2 = x + 7 \in \text{Aut}(QD_{40})$ then $r = 1$, $s = 7 \Rightarrow r + 2s - 1 = 4$ and $1 - r = 0$ then $Q_2 = \{1, a^2, a^3, a^4, a^6, a^8\}$ is non-cyclic.
- b): i): Consider $\theta_3 = 11x + 5 \in \text{Aut}(QD_{40})$ then $r = 11$, $s = 5 \Rightarrow r + 2s - 1 = 0$, $1 - r = 0$, $s = 5 \equiv k$ then $Q_3 = \{1, a^5\}$ is cyclic.
- ii): For $\theta_4 = 7x + 5 \in \text{Aut}(QD_{40})$ then $r = 7$, $s = 5 \Rightarrow r + 2s - 1 = 6$, $1 - r = -6 \not\equiv 0$ then $Q_4 = \{1, a, a^2, \dots, a^9\}$ is cyclic.

Corollary 8.11. Let $\theta = rx + s \in \text{Aut}_v(QD_{2m})$ be of finite order, where $m = 4k$, $n = 2k$ and k is even number. Then Q is isomorphic to a cyclic group as follows:

- (1) If $|Q| = 2$ then $Q = \langle a^k : a^n = 1 \rangle$ and $Q \cong \langle k \rangle$.
- (2) If $|Q| = \frac{k}{2}$ then
 - a): Let $\gcd(r + 2s - 1, n) = d_1$. If $n \nmid r + 2s - 1$ then $Q = \langle a^{d_1} : a^n = 1 \rangle$ and $Q \cong \langle d_1 \rangle$.
 - b): If $n \mid r + 2s - 1$ and $\gcd(r + 2s - 1, n) = n$ then $Q = \langle a^4 : a^n = 1 \rangle$ and $Q \cong \langle 4 \rangle$.
- (3) If $|Q| = k$ and $\gcd(r + 2s - 1, n) = d_1$ then
 - a): if $(r + 2s - 1) \cap (1 - r)j' - s = \emptyset$ then $Q = \langle a^{d_1/2} : a^n = 1 \rangle$ and $Q \cong \langle \frac{d_1}{2} \rangle$.
 - b): if $(r + 2s - 1) \cap (1 - r)j' - s \neq \emptyset$ then $Q = \langle a^{d_1} : a^n = 1 \rangle$ and $Q \cong \langle d_1 \rangle$.

Corollary 8.12. Let $\theta = rx + s \in \text{Aut}_v(QD_{2m})$ be of finite order, if k is odd number. Then the conditions for Q to be cyclic and isomorphism relationship are:

- (1) If s is even, then

- a):** if $r = 1, s = 0 \mid Q \mid = 1$ then $Q = \{1\}$ and $Q \cong \langle 0 \rangle$.
b): if $\mid Q \mid = k$ then $Q = \langle a^2 : a^n = 1 \rangle \cong \langle 2 \rangle$.
 (2) If s is odd, then
a): Let $n \mid 1 - r, n \mid s - k$. If $\mid Q \mid = 2$ then $Q = \langle a^k : a^n = 1 \rangle \cong \langle k \rangle$.
b): if $r + 2s - 1 \not\equiv 0$ and $1 - r \not\equiv 0 \pmod{n}$ and $\mid Q \mid = n$ then $Q = \langle a : a^n = 1 \rangle \cong \langle 1 \rangle$.

Corollary 8.13. Let $\theta = rx + s \in \text{Aut}(QD_{2m})$ be a fixed automorphism. Then for any H and Q , $HQ \neq QD_{2m}$.

Example 8.13.1. Consider the involution $\theta_1 = 17x + 8$ of QD_{48} . Then $H_1 = \{1, a^3, a^6, a^9, a^{12}, a^{15}, a^{18}, a^{21}, a^{24}, a^{27}, a^{30}, a^{33}, a^{36}, a^{39}, a^{42}, a^{45}\}$ and $Q_1 = \{1, a^4, a^8\} \Rightarrow H_1Q_1 \neq QD_{48}$. Now, consider the automorphism $\theta_2 = 5x + 3$ of order 4. Then $H_2 = \{1, a^6, a^{12}, a^{18}, a^{24}, a^{30}, a^{36}, a^{42}\}$ and $Q_2 = \{1, a^2, a^4, a^6, a^8, a^{10}, a^{12}, a^{14}, a^{16}, a^{18}, a^{20}, a^{22}, a^{24}, a^{26}, a^{28}, a^{30}, a^{32}, a^{34}, a^{36}, a^{38}, a^{40}, a^{42}, a^{44}, a^{46}, a^{48}\} \Rightarrow H_2Q_2 \neq QD_{48}$.

Proposition 8.14. Let $\theta = rx + s \in \text{Aut}_v(QD_{2m})$ and for any $a^l \in Q$.

- i):** If l is odd, then $H \backslash Q = \{\{a^j\} \mid j \in \langle r + 2s - 1 \rangle \cup [(1 - r)j' - s] \pmod{n}\}$
ii): If l is even then $H \backslash Q = \{\{a^j, a^{-j}\} \mid j \in \langle r + 2s - 1 \rangle \cup [(1 - r)j' - s] \pmod{n}\}$

where, j' is any even number. In either case $G \backslash Q = \{Q\}$, that is there is only a single G -orbit on Q .

Proof. Since, H is fixed by θ thus the action of H on Q is simply by conjugation. Notice that $Q \subseteq \langle a \rangle \leq QD_{2m}$ so, the action of H on general element a^l of Q . Let $a^i \in H$ then, $a^i \cdot a^l \cdot a^{-i} = a^{i+l-i} = a^l$. So, $\langle a \rangle$ fixes Q point wise.

Suppose $a^{i'}b \in H$, then $(a^{i'}b) \cdot a^l \cdot (a^{i'}b)^{-1} = a^{i'}(ba^l)b^{-1}a^{-i'} = a^{i'}(a^{l(n-1)}b)b^{-1}a^{-i'} = a^{i'+l(n-1)-i'} = a^{l(n-1)} = a^{ln} \cdot a^{-l} = (a^{2n})^{\frac{l}{2}} \cdot a^{-l} = (a^m)^{\frac{l}{2}} \cdot a^{-l} = a^{-l}$ so, $a^{i'}b$ takes a^l to $a^{-l} \Rightarrow \{a^j, a^{-j}\}$ is an orbit $\Leftrightarrow a^{i'}b \in H$ which is true $\Leftrightarrow l$ is even.

Now, for every element of Q is in G -orbit of $1 \in Q$. Since, every element of Q are of the form $a^l \in Q$. and G has two type of elements a^i and a^ib . Now let us supposed that $q_1 = a^l = a^i \cdot \theta(a^i)^{-1}$ for some $a^i \in G$, thus $q_1 = a^i \cdot \theta(a^{-i}) = a^i \cdot a^{-ri+[2-n]y} = a^{i-ri+[2-n]y} = a^{-i(r-1)+(2-n)y} = a^{v(r-1)+(2-n)y}$, where $v = -i$.

Also, $a^l = (a^ib) \cdot \theta(a^ib)^{-1}$ for some $a^ib \in G \Rightarrow a^l = (a^ib) \cdot \theta(a^ib) = (a^ib) \cdot (a^{ri+y}b) = a^i(ba^{ri+y})b = a^i \cdot a^{(ri+y)(n-1)}b \cdot b = a^{i+(ri+y)(n-1)}b^2$ Let us denote this element $q_2 = a^l = a^{-v+(y-vr)(n-1)}$, where $v = -i$.

For $a^{-v} \in G$, we have $a^{-v} \cdot 1 \cdot \theta(a^{-v})^{-1} = a^{-v} \cdot \theta(a^v) = a^{-v} \cdot a^{rv+(2-n)y} = a^{v(r-1)+(2-n)y} = q_1$. Now for $a^{-v}b \in G$, we have $(a^{-v}b) \cdot 1 \cdot \theta(a^{-v}b)^{-1} = (a^{-v}b) \cdot \theta(a^{-v}b) = (a^{-v}b) \cdot a^{-rv+y}b = a^{-v}(ba^{-rv+y})b = a^{-v} \cdot a^{(-rv+y)(n-1)}b \cdot b = a^{-v+(y-rv)(n-1)} = q_2$.

Therefore, for any $q \in Q$ there exist $g \in G$ such that $g \cdot 1 \cdot \theta(g)^{-1} = q$. Hence $G \backslash Q = \{Q\}$, there is a single G -orbit on Q . $\square$

Example 8.14.1. Revisiting θ_1, θ_2 of Example 8.13.1 applying Proposition 8.14 to obtain that for θ_1 . Since all l are even, therefore $H_1 \backslash Q_1 = \{\{1\}, \{a^4, a^{-4}\}, \{a^8, a^{-8}\}\}$,

for θ_2 there exists even and odd powers of elements of Q . For all even powers $H_2 \setminus Q_2 = \{\{1\}, \{a^2, a^{-2}\}, \dots, \{a^{10}, a^{-10}\}, \{a, a^{-1}\}, \{a^5, a^{-5}\}, \{a^9, a^{-9}\}\}$ and odd powers $H_2 \setminus Q_2 = \{\{1\}, \{a^2\}, \dots, \{a^{10}\}, \{a\}, \{a^5\}, \{a^9\}\}$.

9. INVOLUTIONS IN $Aut(QD_{2m})$

If $\theta = rx + s$ is an automorphism such that $\theta^2 = \text{id}$, then Proposition 7.1 gives $r \in R_m^2$ and $s \in Z\text{Div}(r+1)$.

Proposition 9.1. Suppose that $m = 4p_1^{t_1}p_2^{t_2}$, where the p_i are distinct odd primes. Then $|R_m^2| = 8$

Proof. Since, $m = 4p_1^{t_1}p_2^{t_2} \Rightarrow n = 2p_1^{t_1}p_2^{t_2}$. Suppose that $r \in \mathbb{Z}_n$ with $r^2 \equiv 1 \pmod{n}$ $\because \gcd(r, m) = 1 \Rightarrow \gcd(r, 4p_1^{t_1}p_2^{t_2}) = 1$. But m is even therefore, r is odd. Assume that $r = 2q + 1$ for some $0 \leq q < (n-1)$. Now, $r^2 = (2q+1)^2 = 4q^2 + 4q + 1 \Rightarrow 4q^2 + 4q \equiv 0 \pmod{n} \Rightarrow 4q(q+1) = l \cdot (2p_1^{t_1}p_2^{t_2}) \because \gcd(4, 2p_1^{t_1}p_2^{t_2}) = 2 \Rightarrow 2q(q+1) = l \cdot (p_1^{t_1}p_2^{t_2}) \pmod{p_1^{t_1}p_2^{t_2}}$. Either q or $q+1$ is odd.

- (1) If q is even, then $q = \frac{l}{2(q+1)} \cdot p_1^{t_1}p_2^{t_2} \Rightarrow q = h \cdot p_1^{t_1}p_2^{t_2}$ where $1 \leq h \leq 4$. Now, $r = 2q + 1 \Rightarrow r = 2h \cdot p_1^{t_1}p_2^{t_2} + 1 \Rightarrow r \equiv 1 \pmod{n}$
- (2) If $q+1$ is even, then $q+1 = \frac{l}{2q} \cdot p_1^{t_1}p_2^{t_2} \Rightarrow q+1 = h \cdot p_1^{t_1}p_2^{t_2} \Rightarrow q = h \cdot p_1^{t_1}p_2^{t_2} - 1$ where $1 \leq h \leq 4$. Now, $r = 2q + 1 \Rightarrow r = 2h \cdot p_1^{t_1}p_2^{t_2} - 1 \Rightarrow r \equiv -1 \pmod{n} \Rightarrow r \equiv \pm 1 \pmod{n}$,

□

Proposition 9.2. Suppose that $m = 2^\alpha p_1^{t_1}p_2^{t_2} \dots p_q^{t_q}$ where the p_i are distinct odd primes. Then

$$|R_m^2| = \begin{cases} 2^{q+1} & \alpha = 2 \\ 2^{q+2} & \alpha = 3 \\ 2^{q+3} & \alpha \geq 4. \end{cases}$$

Proof. Similar to the Propositions 10.1. □

Example 9.3. Let $G = QD_{192}$, then $U_{96} = \{1, 5, 7, 11, 13, 17, 19, 23, 25, 29, 31, 35, 37, 41, 43, 47, 49, 53, 55, 59, 61, 65, 67, 71, 73, 77, 79, 83, 85, 89, 91, 95\}$ and $R_{96}^2 = \{1, 7, 17, 23, 25, 31, 41, 47, 49, 55, 65, 71, 73, 79, 89, 95\}$ thus, $|R_{96}^2| = 2^4 = 16$

Corollary 9.4.

$$|Aut_2(QD_{2m})| = \sum_{r \in R_m^2} \gcd(r+1, n).$$

Theorem 9.5. Let $r \in R_m^2$, then the followings hold:

- (1) $\langle r-1 \rangle \leq Z\text{Div}(r+1)$
- (2) $N_r = \left| \frac{Z\text{Div}(r+1)}{\langle r-1 \rangle} \right| \leq 2$

(3) $N_r = 2$ if and only if $m = 2^\alpha l$; $\alpha \geq 2$, and l is odd, $r \equiv \pm 1 \pmod{2^{\alpha-1}}$.

Proof. (1) Given that $r \in R_m^2$, then $(r-1).(r+1) = r^2 - 1 \equiv 0 \Rightarrow (r-1) \leq ZDiv(r+1) \Rightarrow \langle r-1 \rangle \leq ZDiv(r+1)$.

(2) Suppose

$$(9.1) \quad \left| \frac{ZDiv(r+1)}{\langle r-1 \rangle} \right| = j$$

It is well known that $|\langle r-1 \rangle| = \frac{n}{\gcd(r-1, n)}$ and $|ZDiv(r+1)| = \gcd(r+1, n)$
put values in 9.1

$$\frac{\gcd(r+1, n)}{\frac{n}{\gcd(r-1, n)}} = j$$

$$(9.2) \quad \Rightarrow \gcd(r+1, n). \gcd(r-1, n) = n.j$$

Now, suppose that $m = 2^\alpha p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}$ where $\alpha \geq 2 \Rightarrow \alpha - 1 \geq 1$

$\Rightarrow n = 2^{\alpha-1} p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}$ where p_i are distinct odd primes and $t_i \geq 1$. Then

$$\gcd(r+1, n) = 2^{\alpha_1} p_1^{a_1} p_2^{a_2} \dots p_q^{a_q}$$

with $\alpha_1 \leq (\alpha - 1)$ and $0 \leq a_i \leq t_i \forall i$ and

$$\gcd(r-1, n) = 2^{\alpha_2} p_1^{b_1} p_2^{b_2} \dots p_q^{b_q}$$

with $\alpha_2 \leq (\alpha - 1)$ and $0 \leq b_i \leq t_i$. Now, for all $i \in \{1, 2, \dots, q\}$ either $a_i = 0$ or $b_i = 0$. Indeed, if $a_i > 0$ and $b_i > 0$ then p_i divides both $(r+1)$ and $(r-1)$ which is impossible. $\therefore (p_i > 2)$

Similarly, either $\alpha_1 \leq 1$ or $\alpha_2 \leq 1$. Otherwise $2^{\min\{\alpha_1, \alpha_2\}}$ divides $(r+1)$ and $(r-1)$ which is impossible.

Since, from 9.2

$$\gcd(r+1, n). \gcd(r-1, n) = n.j$$

$$(2^{\alpha_1} p_1^{a_1} p_2^{a_2} \dots p_q^{a_q}). (2^{\alpha_2} p_1^{b_1} p_2^{b_2} \dots p_q^{b_q}) = j. 2^{\alpha-1} p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}$$

$$(2^{\alpha_1+\alpha_2} p_1^{a_1+b_1} p_2^{a_2+b_2} \dots p_q^{a_q+b_q}) = j. 2^{\alpha-1} p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}$$

Now, since $\forall i$ either $a_i = 0$ or $b_i = 0 \Rightarrow a_i + b_i = t_i$

$$2^{\alpha_1+\alpha_2} = j. 2^{\alpha-1} \frac{p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}}{p_1^{a_1+b_1} p_2^{a_2+b_2} \dots p_q^{a_q+b_q}}$$

$$(9.3) \quad 2^{\alpha_1+\alpha_2} = j. 2^{\alpha-1}.$$

As $\alpha_1 \leq (\alpha - 1)$ and $\alpha_2 \leq (\alpha - 1)$ and since either $\alpha_1 \leq 1$ or $\alpha_2 \leq 1$,
 $\Rightarrow (\alpha - 1) \leq (\alpha_1 + \alpha_2) \leq \alpha$

equation 9.3 $\Rightarrow j = 2^{\alpha_1+\alpha_2-\alpha+1} \Rightarrow 1 \leq j \leq 2$.

(3) By Theorem 7.8, N_r is the number of divisors of

$$\frac{\gcd(r-1, n) \cdot \gcd(r+1, n)}{n}$$

and N_r is the number of divisor of j computed above. Therefore,

$$N_r = 1 \text{ if } j = 1 \text{ and } N_r = 2 \text{ if } j = 2.$$

In either case,

$$\left| \frac{ZDiv(r+1)}{\langle r-1 \rangle} \right| \leq 2.$$

Finally according to the proof of (2), $j = 2$ if and only if

$$\alpha_1 = \alpha - 1, \alpha_2 = 1 \text{ or } \alpha_1 = 1, \alpha_2 = \alpha - 1.$$

If $\alpha_1 = \alpha - 1, \alpha_2 = 1$ then $2^{\alpha-1}$ divides $(r+1)$.

If $\alpha_1 = 1, \alpha_2 = \alpha - 1$ then $2^{\alpha-1}$ divides $(r-1)$.

$$\Rightarrow r \equiv \pm 1 \pmod{2^{\alpha-1}}.$$

□

Example 9.6. Let $G = QD_{48}$ where $m = 2^3 \cdot 3$, $\alpha = 3$, then $U_{24} = \{1, 5, 7, 11, 13, 17, 19, 23\}$ and $R_{24}^2 = \{1, 5, 7, 11, 13, 17, 19, 23\}$.

- (1) Let $r = 5 \in R_{24}^2$ then $\langle r-1 \rangle = \langle 4 \rangle = \{0, 4, 8\} = B_1$ and $ZDiv(r+1) = ZDiv 6 = \{0, 2, 4, 6, 8, 10\} = B_2$ clearly, $\langle 4 \rangle \leq ZDiv 6$.
- (2) For $r = 5$, $\frac{ZDiv(6)}{\langle 4 \rangle} = \frac{\{0, 2, 4, 6, 8, 10\}}{\{0, 4, 8\}} = \frac{B_2}{B_1} = x + B_1 \forall x \in B_2$
 $0 + B_1 = \{0, 4, 8\} = B_1, 2 + B_1 = \{2, 6, 10\} = B_3$
 $4 + B_1 = \{4, 8, 0\} = B_1, 6 + B_1 = \{6, 10, 2\} = B_3$
 $8 + B_1 = \{8, 0, 4\} = B_1, 10 + B_1 = \{10, 2, 6\} = B_3$
 U_{24} action on $\frac{ZDiv 6}{\langle 4 \rangle} = \{B_1, B_2\} \Rightarrow N_r = N_5 = 2$.
- (3) Since, $m = 2^3 \cdot 3 \Rightarrow l = 3 > 1$ is odd and $r = 5 \equiv 1 \pmod{2^2}$.
Thus $N_5 = 2$ as calculated in (2).

Corollary 9.7. Suppose that $m = 2^\alpha p_1^{t_1} p_2^{t_2} \dots p_q^{t_q}$, where the p_i are distinct odd primes and $\alpha \geq 2$. Then the number of equivalence classes C_{4k} of $Aut_2(QD_{8k})$ is given by

$$C_{4k} = \begin{cases} 2^{q+1} & \alpha = 2 \\ 2^{q+2} & \alpha \geq 3. \end{cases}$$

Example 9.8. Let $G = QD_{384}$. Thus $R_{192}^2 = \{1, 17, 31, 47, 49, 65, 79, 95, 97, 113, 127, 143, 145, 161, 175, 191\} \Rightarrow |R_{192}^2| = 2^4$. It is easy to check that the values $r = 1, 65, 97, 161$ satisfy $r \equiv 1 \pmod{2^5}$ and the values $r = 31, 95, 127, 191$ satisfy $r \equiv -1 \pmod{2^5}$. So, $C_m = |\{1, 31, 65, 95, 97, 127, 161, 191\}| = 2^3$.

Now for θ an involution, the set of twisted involutions is given in the followings porosities:

Proposition 9.9. If $\theta^2 = \text{id}$, then the set of twisted involutions is given as $R = \{a^i \mid (r+1)i \equiv -2s \pmod{n}\} \cup \{a^i b \mid (r-1)i \equiv -s \pmod{n}\}$

Proof. Since, θ is an involution therefore, $r \in R_m^2$ and $s \in Z\text{Div}(r+1)$. To find the set R , let $a^i \in G$, then $\theta(a^i) = (a^i)^{-1} \Rightarrow a^{ri+(2-n)s} = a^{-i} \Rightarrow ri + (2-n)s = -i \Rightarrow (r+1)i + (2-n)s = 0 \Rightarrow (r+1)i \equiv -2s \pmod{n}$.

Now for $a^i b \in G$, we have, $\theta(a^i b) = (a^i b)^{-1} \Rightarrow a^{ri+s} b = a^i b \Rightarrow ri + s = i \Rightarrow (r-1)i \equiv -s \pmod{n}$, hence $R = \{a^i \mid (r+1)i \equiv -2s \pmod{n}\} \cup \{a^i b \mid (r-1)i \equiv -s \pmod{n}\}$. $\square$

Corollary 9.10. If $\theta^2 = \text{id}$, $m = 4k$, $n = 2k$ and k is even number, then Q is a subgroup of $\langle a \rangle$ and the natural isomorphism $\psi : \langle a \rangle \rightarrow \mathbb{Z}_n$ gives using Theorem 8.1 and Corollary 8.7 and 8.11

- (1) If $|Q| = 2$, then $\psi(Q) = \langle k \rangle$.
- (2) If $|Q| = \frac{k}{2}$ then

$$\psi(Q) = \begin{cases} \langle d_1 \rangle & \text{If } n \nmid r+2s-1 \text{ and } d_1 = \gcd(r+2s-1, n) \\ \langle 4 \rangle & \text{If } n \mid r+2s-1 \text{ and } n = \gcd(r+2s-1, n). \end{cases}$$

- (3) If $|Q| = k$ and $\gcd(r+2s-1, n) = d_1$ then

$$\psi(Q) = \begin{cases} \langle \frac{d_1}{2} \rangle & \text{If } \{(r+2s-1) \cap [l(1-r)-s]\} = \emptyset \\ \langle d_1 \rangle & \text{If } \{(r+2s-1) \cap [l(1-r)-s]\} \neq \emptyset, \end{cases}$$

where l is an even number.

Corollary 9.11. If $\theta^2 = \text{id}$, $m = 4k$, $n = 2k$ and k is even number, then Q is a subgroup of $\langle a \rangle$ and the natural isomorphism $\psi : \langle a \rangle \rightarrow \mathbb{Z}_n$ gives using Theorem 8.1 and Corollary 8.9 and 8.12

- (1) If s is even, then

$$\psi(Q) = \begin{cases} \langle 0 \rangle & \text{If } r = 1, s = 0, |Q| = 1 \\ \langle 2 \rangle & \text{If } |Q| = k. \end{cases}$$

- (2) If s is odd, then

$$\psi(Q) = \begin{cases} \langle k \rangle & |Q| = 2 \text{ If } n \mid 1-r \text{ and } n \mid s-k \\ \langle 1 \rangle & |Q| = n \text{ If } n \nmid r+2s-1 \text{ and } n \nmid 1-r. \end{cases}$$

Corollary 9.12. In any Case $R \neq Q$.

REFERENCES

- [1] A. Bishop, C. Cyr, J. Hutchens, C. May, N. Schwartz, and B. Turner (2013). "On involutions and generalized symmetric spaces of dicyclic groups". arXiv:1310.0121v1
- [2] Anders Bjorner and Francesco Brenti (2005). "Combinatorics of coxeter groups". Graduate Text in Mathematics, vol.231, Springer, New York.

- [3] Nicolas Bourbaki (2002). "Lie groups and Lie algebras". Chapter 4-6, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, Translated from the (1968) French original by Andrew Pressley.
- [4] K. K. A. Cunningham, T. J. Edgar, A. G. Helminck, B. F. Jones, OH, R. Schwell, J.F. Vasquez. (2012). "On the Structure of Involution and Symmetric Spaces of Dihedral Groups". arXiv:1205.3207v1.
- [5] A. G. Helminck (1994). Symmetric k-varieties. Proc. Sympos. Pure Math. 56, no. 1, 233-279.
- [6] A. G. Helminck (1988). "Algebraic groups with a commuting pair of involutions and semi-simple symmetric spaces". Adv. in Math 71, 21-91.
- [7] A. G. Helminck and S.P. Wang (1993). "On rationality properties of involutions of reductive groups". Adv. in Math. 99, 26-96.
- [8] Gary L. Walls (1986). "Automorphism groups". Amer. Math. Monthly 93, no. 6, 459-462.

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES, UNIVERSITY OF SHARJAH, UAE.
E-mail address: `zraza@sharjah.ac.ae`

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF COMPUTER AND EMERGING SCIENCES, LAHORE, PAKISTAN.
E-mail address: `imran_aimphd@yahoo.com`

DEPARTMENT OF MATHEMATICS, YAZD UNIVERSITY, YAZD, IRAN
E-mail address: `davvaz@yazd.ac.ir`