

Effective Hamiltonians for rapidly driven many-body lattice systems

A.P. Itin^{1,2}, M.I. Katsnelson^{1,3}

¹*Radboud University Nijmegen, Institute for Molecules and Materials (IMM), The Netherlands,*

²*Space Research Institute, Russian Academy of Sciences, Moscow, Russia,*

³*Dept. of Theoretical Physics and Applied Mathematics, Ural Federal University, Ekaterinburg, Russia*

We consider 1D lattices described by Hubbard or Bose-Hubbard models, in the presence of periodic high-frequency perturbations, such as uniform ac force or modulation of hopping coefficients. Effective Hamiltonians for interacting particles are derived using an averaging method resembling classical canonical perturbation theory. As is known, a high-frequency force may renormalize hopping coefficients, causing interesting phenomena such as coherent destruction of tunnelling and creation of artificial gauge fields. We find explicitly additional corrections to the effective Hamiltonians due to interactions, corresponding to non-trivial processes such as single-particle density-dependent tunnelling, correlated pair hoppings, nearest neighbour interactions, etc. Some of these processes arise also in multiband lattice models, and are capable to give rise to a rich variety of quantum phases. The apparent contradiction with other methods, e.g. Floquet-Magnus expansion, is explained. The results may be useful for designing effective Hamiltonian models in experiments with ultracold atoms, as well as in the field of ultrafast nonequilibrium magnetism. An example of manipulating exchange interaction in a Mott-Hubbard insulator is considered, where our corrections play an essential role.

The idea of engineering effective Hamiltonians using high-frequency perturbations probably goes back to the famous Kapitza pendulum [1–4]. In classical and celestial mechanics there are many examples of systems with separation of typical timescales on slow and fast ones, and corresponding perturbation methods were developed long time ago (see, e.g., [5]). It is interesting to adopt these methods to the quantum realm, especially to lattice systems, where high-frequency perturbations are often used, e.g. for construction of quantum simulators (well-controllable quantum systems for simulating complicated condensed matter phenomena [6–8]). Indeed, a suitable driving applied to ultracold atoms in optical lattices allows to realize dynamical localization [9, 10]; mimic photoconductivity [11]; simulate artificial gauge fields [12–14], classical and quantum magnetism [15]; study transport phenomena [16, 17] and phase transitions [18, 19]. In solid-state physics, ac-driven systems are interesting in the context of dynamical localization [22, 23], coherent control of transport [24, 25]; microwave-induced topological insulators [26]; photoinduced quantum Hall insulators [27], metal-insulator transitions [28, 29], and superconductivity [30]. The most of the abovementioned applications are in fact based on modifying single-particle hopping amplitudes by a high-frequency force, a phenomenon that can be derived by averaging a Hamiltonian of the system (i.e., keeping only zeroth order terms in $\frac{1}{\omega}$ (inverse frequency of perturbation) and neglecting all higher-order terms. However, for many realistic applications of such type, it is important to derive accurate effective Hamiltonians taking into account higher-order terms (see, e.g., [31]). Here we determine explicitly higher-order corrections to effective Hamiltonians of driven quantum lattices by elaborating a method [32–34] inspired by canonical perturbation theory. The corrections correspond to non-trivial (many-body) processes such as single-particle density-dependent tunnelling, cor-

related pair hoppings, nonlocal (extended) pair hopping, and so on. Using a suitable driving, we are able to suppress or enhance a particular process in the effective Hamiltonian. Such approach can be very useful for engineering particular Hamiltonians, for simulating solid-state phenomena via optical means, for accurate interpretation of experiments with driven lattice systems, etc. A particular application to a recent insightful proposal of ultrafast and reversible control of exchange interactions [35] is demonstrated.

The method can be seen either as a modification of the method of [34], or as a modification of Magnus expansion approach [36, 37]. Surprisingly, they can lead to different results. Correspondingly, some related methods available in the literature allow to obtain results that are in accord with ours [31, 34], while others [36, 42] may lead to apparently different Hamiltonians. Especially the subtle difference between our approach and modifications of Magnus expansion method [36] is important. Magnus expansion [36, 37] is a popular tool in physics and mathematics with a rapidly growing number of applications. It is interesting that sometimes accuracy of effective Hamiltonians produced by this method can be drastically increased, as explained below and in SI (note that Magnus expansion was actually designed for solving an initial value problem, not for deriving effective Hamiltonians). To clarify our approach, let us first start with Schrödinger equation for a lattice system (e.g., a single particle in a driven 1D tight-binding model), written in the matrix form:

$$i\dot{X} = \epsilon \mathcal{H}X, \quad (1)$$

where $\epsilon = \frac{1}{\omega}$ is a small parameter, ω is a frequency of perturbation, $\mathcal{H}(t)$ is a time-dependent (matrix) Hamiltonian of the system, X is a column of coefficients of expansion of a quantum state in a certain basis, and fast time was introduced ($t \rightarrow t/\epsilon$), resulting in the small coefficient ϵ in front of r.h.s of Eq.(1), to put high-frequency




FIG. 1: (Color online) Exchange interaction J_{ex} in the driven fermionic two-site model as a function of the driving strength \mathcal{E}_0 . Parameters: $U = 10$, 'bare' tunnelling $J = -1$, $\omega = 16$. Harmonic driving changes effective tunnelling constant ($J \rightarrow J_e = J J_0(\mathcal{E}_0)$) affecting the exchange interaction in the zeroth order in $\frac{1}{\omega}$: $J_{ex}^{(0)} = \frac{2J_0^2(\mathcal{E}_0)}{U}$. Solid curves, from up to down: 'bare' exchange interaction $J_{ex}^{(0)}$; theoretical prediction $J_{ex}^{(2)}$ with corrections up to the second order in $\frac{1}{\omega}$ taken into account; $J_{ex}^{(4)}$ with corrections up to the fourth order in $\frac{1}{\omega}$ taken into account; $J_{ex}^{(\infty)}$ with infinite order of terms in $\frac{1}{\omega}$ expansion taken into account (with leading order in U contribution in each term) (the lowest solid curve). Dashed curve (nearly indistinguishable from the lowest solid curve): numerical value of the exchange interaction (kindly provided by J.Mentink). In the regions where J_e is strongly suppressed by driving (around zeros of $J_{ex}^{(0)}$), the second order corrections become insufficient, and one needs to include fourth and higher orders in $\frac{1}{\omega}$.

dependence of the Hamiltonian in explicit form. We may consider Eq.(1) as a classical dynamical system for variables X_n (components of the column X). We adopt then a classical averaging method [5, 32, 38–40] to this system, in such a way that allows convenient generalization to many-body systems [32]. To this end, one makes a unitary transformation $X = C\tilde{X}$ so that equations for the transformed variables are

$$i\dot{\tilde{X}} = [C^{-1}\epsilon\mathcal{H}C - iC^{-1}\dot{C}]\tilde{X}. \quad (2)$$

The expression in the square brackets is the new Hamiltonian. To get an effective time-independent Hamiltonian, the transformation is sought in the form $C = \exp[\epsilon K_1 + \epsilon^2 K_2 + \epsilon^3 K_3 + \dots]$, where K_i are skew-Hermitian time-periodic matrices (with zero mean), which would remove time-dependent terms from the Hamiltonian, leaving only time-independent ones. An iterative procedure analogous to the Hamiltonian averaging method give us [32]

$$\begin{aligned} i\dot{K}_1 &= \mathcal{H}(t) - \langle \mathcal{H}(t) \rangle \equiv \{\mathcal{H}\}, & iK_1 &= \int \{\mathcal{H}\} dt, \\ i\dot{K}_2 &= \left\{ \mathcal{H}K_1 - K_1\mathcal{H} - \frac{i}{2}(K_1\dot{K}_1 - K_1\dot{K}_1) \right\}, \\ \epsilon\mathcal{H}_{eff} &= [C^{-1}\epsilon\mathcal{H}C - iC^{-1}\dot{C}] = \epsilon\mathcal{H}_0 + \epsilon^2\mathcal{H}_1 + \dots, \end{aligned}$$

where curly brackets denote taking the time-periodic part of a time-dependent function: $\{\mathcal{H}\} \equiv \mathcal{H} - \langle \mathcal{H}(t) \rangle$, where $\langle \mathcal{H}(t) \rangle \equiv \frac{1}{2\pi} \int_0^{2\pi} \mathcal{H}(t') dt'$. Indefinite integrals above are defined up to an additive constant, which is chosen in such a way that $\langle K_i \rangle = 0$. We have

$$\begin{aligned} \mathcal{H}_0 &= \langle \mathcal{H} \rangle, & \mathcal{H}_1 &= \frac{1}{2} \langle \{\mathcal{H}\}, K_1 \rangle, \\ \mathcal{H}_2 &= \frac{1}{2} \langle \{\mathcal{H}\}, K_2 \rangle + \frac{1}{12} \langle \{\{\mathcal{H}\}, K_1\}, K_1 \rangle, \dots \end{aligned} \quad (3)$$

where square brackets denote matrix commutation: $[A, B] = AB - BA$. This procedure resembles Floquet-Magnus expansion [36], however there is an important difference due to lifting unnecessary requirement $K_i(0) = 0$ present in that method. As detailed in SI, it often allows to remove the correction \mathcal{H}_1 , therefore obtaining much more accurate effective Hamiltonians, where corrections to \mathcal{H}_0 in the expansion start from \mathcal{H}_2 .

Eqs. (3) allow to consider, e.g., a particle in driven tight-binding models with various boundary conditions, as well as with additional external potentials (see [32] and SI). The general Eqs. (3) (which are in agreement with method of Ref. [33, 34]) are also convenient for studying many-body lattice systems. Indeed, in the case of many particles, one can construct a corresponding Hamiltonian matrix and fulfil the same transformations. Moreover, it is not necessary to consider Hamiltonians (3) in the matrix representation: one can use, e.g., creation and annihilation operators. Indeed, consider now 1D Bose-Hubbard model with a strong high-frequency driving:

$$H = H_{BH} + H_d(t), \quad H_d(t) = \omega\mathcal{E}(\omega t) \sum_j j n_j,$$

$$H_{BH} = J \sum_i (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) + U \sum_i n_i (n_i - 1) \quad (4)$$

where J is the hopping parameter, U is the interaction strength, c^\dagger, c are bosonic creation and annihilation operators. Following the approach we used in the single-particle case, we make a preliminary transformation $U^{(0)}(t) = \exp[i f(\omega t) \sum_j n_j]$, $f(\omega t) \equiv \int_0^t \omega\mathcal{E}(\omega t') dt'$, and make rescaling of time, $t' = \omega t = t/\epsilon$, so that the new Hamiltonian is $\epsilon\mathcal{H}(t) = \epsilon(\mathcal{H}_0 + \delta H(t))$, where

$$\begin{aligned} \mathcal{H}_0 &= \sum_i J_{eff} (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) + U \sum_i n_i (n_i - 1), \\ \delta H(t) &= \sum_i [\delta^+(t) c_i^\dagger c_{i+1} + \delta^-(t) c_{i+1}^\dagger c_i], \\ J_{eff} &= J \langle e^{if(t)} \rangle, \quad \delta^\pm(t) = J [e^{\pm if(t)} - \langle e^{if(t)} \rangle] \end{aligned} \quad (5)$$

In other words, the new Hamiltonian is $\epsilon\mathcal{H}(t) = \epsilon \left[\sum_i (\delta_0^+(t) c_i^\dagger c_{i+1} + \delta_0^-(t) c_{i+1}^\dagger c_i) + U \sum_i n_i (n_i - 1) \right]$, where $\delta_0^\pm = J e^{\pm if(t)}$.

The unitary transformations we fulfilled in the single-particle case should be done here as well.

However, to find operators $K_1, K_2, \mathcal{H}_1, \mathcal{H}_2$ explicitly, we fulfil more complicated many-body calculations. In the first order, we have $\dot{K}_1 = -i\{\mathcal{H}\} = -i\epsilon \sum_i [\delta^+(t)c_i^\dagger c_{i+1} + \delta^-(t)c_{i+1}^\dagger c_i]$, and therefore $K_1 = -i\epsilon \sum_i [\delta_1^+(t)c_i^\dagger c_{i+1} + \delta_1^-(t)c_{i+1}^\dagger c_i]$, $\delta_1^\pm = \int \delta^\pm(t')dt'$. Commutators $[\mathcal{H}, K_1], [[\mathcal{H}, K_1], K_1]$, etc, are derived in SI. The first-order terms in the new Hamiltonian (before averaging) are $[\mathcal{H}, K_1] = -2\epsilon iU \sum_j \delta_1^+ c_j^\dagger (n_j - n_{j+1})c_{j+1} - \delta_1^- c_{j+1}^\dagger (n_j - n_{j+1})c_j$. These terms resemble somehow the result of calculations of [42] (the structure of Hamiltonian is the same, but the coefficients δ_1^\pm are different). However, in our approach this contribution disappears during time-averaging (being averaged to zero: $\mathcal{H}_1 = \frac{1}{2}\langle\{\mathcal{H}\}, K_1\rangle = 0$), and therefore to find the non-vanishing contribution to the effective Hamiltonian, we need to consider the next orders of perturbation, exactly as in the single-particle case. This gives us (see SI)

$$\mathcal{H}_2 = -2U(\Delta^+ \hat{a}_1 + h.c.) - 2U\Delta_0 \hat{a}_2, \quad (6)$$

where

$$\begin{aligned} \Delta^+ &= \frac{1}{2}\langle\delta_2^+ \delta^+\rangle, \quad \Delta_0 = \frac{1}{2}\langle\delta^+ \delta_2^- + \delta_2^+ \delta^-\rangle, \\ \hat{a}_1 &= \sum_j \left(c_{j-1}^\dagger (4n_j - n_{j+1} - n_{j-1})c_{j+1} \right. \\ &\quad \left. - 2c_j^\dagger c_j^\dagger c_{j+1} c_{j+1} \right), \quad \delta_2^\pm = \int \delta_1^\pm(t')dt', \\ \hat{a}_2 &= \sum_j \left[4n_j n_{j+1} - 2n_j(n_j - 1) - (c_{j-1}^\dagger c_{j+1}^\dagger c_j^2 + H.c.) \right] \end{aligned} \quad (7)$$

The effective Hamiltonian contains nearest-neighbour interactions and several types of correlated tunnelling processes (pair tunnelling $c_j^\dagger c_j^\dagger c_{j+1} c_{j+1}$ and pair "dissociation"/"association" process $c_{j-1}^\dagger c_{j+1}^\dagger c_j c_j / c_j^\dagger c_j^\dagger c_{j+1} c_{j-1}$). Due to these terms, one may expect a rich phase diagram of the driven system. In particular, it is known that extended Bose-Hubbard model, obtained by adding nearest-neighbour interactions to the Bose-Hubbard Hamiltonian, poses a supersolid phase.

We note that in the particular case of harmonic perturbation $\mathcal{E} = -\mathcal{E}_0 \sin t$, we have $\Delta^+ = -\sum_{k=1}^{\infty} (-1)^k \frac{J_k^2(\mathcal{E}_0)}{k^2}$. This is a decaying oscillatory function of \mathcal{E}_0 which can be either positive or negative, and by varying the amplitude \mathcal{E}_0 one can either maximize its absolute value (e.g., at $\mathcal{E}_0 = 1.77$), or put it to zero (e.g., at $\mathcal{E}_0 = 3.33$). At the same time, $\Delta_0 = -\sum_{k=1}^{\infty} \frac{J_k^2(\mathcal{E}_0)}{k^2}$, which is a non-vanishing oscillatory function of \mathcal{E}_0 with local minima at $\mathcal{E}_0 = 1.93, 5.32$, etc. We see that using a suitable driving, it is possible to suppress or enhance particular processes in the effective Hamiltonian.

It is worth to consider a classical limit $n_i \gg 1$, where one gets a driven Discrete Nonlinear Schrödinger Equation (DNLSE). Applying canonical perturbation theory to the driven DNLSE we get an effective Hamiltonian which is indeed the classical limit of Eq.(6), i.e. it can be obtained from the Eq.(6) by replacing operators with c -numbers (SI).

Consider now the driven Hubbard model

$$H = H_H + H_d(t), \quad H_d(t) = \omega \mathcal{E}(\omega t) \sum_j j n_j, \quad (8)$$

$$H_H = J \sum_{i,\sigma} (c_{i,\sigma}^\dagger c_{i+1,\sigma} + c_{i+1,\sigma}^\dagger c_{i,\sigma}) + U \sum_i n_{i,\sigma} n_{i,-\sigma}$$

(with $c_{i,\sigma}^\dagger, c_{i,\sigma}$ being fermionic creation and annihilation operators) which becomes, after the preliminary transformation discussed above,

$$\mathcal{H} = \sum_{i,\sigma} \left(\delta_0^+ c_{i,\sigma}^\dagger c_{i+1,\sigma} + \delta_0^- c_{i+1,\sigma}^\dagger c_{i,\sigma} \right) + U \sum_i n_{i,\sigma} n_{i,-\sigma}.$$

Calculations analogous to the Bose-Hubbard model case give us (see SI)

$$\begin{aligned} \mathcal{H}_1 &= 0, \quad \mathcal{H}_2 = U\Delta_0(-2\mathcal{S} + \mathcal{A} + \mathcal{A}^\dagger + 4\mathbf{V} - 2V) \\ &\quad + U[\Delta^-(4\mathbf{R} - 2\mathcal{R} + R_2) + H.c.], \end{aligned} \quad (9)$$

where $\Delta^- = (\Delta^+)^*$, and several types of tunnelling and interaction processes were denoted as $\mathcal{S}, \mathcal{A}, \mathcal{R}, \mathbf{R}, R_2, V$, and \mathbf{V} . \mathcal{S} is the kinetic exchange interaction. \mathcal{A}^\dagger denotes a tunnelling process where a pair of atoms at j th site is dissociated into two atoms at neighbouring sites ($j-1$ and $j+1$). \mathcal{A} denotes a corresponding correlated tunnelling process of association of two atoms into a pair. \mathcal{R} (\mathcal{R}^\dagger) denotes a correlated tunnelling process where an *extended pair* of atoms is tunnelling to the right (left) neighbouring sites. \mathbf{R} (\mathbf{R}^\dagger) denotes tunnelling of a localised pair of atoms to the right (left). R_2 (R_2^\dagger) denotes single-particle next-nearest-neighbour density-dependent tunnelling to the right (left); V is the nearest-neighbour interaction, and \mathbf{V} is the usual local interaction.

$$\begin{aligned} \mathcal{S} &= \sum_{j,\sigma} c_{j+1,\sigma}^\dagger c_{j,\sigma} c_{j,-\sigma}^\dagger c_{j+1,-\sigma}, \\ \mathcal{A} &= \mathcal{A}_{\uparrow,\downarrow} + \mathcal{A}_{\downarrow,\uparrow}, \quad \mathcal{A}_{\sigma,-\sigma} = \sum_j c_{j,\sigma}^\dagger c_{j-1,\sigma} c_{j,-\sigma}^\dagger c_{j+1,-\sigma}, \\ \mathcal{R} &= \mathcal{R}_{\uparrow\downarrow} + \mathcal{R}_{\downarrow\uparrow}, \quad \mathcal{R}_{\sigma,-\sigma} = \sum_j c_{j,\sigma}^\dagger c_{j-1,\sigma} c_{j+1,-\sigma}^\dagger c_{j,-\sigma}, \\ \mathbf{R} &= \sum_j c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j+1,\uparrow}^\dagger c_{j,\uparrow}, \\ R_2 &= \sum_{j,\sigma} c_{j+1,\sigma}^\dagger c_{j-1,\sigma} (n_{j-1,-\sigma} - 2n_{j,-\sigma} + n_{j+1,-\sigma}), \\ V &= \sum_{j,\sigma} n_{j,\sigma} n_{j+1,-\sigma}, \quad \mathbf{V} = \sum_j n_{j,\downarrow} n_{j,\uparrow} \end{aligned} \quad (10)$$

After returning to the physical time (remember that we did rescaling $t \rightarrow t/\epsilon$), the second-order correction is $\frac{1}{\omega^2}\mathcal{H}_2$ (Eq. 9) (see also [43]).

All the second-order corrections we found enter the effective Hamiltonian with a prefactor $\frac{UJ^2}{\omega^2}$. We therefore require not only $\omega \gg J$, but also $\frac{UJ^2}{\omega^2} \ll 1$ for our theory to be applicable. It is clear that the larger U makes the corrections more important. Moreover, it is possible to find situations where our "corrections" drastically change effective low-energy Hamiltonians. E.g., for $U \gg 1$ and half-filling, the Hubbard model describes a Mott insulator where remaining spin degrees of freedom are coupled by an antiferromagnetic exchange interaction $J_{ex} = \frac{2J^2}{U}$ (this is captured already in a two-site Hubbard model: among the four states (for total $S_z = 0$), the two low-lying states are singlet and triplet states with one electron per site and energies $E_S = -\frac{4}{J}U$, $E_T = 0$). The spectrum at low energies is described by a spin Hamiltonian $2J_{ex}\mathbf{s}_1\mathbf{s}_2$ with $J_{ex} = (E_T - E_S)/2 = \frac{2J^2}{U}$). If parameters in the driven Hubbard model are such that $\frac{J^2}{U} \sim U\frac{J^2}{\omega^2}$, i.e. $U \sim \omega$, then high-frequency corrections are of the same order as the effective exchange interaction itself, and therefore play a crucial role. Very recently, it was proposed to control exchange interactions by time-periodic modulation of an electric field [35]. The corresponding results of [35] can be reproduced and enlightened in our approach. For the two-site model, the correction to the effective Hamiltonian simplifies to

$A(\mathcal{E}_0)\left[(n_{1\downarrow} - n_{2\downarrow})(n_{1\uparrow} - n_{2\uparrow}) - (c_{2\uparrow}^\dagger c_{1\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} c_{1\uparrow}^\dagger c_{2\uparrow})\right] - C(\mathcal{E}_0)(d_2^\dagger d_1 + h.c.)$, where d_k are annihilation operators of pairs (doublons), A and C are functions of driving that in the particular case of harmonic driving are equal to $-\frac{4UJ^2}{\omega^2} \sum_1^\infty \frac{J_m^2(\mathcal{E}_0)}{m^2} + O(\frac{1}{\omega^4})$ and $\frac{4UJ^2}{\omega^2} \sum_1^\infty \frac{(-1)^m J_m^2(\mathcal{E}_0)}{m^2} + O(\frac{1}{\omega^4})$, correspondingly. Parameter A (which gives the strength of exchange interaction) can be calculated taking into account infinite number of terms in $\frac{1}{\omega}$ expansion, if we leave only the leading order contribution in U in every term. Then, $A = -4J^2U \sum_{m=1}^\infty \frac{J_m^2(\mathcal{E}_0)}{m^2\omega^2 - U^2}$. It is interesting that this result, obtained under assumption $\omega > U$, remains valid also for $\omega < U$ [35]. We can obtain even more accurate expression for the effective interaction, which includes influence of other terms in the effective Hamiltonian: eigenvalues of the effective two-site model can be obtained analytically (see SI). The half-distance between the two lowest levels is equal to $\Delta E/2 \equiv J_{ex} = \frac{1}{4} \left(A + C - U + \sqrt{16J_e^2 + (U + 3A - C)^2} \right) \approx \frac{2J_e^2}{U} + A + \frac{2J_e^2}{U^2}(C - 3A) + \dots$ For parameters used in [35], J_{ex} is shown on Fig. 1. It reproduces calculations of [35] amazingly well (see Fig.1 and SI). Moreover, for longer lattices, our theory provides a value of the effective exchange interaction, which is free from (possible) finite-size effects of the two-site model. Indeed, for large U , one can see that

the most important contribution to the effective Hamiltonian comes from \mathcal{S} , other terms increase the number of double occupancies and therefore are suppressed by the (original) interaction term of the Hubbard model. One can eliminate them via a standard approach, by a canonical transformation which renormalizes coefficients of the remaining terms, i.e. exchange interaction ([46]). When $\frac{J^2}{U} \sim U\frac{J^2}{\omega^2}$, with a good accuracy we find the effective exchange interaction is composed of the usual $\frac{J^2}{U}$ part and our correction from \mathcal{S} . So the driving not only modifies the $\frac{J^2}{U}$ part (by renormalizing $J \rightarrow J_e$), but also adds a 'correction' \mathcal{S} which can be of the same order. Suppression of the other terms in the effective Hamiltonian at large U justify usage of the two-site model result in the extended system: we see that at least in the 1D case, the change of the exchange interaction is just the same as in the two-site system. Recently, the fermionic two-site system was realised experimentally [47]. For a long 1D lattice, the ability to switch exchange interaction to the ferromagnetic type imply an interesting possibility of simulating an unusual spin-polaron excitation in driven lattices: a bound state of an extra fermion and a magnon [45]. Such a quasiparticle has peculiar properties, e.g. large effective mass ([45],SI). We can also switch on and off single-particle and doublon hopping by varying parameters of driving (SI).

To conclude, a convenient method based on canonical (unitary) transformations has been applied to two different lattice systems: driven 1D Hubbard and Bose-Hubbard models. For a general high-frequency driving, we derive explicitly effective Hamiltonians including corrections from interactions. For a Hubbard model, a very interesting regime is found, where the corrections drastically influence the effective (Heisenberg) Hamiltonian.

The results should be useful for forthcoming experiments with cold atoms in driven optical lattices. In particular, presently in experiments with shaken optical lattices utilising effective (averaged) tunnelling constants, a frequency of driving is typically chosen in kHz regime, in order to be much higher than all timescales related to trapping potential and interactions, and much lower than interband transition frequency. Explicit knowledge of corrections from trapping potentials [32] and many-body interactions we derived here allow to extend area of applicability of shaken lattice simulations to sub-kHz regime, by choosing driving protocols that cancel second-order corrections. On the other hand, one can amplify particular contributions from second-order corrections, thereby engineering new model Hamiltonians. Another inspiring direction of applications is photoinduced superconductivity and metal-insulator transitions [28–30], where recent studies show importance of induced Kapitza-like effective potentials [29].

Financial support by NWO via Spinoza Prize is acknowledged. A.P.I is grateful to A.I.Neishtadt, M. Eckstein, J.Mentink, A.Polkovnikov, S.Jochim, A.Lichtenshtein, A.Nocera, J.Simonet, M.Thorwart,

T.Rexin for inspiring discussions, and thanks K.Sengstock for hospitality during his stay in Hamburg.

Partial support from RFBR (project no. 13-01-00251) and LEXI (Hamburg) is acknowledged.

-
- [1] N.N. Bogolyubov, The Theory of Perturbations in Non-linear Mechanics, Proc. Inst. Struct. Mech., no. 14, 9 (1950) [In Russian: Teoriya vozmusheniyy v nelineynoy mekhaniki AN USSR, no. 14, 9 (1950)].
- [2] P.L. Kapitza, Soviet Phys. JETP 21, 588 (1951).
- [3] A. Stephenson, Phil. Mag. 15, 233 (1908).
- [4] Stabilization of unstable equilibrium of a pendulum with high-frequency driving was studied firstly in [3] using a linearization procedure, i.e. without introducing effective potentials (being elaborated in Refs. [1, 2]).
- [5] V. I. Arnold, V. V. Kozlov, and A. I. Neishtadt, Mathematical Aspects of Classical and Celestial Mechanics, 3rd ed. (Springer, Berlin, 2006).
- [6] R. Feynman, "Simulating Physics with Computers", International Journal of Theoretical Physics 21, 467 (1982).
- [7] J. I. Cirac, P. Zoller, Nat. Phys. 8, 264 (2012); I. Bloch, J. Dalibard, S. Nascimbene, Nat. Phys. 8, 267 (2012).
- [8] M. Lewenstein et.al., Ultracold Atoms in Optical Lattices: Simulating Quantum Many-Body Systems (Oxford Univ. Press, Oxford, 2012).
- [9] H. Lignier et. al., Phys. Rev. Lett. 99, 220403 (2007).
- [10] A. Eckardt et. al., Phys. Rev. A 79, 013611 (2009).
- [11] J. Heinze et. al., Phys. Rev. Lett. 110, 085302 (2013).
- [12] J. Struck et.al., Phys. Rev. Lett. 108, 225304 (2012).
- [13] S. Greschner et.al., Phys. Rev. Lett. 113, 215303 (2014); N. Goldman, J. Dalibard, Phys. Rev. X 4, 031027 (2014).
- [14] Artificial gauge fields were also simulated using mechanical deformations in graphene: M.A.H. Vozmediano, M.I. Katsnelson, F. Guinea, Phys. Rep. 496 109 (2010); F. Guinea, M. I. Katsnelson, A. K. Geim, Nat. Phys. 6, 30 (2010); Levy et al., Science 329, 544 (2010); P. San-Jose, J. Gonzalez, and F. Guinea, Phys. Rev. Lett. 108 216802 (2012).
- [15] J. Struck et.al., Science 333, 996 (2011).
- [16] N. Strohmaier et.al., Phys. Rev. Lett. 99, 220601 (2007); A. Alberti et al., Nat. Phys. 5, 547 (2009); A. Zenesini et al., Phys. Rev. Lett. 102, 100403 (2009); A. Eckardt et al., Europhys. Lett. 89, 10010 (2010).
- [17] K. Kudo, T. Boness, and T. S. Monteiro, Phys. Rev. A 80, 063409 (2009); K. Kudo and T. S. Monteiro, ibid. 83, 053627 (2011); A. R. Kolovsky, E. A. Gomez, and H. J. Korsch, ibid. 81, 025603 (2010).
- [18] A. Eckardt, C. Weiss, and M. Holthaus, Phys. Rev. Lett. 95, 260404 (2005).
- [19] A. Zenesini et.al., Phys. Rev. Lett. 102, 100403 (2009).
- [20] W. Zwerger, J. Opt. B: Quantum Semiclass. Opt. 5, S9 (2003).
- [21] I. Bloch, J. Dalibard, W. Zwerger, Rev. Mod. Phys. 80, 885 (2008).
- [22] D.H. Dunlap and V.M. Kenkre, Phys. Rev. B 34, 3625 (1986).
- [23] M. Holthaus, Phys. Rev. Lett. 69, 351 (1992).
- [24] M. Holthaus, D.W. Hohe, Philos. Mag. B 74, 105 (1996).
- [25] M. Glück, A.R. Kolovsky, and H.J. Korsch, Phys. Rep. 366, 103 (2002).
- [26] N.H. Lindner, G. Refael, and V. Galitski, Nat. Phys. 7, 490 (2011).
- [27] T. Kitagawa et.al., Phys. Rev. B 84, 235108 (2011); M.S. Rudner, et.al., Phys. Rev. X 3, 031005 (2013).
- [28] M. Rini et.al., Nature 449, 72 (2007).
- [29] A. Subedi, A. Cavalleri, and A. Georges arXiv:1311.0544v1 (2013).
- [30] D. Fausti et. al., Science 331, 189 (2011); S. Kaiser, et al., arXiv e-print 1205.4661 (2013).
- [31] S. Longhi, Phys. Rev. B 77, 195326 (2008).
- [32] A.P. Itin, A.I. Neishtadt, Phys. Lett. A 378, 822 (2014).
- [33] T. Grozdanov and M. Rakovic, Phys. Rev. A 38, 1739 (1988).
- [34] S. Rahav, I. Gilary, and S. Fishman, Phys. Rev. A 68, 013820 (2003).
- [35] J. H. Mentink, K. Balzer, and M. Eckstein, arXiv:1407.4761 (2014).
- [36] S. Blanes, F. Casas, J.A. Oteo, J. Ros, Phys. Rep. 470, 151 (2009).
- [37] W. Magnus, Comm. Pure and Appl. Mathematics (1954).
- [38] A. P. Itin, R. de la Llave, A. I. Neishtadt, and A. A. Vasiliev, Chaos 12, 1043 (2002); X. Leoncini, A. I. Neishtadt, and A. A. Vasiliev, Phys. Rev. E 79, 026213 (2009).
- [39] A. P. Itin and A. I. Neishtadt, Phys. Rev. E 86, 016206 (2012).
- [40] A.P. Itin et al., in preparation.
- [41] J. Heinze et.al., Phys. Rev. Lett. 107, 135303 (2011).
- [42] A. Verdeny et.al., Phys. Rev. Lett. 111, 175301 (2013).
- [43] Note that a two-particle problem in a driven *asymmetric* Hubbard model was recently considered in [44], where formation of a doublon of two fermions (an *extended pair*) residing on neighbouring sites and co-tunnelling along the lattice was found (bare tunnelling constants of the two fermions being different, $J_{\uparrow} \neq J_{\downarrow}$). The first-order correction found in [44] is proportional to $J_{\uparrow} - J_{\downarrow}$, i.e. it completely disappears in the symmetric limit $J_{\uparrow} = J_{\downarrow}$, in accordance with our findings.
- [44] S. Longhi and G.D. Valle, Phys. Rev. A 87, 013634 (2013).
- [45] M.I. Katsnelson, Fizika Metallov i Metallovedenie 54, 396 (1981) [In Russian].
- [46] A.H. MacDonald, S.M. Girvin, D. Yoshioka, Phys. Rev. B 37, 9753 (1988).
- [47] S. Murmann et. al., Phys. Rev. Lett. 114, 080402 (2015).

Supplementary Information

A. Comparison with Floquet-Magnus expansion

Consider the linear differential equation

$$Y'(t) = A(t)Y(t), \quad (11)$$

where $A(t)$ is, for example, a $n \times n$ matrix and $Y(t)$ is a n -dimensional vector function (more general objects can also be treated [2], e.g. $A(t)$ and $Y(t)$ being operators in Hilbert space). Initial condition $Y(0) = Y_0 = I$ is imposed. Magnus' proposal is to search for a solution of this initial value problem in the form of

$$Y(t) = \exp(\Omega(t)) Y_0, \quad (12)$$

where $\Omega(t)$ is $n \times n$ matrix.

In the context of quantum physics, this approach is in contrast with the representation $Y(t) = \mathcal{T} \left(\exp \int_0^t A(s) ds \right)$, where \mathcal{T} is the Dyson time-ordering operator.

Magnus found that

$$\frac{d\Omega(t)}{dt} = \sum_0^{\infty} \frac{B_n}{n!} ad_{\Omega}^n A, \quad (13)$$

where $ad_{\Omega}^n A$ is a shorthand for a nested commutator ($ad_{\Omega} A = [\Omega, A]$, $ad_{\Omega}^k A = [\Omega, ad_{\Omega}^{k-1} A]$), and B_n are the Bernoulli numbers.

Integration of (13) by means of iteration leads to a series for Ω . In the original formulation, this approach does not immediately serve for finding time-independent Hamiltonians. However, one can modify it in the following way [2]. In the case of periodically time-dependent matrix $A(t)$, according to Floquet and Lyapunov, a solution can be factorized on periodic part and a purely exponential factor:

$$Y(t) = P(t) \exp(tF), \quad (14)$$

where F and $P(t)$ are $n \times n$ matrices, the latter is periodic $P(t+T) = P(t)$, and F is constant. Now, one can interpretate Eq.14 as following: after transformation of the solution by $P^{-1}(t)$, one get in the new representation an equation of motion determined by constant matrix F instead of $A(t)$. Therefore, $P(t)$ is the analog of our $C = \exp(\epsilon K_1(t) + \dots)$ from the main text, while F is the analog of our time-independent Hamiltonian H_{eff} .

Naively, F and H_{eff} should merely coincide with each other. However, there is a subtle but important difference stemming from different initial conditions.

Let us remind the derivation of Floquet-Magnus expansion [2]. Introducing the factorized form 14 into the differential equation $Y' = A(t)Y$, one gets equation of motion for $P(t)$.

$$P'(t) = A(t)P(t) - P(t)F, \quad P(0) = I. \quad (15)$$

For $P(t)$ one uses the exponential ansatz

$$P(t) = \exp(\Lambda(t)), \quad \Lambda(0) = 0, \quad (16)$$

and obtains

$$\Lambda' = \sum_{k=0}^{\infty} \frac{B_k}{k!} ad_{\Lambda}^k (A + (-1)^{k+1} F) \quad (17)$$

Then, one considers series expansion for F and Λ

$$\Lambda(t) = \sum_{k=1}^{\infty} \Lambda_k(t), \quad F = \sum_{k=1}^{\infty} F_k, \quad (18)$$

with all $\Lambda_k(0) = 0$. Λ_k is supposed to be of order ϵ^k . Alternatively, one can replace A by ϵA , obtaining powers of ϵ explicitly in the expansion $\Lambda(t) = \sum_{k=1}^{\infty} \epsilon^k \Lambda_k(t)$.

Evaluating terms with the same order, one gets

$$\begin{aligned}\Lambda_1(t) &= \int_0^t A(x) dx - tF_1, \\ F_1 &= \frac{1}{T} \int_0^T A(x) dx, \\ \Lambda_2(t) &= \frac{1}{2} \int_0^t [A(x) + F_1, \Lambda_1] dx - tF_2, \\ F_2 &= \frac{1}{2T} \int_0^T [A(x) + F_1, \Lambda_1(x)] dx\end{aligned}\tag{19}$$

One can see that while F_1 coincide with our H_0 , F_2 is different from H_1 . The reason is that the mean value of Λ_k is not zero. Instead, a condition $\Lambda_k = 0$ is imposed. In fact, this condition is unnecessary for our purposes (for derivation of effective Hamiltonians). Let us return back to the original Magnus expansion. In the equation (11), the initial condition $Y(0) = I$ was imposed. General solution of (11) with other initial conditions can be obtained by multiplying $Y(t)$ by a constant matrix. The chosen initial condition transforms to $P(0) = I$ when considering the periodic system, and then to $\Lambda(0) = 0$. However, generally we can admit transformations where $P(0) \neq I$, since we actually interested not in constructing the fundamental matrix for the solution of the initial value problem, but in constructing effective Hamiltonians. So, its not a problem if $\bar{X}(0) \neq X(0)$ in the main text. Our main priority is that transformed \bar{X} evolves according to time-independent effective Hamiltonian which captures averaged behaviour of the system on long times as good as possible. This freedom ($\bar{X}(0) \neq X(0)$) allows for tremendous improvement in the construction of the effective Hamiltonian: namely, cancellation of the first-order corrections. The remaining second-order corrections cannot be removed by further linear transformations and therefore essential for understanding averaged properties of the system. One necessarily need to truncate expansion of the time-independent matrix F at some order. Clearly, it is insufficient to truncate it on the first order corrections (F_2), since this part can be removed as described above. We see that while our expansion described in the main text is in accord with Magnus expansion, it is slightly different due to initial conditions, and is actually more effective for revealing long-time features of the system: it completely removes the lowest order correction, so that only higher-order corrections remains.

B. A single-particle tight-binding model

Here and in the next two sections we consider particular models and calculate commutators for the older version of Eqs. (3) of the main text, that is for expressions available in [3]:

$$\begin{aligned}\mathcal{H}_0 &= \langle \mathcal{H} \rangle, \quad \mathcal{H}_1 = \frac{1}{2} \langle [\{\mathcal{H}\}, K_1] \rangle, \\ \mathcal{H}_2 &= \langle [\mathcal{H}, K_2] + \frac{1}{2} [[\mathcal{H}, K_1], K_1] - \frac{i}{2} ([\dot{K}_1, K_2] \\ &\quad + [\dot{K}_2, K_1]) - \frac{i}{6} [[\dot{K}_1, K_1], K_1] \rangle,\end{aligned}\tag{20}$$

Despite being less compact, they allow somehow deeper insight in the structure of various terms of the expansion.

In the case of a particle in a fastly driven tight-binding chain, one has (see [3]) $H = J \sum (|n\rangle \langle n+1| + |n+1\rangle \langle n|) - ed\omega E(\omega t) \sum_n n |n\rangle \langle n|$, where d is the intersite distance, ωE is the applied electric field (here, we explicitly place $\omega \equiv 1/\epsilon$ in the definition to emphasize strong driving), e is the charge of the particle, J is the tunnelling constant. The same model can be realized also with neutral particles (in the co-moving frame), by appropriate shaking of the lattice [4]. Expanding a quantum state as $|\psi(t)\rangle = \sum c_n |n\rangle$, we get a system of equations

$$i\dot{c}_n = J(c_{n+1} + c_{n-1}) - \omega \mathcal{E}(\omega t) n c_n.\tag{21}$$

We make a transformation $c_n(t) = X_n(t) \exp\left[-in \int_0^t \omega \mathcal{E}(\omega t') dt'\right]$, so that equations of motion are

$$i\dot{X}_n = J(X_{n+1}F(\omega t) + X_{n-1}F^*(\omega t)), \quad (22)$$

where $F(\omega t) = \exp[-i \int_0^t \omega \mathcal{E}(\omega t') dt'] = F_0 + \sum F_l \exp(-il\omega t)$, $\mathcal{E} = edE$.

Introducing fast time $t' = \omega t \equiv t/\epsilon$, we get, in the matrix form, Eq.(1) of the main text, with the Hamiltonian \mathcal{H} corresponding to Eq.(22).

In a more general setting, a particle in a driven tight-binding chain subject to an additional external potential $V(n)$, we have

$$i\dot{X}_n = \epsilon[J(X_{n+1}F(t) + X_{n-1}F^*(t)) + V(n)X_n], \quad (23)$$

or

$$i\dot{X} = \epsilon H, \quad (24)$$

In the spirit of classical canonical perturbation theory, we are making a unitary transformation $X = C\tilde{X}$ so that equations for the transformed variables are

$$i\dot{\tilde{X}} = [C^{-1}\epsilon HC - iC^{-1}\dot{C}]\tilde{X}. \quad (25)$$

We are searching for a transformation of the form $C = \exp[\epsilon K_1 + \epsilon^2 K_2 + \epsilon^3 K_3]$, where K_i are skew-Hermitian time-periodic matrices, which would remove time-dependent terms from the Hamiltonian, leaving only time-independent terms.

We have

$$\begin{aligned} C &\approx I + \epsilon K_1 + \epsilon^2 \left(\frac{1}{2}K_1^2 + K_2\right) + \epsilon^3 \left(\frac{1}{6}K_1^3 + \frac{1}{2}(K_1K_2 + K_2K_1) + K_3\right), \\ C^\dagger &\approx I - \epsilon K_1 + \epsilon^2 \left(\frac{1}{2}K_1^2 - K_2\right) + \epsilon^3 \left(-\frac{1}{6}K_1^3 + \frac{1}{2}(K_1K_2 + K_2K_1) - K_3\right), \end{aligned} \quad (26)$$

where I is the unity matrix.

In the first order, we have

$$i\dot{K}_1 = H(t) - \langle H(t) \rangle \equiv \{H\}, \quad (27)$$

and therefore $iK_1 = \int (H - \langle H \rangle) dt = \int \{H\} dt$. We introduce here curly brackets as taking time-periodic part of a time-dependent function: $\{X\} \equiv X - \langle X(t) \rangle$, where $\langle X(t) \rangle \equiv \frac{1}{2\pi} \int_0^{2\pi} X(t') dt'$.

In the second order,

$$i\dot{K}_2 = \left\{ HK_1 - K_1H - \frac{i}{2}(\dot{K}_1K_1 - K_1\dot{K}_1) \right\}, \quad (28)$$

and the effective Hamiltonian is

$$\epsilon H_{eff} = \epsilon \mathcal{H}_0 + \epsilon^2 \mathcal{H}_1 + \epsilon^3 \mathcal{H}_2, \quad (29)$$

where

$$\begin{aligned} \mathcal{H}_0 &= \langle H \rangle \\ \mathcal{H}_1 &= \langle HK_1 - K_1H - \frac{i}{2}(\dot{K}_1K_1 - K_1\dot{K}_1) \rangle \\ \mathcal{H}_2 &= \langle HK_2 - K_2H + \frac{1}{2}(HK_1^2 + K_1^2H) - K_1HK_1 - \frac{i}{2}(\dot{K}_1K_2 - K_1\dot{K}_2 + \dot{K}_2K_1 - K_2\dot{K}_1) \\ &\quad - \frac{i}{6}(\dot{K}_1K_1^2 + K_1^2\dot{K}_1 - 2K_1\dot{K}_1K_1) \rangle \end{aligned} \quad (30)$$

In the case of the uniform tight-binding model (without external potential) and open boundary conditions the effective Hamiltonians have the following simple form

$$\mathcal{H}_1 = J^2 D \mathcal{Z}_1, \quad \mathcal{H}_2 = -\frac{J^3}{3} (L_3 \mathcal{U}_1 + L_3^* \mathcal{B}_1), \quad (31)$$

where $\mathcal{Z}_1 = \delta_{i,j} \delta_{i,1} - \delta_{i,j} \delta_{i,N}$, N is the number of sites, $D = \sum_{l=1}^{\infty} (|F_l|^2 - |F_{-l}|^2)/l$, $\mathcal{U}_1 = \delta_{i,i+1} \delta_{i,1} + \delta_{i,i+1} \delta_{i,N-1}$ (the upper co-diagonal with '1' on its ends, and zeros elsewhere), $\mathcal{B}_1 = \delta_{i,i-1} \delta_{i,2} + \delta_{i,i-1} \delta_{i,N}$ (the lower co-diagonal with '1' on its ends), L_3 is a function of driving defined in [3]. In other words, the first-order correction \mathcal{H}_1 is non-zero only if perturbation has certain broken time symmetry (for a single harmonic perturbation, $\cos t$, H_1 disappears), and it localizes near the ends of the chain (near boundaries). More generally, it requires non-uniformity of the coupling constant J , so e.g. in a model with alternating coupling constants $J_1 - J_2 - J_1$, or with a more general non-uniform coupling $J(n)$, the correction will be non-zero throughout the whole chain [5]. For the uniform coupling $J(n)$, \mathcal{H}_2 is also localized near the boundaries of the open chain. The method of multiple time scales [5] produces the same results in these cases.

A very interesting new result can be obtained for a parabolic external potential $V(n) = Vn^2/2$: the second-order correction to the averaged Hamiltonian corresponds to an induced uniform next-nearest neighbour coupling, which for the harmonic perturbation with amplitude K is explicitly given by $J' = -\epsilon^2 J^2 V \sum_{l=1}^{\infty} \frac{(-1)^l J_l^2(K)}{l^2}$ [3].

In the case of infinite uniform lattice, or periodic boundary conditions, \mathcal{H}_1 and higher-order corrections are absent.

C. Bose-Hubbard model

We have

$$\dot{K}_1 = -i\{H\} = -i \sum_i \left(\delta^+ c_i^\dagger c_{i+1} + \delta^- c_{i+1}^\dagger c_i \right), \quad (32)$$

$$K_1 = -i \sum_i \left(\delta_1^+ c_i^\dagger c_{i+1} + \delta_1^- c_{i+1}^\dagger c_i \right) \quad (33)$$

$$HK_1 - K_1H = -i2U \sum_j \left(\delta_1^+ c_j^\dagger (n_j - n_{j+1}) c_{j+1} + \delta_1^- c_{j+1}^\dagger (n_{j+1} - n_j) c_j \right) \quad (34)$$

Since $[\dot{K}_1, K_1] = 0$ if we impose periodic boundary conditions,

$$\begin{aligned} i\dot{K}_2 &= \{HK_1 - K_1H\}, \\ K_2 &= -2U \sum_j \left(\delta_2^+ c_j^\dagger (n_j - n_{j+1}) c_{j+1} + \delta_2^- c_{j+1}^\dagger (n_{j+1} - n_j) c_j \right) \end{aligned} \quad (35)$$

For the second-order corrections we have

$$\begin{aligned}
HK_2 - K_2H &= 2U \sum_{\alpha=1}^6 A_\alpha, \\
A_1 &= -\delta_2^+ \delta_0^+ \sum_j \left(c_{j-1}^\dagger (4n_j - n_{j+1} - n_{j-1}) c_{j+1} - 2c_j^\dagger c_j^\dagger c_{j+1} c_{j+1} \right) = -\delta_2^+ \delta_0^+ a_1 \\
A_2 &= -\delta_2^+ \delta_0^- \sum_j \left(4n_j n_{j+1} - 2n_j (n_j - 1) - c_{j-1}^\dagger c_{j+1}^\dagger c_j c_j - c_j^\dagger c_j^\dagger c_{j+1} c_{j-1} \right) = -\delta_2^+ \delta_0^- a_2, \\
A_3 &= \sum_j \left(4n_j n_{j+1} - 2n_j (n_j - 1) - c_{j-1}^\dagger c_{j+1}^\dagger c_j c_j - c_j^\dagger c_j^\dagger c_{j+1} c_{j-1} \right) = -\delta_2^- \delta_0^+ a_3, \\
A_4 &= -\delta_2^- \delta_0^- \sum_j \left(c_{j+1}^\dagger (4n_j - n_{j+1} - n_{j-1}) c_j - 2c_{j+1}^\dagger c_{j+1}^\dagger c_j c_j \right) = -\delta_2^- \delta_0^- a_4, \\
A_5 &= -2U \delta_2^+ \sum_j \left(c_j^\dagger (n_j - n_{j+1})^2 c_{j+1} \right) = -2U \delta_2^+ a_5, \\
A_6 &= -2U \delta_2^- \sum_j \left(c_{j+1}^\dagger (n_j - n_{j+1})^2 c_j \right) = -2U \delta_2^- a_6,
\end{aligned} \tag{36}$$

$$\begin{aligned}
-\frac{i}{2} \left(\dot{K}_1 K_2 - K_2 \dot{K}_1 \right) &= U \sum_{\alpha=1}^4 B_\alpha, \\
B_1 &= \delta^+ \delta_2^+ a_1, \\
B_2 &= \delta^+ \delta_2^- a_2, \\
B_3 &= \delta^- \delta_2^+ a_3, \\
B_4 &= \delta^- \delta_2^- a_4,
\end{aligned} \tag{37}$$

$$\begin{aligned}
-\frac{i}{2} \left(\dot{K}_2 K_1 - K_1 \dot{K}_2 \right) &= U \sum_{\alpha=1}^4 C_\alpha, \\
C_1 &= -(\delta_1^+)^2 a_1, \\
C_2 &= -\delta_1^+ \delta_1^- a_2, \\
C_3 &= -\delta_1^- \delta_1^+ a_3, \\
C_4 &= -(\delta_1^-)^2 a_4,
\end{aligned} \tag{38}$$

$$\begin{aligned}
\frac{1}{2} [[H, K_1], K_1] &= U \sum_{\alpha=1}^4 D_\alpha, \\
D_1 &= (\delta_1^+)^2 a_1, \\
D_2 &= \delta_1^+ \delta_1^- a_2, \\
D_3 &= \delta_1^- \delta_1^+ a_3, \\
D_4 &= (\delta_1^-)^2 a_4,
\end{aligned} \tag{39}$$

We note that $D_k = -C_k$ and time-averages of the coefficients A_5 and A_6 are equal to zero. Also, $[[\dot{K}_1, K_1], K_1] = 0$. We therefore get the effective second-order correction

$$\begin{aligned}
\mathcal{H}_2 &= U \sum_{\alpha=1}^4 \Delta_\alpha a_\alpha, \\
\Delta_1 &= \langle \delta_2^+ \delta^+ - 2\delta_2^+ \delta_0^+ \rangle, \\
\Delta_2 &= \langle \delta^+ \delta_2^- - 2\delta_2^+ \delta_0^- \rangle, \\
\Delta_3 &= \langle \delta^- \delta_2^+ - 2\delta_2^- \delta_0^+ \rangle, \\
\Delta_4 &= \langle \delta^- \delta_2^- - 2\delta_2^- \delta_0^- \rangle,
\end{aligned} \tag{40}$$

where a_α are given by

$$\begin{aligned}
a_1 &= \sum_j \left(c_{j-1}^\dagger (4n_j - n_{j+1} - n_{j-1}) c_{j+1} - 2c_j^\dagger c_{j+1}^\dagger c_{j+1} \right) \\
a_2 &= \sum_j \left(4n_j n_{j+1} - 2n_j (n_j - 1) - c_{j-1}^\dagger c_{j+1}^\dagger c_j c_j - c_j^\dagger c_{j+1}^\dagger c_{j+1} c_{j-1} \right) \\
a_3 &= \sum_j \left(4n_j n_{j+1} - 2n_j (n_j - 1) - c_{j-1}^\dagger c_{j+1}^\dagger c_j c_j - c_j^\dagger c_{j+1}^\dagger c_{j+1} c_{j-1} \right) \\
a_4 &= \sum_j \left(c_{j+1}^\dagger (4n_j - n_{j+1} - n_{j-1}) c_{j-1} - 2c_{j+1}^\dagger c_{j+1}^\dagger c_j c_j \right)
\end{aligned} \tag{41}$$

We can simplify it further:

$$\mathcal{H}_2 = U(\Delta_1 a_1 + h.c) + U(\Delta_2 + \Delta_2^*) a_2 \tag{42}$$

Its not difficult to see that $(\Delta_2 + \Delta_2^*) = -2\Delta_0$ as given in the main text, while $\Delta_1 = -2\Delta^+$.

D. Hubbard model

We have:

$$\begin{aligned}
H &= \sum_{i,\sigma} \left(\delta_0^+ c_{i,\sigma}^\dagger c_{j+1,\sigma} + \delta_0^- c_{i+1,\sigma}^\dagger c_{i,\sigma} \right) + U \sum_i n_{i,\sigma} n_{i,-\sigma} \\
&= T^\uparrow + T^\downarrow + \mathcal{U} \\
\dot{K}_1 &= -i\{H\} = -i \sum_{j,\sigma} \left(\delta^+ c_{j,\sigma}^\dagger c_{j+1,\sigma} + \delta^- c_{j+1,\sigma}^\dagger c_{j,\sigma} \right), \\
K_1 &= -i \sum_{j,\sigma} \left(\delta_1^+ c_{j,\sigma}^\dagger c_{j+1,\sigma} + \delta_1^- c_{j+1,\sigma}^\dagger c_{j,\sigma} \right) \equiv K_1^\uparrow + K_1^\downarrow,
\end{aligned}$$

$$\begin{aligned}
HK_1 - K_1H &= [T^\uparrow, K_1^\uparrow] + [T^\downarrow, K_1^\downarrow] + [\mathcal{U}, K_1^\uparrow] + [\mathcal{U}, K_1^\downarrow], \\
T^\uparrow K_1^\uparrow - K_1^\uparrow T^\uparrow &= -i(\delta_0^- \delta_1^+ - \delta_0^+ \delta_1^-) \sum_i (n_{i+1,\uparrow} - n_{i,\uparrow}) \\
T^\downarrow K_1^\downarrow - K_1^\downarrow T^\downarrow &= -i(\delta_0^- \delta_1^+ - \delta_0^+ \delta_1^-) \sum_i (n_{i+1,\downarrow} - n_{i,\downarrow}) \\
\mathcal{U} K_1^\uparrow - K_1^\uparrow \mathcal{U} &= -iU \sum_j (\delta_1^+ c_{j,\uparrow}^\dagger c_{j+1,\uparrow} (n_{j,\downarrow} - n_{j+1,\downarrow}) - \delta_1^- c_{j+1,\uparrow}^\dagger c_{j,\uparrow} (n_{j,\downarrow} - n_{j+1,\downarrow})) \\
\mathcal{U} K_1^\downarrow - K_1^\downarrow \mathcal{U} &= -iU \sum_j (\delta_1^+ c_{j,\downarrow}^\dagger c_{j+1,\downarrow} (n_{j,\uparrow} - n_{j+1,\uparrow}) - \delta_1^- c_{j+1,\downarrow}^\dagger c_{j,\downarrow} (n_{j,\uparrow} - n_{j+1,\uparrow}))
\end{aligned} \tag{43}$$

We have also $[\dot{K}_1, K_1] = 0$, therefore

$$\begin{aligned} i\dot{K}_2 &= \{HK_1 - K_1H\}, \\ \dot{K}_2 &= \delta_{01} \sum_{j,\sigma} (n_{j,\sigma} - n_{j+1,\sigma}) + U \sum_{j,\sigma} (\delta_1^- c_{j+1,\sigma}^\dagger c_{j,\sigma} - \delta_1^+ c_{j,\sigma}^\dagger c_{j+1,\sigma}) (n_{j,-\sigma} - n_{j+1,-\sigma}) = K_\delta^\uparrow + K_\delta^\downarrow + K_u^\uparrow + K_u^\downarrow \end{aligned} \quad (44)$$

To calculate commutator of H and K_2 , we proceed as follows:

$$\begin{aligned} [H, K_2] &= [T^\uparrow, K_u^\uparrow] + [T^\uparrow, K_u^\downarrow] + [T^\downarrow, K_u^\uparrow] + [T^\downarrow, K_u^\downarrow] + [\mathcal{U}, K_u^\uparrow] + [\mathcal{U}, K_u^\downarrow] \\ T^\uparrow K_u^\uparrow - K_u^\uparrow T^\uparrow &= U \sum_j (\delta_0^+ \delta_2^- + \delta_0^- \delta_2^+) (2n_{j,\uparrow} n_{j,\downarrow} - n_{j,\uparrow} n_{j+1,\downarrow} - n_{j,\downarrow} n_{j+1,\uparrow}) \\ &\quad + U \sum_j (\delta_0^- \delta_2^- c_{j+1,\uparrow}^\dagger c_{j-1,\uparrow} + \delta_0^+ \delta_2^+ c_{j-1,\uparrow}^\dagger c_{j+1,\uparrow}) (n_{j-1,\downarrow} - 2n_{j,\downarrow} + n_{j+1,\downarrow}) \\ T^\downarrow K_u^\downarrow - K_u^\downarrow T^\downarrow &= \mathcal{P}(\uparrow \rightarrow \downarrow, \downarrow \rightarrow \uparrow) (T^\uparrow K_u^\uparrow - K_u^\uparrow T^\uparrow) \\ T^\uparrow K_u^\downarrow - K_u^\downarrow T^\uparrow &= U \sum_j \delta_0^+ \delta_2^- (-2c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j,\uparrow}^\dagger c_{j+1,\uparrow} + c_{j,\downarrow}^\dagger c_{j-1,\downarrow} c_{j,\uparrow}^\dagger c_{j+1,\uparrow} + c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j-1,\uparrow}^\dagger c_{j,\uparrow}) \\ &\quad - U \sum_j \delta_0^+ \delta_2^+ (-2c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j,\uparrow}^\dagger c_{j+1,\uparrow} + c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j-1,\uparrow}^\dagger c_{j,\uparrow} + c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j+1,\uparrow}^\dagger c_{j+2,\uparrow}) \\ &\quad + U \sum_j \delta_0^- \delta_2^- (2c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j+1,\uparrow}^\dagger c_{j,\uparrow} - c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j+2,\uparrow}^\dagger c_{j+1,\uparrow} - c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j,\uparrow}^\dagger c_{j-1,\uparrow}) \\ &\quad - U \sum_j \delta_0^- \delta_2^+ (2c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j+1,\uparrow}^\dagger c_{j,\uparrow} - c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j+2,\uparrow}^\dagger c_{j+1,\uparrow} - c_{j,\downarrow}^\dagger c_{j+1,\downarrow} c_{j,\uparrow}^\dagger c_{j-1,\uparrow}) \\ T^\downarrow K_u^\uparrow - K_u^\uparrow T^\downarrow &= U \sum_j \delta_0^+ \delta_2^- (-2c_{j+1,\uparrow}^\dagger c_{j,\uparrow} c_{j,\downarrow}^\dagger c_{j+1,\downarrow} + c_{j+1,\uparrow}^\dagger c_{j,\uparrow} c_{j-1,\downarrow}^\dagger c_{j,\downarrow} + c_{j,\uparrow}^\dagger c_{j-1,\uparrow} c_{j,\downarrow}^\dagger c_{j+1,\downarrow}) \\ &\quad - U \sum_j \delta_0^+ \delta_2^+ (-2c_{j,\uparrow}^\dagger c_{j+1,\uparrow} c_{j,\downarrow}^\dagger c_{j+1,\downarrow} + c_{j,\uparrow}^\dagger c_{j+1,\uparrow} c_{j-1,\downarrow}^\dagger c_{j,\downarrow} + c_{j-1,\uparrow}^\dagger c_{j,\uparrow} c_{j,\downarrow}^\dagger c_{j+1,\downarrow}) \\ &\quad + U \sum_j \delta_0^- \delta_2^- (2c_{j+1,\uparrow}^\dagger c_{j,\uparrow} c_{j+1,\downarrow}^\dagger c_{j,\downarrow} - c_{j,\uparrow}^\dagger c_{j-1,\uparrow} c_{j+1,\downarrow}^\dagger c_{j,\downarrow} - c_{j+1,\uparrow}^\dagger c_{j,\uparrow} c_{j,\downarrow}^\dagger c_{j-1,\downarrow}) \\ &\quad - U \sum_j \delta_0^- \delta_2^+ (2c_{j,\uparrow}^\dagger c_{j+1,\uparrow} c_{j+1,\downarrow}^\dagger c_{j,\downarrow} - c_{j-1,\uparrow}^\dagger c_{j,\uparrow} c_{j+1,\downarrow}^\dagger c_{j,\downarrow} - c_{j,\uparrow}^\dagger c_{j+1,\uparrow} c_{j,\downarrow}^\dagger c_{j-1,\downarrow}) \\ &= \mathcal{P}(\uparrow \rightarrow \downarrow, \downarrow \rightarrow \uparrow) (T^\uparrow K_u^\downarrow - K_u^\downarrow T^\uparrow) \\ \mathcal{U} K_u^\uparrow - K_u^\uparrow \mathcal{U} &= -U^2 \sum_j (\delta_2^- c_{j+1,\uparrow}^\dagger c_{j,\uparrow} - \delta_2^+ c_{j,\uparrow}^\dagger c_{j+1,\uparrow}) (n_{j,\downarrow} - n_{j+1,\downarrow})^2 \\ \mathcal{U} K_u^\downarrow - K_u^\downarrow \mathcal{U} &= -U^2 \sum_j (\delta_2^- c_{j+1,\downarrow}^\dagger c_{j,\downarrow} - \delta_2^+ c_{j,\downarrow}^\dagger c_{j+1,\downarrow}) (n_{j,\uparrow} - n_{j+1,\uparrow})^2 \end{aligned} \quad (46)$$

The last two terms in Eq.(45) will produce zero time-average and are not important for us. Let us introduce $[H, K_2]_{\mathcal{U}} \equiv [T^\uparrow, K_u^\uparrow] + [T^\uparrow, K_u^\downarrow] + [T^\downarrow, K_u^\uparrow] + [T^\downarrow, K_u^\downarrow]$, i.e. the expression (45) without the last two terms. Expressions above represent several fundamental processes. We introduce the following notation for them:

$$\begin{aligned}
\mathcal{S} &= \mathcal{S}_{\uparrow,\downarrow} + \mathcal{S}_{\downarrow,\uparrow}, & \mathcal{S}_{\sigma,-\sigma} &= \sum_j c_{j+1,\sigma}^\dagger c_{j,\sigma} c_{j,-\sigma}^\dagger c_{j+1,-\sigma}, \\
\mathcal{A} &= \mathcal{A}_{\uparrow,\downarrow} + \mathcal{A}_{\downarrow,\uparrow}, & \mathcal{A}_{\sigma,-\sigma} &= \sum_j c_{j,\sigma}^\dagger c_{j-1,\sigma} c_{j,-\sigma}^\dagger c_{j+1,-\sigma}, \\
\mathcal{R} &= \mathcal{R}_{\uparrow\downarrow} + \mathcal{R}_{\downarrow\uparrow}, & \mathcal{R}_{\sigma,-\sigma} &= \sum_j c_{j,\sigma}^\dagger c_{j-1,\sigma} c_{j+1,-\sigma}^\dagger c_{j,-\sigma}, \\
\mathbf{R} &= \sum_j c_{j+1,\downarrow}^\dagger c_{j,\downarrow} c_{j+1,\uparrow}^\dagger c_{j,\uparrow}, \\
R_2 &= R_{2,\uparrow} + R_{2,\downarrow}, & R_{2,\sigma} &= \sum_j c_{j+1,\sigma}^\dagger c_{j-1,\sigma} (n_{j-1,-\sigma} - 2n_{j,-\sigma} + n_{j+1,-\sigma}), \\
V &= \sum_\sigma V_\sigma, & V_\sigma &= \sum_j n_{j,\sigma} n_{j+1,-\sigma}, \\
\mathbf{V} &= \sum_j n_{j,\downarrow} n_{j,\uparrow}
\end{aligned} \tag{47}$$

There are several different types of tunnelling and interaction processes here (\mathcal{S} , \mathcal{A} , \mathcal{R} , \mathbf{R} , R_2 , V , and \mathbf{V}). \mathcal{S} denotes a correlated tunneling process of two atoms on neighboring sites exchanging their positions ('superexchange'). \mathcal{A}^\dagger denotes a tunnelling process where a pair of atoms at j th site is dissociated into two atoms at neighboring sites ($j-1$ and $j+1$). \mathcal{A} denotes a corresponding correlated tunneling process of association of two atoms into a pair. \mathcal{R} denotes a correlated tunnelling process where an extended pair of atoms is tunnelling to the right neighboring sites, and \mathcal{R}^\dagger denotes tunnelling to the left. \mathbf{R} denotes tunnelling of a localized pair of atoms to the right, and \mathbf{R}^\dagger denotes tunnelling to the left. R_2 denotes single-particle next-nearest-neighbour density-dependent tunnelling to the right, and R_2^\dagger denotes corresponding tunnelling to the left. V denotes nearest-neighbour interaction, and \mathbf{V} denotes usual local interaction. We can rewrite $[H, K_2]_{\mathcal{U}}$ in terms of these fundamental processes:

$$[H, K_2]_{\mathcal{U}} = U[(\delta_0^- \delta_2^- (4\mathbf{R} - 2\mathcal{R} + R_2) + H.c.) + (\delta_0^+ \delta_2^- + \delta_0^- \delta_2^+) (-2\mathcal{S} + \mathcal{A} + \mathcal{A}^\dagger + 4\mathbf{V} - 2V)] \tag{49}$$

Now, let us consider the term $\frac{1}{2}[[H, K_1], K_1]$. It can be shown $\frac{1}{2}[[H, K_1], K_1] = \frac{1}{2}\mathcal{P}(\delta_0^\pm \rightarrow \delta_1^\pm, \delta_2^\pm \rightarrow -\delta_1^\pm)[H, K_2]_{\mathcal{U}}$. Also,

$$-\frac{i}{2}[\dot{K}_2, K_1] = \frac{1}{2}\mathcal{P}(\delta_0^\pm \rightarrow \delta_1^\pm, \delta_2^\pm \rightarrow \delta_1^\pm)[H, K_2]_{\mathcal{U}} = -\frac{1}{2}[[H, K_1], K_1], \tag{50}$$

and

$$-\frac{i}{2}[\dot{K}_1, K_2] = -\frac{1}{2}\mathcal{P}(\delta_0^\pm \rightarrow \delta^\pm)[H, K_2]_{\mathcal{U}} \tag{51}$$

We have finally

$$\mathcal{H}_2 = U\Delta_0(-2\mathcal{S} + \mathcal{A} + \mathcal{A}^\dagger + 4\mathbf{V} - 2V) + U[\Delta^- (4\mathbf{R} - 2\mathcal{R} + R_2) + H.c.],$$

where

$$\begin{aligned}
\Delta_0 &= \langle \delta_2^- \left(\delta_0^+ - \frac{1}{2}\delta^+ \right) + \delta_2^+ \left(\delta_0^- - \frac{1}{2}\delta^- \right) \rangle = \frac{1}{2} \langle \delta_2^- \delta^+ + \delta^- \delta_2^+ \rangle \\
\Delta^- &= \langle \delta_2^- \left(\delta_0^- - \frac{1}{2}\delta^- \right) \rangle = \frac{1}{2} \langle \delta_2^- \delta^- \rangle = (\Delta^+)^*
\end{aligned} \tag{52}$$

E. Driven DNLSE as a classical limit of driven Bose-Hubbard model

Higher-order corrections for the driven Bose-Hubbard model discussed above can be compared with the corrections for a driven DNLSE obtained using canonical perturbation theory. In the limit of large occupation numbers one would expect to obtain similar results for both systems. Indeed, consider a fastly driven DNLSE

$$i\dot{\psi}_n = -J(\psi_{n-1} + \psi_{n+1}) + g|\psi_n|^2\psi_n - \omega E(\omega t)n\psi_n \quad (53)$$

We remove the last term using the transformation $\psi_n = v_n \exp[-in\omega \int_0^t E(\omega t')dt']$, obtaining in new variables

$$i\dot{v}_n = -J(v_{n+1}F(\omega t) + v_{n-1}F^*(\omega t)) + g|v_n|^2v_n, \quad (54)$$

$F(\omega t) = \exp[-i\omega \int_0^t E(\omega t')dt']$ This is an equation corresponding to a classical Hamiltonian

$$H = -J \sum_n [Fv_n^*v_{n+1} + F^*v_n^*v_{n-1}] + \frac{g}{2} \sum_n |v_n|^4, \quad (55)$$

introducing the fast time $\tau = \omega t \equiv t/\epsilon$, we get the Hamiltonian ϵH , where $F(\tau) = \exp[-i\omega \int_0^t E(\omega t')dt'] = \exp[-i \int_0^\tau E(\tau)d\tau] = F_0 + \sum F_l \exp(-il\tau)$. Obviously, $|v_n|^2 = |\psi_n|^2$, so the physical meaning of new and old variables is very similar.

Now, we apply a canonical transformation using the generating function

$$W = i \sum_n v_n V_n^* + i\epsilon S(\mathbf{v}, \mathbf{V}^*, \tau), \quad S = S_1(\mathbf{v}, \mathbf{V}^*, \tau) + \epsilon S_2(\mathbf{v}, \mathbf{V}^*, \tau) + \dots, \quad (56)$$

where $\{\mathbf{v}, \mathbf{V}^*\}$ denote the whole set of variables v_n, V_n^* , $n = 1, \dots, N$

The new and the old variables are related by

$$\begin{aligned} v_n^* &= -i \frac{\partial W}{\partial v_n} = V_n^* + \epsilon \frac{\partial S_1}{\partial v_n} + \epsilon^2 \frac{\partial S_2}{\partial v_n} \\ V_n &= -i \frac{\partial W}{\partial V_n^*} = v_n + \epsilon \frac{\partial S_1}{\partial V_n^*} + \epsilon^2 \frac{\partial S_2}{\partial V_n^*} \end{aligned} \quad (57)$$

We want to obtain an effective time-independent Hamiltonian $\epsilon \mathcal{H}$ using this transformation. The effective Hamiltonian can be expanded in series in ϵ

$$\mathcal{H}(\mathbf{V}, \mathbf{V}^*) = \mathcal{H}_0(\mathbf{V}, \mathbf{V}^*) + \epsilon \mathcal{H}_1(\mathbf{V}, \mathbf{V}^*) + \dots \quad (58)$$

After tedious calculations determining appropriate generating function, one obtains [6]

$$\begin{aligned} \mathcal{H}_1 &= 0, \\ \mathcal{H}_2 &= -J^2 g \Phi (v_n^2 v_{n-1}^* v_{n+1}^* + (v_n^*)^2 v_{n+1} v_{n-1} + 2|v_n|^4 - 4|v_n|^2 |v_{n+1}|^2) \\ &\quad + J^2 g T ((|v_{n-1}|^2 - 2|v_n|^2) v_{n+1} v_{n-1}^* + v_{n+1}^2 (v_n^*)^2) \\ &\quad + J^2 g T^* ((|v_{n+1}|^2 - 2|v_n|^2) v_{n+1}^* v_{n-1} + v_n^2 (v_{n+1}^*)^2) \\ &\quad + J^3(\dots), \end{aligned} \quad (59)$$

where $\Phi \equiv \langle |\tilde{F}|^2 \rangle = \sum_{l \neq 0} \frac{|F_l|^2}{l^2}$, $\tilde{F} = i \sum_{l \neq 0} \frac{F_l}{l} \exp(-ilt)$, $T \equiv - \sum_{l \neq 0} \frac{F_l F_{-l}}{l^2}$.

If we now consider expressions obtained earlier for Bose-Hubbard models, and assume Fourier expansion $\delta_0^+ = F_0 + \sum_{l \neq 0} F_l \exp(-ilt)$, we obtain exactly $\Delta_0 = -J^2 \Phi$, $\Delta^+ = \frac{1}{2} J^2 T$. Therefore, the semiclassical limit of our averaging procedure for the Bose-Hubbard model produces the result analogous to the canonical perturbation theory for DNLSE (where variables v_n are considered as c-number analogs of the creation/annihilation operators of the Bose-Hubbard model).




FIG. 2: (Color online) Parameters Δ_0 (left panel), Δ (right panel) as a function of $\mathcal{E}_0, \mathcal{E}_2$. Upper panel: two-colour driving with double frequency $f(t) = \mathcal{E}_0 \sin t + \mathcal{E}_2 \sin 2t$, bottom panel: two-colour driving with tripled frequency $f(t) = \mathcal{E}_0 \sin t + \mathcal{E}_2 \sin 3t$. Dotted (green) line: zeros of $\Delta = \Delta^+ + \Delta^-$ which controls pair tunnelling. Dashed (red) line: zeros of effective single-particle tunnelling J_e .

F. Engineering the parameters of the effective Hamiltonian.

Here we show how a simple non-monochromatic perturbation allows to vary parameters Δ_0, Δ^\pm of the effective Hamiltonian.

Let $f(t) = \mathcal{E}_0 \sin t + \mathcal{E}_2 \sin 2t$, i.e. a two-parameter, bichromatic driving. Then,

$$\begin{aligned} \delta_0^+ &= J \exp\left[i(\mathcal{E}_0 \sin t + \mathcal{E}_2 \sin 2t)\right] = J \sum_n J_n(\mathcal{E}_0) e^{int} \sum_m J_m(\mathcal{E}_2) e^{i2mt}, \\ \langle \delta_0^+ \rangle &= J \sum_m J_m(\mathcal{E}_2) J_{-2m}(\mathcal{E}_0), \quad \delta^+ = J \sum_{n \neq -2m} \sum_m J_n(\mathcal{E}_0) J_m(\mathcal{E}_2) e^{i(n+2m)t}, \end{aligned} \quad (60)$$

$$\langle \delta_2^+ \delta^- \rangle = -J^2 \sum_{n_1 \neq -2m_1} \sum_{m_1, m_2} \frac{J_{n_1}(\mathcal{E}_0) J_{m_1}(\mathcal{E}_2) J_{m_2}(\mathcal{E}_2) J_{n_1+2m_1-2m_2}(\mathcal{E}_0)}{(n_1 + 2m_1)^2} \quad (61)$$

Plotting $\Delta_0, \Delta = \Delta^- + \Delta^+$ as a function of parameters $\mathcal{E}_0, \mathcal{E}_2$ (Fig 2-5), one notes that it is possible to (i) nullify single-particle and doublon tunnelling simultaneously, while keeping exchange interaction large (ii) nullify single-particle tunnelling keeping only two-particle tunnelling and exchange interactions (iii) nullify doublon tunnelling while keeping single-particle tunnelling finite.

G. Driven fermionic two-site model

For the fermionic two-site model we wish to obtain even higher order (fourth order and beyond) corrections. Expressions for them become very bulky. However, in the case of strong interaction (large U), we can keep only the terms of the highest order in U in all terms of fourth order in $1/\omega$ and higher. Such diagrammatic-like technique allows to obtain very accurate expressions for the effective Hamiltonian. In the expressions below the subscript U denotes the leading term in U . So, starting with

$$H = \delta_0^+(t)(c_{1\uparrow}^\dagger c_{2\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow}) + \delta_0^-(t)(c_{2\uparrow}^\dagger c_{1\uparrow} + c_{2\downarrow}^\dagger c_{1\downarrow}) + U(n_{1\uparrow}n_{1\downarrow} + n_{2\uparrow}n_{2\downarrow}) \quad (62)$$

we have

$$\mathcal{H}_0 = \langle H \rangle = J_0(c_{1\uparrow}^\dagger c_{2\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow}) + J_0(c_{2\uparrow}^\dagger c_{1\uparrow} + c_{2\downarrow}^\dagger c_{1\downarrow}) + U(n_{1\uparrow}n_{1\downarrow} + n_{2\uparrow}n_{2\downarrow}) \approx U(n_{1\uparrow}n_{1\downarrow} + n_{2\uparrow}n_{2\downarrow}) \equiv \langle H \rangle_U \quad (63)$$

$$i\dot{K}_1 = \{H\}, \quad K_1 = -i \left(\delta_1^+(c_{1\uparrow}^\dagger c_{2\uparrow} + c_{1\downarrow}^\dagger c_{2\downarrow}) + \delta_1^-(c_{2\uparrow}^\dagger c_{1\uparrow} + c_{2\downarrow}^\dagger c_{1\downarrow}) \right) \sim U^0 \quad (64)$$

$$[\{H\}, K_1] = (\delta_1^+ \delta^- - \delta_1^- \delta^+)(n_1 - n_2) \quad (65)$$

$$\begin{aligned} i\dot{K}_2 &= \left\{ \left[H - \frac{\{H\}}{2}, K_1 \right], K_1 \right\} = \left\{ \left[\langle H \rangle + \frac{\{H\}}{2}, K_1 \right], K_1 \right\} \\ &= \left[\langle H \rangle + \frac{\{H\}}{2}, K_1 \right] - \frac{1}{2} \langle [\{H\}, K_1] \rangle = \left[\langle H \rangle + \frac{\{H\}}{2}, K_1 \right] \approx [\langle H \rangle_U, K_1] \end{aligned} \quad (66)$$

$$\mathcal{H}_1 = \frac{1}{2} \langle [\{H\}, K_1] \rangle = 0 \quad (67)$$

$$\begin{aligned} K_2 &= -\delta_{02}(n_{2\uparrow} - n_{1\uparrow} + n_{2\downarrow} - n_{1\downarrow}) + U \sum_{\sigma} (n_{1,\sigma} - n_{2,-\sigma})(\delta_2^- c_{2,\sigma}^\dagger c_{1,\sigma} - \delta_2^+ c_{1,\sigma}^\dagger c_{2,\sigma}) \\ &\approx U \sum_{\sigma} (n_{1,\sigma} - n_{2,-\sigma})(\delta_2^- c_{2,\sigma}^\dagger c_{1,\sigma} - \delta_2^+ c_{1,\sigma}^\dagger c_{2,\sigma}) \equiv K_{2U} \end{aligned} \quad (68)$$

$$\mathcal{H}_2 = \frac{1}{2} \langle [\{H\}, K_2] \rangle + \frac{1}{12} \langle [\{[\{H\}, K_1]\}, K_1] \rangle \quad (69)$$

Since

$$\langle [\{[\{H\}, K_1]\}, K_1] \rangle = 2i \langle \delta_1^- (\delta_1^+ \delta^- - \delta_1^- \delta^+) \rangle (c_{2\downarrow}^\dagger c_{1\downarrow} + c_{2\uparrow}^\dagger c_{1\uparrow}) + h.c., \quad (70)$$

the last term in Eq.(69) adds a small correction to effective tunnelling constant $J_{eff} = J_0$. This is a finite size effect which is absent in infinite 1D lattice or 1D lattice with periodic boundary conditions. The first term gives

$$\begin{aligned} \frac{1}{2} \langle [\{H\}, K_2] \rangle &= \left[\langle \delta_{02} \delta^- \rangle (c_{2\downarrow}^\dagger c_{1\downarrow} + c_{2\uparrow}^\dagger c_{1\uparrow}) + h.c. \right] \\ &+ \langle (\delta_2^+ \delta^- + \delta_2^- \delta^+) \rangle U \left[(n_{1\downarrow} - n_{2\downarrow})(n_{1\uparrow} - n_{2\uparrow}) - (c_{2\uparrow}^\dagger c_{1\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} c_{1\uparrow}^\dagger c_{2\uparrow}) \right] \\ &- 2 \left[\langle \delta_2^- \delta^- \rangle U d_2^\dagger d_1 + h.c. \right], \end{aligned} \quad (71)$$

where we introduce doublon operators $d_k = c_{k\downarrow}c_{k\uparrow}$, $k = 1, 2$.

While general expressions for \mathcal{H}_k, K_k become very complicated with increasing k , leading term in U can be calculated:

$$\mathcal{H}_3 \approx \langle \delta_1^+ \delta_2^- - \delta_1^- \delta_2^+ \rangle (n_{1\downarrow} n_{1\uparrow} - \dots) = 0, \quad \text{if } |F_l|^2 = |F_{-l}|^2 \quad (72)$$

$$\begin{aligned} \mathcal{H}_4 \approx & -U^3 \langle \delta_4^- \delta^+ + \delta_4^+ \delta^- \rangle \left[(n_{1\downarrow} - n_{2\downarrow})(n_{1\uparrow} - n_{2\uparrow}) - (c_{2\uparrow}^\dagger c_{1\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} c_{1\uparrow}^\dagger c_{2\uparrow}) \right] \\ & + 2U^3 \left[\langle \delta_4^- \delta^- \rangle d_2^\dagger d_1 + h.c. \right] \end{aligned} \quad (73)$$

I.e., the leading term in \mathcal{H}_4 has exactly the same structure as in \mathcal{H}_2 , just with different coefficients. It is not difficult to sum up infinite number of leading terms of \mathcal{H}_k with the same structure.

For the harmonic perturbation used in [7] one has

$$\begin{aligned} \langle \delta_4^+ \delta^- \rangle &= \langle \delta_4^- \delta^+ \rangle = \frac{A_4}{2} = 2J^2 \sum_{m=1}^{\infty} \frac{J_m^2(\mathcal{E}_0)}{m^4} \\ \langle \delta_4^- \delta^- \rangle &= \langle \delta_4^+ \delta^+ \rangle = \frac{C_4}{2} = 2J^2 \sum_{m=1}^{\infty} \frac{(-1)^m J_m^2(\mathcal{E}_0)}{m^4} \\ \langle \delta_2^+ \delta^- \rangle &= \langle \delta_2^- \delta^+ \rangle = \frac{A_2}{2} = -2J^2 \sum_{m=1}^{\infty} \frac{J_m^2(\mathcal{E}_0)}{m^2} \\ \langle \delta_2^- \delta^- \rangle &= \langle \delta_2^+ \delta^+ \rangle = \frac{C_2}{2} = -2J^2 \sum_{m=1}^{\infty} \frac{(-1)^m J_m^2(\mathcal{E}_0)}{m^2} \end{aligned} \quad (74)$$

$$\begin{aligned} \mathcal{H}_{eff} \approx & \mathcal{H}_0 + \left(A_2 \frac{U}{\omega^2} - A_4 \frac{U^3}{\omega^4} + \dots \right) \left[(n_{1\downarrow} - n_{2\downarrow})(n_{1\uparrow} - n_{2\uparrow}) - (c_{2\uparrow}^\dagger c_{1\uparrow} c_{1\downarrow}^\dagger c_{2\downarrow} + c_{2\downarrow}^\dagger c_{1\downarrow} c_{1\uparrow}^\dagger c_{2\uparrow}) \right] \\ & - \left(C_2 \frac{U}{\omega^2} - C_4 \frac{U^3}{\omega^4} + \dots \right) (d_2^\dagger d_1 + h.c.) \end{aligned} \quad (75)$$

Denoting expressions in round brackets as A and C , correspondingly, we get the following matrix elements of the effective Hamiltonian in the basis of 4 states of the two-site system:

$$\mathcal{H}_{eff} = \begin{pmatrix} -A & -A & J_e & J_e \\ -A & -A & J_e & J_e \\ J_e & J_e & U+A & -C \\ J_e & J_e & -C & U+A \end{pmatrix}, \quad (76)$$

where J_e denotes effective tunnelling.

The half-difference between the two lowest levels is

$$\frac{\Delta E}{2} = J_{ex} = \frac{1}{4} \left(A + C - U + \sqrt{16J_e^2 + (U + 3A - C)^2} \right) \approx \frac{2J_e^2}{U} + A + \frac{2J_e^2}{U^2} (C - 3A) + \dots \quad (77)$$

For the harmonic perturbation summation of infinite number of leading order in U terms leads to

$$A = -4J^2 \sum_{m=1}^{\infty} J_m^2(\mathcal{E}_0) \left(\frac{U}{\omega^2 m^2} + \frac{U^3}{\omega^4 m^4} + \dots \right) = -4J^2 U \sum_{m=1}^{\infty} \frac{J_m^2(\mathcal{E}_0)}{m^2 \omega^2 - U^2} \quad (78)$$

Note that although expression (78) can change its sign at $\omega < U$, it was obtained as a summation of a geometric series with a multiplier $\frac{U^2}{m^2 \omega^2}$, therefore it is assumed $\omega > U$. The case $\omega < U$ deserves further consideration,




FIG. 3: (Color online) Induced exchanged interaction in the driven fermionic two-site model. Solid curves, from up to down: theoretical prediction Eq. (77) with corrections up to the second order in $\frac{1}{\omega}$ taken into account (upper curve); with corrections up to the fourth order in $\frac{1}{\omega}$ taken into account (lower curve). Dashed curve: numerical value of the exchange interaction (numerical data kindly provided by J.Mentink). Dotted curve: effective tunnelling J_e . It is seen that in the regions where J_e is strongly suppressed due to driving, second order corrections become insufficient, and one needs to include fourth and higher orders in $\frac{1}{\omega}$.

in particular because resonances between $m\omega$ and U can happen. It is a remarkable fact however that this formal expression is correct also for $\omega < U$, according to [7]

-
- [1] W. Magnus, *Comm. Pure and Appl. Math.* VII, 649 (1954).
 - [2] S. Blanes, F. Casas, J.A. Oteo, J. Ros, *Physics Reports* 470, 151 (2009).
 - [3] A.P. Itin, A.I. Neishtadt, *Phys. Lett. A* 378, 822 (2014)
 - [4] A. Eckardt, C. Weiss, and M. Holthaus, *Phys. Rev. Lett.* 95, 260404 (2005).
 - [5] S. Longhi, *Phys. Rev. B* 77, 195326 (2008)
 - [6] A.P. Itin et al., in preparation.
 - [7] J. H. Mentink, K. Balzer, and M. Eckstein, arXiv:1407.4761 (2014).