

# A THEORY OF EMERGENT QUANTUM MECHANICS

Ricardo Gallego Torromé<sup>1</sup>

Departamento de Matemática, Universidade Federal de São Carlos, Rodovia Washington Luís, km 235, SP, Brazil

**ABSTRACT.** Cartan-Randers dynamical systems, a particular class of first order ordinary differential dynamical systems, are considered as mathematical description for deterministic emergent quantum mechanics. We show that in such framework local diffeomorphism invariance, reversibility of the effective quantum dynamics and a covariant maximal universal acceleration emerge from a fundamental classical, deterministic and local (in a novel sense) dynamics. Starting from the elements of Cartan-Randers models, a phenomenological Hilbert space is constructed and associated with the space of wave functions of quantum mechanics. A geometric description of spontaneous reduction of the quantum state is described. The mechanism is applied heuristically to the problem of a quantum particle passing through two small slits. Furthermore, the same geometric mechanism is shown to imply the boundness from below of the matter Hamiltonian. The mechanism also provides an explanation for the weak equivalence principle. This fact, together with the local Lorentz invariance of the models and diffeomorphism invariance of the interaction driving the reduction, suggest that the reduction of the quantum state is driven by a gravitational type interaction. Moreover, since such interaction appears only in the concentration domain of the internal dynamics, it must be associated with a classical interaction.

**Keywords.** Emergent quantum mechanics, emergence of weak equivalence principle, emergence of diffeomorphism invariance, Cartan spaces, Randers spaces, maximal acceleration, concentration of measure, dissipative dynamics, ergodicity.

## CONTENTS

1. Introduction	2
1.1. Motivation	2
1.2. Structure of the work	3
2. Assumptions for deterministic models at the Planck scale	4
2.1. Some remarks on the assumptions	5
Maximal acceleration	5
3. Deterministic Cartan-Randers systems	6
3.1. Geometric background	6
3.2. Notion of Cartan-Randers Space	7
3.3. Some properties of the $U_t$ dynamics	9
t-time inversion	10
3.4. $U_t$ is an information loss dynamics	11
3.5. The $U_\tau$ dynamics	12
4. Hilbert space formulation of DCRS	13
5. Quantum mechanical structures from DCRS	15
5.1. Construction of local quantum phases	15
5.2. The bundle associated with local quantum phases	16
5.3. Definition of wave function in DCRS	16

---

<sup>1</sup>email: rigato39@gmail.com

5.4. Hilbert space associated with a DCRS	17
6. Concentration of measure and measurement processes in DCRS	17
6.1. Concentration of measure	17
6.2. An application to dynamical reduction of quantum states	18
6.3. Description of the two slit quantum experiment in DCRS	19
7. Hamiltonian boundeness and emergence of the weak equivalence principle and gravitational interaction	20
7.1. Boundeness of the Hamiltonian	20
7.2. Emergence of the weak equivalence principle	23
7.3. Emergence of the gravitational interaction	25
8. Discussion	25
8.1. Cartan-Randers systems and other deterministic models for emergent quantum mechanics	25
8.2. Open problems and further developments	27
Acknowledgements	29
References	29

## 1. INTRODUCTION

**1.1. Motivation.** The fundamental laws of physics assume the existence of a geometric spacetime where *events happen*. This applies also to the standard formulation of the quantum theory, including the description of the measurement processes, where the spacetime is the arena where they happen. After the advent of general relativity, the spacetime structure becomes dynamical and it is not only a geometric back-ground for other processes to happen. This is in sharp contrast with the quantum theory, where the geometric structure is fixed and non-dynamical. Given this dichotomy on the foundations of these two universal theories of physics, it appears as a fundamental problem to find a theory that merges both or to have a unified framework for a the relativistic theory of gravitation and the quantum theory.

Even if only as an approximate description to a more accurate theory, to find a consistent spacetime representation of measurement process, non-local quantum correlations and a geometric understanding of the notion of entanglement is a very difficult task. In the case of the quantum measurement processes this is not surprising, since the standard *collapse postulate* of quantum mechanics involves the instantaneous reduction of the wave packet, an interpretation which is clearly against the spirit of the special theory of the relativity. The formal unification of quantum mechanics with special relativity as are the relativistic quantum field theories is not a consistent solution, since measurement processes are not considered in the relativistic quantum field framework.

Due to the difficulties with the spacetime interpretation of the quantum mechanical processes and assuming that fundamental physical phenomena should be representable in a geometric framework, it is reasonable to doubt that the quantum theory as currently stands is the ultimate theoretical framework for the description of physical processes. Clearly, the problem of the spacetime representation by a macroscopic observer of quantum entanglement and quantum measurement processes is related with the logical and mathematical structure of the quantum theory itself and with the ontological interpretation of the fundamental physical processes.

A general type of pre-quantum framework are recognized under the name of *emergent quantum mechanics*. Such frameworks share the idea that the quantum formalism arises from an underlying, more fundamental theory. We cannot make

full justice here to the whole contribution to these research lines, but let us mention as examples of emergent quantum frameworks the theories developed in [2, 9, 10, 13, 16, 24].

A fundamental difficulty for deterministic quantum models in the line of Hooft's approach is that the associated Hamiltonian operators, being linear in momentum variables, are not bounded from below. One of the mechanism proposed in the literature to solve this problem involves a dissipative dynamics [9, 13, 24]. In those proposals, the gravitational interaction plays an essential role as the origin of the information loss dynamics and must be present at the level of the fundamental dynamics. However, gravity could be a classical and emergent phenomenon, not present at the scale energy where the dynamics of the microscopic fundamental degrees of freedom happens. If this is the case, it is not natural to appeal from the beginning to gravity as the origin of dissipation of information.

Another fundamental difficulty associated with deterministic quantum models is the relation between the degrees of freedom at the fundamental scale with the degrees of freedom at subatomic, atomic and classical scales. However, some relevant results have been obtained. In particular, it has been shown that some  $(1+1)$ -bosonic quantum field models and some string models can be interpreted as deterministic quantum models [27, 28]. Also, it was shown by Hooft's that massless non-interacting neutrino's field can be identified with a deterministic system [24]. Such examples show that there is the possibility to describe the dynamics of non-trivial quantum systems from the dynamics of deterministic systems.

In this work, that supersedes the previous work of the author on the same topic [16], we develop several new ideas for an underlying deterministic theory beneath quantum mechanics. In this work it is shown how the quantum mechanical phenomenology can emerge from an underlying classical dynamics. We start describing the relation between the classical Hamiltonian formulation of first order dynamical systems and *symmetrized* Cartan-Randers spaces. The correspondence is based on the relation between Hamiltonian systems and Cartan spaces and is resumed as a correspondence,

*First order Hamiltonian dynamical system with maximal acceleration and speed*  
are in relation one to one with *deterministic Cartan-Randers dynamical systems*.

The aim of this work is to explain and prove this result and show its implications for a theory beneath quantum theory and its implications suggesting that gravity is a classical, emergent theory.

**1.2. Structure of the work.** The structure of this work is the following. In *section 2* we introduce several assumptions that we make on the nature of the physical systems at the Planck scale. From such assumptions we argue that there must exist a maximal acceleration. In *section 3* is provided a short exposition of Cartan spaces of Randers type and, after time symmetrization, we will show the relation of Cartan-Randers geometry with dynamical systems. We will describe the two dynamics  $U_t$  and  $U_\tau$  associated with DCRS. This is a mathematical fact of fundamental relevance for our theory, since the interpretation of some quantum phenomena depends on the correct interpretation of the mathematics of DCRS, specially the notion of two dimensional time associated to the two dimensional parameter  $(t, \tau)$  that appear in the theory. The existence of maximal acceleration and speed for DCRS systems is shown. In *Section 4* we will explain an idea from G.'t Hooft on why it is useful to consider a quantized version formulation of classical dynamical systems. The underlying motivation for this was to show the difficulties one has to differentiate between classical and quantum systems, specially when the dimensionality of the

configuration space. After this, we will formulate *DCRS* using the quantum formalism of Hilbert spaces and operators. The receipt is clear and taken from Hooft work. *Section 5* is dedicated to construct the *quantum local phase* from the original degrees of freedom of a DCRS and to make the transition from discrete degrees of freedom (associated with a DCRS) to continuous degrees of freedom (associated with wave functions). It is provided the construction of the quantum wave function and quantum Hilbert space. In *Section 6* several fundamental mathematical notions of the quantum theory are derived from DCRS and from the mathematical theory of *asymptotic geometric analysis*. We apply the concentration of measure to explain the spontaneous reduction of the wave function. Moreover, a description of the quantum measurement process for a quantum system, based one more time on the concentration of measure is developed. In *section 7* it is shown how the existence of an equilibrium domain provides a natural solution to the problem of the un-boundedness from below of the linear quantum Hamiltonian for deterministic models. We also discuss how the weak equivalence principle is emergent in DCRS and as a consequence, why gravity is emergent in DCRS theory. A short discussion of the DCRS framework is presented in *section 8* and several open problems of our approach are discussed.

## 2. ASSUMPTIONS FOR DETERMINISTIC MODELS AT THE PLANCK SCALE

Our goal is to find mathematical models for emergent quantum mechanics systems that are deterministic, local and causal. Certain assumptions must be done, in order to constraint the class of dynamical systems. We have found that the following assumptions hold naturally for this purpose:

- A.0.

**Definition 2.1.** *An asymmetric metric is a function  $d : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$  such that*

- $d(u, v) \geq 0$ , for each  $u, v \in \mathcal{M}$ .
- $d(u, u) = 0$ , for each  $u \in \mathcal{M}$
- $d(u_a, u_b) + d(u_b, u_c) \geq d(u_a, u_c)$ , for each  $u_a, u_b, u_c \in \mathcal{M}$ .

The configuration space  $\mathcal{M}$  for the fundamental degrees of freedom is a topological space endowed with an *asymmetric metric*,

The symmetry condition has been dropped from the usual notion of metric function. The need for a metric structure is necessary because we will consider metric properties of high dimensional spaces.

- A.1. The sub-quantum degrees of freedom are deterministic. Their dynamics is described by first order ordinary differential equations respect to a macroscopic time parameter  $\tau$ , if the time parameter  $\tau$  can be approximated by a continuous parameter without significantly affecting the macroscopic description. The dynamics is described by finite difference equations, if the time parameter  $\tau$  is discrete.
- A.2. The sub-quantum degrees of freedom are identical, undistinguishable degrees of freedom composed by two *sub-quantum atoms*. One of such sub-quantum atoms evolves toward the future, while the companion evolves towards the past in the external time  $\tau$ . We assume for simplicity that they can be described as point particles.
- A.3. The following *locality condition* holds: given a *system*  $\mathcal{S}$  corresponding to a collection of sub-quantum molecules, there is a small neighborhood  $\mathcal{U}$  with  $\mathcal{S} \subset \mathcal{U}$  such that only  $\mathcal{U}$  acts dynamically on the system  $\mathcal{S}$  in the sense that for any larger  $\tilde{\mathcal{U}} \supset \mathcal{U}$ , the dynamical effect of any action of  $\mathcal{U}$  and  $\tilde{\mathcal{U}}$  on  $\mathcal{S}$  are the same.

- A.4. The following *causality condition* holds: there is a maximal, universal bound for the speed of the sub-quantum atoms and sub-quantum molecules. We assume that the maximal relative speed between different sub-quantum atoms (and sub-quantum molecules) is the speed of the light in vacuum.

The above assumptions constraint significantly the possible mathematical models. In this work we will discuss a category of dynamical systems (that we have called Cartan-Randers models) that completely fulfill these requirements and that contain several relevant additional features that makes them suitable candidates for sub-quantum physics models.

**2.1. Some remarks on the assumptions.** There are some remarks on the assumptions that are worthily to point out. The assumption A.0 on the existence of an asymmetric metric structure will be applied in two ways. First, note that if an asymmetric metric structure is given, there is a notion of proximity. Indeed, given an asymmetric metric  $d : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}$ , a conventional metric function can be constructed by symmetrization and is given by the expression

$$(2.1) \quad d_S : \mathcal{M} \times \mathcal{M} \rightarrow \mathbb{R}, \quad d_S(u, v) = \frac{1}{2} (d(u, v) + d(v, u)).$$

There is a topology associated with  $d$ , which is equivalent to the topology associated to  $d_S$ . Moreover, one have a notion of Cauchy's sequence and completeness. The underlying reason to start with an asymmetric metric is because it can be associated with an underlying non-reversible dynamics, from which a reversible dynamics quantum dynamics emerges. The second use of the asymmetric metric  $d$  is to define a probability measure on  $\mathcal{M}$ . It is with this probability measure that notions as ergodicity and concentration dynamics are conveniently introduced.

A relevant example of asymmetric metric structure is a Randers space, originally described by G. Randers to describe in a geometric way the irreversible nature of the physical world [39]. We will introduce related Randers structures but with a different aim, that is, as models for deterministic, local, non-reversible dynamics at the Planck scale.

**Maximal acceleration.** A direct consequence of the assumptions A.1, A.3 and A.4 is the existence of a *maximal universal acceleration* for sub-quantum atoms and sub-quantum molecules. Since  $\mathcal{U}$  is the maximal region in the spacetime that can affect the dynamics of  $\mathcal{S}$ , there is associated with the size of  $\mathcal{U}$  a minimal length. Then the maximal elementary mechanical work as a result of the action of  $\mathcal{U}$  on  $\mathcal{S}$  is

$$L_{min} m a \simeq \frac{1}{2} (\delta m v_{max}^2 + m(\delta v)_{max}^2),$$

where  $a$  is the value of the acceleration in the direction of the total exterior effort is done and the parameter  $m$  is the inertial mass of the sub-quantum atom. Since the speed of any physical degree of freedom is bounded by the hypothesis of locality A.3,  $\delta v_{max} = v_{max} = c$ , where we can assume that  $c$  is the speed of light in vacuum. The maximal work at rest produced by the system on a point particle is  $\delta m \simeq m$ , because the most natural way to satisfy assumption A.4 is to assume that the theory is local Lorentz invariant, in which case the relation  $E = mc^2$  holds. These relations imply an universal bound for the value of the coordinate acceleration  $a$  for the sub-quantum atoms,

$$(2.2) \quad a_{max} \simeq \frac{c^2}{L_{min}}.$$

Therefore, if the above assumptions A.1, A.3 and A.4 hold, the mathematical formalism for the deterministic models should be constrained to have a maximal universal proper acceleration. It must be universal, since (2.2) does not depend on the details of the system. It also must be proper, because we have use Einstein's relativistic relation for the energy in a co-moving frame.

This heuristic derivation of an universal maximal acceleration for fundamental systems differs from Caianiello's original derivation of the existence of maximal accelerations [11]. Although Caianiello's derivation was based on the causality condition of a limit speed for matter, the second fundamental assumption was the non-degeneracy of a associated Fisher-type pseudo-Riemannian metric in a cotangent space of a point particle. Furthermore, in Caianiello's theory, maximal acceleration depends on the mass of the system, while in the dynamical systems that we are considering, the maximal acceleration is universal and independent of the mass of the system.

From the above argument, it is natural that we require the following additional assumptions for our dynamical systems,

- A.5. There is a maximal, universal acceleration for both sub-quantum atoms and di-atomic sub-quantum molecules.

A framework for deterministic systems with maximal acceleration and maximal speed such that the above assumptions hold is *deterministic Cartan-Randers systems*<sup>2</sup> (or in short, DCRS). Deterministic Cartan-Randers systems are based on a general correspondence between a particular class of first order ordinary differential equations systems and *time oriented symmetrized Cartan-Randers metrics*, defined on a convenient cone of the cotangent bundle  $T^*TM$  (see for instance for the notion of Cartan spaces and [33, 39] for the notion of Randers space in different contexts).

### 3. DETERMINISTIC CARTAN-RANDERS SYSTEMS

**3.1. Geometric background.** We assume the existence of a model *four-manifold spacetime*  $M_4$  that corresponds to the manifold of observable macroscopic events. The configuration manifold is a tangent space  $TM$  with  $\dim(TM) = 2 \dim(M) = 8N$ , such that  $M$  has a product structure of the form

$$(3.1) \quad M = \prod_{k=1}^N M_4^k,$$

with each  $M_4^k$  diffeomorphic to  $M_4$ . The tangent configuration space for a classical gas of point particles is a smooth manifold  $M$  of the form

$$(3.2) \quad TM = \prod_{k=1}^N TM_4^k.$$

We assume that the dimension  $\dim(TM) = 8N$  is large compared with  $2 \dim(M_4) = 8$ . Each of the sub-quantum molecules is labeled by a natural number  $k \in \{1, \dots, N\}$ . The configuration manifold  $TM_4^k$  is the configuration space for the  $k$ -fundamental molecule. The dimension  $\dim(TM_4^k) = 8$  corresponds to four spacetime coordinates  $(\xi^1, \xi^2, \xi^3, \xi^4)$  for the point  $\xi(k) \in M_4^k$  and four independent velocities coordinates  $T_\xi M_4^k \ni (\dot{\xi}^1, \dot{\xi}^2, \dot{\xi}^3, \dot{\xi}^4)$ . There will not be spin degrees of freedom at the fundamental scale and for fundamental dynamics. The present view of spin and other quantum numbers is that they are not fundamental properties an are better associated with phenomenological constructions theory. However, there is nothing

---

<sup>2</sup>In ref. [16] such dynamical systems were named *deterministic Finslerian models*. In the present work we have called such dynamical systems deterministic Cartan-Randers systems, since they are constructed using as a back-ground Cartan spaces [33] of *Randers type* [39].

prohibiting to modified our formalism to introduce spin or other quantum numbers from the beginning in the parametrization of the fundamental systems.

The model 4-manifold  $M_4$  is endowed with a Lorentzian metric  $\eta_4$  of signature  $(1, -1, -1, -1)$ . We assume that the metric  $\eta_4$  is a background, non-dynamical structure. This preliminary treatment of the metric structure is useful to fix many of the concepts of the foundations without deviating from the main problem considered. Moreover, for each  $k \in \{1, \dots, N\}$  there is a Lorentzian metric  $\eta_4(k)$  on  $M_4^k$ . Moreover, we assume that the structures

$$\{(M_4^k, \eta_4(k)), k = 1, \dots, N\}$$

are isometric to  $(M_4, \eta_4)$ . Then we have a collection of diffeomorphisms  $\{\phi_k : M_4^k \rightarrow M_4, k = 1, \dots, N\}$ . This is in accordance with the assumption that the  $N$  degrees of freedom are indistinguishable and identical. The algebra of functions  $\mathcal{F}_D(T^*TM)$  on  $T^*TM$  that we consider is the algebra *diagonal functions*  $\mathcal{F}_D(T^*TM)$ , obtained by an algebra embedding

$$(3.3) \quad \begin{aligned} \theta : \mathcal{F}(T^*TM_4) &\rightarrow \mathcal{F}(T^*TM), \quad f \mapsto (f_1, \dots, f_N) \\ f_k &= f_0, \quad f_0 : T^*TM_4 \rightarrow \mathbb{R} \quad k = 1, \dots, N. \end{aligned}$$

It is composed by elements of  $\mathcal{F}(T^*TM)$  of the form

$$((u_1, p_1), \dots, (u_N, p_N)) \mapsto (f(u_1, p_1), \dots, f(u_N, p_N)).$$

It is of relevance for our considerations that the average values of the functions  $f \in \mathcal{F}_D(T^*TM)$  are well defined, that is, independent of the diffeomorphisms  $\{\phi_k : M_4^k \rightarrow M_4, k = 1, \dots, N\}$ . This can be achieved if the measure used in the averages  $\mu_P$  in  $T^*TM$  is the product measure

$$(3.4) \quad \mu_P = \prod_{k=1}^N \mu(k)_P,$$

where  $\mu(k)_P$  is a probability measure in  $T^*TM_4^k$ .

Given the above geometric structures, there is a pseudo-Riemannian metric  $\eta_S^*(k)$  defined on  $TM_4^k$  (the Sasaki-type metric), which is a lift of the metric  $\eta_4^k$  on  $M_4^k$  to  $TM_4^k$ , using the Levi-Civita connection of  $\eta_4(k)$  to determine the horizontal piece. The dual metric of  $\eta_S^*$  is the dual pseudo-Riemannian metric  $\eta_S(k) = (\eta_S^*(k))^*$ . The dual Sasaki-type metric  $\eta_S$  allows to define the dual pseudo-Riemannian metric<sup>3</sup>

$$(3.5) \quad \eta = \oplus_{k=1}^N \eta_S(k)$$

on the  $8N$ -dimensional cotangent space  $T^*TM$ .

**3.2. Notion of Cartan-Randers Space.** Let  $\widetilde{M}$  be a smooth manifold and  $\mathcal{C} \subset T^*\widetilde{M}$  a connected open cone of  $T^*M$ . We introduce the notion of *Cartan spacetime* as a direct generalization from the notion in the case positive definite metrics [33],

**Definition 3.1.** A *Cartan space* is a triplet  $(\widetilde{M}, \widetilde{F}, \mathcal{C})$  with  $\widetilde{F} : \mathcal{C} \rightarrow \mathbb{R}^+$  smooth on the open cone  $\mathcal{C} \hookrightarrow T^*\widetilde{M}$  such that

- $\widetilde{F}$  is homogeneous of degree one on the momentum variables and

---

<sup>3</sup>Note that the use of  $*$ -notation for metrics and norms here is inverted respect to the usual notation in Riemannian geometry. For instance  $\eta_S^*$  is a metric on  $TM_4$ , while  $\eta_S$  is a metric in  $T^*M_4$ . Our particular deviated convention is motivated to do not cluster the mathematical expressions in the text, since we will consider geometric constructions on co-tangent spaces.



- The vertical Hessian (or fundamental tensor  $g$ ) of the function  $(\tilde{F})^2$  in natural coordinates  $\{(u^i, p_j), i, j = 1, \dots, \dim(\widetilde{M})\}$

$$(3.6) \quad g^{ij}(u, p) = \frac{1}{2} \frac{\partial^2 (\tilde{F}(u, p))^2}{\partial p_i \partial p_j}.$$

is non-degenerate for each  $u \in \widetilde{M}$ .

The function  $\tilde{F}$  is the Cartan function.

There is a direct relation between Cartan structures and Hamiltonian dynamical systems because, if the fundamental tensor  $g^{ij}(u, p)$  is non-degenerated, the function  $\tilde{F}$  has associated a geodesic and a Hamiltonian flow in  $T\widetilde{M}$  and  $T^*\widetilde{M}$  respectively, the respective projections to  $M$  of both flows coincide.

A relevant example of Finsler space is a Randers space [39],

**Definition 3.2.** A Randers space is a Finsler space whose Finsler function  $F^* : T\widetilde{M} \rightarrow \mathbb{R}$  is  $F^* = \alpha^*(x, y) + \beta^*(x, y)$ , with  $\alpha^*(x, y)$  the Riemannian norm of  $y \in T\widetilde{M}$ ,  $\beta^*(x, y)$  the result of the action 1-form  $\beta^* \in T^*\widetilde{M}$  on  $y$  and such that

$$(3.7) \quad \eta(\beta, \beta) < 1.$$

holds good.

We now consider a similar structure in the category  $\mathcal{F}_C^*(TM)$  of Cartan spaces on  $TM$  with non-definite signature. It is well known that in order to consider Randers-type structures in the category of non-definite metric structures it is necessary to restrict the domain of definition of  $F$ , since it is not possible to have a well defined Randers function on the whole  $T^*TM$ . Given the background metric  $\eta$ , the domain of a Cartan-Randers space will be restricted to the open cones  $\mathcal{C}_u \hookrightarrow T_u^*TM$ , defined by the set of co-vectors  $p \in T_u^*TM$  such that

$$(3.8) \quad \alpha^2(u, p) = \sum_{i,j=1}^{8N} \eta^{ij}(u) p_i p_j \geq 0.$$

Let us consider a vector field  $\beta \in \Gamma TTM$  such that the following condition holds,

$$(3.9) \quad 0 \leq |\eta^*(\beta, \beta)| < 1.$$

The kinematical significance for the dynamical systems that we consider will be clarified later.

**Definition 3.3.** A Cartan-Randers space is a Cartan space whose Cartan function  $\tilde{F}$  is of the form

$$(3.10) \quad F : \mathcal{C} \rightarrow \mathbb{R}^+, \quad (u, p) \mapsto F(u, p) = \alpha(u, p) + \beta(u, p).$$

with  $\alpha$  real and where  $\beta$  is constrained by the condition (3.9).

The space of Cartan-Randers structures on  $TM$  is denoted by  $\mathcal{F}_{CR}(TM)$  and plays a fundamental role in the theory of DCRS.

A Cartan-Randers structure can be seen as a *linearly perturbed* dual Riemannian structure on the tangent space  $TM$ . In analogy with the positive definite case, the requirement (3.9) should imply that the linear perturbation  $\beta(u, p)$  does not introduce a degeneracy in the fundamental tensor (3.6). Note that although controlled by the linear perturbation (3.9), the linear perturbation is not necessary small.

Let us consider the pre-configuration manifold (3.1). At a given point  $u = (x, y) \in TM$  the  $\beta$ -term is given by the expression

$$(3.11) \quad \beta(u, p) = \sum_{i=1}^{8N} \beta^i(u) p_i.$$



where  $\{\beta^i(u)\}_{i=1}^{8N}$  are the components of the vector  $\beta \in \Gamma TTM$ .  $\beta$  acts on the 1-form  $p \in \mathcal{C} \subset T^*TM$ . The boundness condition (3.9) and the restriction to the open cone  $\mathcal{C}$  implies that the Cartan-Randers structure is positive definite and non-degenerate on  $\mathcal{C}$  and therefore, that (3.10) defines a Cartan space on  $\mathcal{C}$ .

One can consider natural geometric flows in space of Cartan-Randers structures  $\mathcal{F}_{CR}(TM)$ . The analogy with Riemannian geometric flows as the celebrated Ricci flow and mean curvature flow are examples of dynamical geometric flows where the geometry evolves. This is in contrast with the situation in General Relativity, where the geometry is fixed by Einstein's equations and where time diffeomorphism invariance implies that there is not notion of time. We will call this dynamics  $t$ -dynamics or  $U_t$  dynamics. It is required that for such dynamics the following to properties hold,

- There is a 1-parameter family of connected submanifolds

$$\{\mathcal{E}_u(r) \hookrightarrow T_u^*TM, u \in TM\}$$

that are stable by the geometric flow.

- The dynamics of the connected components of the  $r$ -hyperboloid

$$\Sigma_u(r) := \{p \in T^*TM \text{ s.t. } F^2(u, p) = r^2\}$$

by the  $t$ -dynamics has as limit the corresponding  $\mathcal{E}_u(r)$ . This limit is reached in a finite time evolution  $\delta t = T$ .

- The equilibrium at  $t = T$  is instantaneously broken, mainly because the properties of maximal acceleration, that prevents the system to stay for a long  $t$ -time collapsed. Then a process of expansion starts again, until the system becomes ergodic.

Therefore, it is of relevance to provide a geometric/number theoretical flow for the dynamics associated with the  $U_t$  dynamics. The relation with number theory seems a bit odd at first sight, but the requirement of coherence in synchronization for different quantum systems and in particular, for different values of the period  $T$ , implies a deep relation between the  $t$ -dynamics and the theory of congruences in Number Theory. This will be provided in a separate paper, where we will develop the relation between Cartan-Randers models and a specific model of  $xp$ -dynamics.

The time scale parameter  $T$ , that has the properties of a time period, depends upon the physical system described. It can be thought as a *measure* of the complexity of the system in the following way. Since the sub-quantum particles are identical, the limit  $t \rightarrow T$  is reached faster for larger systems. Therefore,  $T^{-1}$  is a measurement of the number of components of the system.

**3.3. Some properties of the  $U_t$  dynamics.** Given the Cartan-Randers space  $(TM, F, \mathcal{C})$ , it is possible to define a Riemannian structure  $h \in \Gamma T^{(2,0)}TM$  by averaging the fundamental tensor components  $g_{ij}(u, p)$  on each  $\Sigma_u(1)$  [18]. The tensor components  $h^{ij}$  are obtained by averaging the metric coefficients  $g^{ij}(u, p)$  on the open cone  $\Sigma_u(1)$  over  $u \in TM$ ,

$$(3.12) \quad h^{ij}(u) := \frac{1}{\int_{\Sigma_u(1)} dvol_u(p)} \int_{\Sigma_u(1)} dvol_u(p) g^{ij}(u, p),$$

$i, j = 1, \dots, 8N$ , where non-degenerate volume form on the unit hyperboloid  $\Sigma_u(1)$ . Although the sub-manifold  $\Sigma_u(1)$  is not compact, the volume form  $vol_u(p)$  is chosen such that the integrals in (3.12) are defined. It can be proven that the volume function  $dvol_u(p)$  is invariant under local isometries of the Cartan-Randers function  $F$ . Then the squared norms  $h(u, p)$  and  $g(u, p)$  are defined by

$$h(u, p) = h^{ij}(u) p_i p_j, \quad g(u, p) = g^{ij}(u, p) p_i p_j = F^2(u, p).$$

Note that for  $p$  and  $P$  is in the open cone  $\mathcal{C}_u$  over  $u$  it could happen that  $g^{ij}(u, p)P_i P_j$  is not positive, because the pseudo-Riemannian signature of the fundamental tensor  $g^{ij}(x, p)$ . However, for a Cartan-Randers spacetime it is easily checked that this does not happen for  $\beta$  timelike and if  $(P_i p^i)(p_i \beta^i) > 0$ , which is a compatible condition with timelike worldline for the fundamental degrees of freedom.

**Definition 3.4.** *The  $U_t$  dynamics in the interval  $[0, T] \subset \mathbb{R}$  is a geometric flow of the form*

(3.13)

$$U_t : \mathcal{F}_{CR}(TM) \rightarrow \mathcal{F}_{CR}(TM), F \mapsto F_t = \sqrt{\kappa(g, t) |h(u, p)| + (1 - \kappa(g, t)) |g(u, p)|},$$

such that the function  $\kappa : \mathcal{F}_{CR}(TM) \times [0, T] \rightarrow [0, T] \subset \mathbb{R}$  satisfies

$$(3.14) \quad \lim_{t \rightarrow 0} \kappa(g, t) = 0, \quad \lim_{t \rightarrow T} (\kappa(g, t) - 1) = 0.$$

Note that in defining this dynamics  $t \in [0, T]$ , instead of being in  $\mathbb{R}$ . This is because the dynamics becomes almost periodic, with period  $2T$  (that encompasses an expanding phase, an ergodic phase and a concentration phase). Thus we concentrate our attention in one elementary semi-cycle  $t \in [0, T]$ . The  $U_t$  dynamics is dissipative. Consider the convex hull  $C_{hg} \subset \mathcal{F}_{CR}(TM)$  containing  $g$  and  $h$ ,

$$C_{hg} := \{F \in \mathcal{F}_{CR}(TM) \text{ s.t. } g_F = t_1 g + t_2 h, t_1 + t_2 = 1, t_1, t_2 \geq 0\},$$

where  $g_F$  is the fundamental tensor of  $F$ .

**Proposition 3.5.** *Every Cartan-Randers structure  $F$  in the convex hull  $C_{hg}$  containing  $g$  and  $h$  evolves towards the averaged structure  $h \in \Gamma T^{(2,0)}TM$ ,*

$$(3.15) \quad \lim_{t \rightarrow T} U_t(F) = \sqrt{h(u, p)}, \quad \forall F \in C_{hg}.$$

*Proof.* Note that any  $F \in C_{hg}$  has the same averaged metric,

$$\langle g_F \rangle = \langle (t_1 g + t_2 h) \rangle = t_1 h + t_2 h = h.$$

Then we have

$$\begin{aligned} \lim_{t \rightarrow T} U_t(F) &= \lim_{t \rightarrow T} \sqrt{\kappa(g, t) h(u, p) + (1 - \kappa(g, t)) g_F(u, p)} \\ &= \lim_{t \rightarrow T} \sqrt{\kappa(g, t) h(u, p) + (1 - \kappa(g, t)) g_F(u, p)} \\ &= \sqrt{h(u, p)}. \end{aligned}$$

□

Therefore, the  $U_t$  dynamics is a dissipative dynamics in  $\mathcal{F}_{CR}(TM)$ , since  $h$  is the limit of the convex hull  $C_{hg}$ . Therefore, one can classify the elements of  $\mathcal{F}_{CR}(TM)$  in equivalence classes, each equivalence class having the same averaged structure.

**t-time inversion.** The parameter  $t \in \mathbb{R}$  is interpreted as the time parameter for an *internal dynamics* of the system. The *time inversion operation*  $\mathcal{T}_t$  is defined in local natural coordinates on  $T^*TM$  by the expression

(3.16)

$$\mathcal{T}_t : T^*TM \rightarrow T^*TM, (u, p) = (x, y, p_x, p_y) \mapsto (\mathcal{T}_t(u), \mathcal{T}_t^*(p)) = (x, -y, -p_x, p_y).$$

There is an induced action of  $\mathcal{T}_t$  on the space  $\mathcal{F}_{CR}(TM)$  by the expression

$$\mathcal{T}_t(F)(u, p) := F(\mathcal{T}_t(u), \mathcal{T}_t(p)).$$

Note that a Cartan-Randers metric is non-reversible in the sense that

$$F(u, p) \neq F(\mathcal{T}_t(u), \mathcal{T}_t(p))$$

except for a subset of measure zero in  $(u, p) \in T_u^*TM$ . From this relation it follows the intrinsic irreversible character of the Randers geometry [39]. However, we assume the natural condition that such action commutes with the  $U_t$  dynamics,

$$(3.17) \quad [U_t, \mathcal{T}_t] = 0, \quad \forall t \in [0, T] \subset \mathbb{R}.$$

This commutation relation guarantees that  $\mathcal{T}_t(F)$  and  $F$  are in the same class of equivalence  $[h]$ , if  $h$  is invariant under  $\mathcal{T}_t$ ,

$$\mathcal{T}_t h = h,$$

since equivalence classes are disjoint sets in  $\mathcal{F}_{CR}(TM)$ . Indeed, one can prove by a continuity argument on the parameter  $t$  that for  $t = n \in \mathbb{N}$  there is an  $\epsilon \in \mathbb{R}$  such that the condition (3.17) holds for each  $\tilde{t} \in (n - \epsilon, n]$ ,  $n \in \mathbb{N}$ . We can perform the explicit calculation of  $\mathcal{T}_t h$ ,

$$\begin{aligned} \mathcal{T}_t(h) &= \mathcal{T}_t(\lim_{t \rightarrow T} U_t^2(F)) \\ &= \mathcal{T}_t(\lim_{t \rightarrow T} \kappa(g, t) h(u, p) + (1 - \kappa(g, t)) g_F(u, p)) \\ &= \lim_{t \rightarrow T} \mathcal{T}_t(\kappa(g, t) h(u, p) + (1 - \kappa(g, t)) g_F(u, p)) \\ &= \lim_{t \rightarrow T} (\mathcal{T}_t(\kappa(g, t)) \mathcal{T}_t(h)(u, p) + \mathcal{T}_t(1 - \kappa(g, t)) \mathcal{T}_t(g_F)(u, p)) \\ &= \lim_{t \rightarrow T} (\mathcal{T}_t(\kappa(g, t)) h(u, p) + \mathcal{T}_t(1 - \kappa(g, t)) g_F(u, p)). \end{aligned}$$

Therefore, a sufficient condition for the last line coincides with  $h$  is that

$$(3.18) \quad \mathcal{T}_t(\kappa(g, t)) = \kappa(g, t).$$

Furthermore, it is clear that  $\mathcal{T}_t$  is idempotent,

$$(3.19) \quad (\mathcal{T}_t)^2 = Id, \quad \forall t \in I \subset \mathbb{R}.$$

**3.4.  $U_t$  is an information loss dynamics.** If  $(TM, F, \mathcal{C})$  is a Cartan-Randers space that evolves towards the final *averaged structure*  $(TM, h)$  by the  $U_t$  operation, for each value of  $t$  there is an element  $(TM, F_t)$  of  $\mathcal{F}_{CR}(TM)$ . Applying the time inversion operation  $\mathcal{T}_t$  to  $F_t$  and taking into account that the function  $\kappa(g, t)$  is invariant under  $\mathcal{T}_t$ , one obtains the corresponding Hamiltonian of a DCRS at the instant  $(t, \tau)$ ,

$$\begin{aligned} H_t(u, p) &= \frac{1}{2} F_t^2(u, p) - \frac{1}{2} F_t^2(\mathcal{T}_t(u), \mathcal{T}_t^*(p)) \\ &= \frac{1}{2} (1 - \kappa(g, t)) g^{ij}(u, p) p_i p_j + \frac{1}{2} \kappa(g, t) \langle g^{ij} \rangle p_i p_j \\ &\quad - \frac{1}{2} (1 - \kappa(g, t)) g^{ij}(\mathcal{T}_t(u), \mathcal{T}_t^*(p)) p_i p_j - \frac{1}{2} \kappa(g, t) \langle g^{ij} \rangle p_i p_j \\ &= \frac{1}{2} (1 - \kappa(g, t)) g^{ij}(u, p) p_i p_j - \frac{1}{2} (1 - \kappa(g, t)) g^{ij}(\mathcal{T}_t(u), \mathcal{T}_t^*(p)) p_i p_j \\ &= 2(1 - \kappa(g, t)) \sum_{k=1}^{8N} \beta^k(u) p_k. \end{aligned}$$

Therefore, the Hamiltonian associated with a DCRS at the instant  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$  is

$$(3.20) \quad H_t(u, p) = 2(1 - \kappa(g, t)) \sum_{k=1}^{8N} \beta^k(u) p_k.$$

Then by the limit conditions (3.14), the *equilibrium Hamiltonian* (or completely averaged Hamiltonian) of a DCRS is identically zero,

$$(3.21) \quad \lim_{t \rightarrow T} H_t(u, p) = 0.$$

**3.5. The  $U_\tau$  dynamics.** The kinematics and dynamics of the sub-quantum degrees of freedom respect to the external time parameter  $\tau \in \mathbb{R}$  is described as follows. First, the speed components  $\beta_x$  and the acceleration components  $\beta_y$  of the vector field  $\beta \in \Gamma TTM$  are defined with the aid of the time inversion operator  $\mathcal{T}_t$ ,

$$(3.22) \quad \beta_x := \frac{1}{2}(\beta - \mathcal{T}_t(\beta)), \quad \beta_y := \frac{1}{2}(\beta + \mathcal{T}_t(\beta)).$$

For DCRS, the non-degeneracy of the fundamental tensor  $g$  of the underlying Cartan-Randers space is ensured if the vector field  $\beta$  is constrained by the condition (3.9). Such condition implies:

- (1) The velocity vector of the sub-quantum atoms is normalized by the condition  $\eta_4^k(\dot{\xi}_k, \dot{\xi}_k) \leq v_{\max}^2$ ,
- (2) There is a bound for the maximal acceleration  $\eta_4^k(\ddot{\xi}_k, \ddot{\xi}_k) \leq a_{\max}^2$ ,
- (3) The acceleration and speed are bounded.

For the Newtonian speed and accelerations are bounded in the sense that

$$(3.23) \quad \eta_4(\beta_{kx}, \beta_{kx}) \leq c, \quad \eta_4(\beta_{ky}, \beta_{ky}) \leq a_{\max}, \quad k = 1, \dots, N.$$

A geometric theory with maximal speed and maximal acceleration can be found in [17]. This theory uses a background Lorentzian metric structure. The conditions (3.23) are 4-covariant conditions. There are coordinate systems where these can be read as implying that the Newtonian speed and acceleration are bounded.

The classical *Hamiltonian function* of a DCRS is defined by the *partial average*

$$(3.24) \quad H(u, p) := \left( \frac{1}{2} F^2(u, p) - \frac{1}{2} F^2(\mathcal{T}_t(u), \mathcal{T}_t^*(p)) \right) = 2 \sum_{k=1}^{8N} \beta^k(u) p_k, \quad k = 1, \dots, 8N.$$

We remark that the motivation to have non-degeneracy for the underlying metric structure comes from the possibility to interpret the Hamiltonian (3.24) as the sum (or better, difference) of two pieces: one evolving to the future and second one to the past and that both corresponds to geodesic evolutions. If this interpretation must be correct, then the need of the condition (3.9) is justified. As we have seen, the consequences of this interpretation is that Cartan-Randers models are local Lorentz invariant and have maximal acceleration.

The Hamiltonian function (3.24) corresponds to a *time orientation average* of the Cartan-Randers function associated with a particular form of classical Hamiltonian. It defines the  $U_\tau$  dynamics of the sub-quantum degrees of freedom. It is the first stage towards the *total averaged Hamiltonian*. The Hamilton equations for  $H(u, p)$  are

$$(3.25) \quad \dot{u}^i = \frac{\partial H(u, p)}{\partial p_i} = 2 \beta^i(u), \quad \dot{p}_i = -\frac{\partial H(u, p)}{\partial u^i} = -2 \sum_{k=1}^{8N} \frac{\partial \beta^k(u)}{\partial u^i} p_k, \quad i, k = 1, \dots, 8N,$$

where the time derivatives are taken respect to the non-compact time parameter  $\tau$ . We have the following theorem,

**Theorem 3.6.** *For each dynamical system as given by the equations (3.25), there exists a Cartan-Randers system whose Hamiltonian function is (3.24).*

If in addition one requires that the *on-shell* conditions

$$(3.26) \quad \{\dot{x}^k = y^k, k = 1, \dots, 4N\}$$

hold, the Hamilton equation must be constrained. Thus the first equations of Hamilton imply

$$\beta_y^k(x, \dot{x}) = \dot{\beta}_x^k(x, \dot{x}), \quad k = 1, \dots, 8N$$

Note that in *Theorem 3.6* the metric  $\alpha$  is not fixed. Therefore, there are many Cartan-Randers systems for the same dynamical Hamiltonian system.

A relevant consistent property of the Hamiltonian dynamics is that the  $U_\tau$  dynamics is reversible,

**Proposition 3.7.** *The  $U_\tau$  dynamics is reversible,*

$$(3.27) \quad H(u, p) = H(\mathcal{T}_t(u), \mathcal{T}_t(p)).$$

*Proof.* It is direct from the definition of  $\mathcal{T}_t$  by equation (3.16), the property that  $\mathcal{T}_t$  is idempotent and the definition of the family of  $\beta$ -functions (3.22).  $\square$

#### 4. HILBERT SPACE FORMULATION OF DCRS

Every DCRS has associated a Hilbert space formulation. The algebra of functions  $\mathcal{F}_D(T^*TM)$  on  $T^*TM$  that we consider are the diagonal functions. The quantum operators that we will consider are obtained by a quantization of functions in  $\mathcal{F}_D(T^*TM)$ . However, since in DCRS the time parameters are associated with two-dimensional parameters, we need to specify the commutation relations at each fixed value of the pair  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . Following closely the procedure of standard quantization we adopt the following quantization rules defined as follows,

- The values of the position coordinates  $\{x_k^\mu, k = 1, \dots, N, \mu = 1, 2, 3, 4\}$  and the velocity coordinates  $\{y_k^\mu, k = 1, \dots, N, \mu = 1, 2, 3, 4\}$  of the fundamental degrees of freedom appear as eigenvalues of certain hermitian operators  $\{\hat{x}_k^\mu, \hat{y}_k^\mu, \mu = 1, 2, 3, 4; k = 1, \dots, N, \}$ ,

$$(4.1) \quad \begin{aligned} \hat{x}_k^\mu |x_l^\mu, y_l^\nu, \mu, \nu = 1, 2, 3, 4\rangle &= \sum_l \delta_{kl} x_l^\mu |x_l^\mu, y_l^\nu, \mu, \nu = 1, 2, 3, 4\rangle, \\ \hat{y}_k^\nu |x_l^\mu, y_l^\nu, \mu, \nu = 1, 2, 3, 4\rangle &= \sum_l \delta_{kl} y_l^\nu |x_l^\mu, y_l^\nu, \mu, \nu = 1, 2, 3, 4\rangle. \end{aligned}$$

where  $\delta_{kl}$  is Kronecker's delta function.

- There is a set of Hermitian operators  $\{\hat{p}_{\mu x_k}, \hat{p}_{\mu y_k}, k = 1, \dots, N, \mu = 1, 2, 3, 4\}$  that generates local diffeomorphism on  $TM$  along the integral curves of the local vector fields  $\{\frac{\partial}{\partial x_k^\mu} \frac{\partial}{\partial y_k^\nu}, \in \Gamma TTM \quad \mu, \nu = 1, 2, 3, 4; k = 1, \dots, N\}$ .
- The *canonical commutation relations* at each fixed 2-time  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$  hold good,

$$(4.2) \quad [\hat{x}_k^\mu, \hat{p}_{\nu x_l}] = i \hbar \delta_\nu^\mu \delta_{kl}, \quad [\hat{y}_k^\mu, \hat{p}_{\nu y_l}] = i \hbar \delta_\nu^\mu \delta_{kl}$$

Any other canonical commutator is equal to zero.

The collection  $\{|x_l^\mu, y_l^\nu, \mu, \nu = 1, 2, 3, 4\rangle, l = 1, \dots, N\}$  generates a complex vector space  $\mathcal{H}_{Planck}$  that can be promoted to a complex pre-Hilbert space by defining the following scalar product,

$$(4.3) \quad \langle x_l^\mu, y_l^\nu | x_k^\rho, y_k^\lambda \rangle = \delta_l^k \delta_\rho^\mu \delta_\lambda^\nu$$

and extended by assuming bilinear property of the product for arbitrary linear combinations with finite norm. Following this quantization scheme, the Hamiltonian (3.20) can be quantized obtaining the Hermitian operator,

$$(4.4) \quad \hat{H}(t, \tau, \hat{u}, \hat{p}) = \frac{1}{2} \sum_{k=1}^{8N} (\beta^k(t, \tau, \hat{u}) \hat{p}_k + \hat{p}_k \beta^k(t, \tau, \hat{u})).$$

When the quantum conditions (4.2) are applied to the Heisenberg's equation of the Hamiltonian (4.4), the equations (3.25) for the eigenvalues  $\{x_k^\mu, y_k^\mu \mid \mu = 1, 2, 3, 4; k = 1, \dots, N\}$  in the eigenbasis (4.1) are obtained. Furthermore, in order to have consistency between the Hilbert formalism and the geometric formalism, the constraints

$$(4.5) \quad \hat{y}_k^\mu = \frac{d\hat{x}_k^\mu}{d\tau}, \quad \forall \mu = 1, 2, 3, 4; \quad k = 1, \dots, N$$

need to be imposed. These constraints define trajectories *on-shell* sub-quantum degrees of freedom.

The commutation relations (4.2) are mathematically and physically consistent, despite the fact that in quantum mechanics the canonical coordinate operators  $\{\hat{X}^a, a = 2, 3, 4\}$  and the velocity operators  $\{\dot{\hat{X}}^a, a = 2, 3, 4\}$  do not commute for each quantum degree of freedom  $a$ . However, let us remark that the operators  $\{(\hat{x}_k^a, \hat{y}_k^a)\}_{k=1, a=2}^{N,4}$  do not coincide with the quantum operators  $\{\hat{X}^a, \dot{\hat{X}}^a, a = 2, 3, 4\}$ . The operators  $\{\hat{X}^a, \dot{\hat{X}}^a, a = 2, 3, 4\}$  have as spectrum the possible outcomes of measurements of position and velocity for each component of the cartesian macroscopic position and newtonian macroscopic velocity. The operators  $\{\hat{X}^a, \dot{\hat{X}}^a, a = 2, 3, 4\}$  should emerge in DCRS together with the wave function for the quantum state, that appears as phenomenological description of the ergodic regime when one considers the projection  $(t, \tau) \mapsto \tau$ . The general relation between the emergent operators  $\{\hat{X}^a, \dot{\hat{X}}^a, a = 2, 3, 4\}$  and the entire family of operators  $\{(\hat{x}_k^a, \hat{p}_k^a)\}_{k=1, a=2}^{N,4}$  is under current investigation.

The quantum states that are obtained from DCRS are generically non-localized in both position and speed (or canonical momentum) and both  $\hat{X}^a$  and  $\dot{\hat{X}}^a$  have generally non-zero dispersion. Therefore, the quantum states must hold a representation of a non-commutative algebra,

$$(4.6) \quad [\hat{X}_a^\mu, \hat{X}_b^\nu] = A^{\mu\nu} \delta_{ab}, \quad [\dot{\hat{X}}_a^\mu, \dot{\hat{X}}_b^\nu] = B^{\mu\nu} \delta_{ab}, \quad [\hat{X}_a^\mu, \dot{\hat{X}}_b^\nu] = C^{\mu\nu} \delta_{ab},$$

with  $\mu, \nu = 1, 2, 3, 4, a, b = 1, \dots, N$ . The requirement of local invariance under the Lorentz group of this algebra implies that the spacetime must be a quantum spacetime compatible with an universal maximal acceleration and a universal maximal speed.

The *symmetric quantization* of the Hamiltonian (3.20) gives the operator

$$(4.7) \quad \hat{H}_t(\tau, u, \hat{p}) := \frac{1}{2} (1 - \kappa(g, t, \tau)) \left( \sum_{k=1}^N (\beta^k(\tau, \hat{u}) \hat{p}_k + \hat{p}_k \beta^k(\tau, \hat{u})) \right),$$

with  $(t, \tau) \in I \times \mathbb{R}$ ,  $k = 1, \dots, 8N$ .  $\hat{H}_t(\tau, u, p)$  is uniquely defined, Hermitian and is determined by elements from the classical version of a DCRS.

Although it will be defined later on, let us assume first the existence of a Hilbert space  $\mathcal{H}_{Planck}$  whose elements  $|\psi\rangle \in \mathcal{H}_{Planck}$  are expanded by fundamental states associated with Cartan-Randers system. Such fundamental systems describe individual quantum systems as it could be an electron, for example. Let us introduce the quantum dynamics operator  $U_\tau$  by the expression

$$(4.8) \quad U_{\tau_0}^\tau |\psi\rangle(\tau_0) := |\psi\rangle(\tau_0) - i \delta\tau \hat{H} |\psi\rangle(\tau_0), \quad \delta\tau = \tau - \tau_0.$$

In this expression  $\delta\tau$  is a very short interval of time measured by a macroscopic external observer. However, since it is very small compared with any measurement of interval of time performed by macroscopic observer, we can take the approximation that  $|\delta\tau|$  can be arbitrarily small. Therefore, it follows from the definition of the classical Hamiltonian as generator of the dynamics of the system that the unitary

operator  $U_{\tau_0}^\tau$  is such that

$$(4.9) \quad U_{\tau_0}^\tau |\psi\rangle(\tau_0) = |\psi\rangle(\tau).$$

Therefore, one has

**Theorem 4.1.** *The Schrödinger equation holds*

$$(4.10) \quad i \frac{d}{d\tau} |\psi\rangle(\tau) = \hat{H} |\psi\rangle(\tau).$$

*Proof.* It is direct from the relation (4.9) and (4.1).  $\square$

Equation (4.10) is the Schrödinger's equation for degrees of freedoms whose dynamics happens at the Planck scale. In order to promote a similar equation for quantum states we need first to introduce the Hilbert space structure (approximately) corresponding to quantum systems and then extend the equation (4.10) to the quantum Hilbert space. The construction of the quantum Hilbert space is presented in the next section.

## 5. QUANTUM MECHANICAL STRUCTURES FROM DCRS

**5.1. Construction of local quantum phases.** Given a Cartan-Randers space, the construction of a symmetric metric structure as in (2.1) from the Randers structure data (that define an irreversible metric) is defined as follows. If  $M_4$  is endowed with a Lorentzian metric  $\eta_4$  with signature  $(1, -1, -1, -1)$  and with a timelike vector field  $W \in \Gamma TM_4$ , there is associated a Riemannian metric on  $M_4$  defined by the expression

$$(5.1) \quad \bar{\eta}_4(U, V) = \eta_4(U, V) - 2 \frac{\eta_4(U, W)\eta_4(V, W)}{\eta_4(W, W)}, \quad U, V \in \Gamma TM.$$

The distance function  $d_W : M_4 \times M_4 \rightarrow \mathbb{R}$  is associated with the Riemannian norm of  $\bar{\eta}_4$ . By the embeddings  $\varphi_k : M_4^k \rightarrow M_4$ , each manifold  $M_4^k$  is diffeomorphic to the manifold of observable macroscopic events  $M_4$ . Furthermore, the manifolds  $M_4^k$  and  $M_4$  are endowed with Lorentzian metrics that are isometric to the Lorentzian metric for each  $k \in \{1, \dots, N\}$ . Then one has the embeddings of the worldlines of the *sub-quantum molecules*  $\{1, \dots, N\}$  from  $M_4^k$  in  $M_4$   $\varphi_k(\xi_k) = \hat{\xi}_k \hookrightarrow M_4$ ,

$$\hat{\xi}(k) : \{t\} \times M_4^k \rightarrow M_4, \quad (t, \tau) \mapsto (t, \tau, \vec{\xi}(k)),$$

for each fixed value of the parameter  $t \in I$ .

If we assume as valid the axiom of choice, the following function can be defined: given  $x \in M_4$  and  $\xi(k)$ , the *distance function* between  $x$  and  $\hat{\xi}(k)$  is given by the expression

$$(5.2) \quad d_4(x, \hat{\xi}(k)) := \inf \{d_W(x, \tilde{x}), \tilde{x} \in \hat{\xi}(k)\}.$$

Then we can consider  $\min\{d_4(x, \hat{\xi}(k)), k = 1, \dots, N\}$ . This depend upon  $t \in I$ . Let us assume that such distance is realized for the fundamental molecule denoted by the integer  $\bar{k}$  at the instant  $t \in I$ . Given a curve  $z : \mathbb{R} \rightarrow M_4$ , one can consider the following definition,

**Definition 5.1.** *Let  $W \in \Gamma TM$  be a non-zero, timelike, unitary vector. Then the phase  $\theta \in \mathbb{R}$  is defined by*

$$(5.3) \quad \cosh(\theta)(x, W, t) := \eta_4(W, \dot{\hat{\xi}}(\bar{k})) \frac{1}{\sqrt{\eta_4(\dot{\hat{\xi}}(\bar{k}), \dot{\hat{\xi}}(\bar{k}))}}.$$



We call this function the *almost local phase*, since although it depends on the internal parameter  $t$ , it is defined on each  $u \in TM_4$ .  $\theta(x, W, t)$  depends upon the choice of the vector field  $W$ , that apart from being timelike and unitary, is otherwise arbitrary. Different choices of  $W$  imply different choices of almost-local phases (5.3) at each point  $x \in M_4$ .

At each point  $x \in M_4$  we define the *average phase* by the relation

$$(5.4) \quad \bar{\theta}(x) = \frac{\int_I dt \theta(x, t)}{\int_I dt}.$$

The phase (5.4) is independent of smooth, positive reparameterizations  $\mathbb{R} \rightarrow \mathbb{R}$ ,  $t \mapsto s(t)$ .

**5.2. The bundle associated with local quantum phases.** Let us consider the product

$$\mathcal{P} := \widetilde{M}_4 \times \mathbb{R}.$$

An element in  $\mathcal{P}$  is of the form  $(x, \theta)$ .  $\mathcal{P}$  is topologized with the product topology. Then we have the following

**Proposition 5.2.** *Let  $\mathcal{P}$  and  $\widetilde{M}_4$  be as before with the standard topologies. Then the canonical projection  $\pi_{\mathcal{P}} : \mathcal{P} \rightarrow \widetilde{M}_4$  is continuous and onto.*

Therefore,  $(\mathcal{P}, \widetilde{M}_4, \mathbb{R}, \pi_{\mathcal{P}})$  is a trivial fibre bundle such that

- The fibers  $\pi_{\mathcal{P}}^{-1}(\xi)$  are homomorphic to the abelian group  $\mathbb{R}$ ,
- The base space is the four manifold  $\widetilde{M}_4$ .

There is an action of the Lorentz group  $O(1, 3)$  on  $\mathcal{P}$  defined as follows. To each Lorentz transformation  $O \in O(1, 3)$ , the action is defined by the relation

$$(5.5) \quad \rho : \mathcal{P} \times O(1, 3) \rightarrow \mathcal{P}, \quad (x, \bar{\theta}, O) \mapsto (x, \rho_O(\bar{\theta})),$$

such that if  $\bar{\theta}$  are determined by the unitary time-like vector field  $V$ , then the phases  $\rho_O(\bar{\theta})$  are determined by the vector field  $O^{-1}(V) \in \Sigma$ . This action is a right, smooth action of  $O(1, 3)$  on  $\mathcal{P}$ .

**5.3. Definition of wave function in DCRS.** Let us consider a point  $x \in M_4$ . The number of worldlines  $\hat{\xi} : I \rightarrow M_4$ , for fixed  $\tau$ , such that they pass by  $x$  close to  $x$  is the density of lines  $n(x)$ . The density of lines is normalized by the condition

$$\int_{M_4} n(x) d\text{vol}_{\bar{\eta}_4} = N.$$

**Definition 5.3.** *A fundamental wave function  $\psi$  is a smooth section of  $\mathcal{P}$  defined pointwise by the expression*

$$(5.6) \quad \psi(x) = \frac{n(x)}{\sqrt{N}} \exp(i \bar{\theta}(x)).$$

A fundamental state is described by a fundamental wave function (5.6). The physical fundamental densities are normalized,

$$\int_{M_4} d\text{vol}_{\bar{\eta}_4} |\psi|^2 = 1.$$

If we consider a combination of two fundamental wave functions

$$\psi(x)_{A,B} = \frac{n_{A,B}(x)}{\sqrt{N}} \exp(i \bar{\theta}_{A,B}(x)),$$

the module square is

$$(5.7) \quad |\psi_A + \psi_B|^2 = \frac{n_A^2(x)}{N_A} + \frac{n_B^2(x)}{N_B} + 2 \frac{n_A(x)}{\sqrt{N_A}} \frac{n_B(x)}{\sqrt{N_B}} \cos(\bar{\theta}_A(x) - \bar{\theta}_B(x))$$

**5.4. Hilbert space associated with a DCRS.** The complex, linear hull generated by finite combinations of the fundamental states  $\psi$  is a complex vector space  $\mathcal{H}$ . Thus an element  $\Psi \in \mathcal{H}$  is of the form

$$(5.8) \quad \Psi = \sum_{k=1}^N \lambda_k \psi_k$$

where  $\{\psi_k\}$  are a collection of fundamental wave functions (5.6).

There is a  $L^2$ -norm defined in  $\mathcal{H}$ ,

$$(5.9) \quad \|\psi_A + \lambda\psi_B\|_2^2 = \int_{M_4} d\text{vol}_{\bar{\eta}_4} |\psi_A + \lambda\psi_B|^2$$

for each combination  $\psi_A + \lambda\psi_B$  fundamental states of the form (5.6) with  $\lambda \in \mathbb{C}$ .

**Proposition 5.4.** *The space  $\mathcal{H}$  with the norm  $\|\cdot\|_2$  is a complex Hilbert space.*

*Proof.* The set of functions  $L(M_4)$  of smooth square integrable functions on the four manifold  $M_4$  is a pre-Hilbert space. Moreover, since the elements of  $\mathcal{H}$  are finite combinations as in (5.8), it is clear that any Cauchy sequence  $\{\lambda_{kn}\}_{k,n=1}^{N,+\infty}$  converges in  $\mathbb{R}^N$ .  $\square$

Taking into account that if  $\hat{H} = 0$  in  $\mathcal{H}_{Planck}$  implies  $\hat{H} = 0$  in  $\mathcal{H}$ , we have that the quantum version of the diffeomorphism invariant condition (3.21) can be stated as the following

**Proposition 5.5.** *For any physical state  $|\psi\rangle \in \mathcal{H}$  the constraint*

$$(5.10) \quad \left( \lim_{t \rightarrow T} \hat{H}_t(u, p) \right) |\psi\rangle = 0$$

*holds good.*

This constraint is repeated periodically (with a period  $T$  depending on the quantum system) and constitute a version of de Wheeler-de Witt equation. The main difference with the standard version of the Wheeler-de Witt equation is that it is usually written in terms of operators of macroscopic operators  $\{\hat{X}, \hat{P}\}$ , while (5.10) is given in terms of microscopic operators. Due to the complexity of the systems, to relate both formulations is a difficult task, that we do not pursue in this work.

## 6. CONCENTRATION OF MEASURE AND MEASUREMENT PROCESSES IN DCRS

**6.1. Concentration of measure.** In this *section* we discuss the relation between the theory of functional analysis known by *concentration of measure* [23, 32, 42] and emergent quantum mechanics. The concentration of measure is a general property of high dimensional normed spaces with a metric and measure structures, which is generally stated as

*In a metric measure space, real 1-Lipschitz functions of many variables are almost constant almost everywhere.*

The precise meaning of *almost constant* and *almost everywhere* make use of the metric and measure properties of the space. The concentration of measure can be considered as a broad generalization of the *isoperimetric inequality* in spheres [23] or as a generalization of the *Central Limit Theorem* in probability theory [42].

**Example 6.1.** In a general measure metric space  $(M, \eta_P, d)$ , the concentration function  $\alpha(\eta_P) : \mathbb{R} \rightarrow \mathbb{R}$  is defined as the minimal real number  $\alpha(\eta_P, \rho)$  such that

$$(6.1) \quad \eta_P(|f - M_f| > \rho) \leq 2\alpha(\eta_P, \rho),$$

for any real function  $f : M \rightarrow \mathbb{R}$ . Therefore, the probability that the function  $f$  differs from the median  $M_f$  for more than the given value  $\rho \in \mathbb{R}$ , using the measure of probability  $\mu$ , is bounded by the concentration function  $\alpha(P, \rho)$ . A typical example of concentration of measure is provided by the concentration of measure in spheres  $\mathbb{S}^N \subset \mathbb{R}^{N+1}$ . For spheres, the concentration function is of the form

$$(6.2) \quad \alpha(P_M, \rho) \leq c \exp\left(-\frac{(N-1)\rho^2}{2\rho_P^2}\right),$$

where  $\rho_P$  has the physical interpretation of being the minimal value attainable for the physical observable described by the function  $f : \mathbb{S}^N \rightarrow M$ .

For Lipschitz functions on space of dimension  $N$  there are similar *Chernov's type bounds*. This generic feature, which is a consequence of the regularity conditions is useful when one is dealing with higher order dimensions.

**6.2. An application to dynamical reduction of quantum states.** Let us consider an application of the concentration of measure in  $R^q$  with  $q \gg 1$ .  $\eta_P$  is a measure and  $f : \mathbb{R}^q \rightarrow \mathbb{R}$  a real 1-Lipschitz function. Then there is concentration given by (see for instance [42], pg. 8)

$$(6.3) \quad \eta_P(|f - M_f| > \rho) \leq \frac{1}{2} \exp\left(-\frac{\rho^2}{2\rho_P^2(f)}\right),$$

where we have adapted the example from [42] to a Gaussian distribution  $\eta_P$  with mean  $M_f$  and standard deviation  $\rho_P(f)$ .  $\rho_P(f)$  has the physical interpretation of being the minimal resolution attainable when measuring the observable associated with the classical function  $f : \mathbb{R}^q \rightarrow \mathbb{R}$ .

We apply this *example* of concentration to the function  $f \in \mathcal{F}_D(T^*TM)$ . We assume first that for  $f$  there is a maximal resolution  $\rho_p(f)$  in their possible measurement outcome values. In the 1-Lipschitz dynamical regime of the dynamics operator  $U_t$ , the function  $f$  must be constant almost everywhere, since  $f$  is 1-Lipschitz in  $(u, p)$  and  $t$ . Moreover, for macroscopic observable effects one expect a relation of the type

$$(6.4) \quad \frac{\rho^2}{\rho_P^2(\tilde{f})} \simeq N^2, \quad 1 \ll N,$$

which defines relative scales in terms of observable quantities. The two scales in deterministic Cartan-Randers systems are on a relation  $N : 1 \sim 10^2$  or higher. This ratio is motivated by the ratio between the GUT scale and the Planck scale. The GUT scale can be thought as the higher scale where the fundamental particle dynamics is well described by a quantum gauge field theory. However, this possibility is only an assumption and it could be that  $N$  is bigger. However, our following arguments does not depend on the exact value of the high energy quantum regime except that must be below the Planck or fundamental classical scale.

Let us restrict our considerations to the case when the quantum system corresponds to a fundamental particle which quantum fields appear in the Standard Model. The natural number  $N$  provides a measure of the *complexity* of the fundamental system compared with the complexity of the associated quantum system. The degree of complexity of a quantum state is of order 1, since there is one quantum particle involved compared with the degree of complexity of a DCRS, which is of order  $N$ . This order of complexity 1 is of the same order than the dimension

of the model spacetime manifold  $M_4$ , the number of spin degrees of freedom, and other quantum numbers associated with the quantum mechanical description of a fundamental quantum particle. Let us consider the case  $TM \cong \mathbb{R}^{8N}$ . By the relation (6.4), the concentration relation (6.3) when is applied to the function  $f$  in the 1-Lipschitz dominated regime of  $U_t$  reduces to

$$(6.5) \quad \eta_P(|f - M_f| > \rho_P(f)) \leq \frac{1}{2} \exp(-32 N^2).$$

$M_f$  depends upon the initial conditions  $(u_1(0), \dots, u_{8N}(0), p_1(0), \dots, p_{8N}(0))$ . Since by assumption  $\rho_P(f)$  is small compared with  $|f - M_f|$  and  $N \gg 1$ , there is concentration of measure around the mean  $M_f$ . Thus, if a measurement of an observable associated with  $f$  is performed, the result  $M_f$  will be with probability very close to 1. This phenomenon of concentration of measure corresponds with a process of reduction of the quantum state. Note that the interaction responsible for this concentration is purely classical, since it appears in the limit  $t \rightarrow T = 1$  only, in contrast with other interactions that happen also during the non-local ergodic regime of the  $U_t$  interaction.

The concentration of measure that produces the reduction of the quantum state are determined by the initial conditions of the system. Indeed, these reduction of the quantum state not only happen when the system is being *measured*, but it is an *spontaneous process, happening all the time*. However, we do not have at the moment which are the possible initial conditions of the systems. However, this is not in favour or against super-determinism in DCRS.

There is an important consequence from this process. The parameter  $\tau$  has been considered to be a continuous, real parameter. However, the observation that the  $U_t$  is periodic on  $t$  and that any measurement is done in a regime of equilibrium given by (5.10) implies that the time parameter  $\tau$  must indeed be integer. This is because any physical clock is linked to a series of observable processes, that can happen only on the equilibrium domains. Therefore, we arrive to the conclusion that any notion of external time, linked with a physical quantum or macroscopic process, is emergent and quantized. As a consequence, the continuous description of the clock must be regarded as an idealization. Thus, we will use such idealization in this work and post-pone for further analysis the most exact description in terms of discrete external time parameters.

**6.3. Description of the two slit quantum experiment in DCRS.** Let us consider the quantum two slit experiment as prototype of quantum interference phenomena [15]. In a simplified description of the two slit experiment (slit A and slit B), the theoretical setting is two-dimensional, with the  $x$ -direction being the direction of propagation of the beam of quantum particles and the  $z$ -axis the vertical direction of orientation for the slits and the measurement screen. Furthermore, the states are pure quantum states, representing individual quantum particles each time that an experiment is done. The experiment is done many times with different quantum particles, with the constraint that the macroscopic initial momenta before reaching the slits is the same. One assumes that other conditions on the experiment (like the value of the gravitational field, interactions with the ambient, spin states) either are the same for all the experiments or not affect the experiment at all.

If there are two slits  $A$  and  $B$ , each time an individual experiment is done, we can ask by which of the slits the individual system pass through. Because the process of instantaneous collapse described before, this question is legitim. Then one says that after passing the slit, the system is generated by the slit  $A$  if it passed through the slit  $A$  at some time  $\tau_A$  and analogously, the system has been generated by the slit  $B$  if the system passed through the slit  $B$  at some instant  $\tau_B$ . Let the state with source at the slit  $A$  be  $\psi_A$ . The slit  $B$  generates another state that we denote

by  $\psi_B$ . It is clear that the states  $\psi_A$  and  $\psi_B$  are fundamental states of the type described by equation (5.6). Now, the evolution after passing the slit is linear and given by a Laplacian operator, characterized by the boundary conditions. Then the effective state just after the system passes the slits is described by a vector  $\psi \in \mathcal{H}$  of the form

$$\psi(x, z) = C (\psi_A(x, z) + \psi_B(x, z)),$$

with  $C$  a normalization real constant such that  $\|\psi\|_{M_4} = 1$ . This state has an epistemic nature, but is not ontological, that is, there is not described directly in terms of sub-quantum degrees of freedom.

Since the slits  $A$  and  $B$  are different, the states  $\psi_A$  and  $\psi_B$  produce different screen patterns, after a long time exposition. Therefore, it is natural to adopt the hypothesis that  $\psi_A \neq \psi_B$  in the vicinity of the detection screen. Furthermore, symmetry considerations imply that  $\psi_A = \psi_B$  in the central region of the screen, generating a maximum for  $\|\psi_A + \psi_B\|^2$ . Out of the central axis one expects that  $|\psi_A| \neq |\psi_B|$  or  $\arg(\psi_A) \neq \arg(\psi_B)$  or both conditions. If there is a relative phase between  $\psi_A$  and  $\psi_B$ , one will find an interference pattern depending on the geometric arrangement of the experiment and according to the equation (5.7). As long as we move out from the axis  $z = 0$ , at least a difference on phase will appear. Otherwise, the phases  $(\theta_A, \theta_B)$  should be constant along the  $z$ -axis, which is not possible. Moreover, one also expects to find a variability on the norms  $\|\psi_A\|$  and  $\|\psi_B\|$  along the  $z$ -axis.

There is a physical interpretation for existence of nodes or regions where  $|\psi|^2$  is zero in the two state interference. In a measurement of a quantum system, we have two time flows,

- The flow in the same direction than the flow of any external macroscopic observer.
- The flow in time of the quantum system due to the internal dynamics.

If these two flows coincide, there is a constructive contribution to interferences. However, if the two flows do not coincide, that can happen in some occasions because the evolution of the sub-quantum degrees of freedom in the  $t$ -time can happen in both directions respect to the  $\tau$ -time orientation, there is a *lack of presence* of the quantum state. It is in this sense that the observer observes a node. The node appears even if the modules  $|\psi_A|^2$  and  $|\psi_B|^2$  are different from zero.

## 7. HAMILTONIAN BOUNDEDNESS AND EMERGENCE OF THE WEAK EQUIVALENCE PRINCIPLE AND GRAVITATIONAL INTERACTION

**7.1. Boundedness of the Hamiltonian.** A fundamental problem in deterministic schemes for emergent quantum mechanics is that the Hamiltonian is linear in the momenta operators and therefore, it is not trivial to have a mechanism stabilizing the dynamical systems against decays towards arbitrary large negative energy states. One needs to impose extra conditions or introduce a mechanism to stabilize the vacuum. We show that a natural mechanism exists in DCRS theory.

The basic idea is to decompose  $\hat{H}$  in a 1-Lipschitz component and a non-Lipschitz component,

$$(7.1) \quad \hat{H}_t(\hat{u}, \hat{p}) = \hat{H}_{matter,t}(\hat{u}, \hat{p}) + \hat{H}_{Lipschitz,t}(\hat{u}, \hat{p}).$$

Clearly, this type of decomposition is not unique and even it is not evident that it exists. However, one can obtain a characterization of the decomposition under several additional assumptions. In particular, let us assume that the classical Hamiltonian function (3.24) is  $\mathcal{C}^2$  - *smooth* in the variables  $(x, y, p_x, p_y)$ . In this case, we can prove the following

**Lemma 7.1.** *Let  $H$  be a  $\mathcal{C}^2$ -smooth Hamiltonian function. Then there exists a compact domain  $K \subset TM$  where  $H|_K$  is 1-Lipschitz.*

*Proof.* By Taylor's expansion, one has formally (the notation does not introduce indices),

$$\begin{aligned} H(u, p) &= H_0 + \sum_{k=1}^{8N} \frac{\partial H}{\partial u^k} \big|_{(\xi, \chi)} (u^k - \xi^k) + \sum_{k=1}^{8N} \frac{\partial H}{\partial p_k} \big|_{(\xi, \chi)} (p_k - \chi_k) \\ &\quad + \sum_{k=1}^{8N} R_k (u^k - \xi^k)^2 + \sum_{k=1}^{8N} Q_k (p_k - \chi_k)^2 \\ &= H_0 + \sum_{k=1}^{8N} \frac{\partial H}{\partial u^k} \big|_{(\xi, \chi)} (u^k - \xi^k) + \sum_{k=1}^{8N} \beta^k(\chi) (p_k - \chi_k) \\ &\quad + \sum_{k=1}^{8N} R_k (u^k - \xi^k)^2 + \sum_{k=1}^{8N} Q_k (p_k - \chi_k)^2 \end{aligned}$$

The differenc for the values at two different points is

$$\begin{aligned} |H(u(1), p(1)) - H(u(2), p(2))| &= \left| \sum_{k=1}^{8N} \frac{\partial H}{\partial u^k} \big|_{(\xi, \chi)} (u^k(1) - \xi^k) \right. \\ &\quad + \sum_{k=1}^{8N} \beta^k(\chi) (p_k(1) - \chi_k) + \sum_{k=1}^{8N} R_k(1) (u^k(1) - \xi^k)^2 + \sum_{k=1}^{8N} Q^k(1) (p_k(1) - \xi_k)^2 \\ &\quad - \sum_{k=1}^{8N} \frac{\partial H}{\partial u^k} \big|_{(\xi, \chi)} (u^k(2) - \xi^k) - \sum_{k=1}^{8N} 2 \beta^k(\chi) (p_k(2) - \chi_k) \\ &\quad \left. - \sum_{k=1}^{8N} R_k(2) (u^k(2) - \xi^k)^2 - \sum_{k=1}^{8N} Q^k(2) (p_k(2) - \chi_k)^2 \right| \\ &\leq \left| \sum_{k=1}^{8N} \frac{\partial H}{\partial u^k} \big|_{(\xi, \chi)} (u^k(1) - u^k(2)) \right| + \left| \sum_{k=1}^{8N} \beta^k(\chi) (p_k(1) - p_k(2)) \right| \\ &\quad + \left| \sum_{k=1}^{8N} R_k(1) (u^k(1) - \xi^k)^2 - R_k(2) (u^k(2) - \xi^k)^2 \right| \\ &\quad + \left| \sum_{k=1}^{8N} Q^k(1) (p_k(1) - \chi_k)^2 - Q^k(2) (p_k(2) - \chi_k)^2 \right|. \end{aligned}$$

For each compact set  $\tilde{K} \subset T^*TM$ , there are two constants  $C_R(\tilde{K}) > 0$  and  $C_Q(\tilde{K}) > 0$  such that  $|R_k(1)|, |R_k(2)| < C_R(\tilde{K})$  and  $|Q_k(1)|, |Q_k(2)| < C_Q(\tilde{K})$ , for each  $k = 1, \dots, 8N$  and with

$$\lim_{1 \rightarrow 2} C_Q(\tilde{K}) = 0, \quad \lim_{1 \rightarrow 2} C_R(\tilde{K}) = 0,$$

This is a consequence of Taylor's theorem. Therefore, the constants  $C_Q(\tilde{K})$  and  $C_R(\tilde{K})$  can be taken finite on  $\tilde{K}$ . Furthermore, one has that without losing generality

$$(7.2) \quad C_R(\tilde{K})(u^k(1) - u^k(2)) \leq 1/2, \quad C_Q(\tilde{K})(p_k(1) - p_k(2)) \leq 1/2.$$

Therefore, the last two lines are rewritten as

$$\begin{aligned} \left| \sum_{k=1}^{8N} R_k(1) (u^k(1) - \xi^k)^2 - R_k(2) (u^k(2) - \xi^k)^2 \right| &= C_R(\tilde{K}) \left| \sum_{k=1}^{8N} (u^k(1) - u^k(2))^2 \right| \\ \left| \sum_{k=1}^{8N} Q^k(1) (p_k(1) - \chi_k)^2 - Q^k(2) (p_k(2) - \chi_k)^2 \right| &= C_Q(\tilde{K}) \left| \sum_{k=1}^{8N} (p_k(1) - p_k(2))^2 \right|. \end{aligned}$$

Now, let us consider a compact set  $K \subset T^*TM$  such that for each  $(\xi, \chi) \in K$ .

$$\left| \frac{\partial H}{\partial u^k} \right|_{(\xi, \chi)} \leq C_U$$

for some given constant  $C_U$ . Also, on  $K$  it must hold that

$$\begin{aligned} &C_R(K) \left| \sum_{k=1}^{8N} (u^k(1) - u^k(2))^2 \right| + C_Q(K) \left| \sum_{k=1}^{8N} (p_k(1) - p_k(2))^2 \right| \\ &\leq 1/2 \sum_{k=1}^{8N} |(u^k(1) - u^k(2))| + 1/2 \sum_{k=1}^{8N} |(p_k(1) - p_k(2))|. \end{aligned}$$

That there is such compact set  $K$  follows from the fact that it is defined by close relations. Then we have that

$$\begin{aligned} |H(u(1), p(1)) - H(u(2), p(2))| &\leq C_U \sum_{k=1}^{8N} |(u^k(1) - u^k(2))| + \sum_{k=1}^{8N} |(p_k(1) - p_k(2))| \\ &+ 1/2 \sum_{k=1}^{8N} |(u^k(1) - u^k(2))| + 1/2 \sum_{k=1}^{8N} |(p_k(1) - p_k(2))|. \end{aligned}$$

This proves that  $H$  is Lipschitz with Lipschitz constant  $\max\{2, C_U\}$ , which is necessarily finite. Now we can redefine the Hamiltonian dividing by such a constant, which is equivalent to redefine the vector field  $\beta$  only, an operation that does not changes the equations of motion. Then we obtain a 1-Lipschitz Hamiltonian on  $K$  equivalent to the previous one.  $\square$

The compact domain  $K$  is not unique and non-empty. However, extensions from  $K$  to the whole phase space can be constructed as follows. Let us consider the projection on  $K$

$$(7.3) \quad \pi_K : TM \rightarrow K, (u, p) \mapsto (\bar{u}, \bar{p})$$

where  $(\bar{u}, \bar{p})$  is defined by the condition that the distance from  $(u, p)$  to  $K$  is achieved at  $(\bar{u}, \bar{p})$ . Then one defines the *radial decomposition* of  $H$  by

$$(7.4) \quad H(u, p) = R(d((u, p), (u_0, p_0))) H(\bar{u}, \bar{p}) + \delta \mathcal{H}(u, p).$$

The positive function  $R(d((u, p), (u_0, p_0)))$  is chosen such that decreases with the distance function  $d((u, p), (u_0, p_0))$  and is bounded in a way that the first piece of the Hamiltonian is still 1-Lipschitz. The second contribution is not Lipschitz, although it is associated with the matter Hamiltonian  $H_{matter}$ ,

$$(7.5) \quad H_{matter}(u, p) = \delta \mathcal{H}(u, p).$$

Therefore, we have the following

**Theorem 7.2.** *Every Hamiltonian (3.24) has a normalization such that the decompositions (7.1) hold good.*



Now we can read the Hamiltonian constraint (5.10) in the following way. From the  $U_t$  dynamics it follows that

$$\lim_{t \rightarrow T} (\hat{H}_{matter,t} + \hat{H}_{Lipschitz,t})|\psi\rangle = 0$$

However, each of the individual terms can be different from zero in the equilibrium regime or near the equilibrium regime

$$\lim_{t \rightarrow T} \hat{H}_{matter,t}|\psi\rangle \neq 0, \quad \lim_{t \rightarrow T} \hat{H}_{Lipschitz,t}|\psi\rangle \neq 0.$$

This implies that in order to have a stable equilibrium regime  $t \rightarrow T$ , in addition with the matter Hamiltonian (7.5), an additional interaction described by the Hamiltonian  $\hat{H}_{Lipschitz,t}$  is needed.

In presence of matter, if we assume that the 1-Lipschitz Hamiltonian has negative eigenvalues only, then the matter Hamiltonian (7.5) must be positive definite on physical states in the equilibrium and near the equilibrium domain. However, because the  $U_t$  evolution generates a geometric flow in the cotangent space  $T^*TM$  and induces an evolution for the Hamiltonian function, the  $t$ -dependence of the Hamiltonian can be parameterized by the expression (4.7). Because the geometric character of the Hamiltonian, constructed from the difference between two positive Hamiltonian functions at any instant  $(t, \tau)$ , the function  $\kappa(g, t, \tau)$  is in the interval  $[0, 1]$ . Therefore, the positiveness of the matter Hamiltonian is extended to all  $t \in [0, T]$ . This implies the positiveness of the energy level for the matter quantum Hamiltonian (4.4).

**7.2. Emergence of the weak equivalence principle.** We denote by  $X^\mu(\mathcal{S}_i)$ ,  $i \equiv \mathcal{S}, A, B$  the *macroscopic observable  $\mu$ -coordinate* associated to the system  $\mathcal{S}_i$ . This symbol represents the corresponding coordinates of a quantum system that a macroscopic observer can associate to the system. We assume a smooth dependence of  $X^\mu$  on  $(u^{k_1}, \dots, u^{k_N}, p^{k_1}, \dots, p^{k_N}, t, \tau)$ . Under the additional assumption that in the same coordinate system where the fundamental degrees of freedom have bounded maximal acceleration and speed there is also a constrain on the corresponding macroscopic observable quantities, in the equilibrium domain  $t \rightarrow T$  the functions  $X^\mu(t, \tau) \mapsto X^\mu(\tau)$  are 1-Lipschitz. In order to show this we consider the relations

$$(7.6) \quad \lim_{t \rightarrow T} \frac{\partial X^\mu(u, p, t, \tau)}{\partial t} = 0,$$

$$(7.7) \quad \frac{\partial X^\mu(u, p, t, \tau)}{\partial \tau} = \sum_{k=1}^{8N} \frac{\partial X^\mu}{\partial u_k^\rho} \frac{du_k^\rho}{d\tau} + \sum_{k=1}^{8N} \frac{\partial X^\mu}{\partial p_{\rho k}} \frac{dp_{\rho k}}{d\tau}.$$

If all the macroscopic observable coordinates speed coordinates are bounded (by the speed of light), then the derivatives  $\{\frac{\partial X^\mu}{\partial u_k^\rho}, \frac{\partial X^\mu}{\partial p_{\rho k}}, \mu = 1, 2, 3, 4; k = 1, \dots, N\}$  must be bounded as well. This fact implies that the coordinate functions  $\{X^\mu(\tau)\}_{\mu=0}^3$  have the Lipschitz property. By fixing conveniently the units and normalizing as we did in the proof of *Lemma 7.1*, the Lipschitz constant for the coordinate functions  $X^\mu$  are bounded by 1, which is equivalent to have a maximal acceleration and maximal speed for the observable macroscopic coordinates, in the reference frames where the speeds and accelerations  $\{\frac{du_k^\rho}{d\tau}\}_{k=1}^{8N}$  are bounded.

Let us assume that the subsystems  $A$  and  $B$  are coordinated in a given local coordinates on  $TM$  by

$$A \equiv (u_1(\tau), \dots, u_{N_A}(\tau), 0, \dots, 0) \quad \text{and} \quad B \equiv (0, \dots, 0, v_1(\tau), \dots, v_{N_B}(\tau)),$$

with  $N = N_A + N_B, N_A, N_B \gg 1$ . Locally, the full system  $\mathcal{S}$  can be represented as

$$\mathcal{S} \equiv (u_1(\tau), \dots, u_{N_A}(\tau), v_1(\tau), \dots, v_{N_B}(\tau)).$$

By the concentration property (6.3) for the  $U_t$  dynamics, the  $\tau$ -evolution of the coordinates  $X^\mu(\mathcal{S}(\tau))$ ,  $X^\mu(A(\tau))$  and  $X^\mu(B(\tau))$  that started with the same initial conditions will differ after the dynamics at time  $\tau$  such that the constraint

$$\eta_P(|X^\mu(\mathcal{S}_i(\tau)) - M^\mu(\tau)| > \rho)_{t \rightarrow T} \sim C_1 \exp(-C_2 \frac{\rho^2}{2L_p^2})$$

holds good. The constant  $C_2$  are of order 1 and is independent of the system  $A, B, \mathcal{S}$ . Note that there is no dependence on the  $t$ -parameter, since we are in the limit  $t \rightarrow T$ . The mean  $M^\mu(\tau)$  is determined by the expression

$$(7.8) \quad M^\mu(\tau) = \frac{1}{N} \lim_{t \rightarrow T} \sum_{k=1}^N \eta_k(t, \tau) x_k^\mu(t, \tau).$$

From this definition of the mean value  $M^\mu(\tau)$  it follows the theoretical meaning of the functions  $X^\mu(\mathcal{S}_i(\tau))$ . Because of the diffeomorphisms  $\varphi_k : M_4^k \rightarrow M_4$ , one can consider the world lines of the  $k$ -molecules on  $M_4$  constant value of  $t$ . Then  $X^\mu$  identifies the *total presence of sub-quantum matter* at a given point of the spacetime with the potential observability of quantum matter at such point. The exact nature of this relation is supposed to involve a rather intricate non-linear relations, with complexity increasing with  $N$ . We do not address this question in this work.

An interchange along the  $\tau$ -time of fundamental  $k$ -degrees of freedom with the environment can happen even at equilibrium. This can affect the motion of the center of mass  $M^\mu(\mathcal{S}_i(\tau))$  in a rather intricate way, because the derivatives

$$\frac{d}{d\tau} \eta_k(t, \tau), \quad k = 1, \dots, N.$$

depend upon the details of the system  $\mathcal{S}_i(\tau)$ . However, if in the equilibrium limit  $t \rightarrow T$  it holds that

$$(7.9) \quad \frac{d}{d\tau} \eta_k(t, \tau) = 0, \quad k = 1, \dots, N,$$

then the center of mass coordinates  $M^\mu(\mathcal{S}_i(\tau))$  follow a well defined ordinary differential equation,

$$(7.10) \quad \frac{d}{d\tau} M^\mu(\tau) = h^\mu(\tau),$$

where the functions  $h^\mu$  are in theory fixed by the equations of motion of the  $8N$ -degrees of freedom and by the properties of the measure  $\eta_k$  and whose solutions depend only on the initial conditions for  $M^\mu(\tau = 0)$ . Let us consider geometric configurations for the systems  $\mathcal{S}_i$ ,  $i = 1, 2, 3$  such that  $M^\mu(\tau)$  do not depend on the system  $A, B$  or  $\mathcal{S}$  at  $\tau = 0$ . Then the center of mass coordinates  $M^\mu(\tau)$  will not depend on the system  $\mathcal{S}_i$  as well. For example, if  $N = 1$  the dispersion of finding  $X^\mu$  respect to the value of the center of mass is  $M^\mu$  is of order  $\exp(-\frac{1}{2}(8+1)^2) \simeq \exp(-40)$ , while for  $N = 2$  it should be of order  $\exp(-128)$ . Since  $N \gg 1$ , a weak equivalence principle applied to the observable coordinates  $\{X^\mu\}_{\mu=1}^4$  emerges. It is exact up to an error of order  $\exp(-40N^2)$ .

Therefore, in the equilibrium regime, there is a strong concentration for the value the functions  $\{X^\mu(\tau)\}_{\mu=1}^4$  around  $\{M^\mu(\tau)\}_{\mu=1}^4$ . Note that this universality is up to fixing the initial conditions of the center of mass  $M^\mu$ , which is equivalent to fix the initial conditions  $\{x_k^\mu\}_{k=1}^{8N}$ . That is, it depends in the general case, of a set of un-controllable number of initial conditions. This does not rest relevance to the

main message: that for the concentration dynamics  $U_t$  in the equilibrium domain  $t \rightarrow T$ , for the same initial conditions, the center of mass coordinates  $M^\mu$  evolves in the same way, irrespective of the nature of the system.

**7.3. Emergence of the gravitational interaction.** If we collect all the characteristics of the 1-Lipschitz interaction  $U_t$  in the regime  $t \rightarrow T$  we have that,

- (1) It is described by a theory invariant under infinitesimal diffeomorphism transformations, since the constraint (3.21) holds good,
- (2) The weak equivalence principle for the center of mass functions  $M^\mu(\mathcal{S}(\tau))$  holds good,
- (3) There is a local maximal speed for the fundamental degrees of freedom of a DCRS and local Lorentz invariance holds,
- (4) It is a classical interaction,
- (5) It must be compatible with the existence of a maximal acceleration.
- (6) The associated quantum Hamiltonian operator has negative energy eigenvalues.

In view of the strong resemblance of these properties with the weak equivalence principle, one can make an strong assumption and postulate

*In the equilibrium domain  $\hat{H}|\psi\rangle = 0$ , the 1-Lipschitz dynamics associated with  $\hat{H}_{Lipshitz, t=T}$  is the gravitational interaction.*

That there is a domain where the gravitational interaction is concentration can be easily seen in the Newtonian gravity limit. Let us consider a Newtonian gravitational potential

$$V(\vec{x}) = -\frac{m M G}{r}, \vec{x} \in \mathbb{R}^3$$

and  $r = |\vec{x}|$  the distance to the origin in  $\mathbb{R}^3$  of  $\vec{x}$ . Then we have the relation

$$|V(\vec{x}_1) - V(\vec{x}_2)| = m M G \left| \frac{1}{r_1} - \frac{1}{r_2} \right| = m M G \left| \frac{r_1 - r_2}{r_1 r_2} \right|$$

Let  $m = M = m_e$  be of the size of the electron  $r_1 = r_2$  of the scale of the upper bound for the quantum radius of the electron  $10^{-22}m$ . Then there is a natural norm for the gravitational field,

$$|\vec{F}_G| = \frac{|V(\vec{x}_1) - V(\vec{x}_2)|}{|r_1 - r_2|} \sim m M G \frac{1}{r_1 r_2}.$$

This scale of force must be compared with the associated Planck's force  $F_P = \hbar c$ , where  $c$  is the speed of light in vacuum,

$$\frac{|\vec{F}_G|}{F_P} = \frac{m M G}{\hbar c} \frac{1}{r_1 r_2}.$$

For an electron this quotient is  $\frac{|\vec{F}_G|}{F_P} \approx 2.7 \times 10^{-2}$ . Thus, it is clear that the gravitational force is 1-Lipschitz at these scales. Similar arguments should applied to any massive quantum system.

## 8. DISCUSSION

**8.1. Cartan-Randers systems and other deterministic models for emergent quantum mechanics.** In Cartan-Randers systems time is a two-dimensional parameter  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . The parameter  $\tau$  is the external time used to describe the dynamics of a quantum state  $|\psi\rangle$  when interacting with the environment. This time parameter corresponds with the usual notion of *macroscopic time* as it appears

in quantum mechanics and quantum field theory. The internal time parameter  $t$  in DCRS describes the  $U_t$  internal dynamics of the Hamiltonian  $H_t(u, p)$  and appears in the DCRS formalism as the parameter of the homotopy (3.13) of Cartan-Randers dynamical systems. This notion of *two dimensional time* sharply contrast with the usual notion of time in other emergent deterministic approaches to quantum mechanics [2, 9, 10, 13, 24, 34]. In such approaches the notion of time is the usual as in quantum mechanics and field theory, except for the fact that in some of these approaches the time parameter  $\tau$  can be discrete.

It is this notion of two dimensional time parameter  $(t, \tau)$  which is beneath our interpretation of quantum mechanics as *geometric dynamics*: the sub-quantum degrees of freedom experience a doubly dynamics, one associated with the geometric flow  $U_t$  and the second one described by the canonically quantized of the Hamiltonian (3.24). The essence of our interpretation of the fundamental non-local description of the quantum mechanics is based in the following argument. In order to speak of the state of a system at an instant of time in the DCRS framework, we need to specify two parameters,  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . If only the parameter  $\tau$  is specified in a dynamical description of a physical system (as it is done in usual field theory and quantum mechanics) and the system is not in an localized state for the  $U_t$  dynamics, an intrinsic non-local behavior of the state associated with the ergodicity of the dynamics emerges. We believe that this is the origin of the non-locality description of an individual system in quantum mechanics.

In DCRS framework, the existence of the spontaneous collapse explains the measurement problem because the system is in a localized state prior to any measurement. In an localized state, all the possible classical observables have a definite value. This is applicable to spin measurements. Thus, we have that a picture resembling a Bohmian system, with well defined observable properties at each macroscopic instant  $\tau$  emerges from a deterministic system, whose time parameterization requires two time coordinates  $(t, \tau)$ . The idea that Bohmian mechanics is emergent in DCRS is therefore, very attractive and could support independent investigations [22]. However, it is need to investigate further if effectively, Bohmian mechanics emerges from Cartan-Randers systems. The main obstruction is the ontological character of the wave function, compared with the (in general) epistemological character ascribed in DCRS. In Cartan-Randers systems, only for fundamental states the wave function has an ontological nature, associated with *the density of presence of matter*.

One expects that the weak equivalence principle has an emergent interpretation from systems with large number of degrees of freedom. Such emergent interpretation is based on the same principle generating the macroscopically observable states, which is the concentration of measure phenomena. If our mechanism proposed for the emergence of WEP is correct, the weak equivalence principle will not be gradually violated in quantum physics: for all possible energies where the degrees of freedom of the Standard Model are presented, the WEP will be valid, with no observable violation of it before reaching the fundamental scale. However, it will be completely violate at the energy scale where the degrees of freedom correspond to sub-quantum molecules of a DCRS. Therefore, we should not expect violation of the WEP up to energies of the order of the fundamental scale.

Another consequence from DCRS is that the covariant accelerations vectors  $D_\xi \dot{\xi}$  are universally bounded for all physical degrees of freedom. If the fundamental scale is of order of the Planck scale, then the maximal acceleration associated should be of order  $a_{planck} = 10^{50} m/s^2$ . Maximal acceleration should have relevant implications for the absence of spacetime curvature singularities (see for instance

[11]). It is in this direction where we expect that an universal acceleration as the Planck acceleration  $a_{Planck}$  could have falsifiable predictions.

**8.2. Open problems and further developments.** Many important points need to be clarified in our proposal. Between the most relevant are to find a dynamics  $U_t$  full-filling all the properties that we have discovered. Several candidates are being investigated by the author, among them, models for the so called *Riemann dynamics*. It can be shown that Cartan-Randers models are relativistic versions of the so called  $xp$ -Hamiltonian [6, 7, 12, 41]. It is stimulating that the properties of DCRS are found in correlation with many properties found in the quantum mechanical approach to the Riemann Hypothesis and that they contain, not only the standard  $xp$ -Hamiltonian but also further exquisite Hamiltonians of  $xp$ -type. Thus, we are investigating the possibility that the Hilbert-Polya conjecture is realized, not at the atomic or quantum level [40], but at a more fundamental level of the physical reality. It is worthily to mention that the work of Sierra [41] investigates a  $xp$ -model with many features that the Riemann dynamics should have and that has an (ad hoc) maximal acceleration and maximal speed (the speed of light in vacuum). Therefore, it should not be difficult to re-write Sierra's model as a DCRS. Moreover, one needs to construct the Hamiltonian operator in terms of macroscopic operators. This task is difficult, since the systems that we are considering are rather involved. Passing from  $\hat{H}(\hat{u}, \hat{p})$  to  $\hat{H}(\hat{X}, \hat{P})$  requires to know the structure of the operators  $\{\hat{X}, \hat{P}\}$  in terms of the operators  $\{\hat{u}, \hat{p}\}$  or a formal argument to identify the quantum Hamiltonian. We think that the work of Hooft and others on quantum automaton models [29] should provide important insights.

We find rather unsatisfactory the introduction in our theory of several geometric structures as background. Instead, we believe that a more complete treatment of the theory will rely in very few fundamental elements and that the geometric structures will become either emergent or dynamical, specially the metric structures that we use. This must be the case, since the theory should contain gravity and being geometrically background independent. Thus, the part of giving a dynamics to the metric should provide a theory where quantum mechanics and general relativity marriage in a different way that quantization of gravity proposes [36, 37].

We have not discussed how to interpret quantum non-locality and quantum entanglement in detail and we did not consider a detailed treatment of Bell's inequalities and Kochen-Specker's theorem in the DCRS framework. Moreover, the origin of the Born's rule should be also addressed in DCRS. Furthermore, because the direct relation between the  $U_t$  dynamics and the quantum dynamics could be extremely complex, it is useful to provide phenomenological quantum models compatible with DCRS models. These are problems that we will address in future work.

There are some relations between Cartan-Randers theory and other theories and interpretations of quantum mechanics. For instance, some resemblance between our proposal and de Broglie-Bohm approach to quantum theory [8], since the observables of the system are defined prior to measurement, according to de Broglie-Bohm theory. Also, it seems natural to interpret the wave function as a driving field, although at this point we need to remark that in de Broglie-Bohm theory the pilot field is real, while in our theory it is epistemological, a convenient tool to deal with the notion of the *presence of matter*. Thus, we think that essentially our theory will not reproduce de Broglie-Bohm theory and instead it will provide a genuinely new interpretation of the quantum theory.

Cartan-Randers systems, should have however, a relation with the two-state formulation of quantum mechanics and protective measurements [3, 4, 43, 44]. This is related with the existence of two directions of evolution for the sub-quantum

atoms, that we use to describe the Hamiltonian (3.24). Thus, although this is rather similar to the idea of the two-vector formalism, it is formulated at the level of the Planck's scale.

Cartan-Randers theory should be related with emergent gravity of form developed in some related investigations [30, 31, 35]. In these theories gravity emerges in an *equilibrium regime* from a dynamics of quantum gravity at the Planck scale. This situation is partially similar to what happens in Cartan-Randers theory, where a gravitational interaction emerges only in the thermodynamical equilibrium regime  $t \rightarrow T$ , where the macroscopic properties of the system are defined and are measurable. Moreover, if classicality is seen as an equilibrium regime, it is natural to think that quantum mechanical behavior is related with the non-equilibrium regime. Then a relation of Cartan-Randers models with the thermodynamical approach vindicated in [38, 1] eventually can emerge. This is satisfactory from the point of view that we are considering a microscopic dynamics and the corresponding thermodynamics is associated with classicality, while non-equilibrium thermodynamics should be associated to an intermediate description of the real world as is quantum mechanics.

The theory of Cartan-Randers spaces have clear similarities with Hooft's approach to the foundations of quantum mechanics. There are formal similarity in several conceptual issues [24, 25]. In this sense, we believe that our models are a particular realization of many features that Hooft's theory advocates. Among them, DCRS are deterministic, local dynamical systems, there is an information loss dynamics, remarkably, they include gravitational interaction, although in a very novel way, there is Hamiltonian which is bounded from below by a natural mechanism and there is a distinction between fundamental (quantum) states and their quantum superpositions, which in DCRS are considered phenomenological devices only. In DCRS there is a natural mechanism that could provide an explanation for non-locality in quantum mechanics. Bell's inequalities are applicable, but not necessarily to the observables that are currently being tested in experiments. They can be applied to the degrees of freedom at the Planck's scale. The applicability for quantum observable operators is lost when equivalence classes are considered. All these features are present in both Hooft's proposal and in Cartan-Randers theory.

Despite these common features between both approaches to emergent quantum mechanics, there are also important differences. The first one is on the notion of time parameter, being in our proposal two-dimensional. This notion is useful in a potential explanation of quantum non-locality from the projection two-dimensional to one-dimensional time in the mathematical description. However, in the explanation that we propose for non-locality it is not necessary to introduce super determinism in the sense of [25, 26]. Our theory is compatible with super determinism, but this is not a necessary element of the theory and indeed, it is not used to explain non-locality and quantum correlations.

The second fundamental difference with Hooft's theory is on the notion of collapse of the wave function, being absent in Hooft proposal [26] but that happens spontaneously in Cartan-Randers systems. Indeed, in DCRS the phenomenon contrast with the theory of collapse models of quantum mechanics [19, 20, 21, 5]. While in collapse models the collapse is induced by the effect of a macroscopic measurement body, in DCRS the collapse is spontaneous and is not necessarily induced by any external body. In DCRS the collapse (or concentration regime as we called) is caused by the internal dynamics of the quantum system. Moreover, it is not a complete collapse as it happens in collapse models. After a concentration regime there is an expanding regime, a phenomenon completely absent in collapse models



of quantum mechanics. Moreover, the scale and frequency where the collapse happens in DCRS is different than in collapse models: in DCRS such scales are tied with the GUT scale or very close to the Planck's scale.

Our models are relativistic, in the sense that there is a maximal speed for the fundamental degrees of freedom. This also seems to be a feature compatible with Hooft's approach. Indeed, several relativistic examples of relativistic dynamical systems as cellular automaton quantum mechanics [27, 28] exist, although further study is required to see if they also have maximal acceleration. However, in DCRS there is in addition a maximal value for acceleration, that contrast in principle with the models proposed by Hooft.

**Acknowledgements.** This work was supported by PNPd-CAPES n. 2265/2011, Brazil. I would like to thank Prof. J. M. Isidro for several interesting comments.

#### REFERENCES

- [1] D. Acosta, P. Fernández de Córdoba, J. M. Isidro, J. L. G. Santander, *Emergent quantum mechanics as a classical, irreversible thermodynamics*, International Journal of Geometric Methods in Modern Physics, **10**, No. 04 1350007 (2013).
- [2] S. L. Adler, *Statistical Dynamics of Global Unitary Invariant Matrix Models as Pre-Quantum Mechanics*, hep-th/0206120; S. L. Adler, *Probability in Orthodox Quantum Mechanics: Probability as a Postulate Versus Probability as an Emergent Phenomenon*, quant-ph/0004077.
- [3] Y. Aharonov, P. G. Bergmann and J. L. Lebowitz, *Time symmetry in the quantum process of measurement*, Physical Review B **134**, no. 6, pp. 14101416 (1964).
- [4] Y. Aharonov and L. Vaidman, *The Two-State Vector Formalism of Quantum Mechanics an Updated Review*, In: Juan Gonzalo Muga, Rafael Sala Mayato, Iigo Egusquiza (eds.): Time in Quantum Mechanics, Volume 1, Lecture Notes in Physics 734, pp. 399-447, 2nd ed., Springer (2008).
- [5] A. Bassi and G.C. Ghirardi, *Dynamical Reduction Models*, Physics Reports **379**, 257 (2003).
- [6] M. V. Berry, *The Bakerian Lecture, 1987: Quantum Chaology*, Proc. R. Soc. Lond. A **413**, 183-198 (1987).
- [7] M. V. Berry and J. P. Keating,  *$H = xp$  and the Riemann zeros*, in *Supersymmetry and Trace Formulae: Chaos and Disorder*, ed. J. P. Keating, D. E. Khemelnitskii, L. V. Lerner, Kowler 1999; M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, SIAM REVIEW **41** (2), 236 (1999).
- [8] D. Bohm, *A suggested Interpretation of the Quantum Theory in Terms of Hidden variables. I*, Phys. Rev. **85**, 166-179 (1952); *A suggested Interpretation of the Quantum Theory in Terms of Hidden variables. II*, Phys. Rev. **85**, 180-193 (1952); D. Bohm, *Wholeness and the Implicate Order*, London: Routledge, (1982).
- [9] M. Blasone, P. Jizba and G. Vitiello, *Dissipation, Emergent Quantization and Quantum Fluctuations*, in *Decoherence and Entropy in Complex Systems, Selected Lectures from DICE 2002*, H.-T. Elze (ed.), Lecture Notes in Physics 633, Springer, Berlin (2004).
- [10] M. Blasone, P. Jizba and F. Scardigli, *Can Quantum Mechanics be an Emergent Phenomenon?*, J. Phys. Conf. Ser. **174** (2009) 012034, arXiv:0901.3907[quant-ph].
- [11] E. R. Caianiello, *Is there a Maximal Acceleration*, Lett. Nuovo Cimento **32**, 65 (1981); E. R. Caianiello, *Quantum and Other Physics as Systems Theory*, La Rivista del Nuovo Cimento, Vol. **15**, Nr 4 (1992).
- [12] A. Connes, *Trace formula in noncommutative geometry and the zeros of the Riemann zeta function*, Selecta Mathematica (New Series) **5**, 29 (1999), math.NT/9811068.
- [13] H.T. Elze *Quantum mechanics emerging from "timeless" classical dynamics*, Trends in General Relativity and Quantum Cosmology, ed. C.V. Benton (Nova Science Publ., Hauppauge, NY, 79-101 (2006); H.T. Elze, *The Attractor and the Quantum States*, Int. J. Qu. Info. **7**, 83 (2009); H.T. Elze, *Symmetry Aspects in Emergent Quantum Mechanics*, J. Phys. Conf. Ser. **171**, 012034 (2009).
- [14] H. T. Elze, *Action principle for cellular automata and the linearity of quantum mechanics*, arXiv:1312.1615 [quant-ph].
- [15] R. Feynman, R. B. Leighton and M. Sands, *The Feynman lectures on Physics*, Ad. Wesley (1964).
- [16] R. Gallego Torromé, *A Finslerian version of 't Hooft Deterministic Quantum Models*, J. Math. Phys. **47**, 072101 (2006).



- [17] R. Gallego Torromé, *On a covariant version of Caianiello's Model*, Gen.Rel.Grav. **39**, 1833-1845 (2007).
- [18] R. Gallego Torromé, *On the convex invariance in Finsler geometry*, Symmetry: Culture and Science **23**, Number 2, 133-140, 2012.
- [19] G.C. Ghirardi, A. Rimini and T. Weber, *Unified dynamics for microscopic and macroscopic systems*, Physical Review, D **34**, 470 (1986).
- [20] G.C. Ghirardi, P. Pearle and A. Rimini, *Markov-processes in Hilbert-space and continuous spontaneous localization of systems of identical particles*, Phys. Rev. A **42**, 78 (1990).
- [21] G. Ghirardi, R. Grassi and A. Rimini, *Continuous-spontaneous-reduction model involving gravity*, Phys. Rev. A **42**, 1057-1064.
- [22] G. Gröessing, *Emergence of Quantum Mechanics from a Sub-Quantum Statistical Mechanics*, Int. J. Mod. Phys. B, **28**, 1450179 (2014).
- [23] M. Gromov, *Riemannian structures for Riemannian and non-Riemannian spaces*, Birkhäuser (1999).
- [24] G. 't Hooft, *Determinism and Dissipation in Quantum gravity*, hep-th/0003005; G. 't Hooft, *The mathematical basis for deterministic quantum mechanics*, ITP-UU-06/14, SPIN-06/12, quant-ph/0604008; G. 't Hooft, *Emergent Quantum Mechanics and Emergent Symmetries*, 13th International Symposium on Particles, Strings, and Cosmology-PASCOS 2007. AIP Conference Proceedings **957**, pp. 154-163 (2007).
- [25] G. 't Hooft, *Entangled quantum states in a local deterministic theory*, arXiv:0908.3408 [quant-ph].
- [26] G. 't Hooft, *How a wave function can collapse without violating Schrödinger's equation and how to understand Born's rule*, arXiv:1112.1811v3 [quant-ph].
- [27] G. 't Hooft, *Duality between a deterministic cellular automaton and a bosonic quantum field theory in 1+1 dimensions*, TP-UU-12/18; SPIN-12/16, arXiv:1205.4107.
- [28] G. 't Hooft, *Discreteness and Determinism in Superstrings*, ITP-UU-12/25; SPIN-12/23, arXiv:1207.3612.
- [29] G. 't Hooft, *The Cellular Automaton Interpretation of Quantum Mechanics: a View on the Quantum Nature of our Universe, Compulsory or Impossible?*, arXiv:1405.1548 v3.
- [30] B. L. Hu, *Emergent/quantum gravity: macro/micro structures of spacetime*, J. Phys.: Conf. Ser. **174** 012015 (2009).
- [31] T. Jacobson, *Thermodynamics of space-time: The Einstein equation of state*, Phys. Rev. Lett. **75**, 1260 (1995).
- [32] V. D. Milman and G. Schechtman, *Asymptotic theory of Finite Dimensional normed spaces*, Lecture notes in Mathematics 1200, Springer (2001).
- [33] R. Miron, D. Hrimiuc, H. Shimada and V. Sabau, *The Geometry of Hamilton and Lagrange Spaces*, Fundamental Theories in Physics 118, Kluwer (2001).
- [34] E. Nelson, *Derivation of the Schrödinger equation from Newtonian Mechanics*, Phys. Rev. **150** 1079-1083 (1964); *Dynamical theories of brownian motion*, Princeton University Press (1967); *Quantum Fluctuations*, Princeton University Press (1985).
- [35] T. Padmanabhan, *Gravity from space-time thermodynamics*, Astrophys. Space Sci. **285**, 407 (2003).
- [36] R. Penrose, *On Gravity's Role in Quantum State Reduction*, General Relativity and Gravitation, Vol. **8**, No. 5, 581 (1996); R. Penrose, *The Road to Reality*, Vintage, London (2005).
- [37] R. Penrose, *On the gravitization of quantum mechanics 1: quantum state reduction*, Found. Phys. **44**, 557-575 (2014).
- [38] N. Olah, *Einsteins Trojanisches Pferd: eine Thermodynamische Deutung der Quantentheorie*, Springer, Wien (2011).
- [39] G. Randers, *On an Asymmetrical Metric in the Four-Space of General Relativity*, Phys. Rev. **59**, 195-199 (1941).
- [40] G. Sierra and P. K. Townsend *Landau levels and Riemann zeroes*, Phys. Rev. Lett. **101**, 110201 (2008).
- [41] G. Sierra, *The Riemann zeros as energy levels of a Dirac fermion in a potential built from the prime numbers in Rindler spacetime*, J. Phys. A: Math. Theor. **47** 325204 (2014).
- [42] M. Talagrand, *A new look at independence*, Ann. Probab. **24**, Number 1, 1-34 (1996).
- [43] L. Vaidman, *The two-state vector Formalism*, arXiv:0706.1347 [quant-ph].
- [44] S. Watanabe, *Symmetry of physical laws. Part III. Prediction and retrodiction*, Reviews of Modern Physics **27** 2, 179 (1955).