### A THEORY OF EMERGENT QUANTUM MECHANICS

Ricardo Gallego Torromé

Departamento de Matemática, Universidade Federal de São Carlos, Rodovia Washington Luís, km 235, SP, Brazil

ABSTRACT. A particular class of non-regular Hamiltonian dynamical systems is considered as deterministic mathematical models for the description of physical systems at a more fundamental level description that the one provided by quantum mechanics. After introducing the geometric description for such physical systems, that is around our notion of Hamilton-Randers space, and after reformulating the theory using Hilbert space theory, a Hilbert space structure is constructed and associated with the space of wave functions of quantum mechanics from the underlying Hamilton-Randers model. In the general framework of this theory we can prove the emergence of the Born's rule from ergodic considerations and the emergence of the Heisenberg's picture of quantum dynamics. A geometric description for a spontaneous reduction of the quantum states based on the concentration of measure phenomena as it appears in metric geometry is described. The conditions for the existence of concentration are also fundamental to show the existence of stable vacua states for the matter Hamiltonian. Another consequence of the concentration is the emergence of a weak equivalence principle. This fact, together with the existence in the theory of a maximal speed and the property of diffeomorphism invariance of the interaction driving the reduction of the quantum state, this suggests that the reduction of the quantum state is driven by a gravitational type interaction. Moreover, since such interaction appears only in the dynamical domain when localization happens, it must be associated with a classical interaction. Then it is shown that the operators associated with observables follow the Heisenberg's equation. We discuss the double slit experiment in the context of the models that we propose, as well as an argument in favour of an emergent interpretation of the macroscopic time.

**Keywords**. Emergent quantum mechanics, emergence of the weak equivalence principle, diffeomorphism invariance, generalized Hamiltonian spaces, Randers spaces, maximal acceleration, concentration of measure, ergodicity, break of ergodicity, natural spontaneous collapse, quantum measurement, quantum nonlocality, superluminal motion.

#### Contents

1. Introduction	2
1.1. Motivation	2
1.2. Structure of the work	5
2. Assumptions for deterministic dynamical models at the Planck scale	6
2.1. Remarks on the assumptions	7
2.2. Maximal proper acceleration	9
3. Hamilton-Randers systems	11
3.1. Geometric background	11

## EMERGENT QUANTUM MECHANICS

3.2. Notion of Hamilton-Randers space	14
3.3. Interpretation of the parameter $t$ and the semi-period $T$	16
3.4. Properties of the $U_t$ flow	16
3.5. t-time inversion operation	18
3.6. The $U_t$ flow is information loss	18
3.7. Notion of external time parameter $\tau$	19
3.8. The $U_{\tau}$ -dynamics	21
3.9. $U_t$ -flow and redefinition of the $\tilde{\tau}$ -time parameter	22
3.10. Observers and metric structures	23
4. Hilbert space formulation of Hamilton-Randers systems	24
4.1. Hilbert formulation of classical systems applied to HR-systems	24
4.2. Quantum Hamiltonian associated to a HR-system	26
4.3. Heisenberg dynamics of the ontological states is the classical dynamics	26
4.4. Sub-quantum operators and quantum operators	27
5. Quantum mechanical elements from Hamilton-Randers systems	27
5.1. Quantum Hilbert space from HR-systems	27
5.2. Representations of the sub-quantum degrees of freedom	29
5.3. Emergence of the Born rule in the Heisenberg representation of the	
dynamics	29
5.4. Emergence of the $\tau$ -time diffeomorphism invariant constraint	30
6. Concentration of measure and measurement processes in Hamilton-	
Randers systems	31
6.1. Concentration of measure	31
6.2. Spontaneous reduction of the quantum state as concentration of	
measure phenomena	33
6.3. Notion of natural spontaneous reduction	35
7. Hamiltonian bound and emergence of the weak equivalence principle	
and gravitational interaction	36
7.1. Bound of the matter Hamiltonian	36
7.2. Emergence of a weak equivalence principle	39
7.3. An heuristic argument in favour of the emergent origin of the	
gravitational interaction	41
7.4. The Heisenberg dynamics of quantum observables	43
8. Conceptual issues in quantum mechanics from the point of view of	
Hamilton-Randers theory	44
8.1. The two slit quantum experiment and its interpretation in Hamilton-	
Randers Theory	45
8.2. On the non-local quantum correlations	46
8.3. On the emergence and ontological nature of the $\tau$ -time	49
9. Concluding remarks	50
9.1. Relation with others theories of emergent quantum mechanics	50
9.2. Relation of Hamilton-Randers theory and de Broglie-Bohm theory	51
9.3. Open problems in Hamilton-Randers theory	51
References	53

# 1. INTRODUCTION

1.1. Motivation. Current theories in physics assume the existence of a geometric spacetime where *happen*. Indeed, with the advent of general relativity spacetime becomes a dynamical entity, which is affected by the presence and dynamics of matter and it is not only a geometric back-ground structure

 $\mathbf{2}$ 

where other processes happen. This is in sharp contrast with the quantum mechanical description of reality, where the back-ground spacetime arena is fixed. Being general relativity and quantum mechanics theories aimed to be of universal applicability, it is to expect that both theories clash on its predictions at some domain of experience. It is also commonly believe that a quantum mechanical theory of gravity will resolve the singularities of general relativity. Several fields of research have emerged as programs to solve the incompatibility problem. Among them string theory, loop quantum gravity and many others quantum theories of gravity. Quantum gravity phenomenology is a new feel that explores general implications of the quantization of gravity. All these attempts, despite based in very interesting ideas, are at least unconcluded. Indeed, it is possible that to find a theory merging general relativity and quantum mechanics or suitable modifications of them in a unique scheme is the most fundamental problem in physics.

Even if only as an approximated description of a more accurate theory, to find a consistent spacetime representation for the quantum measurement processes, the non-local quantum correlation phenomena and a geometric understanding of quantum entanglement appears as a remarkable difficult task. In the case of the quantum measurement processes this is not surprising, since the standard *collapse postulate* of quantum mechanics involves the instantaneous reduction of the wave packet in each quantum measurement, a postulate that assumes processes clearly against the spirit of the special and general theories of the relativity. Other alternative descriptions of the measurement processes have been found not strictly compatible with a geometric picture of physical arena [37]. Also rather puzzling is the appearance of non-local phenomena. In particular, as it currently stands the quantum formalism, the explanation of quantum correlations in terms of direct action implies supraluminal communication or a fundamental non-locality [21].

The formal unification of quantum mechanics and special relativity achieved by relativistic quantum field theory, apart from the serious problems of mathematical consistency and rigourous formulation of the theory itself, the description of quantum measurement processes is excluded in quantum field theory. Moreover, the principles of general relativity are absent in formulations of quantum field theory.

Furthermore, because the difficulties in finding a consistent spacetime picture of the quantum mechanical non-local phenomena and assuming that fundamental physical systems are represented in a geometric and relativistic form, it is reasonable to doubt that the current quantum theory is the ultimate theoretical framework for the description of physics. From such perspective, it is clear that the problem of the spacetime representation by a macroscopic observer of quantum entanglement phenomena and quantum measurement processes is related with the logical and mathematical structure of the quantum theory and general relativity theory and with the ontological interpretation of the quantum physical processes. This suggests certain urgency to investigate alternative ways of understanding quantum mechanical phenomenology and gravity.

Partially justified by the above considerations, a general class of theories for physics beneath quantum mechanics generically known by *emergent*  quantum mechanics have been proposed in the literature. These theories share the common assumption that the quantum formalism should arise from an underlying, more fundamental theory. We cannot make full justice here to the whole contribution to this research area, but let us mention as examples of emergent quantum frameworks the theories have been developed in [2, 10, 15, 18, 23, 26, 32, 36, 41, 44]. Thermodynamical arguments in favour of an emergent origin of quantum mechanics have been provided by several authors [1, 17].

An specific sub-class of theories for emergent quantum mechanics assume that the degrees of freedom at the Planck scale are deterministic and local [10, 15, 18, 26]. However, a fundamental difficulty in such approaches is that the associated Hamiltonian operators, being linear in the momentum operators, are not bounded from below.

Therefore, in order to ensure the existence of stable vacuum states, a mechanism is need to stabilize the vacuum. One of the mechanism proposed in the literature involves dissipative dynamics at the fundamental Planck scale [10, 15, 26]. What is the origin of the dissipation of information? It has been argued that the gravitational interaction plays an essential role as the origin of the information loss dynamics and must be present at the level of the fundamental Planck scale. However, gravity could also be a classical and emergent phenomenon, and therefore, absent at the Planck scale energy where it is supposed that the dynamics of the microscopic fundamental degrees of freedom takes place. If this is the case, it is not natural to appeal from the beginning to gravity as the origin of the dissipation of information.

Another fundamental difficulty associated with deterministic quantum models for physics at the Planck scale is the relation between the degrees of freedom at the fundamental scale and the degrees of freedom at the quantum scales, that includes the degrees of freedom of the standard model of particles or unification quantum field theories but also, atomic and hadronic scales, for instance. Examples of field models that can be described as deterministic models have been discussed in the literature. In particular, it has been shown that free (1 + 1)-bosonic quantum field models and several string models can be interpreted as deterministic quantum models [30]. Also, it was proved that the massless non-interacting 4-dimensional Dirac field can be identified with a deterministic system [27]. Such examples show that to describe the dynamics of non-trivial quantum systems as deterministic dynamical systems is at least feasible and a extended theory of emergent quantum mechanics has been constructed based on cellular automaton models [16, 30].

The present paper supersedes the previous work by the author on deterministic systems as models for quantum mechanics [18]. In addition to further analysis and improvement of our previous proposal, several new developments are presented here. It is shown how fundamental elements of the quantum mechanical formalism are derived from the formalism of a specific type of dynamical models. Moreover, an interaction with formal similarities with classical gravity emerges in Hamilton-Randers systems in combination with a general principle of metric geometry known as concentration of measure arises. This interaction causes the natural spontaneous collapse. In such theory the collapse of the quantum state is not necessarily induced by the measurement device.

Although there are several serious constraints for local, deterministic and causal view for theories beneath quantum mechanics, we think that such theories can be logically consistent, although deviations from quantum mechanics should be expected. Further research on clarify such deviations is postponed for future work.

1.2. Structure of the work. This work is organized in three parts. The first part is contained in section 2. In this part, there is an heuristic introduction of the fundamental requirements that are used in our theory, as well as the introduction of the idea of maximal acceleration. The second part starts from section 3 and expands until section 5. It describes the fundamental mathematical structure of Hamilton-Randers dynamical systems and the reconstruction of the fundamental notions of the mathematical formalism of quantum mechanics from Hamilton-Randers systems. Section 3 provides a short exposition of generalized Hamilton space of Randers type. After time symmetrization, we show the relation of such Hamilton-Randers geometry with a specific type of dynamical systems that we call Hamilton-Randers dynamical systems (in short, HR-systems). We describe the two dynamics  $U_t$  and  $U_{\tau}$  associated with HR-systems. This is of fundamental relevance for our theory, since our interpretation of the quantum phenomena relies on the correct interpretation of the mathematics of HR-systems. Special care has been taken to explain the notion of two dimensional time associated to the two dimensional parameter  $(t, \tau)$ . The existence of maximal acceleration and speed in each HR-system is shown. In Section 4, a formulation of HR-systems by means of Hilbert space theory is described. Section 5 is dedicated to the theoretical construction of the quantum local phase from the original degrees of freedom of the underlying HR-systems. The transition from the description in terms of discrete degrees of freedom associated with HR-systems to continuous degrees of freedom associated with quantum wave functions is considered. Furthermore, a constructive approach to the quantum wave function, that admits a natural probabilistic interpretation and the associated quantum Hilbert space from the underlying HR-system are described. A derivation of the Born rule in the Heisenberg picture of the dynamics is provided.

The third part of the paper presents mainly the application of fundamental notions of metric geometry to the dynamics of HR-systems. In particular, in section 6 we apply the concentration of measure to explain the natural spontaneous reduction of the wave function. In section 7 it is shown the existence of an equilibrium domain where the dynamics is 1-Lipschitz. This property is used to discuss a natural mechanism to bound from below the quantum Hamiltonian operator for matter in HR-systems. We also discuss how a weak equivalence principle emerges in Hamilton-Randers theory by applying concentration of measure in the 1-Lipschitz regime of the dynamics. As a consequence we argue that classical gravity can be interpreted as an emergent phenomena. It is also argued that the dynamics of observables is determined by the Heisenberg equation for the matter Hamiltonian. An

#### EMERGENT QUANTUM MECHANICS

heuristic discussion of the quantum double slit interference theory is presented in *section 8*, where we also analyze the consequences for the notion of external time as an emergent concept from the dynamics of HR-systems. Finally, the relation of our theory with other emergent quantum mechanics frameworks and several open problems of our approach are briefly discussed in *section 9*.

# 2. Assumptions for deterministic dynamical models at the Planck scale

We assume a topological structure on the configuration space of the models. In particular, we consider the following type of structures,

**Definition 2.1.** Let **T** be a set. An asymmetric topological metric is a function  $\rho : \mathbf{T} \times \mathbf{T} \to \mathbb{R}$  such that

- $\varrho(u,v) \ge 0$ , for each  $u,v \in \mathbf{T}$ .
- $\varrho(u,v) = 0$  iff  $u = v \in \mathbf{T}$
- $\varrho(u_a, u_c) \leq \varrho(u_a, u_b) + \varrho(u_b, u_c)$ , for each  $u_a, u_b, u_c \in \mathbf{T}$ .

The main difference of an asymmetric metric with an usual metric function is that the symmetry condition has been dropped out. The notion of asymmetric metric is useful for encapsulate formally the concept of nonreversible evolution for dynamical systems. In particular, the dynamics of the systems that we investigate is associated to an asymmetric metric structure of a special type.

The following assumptions are useful in the search of the properties of dynamical systems at the Planck scale. They reflect an scenario which is significatively different than the standard scenarios, namely, quantum gravity theories. Rather than a set of formal axioms, the following assumptions must be considered as formal requirements that help to fix the structure of the dynamical systems with the characteristics that we consider fundamental, that is, causal, deterministic and local systems and that follow an irreversible dynamics. The assumptions are tried to be divided in three categories, attending the level of description that they need to be incorporated.

#### • Topological and measure assumptions:

- A.1. There is a discrete topological space model  $M_4$  which is the arena where macroscopic observers can locate macroscopic phenomena, characterized as events.
- A.2. There is a topological configuration space  $\mathcal{M}$  which is endowed with an asymmetric metric structure (2.1) and with a probability measure  $\mu_P$ .
- Ontological assumptions:
  - A.3. The sub-quantum degrees of freedom are identical degrees of freedom composed by two *sub-quantum atoms*. Since the sub-quantum degrees of freedom are composed, we call them *sub-quantum molecules*.
  - A.4. There is a natural minimal coordinate scale  $L_{min}$ , which is of the order of the Planck length. It is the universal minimal coordinate difference for events measured in any instantaneous inertial co-moving frame with a given sub-quantum molecule.

6

#### • Dynamical assumptions:

- A.5. The fundamental dynamics, that is, the dynamics of the sub-quantum molecules, is described by deterministic, finite difference equations, where the time parameter  $\tau$  is discrete. In each sub-quantum molecule, one of such sub-quantum atoms evolves toward the future, while the companion evolves towards the past in the external time  $\tau$ .
- A.6. The following *locality condition* holds: given a system S corresponding to a collection of sub-quantum molecules, there is a smallest neighborhood  $\mathcal{U}$  with  $S \subset \mathcal{U}$  such that for any  $\widetilde{\mathcal{U}} \supset \mathcal{U}$  the dynamical effect of any action of  $\mathcal{U}$  and  $\widetilde{\mathcal{U}}$  on S are the same.
- A.7. The following *causality condition* holds: there are local coordinate systems where there is a maximal, universal bound for the coordinate speed of the sub-quantum atoms and sub-quantum molecules. We assume that the maximal speed is the speed of light in vacuum at each given direction.
- A.8. The fundamental dynamics at the Planck scale is irreversible, that is, there is a function  $\Omega : \mathcal{M} \to \mathcal{M} \to \mathbb{R}$  such that for a possible evolution from  $A \in \mathcal{M}$  to  $B \in \mathcal{M}$  and from B to A, the function  $\Omega(A, B) \neq \Omega(B, A)$ .
- A.9. The fundamental dynamics is sensitive to initial conditions (it is a chaotic dynamics), with the appearance of unstable attractors in the long time evolution.

In this work, a category of dynamical systems (Hamilton-Randers systems) that fulfill the above assumptions and the application to reconstruct several aspects of the quantum mechanical formalism are investigated. Indeed, we show that Hamilton-Randers systems are indeed suitable candidates for deterministic models for describing systems at a fundamental scale (we assume that corresponds to the Planck scale), since from the dynamics of the Hamilton-Randers systems one can recover aspects of the quantum mechanics.

2.1. **Remarks on the assumptions.** There are several remarks on the assumptions that are worthwhile to mention here. The assumption A.2 on the existence of an asymmetric metric structure is useful because has several applications. First, note that if an asymmetric metric structure is given, then there is a well defined topological structure on  $\mathcal{M}$ . There is a natural topology associated with the asymmetric metric function  $\rho$ . Also, the notions of Cauchy sequence and completeness associated to the asymmetric metric  $\rho$  are well defined, although not symmetric (similar to what happens in Finsler geometry [4]). Second, An asymmetric metric as fundamental metric structure of the configuration space can be used to describe a fundamental irreversible dynamics and hence accommodate assumption A.8 by linking the dynamics with the metric structure of the space. The typical example of asymmetric metric in the category of smooth manifolds are Randers spaces [38], originally introduced by G. Randers in an attempt to describe the irreversibility of our world as a fundamental property of the spacetime arena. In this paper we introduce and apply a theory of dual Randers spaces

as models for deterministic, local, causal and non-reversible dynamics at the Planck scale in the continuous geometry approximation.

Although the formulation that we present is in terms of continuous models, the degrees of freedom define a discrete set. Hence the probability measure is discrete and in some sense equivalent to counting fundamental degrees of freedom. Moreover, the measure properties and the metric properties are logically separated concepts. The fact that the metric and measure structures are separated is a distinctive characteristic of the mm-Gromov spaces [24]. It is in this category of geometric spaces, with a class of specific regular functions (1-Lipschitz functions), that the formulation of the theory of Hamilton-Randers dynamical systems appears natural.

Let us consider assumption A.3. That the degrees of freedom at the fundamental scale are deterministic and localized is against the main stream approaches to the problem of quantum gravity. However, in defense of our deviated point of view, we have to admit that very few is known with certainty of the dynamics at the Planck scale and therefore, the possibility to have a deterministic description is at least technically rather appealing. In deterministic approaches the main problem is to recover the quantum description from the given fundamental dynamics in an appropriate limit.

It is of relevance the issue of the local/non-local character of the interactions of the degrees of freedom at the Planck scale. In favour of locality, we are supported by a natural geometric appealing, since it is possible to imagine local interactions, but a non-local interaction is far from being integrable in a geometric picture. We also think that a theory which is fundamental needs to be of local character, since in any pretended fundamental theory, there is no room left for explaining non-locality at a more fundamental level and a theory that aims to explain non-local phenomena needs of fundamental local explanations to be falsified by local experiments. However, deep difficulties accompany the search for local descriptions of a sub-quantum theory, as it is Bell's inequalities violation. A possible explanation of how to explain the violation of Bell's inequalities is discussed in this work, although a complete development of the ideas is still missing.

Assumptions A.4 and A.7 are in concordance with many current approaches to the problem of quantum gravity. Assumption A.4 implies that the Planck scale is fundamental in our approach, while assumption A.7 requires that the speed of light is maximal and therefore, independent of the observer for each particular direction (hence allowing for anisotropic structures and other local relativity groups other than Poincaré's group). The values of these two scales cannot be determined in our theory and are imposed by consistency with the current idea that the Planck scale is fundamental and with experimental evidence regarding the independence of the speed of light in vacuum.

At this point we should state clearly that because of how the external time parameters (denoted by  $\tau$ -time) is defined, they are discrete parameters. The degrees of freedom appear as discrete too, which is compatible with the existence of a minimal inertial coordinate difference  $L_{min}$ . Despite the discreteness that the assumptions require for the physical systems at the Planck scale, we consider in this paper continuous models for the dynamics. This is motivated by the smallness of the Planck scale compared with any other scale that appear in physics but also by mathematical convenience. In the continuous approximation, where  $\tau$  is continuous instead of discrete, several of the assumptions should be re-written as follows:

- A.1.bis. There is smooth manifold  $M_4$  which is the spacetime manifold.
- A.2.bis. There is a configuration smooth manifold  $\mathcal{M}$  which is endowed with an asymmetric metric structure (2.1) which is at least  $C^2$ -smooth on  $\mathcal{M}$ .
- A.5.bis. The dynamics respect to the  $\tau$ -time parameter is deterministic and given by a system of first order ordinary differential equations.

In the continuous approximation the rest of the assumptions remain formally the same than in the original formulation. However, because the different categories (smooth manifold category versus discrete topological spaces category), the implementation and the techniques that we can use are different than in the discrete case. In the continuous limit, the assumption of the existence of minimal length must be interpreted as a theoretical constraint on the kinematics, despite that there is no indication that in the continuous limit such constraint is necessary. Hence our continuous theory can only be considered as an effective approximation to a fundamental discrete theory. We postpone to the future the development of a complete discrete theory for dynamical models at the fundamental scale.

In the continuous approximation that we can speak directly of topological manifold and dimension. The fact that all our observations are linked to macroscopic devices and can be represented with consistent accuracy in a 4-dimensional spacetime is motivation enough to consider a manifold structure as the arena to represent quantum and classical phenomenology.

We do not introduce a metric or pseudo-Riemannian structure at this stage:  $M_4$  is only the arena were each possible measurement performed by any possible macroscopic observers can happen. Moreover, since the dimension of the spacetime manifold is four, the existence of a smooth structure is not a trivial requirement if we provide only the topology associated to an asymmetric metric  $\rho$ . In the present paper we assume a differential structure on  $M_4$  and on  $\mathcal{M}$ .

2.2. Maximal proper acceleration. A direct consequence of the assumptions A.3, A.4, A.6 and A.7 is the existence of a maximal universal proper acceleration for sub-quantum atoms and sub-quantum molecules. An heuristic argument considers one space dimensional problems. To show such direct consequence we follow the theory developed in reference [19], where an effective kinematical theory of spacetimes with metrics of maximal acceleration was constructed. Due to the fact that there is a lower bound for the difference between coordinates in any instantaneous inertial system (by assumption A.4), it holds that for any elementary work the relation

$$\delta \mathcal{W} := \vec{F} \cdot \delta \vec{L} = \delta L m a,$$

where  $\vec{F}$  is the external mechanical action on the sub-quantum molecule caused by the rest of the system. It is defined by the quotient  $\vec{F} := \frac{\delta W}{\delta \vec{L}}$ , the symbol  $\delta \vec{L}$  representing the infinitesimal displacement of the sub-quantum molecule caused by the rest of the system in the instantaneous coordinate system associated to the sub-quantum molecule at the instant just before the sub-quantum molecule suffers the interaction, a is the value of the acceleration in the direction of the total exterior effort is done and the parameter m is the inertial mass of the sub-quantum molecule S. By hypothesis A.4, it holds that  $\delta L = L_{min}$ . Hence we have that  $\delta W = L_{min} m a$ . For an infinitesimal work, the change in speed is much smaller than the speed of light, since an infinitesimal work is very small in magnitude by definition. Therefore we can use a low speed Newtonian approximation and trust Newtonian mechanics of point particles. Then in a instantaneous coordinate systems with the sub-quantum particle, one has the expression

$$\delta \mathcal{W} = \frac{1}{2} \left( \delta m \, v_{max}^2 + \, m (\delta v)_{max}^2 \right) = L_{min} \, m \, a.$$

Since according to assumption A.6 there is a maximal domain  $\mathcal{U}$  that determines the effect on dynamics of  $\mathcal{S}$ , we have that  $\delta m \leq C m$ , with C a constant of dimension 1 that depends on the size of  $\mathcal{U}$ . One can imagine  $\mathcal{U}$  as a lattice surrounding the sub-quantum molecule. Hence C is a measure of the size of  $\mathcal{U}$  respect to the size of the sub-quantum molecule.

Assume that there is no change in the matter content of  $\mathcal{S}$ . Hence the relation

$$\delta m = 0$$

holds good. Since the speed of any physical degree of freedom is bounded by the speed of light by the assumption of locality A.7, one has that

$$\delta v_{max} \le v_{max} = c$$

Hence the maximal infinitesimal work at instantaneous rest with the particle produced by the system on a point particle is

$$m L_{min} a \le m c^2.$$

This relation implies an universal bound for the value of the proper acceleration a for the sub-quantum system S as

$$a \leq \frac{c^2}{L_{min}}$$

Therefore, if assumptions A.3, A.4, A.6 and A.7 hold good, it is natural to require that the following assumption holds,

• A.10. There is a maximal, universal proper acceleration for both subquantum atoms and di-atomic sub-quantum molecules. The value of the maximal acceleration is

(2.1) 
$$a_{max} = \frac{c^2}{L_{min}}$$

In the continuous approximation, where the degrees of freedom follow a continuous dynamics instead of a discrete dynamics, the assumption A.10 cannot be derived heuristically as we did. Hence in the continuous case,

the assumption A.10 is an independent constraint on the kinematics of the theory. That such kinematical theory exists and is indeed compatible with the Lorentz invariance group has been shown by the author in [19]. The leading order term of the geometry of maximal acceleration is a Lorentzian structure. In this work the Lorentzian structure has been adopted, living the full treatment of spaces with maximal acceleration for future investigations.

#### 3. HAMILTON-RANDERS SYSTEMS

3.1. Geometric background. According to assumption A.1.bis, let us consider the spacetime four-manifold  $M_4$  as the geometric arena where every macroscopic event happens.  $M_4$  does not depend on the observer. On the other hand, the configuration manifold  $\mathcal{M}$  of the Hamilton-Randers system is a tangent space TM such that the base manifold M is diffeomorphic to a product structure of the form

$$(3.1) M \cong \prod_{k=1}^{N} M_4^k$$

with each  $M_4^k$  diffeomorphic to  $M_4$ . Then the tangent configuration manifold for a classical gas of point particles is a smooth manifold M of the form

$$TM \cong \prod_{k=1}^{N} TM_4^k.$$

and the dimension  $\dim(TM) = 2 \dim(M) = 8 N$ . The canonical projections are  $\pi_k : TM_4^k \to M_4^k$ . The vertical fiber over  $x(k) \in M_4^k$  is  $\pi_k^{-1}(x(k))$ . For the dynamical systems as what we are interested in this work we as-

For the dynamical systems as what we are interested in this work we assume that the dimension dim(TM) = 8N is large compared with  $dim(TM_4) =$ 8. Note that choosing the configuration space  $\mathcal{M}$  as a tangent space TMinstead than the base manifold M allows to implement geometrically second order differential equations in a straight way.

Each of the sub-quantum molecules is labeled by a natural number  $k \in \{1, ..., N\}$ . The configuration manifold  $TM_4^k$  is the configuration manifold for the k-th sub-quantum molecule. The dimension  $\dim(TM_4^k) = 8$  and locally each point in the tangent space  $TM_4^k$  is described by using four spacetime coordinates  $(\xi^1, \xi^2, \xi^3, \xi^4)$  of the point  $\xi(k) \in M_4^k$  and four independent velocities coordinates  $T_{\xi}M_4^k \ni (\dot{\xi}^1, \dot{\xi}^2, \dot{\xi}^3, \dot{\xi}^4)$ .

It is possible to consider more general configuration manifold for the description of the dynamics of other degrees of freedom of the sub-quantum molecules. However, in this work only spacetime configuration manifold are investigated. This attitude is based on the following grounds. Although, it is remarkable that when considering point particles, other properties as spin are associated with the quantum description, it is a reasonable assumption that in quantum mechanics all measurements are ultimately reduced to local position and time measurements. Hence we can adopt such reductionist point of view first. Moreover, one of our goals is to investigate the possibility of emergence of spacetime. Macroscopically, spacetime is a four-dimensional manifold. Therefore, it appears natural that the configuration manifold  $M_4^k$ for each sub-quantum molecule is a four-manifold. Henceforth, the geometry of four-manifolds is our basis for the formulation of the dynamical models, as a first treatment towards a fully discrete theory.

Given  $M_4$  and the collection of four-manifolds  $\{M_4^k\}_{k=1}^N$ , there is also a collection of diffeomorphisms

(3.2) 
$$\Upsilon := \{ \varphi_k : M_4^k \to M_4, \, k = 1, ..., N \}$$

This is in accordance with the assumption A.4, stating that each of the N degrees of freedom are indistinguishable and identical. We denote by

$$\varphi_{k*}: T_{x_k}M_4^{\kappa} \to T_{\varphi_k(x)}M_4$$

the differential map at  $x_k$  and by

$$\varphi_k^*: T_{\varphi(x_k)}T^*M \to T_{x_k}^*M_4$$

the pull-back of 1-forms at  $\varphi_k(x) \in M_4$ .

Given  $M_4^k$  and  $M_4$ , there are is a non-denumerable set of possible diffeomorphisms  $\varphi_k : M_4^k \to M_k$ . However, any two of such diffeomorphisms  $\varphi_k, \tilde{\varphi}_k$  are related by a diffeomorphism  $\varphi_{tran} \in \text{Diff}(M_4)$ , where  $\text{Diff}(M_4)$ is the group of diffeomorphims of  $M_4$ . Therefore, any particular choice of the diffeomorphisms  $\varphi_k$  can be seen as an ambiguity of our theory. The natural way to solve this problem is to require for all observable predictions of the theory to be invariant under  $\text{Diff}(M_4)$ , in which case  $\text{Diff}(M_4)$  is a symmetry of the theory. This consistence requirement must constraint the mathematical formalism.

The model four-manifold  $M_4$  is endowed with a Lorentzian metric  $\eta_4$  of signature (1, -1, -1, -1). Moreover, for each  $k \in \{1, ..., N\}$  there is a Lorentzian metric  $\eta_4(k)$  on  $M_4^k$  and we assume that each of the Lorentzian structures

$$\mathbf{M}_4 := \{ (M_4^k, \eta_4(k)), \ k = 1, ..., N \}$$

is isometric to the Lorentzian model  $(M_4, \eta_4)$ .

The algebra of functions  $\mathcal{F}_D(T^*TM)$  on  $T^*TM$  that we consider is the algebra of *diagonal smooth functions*, obtained by an algebra embedding

(3.3) 
$$\theta: \mathcal{F}(T^*TM_4) \to \mathcal{F}(T^*TM), \quad f \mapsto (f_1, ..., f_N)$$
$$f_k = f_0, \quad f_0: T^*TM_4 \to \mathbb{R}, \quad k = 1, ..., N.$$

Hence the algebra of functions  $\mathcal{F}_D(T^*TM)$  is the subset of  $\mathcal{F}(T^*TM)$  whose elements are functions of the form

$$\mathcal{F}(T^*TM) \to \mathcal{F}(T^*TM), ((u_1, p_1), ..., (u_N, p_N)) \mapsto (f_0(u_1, p_1), ..., f_0(u_N, p_N))$$

In this way, the admissible functions on  $T^*TM$  are modeled on the functions defined on  $T^*TM_4$ .

Measure and metric structures. The expectation values of diagonal functions  $f \in \mathcal{F}_D(T^*TM)$  are of physical relevance, since such averages describe macroscopic observables, they must be invariant of the particular choice of the collection of diffeomorphisms  $\varphi$ . Such invariance is achieved if the measure  $\mu_P$  used in the averages in  $T^*TM$  is a product measure,

(3.4) 
$$\mu_P = \prod_{k=1}^{N} \mu_P(k),$$

where  $\mu_P(k)$  is a Diff $(M_4)$ -invariant probability measure in  $T^*TM_4^k$ . The construction of an invariant measure  $\mu_P$  on TM and other associated measures is presented below.

The Levi-Civita connection of  $\eta_4(k)$  determines the horizontal distribution in a canonical way [33]. Given this standard connection, here is defined a pseudo-Riemannian metric  $\eta_S^*(k)$  on  $TM_4^k$  (the Sasaki-type metric), which is the Sasaki-type lift of the metric  $\eta_4^k$  on  $M_4^k$  to  $TM_4^k$  by the distribution associated with the Levi-Civita connection. The dual metric to  $\eta_S^*(k)$  is the dual pseudo-Riemannian metric  $\eta_S(k) = (\eta_S^*(k))^*$ . The dual Sasaki-type metrics  $\{\eta_S^*(k)\}_{k=1}^N$  allows to define the dual pseudo-Riemannian metric<sup>1</sup>

(3.5) 
$$\eta = \oplus_{k=1}^N \eta_S(k)$$

on the 8N-dimensional cotangent space  $T^*TM$ .

**Measures**. The Lorentzian metric  $\eta_4$  allows to define a Diff $(M_4)$ -invariant volume form  $dvol_{\eta_4}$  on  $M_4$  in a canonical way,

(3.6) 
$$dvol_{\eta_4} = \sqrt{-\det \eta_4} \, dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4.$$

Moreover, it is possible to define an Diff $(M_4)$ -invariant vertical form  $dvol_k(y_k)$  on each fiber  $\pi_k^{-1}(x(k))$ ,

(3.7) 
$$d^4 z_k = \sqrt{-\det \eta_4} \, dy^1 \wedge \, dy^2 \wedge \, dy^3 \wedge \, dy^4,$$

Then we can give a concrete expression for the measure  $\mu_P$ ,

(3.8) 
$$\mu_P = \prod_{k=1}^N d^4 z_k \wedge dvol_{\eta_4}.$$

The measure (3.8) is invariant under diffeomorphisms of  $M_4$ . Also important, the measure can be pull-back to convenient to sub-manifolds of TM. **Deterministic dynamics for the ontological degrees of freedom**. Given this differential geometric setting, the dynamical systems that we consider are systems of ordinary first order differential equations of the form (3.9)

$$\frac{d\xi_k^{\mu}}{d\tilde{\tau}} = \beta_{kx}^{\mu}(\xi, \dot{\xi}, \tilde{\tau}), \quad \frac{d\dot{\xi}_k^{\mu}}{d\tilde{\tau}} = \beta_{ky}^{\mu}(\xi, \dot{\xi}, \tilde{\tau}), \qquad \mu = 1, 2, 3, 4, \ k = 1, ..., N.$$

This is a coupled 8*N*-dimensional system of special type called semi-spray [35]. In order to determine locally the solutions, it is necessary and sufficient to provide the set of initial conditions  $\{\xi_k^{\mu}(0), \dot{\xi}_k^{\mu}(0)\}_{k=1,\mu=1}^{N,4}$ . We will see that there are two mathematical theories, related with different mathematical formalism, that describe dynamical systems of the type (3.9). One is a geometric theory of Hamilton-Randers spaces and the second is the quantized theory of Hamilton-Randers spaces.

It is remarkable that the equations of motion (3.9) are equivalent to 4N second order differential equations of semi-spray type. Our formalism has been set up in this way to accommodate the first Newton law of the dynamics, the law of inertia, for each of the sub-quantum molecules.

<sup>&</sup>lt;sup>1</sup>Note that the use of \*-notation for dual metrics and norms here is partially inverted respect to the usual notation in Riemannian geometry. For instance,  $\eta_S^*$  is a metric on  $TM_4$ , while  $\eta_S$  is a metric in  $T^*M_4$ .

3.2. Notion of Hamilton-Randers space. The first formalism for the dynamical systems that we consider in this work is a geometric formalism, based on the two notions of Hamilton space and Randers space.

Notion of generalized Hamiltonian function. Let M be a smooth manifold and  $\mathcal{C} \subset T^*\widetilde{M}$  a connected open cone of  $T^*\widetilde{M}$ . Let the topological closure of  $\mathcal{C}$  respect to the manifold topology to be  $\overline{\mathcal{C}}$ .

**Definition 3.1.** A generalized Hamiltonian function is a triplet  $(\widetilde{M}, F, C)$ with  $F : \overline{C} \to \mathbb{R}^+ \cup \{0\}$  smooth on the open cone  $C \hookrightarrow T^*\widetilde{M}$ .

The vertical Hessian (or fundamental tensor g) of the function  $F^2$  in natural coordinates  $\{(u^i, p_j), i, j = 1, ..., dim(\widetilde{M})\},\$ 

(3.10) 
$$g^{ij}(u,p) = \frac{1}{2} \frac{\partial^2 F^2(u,p)}{\partial p_i \partial p_j}.$$

The function F is the generalized Hamiltonian function. Only points where F is positive or zero are of physical relevance. Direct from this definition is that the topological closure of C is

$$\bar{\mathcal{C}} = \{(u, p) \in T^*TM \ s.t. \ F(u, p) = 0\}.$$

Moreover, if one does not fix the external time parameter and we want to recover equations (3.9) from a dynamics related with F(u, p), the domain of definition should allow for global, positive re-scaling on the momenta coordinates  $p \mapsto \lambda p$ . Hence the domain of definition of F must be a cone.

Notion of pseudo-Randers space. In the category of positive definite Finsler geometry, a Randers structure on a manifold  $\widetilde{M}$  is a Finsler structure such that the associated Finsler function is of the form

$$F^*: TM \to \mathbb{R}, \quad (u, \tilde{y}) \mapsto \alpha^*(u, \tilde{y}) + \beta^*(u, \tilde{y}),$$

with  $\alpha^*(u, \tilde{y})$  the Riemannian norm of  $\tilde{y} \in T_u \widetilde{M}$ ,  $\beta^*(u, \tilde{y})$  the result of the action 1-form  $\beta^* \in \Gamma T^* \widetilde{M}$  on  $\tilde{y}$ . The condition

$$(3.11) \qquad \qquad \alpha^*(\beta,\beta) < 1.$$

must be satisfied. The condition (3.11) implies the non-degeneracy and the positiveness of the associated fundamental tensor [4].

We now consider the analogous to a Randers structure in the category  $\mathcal{F}_{HR}(TM)$  of generalized Hamiltonian functions on TM whose fundamental tensors (3.10) have non-definite signature. In this case, it is necessary to restrict the domain of definition of the Hamiltonian function F, since it is not possible to have a well defined Hamilton-Randers function on the whole  $T^*TM$ . This is because being  $\eta$  a pseudo-Riemannian metric, it can take negative values on certain domains of  $T_u^*TM$ , in which case the function  $\alpha(u, p)$  is purely imaginary. Hence the domain of a Hamilton-Randers function is restricted to be the topological closure of the open cone  $\mathcal{C}_u \hookrightarrow T_u^*TM$  over u timelike momenta. The cone of timelike momenta over  $u \in TM$  is defined by the set of co-vectors  $p \in T_u^*TM$  such that

(3.12) 
$$\alpha(u,p) = \sum_{i,j=1}^{8N} \eta^{ij}(u) p_i p_j > 0.$$

14

That it is a cone is direct since if  $p \in C_u$  then  $\lambda p \in C_u$ ; that  $C_u$  is open is because it is the domain of an open set  $(0, +\infty)$  by a continuous map  $F = \alpha(u, p) + \beta(u, p)$ .

Notion of Hamilton-Randers space Let  $\beta \in \Gamma TTM$  be a vector field such that a dual condition to the Randers condition (3.11)

(3.13) 
$$\eta^*(\beta,\beta) < 1, \quad \beta \in \Gamma TTM$$

holds good.

**Definition 3.2.** A Hamilton-Randers space is a generalized Hamiltonian space whose Hamiltonian function  $F : \mathcal{C} \to TM$  is of the form

(3.14) 
$$F: \mathcal{C} \to \mathbb{R}^+, \quad (u, p) \mapsto F(u, p) = \alpha(u, p) + \beta(u, p).$$

with  $\alpha = \sqrt{\eta_{ij}(u)p^ip^j}$  real on  $\mathcal{C}$  and where

$$\beta(u,p) = \sum_{i=1}^{8N} \beta^i p_i,$$

with  $\beta$  constrained by the condition (3.13).

The space of Hamilton-Randers structures on TM is denoted by  $\mathcal{F}_{HR}(TM)$ . One can consider natural geometric flows in the space of Hamilton-Randers structures  $\mathcal{F}_{HR}(TM)$ . The geometric flow responsible for the dynamics in Hamilton-Randers systems is denoted by  $U_t$ ,

$$U_t: \mathcal{F}_{HR}(TM) \to \mathcal{F}_{HR}(TM).$$

There is also an induced flow on  $\tilde{\mathcal{C}}$ , the topological closure of  $\mathcal{C}$ . It is required that for the  $U_t$ -flow holds the following general properties,

(1) There is a 1-parameter family of connected submanifolds

$$\{\mathcal{E}_u(r) \hookrightarrow T_u^*TM, u \in TM\}$$

which are invariant under the action of  $U_t$ . These manifolds define metastable domains.

(2) The dynamics of the connected components of the r-hyperboloid

$$\Sigma_u(r) := \{ p \in T^*TM \, s.t. \, F^2(u, p) = r^2 \}$$

by the *t*-dynamics has as limit the corresponding  $\mathcal{E}_u(r)$ . This limit is reached in a finite time evolution  $t_b - t_a = T$ .

(3) The domains

$$\mathbf{D}_0 := \{ t = (2n+1) T, n \in \mathbb{Z} \}$$

are metastable. The properties of maximal acceleration and maximal speed of HR-systems prevents the system to stay for a long *t*-time in the metastable equilibrium regime. Then a process of expansion in phase space  $T^*TM$  starts again, until the dynamical system becomes ergodic again.

These flows contrast with the situation in general relativity, where the geometry is fixed by Einstein's equations without the need of an external time evolution. 3.3. Interpretation of the parameter t and the semi-period T. The time scale parameter T, that has the properties of a time semi-period depends on the size of the physical system. Since the sub-quantum particles are identical, the metastable domains  $\mathbf{D}_0$  are reached faster for large systems. In this way,  $T^{-1}$  is an indicator of the size of the system.

The meaning of the t-time parameter is not direct, since there is no geometric structure on TM that allows us to define an internal t-time parameter. Hence the theory must be invariant under t-time re-parameterizations and the parameter t should not have direct physical meaning.

On the other hand, the *t*-parameter cannot be fully arbitrary, since the inverse of the semi-period T for the  $U_t$  dynamics is related with the size of the system. This dichotomy is resolved if we make precise the relation between the semi-period T and a measure of the size of the system. This relation could be written in the form

(3.15) 
$$T = \alpha \frac{\hbar}{Mc^2},$$

where M is a coefficient that we can associate to the mass at rest of the quantum system and  $\alpha$  is a constant that depends on the choice of the arbitrary parameter t but that is the same for all the quantum systems.

The relation (3.15) is invariant under *t*-time re-parameterizations,

$$\frac{T_1}{\alpha_1} = \frac{T_2}{\alpha_2} = \frac{1}{M c^2} \hbar$$

for two arbitrary quantum systems.

Note that M depends on the quantum system and is not the mass of the sub-quantum molecule. The minimal mass that one can think associated with a quantum particle is

$$(3.16) M = m.$$

This relation corresponds to quantum massless particles. This must be understood as an asymptotic limit, in the sense that for any other quantum particle  $M \gg m$ . It corresponds to the quantum particle of maximal period  $T_{max}$ , which is very large and can be approximated as bounded for many considerations. The maximal mass associated to a quantum particle corresponds to the minimal period  $T_{min}$ .

Despite the resemblance with the energy-time uncertainty relation, the relation (3.15) is not a quantum energy-time relation, since it relates semiperiods of the *t*-time with energy of the system and also it is an equality, instead of an inequality.

3.4. **Properties of the**  $U_t$  flow. Given the Hamilton-Randers space (TM, F, C), it is possible to define a Riemannian structure  $h \in \Gamma T^{(2,0)}TM$  by averaging the fundamental tensor components  $g_{ij}(u, p)$  on each  $\Sigma_u(1)$ . The tensor components  $h^{ij}$  are obtained by averaging the metric coefficients  $g^{ij}(u, p)$ on the open cone  $\Sigma_u(1)$  at  $u \in TM$ ,

(3.17) 
$$h^{ij}(u) := \frac{1}{\int_{\Sigma_u} dvol_{\Sigma_u}(p)} \int_{\Sigma_u} dvol_{\Sigma_u}(p) g^{ij}(u,p),$$

i, j = 1, ..., 8N, where  $dvol_{\Sigma_u}$  is the volume form on the unit hyperboloid  $\Sigma_u$ . Although the sub-manifold  $\Sigma_u$  is not compact, the volume form  $dvol_{\Sigma_u}(p)$  is chosen such that the integrals in (3.17) are well defined. Then the squared norms h(u, p) and g(u, p) are defined by

$$h(u,p) = h^{ij}(u) p_i p_j, \quad g(u,p) = g^{ij}(u,p) p_i p_j = F^2(u,p).$$

**Definition 3.3.** The  $U_t$  dynamics in the interval  $[0,T] \subset \mathbb{R}$  is a geometric flow of the form

(3.18) 
$$U_t: \mathcal{F}_{HR}(TM) \to \mathcal{F}_{HR}(TM),$$
$$F \mapsto F_t = \sqrt{\kappa(u, p, t, \tilde{\tau}) |h(u, p)| + (1 - \kappa(u, p, t, \tilde{\tau})) |g(u, p)|},$$

such that the function  $\kappa : \mathcal{F}_{HR}(TM) \times [0,T] \to [0,T] \subset [0,1]$  satisfies the boundary conditions

(3.19) 
$$\lim_{t \to 0^+} \kappa(u, p, t, \tilde{\tau}) = 0, \quad \lim_{t \to T^-} (\kappa(u, p, t, \tilde{\tau}) - 1) = 0.$$

Hence the  $U_t$ -flow determines an homotopy transformation in the space  $\mathcal{F}_{HR}(TM)$  compatible with the conformal transformations

(3.20) 
$$g \to (1 - \kappa(u, p, t, \tilde{\tau})) g.$$

Such interpolation and conformal transformations are compatible with the properties (1) and (2) of the flow  $U_t$ . Note that in this definition  $t \in [0, 2T]$ , instead of  $t \in \mathbb{R}$ , in concordance with the assumption that the dynamics becomes almost periodic, with period 2T (that encompasses a sequence of an expanding phase, an ergodic phase and a concentration phase).

Let us consider the convex hull  $C_{hq} \subset \mathcal{F}_{HR}(TM)$  containing g and h,

$$C_{hg} := \{ F \in \mathcal{F}_{HR}(TM) \, s.t. \, g_F = t_1 \, g + t_2 \, h, \, t_1 + t_2 = 1, \, t_1, t_2 \ge 0 \},$$

where  $g_F$  is the fundamental tensor of F. Then we have that

**Proposition 3.4.** Every Hamilton-Randers structure F in the convex hull  $C_{hg}$  containing g and h evolves towards the averaged structure  $h \in \Gamma T^{(2,0)}TM$  under the action of the flow (3.18),

(3.21) 
$$\lim_{t \to T} U_t(F) = \sqrt{h(u, p)}, \quad \forall F \in C_{hg}$$

*Proof.* Note that any  $F \in C_{hq}$  has the same averaged metric,

$$\langle g_F \rangle = \langle t_1 g + t_2 h \rangle = t_1 h + t_2 h = h.$$

Then we have

$$\lim_{t \to T} U_t(F) = \lim_{t \to T} \sqrt{\kappa(u, p, t, \tilde{\tau}) h(u, p) + (1 - \kappa(u, p, t, \tilde{\tau})) g_F(u, p)}$$
$$= \lim_{t \to T} \sqrt{\kappa(u, p, t, \tilde{\tau}) h(u, p) + (1 - \kappa(u, p, t, \tilde{\tau})) g_F(u, p)}$$
$$= \sqrt{h(u, p)}.$$

Hence the  $U_t$ -flow is information loss in  $\mathcal{F}_{HR}(TM)$ , since the structure h is the limit of the convex hull  $C_{hg}$  and there are many structures F evolving towards the same metastable limit h.

 $\square$ 

3.5. **t-time inversion operation.** The parameter  $t \in \mathbb{R}$  is interpreted as the time parameter for an *internal dynamics* of the system. The *time inversion operation*  $\mathcal{T}_t$  is defined in local natural coordinates on  $T^*TM$  by the operator

(3.22) 
$$\begin{aligned} \mathcal{T}_t : T^*TM \to T^*TM, \\ (u,p) = (x, y, p_x, p_y) \mapsto (\mathcal{T}_t(u), \mathcal{T}_t^*(p)) = (x, -y, -p_x, p_y). \end{aligned}$$

The induced action of  $\mathcal{T}_t$  on elements  $F \in \mathcal{F}_{HR}(TM)$  is given by the expression

$$\mathcal{T}_t(F)(u,p) := F(\mathcal{T}_t(u), \mathcal{T}_t(p)).$$

Note that a Hamilton-Randers metric is non-reversible in the sense that

$$F(u,p) \neq F(\mathcal{T}_t(u),\mathcal{T}_t(p))$$

except for a subset of measure zero in  $(u, p) \in T_u^*TM$ . From this relation the intrinsic irreversible character of the Randers geometry follows. However, we assume the natural condition that  $\mathcal{T}_t$  commutes with the  $U_t$  dynamics,

$$(3.23) [U_t, \mathcal{T}_t] = 0, \forall t \in [0, T] \subset \mathbb{R}$$

This commutation relation guarantees that  $\mathcal{T}_t(F)$  and F are in the same equivalence class [h], if h is invariant under  $\mathcal{T}_t$ . Indeed, one can prove by a continuity argument on the parameter t that for  $t = n \in \mathbb{N}$  there is an  $\epsilon \in \mathbb{R}$  such that the condition (3.23) holds for each  $\tilde{t} \in (n - \epsilon, n], n \in \mathbb{N}$ .

We can perform the explicit calculation of  $\mathcal{T}_t h$ ,

$$\begin{split} \mathcal{T}_t(h) &= \mathcal{T}_t(\lim_{t \to T} U_t(F)) \\ &= \mathcal{T}_t(\lim_{t \to T} \kappa(u, p, t, \tilde{\tau}) h(u, p) + (1 - \kappa(u, p, t, \tilde{\tau})) g_F(u, p)) \\ &= \lim_{t \to T} \mathcal{T}_t(\kappa(u, p, t, \tilde{\tau}) h(u, p) + (1 - \kappa(u, p, t, \tilde{\tau})) g_F(u, p)) \\ &= \lim_{t \to T} \left( \mathcal{T}_t(\kappa(u, p, t, \tilde{\tau})) \mathcal{T}_t(h)(u, p) + \mathcal{T}_t(1 - \kappa(u, p, t, \tilde{\tau})) \mathcal{T}_t(g_F)(u, p)) \right) \\ &= \lim_{t \to T} \left( \mathcal{T}_t(\kappa(u, p, t, \tilde{\tau})) h(u, p) + \mathcal{T}_t(1 - \kappa(u, p, t, \tilde{\tau})) g_F(u, p) \right). \end{split}$$

Therefore, a sufficient condition for the invariance  $\mathcal{T}_t h = h$  is that

(3.24) 
$$\mathcal{T}_t(\kappa(u, p, t, \tilde{\tau})) = \kappa(u, p, t, \tilde{\tau}).$$

Furthermore, it is clear that  $\mathcal{T}_t$  is an idempotent operator,

(3.25) 
$$(\mathcal{T}_t)^2 = Id, \quad \forall t \in I \subset \mathbb{R}.$$

3.6. The  $U_t$  flow is information loss. Hamilton-Randers structures and linear Hamiltonian functions on momentum variables are strongly related. Let us prove in detail this fact. If (TM, F, C) is a Hamilton-Randers space that evolves towards the final averaged structure (TM, h) by the  $U_t$  flow, for each value of t there is an element  $(TM, F_t)$  of  $\mathcal{F}_{HR}(TM)$ . Applying the time inversion operation  $\mathcal{T}_t$  to  $F_t$  and taking into account that the function  $\kappa(u, p, t, \tilde{\tau})$  is invariant under  $\mathcal{T}_t$ , one obtains the corresponding Hamiltonian function of a HR-systems at the instant  $(t, \tau)$ ,

$$\begin{aligned} H_t(u,p) &= \frac{1}{2} F_t(u,p) - \frac{1}{2} F_t(\mathcal{T}_t(u),\mathcal{T}_t^*(p)) \\ &= \frac{1}{2} (1 - \kappa(u,p,t,\tilde{\tau})) \left( g^{ij}(u,p) p_i p_j \right)^{1/2} + \frac{1}{2} \kappa(u,p,t,\tilde{\tau}) \left( \langle g^{ij} \rangle p_i p_j \right)^{1/2} \\ &- \frac{1}{2} (1 - \kappa(u,p,t,\tilde{\tau})) \left( g^{ij}(\mathcal{T}_t(u),\mathcal{T}_t^*(p)) p_i p_j \right)^{1/2} - \frac{1}{2} \kappa(u,p,t,\tilde{\tau}) \left( \langle g^{ij} \rangle p_i p_j \right)^{1/2} \\ &= \frac{1}{2} (1 - \kappa(u,p,t,\tilde{\tau})) \left( g^{ij}(u,p) p_i p_j \right)^{1/2} \\ &- \frac{1}{2} (1 - \kappa(u,p,t,\tilde{\tau})) \left( g^{ij}(\mathcal{T}_t(u),\mathcal{T}_t^*(p)) p_i p_j \right)^{1/2} \\ &= (1 - \kappa(u,p,t,\tilde{\tau})) \sum_{k=1}^{8N} \beta^k(u) p_k. \end{aligned}$$

Hence the Hamiltonian function associated with a HR-system at  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$  is

(3.26) 
$$H_t(u,p) = (1 - \kappa(u,p,t,\tilde{\tau})) \sum_{k=1}^{8N} \beta^k(u) p_k.$$

Then by the limit conditions (3.19) it holds the following condition,

**Proposition 3.5.** The Hamiltonian in the metastable equilibrium regime is the average Hamiltonian of a HR-system, which is identically zero,

(3.27) 
$$\lim_{t \to (2n+1)T} H_t(u,p) = 0.$$

3.7. Notion of external time parameter  $\tau$ . The  $U_t$  flow is almost cyclic. By this we mean that the total  $U_t$  evolution is sub-divided in fundamental cycles with

$$t \in [(2n+1)T, (2n+3)T], n \in \mathbb{Z}.$$

Each of these fundamental cycles is composed by a ergodic regime for the evolution of the sub-quantum molecules defining the system, followed by a contractive regime, followed by an expanding regime. Then a new ergodic regime start, defining the next fundamental cycle. The period of each fundamental cycle, if the quantum particle description of the system does not change, is constant and equal to 2T. This period depends on the physical system (quantum system). Such ergodic-contractive-expansion cycles are universal, that is, they happen for every quantum system. This includes the standard model degrees of freedom but also atomic and some molecular systems. Since physical clocks are based on stable, periodic processes, that can be reduce to the analysis of periodic quantum processes, they are also periodic in the number of fundamental periods. Hence the notion of exterior time  $\tau$  emerges from the processes of the  $U_t$  dynamics,

**Definition 3.6.** An external time parameter  $\tau$  is a  $\mathbb{Z}$ -multiple function of the number of fundamental cycles of a  $U_t$  flow.

Two different species of elementary particles a and b can have different fundamental semi-periods  $T_a$  and  $T_b$ . Therefore, consistence between possible definitions of the  $\tau$ -time parameter using system a or system b as clocks implies naturally the congruence condition

$$T_a = q T_b, \quad q \in \mathbb{N}.$$

Since the species of elementary particles is finite, there is a minimal semiperiod  $T_{min}$  and all other fundamental semi-periods are multiples of  $T_{min}$ . Being  $T_a$  and  $T_b$  non-primes between each other it is possible to define classical and quantum clocks that preserve the congruence between the different constitutive fundamental semi-periods. More generally, these compatibility conditions between each quantum systems suggest that the mathematics describing how different quantum systems interact is the theory of congruences in arithmetics. If fundamental periods are associated with prime numbers, and by the relation (3.15), the minimal semi-period is T = 2 and corresponds to the system describing the whole universe.

Defined in this way, the time  $\tau$ -time parameters are necessarily discrete parameters, describing discrete dynamical processes. However, when used to measure time durations of quantum or macroscopic processes, it can be considered continuous, since the usual quantum and classical scales are much larger than the fundamental scale. Indeed, this is the working hypothesis in this paper, where we describe time as real parameter,  $\tau \in \mathbb{R}$ . Then a direct implication of 3.5 is that the physical spacetime manifold is a topological space  $M_4 \times \mathbb{Z}$  such that the *time topological space*  $\mathbb{R} \times \mathbb{Z}$  in the continuous limit is instead replaced by the *time manifold*  $\mathbb{R} \times \mathbb{R} \equiv \mathbb{C}$  and embedded in a five manifold  $M_5$ . The set of slits

$$\{t = (2n+1)T, n \in \mathbb{Z}\}\$$

defines a 4-dimensional submanifold  $M_4 \hookrightarrow M_5$ , that it is associated with the usual 4-dimensional spacetime manifold.

Let us assume that t-time parameter is continuous. Within the general idea advocated in Hamilton-Randers theory, the degrees of freedom and dynamics of the sub-quantum degrees of freedom are discrete, as well as the  $\tau$ -time parameters. It is natural to doubt the correctness of the continuity hypothesis for the t-time parameter. However, it is particularly useful that t can be treated as continuous, due to the difference in scales between the Planck scale and the quantum scale, as part of an approximation in the description of time and evolution. This is the aptitude advocated in the present work, leaving a more deep and detailed discrete formalization for future work.

A  $\tau$ -slit consists of the domain of the full phase space filled by the  $U_t$  flow between an interval [n T, 2(n+1) T] that corresponds to a given fundamental cycle. Then the *t*-time parameter appears as an internal time parameter: it is *a time inside a time*, since *t*-time parameters describe the evolution of the sub-quantum molecules even between two consecutive instants of  $\tau$ -time.

An alternative interpretation of the pair  $(t, \tau)$  as corresponding to a faster/slow time parameters is not adequate. It implies to identify the  $\tau$ -time parameter with a one to one correspondence with the values of the *t*-time parameter. However, this is not possible, since each  $\tau$ -time parameter is

defined as an integer multiple of a integer number of fundamental cycles, while the parameter t also parameterizes the dynamics on each internal cycle. Identifying the parameter  $\tau$  with a diffeomorphism of the parameter t, in the present context, implies to use a large scale (quantum or classical scale) for describing a sub-quantum scale, which is a contradiction. Hence Hamilton-Randers theory is based on the nobel idea of a two-dimensional time, which is constructive and that emerges from the process of evolution of natural systems.

3.8. The  $U_{\tau}$ -dynamics. The kinematics and dynamics of the sub-quantum molecules respect to the external time parameter  $\tau \in \mathbb{R}$  is described as follows. First, the speed components  $\beta_x$  and the acceleration components  $\beta_y$  of the vector field  $\beta \in \Gamma TTM$  are defined in terms of the time inversion operator  $\mathcal{T}_t$  by the expressions

(3.28) 
$$\beta_x := \frac{1}{2}(\beta - \mathcal{T}_t(\beta)), \quad \beta_y := \frac{1}{2}(\beta + \mathcal{T}_t(\beta)).$$

The set of pre-word lines is

$$\xi_{tk} : \mathbb{R} \to M_4^k, \, \tau \mapsto \xi_{tk}(\tau) \in M_4^k,$$

where the value of the parameter t has been fixed. They determine the loci of the sub-quantum degrees of freedom at a given frozen t. That is, for each of the fundamental cycles, we consider the loci of each k-molecule at  $\{t, 2T + t, 4T + t, ...\}$ .

For HR-systems, the non-degeneracy of the fundamental tensor g of the underlying Hamilton-Randers space is ensured if the vector field  $\beta$  is constrained by the condition (3.13). Such condition implies

- (1) The velocity vector of the sub-quantum atoms is normalized by the condition  $\eta_4^k(\dot{\xi}_{tk}, \dot{\xi}_{tk}) \leq v_{\max}^2 = c^2$ . That is the sub-quantum molecules evolve on the time  $\tau$  in a sub-luminal or luminal world lines.
- (2) If the *on-shell* conditions  $\{\dot{x}^k = y^k, k = 1, ..., 4N\}$  hold, then there is a maximal bound for the proper acceleration  $\eta_4^k(\ddot{\xi}_{tk}, \ddot{\xi}_{tk}) \leq A_{\max}^2$ .

Hence the speed vector and accelerations of the sub-quantum molecules respect to the  $U_{\tau}$  evolution are bounded,

(3.29) 
$$\eta_4(\beta_{kx}, \beta_{kx}) \le c, \quad \eta_4(\beta_{ky}, \beta_{ky}) \le A_{max}^2, \quad k = 1, ..., N.$$

The conditions (3.29) are 4-dimensional covariant.

The classical Hamiltonian function of a HR-system can also be defined by the *partially averaged Hamiltonian function* 

$$H_t(u,p) := \left(\frac{1}{2} F(u,p) - \frac{1}{2} F(\mathcal{T}_t(u), \mathcal{T}_t^*(p))\right).$$

The non-degeneracy for the Hamilton-Randers structure is required to have the following interpretation for the Hamiltonian (3.26). Indeed, under such condition the Hamiltonian is the difference of two individual Hamiltonian functions: one that corresponds to sub-quantum atoms evolving towards the future  $\tau \to +\infty$  and second one that corresponds to sub-quantum atoms evolving back-wards  $\tau \to -\infty$ . This physical interpretation justifies the need of the condition (3.13), in order to have non-degenerate Hamilton-Randers structures and to provide a Hamiltonian interpretation for each of them. Therefore, the Hamiltonian function (3.26) corresponds to a *time orientation average* of the Hamilton-Randers function associated with a particular form of classical Hamiltonian.

The Hamilton equations for H(u, p) are

$$u'^{i} = \frac{\partial H(u,p)}{\partial p_{i}} = 2(1 - \kappa(u,p,t,\tilde{\tau}))\beta^{i}(u) - \frac{\partial \kappa(u,p,t,\tilde{\tau})}{\partial p_{i}} \left(\sum_{k=1}^{8N} \beta^{k}(u)p_{k}\right)$$
$$p'^{i} = -\frac{\partial H(u,p)}{\partial u^{i}} = -2(1 - \kappa(u,p,t,\tilde{\tau}))\sum_{k=1}^{8N} \frac{\partial \beta^{k}(u)}{\partial u^{i}}p_{k}$$
$$+ \frac{\partial \kappa(u,p,t,\tilde{\tau})}{\partial u^{i}} \left(\sum_{k=1}^{8N} \beta^{k}(u)p_{k}\right).$$

3.9.  $U_t$ -flow and redefinition of the  $\tilde{\tau}$ -time parameter. The  $U_t$ -flow has been parameterized by the conformal factor  $\kappa(u, p, t, \tilde{\tau})$ . However, in order to obtain dynamical equations of motion(3.9) it is necessary to reparameterized the  $\tilde{\tau}$ -time parameter. Indeed, it is easy to see that the Hamiltonian (3.26) does not have as Hamilton equations (3.9), except if the following re-parameterizations is allowed:

(3.30) 
$$\tilde{\tau} = (1 - \kappa(u, p, t, \tilde{\tau}))\tau, \quad \frac{d}{d\tilde{\tau}} = \frac{1}{1 - \kappa(u, p, t, \tilde{\tau})}\frac{d}{d\tau}.$$

As it stands (3.30) is ill-defined, since  $\kappa(u, p, t, \tilde{\tau})$  depends on  $u \in TM$ (there is not such a problem with the *p*-dependence, since it is a dummy variable, that it is completely fixed in function of  $\tilde{\tau}$ , once the equations are solved). Hence we are constrained to postulate that  $\kappa(u, p, t, \tilde{\tau})$  is of the form  $\kappa(p, t, \tilde{\tau})$ ,

(3.31) 
$$\frac{\partial \kappa(u, p, t, \tilde{\tau})}{\partial p_i} = 0, \quad i = 1, ..., 8N.$$

This constraint is possible if the structure (TM, F) is of *Berwald-type*. In the positive definite case of Finsler geometry, a Berwald space is such that all the unit spheres are linearly isometric [4]. In the category of Hamilton-Randers structures as defined in this work, we guess that the analogous condition holds and that for a Berwald-type Hamilton-Randers space the unit hyperboloids  $\{\Sigma_u(1), u \in TM\}$  are all linear isometric. This can be taken as the definition os such structures. Then following the analogy with the positive case, the Berwald condition of a Randers space is that

$$(3.32) \nabla\beta = 0,$$

where  $\nabla$  is the Levi-Civita connection of  $\eta$ . The constraint (3.32) is a conservation law for  $\beta$  in the configuration space TM. This conservation law is of relevance for partially explaining quantum correlations.

The Randers category is a good category in the sense that the Randers condition heritages to submanifolds. On the other hand, the Berwald condition is not heritage. This means that the condition (3.32) does not need to hold for each of the individual products  $TM_4^k$ .

In the metastable equilibrium regime  $t \to (2n+1)T$ , the  $\tilde{\tau}$ -parameter is indeed a slow time. Respect to this  $\tau$ -time parameter and in the metastable, the Hamilton equation of motion are

$$\dot{u}^{i} = \frac{\partial H(u,p)}{\partial p_{i}} = 2\beta^{i}(u),$$

$$\dot{p}_{i} = -\frac{\partial H(u,p)}{\partial u^{i}} = -2\sum_{k=1}^{8N} \frac{\partial \beta^{k}(u)}{\partial u^{i}} p_{k}, \quad i,k = 1,...,8N,$$

where the time derivatives are taken respect to the non-compact slow time parameter  $\tilde{\tau}$ . Since this is the domain of interest for the observable  $U_{\tau}$  dynamics, we adopt this slow time for the description of the  $U_{\tau}$  dynamics. Note that our remark on the physical meaning of  $\tau$  still applies to the slow time  $\tilde{\tau}$ . Also note that the equations for  $u^i$  determine an autonomous dynamical system independent of the time  $\tau$ .

The above results can be compiled in the form of a

**Theorem 3.7.** For each dynamical system as given by the equations (3.33), there exists a Hamilton-Randers system whose Hamiltonian function is (3.26).

If the constraints  $y^k = \dot{x}^k, \, k = 1, ..., N$  hold, then the first equations of Hamilton imply

$$y^{k} = \dot{\beta}_{x}^{k}(x, \dot{x}, y, \dot{y}), \quad k = 1, ..., 8N.$$

It is remarkable and of relevance for the physical interpretation of the theory that the  $U_{\tau}$  dynamics defined in this way does not depend on the particular value of  $t \in (2n+1)T$ , (2n+3)T). This is of relevance for the definition od the evolution of densities and observables. Moreover, the dynamical systems that we are considering there is not a relation as in classical mechanics of the standard form  $\dot{x}^k = 1/m p^k$ , since the Hamiltonian function (3.26) that we are considering is linear on the canonical momentum variables.

3.10. Observers and metric structures. Since  $M_4$  is endowed with a Lorentzian metric  $\eta_4$  with signature (1, -1, -1, -1), there is natural definition of ideal, macroscopic observer,

**Definition 3.8.** An ideal, macroscopic observer is a timelike vector field  $W \in \Gamma TM_4$ ,  $\eta(W, W) > 0$ .

Given an observer W, there is associated a Riemannian metric on  $M_4$  given by the expression

$$\bar{\eta}_4(u,v) = \eta_4(u,v) - 2\frac{\eta_4(u,W(x))\eta_4(v,W(x))}{\eta_4(W(x),W(x))}, \quad u,v \in T_x M_4, x \in TM_4.$$

 $d_W: M_4 \times M_4 \to \mathbb{R}$  is the distance function associated with the Riemannian norm of  $\bar{\eta}_4$ . By the embeddings  $\varphi_k: M_4^k \to M_4$  each manifold  $M_4^k$  is diffeomorphic to the manifold of observable macroscopic events  $M_4$ . Then for each fixed value of the parameter  $t \in I$  one can also consider the embedding of the world lines of the sub-quantum molecules  $\{1, ..., N\}$  from  $M_4^k$  in  $M_4$ ,  $\varphi_k(\xi_{tk}) = \hat{\xi}_{tk} \hookrightarrow M_4$ ,

$$\hat{\xi}_t(k): \{t\} \times \mathbb{R} \to M_4, \quad (t,\tau) \mapsto (t, \hat{\xi}_{tk}(\tau)).$$

The manifolds  $M_4^k$  are endowed with Lorentzian metrics that are isometric to the Lorentzian metric  $\eta_4$  on  $M_4$ , for each  $k \in \{1, ..., N\}$ . Given a point  $x \in M_4$  and  $\xi_t(k)$ , the *distance function* between x and  $\hat{\xi}_t(k)$  is given by the expression

(3.35) 
$$d_4(x, \tilde{\xi}_t(k)) := \inf \left\{ d_W(x, \tilde{x}), \, \tilde{x} \in \tilde{\xi}_t(k) \right\}.$$

It is clear that  $\min\{d_4(x, \hat{\xi}_t(k)), k = 1, ..., N\}$  depends on  $t \in I$ . Such distance is realized, at the instant  $t \in [0, T]$ , by the sub-quantum molecule denoted by the integer  $\bar{k}$ ,

$$d_4(x, \xi_t(\bar{k})) = \min\{d_4(x, \xi_t(k)), k = 1, ..., N\}.$$

#### 4. HILBERT SPACE FORMULATION OF HAMILTON-RANDERS SYSTEMS

In order to provide a direct link between quantum mechanics and Hamilton-Randers theory, it is useful to formulate HR-systems using the same formalism than the one used in quantum mechanics, that is, the theory of Hilbert spaces and operators on Hilbert spaces. This formulation follow the lines proposed by Hooft [26].

4.1. Hilbert formulation of classical systems applied to HR-systems. Each HR-system has associated a Hilbert space and there is also a Hilbert space formulation of the dynamics. To show this correspondence, let us start considering the algebra of diagonal functions  $\mathcal{F}_D(T^*TM)$  and the canonical quantization of this algebra. Since in Hamilton-Randers theory the time parameter  $(t, \tau)$  is 2-dimensional, the specification of the commutation relations of the algebra is done at each fixed value of the pair  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . Therefore, following closely the procedure of standard canonical quantization, we adopt the following canonical quantization rules,

• The values of the position coordinates  $\{x_k^{\mu}, k = 1, ..., N, \mu = 1, 2, 3, 4\}$ and the velocity coordinates  $\{y_k^{\mu}, k = 1, ..., N, \mu = 1, 2, 3, 4\}$  of the sub-quantum molecules are the eigenvalues of certain self-adjoint linear operators on a given Hilbert space  $\mathcal{H}_{Planck}$ ,

$$\{\hat{x}_{k}^{\mu}, \hat{y}_{k}^{\mu}, \, \mu = 1, 2, 3, 4; \, k = 1, ..., N\}.$$

Hence the operators  $\{\hat{x}_k^{\mu}, \hat{y}_k^{\mu}, \mu = 1, 2, 3, 4; k = 1, ..., N\}$  are characterized by the relations

(4.1) 
$$\hat{x}_{k}^{\mu} |x_{l}^{\mu}, y_{l}^{\nu}\rangle = \sum_{l} \delta_{kl} x_{l}^{\mu} |x_{l}^{\mu}, y_{l}^{\nu}\rangle,$$
$$\hat{y}_{k}^{\nu} |x_{l}^{\mu}, y_{l}^{\nu}\rangle = \sum_{l} \delta_{kl} y_{l}^{\nu} |x_{l}^{\mu}, y_{l}^{\nu}\rangle,$$

where  $\delta_{kl}$  is the Kronecker delta function.

• There is a set of self-adjoint linear operators

$$\{\hat{p}_{\mu x_k}, \hat{p}_{\mu y_k}, k = 1, ..., N, \mu = 1, 2, 3, 4\}$$

that generates local diffeomorphism on TM along the integral curves of the local vector fields

$$\{\frac{\partial}{\partial x_k^{\mu}}, \frac{\partial}{\partial y_k^{\nu}} \in \Gamma TTM \quad \mu, \nu = 1, 2, 3, 4; \ k = 1, ..., N\}.$$

• The following commutation relations at each fixed 2-time  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$  hold good,

 $[\hat{x}_{k}^{\mu}, \hat{p}_{\nu x_{l}}] = \imath \hbar \, \delta_{\nu}^{\mu} \, \delta_{kl}, \quad [\hat{y}_{k}^{\mu}, \hat{p}_{\nu y_{l}}] = \imath \hbar \, \delta_{\nu}^{\mu} \, \delta_{kl}, \quad \mu, \nu = 1, 2, 3, 4, \ k = 1, ..., N.$ 

• The phase space  $T^*TM$  is commutative,

(4.3) 
$$[\hat{x}_k^{\mu}, \hat{x}_l^{\nu}] = 0, \quad [\hat{p}_{\mu k}, \hat{p}_{\nu l}] = 0, \quad \mu, \nu = 1, 2, 3, 4, \ k = 1, ..., N.$$

The collection

$$\{|x_l^{\mu}, y_l^{\nu}\rangle, \mu, \nu = 1, 2, 3, 4, l = 1, ...N\}$$

generates a complex vector space  $\bar{\mathcal{H}}_{Planck}$  that can be promoted to a complex pre-Hilbert space by defining the following scalar product,

(4.4) 
$$\langle x_l^{\mu}, y_l^{\nu} | x_k^{\rho}, y_k^{\lambda} \rangle = \delta_l^k \,\delta(x^{\mu} - x^{\rho}) \,\delta(y^{\nu} - y^{\lambda})$$

and extended by assuming bilinear property of the product for arbitrary linear combinations. Note the symmetric property of this product rule,

$$\langle x_l^{\mu}, y_l^{\nu} | x_k^{\rho}, y_k^{\lambda} \rangle = \langle x_k^{\rho}, y_k^{\lambda} | x_l^{\mu}, y_l^{\nu} \rangle.$$

Let us consider combinations of the form

$$\Psi = \sum_{k} \int_{\varphi_{k}^{-1}(T_{x}M_{4})} d^{4}z_{k} \, \alpha(\varphi_{k}^{-1}(x_{k}), z_{k}) \, |\varphi_{k}^{-1}(x), z_{k}\rangle.$$

In the case of simple systems, associated with free particles, one expects that the fundamental cycles are indeed exactly modular. In this case, the functions on  $\Psi$  must be *T*-module invariant. The natural way to satisfy this is by complex modularity of the coefficients  $\alpha(\varphi_k^{-1}(x_k), z_k)$ , in accordance with the congruence module *n* idea described in *subsection* 3.7. Hence the general combinations of fundamental states  $|\varphi_k^{-1}(x), z_k\rangle$  that we should consider are complex,  $\alpha(\varphi_k^{-1}(x_k), z_k) \in \mathbb{C}$ .

The norm of  $\Psi$  is calculated by using the product rule and bilinear extension of (4.4). The norm coincides with the  $L_2$  norm of  $\Psi$ ,

$$\langle \Psi | \Psi \rangle = \| \Psi \|_{L_2} = \sum_{k=1}^N \int_{M_4} dvol_{\eta_4} |\alpha|^2.$$

The linear closure of the states with finite norms is denoted by  $\mathcal{H}_{Planck}$  and with the scalar product associated to the  $L_2$  norm,  $\mathcal{H}_{Planck}$  is a complex pre-Hilbert space.

4.2. Quantum Hamiltonian associated to a HR-system. By applying the canonical quantization of the Hamiltonian (3.26) one obtains the operator

$$H_t(\hat{u}, \hat{p}) : \mathcal{H}_{Planck} \to \mathcal{H}_{Planck}$$

given by the expression

(4.5) 
$$\widehat{H}_t(\hat{u}, \hat{p}) := (1 - \kappa(t)) \frac{1}{2} \Big( \sum_{k=1}^N \left( \beta^k(\tau, \hat{u}) \hat{p}_k + \hat{p}_k \beta^k(\tau, \hat{u}) \right) \Big),$$

with  $(t,\tau) \in [0,T] \times \mathbb{R}$ , k = 1, ..., 8N.  $\widehat{H}_t(\tau, u, p)$  is uniquely defined, Hermitian and is determined by elements from the classical version of a HR-system.

4.3. Heisenberg dynamics of the ontological states is the classical dynamics. The Heisenberg's equations associated with  $\hat{x}_k^{\mu}$  and  $\hat{y}_k^{\mu}$  reproduces the Hamiltonian equations for the slow time  $\tau$ -time parameter. Let us consider the Heisenberg representation for the  $U_{\tau}$  dynamics, where the Hermitian operators evolve in the  $U_{\tau}$  time by the Heisenberg equations determined by the Hamiltonian (4.5). Then

**Theorem 4.1.** The Heisenberg's equations associated with  $\hat{x}_k^{\mu}$  and  $\hat{y}_k^{\mu}$  reproduces the Hamiltonian equations for the slow time  $\tau$ -time parameter.

The proof of this result is direct if we take into account the definition of slow time (3.30), the canonical theory developed in this section (the commutations relations (4.2) and (4.3)) and the structure of the Hamiltonian (4.5). These line of thought is formally similar to the original pictured developed in the work of G. 't Hooft [26].

Moreover, we consider the constraints

(4.6) 
$$\hat{y}_k^{\mu} = \frac{d\hat{x}_k^{\mu}}{d\tau}, \quad \mu = 1, 2, 3, 4; \quad k = 1, ..., N.$$

These constraints define the *on-shell evolution* of the sub-quantum molecules. However, the constraints (4.6) are not strictly necessary from the point of view of the Hamiltonian dynamics.

We need to emphasize that adopting (4.1) the dynamical picture has been chosen, namely, the Heisenberg picture of dynamics is adopted in this theory as preferable to describe the dynamics of the ontological degrees of freedom. In these picture, the ket space of sub-quantum states  $\{|x_l^{\mu}, y_l^{\mu}\rangle\}_{k=1,\mu=1}^{N,4}$  do not change on  $\tau$ -time, but are the operators associated to the quantized generalized position and canonical momenta that change. An state  $\Psi \in$  $\mathcal{H}_{Planck}$  does not change on time  $\tau$  in this picture, while operators that are functional dependent of the canonical operators  $\{\hat{u}_k^{\mu}, \hat{p}_k^{\mu}, \mu = 1, 2, 3, 4, N =$  $1, ..., N\}$  change on  $\tau$ -time as the Heisenberg equation dictates. Also note that although in the usual Heisenberg picture of the dynamics the whole Hilbert space generated by the ket vectors changes with time [14], in our case the space  $\mathcal{H}_{Planck}$  is the same. Still are the operators and not the fundamental states what change on time, following the relation described in *Theorem* 4.1. This is in concordance with the fact that we are considering

26

a four dimensional representation from the beginning, although it does not follow from that fact.

Finally, let us note that the meta-stability condition (3.27) is translated in the quantum formulation of HR-systems as

(4.7) 
$$\lim_{t \to (2n+1)T} \widehat{H}_t(\hat{u}, \hat{p}) |\Psi\rangle = 0, \quad n \in \mathbb{N}.$$

4.4. Sub-quantum operators and quantum operators. The commutation relations (4.2) are mathematically and physically consistent, despite the fact that in quantum mechanics the canonical coordinate operators  $\{\hat{X}^a, a = 2, 3, 4\}$  do not commute in general with the velocity operators  $\{\hat{X}^a, a = 2, 3, 4\}, [\hat{X}^a, \hat{X}^a] \neq 0$ . In order to clarify this point, let us remark that the operators  $\{(\hat{x}^a_k, \hat{y}^a_k)\}_{k=1,a=2}^{N,4}$  do not coincide with the quantum operators  $\{\hat{X}^a, \hat{X}^a = 2, 3, 4\}$ . By definition, the operators  $\{\hat{X}^a, \hat{X}^a = 2, 3, 4\}$ have as spectrum the possible outcomes of measurements on a quantum system for the cartesian position coordinates and the corresponding speeds.

In a complete theory, the operators  $\{\hat{X}^a, \hat{X}^a, a = 2, 3, 4\}$  should emerge in HR-systems together with the wave function for the quantum state, they constitute a phenomenological description of the ergodic dynamics of the sub-quantum molecules, when the projection  $(t, \tau) \mapsto \tau$  is considered. The exact relation between the emergent operators  $\{\hat{X}^a, \hat{X}^a, a = 2, 3, 4\}$  and the entire family of operators  $\{(\hat{x}^a_k, \hat{y}^a_k, \hat{p}^a_{kx_k}, \hat{p}^a_{ky_k})\}_{k=1,a=2}^{N,4}$  is still missing, but we expect the following generic fact. The quantum states obtained from HR-systems are generically non-localized in both position and speed (or canonical momentum) and both, the operators  $\hat{X}^a$  and  $\hat{X}^a$  have generically non-zero dispersion when applied to physical states. Therefore, the quantum states must hold a representation of a non-commutative algebra,

(4.8) 
$$[\hat{X}^{\mu}_{a}, \hat{X}^{\nu}_{b}] = A^{\mu\nu} \,\delta_{ab}, \quad [\dot{\hat{X}}^{\mu}_{a}, \dot{\hat{X}}^{\nu}_{b}] = B^{\mu\nu} \,\delta_{ab}, \quad [\hat{X}^{\mu}_{a}, \dot{\hat{X}}^{\nu}_{b}] = C^{\mu\nu} \,\delta_{ab},$$

with  $\mu, \nu = 1, 2, 3, 4, a, b = 1, ..., N$ . This algebra contrasts with the related commutative algebra (4.3). Moreover, it is evident that for our theory it is more natural to have a non-commutative spacetime where  $A^{\mu\nu} \neq 0$  holds than a commutative spacetime and that the commutativity assumption of the manifold  $M_4$  must be considered an approximation. Hence our theory must be associated naturally to generalized uncertainty relations.

# 5. Quantum mechanical elements from Hamilton-Randers systems

5.1. Quantum Hilbert space from HR-systems. It is of relevance to relate a generic quantum mechanical wave functions with the ontological states  $\{|x_k^{\mu}, y_k^{\mu}\rangle\}_{k=1,\mu=1}^{N,4}$ . First, we consider the state  $\Psi \in \mathcal{H}_{Planck}$  given by

(5.1) 
$$\Psi(u) = \sum_{k=1}^{N} e^{i \vartheta_{k\Psi}(\varphi_{k}^{-1}(x), z_{k})} \frac{1}{\sqrt{N}} n_{k\Psi}(\varphi_{k}^{-1}(x), z_{k}) |\varphi_{k}^{-1}(x), z_{k}\rangle.$$

We call these specific combinations of ontological states *predecessor states*. Our ansatz is that any quantum state  $\psi$  is obtained from a predecessor state by averaging speed components,

(5.2)  

$$\psi(x) = \sum_{k=1}^{N} \int_{\varphi_{k*}^{-1}(T_{x}M_{4})} d^{4}z_{k} e^{i\vartheta_{k\Psi}(\varphi_{k}^{-1}(x),z_{k})} \frac{1}{\sqrt{N}} n_{k\Psi}(\varphi_{k}^{-1}(x),z_{k}) |\varphi_{k}^{-1}(x),z_{k}\rangle.$$

Hence a quantum state is a class of equivalence respect to averaging along speed components. Many predecessor states have associated the same equivalence class  $\psi$ . The physical significance of the averaging operation along the speed components comes from the ergodicity property of the  $U_t$ -flow in the ergodic dynamics. Once fixed the point x, the ergodic theorem implies that averaging on time t along a semi-period [0, T] is equivalent to average along the space of speeds.

Let us make the assumption that the collection of product of local phases

$$\left\{ e^{i \vartheta_{kA}(\varphi_k^{-1}(x), z_k)} e^{-i \vartheta_{kB}(\varphi_k^{-1}(x), z_k)}, \quad x \in M_4, \, z_k \in T_{\varphi_k^{-1}(x)} M_4^k \right\}$$

are highly oscillating compared to an universal characteristic frequency  $1/T_{min}$ . This defines a subset of averaged quantum spaces. Formally, this highly oscillating property is equivalent to

(5.3) 
$$\int_{\varphi_{k*}^{-1}(T_x M_4)} d^4 z_k \, e^{i \,\vartheta_{kA}(\varphi_k^{-1}(x), z_k)} \, e^{-i \,\vartheta_{kB}(\varphi_k^{-1}(x), z_k)} \equiv \,\delta_{AB},$$

where if  $A \neq B$ , then  $\delta_{AB} = 0$  and if A = B it is defined to be equivalent to  $\delta_{AB} = 1$ . What this last statement means is that the measure  $d^4 z_k$  is such that on any normalized test function  $\chi \in T_{\varphi_k^{-1}(x)}M_4$ ,

$$\int_{\varphi_{k*}^{-1}(T_x M_4)} d^4 z_k \, |\chi(\varphi_k^{-1}(x), z_k)|^2 = 1$$

From a physical point of view, condition (5.3) is a strong form of locality for the quantum states defined on the spacetime manifold  $M_4$ . The phases of the quantum state  $\psi$  contains information of the sub-quantum degrees of freedom location in the corresponding four-manifold  $M_4^k$ .

In terms of the ontological states  $\{|x_k^{\mu}, y_k^{\mu}\rangle\}_{k=1,\mu=1}^{N,4}$  and using the product rule (4.4), one finds the following expression for the isometrically induced scalar product from  $(\mathcal{H}_{Planck}, \langle \rangle)$ ,

(5.4) 
$$\langle \psi_A | \psi_B \rangle = \sum_{k=1}^N \frac{1}{\sqrt{N}} \int_{M_4} \int_{\varphi_{k*}^{-1}(T_x M_4)} dvol_{\eta_4} \wedge d^4 z_k \, e^{\imath \vartheta_{kA}(\varphi_k^{-1}(x), z_k)} \\ e^{-\imath \vartheta_{kB}(\varphi_k^{-1}(x), z_k)} \cdot n_{Ak}(\varphi_k^{-1}(x), z_k) \, n_{Bk}(\varphi_k^{-1}(x), z_k).$$

After a short calculation and using the product orthogonality condition for the ontological states (4.4), the following orthogonality condition for the wave functions (5.2) is obtained,

(5.5) 
$$\langle \psi_A | \psi_B \rangle = 0, \quad \text{if } A \neq B.$$

Moreover, the same procedure implies the normalization rule

**N** 7

(5.6) 
$$\langle \psi | \psi \rangle = \int_{M_4} dvol_{\eta_4} \sum_{k=1}^N \int_{\varphi_{k*}^{-1}(T_x M_4)} d^4 z_k \ n_k^2(\varphi_k^{-1}(x), z_k) = 1.$$

The complex, linear bulk generated by finite combinations of fundamental states  $\psi$  is a complex vector space  $\mathcal{H}$ . Note that because the elements of  $\mathcal{H}$  have finite norm, they are normalizable:

(5.7) 
$$\|\psi\|_{L_2} = 1.$$

**Proposition 5.1.** The space  $\mathcal{H}$  with the  $L_2$  norm is a complex Hilbert space.

*Proof.* The set  $\mathcal{H}$  with the product

$$\langle \psi | \chi \rangle := \int_{M_4} dvol_{\eta_4} \, \psi^* \, \chi$$

is a pre-Hilbert space. Moreover, since the elements of  $\mathcal{H}$  are finite combination of ontological states, that define a Hilbert space, any Cauchy sequence of  $\psi$  can be decompose in a finite sum of Cauchy sequences of ontological states, hence it is a convergent sequence.

5.2. Representations of the sub-quantum degrees of freedom. From the way the space  $\mathcal{H}$  it is defines there is an embedding

$$(\mathcal{H},\langle,\rangle^*) \hookrightarrow (\mathcal{H}_{Planck},\langle,\rangle)$$

holds. By applying linearity of the product  $\langle,\rangle$  and the above isometric embedding, one obtains the relation

(5.8) 
$$\langle x_k, z_k | \psi \rangle = e^{i \vartheta(\varphi_k^{-1}(x), z_k)} n_l(\varphi_k^{-1}(x), z_k)$$

This expression is analogous to the quantum mechanical relation between the space representation and the momentum representation in quantum mechanics,

$$\langle \vec{x} | \vec{p} 
angle = e^{i \, \vec{x} \cdot \vec{p}} \frac{1}{(2\pi \, \hbar)^{3/2}}.$$

This representation is obtained in quantum mechanics from the coordinate representation of the translation operator, by solving a simple differential equation [14] and determines an unitary transformation.

5.3. Emergence of the Born rule in the Heisenberg representation of the dynamics. The density squared of the fundamental wave function  $n^2(x)$  is given by the expression

(5.9) 
$$n^{2}(x) = \sum_{k=1}^{N} \int_{\varphi_{k*}^{-1}(T_{x}M_{4})} d^{4}z_{k} n_{k}^{2}(\varphi_{k}^{-1}(x), z_{k}).$$

 $n_k^2(\varphi_k^{-1}(x), z_k)$  is the number of world lines passing at  $\varphi_k^{-1}(x) \in M_4^k$  with velocity speed vector  $z_k$ . Therefore,  $n^2(x)$  is the *total density* of world-lines of sub-quantum molecules close to the point x. This is in accordance with our definition of density of lines, showing that the ansatz (5.2) provides a statistical interpretation of a quantum state in terms of sub-quantum degrees of freedom degrees. It is remarkable that such statistical interpretation applies to a pure state  $\psi \in \mathcal{H}$  that describes an individual quantum system.

The density of lines is normalized by the condition

$$\int_{M_4} n^2(x) \, dvol_{\eta_4} = N$$

and since in our models the number of degrees of freedom  ${\cal N}$  are preserved, we have that

$$\frac{d}{d\tau} \left( \int_{M_4} n^2(x) \, dvol_{\eta_4} \right) = 0.$$

The derivative can be pass inside the integration,

$$\frac{d}{d\tau} \left( \int_{M_4} n^2(x) \, dvol_{\eta_4} \right) = \int_{M_4} \frac{d}{d\tau} \left( n^2(x) \, dvol_{\eta_4} \right)$$
$$= \int_{M_4} \frac{d}{d\tau} \left( n^2(x) \right) \, dvol_{\eta_4}$$
$$= \int_{M_4} \left( \frac{\partial n^2(x)}{\partial x^0} - \vec{v} \vec{\nabla} \, n^2(x) \right) \, dvol_{\eta_4}$$

Recall that we are using the Heisenberg picture of the dynamics. Hence the states  $\psi(x)$  given by the expression (5.2) do not change with the time parameter  $\tau$ . Moreover, the density  $n^2(x)$  can be written as the square of the wave function,

(5.10) 
$$n^2(x) = \langle \psi | \psi \rangle(x),$$

by the relations (5.6) and (5.9). If we associate  $n^2(x)$ , that is a probability of presence, with the probability density to find the particle at  $x \in M_4$ , then the Born rule is obtained and not postulated from dynamical and statistical considerations in Hamilton-Randers theory, in the Heisenberg picture of the dynamics.

Physical interpretation of the density. Equipped with the distance structure (3.35) we can interpret the density  $n^2(x)$  as follows. Let us consider an arbitrary point  $x \in M_4$ . By the ergodic theorem, the density of lines  $n^2(x)$  is the number of world lines  $\varphi_k(\xi_{tk}) : I \to M_4$  that pass close to x for a fixed internal time t(n) on each fundamental cycle  $t(n) \in [(2n + 1)T, (2n + 3)T]$ . We define to pass close to x if it image on  $M_4$  by the isometry  $\varphi_k : M_4^k \to M_4$  is in the interior of an open set  $\mathcal{U}(x, L_{min}, \tilde{L})$  whose points are at a distance  $d_W$  between  $\mathcal{L}_{min}$  and  $\tilde{L}$  from the point  $x \in M_4$ .  $\tilde{L}$  is large compared with  $\mathcal{L}_{min}$  but small compared with any macroscopic observable variation in coordinates. We assume that there is independence in the definition of  $n^2(x)$  respect to  $\tilde{L}$ . Note that in this interpretation we have assumed that a local concentration around each  $x \in M_4$  in the given open set  $\mathcal{U}(x, L_{min}, \tilde{L})$  happens.

This construction is not  $\text{Diff}(M_4)$ -invariant. This property is consistent with usual models in quantum mechanics, which are also not invariant. However, the existence of measures that are  $\text{Diff}(M_4)$ -invariant in principle allows to construct invariant theories.

5.4. Emergence of the  $\tau$ -time diffeomorphism invariant constraint. The relation (4.7) is a constraint on  $\mathcal{H}_{Planck}$ . Since  $\mathcal{H} \subset \mathcal{H}_{Planck}$ , the same constrain could be applied to the sub-class of elements in  $\mathcal{H}$  such that (4.7) holds good. Such constraint on the Hilbert space  $\mathcal{H}$  is a quantum version of the  $\tau$ -time diffeomorphism invariant condition (3.27),

30

**Proposition 5.2.** For any physical state  $|\psi\rangle \in \mathcal{H}$  the constraint

(5.11) 
$$\lim_{t \to (2n+1)T} \widehat{H}_t(u,p) |\psi\rangle = 0$$

holds good.

This constraint holds periodically in the parameter t, with a periodicity 2T. One could tempted to think that the constrain (5.11) has some relation with the Wheeler-DeWitt equation. However, this is not the case, since the constraint (5.11) only holds in the metastable regime t = (2n+1)T and not for the whole evolution. Indeed, this suggests that  $\tau$ -time reparetrization invariance only in that regime. From here it follows the emergent nature of the  $\tau$ -time re-parameterizations invariance, according with the also emergent nature of any physical  $\tau$ -time parameter.

### 6. Concentration of measure and measurement processes in Hamilton-Randers systems

6.1. Concentration of measure. In this *section* we discuss the relation between a general type of mathematical phenomena that appears in functional analysis and metric geometry known as *concentration of measure* [24, 34, 42] from one side and Hamilton-Randers theory from the other side. The concentration of measure is a general property of regular enough functions in high dimensional topological spaces endowed with a metric and measure structures. In a nut-shell, the phenomena of concentration of measure for the category of topological spaces admitting a notion of dimension can be stated as follows,

In a measure metric space of large dimension every real 1-Lipschitz function of many variables is almost constant almost everywhere.

The formalization of concept of concentration makes use of both, the metric properties and the measure properties, providing a precise meaning of *almost constant* and *almost everywhere*. Remarkably, both concepts, the measure structure and the metric structure are independent from each other. Hence concentration happens in the category of Gromov mm-spaces [24] and for 1-Lipschitz functions.

In a general measure metric space  $(\mathbf{T}, \mu_P, d)$ , the concentration function

$$\alpha(\mu_P): \mathbb{R} \to \mathbb{R}, \quad \rho \mapsto \alpha(\mu_P, \rho),$$

defined by the condition that  $\alpha(\mu_P, \rho)$  is the minimal real number such that

(6.1) 
$$\mu_P(|f - M_f| > \rho) \le 2\,\alpha(\mu_P, \rho)$$

for any real function  $f : \mathbf{T} \to \mathbb{R}$ .  $M_f$  is the median or Levy's mean of f, which is defined as the value attained by  $f : \mathbf{T} \to \mathbb{R}$  such that

$$\mu_P(f > M_f) = 1/2$$
 and  $\mu_P(f < M_f) = 1/2$ .

Therefore, the probability that the function f differs from the median  $M_f$ in the sense of the defined measure  $M_f$  by more than the given value  $\rho \in \mathbb{R}$ is bounded by the concentration function  $\alpha(P, \rho)$ . **Example 6.1.** A typical example of concentration of measure is provided by the concentration of measure in spheres  $\mathbb{S}^N \subset \mathbb{R}^{N+1}$ . Let  $(\mathbb{S}^N, \mu_S, d_S)$  be the N-dimensional sphere with the standard measure and the round metric distance function. As a consequence of the isoperimetric inequality [34] it holds that for each  $A \in \mathbb{S}^N$  with  $\mu(A) \geq 1/2$  and  $\epsilon \in (0, 1)$ , the set

$$A_{\epsilon} := \{ x \in \mathbb{S}^N \ s.t. \ d_S(x, A) \le \epsilon \}$$

is such that

$$\mu_{\mathbb{S}}(A_{\epsilon}) \le 1 - \sqrt{\pi/8} \exp(-\epsilon^2 (N-1)/2)$$

In particular, Levy's lemma holds: and therefore, for any smooth function

$$\mu_{\mathbb{S}}(A) := \{ x \in \mathbb{S}^N \ s.t. \ f(x) = M_f \} \ge 1/2$$

the concentration inequality

(6.2) 
$$\mu_{\mathbb{S}}(A_{\epsilon}) \leq 1 - \sqrt{\pi/2} \exp(-\epsilon^2 (N-1)/2).$$

holds.

The direct significance of the relation (6.2) is that for high dimensional spaces  $N \to +\infty$  and for each  $\epsilon \in (0,1)$  (note that the radius of the sphere is normalized, such that  $\epsilon = 1$  is the maximal distance between points in the sphere), then almost for all the points on the sphere (that is, module a set of measure zero with  $\mu_S$ ) the limit

(6.3) 
$$\lim_{N \to \infty} \mu(A_{\epsilon}) = 1$$

holds good. That is, the function f must be almost constant on  $\mathbb{S}^N$ , concentrating its value at the median  $M_f$ . In particular, for the sphere  $\mathbb{S}^N$  the concentration of 1-Lipschitz functions is of the form

(6.4) 
$$\alpha(P_M,\rho) \le C \exp\left(-\frac{(N-1)}{2}\epsilon\right),$$

with C a constant of order 1.

**Example 6.2.** The second example of concentration that we consider here is for 1-Lipschitz functions on  $\mathbb{R}^N$  (compare with [42], pg. 8). Then there is concentration, determined by the concentration inequality

(6.5) 
$$\mu_P(|f - M_f| > \rho) \le \frac{1}{2} \exp\left(-\frac{\rho^2}{2\rho_P^2(f)}\right),$$

where we have adapted the example from [42] to a Gaussian measure  $\eta_P$  with median  $M_f$  and standard deviation  $\rho_P(f)$ .

For 1-Lipschitz functions on a measure metric space  $\mathbf{T}$  of dimension N there are similar *exponential bounds* as for  $\mathbb{R}^N$  and other manifolds locally homeomorphic to  $\mathbb{R}^N$ . In such cases, the concentration of measure phenomena is an universal feature, which is a consequence of the Lipschitz regularity condition of the function  $f : \mathbf{T} \to \mathbb{R}$  and the higher dimensionality of the space  $\mathbf{T}$ . For dim(M) large, the concentration of 1-Lipschitz functions implies that the values of the function are very picked around a certain constant value.

Heuristically, one can understand the phenomena of concentration of measure as partially originated by the constraint that the 1-Lipschitz regularity condition imposes on f viewed globally. Let  $f : \mathbf{T} \to \mathbb{R}$  be a 1-Lipschitz function on a normed topological space  $(\mathbf{T}, \|, \|_{\mathbf{T}})$  locally homeomorphic to a given  $\mathbb{R}^N$ . The Lipschitz condition is a form of *equipartition* of the variations of f caused by arbitrary variation on the points on the topological space  $\mathbf{T}$ . Hence when the space  $\mathbf{T}$  is highly dimensional, the significance of the 1-Lipschitz condition is that the function f cannot admit large *standard variations* caused by the corresponding standard variations along any of the independent variables. Otherwise, a violation of the Lipschitz condition can happen, since the large dimension provides long contributions to the variation of f. Note that to speak of large and small variations, one needs to have a standard reference.

6.2. Spontaneous reduction of the quantum state as concentration of measure phenomena. Let us consider the Euclidean space  $\mathbb{R}^q$  with  $q \gg 1$ ,  $\mu_P$  the standard product measure in  $\mathbb{R}^q$  and  $f : \mathbb{R}^q \to \mathbb{R}$  a real 1-Lipschitz function. Then there is concentration determined by the inequality

(6.6) 
$$\mu_P\left(\left|f - M_f\right| \frac{1}{\sigma_f} > \frac{\rho}{\rho_P}\right) \le \frac{1}{2} \exp\left(-\frac{\rho^2}{2\rho_P^2}\right)$$

where we have adapted the example from [42] to a Gaussian measure  $\mu_P$  and standard deviation  $\rho_P$  and where the median of f is  $M_f$ .  $\sigma_f$  has the physical interpretation of being the minimal physical resolution of any measurement of observable operators associated to the 1-Lipschitz function f. By applying this concentration to a generic function  $f \in \mathcal{F}_D(T^*TM)$  in the domain where the flow  $U_t$  is 1-Lipschitz, one obtains

$$U_t: \mathcal{F}_D(T^*TM) \to \mathcal{F}_D(T^*TM), \quad f(t=0) \mapsto f(t);$$

 $U_t$  could depend on the  $\tau$ -time parameter. We assume that for f there is a minimal resolution  $\sigma_f$ . In the 1-Lipschitz dynamical regime of the dynamics operator  $U_t$  the function f must be constant almost everywhere, since f is 1-Lipschitz in (u, p) and  $(t, \tau)$ . Moreover, for macroscopic observations associated to any measurement of a quantum system, one expect the existence of scales such that the relation

(6.7) 
$$|f - M_f| \frac{1}{\sigma_f} \simeq N \quad \text{or} \quad |f - M_f| \frac{1}{\sigma_f} \gg N, \quad 1 \ll N,$$

holds. This relation, for what concentration concerns, can be re-written as

(6.8) 
$$\frac{\rho^2}{\rho_P^2} = N^2$$

This condition provides a quantum scale relative to the Planck scale. This is related with ratio between the GUT scale energy and the Planck scale energy. In this context, we make the hypothesis that the GUT scale is the higher energy scale where the dynamics is well described by a quantum gauge field theory. Therefore, let us restrict our considerations to the case when the quantum system corresponds to a elementary particle which quantum field operator appears in the Standard Model. If we adopt such relation, then  $N^2$  is large for the purposes of the application of the concentration phenomena. We see from (6.8) that  $N \in \mathbb{N}$  provides a measure of the *complexity* of the associated HR-system compared with the complexity of the associated quantum system. The degree of complexity of a quantum state is of order 1, since there is one quantum particle involved compared with the degree of complexity of the associated HR-system, which is of order N. This order of complexity 1 is of the same order than the dimension of the model spacetime manifold  $M_4$ , the number of spin degrees of freedom and other quantum numbers associated with the quantum mechanical description of the elementary quantum particle.

Let us consider the case  $TM \cong \mathbb{R}^{8N}$ . By the relation (6.7), the concentration of measure relation (6.6) is applied to the function f in the 1-Lipschitz dominated regime of  $U_t$  obtaining

(6.9) 
$$\mu_P\left(|f - M_f| > \sigma_f(8N)\right) \le \frac{1}{2} \exp\left(-32N^2\right).$$

Note that although the value  $(8N)^2$  is large, the quotient  $|f - M_f|/\sigma_f$  is larger or the same order of magnitude. Indeed,  $\sigma_f$  is the minimal theoretical resolution for f. Extrapolation of how much  $\sigma_f$  when measuring systems at the Planck scale and how much it could be the difference  $|f - M_f|$  when applied to standard model scale to GUT scale systems convince us that (6.9) holds. Hence the concentration relation (6.9) is a strong condition. In our case, the GUT scale, defined in our case by the conditions

$$(6.10) \qquad \qquad |f - M_f| > \sigma_f (8N)$$

provides us with the limit where the inequality starts to loss concentration meaning.

Hence for functions that are associated with measurements of the properties of quantum systems and since  $N \gg 1$  there must be concentration of measure around the median  $M_f$  with probability very close to 1. Thus, if a measurement of an observable associated with f is performed, the result  $M_f$ is with probability very close to 1 and in practice, equal to 1 in the regime where the evolution  $U_t$  evolution is Lipschitz.

Notion of classical and quantum interaction. Depending on the domain where an interaction is dominant, one can make a clear distinction between *classical* and *quantum interactions*, that can serve as definition.

**Definition 6.3.** A classical interaction in Hamilton-Randers theory is an interaction which is dominant only on the metastable domain  $\{t = (2n + 1)T, n \in \mathbb{Z}\}$ . In contrast, a quantum interaction in Hamilton-Randers theory is dominant at least in the interior of the fundamental cycles  $\cup_{n \in \mathbb{Z}} (2n + 1, 2n + 3)$ .

A quantum interaction, since it is extended through the whole cycles, after the projection  $(t, \tau) \mapsto \tau$  is associated with density probability functions and transition amplitudes. Hence it is sensible to speak of *probability density for a given interactions*. A main characteristic of such interactions is that, although they are causal, the description is not local. Example of non-local interactions are gauge theories, whose exact quantum mechanical description is through holonomy variables. The Aharonov-Bohm effect is an example of non-local behaviour of an interaction. Hence we should be

34

open to the possibility that quantum gauge theories described in terms of non-local holonomy variables are emergent interactions.

In contrast, a classical interaction as defined above is local, since it is related to the domain where all the properties of the systems are well defined. A possible example of a classical interaction is gravity, as is discussed in *section* 7.

6.3. Notion of natural spontaneous reduction. The concentration of measure for functions on the Lipschitz regime provides a natural mechanism for the reduction of the quantum state. However, such reduction processes not only happen when the quantum system is being measured by a macroscopic observer. On the contrary, they are spontaneous processes happening at the final domain of each fundamental cycle in the  $U_t$ -evolution. This is the reason of the name natural reduction processes, in contrast with induced reduction of the quantum state by an interaction with a quantum measurement system [13, 20, 37]. A natural reduction process do not destroy the quantum system and it is dissociated from quantum measurement process. In a measurement process the measurement device can change the original quantum state, since there exists an interaction between the system being measured and the apparatus measurement. This is described through Von Neumann models in a quantum mechanical setting.

Note that the interaction responsible for this concentration is purely classical, since it is dominant only in the metastable equilibrium regime  $\{t = (2n + 1)T, n \in \mathbb{N}\}$ , when all the properties of the system are localized by the effect to the same interaction  $U_t$ . As a consequence of this phenomena the properties of the system appear as macroscopically well defined when the system is measured by means of a macroscopic measurement device. In Hamilton-Randers theory there is no entanglement between the measurement device and the quantum system: both have an exact localized state when they interact and the macroscopic measurement device do not induces the reduction of the quantum system.

If the initial conditions of the Hamilton-Randers system are fixed, then the evolution of  $M_f$  is fixed as well. However, it is difficult to determine initial conditions for HR-systems in general. It is also difficult to fix operationally  $\tau_0$  and hence, the long time dynamics of the average value  $M_f$ . Thus although the dynamics is deterministic, if we adopt assumption A.9. in section 2, the system is chaotic and hence unstable and sensitive to initial conditions. In this case, the natural way to describe the systems is by probabilistic methods, with the probability distribution determined by the fundamental dynamics  $U_t$  of the system during the ergodic regime. In particular, the probability function is associated with the density of world lines of sub-quantum molecules.

**Emergence of classicality.** We observe that in the metastable domain associated with the dominance in the dynamics of 1-Lipschitz operators every 1-Lipschitz observable have well defined values at each  $\tau_n = (2n+1)T$ . Under the assumptions that are discussed in the next *section*, the inertial position and speed observable quantities are 1-Lipschitz. The role of uniform bound of the sub-quantum molecules is fundamental to prove the 1-Lipschitz condition without further restrictions for macroscopic observables

that depend smoothly of the coordinates  $(x_k^{\mu}, y_k^{\mu})$ . As a consequence of this argument the trajectory of the quantum system is defined at each  $\tau_n$ . Hence one identifies the metastable equilibrium regime with the classical limit.

## 7. Hamiltonian bound and emergence of the weak equivalence principle and gravitational interaction

7.1. Bound of the matter Hamiltonian. In the following we show that in Hamilton-Randers theory a natural mechanism exists that ensures the existence of a lower bound for the energy level for the total Hamiltonian (4.5). Let us consider the decomposition of  $\hat{H}_t$  in a 1-Lipschitz component  $\hat{H}_{Lipschitz,t}$  and a non-Lipschitz component  $\hat{H}_{matter,t}$ ,

(7.1) 
$$\widehat{H}_t(\hat{u}, \hat{p}) = \widehat{H}_{matter,t}(\hat{u}, \hat{p}) + \widehat{H}_{Lipschitz,t}(\hat{u}, \hat{p}).$$

We have defined in this expression the matter Hamiltonian as precisely the piece which is not 1-Lipschitz. This is according with the idea that matter (including gauge interactions) is quantized. If matter is quantized, the spacetime distribution of energy, by definition, is presented in the form of quantum packets, which is a form of non-regular distribution.

This type of decomposition is not unique and also it is not evident that it exists. However, one can obtain a characterization that allows for such type of decompositions, if we adopt several additional assumptions. In particular, let us assume that the classical Hamiltonian function (3.26) is  $C^2$ -smooth in the variables  $(x, y, p_x, p_y)$ . In this case, we can prove the following

**Lemma 7.1.** Let  $H_t : T^*TM \to \mathbb{R}$  be a  $\mathcal{C}^2$ -smooth Randers Hamiltonian function (4.5). Then there exists a compact domain  $K \subset TM$  where  $H|_K$  is 1-Lipschitz.

*Proof.* By Taylor's expansion up to second order one has

$$H_{t}(u,p) = H_{t_{0}} + \sum_{k=1}^{8N} \frac{\partial H_{t}}{\partial u^{k}}|_{(\xi,\chi)}(u^{k} - \xi_{t}^{k}) + \sum_{k=1}^{8N} \frac{\partial H_{t}}{\partial p_{k}}|_{(\xi,\chi)}(p_{k} - \chi_{k})$$
$$+ \sum_{k=1}^{8N} R_{k} (u^{k} - \xi_{t}^{k})^{2} + \sum_{k=1}^{8N} Q_{k} (p_{k} - \chi_{k})^{2}$$
$$= H_{t_{0}} + \sum_{k=1}^{8N} \frac{\partial H_{t}}{\partial u^{k}}|_{(\xi,\chi)}(u^{k} - \xi^{k}) + \sum_{k=1}^{8N} \beta^{k}(\chi) (p_{k} - \chi_{k})$$
$$+ \sum_{k=1}^{8N} R_{k} (u^{k} - \xi^{k})^{2} + \sum_{k=1}^{8N} Q_{k} (p_{k} - \chi_{k})^{2}.$$

Hence difference for the values of the Hamiltonian  $H_t$  at two different points can be written formally as

$$\begin{aligned} |H_t(u(1), p(1)) - H_t(u(2), p(2))| &= \Big| \sum_{k=1}^{8N} \frac{\partial H_t}{\partial u^k} |_{(\xi, \chi)} (u^k(1) - \xi^k) \\ &+ \sum_{k=1}^{8N} \beta^k(\chi) \left( p_k(1) - \chi_k \right) + \sum_{k=1}^{8N} R_k(1) \left( u^k(1) - \xi^k \right)^2 + \sum_{k=1}^{8N} Q^k(1) \left( p_k(1) - \xi_k \right)^2 \\ &- \sum_{k=1}^{8N} \frac{\partial H_t}{\partial u^k} |_{(\xi, \chi)} (u^k(2) - \xi^k) - \sum_{k=1}^{8N} 2 \beta^k(\chi) \left( p_k(2) - \chi_k \right) \\ &- \sum_{k=1}^{8N} R_k(2) \left( u^k(2) - \xi^k \right)^2 - \sum_{k=1}^{8N} Q^k(2) \left( p_k(2) - \chi_k \right)^2 \Big| \\ &\leq \Big| \sum_{k=1}^{8N} \frac{\partial H_t}{\partial u^k} |_{(\xi, \chi)} (u^k(1) - u^k(2)) \Big| + \Big| \sum_{k=1}^{8N} \beta^k(\chi) (p_k(1) - p_k(2)) \Big| \\ &+ \Big| \sum_{k=1}^{8N} R_k(1) \left( u^k(1) - \xi^k \right)^2 - R_k(2) \left( u^k(2) - \xi^k \right)^2 \Big| \\ &+ \Big| \sum_{k=1}^{8N} Q^k(1) \left( p_k(1) - \chi_k \right)^2 - Q^k(2) \left( p_k(2) - \chi_k \right)^2 \Big|. \end{aligned}$$

Due to the continuity of second derivatives, for each compact set  $K \subset T^*TM$  containing the points 1 and 2, there are two constants  $C_R(K) > 0$  and  $C_Q(K) > 0$  such that  $|R_k(1)|, |R_k(2)| < C_R(K)$  and  $|Q_k(1)|, |Q_k(2)| < C_Q(K)$ , for each k = 1, ..., 8N. Moreover, as a consequence of Taylor's theorem it holds that

$$\lim_{1 \to 2} C_Q(K) = 0, \quad \lim_{1 \to 2} C_R(K) = 0,$$

Therefore, the last two lines in the difference |H(u(1), p(1)) - H(u(2), p(2))| can be rewritten as

$$\left|\sum_{k=1}^{8N} R_k(1) \left(u^k(1) - \xi_t^k\right)^2 - R_k(2) \left(u^k(2) - \xi_t^k\right)^2\right| = C_R(\tilde{K}) \left|\sum_{k=1}^{8N} \left(u^k(1) - u^k(2)\right)^2\right|$$
$$\left|\sum_{k=1}^{8N} Q^k(1) \left(p_k(1) - \chi_k\right)^2 - Q^k(2) \left(p_k(2) - \chi_k\right)^2\right| = C_Q(\tilde{K}) \left|\sum_{k=1}^{8N} \left(p_k(1) - p_k(2)\right)^2\right|.$$

The constants  $C_Q(K)$  and  $C_R(K)$  can be taken finite on K. Furthermore, by restricting further the domain where the points 1 and 2 are to a smaller compact set  $\tilde{K}$ , one can write the following relations,

(7.2) 
$$C_R(\tilde{K})|(u^k(1) - u^k(2))| \le 1/2, \quad C_Q(\tilde{K})|(p_k(1) - p_k(2))| \le 1/2.$$

Let us consider further restriction on the compact set  $K' \subset T^*TM$  such that for each  $(\xi,\chi) \in K'$ 

(7.3) 
$$\left|\frac{\partial H_t}{\partial u^k}\right|_{(\xi,\chi)}\right| \le C_U, \, k = 1, \dots, 4N$$

holds good for some constant  $C_U$ . Also, on K' it must hold that

$$C_R(K) \Big| \sum_{k=1}^{8N} (u^k(1) - u^k(2))^2 \Big| + C_Q(K) \Big| \sum_{k=1}^{8N} (p_k(1) - p_k(2))^2 \Big|$$
  

$$\leq 1/2 \sum_{k=1}^{8N} |(u^k(1) - u^k(2))| + 1/2 \sum_{k=1}^{8N} |(p_k(1) - p_k(2))|.$$

That there is such compact set K' follows from the fact that it is defined by (7.2) and (7.3). Moreover,  $|\beta^i|$  are bounded as a consequence of Randers condition (3.11). Then we have that

$$\begin{aligned} |H(u(1), p(1)) - H(u(2), p(2))||_{K'} &\leq \tilde{C}_U \sum_{k=1}^{8N} \left| (u^k(1) - u^k(2)) \right| + \sum_{k=1}^{8N} \left| (p_k(1) - p_k(2)) \right| \\ &+ 1/2 \sum_{k=1}^{8N} \left| (u^k(1) - u^k(2)) \right| + 1/2 \sum_{k=1}^{8N} \left| (p_k(1) - p_k(2)) \right| \end{aligned}$$

with  $\tilde{C}_K = \max\{C_K, 1\}$ . This proves that  $H|_{K'}$  is a Lipschitz function, with Lipschitz constant  $M = \max\{\frac{1}{2}, \tilde{C}_U\}$ , which is necessarily finite. Now we can redefine the Hamiltonian dividing by M, which is a constant larger than 1, which is equivalent to redefine the vector field  $\beta \in TTM$  and is operation that does not change the equations of motion and the Randers condition (3.11). Then we obtain a 1-Lipschitz Hamiltonian on K' equivalent to the previous one.

**Remark 7.2.** Note that since the Hamiltonian is supposed to be  $C^2$ , it is not necessary for the proof of 7.1 that the Randers condition (3.11) holds. However, consistence with Randers condition implies that the constant  $M = \max\{\frac{1}{2}, \tilde{C}_U\}$ .

The compact domain K' is not empty. However, it is not unique. Extensions from K to the whole phase space can be constructed as follows. Consider the symmetrized distance

$$\varrho_s(a,b) := \frac{1}{2} \left( \varrho(a,b) + \varrho(b,a) \right)$$

on  $T^*TM$  associated to the underlying Hamilton structure (3.2). Let us assume that K' is a star domain and then consider the projection on K'

(7.4) 
$$\pi_{K'}: T^*TM \to K', (u, p) \mapsto (\bar{u}, \bar{p}),$$

where  $(\bar{u}, \bar{p})$  is defined by the condition that the distance from (u, p) to K is achieved at  $(\bar{u}, \bar{p})$ . Then one defines the *radial decomposition* of  $H_t$  by the expression

(7.5) 
$$H_t(u,p) = R(d((u,p),(\bar{u},\bar{p}))) H_t(\bar{u},\bar{p}) + \delta \mathcal{H}_t(u,p)$$

The positive function  $R(d((u, p), (\bar{u}, \bar{p})))$  is such that decreases with the distance function  $d((u, p), (\bar{u}, \bar{p}))$  and it is bounded in a way that the first piece of the Hamiltonian is 1-Lipschitz. The second contribution is not

1-Lipschitz. By definition, it is associated with the matter Hamiltonian  $H_{matter}$ ,

(7.6)  $H_{matter,t}(u,p) := \delta \mathcal{H}_t(u,p).$ 

With these redefinitions we obtain the following

**Theorem 7.3.** Every Hamiltonian (3.26) admits a normalization such that the decomposition (7.1) holds good.

We can read the Hamiltonian constraint (5.11) in the following way. From the properties of the  $U_t$  flow it follows that

$$\lim_{t \to (2n+1)T} \left( \widehat{H}_{matter,t} + \widehat{H}_{Lipschitz,t} \right) |\psi\rangle = 0.$$

However, each of the individual terms in this relation can be different from zero in the metastable regime or near the metastable equilibrium regime,

$$\lim_{t \to (2n+1)T} \widehat{H}_{matter,t} |\psi\rangle \neq 0, \quad \lim_{t \to (2n+1)T} \widehat{H}_{Lipschitz,t} |\psi\rangle \neq 0.$$

This implies that in order to have a the metastable equilibrium regime at t = (2n+1)T, in addition with the matter Hamiltonian (7.6), an additional piece of dynamical systems described by the Hamiltonian  $\hat{H}_{Lipschitz,t}$  is needed.

If we assume that the matter Hamiltonian (7.6) must be positive definite on physical states, then the 1-Lipschitz Hamiltonian should have negative eigenvalues only. Moreover, the function  $\kappa(u, p, t, \tilde{\tau})$  in the quantum Hamiltonian (3.26) takes values in the interval [0, 1]. Hence the positiveness of the matter Hamiltonian is extended to all  $t \in [0, (2n + 1)T]$ . This implies the consistency of the positiveness of the energy level for the quantum Hamiltonian for matter (7.6), since the quantum mechanics description is associated with the full  $U_t$  dynamics.

7.2. Emergence of a weak equivalence principle. We denote by  $X^{\mu}(S_i)$ ,  $i \equiv S, A, B$  the macroscopic observable coordinates associated to the system  $S_i$ , that is, the value of the coordinates that could be associated to a classical system. We assume that the functions

$$\begin{aligned} X^{\mu} &: T^*TM \times \mathbb{R} \to M_4, \\ (u^{k_1}, ..., u^{k_N}, p^{k_1}, ..., p^{k_N}, t) \mapsto X^{\mu}(u^{k_1}, ..., u^{k_N}, p^{k_1}, ..., p^{k_N}, t) \end{aligned}$$

are smooth. Under the additional constraint of universal bounded acceleration and speed for the sub-quantum molecules, in the metastable equilibrium regime  $t \to (2n+1)T$ , the functions  $X^{\mu}((2n+1)T, \tau) = X^{\mu}(\tau)$  are 1-Lipschitz. In order to show this fact we consider first the relations

(7.7) 
$$\lim_{t \to (2n+1)T} \frac{\partial X^{\mu}(u, p, t)}{\partial t} = 0,$$

which hold because in the metastable equilibrium regime there is no tdependence of the physical observables (they are almost fixed points of the  $U_t$  flow). Second, we consider the regularity relations

(7.8) 
$$\frac{dX^{\mu}(u,p,t)}{d\tau} = \sum_{k=1}^{8N} \frac{\partial X^{\mu}}{\partial u_{k}^{\rho}} \frac{du_{k}^{\rho}}{d\tau} + \sum_{k=1}^{8N} \frac{\partial X^{\mu}}{\partial p_{\rho k}} \frac{dp_{\rho k}}{d\tau}.$$

From the relations (7.7) and (7.8) and since the velocity components  $\{\frac{dX^{\mu}(u,p,t)}{d\tau}\}_{\mu=1}^{4}$  are bounded (by the Randers condition (3.11)), it follows that the coordinate functions  $\{X^{\mu}(\tau)\}_{\mu=0}^{3}$  are 1-Lipschitz in all the arguments, since they are  $C^{1}$ -functions with uniformly bounded derivatives.

Since the system of equations for the configuration coordinates  $\{u^i\}_{k=1}^{8N}$ (3.33) is autonomous for u, the functions  $\{X^{\mu}\}_{\mu}^4 = 1$  can be seen as functions of the *u*-coordinates only. Let us assume that the subsystems A and B have local coordinates

$$A \equiv (u_1(\tau), ..., u_{N_A}(\tau), 0, ..., 0)$$
 and  $B \equiv (0, ..., 0, v_1(\tau), ..., v_{N_B}(\tau)),$ 

with  $N = N_A + N_B, N_A, N_B \gg 1$ . The whole system S can be represented in local coordinates as

$$S \equiv (u_1(\tau), ..., u_{N_A}(\tau), v_1(\tau), ..., v_{N_B}(\tau)).$$

By the concentration property (6.6) for the  $U_t$  dynamics in the Lipschitz dynamical regime, the  $\tau$ -evolution of the coordinates  $X^{\mu}(\mathcal{S}(\tau))$ ,  $X^{\mu}(A(\tau))$ and  $X^{\mu}(B(\tau))$  that started with the same initial conditions differs after the dynamics at  $\tau$ -time such that

$$\mu_P \left( \frac{1}{\sigma_{X^{\mu}}} \left| X^{\mu}(\mathcal{S}_i(\tau)) - M^{\mu}(\tau) \right| > \rho \right)_{t \to (2n+1)T} \sim C_1 \exp\left( -C_2 \frac{\rho^2}{2\rho_p^2} \right),$$

 $\mu = 1, 2, 3, 4, i = A, B, S$  holds. The constants  $C_1, C_2$  are of order 1, where  $C_2$  depends on the dimension of the spacetime  $M_4$ . They are independent of the system A, B, S. Note that there is no dependence on the *t*-parameter, since we are in the limit  $t \to T$ . The median  $M^{\mu}(\tau)$  is assumed to be equal to the mean, which is determined by the expression

(7.10) 
$$M^{\mu}(\tau) = \frac{1}{N} \lim_{t \to T} \sum_{k=1}^{N} \mu_{P}(k)(t,\tau) x_{k}^{\mu}(t,\tau).$$

By the diffeomorphisms  $\varphi_k : M_4^k \to M_4$ , one can consider the world lines of the sub-quantum molecules on  $M_4$  at each constant value of t. Hence for t = (2n + 1)T,  $n \in \mathbb{Z}$  we have a set of (discrete) world lines in  $M_4$ , that shows that the functions  $\{X^{\mu}\}_{\mu=1}^4$  characterizes the average presence of sub-quantum lines at a given point of  $M_4$ .

An interchange of sub-quantum molecules between the system and the environment can happen. This can affect the motion of the center of mass  $M^{\mu}(\mathcal{S}_i(\tau))$  in a rather intricate way, because the derivatives  $\{\frac{d}{d\tau}\mu_P(k)(t,\tau)\}_{k=1}^N$ depend upon the details of the system  $\mathcal{S}_i(\tau)$ . In the metastable equilibrium regime t = (2n + 1)T, of there is absence of interaction with the ambient, there is absence of interchange of degrees of freedom with the ambient, at least in average on time t-parameter. Hence the condition

(7.11) 
$$\frac{d}{d\tau}\mu_P(k)(t,\tau) = 0, \quad k = 1, ..., N$$

holds good. In this case, then the center of mass coordinates  $M^{\mu}(\mathcal{S}_i(\tau))$  are solutions of an ordinary differential equation,

(7.12) 
$$\frac{d}{d\tau}M^{\mu}(\tau) = h^{\mu}(\tau), \quad \mu = 1, 2, 3, 4.$$

40

where the functions  $h^{\mu}: T^*TM \to \mathbb{R}$  are fixed by the equations of motion of the 8*N*-degrees of freedom (as consequence of equation (7.10)) and by the properties of the measure  $\mu(k)$ . The local solutions of the equation (7.12) on  $\tau$  with initial condition ( $\tau = 0, M^{\mu}(\tau = 0)$ ) depend only on the initial conditions for  $M^{\mu}(\tau = 0)$ .

**Proposition 7.4.** Let  $S_i$ , i = 1, 2, 3 *HR-systems with*  $N \gg 1$  such that  $M^{\mu}(\tau = 0)$  do not depend on the system A, B or S at  $\tau = 0$  and assume that the condition (7.11) holds for  $\tau \in [0, +\infty]$ . Then the center of mass coordinates  $X^{\mu}(\tau)$  do not depend on the system  $S_i$  for  $\tau$ .

Proof. The coordinate functions  $X^{\mu}(\tau)$  are 1-Lipschitz in the metastable equilibrium regime  $t \to (2n + 1)$ . Then we can apply the concentration of measure (7.9). Moreover, we assume that the condition (6.8) holds, a condition valid for *HR*-systems. Hence the observable coordinates  $\{X^{\mu}\}_{\mu=1}^{4}$ moves following the common  $M\mu(\tau)$  coordinates with an error bounded by  $\exp(-C_2N^2)$ . Since the condition (7.11) holds, the median coordinates  $M\mu(\tau)$  follow an ordinary differential equation, integrable at every  $\tau \in$  $[0, +\infty]$ .

As we discussed before, the figures of N in Hamilton-Randers theory are of order  $10^2$ . The value of the constant  $C_2$  cannot be fixed by the theory, but does not compensate the abrupt concentration caused by the difference of sub-quantum scale and quantum scale.

We see that in the metastable equilibrium regime t = (2n + 1)T there is a strong concentration for the value the functions  $\{X^{\mu}(\tau)\}_{\mu=1}^{4}$  around  $\{M^{\mu}(\tau)\}_{\mu=1}^{4}$ . Note that this universality is up to fixing the initial conditions of the center of mass  $M^{\mu}$ , which is equivalent to fix the initial conditions for  $\{u_{k}^{\mu}\}_{k=1}^{N}$ . This fact does not rest relevance to the main idea that we would like to present here: that for the concentration dynamics  $U_{t}$  in the metastable equilibrium regime  $t \to (2n+1)T$ , for the same initial conditions, the center of mass coordinates  $\{M^{\mu}(\tau)\}_{\mu=1}^{4}$  evolve in the same way, independently of the nature and composition of the system. This result extends towards large systems, composed by atoms. This is the content of the equivalence principle

Furthermore, note that the explanation of the equivalence principle offered along these lines implies that theoretically, such weak equivalence principle is an almost exact law of Nature, only broken at scales  $1/N^2$  compared with the Planck scale. Otherwise, the principle is valid up to an error of  $\exp(-C_2N^2)$ . Such equivalence principle breaks down abruptly in the transition from quantum to classical description of elementary particles.

7.3. An heuristic argument in favour of the emergent origin of the gravitational interaction. Collecting all the previous characteristics for the 1-Lipschitz interaction induced by the  $U_t$  flow in the domain  $\{\lim_{t\to(2n+1)T} U_t, n \in \mathbb{Z}\}$ , we have the following features:

- $U_{(2n+1)T}$  is invariant under infinitesimal diffeomorphism transformations of  $M_4$ , since the constraint (3.27) holds good,
- A weak equivalence principle for the center of mass functions  $S^{\mu}(S(\tau))$  holds good,

#### EMERGENT QUANTUM MECHANICS

- It is a classical interaction, since it appears only during the collapse of the wave function happens
- There is a local maximal speed for the sub-quantum molecules of a HR-systems and invariance under a local relativity group invariance,

Furthermore, we have found the following two additional restrictions

- It is compatible with the existence of a maximal and universal proper acceleration,
- The associated quantum Hamiltonian operator has negative energy eigenvalues (by hypothesis).

In view of the formal similarity of these properties with the analogous properties of the current mathematical description of the gravitational interaction, one can make an strong assumption and postulate that

In the metastable equilibrium regime the 1-Lipschitz dynamics associated with  $H_{Lipshitz,t=(2n+1)T}$  is the gravitational interaction.

That gravity is intrinsically involved in the process of collapse of the wave function is a common idea with several modern approaches to the description of measurement problem [13, 20, 37]. However, as we discuss explicitly before, there are fundamental differences between the models described here and spontaneous collapse models or collapse models induced by large mass measurement devices.

**Existence of a domain where gravity is indeed Lipschitz**. That there is a domain where the gravitational interaction is 1-Lipschitz can be easily seen in the Newtonian limit. Let us consider a Newtonian gravitational force

(7.13) 
$$F(\vec{x}) = -\frac{m M G}{r^2}, \quad \vec{x} \in \mathbb{R}^3$$

and  $r = |\vec{x}|$  the distance to the origin in  $\mathbb{R}^3$  of  $\vec{x}$ . Note that on the set of Euclidean spheres

$$\hat{S}^2 := \{ S^2(r), r \in (0, +\infty) \},\$$

that is where the potential  $V(\vec{x})$  lives,  $|r_1 - r_2|$  defines a norm function. Moreover, to compare significatively length and forces, it is useful to consider dimensionless expressions, for which we need fundamental scales. The Planck force provides a natural unit of force, from which we can compare any other force. In doing this comparison we are adopting a length scale (the Planck length) and a force scale (the Planck force) and use homogenous quantities for length and force. Therefore, let us consider the expression

$$\frac{|F(\vec{x}_2) - F(\vec{x}_1)|}{F_p} = \alpha \frac{|r_2 - r_1|}{l_p}$$

where  $F_p$  is the Planck force and  $l_p$  is the Planck length. After a bit of algebra, one finds an expression for the coefficient  $\alpha$ ,

$$\alpha = l_p \frac{1}{c^4} G^2 m M \frac{1}{r_2^2 r_1^2} |r_2 + r_1|.$$

In order to simplify the argument, we consider m = M. Furthermore, although the case  $r_2 = r_1$  is singular, in order to work in a given scale, we

42

consider a relation  $r_1 = \lambda r_2$  with  $\lambda \sim 1$  constant. Then after some algebra, one obtains a compact expression for  $\alpha$ ,

(7.14) 
$$\alpha = \frac{1+\lambda}{\lambda^3} \frac{D}{D_p} \frac{E}{E_p},$$

where  $D = \frac{m}{r^3}$  is a characteristic density of the system,  $E = mc^2$ ,  $D_p$  is the Planck density and  $E_p$  is the Planck energy. It follows from the expression (7.14) that for scales of the standard model, atomic physics, or macroscopic systems, for instance, that  $\alpha \ll 1$ . Moreover,  $\alpha \to 1$  and even such limit is surpass such value in the limit of the Planck scale.

If we try to repeat this argument for the static electromagnetic field a similar result is obtained. However, we find that it is the full Q.E.D theory that should be used for calculations at the atomic and sub-atomic scale. Hence we should not extend the argument directly from the Newtonian field to the Coulomb field. Moreover, if we take into account the relative intensity of the classical Coulomb field with the Newtonian field, it is a factor, for instance for the electron, of order  $10^{42}$ . This suggests that at such scales the electromagnetic field cannot be 1-Lipschitz. Another argument in favour of this conclusion is that at such scales the electromagnetic field is quantized, which is a very different regime than a 1-Lipschitz regular dynamics.

Although based on a Newtonian limit and in several approximations and assumptions, the conclusion that we can extract is that there is indeed a regime where classical gravity is a 1-Lipschitz interaction. This is in accordance with our interpretation of gravity as an emergent phenomenon.

Despite the fact that as mathematical models, Newtonian gravity or Einstein gravity can be extrapolated to non-Lipschitz domains, specially close to singularities. Such extrapolations, by the arguments given in this section, should be considered un-physical. A possible way out of this dichotomy is to consider classical theories of gravity with a maximal acceleration. Consistently with a weak equivalence principle, such theories must contain maximal acceleration. This can provide eventually providing a classical resolution of General Relativity singularities.

7.4. The Heisenberg dynamics of quantum observables. For each cycle let us fix the value t-time parameter and the slow time  $\tau(t) \in \mathbb{R}$  defined by (3.30). While in the contractive regime the 1-Lipschitz component is dominant, in the ergodic regime of the  $U_t$  flow, it is the matter Hamiltonian what is expected to be dominant. From one side, this is the evolution induced by the generator  $\hat{H}_{matter}$ . From the other side, this is the dynamics that we should expect for any Hermitian observable when there is no present a natural reduction of wave function. Since each cycle labeled by  $\tau$  has associated an ergodic cycle, the sequences of fundamental cycles defines a dynamics along  $\tau$ -time on the observables.

Following this line of argument, let us consider the  $U_{\tau}$  quantum evolution operator by the expression

(7.15) 
$$\widehat{A}(\tau) := \langle (I - \imath \, \delta \tau \, \widehat{H}_{matter})^{\dagger} \, \widehat{A}(\tau_0) \langle (I - \imath \, \delta \tau \, \widehat{H}_{matter}) \, \delta \tau = \tau - \tau_0.$$

It follows at first order in  $\delta \tau$  the following expression,

$$\frac{\widehat{A}(\tau) - \widehat{A}(\tau)}{\tau - \tau_0} = -\imath \left[\widehat{H}_{matter}, \widehat{A}\right].$$

This expression is interpreted as the Heisenberg equation of motion,

(7.16) 
$$i\hbar \frac{d}{d\tau}\hat{A} = -\left[\hat{H}_{matter}, \hat{A}\right].$$

If we assume the existence between fundamental degrees of freedom and quantum degrees of freedom, then the Hamiltonian  $\hat{H}_{matter}$  should be possible to be written in terms of  $\hat{X}^{\mu}$  and  $\hat{P}^{\mu}$  operators. Then (7.16) applies to quantum observables too.

On the linearity of the quantum description. Let us consider a macroscopic observer device determined by two elements: 1. A sub-set of quantum particles that interact with the quantum system being measured via a quantum interaction, 2. An amplification process, that serves to assign macroscopic properties to the quantum system after the measurement is performed. The amplification process consists of a set of quantum processes. If this description is adopted, then according to the Hamilton-Randers theory the natural spontaneous reduction of the quantum state is originated by the  $U_t$  flow. Therefore, the  $U_t$  flow and the linear dynamics described by Heisenberg equation (7.16) provides a natural description of the dynamics.

# 8. Conceptual issues in quantum mechanics from the point of view of Hamilton-Randers theory

In a very interesting book, C. Isham described what is a quaternity of problems in the conceptual foundations of the quantum theory [31]:

- (1) **The meaning of probability.** We have deal with this problem in *sections 5* and *section 6*, pointing out that in Hamilton-Randers theory the origin of the probability description is on an ergodic behaviour of the sub-quantum degrees of freedom.
- (2) The role of measurement. In section 6 we have described our theory of the measurement, showing how is it related with the notion of natural instantaneous reduction of the wave function.
- (3) **The collapse of the quantum state**. In Hamilton-Randers theory this is associated to the natural instantaneous reduction of the wave function.
- (4) **Quantum entanglement**. We think that the relevant point here is to explain the origin of the quantum correlations; quantum entanglement per se is not problematic.

In this *section* we describe an heuristic mechanism to explain quantum nonlocal correlations. Moreover, we describe within our framework the mechanism explaining quantum interference.

Providing a realistic interpretation for all quantum phenomenology is the objective for every realistic interpretation of quantum mechanics. We provide here such interpretations for the fourth problem. Moreover, we also provide a natural interpretation of quantum interference.

8.1. The two slit quantum experiment and its interpretation in Hamilton-Randers Theory. Let us consider the quantum two slit interference experiment as prototype of quantum interference phenomena. In a simplified description of the experiment, the experimental setting is twodimensional, with the x-direction being the direction of propagation of the quantum particles and the z-axis the vertical direction of orientation for the slits and the detector screen. We assume that the intensity of the beam can be regulated to only allow for one quantum particle on flight each time the experiment is repeated. The quantum states are pure quantum states, representing individual quantum particles each time that an experiment is done. The experiment is repeated many times with different quantum particles, under the constraint that the macroscopic initial momenta before reaching the slits is the same for all the particles. We assume that other conditions on the experiment, as the value of the external gravitational field, interactions with the ambient, spin states, etc, either are the same for all the particles or that the variance of these factors do not affect the outcomes of the experiment. Furthermore, the particles in each ensemble of the experiment are quantum mechanically identical.

The standard quantum mechanical description of the experiment can be summarized as follows. After passing the slits, we say that the system is generated by the slit A if, closing the slit B, the particle is detected at the detection screen. Similarly, it is defined that the particle is generated by the slit B. Let  $\psi_A$  be the state generated by the slit A. By this we mean a wave function  $\psi_A$  such that if we close the slit B, it has all the information to reproduce the statistical properties observed in the detection screen. The slit B determines another state denoted by  $\psi_B$ . The evolution after passing the slits is linear and given by a Schrödinger's equation with some given boundary conditions, that depends on the fact of which slits are open.

The evolution is characterized by how many and which slits are open when the quantum system passes through it. If the two slits are open, then the effective state just after the system goes through the slits is described by a vector  $\psi \in \mathcal{H}$  of the form

$$\Psi(x,z) = C \left( \psi_A(x,z) + \psi_B(x,z) \right),$$

with C a normalization real constant such that  $\|\psi\|_{L_2} = 1$ . Since the slits A and B are different, the states  $\psi_A$  and  $\psi_B$  describe different screen patters after a long time exposition when the experiment is repeated many times with prepared individual identical particle states.

It is natural to adopt the hypothesis that  $\psi_A \neq \psi_B$  in the vicinity of the detector screen. Furthermore, symmetry considerations imply that  $\psi_A = \psi_B$  in the central region of the screen, reaching a maximum value  $||\psi_A + \psi_B||^2$ . Out of the central axis one expects either  $|\psi_A| \neq |\psi_B|$  or  $\arg(\psi_A) \neq \arg(\psi_B)$  or both conditions hold. In the case that there is a relative phase between  $\psi_A$  and  $\psi_B$ , an interference pattern depending on the geometric arrangement of the experiment should be found and as we move out from the axis z = 0, a relative phase between  $\psi_A$  and  $\psi_B$  should appear. Moreover, one also expects to find that  $||\psi_A|| \neq ||\psi_B||$  holds along the z-axis.

The physical interpretation of the double slit experiment from the point of view of Hamilton-Randers theory is the following. First, let us note that in the above quantum mechanical description of the double quantum slit experiment the time parameter is 1-dimensional, since only  $\tau$  has been implicitly considered. Hence the quantum system appears as passing through the two slits A and B at the same time, if both slits are open. Then we have that  $\tau_A = \tau_B$ . However, from the point of view of Hamilton-Randers systems, time is a two-dimensional parameter. Therefore, the dynamical state of the quantum particle should be specified by given the conditions of motion at a two-time instant for each sub-quantum molecule. In the case in question, we can think that the particle has passed through A at the instant  $(t_A, \tau)$ , while the sub-quantum particle has pass by the slit B at the instant  $(t_A, \tau)$ . Hence the system pass by the two slits, at two different 2-time instants  $(t_A, \tau)$  and  $(t_B, \tau)$ , but when the projection  $(t, \tau) \mapsto \tau$  is done, the system appears as if it pass through both slits at the same time parameter  $\tau$ . This idea is compatible with a local description of the phenomena of quantum interference. It is after this projection that apparent quantum mechanical non-locality are present.

In this interpretation,  $\psi_A$ ,  $\psi_B$  and  $\psi$  are epistemic states, for different reasons. From one side, the states described by the vectors  $\psi_A$  and  $\psi_B$ are epistemic, since they describe the partial evolution of an unique quantum state. That is, they describe the state when it pass through A or pass through B respectively. What happens from the point of view of HRsystems, is that the sub-quantum degrees of freedom pass through both slits many times (in the two-dimensional sense), in such a way that the *partial* states  $\psi_A$  and  $\psi_B$  are associated with probability distributions that can be used in the forecast of phenomenological predictions. However, these states do not correspond to individual sub-quantum molecules, and are more a convenient tool to calculate probabilities and expectations values associated frequency probabilities. On the other hand,  $\psi$  is only an epistemic state since the projection  $(t,\tau) \mapsto \tau$  has been taken already. Hence it cannot determine the full dynamics of the associated HR-system. In contrast, it is also a convenient tool to calculate probabilities associated with the quantum system. Hence the epistemic character of the quantum description, in contrast with the ontological character of the Hamilton-Randers theory, follows. Finally, if the two slits A and B are open, each time that an individual experiment is done, we can ask by which of the slits the quantum particle passed through it. This question is legitim, due to the process of natural spontaneous collapse described in section 6.

8.2. On the non-local quantum correlations. The existence of *long* distance quantum correlations is an astonishing realization of quantum phenomenology that claims for explanation. Is it possible, even in a qualitative way, to understand such phenomena within the framework of Hamilton-Randers theory? There are three mechanisms that we can think are a priori involved in the quantum correlations.

A. Projection map  $(t, \tau) \mapsto \tau$ . The ergodic motion of the sub-quantum molecules is constrained by hypothesis to be sub-luminal. Henceforth, the maximal distance achievable for the instantaneous quantum correlations is

bounded by

$$d_{cor} \leq 2 T c$$

where c is the speed of light in vacuum. By the relation (3.15), this distance will depend on the mass of the particle. For massive quantum systems, the distance 2Tc, is small and of the order of the range of the distances where one expect to have quantum interference, in the case of massive quantum particles.

This mechanism has interesting consequences for the correlations associated to massive quantum systems, that is the limitation on the distance for the correlations, but also, the apparent instantaneous character of the correlations. Indeed, for massive particles one has

(8.1) 
$$d_{cor} \le 2 \frac{\alpha \hbar}{c} \frac{1}{M}$$

This inverse law of the maximal distance of correlations with the mass of the system is falsifiable, despite the appearance of the un-known scale factor  $\alpha$ . The prediction of apparent instantaneous correlations is also falsifiable in principle. However, the factor  $\alpha$  must be fixed, for each experimental situation, in order to develope the full predictive power of relation (8.1).

For massless systems M = 0, as in the case of photons, the correlation distance is arbitrarily large and appears as instantaneous non-local correlations in  $M_4$ . This can explain the long distance quantum correlations along several kilometres [43] but also the apparent not bound on the correlations speed in the laboratory frame for photons [22].

**B.** Direct action in a geometry of maximal acceleration. It is possible to re-think the definition of celerity in geometric spaces of maximal proper acceleration and maximal speed as follows. First we make the hypothesis that the correct continuous limit of the discrete dynamics for the sub-quantum molecules is described in a maximal acceleration back-ground geometry as in reference [19]. In particular, it was shown that in a geometry of maximal universal proper acceleration and universal maximal speed, the celerity of an accelerated classical point particle with world line  $\mathbf{x} : I \to M_4$  is given by the expression

(8.2) 
$$v(t) := \frac{1}{\sqrt{1 - \frac{a^2}{A_{max}^2}}} \frac{1}{\sqrt{1 - \frac{\tilde{v}^2}{c_{max}^2}}} \tilde{v}(t),$$

where  $a^2$  is the square of the proper acceleration,

$$a^2 = \eta_4(\ddot{\mathbf{x}}, \ddot{\mathbf{x}}).$$

The speed of light in vacuum can be considered as the standard maximal speed from the point of view of a macroscopic observer for sub-quantum molecules. This is because, for macroscopic observers acceleration is small and there is not perception of  $A_{max}$ . If we adopt this perspective, we have that for a macroscopic observer, that by definition is not subjected in usual situations to very large accelerations, the sub-quantum molecules, although un-observable directly (that is, by the effect on a external gauge field or external gravitational field) can be ascribed supra-luminal direct actions.

Now let us define the apparent celerity by the expression

$$\tilde{v}_{app} := \frac{1}{\sqrt{1 - \frac{a^2}{A_{max}^2}}} \, \tilde{v}.$$

Similarly, the apparent speed of light in vacuum  $c_{app}$  is defined by

$$c_{app} := \frac{1}{\sqrt{1 - \frac{a^2}{A_{max}^2}}} c.$$

Hence one can re-write the expression (8.2) as

(8.3) 
$$v(t) := \frac{1}{\sqrt{1 - \frac{\tilde{v}_{app}^2}{c_{app}^2}}} \tilde{v}_{app}(t)$$

The direct reading of (8.3) is suggestive. For accelerate particles, the maximal speed achievable is  $c_{app}$ . Then let us note two points related with (8.3). Firstly is that  $c_{app} \ge c$ . Secondly, the deduction of (8.2) was done under the assumption of small accelerations compared with  $A_{max}$  [19]. Thus in order to have a relevant effect one needs to expand the domain of validity of (8.2) to large accelerations. However, this is mathematically consistent with the domain of the expression (8.3).

This mechanism, that we should associated to a direct propagation and interaction of sub-quantum molecules, is not independent of the mechanics **A**. It applies to sub-quantum molecules by construction, that corresponds to the  $U_t$  flow, since the mass of the sub-quantum molecules m is small but different from zero, and henceforth, suitable to suffer huge accelerations when the interaction is not Lipschitz, that is, when the interaction is not gravitational. Note that such mechanism does not applies directly to quantum systems.

**C.** Conservation laws. Let us consider the conservation law (3.32), obtained by a geometric consistent argument that allows to define the slow time parameter  $\tau$ . Due to the product structure of the configuration space there are certain constraints as consequence of the heritage conservation law property. Such conservation rules can be written schematically in the form

$$\beta_A + \beta_B + \beta_C + \dots = 0$$

for a finite set of observables  $\{\beta_A, \beta_B, \beta_C, ...\}$  that enter on the heritage conservation law involving different macroscopic parties  $\{A, B, C, ...\}$ . The origin of these conservations laws are in the  $U_t$  flow, since are associated with the dynamics and configuration of sub-quantum degrees of freedom. Such conservation laws are of the type of common cause. They are ruled out by the experimental violation of Bell's inequalities[21], except if the hypothesis of super-determinism is adopted [29]. Such hypothesis is fully consistent with our theory, but probably lacks of the possibility of being falsifiable.

Despite the falsification of the possibility of mechanism Cto explain by its own the nature of quantum correlations by Bell's inequalities, a combination of the heritage conservation laws with the mechanism described in B allows the possibility that direct action of sub-quantum systems can change the conservation laws. This is provided by the  $U_t$  flow and henceforth, subject to the constraint (8.1).

8.3. On the emergence and ontological nature of the  $\tau$ -time. The  $\tau$ time class of parameters has been considered to be within the category real parameters. However, the observation that the  $U_t$  flow is almost periodic in t and that every quantum measurement is performed in the metastable equilibrium regime given by (5.11) implies the discreteness of any  $\tau$ -time parameter.

There are several consequences of this reasoning. The first is that the continuous description of clocks in classical physics and quantum mechanics must be regarded as an idealization. It is however, a very useful one, due to the extreme smallness of the Planck scale, related with the scale of subquantum degrees of freedom, compare with quantum and macroscopic scales. We adopted the real description of the  $\tau$ -time parameter in this work only motivated by practical reasons.

The second consequence is the emergent origin of the notion of external  $\tau$ time parameters. Given a particular clock, the time parameters  $\tau$  appear as determined by the  $U_t$  flow at the Planck scale, as the number of fundamental cycles of the  $U_t$ -flow. It is an emergent time, that arises from physical processes. In this sense, time as used in physical description corresponds to an abstraction.

On the other hand, this emergent character of the  $\tau$ -time is not in contradiction with the requirement of *time diffeomorphism invariance* of general relativity, since our interpretation of the  $\tau$ -time applies to any physical clock as specified in *sub-section* 3.7 and there is the constraint that the theory is diffeomorphic invariant. Hence there is not a privileged notion of  $\tau$ -time.

Despite of the emergence of time, we claim that the physical world is fundamentally irreversible and unrepeatable. The irreversible nature of the dynamical evolution is based on its emergent character and on the physical impossibility to repeat the conditions for the fundamental sub-quantum dynamics from the point of view of the quantum scale. By this we mean that the complexity of dynamics of sub-quantum degrees of freedom and the impossibility to control the  $U_t$  flow using quantum interactions. Since the  $\tau$  evolution corresponds to a larger scale than the  $U_t$  flow, it is not possible to control the fundamental flow. This implies the existence of an irreversible time evolution for the physical systems and also for the whole universe, emergent arrows of  $\tau$ -time.

Our theory suggests the impossibility of *travel back*in the  $\tau$ -time for any standard model elementary particle or for any larger system. Second, as we have remarked before, it only leaves the possibility of  $\tau$ -time re-parametrization invariance to be an emergent property in the classical or metastable equilibrium regime. In this regime, the condition is fully compatible with the Hamiltonian constrain of general relativity.

The unrepeatable property can be argued from a probabilistic point of view, since the universe as a systems must be described by a multitude of atomic and sub-atomic systems, each of which is described in Hamilton-Randers systems by a complex systems of sub-atomic molecules. Hence our universe is even more complex than any quantum picture could provide. From this argument follows the impossibility to repeat the same initial conditions for the whole universe. Moreover, the lack of direct control on the sub-quantum levels make even more credible the thesis of un-repeatability.

# 9. Concluding Remarks

9.1. Relation with others theories of emergent quantum mechanics. The theory presented in this work has several remarkable similarities with other theories of emergent quantum mechanics, in particular with the approach developed extensively by G. 't Hooft and others (see for instance [10, 15, 26, 30]). Among such analog properties, there are the introduction of a dissipative dynamics, in order to account of the quantum states [9, 26] and the fact that the wave function is  $\psi$ -epistemic. In Hooft's approach, unitarity is recover at the level of equivalence classes. In our description, where there is not a dissipation of information or a dissipative dynamics, but instead, an internal  $U_t$  flow which is assumed of a rather intricate structure. There is no preservation of the volume phase element for the  $U_t$  flow, since the dynamics is driven by a time dependent Hamiltonian. In contrast, unitarity holds for the  $U_{\tau}$  evolution.

There are important departures between our theory and the above mentioned theories of emergent quantum mechanics. Let us mention, as a very significant one that in our theory, time is described by a two-dimensional parameter. In the continuous limit, such parameter is  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . The parameter  $\tau$  is the external time, which servers to describe the dynamics of a quantum state  $|\psi\rangle$ . The  $\tau$ -time parameter corresponds with the usual notion of macroscopic time as it appears in quantum mechanics and quantum field theory. The internal t-time parameter in HR-systems describes the  $U_t$ internal dynamics of the Hamiltonian  $H_t(u, p)$  and appears in our theory as the parameter of the homotopic evolution (3.18) of the Hamilton-Randers geometric structure. Therefore, the parameters t and  $\tau$  are qualitatively different, which implies the two-dimensional property of time and that the associated two dynamical evolution  $(U_t, U_{\tau})$  are irreducible to each other.

Our theory is deterministic, since the vector field  $\beta \in TTM$  determines the whole dynamics of the system, together with the initial conditions. Moreover, there is the condition (3.32), which can be a source for  $M_4$  spacelike correlations. This is in accordance with the idea of *super-determinism* as explanation for the quantum correlations [29]. A second source of quantum correlations is on direct action causation, based on a reinterpretation of the geometry of maximal acceleration.

Is it necessary to introduce such foreign notion from physics as it is the notion of 2-dimensional time? It is notable the formal resemblance between the dynamical system  $(U_t, U_\tau)$  along  $(t, \tau)$  and fast-slow dynamical models [3]. In such dynamical systems, there are two time scales for the variation of the dynamical degrees of freedom: there are slow degrees of motion and fast degrees of motion. Usually, the relevant degrees of freedom in the slow motion are the slow degrees of freedom, with an effective dynamics on the slow manifold. One is tented to identify in Hamilton-Randers models the fast degrees of motion with the sub-quantum molecules and the slow degrees of motions with the densities and wave functions described by elements of  $\mathcal{H}$ , which are determined by t-time averages. However, a closer examination of the situations refrains us to make such identification. First, in the fast-slow dynamics, there is a one to one map between the values of the fast time t and the values of the slow  $\tau$ -time. In the case of Hamilton-Randers systems, such bijection fails, since the values of  $\tau$  correspond to a discrete set of values of t, although t is continuous. Moreover, in Hamilton-Randers systems the time t and the  $\tau$ -time are not related. By construction of the dynamics, the  $\tau$ -parameter is independent of the t-parameter, in the mathematical description of each Hamilton-Randers system.

9.2. Relation of Hamilton-Randers theory and de Broglie-Bohm theory. There is certain analogy between the properties that our theory present and the de Broglie-Bohm theory [11, 25]. In Hamilton-Randers theory, the value of the observables of the system are well-defined before any measurement is done and independent of the decisions of the observer takes. Indeed, in Hamilton-Randers theory, the existence of the natural spontaneous collapse processes solves the measurement problem using local notions and without the introduction of a entanglement with the measurement device or with the mind of the observer. The quantum system is in a localized state prior to any measurement performed by a macroscopic observer. In an localized state, all the possible classical observables have a definite value. Hence, in Hamilton-Randers theory, there is the possibility to assign define trajectories to the quantum system.

However, it is natural in Hamilton-Randers theory to interpret the wave function as a presence of matter during the non-contractive phase of the  $U_t$  dynamics. Therefore, in our theory the wave function has an epistemic interpretation, in sharp contrast with de Broglie-Bohm theory (at least in some interpretations, that includes the original formulation of D. Bohm), where the wave function is a real ontological field. Hence it must exist observable differences between the theory proposed here and de Broglie-Bohm theory. Because the nature of the interpretation of the wave function in Hamilton-Randers theory (epistemic) and the de Broglie-Bohm theory (ontological) and since these differences appeared mainly in the core of the measurement processes, we think that they could be on the absence of the wave function as a source for the gravitational interaction. In de Broglie-Bohm theory, on the other hand, since the wave function  $\psi$  has physical existence, it seems necessary that modifies the gravitational field. This problem is under current investigation.

9.3. **Open problems in Hamilton-Randers theory.** The main objective of following investigations is to find more concrete realizations of the concepts first explored here. Several important issues need to be clarified and developed in our proposal.

**Realization of HR-models and relation with number theory** One of the most urgent one is to find a concrete set of dynamical systems full-filling all the properties that we have discussed. Several candidates are being investigated by the author. Among them, the so called *Riemann flow* models of analytical number theory and quantum chaos. It can be shown that

HR-systems are N-dimensional relativistic versions (in the sense of compatibility with the Lorentz group) of xp-Hamiltonian models for the Riemann flow (classical relevant works on this field and several more recent ones are [7, 8, 12, 39, 40]). It is remarkable that several characteristics explicitly proved that hold or theoretically required for HR-systems are indeed found in xp-models of Riemann dynamics and viceversa, that HR-models are candidates to the quantum mechanical approach to the Riemann Hypothesis (it is worth while to compare the Hamiltonian (3.26) with the Berry-Connes Hamiltonian [7, 12]. Moreover, there is a clear correlation between some of the assumptions for Hamilton-Randers systems described in section 2 with the assumptions of the Riemann flow [8]). It could be worthwhile to mention that in a recent work [40] by G. Sierra it was discussed a xp-model with many features that the Riemann flow should have. Moreover, it contains an (ad hoc) maximal acceleration and maximal speed (the speed of light in vacuum). Hence we make the suggestion that Sierra's model is an example of HR-system. To investigate in detail this conjecture is interesting, since it could provide a particular example of a dynamical system that plays an important emergent quantum mechanics and analytical number theory. More in general, it is suggested that the Hilbert-Polya conjecture is realized, not at the atomic or quantum level [39], but at a more fundamental level of the physical reality which can be the Planck scale.

Relation between direct observables and microscopic operators. Another open question in our approach is how to construct quantum mechanical observables in terms of microscopic operators acting on ontological states  $\{|x_k^{\mu}, y_k^{\mu}\rangle\}_{k=1,\mu=1}^{N,4}$ . The transition from the description of the dynamics provided by  $\hat{H}_{matter,t}(\hat{u}, \hat{p})$  to the description by  $\hat{H}_{matter,t}(\hat{X}, \hat{P})$  requires to know the structure of the operators  $\{\hat{X}, \hat{P}\}$  in terms of the operators  $\{\hat{u}, \hat{p}\}$  (or viceversa, if the relations are invertible) or a formal argument to identify the quantum operators.

We should address in this contest the relation between the mass m of each sub-quantum molecule and the spectrum of masses of elementary quantum particles.

Quantum non-locality and entanglement. The notion of two dimensional time parameter  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$  is beneath our interpretation of the fundamental non-local description of the quantum mechanics as follows. In order to describe the state of a system at a given instant, in Hamilton-Randers theory we need to specify the two time parameters  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ . If only the parameter  $\tau$  is specified in a dynamical description of a physical system (as it is done in usual field theory and quantum mechanics) and if the system is not in an localized state for the  $U_t$  dynamics, an intrinsic non-local behavior of the state associated with the ergodicity of the dynamics emerges. We think that this is the origin of the non-local properties of quantum systems. Hence a complete explanation of the quantum non-locality could and should be developed in our framework. Indeed, a theory of entangled states, based on the embedding  $\mathcal{H} \hookrightarrow \mathcal{H}_{Planck}$  that generalize the formula (5.2) can be easily constructed. Although we have envisage a geometric way to understand quantum nonlocal behaviour, a complete mathematical treatment must be investigated: how is it related the projection  $(t, \tau) \mapsto \tau$  with Bell's inequalities [5, 6]?

Our idea to explain the quantum correlations are based on a combination of the mechanisms of **A**, time projection  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ , **B** (direct action) and **C** (common cause) based on direct action implies a tremendous change in perspective respect to the structure of relativity and the geometry of spacetime and the idea of local determinism. As we mention before, our argument is based in four ideas

- Time is two dimensional and an effective quantum description is obtained after the projection  $(t, \tau) \in \mathbb{R} \times \mathbb{R}$ .
- The sub-quantum molecules world lines are such the proper acceleration is bounded,
- The expression for the apparent celerity of the sub-quantum particles is given by (8.3).
- Conservations law  $\nabla \beta = 0$  and the heritage conservations laws.

These four components interact between each other, to provide a potential explanation of quantum non-local correlations.

Absolute structures in HR-theory. It is a rather problematic point the fact that several geometric structures has been introduced as background structures without a corresponding dynamical law in our proposal. In particular, the metric  $\eta_4$  and the collection of metrics  $\{\eta_4^k\}_{k=1}^N$  are background structures. We believe that a more through treatment should provide a natural dynamics for  $U_t$  and  $U_{\tau}$ , hence for the Hamilton-Randers structure. Compatibility with metastable equilibrium regime implies that the  $U_t$  flow provides a thermodynamical limit. Hence it can be possible to derive field equations for the metrics  $\{\eta_4^k\}_{k=1}^N$  valid in the metastable regime as equation of states thermodynamical equilibrium condition.

Acknowledgements. This work was supported by PNPD-CAPES n. 2265/2011, Brazil. I would like to thank to J. M. Isidro for several comments.

#### References

- D. Acosta, P. Fernández de Córdoba, J. M. Isidro, J. L. G. Santander, *Emergent quantum mechanics as a classical, irreversible thermodynamics*, Int. J. Geom. Meth. Mod. Phys. **10**, No. 04 1350007 (2013).
- [2] S. L. Adler, 2004 Quantum Theory as an Emergent Phenomenon: The Statistical Mechanics of Matrix Models as the Precursor of Quantum Field Theory, Cambridge University Press (2004).
- [3] V. I. Arnold, Geometric Methods in the Theory of Ordinary Differential Equations, Springer-Verlag (1983).
- [4] D. Bao, S.S. Chern and Z. Shen, An Introduction to Riemann-Finsler Geometry, Graduate Texts in Mathematics 200, Springer-Verlag.
- [5] J. S. Bell, On the Einstein-Podolski-Rosen paradox, Physica 1, 195 (1964).
- [6] J. S. Bell, Speakable and unspeakable in quantum mechanics, Cambridge University Press (1987).
- M. V. Berry, *The Bakerian Lecture*, 1987: *Quantum Chaology*, Proc. R. Soc. Lond. A 413, 183-198 (1987).
- [8] M. V. Berry and J. P. Keating, H = xp and the Riemann zeros, in Supersymmetry and Trace Formulae: Chaos and Disorder, ed. J. P. Keating, D. E. Khemelnitskii, L. V. Lerner, Kuwler 1999; M. V. Berry and J. P. Keating, The Riemann zeros and eigenvalue asymptotics, SIAM REVIEW 41 (2), 236 (1999).

- [9] M. Blasone, P. Jizba and G. Vitiello, Dissipation, Emergent Quantization and Quantum Fluctuations, in Decoherence and Entropy in Complex Systems, Selected Lectures from DICE 2002, H.-T. Elze (ed.), Lecture Notes in physics 633, Springer, Berlin (2004).
- [10] M. Blasone, P. Jizba and F. Scardigli, Can Quantum Mechanics be an Emergent Phenomenon?, J. Phys. Conf. Ser. 174 (2009) 012034, arXiv:0901.3907[quant-ph].
- D. Bohm, A suggested Interpretation of the Quantum Theory in Terms of Hidden variables. I, Phys. Rev. 85, 166-179 (1952); A suggested Interpretation of the Quantum Theory in Terms of Hidden variables. II, Phys. Rev. 85, 180-193 (1952); D. Bohm, Wholeness and the Implicate Order, London: Routledge, (1982).
- [12] A. Connes, Trace formula in noncommutative geometry and the zeros of the Riemann zeta function, Selecta Mathematica (New Series) 5, 29 (1999).
- [13] L. Diósi, Gravitation and quantum-mechanical localization of macro-objects, Phys. Lett. A 105, 199; L. Diósi Phys. Rev. A 40 1165, (1989).
- [14] P. A. M. Dirac, The Principles of Quantum Mechanics, Fourth Edition, Oxford University Press (1958).
- [15] H.T. Elze Quantum mechanics emerging from "timeless" classical dynamics, Trends in General Relativity and Quantum Cosmology, ed. C.V. Benton (Nova Science Publ., Hauppauge, NY, 79-101 (2006); H.T. Elze, The Attractor and the Quantum States, Int. J. Qu. Info. 7, 83 (2009); H.T. Elze, Symmetry Aspects in Emergent Quantum Mechanics, J. Phys. Conf. Ser. 171, 012034 (2009).
- [16] H. T. Elze, Action principle for cellular automata and the linearity of quantum mechancics, Phys. Rev. A 89, 012111 (2014); H. T. Elze, Are nonlinear discrete cellular automata compatible with quantum mechanics?, J. Phys.: Conf. Ser. 631, 012069 (2015).
- [17] P. Fernández de Córdoba, J. M. Isidro and Milton H. Perea Emergent quantum mechanics as thermal essemble, Int. J. Geom. Meth. Mod. Phys. 11, 1450068 (2014).
- [18] R. Gallego Torromé, A Finslerian version of 't Hooft Deterministic Quantum Models, J. Math. Phys. 47, 072101 (2006).
- [19] R. Gallego Torromé, An effective theory of metrics with maximal acceleration, Class. Quantum Grav. 32 245007 (2015).
- [20] G. Ghirardi, R. Grassi and A. Rimini, Continuous-spontaneous-reduction model involving gravity, Phys. Rev. A 42 1057 (1990).
- [21] N. Gisin, Quantum correlations in Newtonian space and time: arbitrarily fast communication or nonlocality, in Quantum Theory: a two-time success story, Yakir Aharonov Festschrift, pp 185-204, Springer (2014).
- [22] N. Gisin, V. Scarani, W. Tittel and H. Zbinden, {100 years of Q theory}, Proceedings, Annal. Phys. 9, 831-842 (2000).
- [23] G. Gröessing, Emergence of Quantum Mechanics from a Sub-Quantum Statistical Mechanics, Int. J. Mod. Phys. B, 28, 1450179 (2014).
- [24] M. Gromov, Riemannian structures for Riemannian and non-Riemannian spaces, Birkhäuser (1999).
- [25] P. R. Holland, The Quantum Theory of Motion: An Account of the de Broglie-Bohm Causal Interpretation of Quantum Mechanics, Cambridge University Press (2004).
- [26] G. 't Hooft, Determinism and Dissipation in Quantum Gravity, hep-th/0003005.
- [27] G. 't Hooft, Determinism beneath quantum mechanics, arXiv:quant-ph/0212095.
- [28] G. 't Hooft, Emergent Quantum Mechanics and Emergent Symmetries, 13th International Symposium on Particles, Strings, and Cosmology-PASCOS 2007. AIP Conference Proceedings 957, pp. 154-163 (2007).
- [29] G. t'Hooft, How a wave function can collapse without violating Schrödinger's equation, and how to understand the Born's rule, arXiv:1112.1811v3.
- [30] G. 't Hooft, The Cellular Automaton Interpretation of Quantum Mechanics: a View on the Quantum Nature of our Universe, Compulsory or Impossible?, arXiv:1405.1548v3.
- [31] C. Isham, Lectures on Quantum Mechanics: Mathematical and Structural Foundations, Imperial College Press (2001).

54

- [32] A. Khrennikov, "Einstein dream: Quantum Mechanics as Theory of Classical Random Fields, arXiv:1204.5172 [quant-ph].
- [33] I. Kolar, P. W. Michor, J. Slovak, Natural operators in differential geometry, Springer-Verlag (1993).
- [34] V. D. Milman and G. Schechtman, Asymptotic theory of Finite Dimensional normed spaces, Lecture notes in Mathematics 1200, Springer (2001).
- [35] R. Miron, H. Hrimiuc, H. Shimada and S.V. Sabau, The geometry of Hamilton and Lagrange spaces, Kluwer Academic Publishers (2002).
- [36] E. Nelson, Derivation of the Schrödinger equation from Newtonian Mechanics, Phys. Rev. 150 1079-1083 (1964); Dynamical theories of brownian motion, Princeton University Press (1967); Quantum Fluctuations, Princeton University Press (1985).
- [37] R. Penrose, On Gravity's Role in Quantum State Reduction, Gen. Rel. and Gravit.
   8, No. 5, 581 (1996); R. Penrose, The Road to Reality, Vintage, London (2005).
- [38] G. Randers, On an Asymmetrical Metric in the Four-Space of General Relativity, Phys. Rev. 59, 195-199 (1941).
- [39] G. Sierra and P. K. Townsend, Landau levels and Riemann zeroes, Phys. Rev. Lett. 101, 110201 (2008).
- [40] G. Sierra, The Riemann zeros as energy levels of a Dirac fermion in a potential built from the prime numbers in Rindler spacetime, J. Phys. A: Math. Theor. 47 325204 (2014).
- [41] L. Smolin, Could quantum mechanics be an approximation to another theory?, quant-ph/0609109.
- [42] M. Talagrand, A new look at independence, Ann. Probab. 24, Number 1, 1-34 (1996).
- [43] W. Tittel, J. Brendel, H. Zbinden and N. Gisin Violation of Bell inequalities by photons more than 10 km apart, Phys. Rev. Lett. 81, 3563 (1998).
- [44] C. Wetterich, Emergence of quantum mechanics from classical statistics, DICE 2008, J. Phys.: Conf. Ser., 174 012008 (2009).