

Energy dependence of $e^+e^- \rightarrow 6\pi$ and $e^+e^- \rightarrow N\bar{N}$ cross sections near the $N\bar{N}$ threshold

A.E. Obrazovsky^{1,*} and S.I. Serednyakov^{1,2,†}

¹*Budker Institute of Nuclear Physics SB RAS, 630090 Novosibirsk, Russia*

²*Novosibirsk State University, 630090 Novosibirsk, Russia*

(Dated: February 28, 2024)

Using recent BABAR, CMD-3 and SND data, the sum of $e^+e^- \rightarrow 3(\pi^+\pi^-), 2(\pi^+\pi^-\pi^0), p\bar{p}, n\bar{n}$ cross sections is obtained. Unlike $e^+e^- \rightarrow 3(\pi^+\pi^-)$ and $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ processes, no structures in total cross section are found near the $N\bar{N}$ threshold within the limits of measurement errors.

PACS numbers: 13.66.Bc, 14.40.Be

First observation of six-pion production in electron-positron annihilation was performed in $\mu\pi$ experiment at ADONE collider [1]. The dip in $e^+e^- \rightarrow 3(\pi^+\pi^-)$ cross section near the $N\bar{N}$ threshold was found in DM1 experiment at DCI [2]. Later, it was confirmed by more precise measurements in FOCUS [3] and BABAR [4] experiments. The similar structure also was observed in $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ mode [4, 5]. One of the possible theoretical explanations of this dip suggests the existence of $p\bar{p}$ underthreshold bound state [6].

Recently, more precise BABAR and SND data on $e^+e^- \rightarrow p\bar{p}$ [7] and $e^+e^- \rightarrow n\bar{n}$ [8] cross sections were published together with the new CMD-3 data on $e^+e^- \rightarrow 3(\pi^+\pi^-)$ [9] and $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ cross sections [10].

We suggest that the sum of $e^+e^- \rightarrow \text{hadrons}$ cross sections including $N\bar{N}$ final state should contain no structure near the $N\bar{N}$ threshold due to absence of quark-antiquark resonances in this energy range.

For the purposes of this work only $e^+e^- \rightarrow 3(\pi^+\pi^-)$, $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ and $e^+e^- \rightarrow N\bar{N}$ processes can be considered, as far as the cross sections of $e^+e^- \rightarrow 2\pi, KK, 3\pi, KK\pi, 4\pi, KK\pi\pi, 5\pi, \dots$ don't contain structures near the $N\bar{N}$ threshold [11]. We use CMD-3 data on $e^+e^- \rightarrow 6\pi$ processes, BABAR and SND data on $e^+e^- \rightarrow N\bar{N}$ processes. The sum of isovector $e^+e^- \rightarrow 3(\pi^+\pi^-)$ and $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ cross sections with isoscalar $e^+e^- \rightarrow \eta\pi^+\pi^-\pi^0$ background subtraction [12] is shown in fig.1a. The sum of $e^+e^- \rightarrow p\bar{p}$ and $e^+e^- \rightarrow n\bar{n}$ cross sections is shown in fig.1b. These sums are fitted by constants in the energy ranges $2E = 1.85\text{--}1.876$ GeV and $1.876\text{--}1.92$ GeV. The cross section discontinuity values are: $\Delta\sigma = -1.5 \pm 0.2$ nb for the sum of $e^+e^- \rightarrow 6\pi$ processes, $\Delta\sigma = 1.6 \pm 0.25$ nb for the sum of $e^+e^- \rightarrow N\bar{N}$ processes. For the total sum of $e^+e^- \rightarrow 6\pi$ and $e^+e^- \rightarrow N\bar{N}$ processes $\Delta\sigma = 0.1 \pm 0.3$ nb (fig. 1c). The quoted uncertainties include both statistical and systematic errors. The total cross section doesn't contain structures within measurement error limits.

We suggest that negative discontinuity in the cross section of $e^+e^- \rightarrow 6\pi$ processes at the $N\bar{N}$ threshold can be quantitatively explained by the opening of new annihilation channel $e^+e^- \rightarrow N\bar{N}$. However, it is unclear, why it is isovector $e^+e^- \rightarrow 6\pi$ final state that provide nearly full compensation in the total cross section $e^+e^- \rightarrow \text{hadrons}$.

The authors are grateful to V. P. Druzhinin and A. I. Milstein for useful discussions. The work is partially supported by the Ministry of Education and Science of the Russian Federation, scientific school grant NSH-2479.2014.2 and Russian Fund for Basic Research grant RFBR 12-02-00065-a.

-
- [1] M. Grilli, E. Iarocci, P. Spillantini et al., *Nuovo Cimento* **13A**, 593 (1973).
 - [2] D. Bisello, J. C. Bizot, J. Buon et al., *Phys. Lett.* **107B**, 145 (1981).
 - [3] P. L. Frabetti, H. W. K. Cheung, J. P. Cumalat et al., *Phys. Lett.* **B514**, 240 (2001).
 - [4] B. Aubert, R. Barate, D. Boutigny et al., *Phys. Rev.* **D73**, 052003 (2006).
 - [5] R. Baldini et al., reported at the "Fenice" Workshop, Frascati (1988).
 - [6] P. L. Frabetti, H. W. K. Cheung, J. P. Cumalat et al., *Phys. Lett.* **B578**, 290 (2004).
 - [7] J. P. Lees, V. Poireau, V. Tisserand et al., *Phys. Rev.* **D87** 092005 (2013).
 - [8] S. I. Serednyakov, reported at International Workshop on e^+e^- collisions from Phi to Psi (PHIPSI13), Rome (2013).
 - [9] R. R. Akhmetshin, A. V. Anisenkov, S. A. Anokhin et al., *Phys. Lett.* **B723** 82 (2013).
 - [10] I. B. Logashenko, reported at International Workshop on e^+e^- collisions from Phi to Psi (PHIPSI13), Rome (2013).
 - [11] V. P. Druzhinin, S. I. Eidelman, S. I. Serednyakov and E. P. Solodov, *Rev. Mod. Phys.* **83**, 1545 (2011).
 - [12] A. A. Botov, private communication.

*Electronic address: A.E.Obrazovsky@inp.nsk.su

†Electronic address: S.I.Serednyakov@inp.nsk.su




FIG. 1: (a) The sum of $e^+e^- \rightarrow 3(\pi^+\pi^-)$ and $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$ cross sections with $e^+e^- \rightarrow \eta\pi^+\pi^-\pi^0$ background subtraction; (b) The sum of $e^+e^- \rightarrow p\bar{p}$ and $e^+e^- \rightarrow n\bar{n}$ cross sections; (c) The (a)+(b) sum. Only statistical errors are shown. The dashed line is the $N\bar{N}$ threshold.