

# 1D area law with ground-state degeneracy

Yichen Huang

Department of Physics, University of California, Berkeley  
Berkeley, CA 94720, USA  
yichenhuang@berkeley.edu

September 4, 2022

## Abstract

An area law is proved for 1D spin systems with ground-state degeneracy. In particular, consider a chain of  $d$ -dimensional spins governed by a Hamiltonian with nearest-neighbor interaction. Suppose the ground states are  $n$ -fold (exactly) degenerate with  $n = O(1)$ , and are separated from excited states by an energy gap  $\epsilon$ . Then for any given cut there exists a ground state such that the entanglement entropy across the cut is upper bounded by  $\tilde{O}((\log^3 d)/\epsilon)$ .

## 1 Introduction

It is conjectured that the entanglement of a region scales as its boundary (area law) in the ground state of a gapped quantum system. Hastings rigorously proves the area law in 1D (one-dimensional) spin systems by giving the upper bound  $\exp(O((\log d)/\epsilon))$  on entanglement entropy [2], where  $d$  is the dimension of each local spin and  $\epsilon$  is the energy gap. The bound is recently improved to  $\tilde{O}((\log^3 d)/\epsilon^{3/2})^1$  [1], where  $\tilde{O}(x) = O(x \text{ polylog}(x))$  hides a polylogarithmic factor.

The aforementioned proofs of the 1D area law assume a unique (nondegenerate) ground state. This paper proves an area law for 1D spin systems with ground-state degeneracy. Suppose the ground states are  $n$ -fold (exactly) degenerate with  $n = O(1)$ . Then for any given cut there exists a ground state such that the entanglement entropy across the cut is upper bounded by  $\tilde{O}((\log^3 d)/\epsilon)$ , which even slightly improves the best known result (see above) for nondegenerate systems ( $n = 1$ ).

One major technical contribution of this paper is a stronger version (Theorem 1) of the robustness theorem (Theorem 6.1 in [1]), and then the improved bound follows. All complications due to degeneracy need to be taken care of. To simplify the presentation, assume  $n = 2$  (generalization to  $n = O(1)$  is straightforward).

## 2 Approximate ground-state projector

This section generalizes the notion of approximate ground-state projector (AGSP) to nearly degenerate systems. Let  $H$  be the Hamiltonian of a 1D spin system. Let  $0 \leq \epsilon_0 \leq \epsilon_1 \leq \epsilon_2$  be the energies of the ground state  $|\psi_0\rangle$ , the first excited state  $|\psi_1\rangle$ , and the second excited state of  $H$ , respectively. Suppose  $H$  is nearly 2-fold degenerate in the sense that  $\epsilon_0 \approx \epsilon_1$ , and define  $\epsilon = \epsilon_2 - \epsilon_0$  to be the energy gap. Let  $G$  be the 2-dimensional space spanned by  $|\psi_{0,1}\rangle$ . Fix a cut, and  $R(|\psi\rangle)$  denotes the Schmidt rank of a state  $|\psi\rangle$  across the cut.

---

<sup>1</sup>The bound claimed in [1] is  $\tilde{O}((\log^3 d)/\epsilon)$ . In my opinion, the proof in [1] of this claim is incomplete. See Appendix for details.

A linear operator  $K$  is a  $(\delta, \Delta, D)$ -AGSP if (i) for any normalized state  $|\psi\rangle \in G$ ,

$$K|\psi\rangle \in G, \quad 1 - \delta \leq \|K|\psi\rangle\|^2 \leq 1; \quad (1)$$

(ii) for any normalized state  $|\psi\rangle \perp G$ ,

$$K|\psi\rangle \perp G, \quad \|K|\psi\rangle\|^2 \leq \Delta; \quad (2)$$

(iii) for any state  $|\psi\rangle$ ,

$$R(K|\psi\rangle) \leq DR(|\psi\rangle). \quad (3)$$

**Lemma 1.** *Let*

$$\mu_\psi^2 = |\langle\psi_0|\psi\rangle|^2 + |\langle\psi_1|\psi\rangle|^2. \quad (4)$$

*A  $(\delta, \Delta, D)$ -AGSP with  $D\Delta \leq 1/10$  implies the existence of a normalized product state  $|\psi\rangle$  across the cut such that*

$$\mu_\psi^2 \geq (9/10 - \delta)/D. \quad (5)$$

*Proof.* The proof is similar to that of Lemma 2.2 in [1]. Let  $K$  be such an AGSP and  $|\psi\rangle$  be the optimal normalized product state in the sense that  $|\mu_\psi| \geq |\mu_{\psi''}|$  for any normalized product state  $|\psi''\rangle$ . The optimal state  $|\psi\rangle$  exists as the set of all normalized product states is compact. The states  $|\psi\rangle$  and  $|\phi\rangle = K|\psi\rangle$  can be decomposed as

$$|\psi\rangle = \mu_\psi|\psi'\rangle + \sqrt{1 - \mu_\psi^2}|\psi^\perp\rangle, \quad |\phi\rangle = \mu'|\phi'\rangle + \mu^\perp|\phi^\perp\rangle, \quad (6)$$

where  $|\psi'\rangle, |\phi'\rangle \in G, |\psi^\perp\rangle, |\phi^\perp\rangle \in G^\perp$  are normalized states. The definition of  $K$  implies

$$(1 - \delta)\mu_\psi^2 \leq |\mu'|^2 \leq \mu_\psi^2, \quad |\mu^\perp|^2 \leq \Delta, \quad R(|\phi\rangle) \leq D. \quad (7)$$

Let  $|\Phi\rangle = |\phi\rangle/\| |\phi\rangle \|$ . Its Schmidt decomposition is given by  $|\Phi\rangle = \sum_{i=1}^D \lambda_i |L_i\rangle |R_i\rangle$  with  $\sum_i \lambda_i^2 = 1$ . Then the Cauchy-Schwarz inequality implies

$$|\mu'|/\| |\phi\rangle \| = |\langle\phi'|\Phi\rangle| \leq \sum_{i=1}^D \lambda_i |\langle\phi'|L_i\rangle| |R_i\rangle| \leq \sqrt{\sum_{i=1}^D |\langle\phi'|L_i\rangle| |R_i\rangle|^2}. \quad (8)$$

There exists an index  $i$  such that

$$\begin{aligned} \mu_\psi^2 &\geq \mu_{|L_i\rangle|R_i\rangle}^2 \geq |\langle\phi'|L_i\rangle| |R_i\rangle|^2 \geq |\mu'|^2 D^{-1} \| |\phi\rangle \|^2 = |\mu'|^2 D^{-1} (\mu'^2 + |\mu^\perp|^2)^{-1} \\ &\geq (1 - \delta)\mu_\psi^2 D^{-1} (\mu_\psi^2 + \Delta)^{-1} \Rightarrow 1 \geq (1 - \delta) D^{-1} (\mu_\psi^2 + \Delta)^{-1} \geq (1 - \delta) (D\mu_\psi^2 + 1/10)^{-1} \\ &\Rightarrow \mu_\psi^2 \geq (9/10 - \delta)/D. \end{aligned} \quad (9)$$

□

Let  $u$  be an upper bound on  $\|H\|$  (the maximum eigenvalue of  $H$ ).

**Lemma 2.** *There exists a polynomial  $C_\ell$  of degree  $\ell$  such that (i)  $C_\ell(\epsilon_0) = 1$ ; (ii)  $1 - \delta \leq C_\ell(\epsilon_1)^2 \leq 1$ , where*

$$\delta = 4\ell^2(\epsilon_1 - \epsilon_0)/(u - \epsilon_2). \quad (10)$$

(iii)  $|C_\ell(x)|^2 \leq \Delta$  for  $\epsilon_2 \leq x \leq u$ , where

$$\Delta = 4 \exp(-4\ell\sqrt{\epsilon/u}); \quad (11)$$

*Proof.* The Chebyshev polynomial  $T_\ell(x)$  is defined as

$$T_\ell(\cos \theta) = \cos(\ell\theta). \quad (12)$$

Clearly  $|T_\ell(x)| \leq 1$  for  $|x| \leq 1$ . For  $x \geq 1$ , the definition reduces to  $T_\ell(x) = \cosh(\ell t)$ , where  $t = \operatorname{arccosh} x \geq 0$ . Then,

$$T_\ell(x) \geq \exp(\ell t)/2 \geq \exp(2\ell \tanh(t/2))/2 \geq \exp(2\ell \sqrt{(x-1)/(x+1)})/2. \quad (13)$$

Moreover,

$$T'_\ell(x)/T_\ell(x) = \ell \tanh(\ell t)(dt/dx) = \ell \tanh(\ell t)/\sinh t \leq \ell(\ell t)/t = \ell^2. \quad (14)$$

Define

$$C_\ell(x) = T_\ell(f(x))/T_\ell(f(\epsilon_0)), \quad (15)$$

where  $f(x) = (u + \epsilon_2 - 2x)/(u - \epsilon_2)$ . Clearly  $|C_\ell(x)| \leq 1/T_\ell(f(\epsilon_0))$  for  $\epsilon_2 \leq x \leq u$ , and

$$T_\ell(f(\epsilon_0)) \geq \exp(2\ell \sqrt{(f(\epsilon_0) - 1)/(f(\epsilon_0) + 1)})/2 = \exp(2\ell \sqrt{\epsilon/(u - \epsilon_0)})/2 \geq \exp(2\ell \sqrt{\epsilon/u})/2. \quad (16)$$

Furthermore, there exists  $\epsilon_0 \leq \xi \leq \epsilon_1$  such that

$$\begin{aligned} C_\ell(\epsilon_1) &= C_\ell(\epsilon_0) - (\epsilon_1 - \epsilon_0)C'_\ell(\xi) \geq 1 - (\epsilon_1 - \epsilon_0)C'_\ell(\xi)/C_\ell(\xi) \geq 1 - 2\ell^2(\epsilon_1 - \epsilon_0)/(u - \epsilon_2) \\ &\Rightarrow C_\ell(\epsilon_1)^2 \geq 1 - 4\ell^2(\epsilon_1 - \epsilon_0)/(u - \epsilon_2), \end{aligned} \quad (17)$$

as  $C_\ell(x)$  is a decreasing function of  $x$  for  $x < \epsilon_2$ .  $\square$

**Lemma 3.** *For any normalized state  $|\psi\rangle$  satisfying  $\langle\psi|H|\psi\rangle \leq \epsilon_0 + b$ , there exists a normalized state  $|\phi\rangle \in G$  such that*

$$\| |\psi\rangle - |\phi\rangle \|^2 \leq 2b/\epsilon. \quad (18)$$

*Proof.* The proof is similar to that of the Markov lemma (Lemma 6.4 in [1]). The state  $|\psi\rangle$  can be decomposed as

$$|\psi\rangle = a|\phi\rangle + \sqrt{1-a^2}|\phi^\perp\rangle, \quad (19)$$

where  $|\phi\rangle \in G$ ,  $|\phi^\perp\rangle \perp G$  are normalized states, and  $a \geq 0$ . Then,

$$\begin{aligned} \epsilon_0 + (1-a^2)\epsilon &= a^2\epsilon_0 + (1-a^2)\epsilon_2 \leq \langle\psi|H|\psi\rangle \leq \epsilon_0 + b \Rightarrow 1-a^2 \leq b/\epsilon \\ &\Rightarrow \| |\psi\rangle - |\phi\rangle \|^2 = 2-2a \leq 2(1-a^2) \leq 2b/\epsilon. \end{aligned} \quad (20)$$

$\square$

### 3 Perturbation theory

Suppose the original Hamiltonian is  $H' = \sum_{i=1}^n H'_i$ , where  $0 \leq H'_i \leq 1$  acts on the spins  $i$  and  $i+1$ . Define a new Hamiltonian

$$H = H_L + H_1 + H_2 + \cdots + H_s + H_R \quad (21)$$

as (i)

$$H_L = \sum_{i=1}^m H'_i - c, \quad H_R = \sum_{i=m+s+1}^n H'_i - c', \quad (22)$$

where  $c, c'$  are the ground state energies of  $\sum_{i=1}^m H'_i, \sum_{i=m+s+1}^n H'_i$ , respectively; (ii)  $H_i = H'_{m+i}$  for  $i = 1, s$ ; (iii)

$$H_i = H'_{m+i} - c''/(s-2) \quad (23)$$

for  $2 \leq i \leq s-1$ , where  $c''$  is the ground state energy of  $\sum_{i=2}^{s-1} H'_i$ .

Clearly, (a)  $H_{L,R} \geq 0$ , and the ground state energies of  $H_{L,R}$  are 0; (b)  $0 \leq H_i \leq 1$  for  $i = 1, s$ ; (c)  $0 \leq \sum_{i=2}^{s-1} H_i \leq s-2$ , and the ground state energy of  $\sum_{i=2}^{s-1} H_i$  is 0; (c) the ground states (2-fold exactly degenerate) and the energy gaps of  $H$  and  $H'$  are the same as  $H - H'$  is a multiple of identity.

Let  $\epsilon_0 = \epsilon_1 \leq \epsilon_2 \leq \dots$  be the lowest eigenenergies of  $H$ , and define  $\epsilon = \epsilon_2 - \epsilon_0$  to be the energy gap of  $H$ . Let  $G$  be the 2-dimensional ground state space of  $H$ . Let  $P_t^{L,R}$  be the projection onto the space spanned by the eigenstates of  $H_{L,R}$  with energies at most  $t$ . Define

$$H_{L,R}^{\leq t} = P_t^{L,R} H_{L,R} P_t^{L,R} + t(1 - P_t^{L,R}). \quad (24)$$

Let  $\epsilon'_0 \leq \epsilon'_1 \leq \epsilon'_2 \leq \dots$  be the lowest eigenenergies of

$$H^{(t)} = H_L^{\leq t} + H_1 + H_2 + \dots + H_s + H_R^{\leq t}, \quad (25)$$

and define  $\epsilon' = \epsilon'_2 - \epsilon'_0$  to be the energy gap of  $H^{(t)}$ . Let  $|\phi_0^{(t)}\rangle, |\phi_1^{(t)}\rangle, |\phi_2^{(t)}\rangle$  be the ground state, the first excited state, and the second excited state of  $H^{(t)}$ . Clearly  $H^{(t)} \leq H$  and  $H^{(t)} \leq 2t + s$ .

**Lemma 4.**

$$\epsilon'_0 \leq \epsilon_0 \leq 2, \quad \epsilon'_2 \leq \epsilon_2 \leq 20. \quad (26)$$

*Proof.* Let  $|\psi_L\rangle, |\psi_M\rangle, |\psi_R\rangle, |\phi_R\rangle$  be the ground states of  $H_L, \sum_{i=2}^{s-1} H_i, H_R, \sum_{i=4}^s H_i + H_R$ , respectively. Let  $|\Psi\rangle \in G$  be a ground state of  $H$ , and

$$|\psi_{LMR}\rangle = |\psi_L\rangle|\psi_M\rangle|\psi_R\rangle, \quad |\psi_{LR}^{0,1,2}\rangle = |\psi_L\rangle|\phi_M^{0,1,2}\rangle|\phi_R\rangle, \quad (27)$$

where

$$|\phi_M^0\rangle = |00\rangle, \quad |\phi_M^1\rangle = |01\rangle, \quad |\phi_M^2\rangle = |10\rangle \quad (28)$$

are pairwise orthogonal states of the spins  $m+2$  and  $m+3$ . Then,

$$\begin{aligned} \epsilon_0 &\leq \langle \psi_{LMR} | H | \psi_{LMR} \rangle \\ &= \langle \psi_L | H_L | \psi_L \rangle + \langle \psi_M | \sum_{i=2}^{s-1} H_i | \psi_M \rangle + \langle \psi_R | H_R | \psi_R \rangle + \langle \psi_{LMR} | A | \psi_{LMR} \rangle \leq \|A\| \leq 2, \\ \langle \psi_{LR}^{0,1,2} | H | \psi_{LR}^{0,1,2} \rangle &= \langle \psi_L | H_L | \psi_L \rangle + \langle \psi_{LR}^{0,1,2} | H_1 + H_2 + H_3 | \psi_{LR}^{0,1,2} \rangle + \langle \phi_R | \sum_{i=4}^s H_i + H_R | \phi_R \rangle \\ &\leq \langle \Psi | H_L | \Psi \rangle + \langle \Psi | H_1 + H_2 + H_3 | \Psi \rangle + 3 + \langle \Psi | \sum_{i=4}^s H_i + H_R | \Psi \rangle \leq \langle \Psi | H | \Psi \rangle + 3 = \epsilon_0 + 3 \end{aligned} \quad (29)$$

Lemma 3 implies that  $|\psi_{LR}^{0,1,2}\rangle$  are close to the 2-dimensional space  $G$  (this is impossible because they are pairwise orthogonal) unless  $\epsilon_2 \leq 20$ .  $\square$

Let

$$A = H_1 + H_s. \quad (30)$$

Clearly the operators  $H_{L,R}, H - A, H^{(t)} - A$  have a common complete set of eigenstates. Let  $P_t$  be the projection onto the space spanned by the eigenstates of  $H - A$  with energies at most  $t$ . Then,

$$H_{L,R} P_t = H_{L,R}^{\leq t} P_t \Rightarrow H^{(t)} P_t = H P_t. \quad (31)$$

Let  $|\phi^{(r)}\rangle$  be an eigenstate of  $H^{(r)}$  with energy  $\epsilon^{(r)}$ .

**Lemma 5.**

$$\|(1 - P_t)|\phi^{(r)}\rangle\|^2 \leq |\langle\phi^{(r)}|(1 - P_t)AP_t|\phi^{(r)}\rangle|/(\min\{r, t\} - \epsilon^{(r)}). \quad (32)$$

*Proof.*

$$\begin{aligned} \epsilon^{(r)} &= \langle\phi^{(r)}|H^{(r)}|\phi^{(r)}\rangle = \langle\phi^{(r)}|P_t H^{(r)}|\phi^{(r)}\rangle + \langle\phi^{(r)}|(1 - P_t)H^{(r)}(1 - P_t)|\phi^{(r)}\rangle \\ &+ \langle\phi^{(r)}|(1 - P_t)H^{(r)}P_t|\phi^{(r)}\rangle \geq \epsilon^{(r)}\langle\phi^{(r)}|P_t|\phi^{(r)}\rangle + \langle\phi^{(r)}|(1 - P_t)(H^{(r)} - A)(1 - P_t)|\phi^{(r)}\rangle \\ &+ \langle\phi^{(r)}|(1 - P_t)AP_t|\phi^{(r)}\rangle + \langle\phi^{(r)}|(1 - P_t)(H^{(r)} - A)P_t|\phi^{(r)}\rangle \\ &\geq \epsilon^{(r)}\|P_t|\phi^{(r)}\rangle\|^2 + \min\{r, t\}\|(1 - P_t)|\phi^{(r)}\rangle\|^2 + \langle\phi^{(r)}|(1 - P_t)AP_t|\phi^{(r)}\rangle \\ &\geq \epsilon^{(r)}(1 - \|(1 - P_t)|\phi^{(r)}\rangle\|^2) + \min\{r, t\}\|(1 - P_t)|\phi^{(r)}\rangle\|^2 - |\langle\phi^{(r)}|(1 - P_t)AP_t|\phi^{(r)}\rangle|. \end{aligned} \quad (33)$$

The proof is completed by rearrangement.  $\square$

Let  $r, t > 100$  and  $\epsilon^{(r)} < 30$ .

**Lemma 6.**

$$\|(1 - P_t)|\phi^{(r)}\rangle\| \leq 2^{-\Omega(t)}. \quad (34)$$

*Proof.* The proof is similar to that of the truncation lemma (Lemma 6.7 in [1]). In particular, it is an induction on  $t$  by keeping  $r$  fixed. Following the notations of [1], I just point out the difference (the remainder of the two proofs is identical). Lemma 5 implies

$$\|(1 - P_t)|\phi^{(r)}\rangle\|^2 \leq |\langle\phi^{(r)}|(1 - P_t)AP_t|\phi^{(r)}\rangle|/(\min\{r, t\} - \epsilon^{(r)}), \quad (35)$$

where  $t = s + n_0 d$ . Clearly the denominator on the right-hand side is larger than 50.  $\square$

Let

$$|\phi^{(t),t}\rangle = P_t|\phi^{(t)}\rangle/\|P_t|\phi^{(t)}\rangle\|. \quad (36)$$

**Lemma 7.**

$$\langle\phi^{(t),t}|H|\phi^{(t),t}\rangle \leq \epsilon^{(t)} + 2^{-\Omega(t)}. \quad (37)$$

*Proof.*

$$\begin{aligned} \epsilon^{(t)} &= \langle\phi^{(t)}|H^{(t)}|\phi^{(t)}\rangle \geq \langle\phi^{(t)}|P_t H^{(t)}P_t|\phi^{(t)}\rangle + \langle\phi^{(t)}|P_t H^{(t)}(1 - P_t)|\phi^{(t)}\rangle \\ &+ \langle\phi^{(t)}|(1 - P_t)H^{(t)}P_t|\phi^{(t)}\rangle = \langle\phi^{(t)}|P_t H P_t|\phi^{(t)}\rangle + \langle\phi^{(t)}|P_t A(1 - P_t)|\phi^{(t)}\rangle \\ &+ \langle\phi^{(t)}|(1 - P_t)AP_t|\phi^{(t)}\rangle \geq \langle\phi^{(t)}|P_t H P_t|\phi^{(t)}\rangle - 2\|AP_t|\phi^{(t)}\rangle\| \cdot \|(1 - P_t)|\phi^{(t)}\rangle\| \\ &\geq \langle\phi^{(t)}|P_t H P_t|\phi^{(t)}\rangle - 2^{-\Omega(t)} \Rightarrow \langle\phi^{(t),t}|H|\phi^{(t),t}\rangle \leq (\epsilon^{(t)} + 2^{-\Omega(t)})/\|P_t|\phi^{(t)}\rangle\|^2 \\ &= (\epsilon^{(t)} + 2^{-\Omega(t)})/(1 - 2^{-\Omega(t)}) = \epsilon^{(t)} + 2^{-\Omega(t)}. \end{aligned} \quad (38)$$

$\square$

**Theorem 1.** Let  $t \geq c \log(1/\epsilon)$  for sufficiently large  $c = O(1)$ . (a)

$$|\epsilon'_{0,1} - \epsilon_0| \leq 2^{-\Omega(t)}; \quad (39)$$

(b) there exist  $|\psi_{0,1}^0\rangle \in G$  such that

$$\| |\psi_{0,1}^0\rangle - |\phi_{0,1}^{(t)}\rangle \|^2 \leq 2^{-\Omega(t)}; \quad (40)$$

(c)  $\epsilon' \geq \epsilon/10$ .

*Proof.* Lemma 7 implies

$$\epsilon'_{0,1} \leq \epsilon_0 \leq \langle \phi_{0,1}^{(t),t} | H | \phi_{0,1}^{(t),t} \rangle \leq \epsilon'_{0,1} + 2^{-\Omega(t)} \quad (41)$$

$$\langle \phi_2^{(t),t} | H | \phi_2^{(t),t} \rangle \leq \epsilon'_2 + 2^{-\Omega(t)} = \epsilon'_0 + \epsilon' + 2^{-\Omega(t)} \leq \epsilon_0 + \epsilon' + 2^{-\Omega(t)}. \quad (42)$$

(a) follows from (41). Lemma 3 implies the existence of  $|\psi_{0,1,2}^0\rangle \in G$  such that

$$\| |\phi_{0,1}^{(t),t}\rangle - |\psi_{0,1}^0\rangle \|^2 \leq 2^{-\Omega(t)}/\epsilon = 2^{-\Omega(t)-\log(1/\epsilon)}, \quad \| |\phi_2^{(t),t}\rangle - |\psi_2^0\rangle \|^2 \leq \epsilon'/\epsilon + 2^{-\Omega(t)}/\epsilon. \quad (43)$$

Lemma 6 implies

$$\| |\phi_{0,1}^{(t)}\rangle - |\phi_{0,1}^{(t),t}\rangle \|^2 \leq 2^{-\Omega(t)}. \quad (44)$$

(b) follows from (43), (44) as  $t \geq c \log(1/\epsilon)$ . Moreover, as  $|\phi_{0,1,2}^{(t)}\rangle$  are pairwise orthogonal,  $|\psi_{0,1,2}^0\rangle$  are approximately pairwise orthogonal (this is impossible because they are in the 2-dimensional space  $G$ ) unless  $\epsilon' \geq \epsilon/10$ .  $\square$

## 4 Area law

**Lemma 8** ([1]). *Let  $K = C_\ell(H)$ , where  $C_\ell$  is an arbitrary polynomial of degree  $\ell$ . For any state  $|\psi\rangle$ ,*

$$R(K|\psi\rangle) \leq (d\ell)^{O(\max\{\ell/s, \sqrt{\ell}\})} R(|\psi\rangle). \quad (45)$$

To construct an AGSP for  $H^{(t_0)}$ , let  $\ell = s^2$  and  $u = 2t_0 + s$ . Lemmas 2,8 imply

$$D\Delta \leq 4(ds)^{O(s)} \exp(-4s^2\sqrt{\epsilon/u}) \leq 4(ds)^{O(s)} \exp(-4s^2\sqrt{\epsilon/(2t_0 + s)}). \quad (46)$$

A little algebra shows that the condition  $D\Delta < 1/100$  for  $t_0 = O(\log(1/\epsilon))$  can be satisfied with  $s = \tilde{O}((\log^2 d)/\epsilon)$  so that  $\log D = \tilde{O}((\log^3 d)/\epsilon)$ . Let  $t_0 = O(\log(1/\epsilon) + \log \log d)$  so that

$$\delta \leq O(\ell^2 2^{-\Omega(t_0)} / (2t_0 + s)) \leq \tilde{O}(2^{-\Omega(t_0)} \log^6 d / \epsilon^3) < 1/10. \quad (47)$$

Let  $t_i = ic + t_0$  with  $c = O(1)$  so that  $\Omega(t_i) \geq i + 4$  for the  $\Omega$  in Theorem 1.

**Lemma 9.** *There exists a convergent sequence of states  $|\psi_0\rangle, |\psi_1\rangle, \dots$  satisfying: (i)*

$$|\langle \phi_0^{(t_i)} | \psi_i \rangle|^2 + |\langle \phi_1^{(t_i)} | \psi_i \rangle|^2 \geq 1 - 2^{-i-4}; \quad (48)$$

(ii) *Letting  $\ell_j = O(\sqrt{(t_j + s)/\epsilon})$ ,*

$$\log R(|\psi_0\rangle) = \tilde{O}((\log^3 d)/\epsilon), \quad \log R(|\psi_i\rangle) = \log R(|\psi_0\rangle) + \sum_{j=1}^i \tilde{O}(\ell_j \log d); \quad (49)$$

(iii) *Letting  $|\psi_\infty\rangle = \lim_{i \rightarrow +\infty} |\psi_i\rangle \in G$ ,*

$$|\langle \psi_i | \psi_\infty \rangle| \geq 1 - O(2^{-i}). \quad (50)$$

*Proof.* Theorem 1 implies the existence of  $|\psi_{0,1}^{(t_i)}\rangle \in G$  such that

$$\| |\psi_{0,1}^{(t_i)}\rangle - |\phi_{0,1}^{(t_i)}\rangle \|^2 \leq 2^{-(i+4)}. \quad (51)$$

for all  $i$ . The AGSP  $K = C_\ell(H^{(t_0)})$  constructed above implies the existence of a normalized product state  $|\psi\rangle$  such that

$$|\langle \phi_0^{(t_0)} | \psi \rangle|^2 + |\langle \phi_1^{(t_0)} | \psi \rangle|^2 \geq (9/10 - \delta)/D. \quad (52)$$

The condition  $D\Delta < 1/100$  implies that  $|\psi_0\rangle = K|\psi\rangle/\|K|\psi\rangle\|$  satisfies

$$|\langle \phi_0^{(t_0)} | \psi_0 \rangle|^2 + |\langle \phi_1^{(t_0)} | \psi_0 \rangle|^2 \geq 15/16, \quad \log R(|\psi_0\rangle) = \tilde{O}((\log^3 d)/\epsilon). \quad (53)$$

Then  $|\psi_i\rangle$  and (i) can be defined and proved by induction. (51) is a quantitative statement that the space spanned by  $|\phi_{0,1}^{(t_i)}\rangle$  is close to  $G$ . It implies that the spaces spanned by  $|\phi_{0,1}^{(t_i)}\rangle$  and spanned by  $|\phi_{0,1}^{(t_{i-1})}\rangle$  are also close. Thus the induction hypothesis implies

$$|\langle \phi_0^{(t_{i-1})} | \psi_{i-1} \rangle|^2 + |\langle \phi_1^{(t_{i-1})} | \psi_{i-1} \rangle|^2 \geq 1 - 2^{-i-3} \Rightarrow |\langle \phi_0^{(t_i)} | \psi_{i-1} \rangle|^2 + |\langle \phi_1^{(t_i)} | \psi_{i-1} \rangle|^2 \geq 1 - 2^{-i}. \quad (54)$$

Let  $\ell_i = O(\sqrt{(t_i + s)/\epsilon})$ . Lemmas 2,8 imply the existence of a  $(\delta, 1/100, D)$ -AGSP  $K$  for  $H^{(t_i)}$  with  $\log D = \tilde{O}(\ell_i \log d)$  and

$$\delta \leq O(\ell^2 2^{-\Omega(t_i)} / (t_i + s)) = O(2^{-\Omega(t_0)} / \epsilon) 2^{-i} \leq 2^{-i}/10. \quad (55)$$

Let  $|\psi_i\rangle = K|\psi_{i-1}\rangle/\|K|\psi_{i-1}\rangle\|$ . Clearly,

$$|\langle \phi_0^{(t_i)} | \psi_i \rangle|^2 + |\langle \phi_1^{(t_i)} | \psi_i \rangle|^2 \geq 1 - 2^{-i-4}, \quad (56)$$

and (ii) follows. (54) and  $\delta = O(2^{-i})$  imply

$$\| |\psi_i\rangle - |\psi_{i-1}\rangle \|^2 = O(2^{-i}). \quad (57)$$

Thus the sequence  $|\psi_0\rangle, |\psi_1\rangle, \dots$  is convergent, and

$$|\langle \psi_i | \psi_\infty \rangle| \geq 1 - \| |\psi_i\rangle - |\psi_\infty\rangle \|^2 / 2 \geq 1 - O(2^{-i}). \quad (58)$$

□

**Theorem 2.** For any given cut there exists a ground state such that the entanglement entropy across the cut is upper bounded by  $\tilde{O}((\log^3 d)/\epsilon)$ .

*Proof.* Let  $\lambda_i$  be the Schmidt coefficients of  $|\psi_\infty\rangle$  across the cut. Then

$$1 - p_i = \sum_{i=1}^{R(|\psi_i\rangle)} \lambda_i^2 \geq |\langle \psi_i | \psi_\infty \rangle|^2 \geq 1 - O(2^{-i}). \quad (59)$$

The entanglement entropy of  $|\psi_\infty\rangle$  is upper bounded by

$$\begin{aligned} & \sum_i p_i \log(R(|\psi_i\rangle) - R(|\psi_{i-1}\rangle)) - \sum_i p_i \log p_i \leq \sum_i p_i \log R(|\psi_i\rangle) + O(1) \\ &= \sum_i p_i \sum_{j=1}^i \tilde{O}(\ell_j \log d) + \sum_i p_i \log R(|\psi_0\rangle) = \sum_i p_i \sum_{j=1}^i \tilde{O}(\sqrt{(t_j + s)/\epsilon} \log d) \\ &+ O(\log R(|\psi_0\rangle)) = \tilde{O}(\sqrt{(t_0 + s)/\epsilon} \log d) + O(\log R(|\psi_0\rangle)) = \tilde{O}((\log^3 d)/\epsilon). \end{aligned} \quad (60)$$

□

## Acknowledgment

This work was supported by ARO via the DARPA OLE program.

## References

- [1] I. Arad, A. Kitaev, Z. Landau, and U. Vazirani. An area law and sub-exponential algorithm for 1D systems. *arXiv:1301.1162v1*, 2013.
- [2] M. B. Hastings. An area law for one-dimensional quantum systems. *J. Stat. Mech.*, 2007(08):P08024, 2007.

## Appendix: Details on footnote 1

For nondegenerate systems ( $n = 1$ ), the upper bound claimed in [1] on entanglement entropy is  $\tilde{O}((\log^3 d)/\epsilon)$ . In my opinion, the proof in [1] of this claim is incomplete. In particular, in Lemma 6.3  $t_0$  should be at least  $O(\epsilon_0/\epsilon^2 + 1/\epsilon)$  in order that the robustness theorem (Theorem 6.1) applies to  $H^{(t_0)}$  (the robustness theorem does not guarantee that  $H^{(t_0)}$  is gapped for  $t_0 = O(1)$ ). Then  $s = \tilde{O}((\log^2 d)/\epsilon)$  (and  $\ell = s^2$ ) does not give an AGSP for  $H^{(t_0)}$  with  $D\Delta \leq 1/2$ , but  $s = \tilde{O}((\log^2 d)/\epsilon^{3/2})$  does. A straightforward calculation shows that the upper bound  $\tilde{O}((\log^3 d)/\epsilon^{3/2})$  follows from the proof in [1].

However, I have shown that the claim in [1] is correct (Theorem 2 also holds for  $n = 1$ ). This is because Theorem 1 (as a stronger version of the robustness theorem) only requires  $t \geq c \log(1/\epsilon)$  for  $c = O(1)$ .