

ON THE BONGIORNO'S NOTION OF ABSOLUTE CONTINUITY

BEATA RANDRIANANTOANINA AND HUAQIANG XU

ABSTRACT. We show that the classes of α -absolutely continuous functions in the sense of Bongiorno coincide for all $0 < \alpha < 1$.

1. INTRODUCTION

The classical Vitali's definition says that when $\Omega \subseteq \mathbb{R}$, a function $f : \Omega \rightarrow \mathbb{R}$ is *absolutely continuous* if for all $\varepsilon > 0$, there exists $\delta > 0$ so that for every finite collection of disjoint intervals $\{[a_i, b_i]\}_{i=1}^k \subset \Omega$ we have (where \mathcal{L}^n denotes the Lebesgue measure on \mathbb{R}^n)

$$(1.1) \quad \sum_{i=1}^k \mathcal{L}^1[a_i, b_i] < \delta \Rightarrow \sum_{i=1}^k |f(a_i) - f(b_i)| < \varepsilon.$$

The study of the space of absolutely continuous functions on $[0, 1]$ and their generalizations to domains in \mathbb{R}^n is connected to the problem of finding regular subclasses of Sobolev spaces which goes back to Cesari and Calderón [6, 5]. There are several natural ways of generalizing the definition of absolute continuity for functions of several variables (cf. [7, 10, 12, 1, 11, 9]).

Recently Bongiorno [2] introduced a generalization of absolute continuity, which is simultaneously similar to Arzelà's notion of bounded variation for functions on \mathbb{R}^2 , cf. [7], and to Malý's notion of absolute continuity [11].

Definition 1.1. (Bongiorno [2]) *Let $0 < \alpha < 1$. A function $f : \Omega \rightarrow \mathbb{R}^l$, where $\Omega \subset \mathbb{R}^n$ is open, is said to be α -absolutely continuous (denoted α -AC⁽ⁿ⁾(Ω, \mathbb{R}^l) or α -AC) if for all $\varepsilon > 0$, there exists $\delta > 0$, such that for any finite collection of disjoint α -regular intervals $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}_{i=1}^k$ we have*

$$(1.2) \quad \sum_{i=1}^k \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta \Rightarrow \sum_{i=1}^k |f(\mathbf{a}_i) - f(\mathbf{b}_i)|^n < \varepsilon.$$

Here, for $\mathbf{a} \in \mathbb{R}^l$, $|\mathbf{a}|$ denotes the Euclidean norm of \mathbf{a} , and we say that an interval $[\mathbf{a}, \mathbf{b}] \stackrel{\text{def}}{=} \{\mathbf{x} = (x_\nu)_{\nu=1}^n \in \mathbb{R}^n : a_\nu \leq x_\nu \leq b_\nu, \nu = 1, \dots, n\}$ is

α -regular if

$$\frac{\mathcal{L}^n([\mathbf{a}, \mathbf{b}])}{(\max_\nu |a_\nu - b_\nu|)^n} \geq \alpha.$$

The goal of this paper is to prove that the classes α -AC coincide for all $\alpha \in (0, 1)$. We show, however, that the choice of δ in (1.2) cannot be made uniformly for all $\alpha \in (0, 1)$.

Our method uses the class 1-AC which was introduced in [8], see Definition 3.1 below.

2. PRELIMINARIES

In 2002, Hencl [9] introduced the following class of absolutely continuous functions.

Definition 2.1. (Hencl [9]) *A function $f : \Omega \rightarrow \mathbb{R}^l$ ($\Omega \subset \mathbb{R}^n$ open) is said to be in $AC_H^{(n)}(\Omega, \mathbb{R}^l)$ (briefly AC_H) if there exists $\lambda \in (0, 1)$ (equivalently, for all $\lambda \in (0, 1)$) so that for all $\varepsilon > 0$, there exists $\delta > 0$ so that for any finite collection of disjoint closed balls $\{B(\mathbf{x}_i, r_i) \subset \Omega\}_{i=1}^k$ we have*

$$\sum_{i=1}^k \mathcal{L}^n(B(\mathbf{x}_i, r_i)) < \delta \Rightarrow \sum_{i=1}^k \text{osc}^n(f, B(\mathbf{x}_i, \lambda r_i)) < \varepsilon.$$

Hencl proved that $AC_H \subset W_{loc}^{1,n}$ and that all functions in AC_H are differentiable a.e. and satisfy the Luzin (N) property and the change of variables formula.

Bongiorno [3] introduced a modification of the notion of α -absolute continuity in the spirit of Definition 2.1. To define it, we will use the following notation.

Given interval $[\mathbf{x}, \mathbf{y}] \subset \mathbb{R}^n$, we denote $|f([\mathbf{x}, \mathbf{y}])| = |f(\mathbf{y}) - f(\mathbf{x})|$, and for $0 < \lambda < 1$, we denote by ${}^\lambda[\mathbf{x}, \mathbf{y}]$ the interval with center $(\mathbf{x} + \mathbf{y})/2$ and sides of length $\lambda(y_\nu - x_\nu)$, $\nu = 1, \dots, n$.

Definition 2.2. (Bongiorno [3]) *A function $f : \Omega \rightarrow \mathbb{R}^l$ ($\Omega \subset \mathbb{R}^n$ open) is said to be in α - $AC_H^{(n)}(\Omega, \mathbb{R}^l)$ (briefly α - AC_H) if there exists $\lambda \in (0, 1)$ (equivalently, for all $\lambda \in (0, 1)$) so that for all $\varepsilon > 0$, there exists $\delta > 0$ such that for any finite collection of disjoint α -regular intervals $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}_{i=1}^k$ we have*

$$(2.1) \quad \sum_{i=1}^k \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta \Rightarrow \sum_{i=1}^k |f({}^\lambda[\mathbf{a}_i, \mathbf{b}_i])|^n < \varepsilon.$$

Bongiorno [3] proved that for all $0 < \alpha < 1$, the classes α - AC_H and AC_H coincide.

Theorem 2.3. ([3, Theorem 3]) *For every $0 < \alpha < 1$,*

$$\alpha\text{-}AC_H^{(n)}(\Omega, \mathbb{R}^l) = AC_H^{(n)}(\Omega, \mathbb{R}^l).$$

3. THE MAIN RESULT

In [8] we introduced an analog of Bongiorno's notion for $\alpha = 1$, which will be an important tool for the main result of this paper.

Definition 3.1. *We say that a function $f : \Omega \rightarrow \mathbb{R}^l$ ($\Omega \subset \mathbb{R}^n$ open) is 1-absolutely continuous, denoted $1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l)$ or $1\text{-}AC$, if for every $\varepsilon > 0$, there exists $\delta > 0$, such that for any finite collection of disjoint 1-regular intervals $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}_{i=1}^k$ we have*

$$\sum_{i=1}^k \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta \Rightarrow \sum_{i=1}^k |f(\mathbf{a}_i) - f(\mathbf{b}_i)|^n < \varepsilon.$$

Note that for all $\alpha < 1$, every α -regular interval is also 1-regular. Thus for all $\alpha < 1$,

$$\alpha\text{-}AC \subseteq 1\text{-}AC.$$

Properties of the class $1\text{-}AC$ were studied in [8], where it was shown that it contains many pathological functions. In particular functions in $1\text{-}AC$ don't need to be continuous and even if they are differentiable a.e. they do not need to belong to the Sobolev space $W_{loc}^{1,n}$. However, the class $1\text{-}AC$ is very useful for characterizing the classes $\alpha\text{-}AC$.

Theorem 3.2. *For all $0 < \alpha < 1$ we have*

$$\alpha\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) = 1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \cap AC_H^{(n)}(\Omega, \mathbb{R}^l).$$

Proof. By definitions, $\alpha\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \subset 1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l)$, and by [2, Theorem 7], $\alpha\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \subset AC_H^{(n)}(\Omega, \mathbb{R}^l)$, which proves that

$$\alpha\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \subset 1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \cap AC_H^{(n)}(\Omega, \mathbb{R}^l).$$

For the other direction, let $f \in 1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l) \cap AC_H^{(n)}(\Omega, \mathbb{R}^l)$ and $\varepsilon > 0$. Let $\beta = (\frac{\alpha}{2-\alpha})^n$ and $\lambda = 1 - \frac{\alpha}{2}$. By Theorem 2.3, $f \in \beta\text{-}AC_H^{(n)}(\Omega, \mathbb{R}^l)$ and there exists $\delta_1 > 0$ so that for each finite family of disjoint β -regular intervals $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}$

$$\sum_i \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta_1 \Rightarrow \sum_i |f^\lambda([\mathbf{a}_i, \mathbf{b}_i])|^n < \frac{\varepsilon}{3^{n+1}}.$$

Since $f \in 1\text{-}AC^{(n)}(\Omega, \mathbb{R}^l)$, there exists $\delta_2 > 0$ so that for each finite family of disjoint 1-regular intervals $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}$

$$\sum_i \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta_2 \Rightarrow \sum_i |f([\mathbf{a}_i, \mathbf{b}_i])|^n < \frac{\varepsilon}{3^{n+1}}.$$

Let $\delta = \min\{\delta_1, \delta_2\}$ and $\{[\mathbf{a}_i, \mathbf{b}_i] \subset \Omega\}$ be a finite family of disjoint α -regular intervals with $\sum_i \mathcal{L}^n([\mathbf{a}_i, \mathbf{b}_i]) < \delta$. We will show that $\sum_i |f([\mathbf{a}_i, \mathbf{b}_i])|^n < \varepsilon$.

For each $[\mathbf{a}_i, \mathbf{b}_i]$ from this family, let $\eta_i = \max_{1 \leq \nu \leq n} (b_{i\nu} - a_{i\nu})$. Since $[\mathbf{a}_i, \mathbf{b}_i]$ is an α -regular interval, we get that

$$\prod_{j=1}^n \left(\frac{b_{ij} - a_{ij}}{\eta_i} \right) \geq \alpha.$$

Thus, for every $j = 1, \dots, n$,

$$\frac{b_{ij} - a_{ij}}{\eta_i} \geq \alpha.$$

Let

$$\mathbf{c}_i = \left(a_{ij} + \frac{\alpha}{4} \eta_i \right)_{j=1}^n, \quad \mathbf{d}_i = \left(b_{ij} - \frac{\alpha}{4} \eta_i \right)_{j=1}^n.$$

Then, for every $j = 1, \dots, n$, $c_{ij} < d_{ij}$, and for every i

$$\max_j (d_{ij} - c_{ij}) = \max_j \left(b_{ij} - a_{ij} - \frac{\alpha}{2} \eta_i \right) = \left(1 - \frac{\alpha}{2} \right) \eta_i.$$

Hence

$$\begin{aligned} \frac{\mathcal{L}^n([\mathbf{c}_i, \mathbf{d}_i])}{\max_j (d_{ij} - c_{ij})^n} &= \prod_{j=1}^n \left(\frac{b_{ij} - a_{ij} - \frac{\alpha}{2} \eta_i}{\left(1 - \frac{\alpha}{2} \right) \eta_i} \right) \\ &\geq \prod_{j=1}^n \left(\frac{\alpha \eta_i - \frac{\alpha}{2} \eta_i}{\left(1 - \frac{\alpha}{2} \right) \eta_i} \right) = \left(\frac{\alpha}{2 - \alpha} \right)^n = \beta. \end{aligned}$$

Thus interval $[\mathbf{c}_i, \mathbf{d}_i]$ is β -regular.

Let

$$\begin{aligned} \overline{\mathbf{c}}_i &= \left(\frac{a_{ij} + b_{ij}}{2} - \frac{1}{\lambda} \left(\frac{a_{ij} + b_{ij}}{2} - a_{ij} - \frac{\alpha}{4} \eta_i \right) \right)_{j=1}^n \\ &= \left(\frac{c_{ij} + d_{ij}}{2} - \frac{1}{\lambda} \left(\frac{d_{ij} - c_{ij}}{2} \right) \right)_{j=1}^n, \\ \overline{\mathbf{d}}_i &= \left(\frac{a_{ij} + b_{ij}}{2} + \frac{1}{\lambda} \left(b_{ij} - \frac{\alpha}{4} \eta_i - \frac{a_{ij} + b_{ij}}{2} \right) \right)_{j=1}^n \\ &= \left(\frac{c_{ij} + d_{ij}}{2} + \frac{1}{\lambda} \left(\frac{d_{ij} - c_{ij}}{2} \right) \right)_{j=1}^n. \end{aligned}$$

Then $[\overline{\mathbf{c}}_i, \overline{\mathbf{d}}_i] \subset [\mathbf{a}_i, \mathbf{b}_i]$, and

$$\lambda[\overline{\mathbf{c}}_i, \overline{\mathbf{d}}_i] = [\mathbf{c}_i, \mathbf{d}_i],$$

$$|f(\mathbf{a}_i, \mathbf{b}_i)|^n \leq 3^n [|f(\mathbf{a}_i, \mathbf{c}_i)|^n + |f(\mathbf{c}_i, \mathbf{d}_i)|^n + |f(\mathbf{d}_i, \mathbf{b}_i)|^n],$$

some $\alpha \in (0, 1)$. Thus (1.2) holds for any finite collection of nontrivial intervals. By [8, Theorem 3.1] this implies that f is constant. \square

Remark 3.4. In [4] Bongiorno introduced another generalization of absolute continuity, a class $AC_\Lambda^n(\Omega, \mathbb{R}^m)$, briefly AC_Λ , which strictly contains the class AC_H , and such that all functions in AC_Λ are differentiable a.e. and satisfy the Luzin (N) property, but $AC_\Lambda^n(\Omega, \mathbb{R}^l) \not\subset W_{loc}^{1,n}(\Omega, \mathbb{R}^l)$. It would be interesting to determine what are the classes $1-AC \cap AC_\Lambda$ and $1-AC \cap AC_\Lambda \cap W_{loc}^{1,n}$ (since, by [8], $1-AC \not\subset W_{loc}^{1,n}$).

REFERENCES

- [1] D. APATSIDIS, S. A. ARGYROS, AND V. KANELLOPOULOS, *Hausdorff measures and functions of bounded quadratic variation*, Trans. Amer. Math. Soc., 363 (2011), pp. 4225–4262.
- [2] D. BONGIORNO, *Absolutely continuous functions in \mathbb{R}^n* , J. Math. Anal. Appl., 303 (2005), pp. 119–134.
- [3] D. BONGIORNO, *On the Hencl’s notion of absolute continuity*, J. Math. Anal. Appl., 350 (2009), pp. 562–567.
- [4] D. BONGIORNO, *On the problem of regularity in the Sobolev space $W_{loc}^{1,n}$* , Topology Appl., 156 (2009), pp. 2986–2995.
- [5] A. P. CALDERÓN, *On the differentiability of absolutely continuous functions*, Rivista Mat. Univ. Parma, 2 (1951), pp. 203–213.
- [6] L. CESARI, *Sulle funzioni assolutamente continue in due variabili*, Ann. Scuola Norm. Super. Pisa (2), 10 (1941), pp. 91–101.
- [7] J. A. CLARKSON AND C. R. ADAMS, *On definitions of bounded variation for functions of two variables*, Trans. Amer. Math. Soc., 35 (1933), pp. 824–854.
- [8] M. DYMOND, B. RANDRIANANTOANINA AND H. XU, *On interval based generalizations of absolute continuity for functions on \mathbb{R}^n* , preprint.
- [9] S. HENCL, *On the notions of absolute continuity for functions of several variables*, Fund. Math., 173 (2002), pp. 175–189.
- [10] E. W. HOBSON, *The theory of functions of a real variable and the theory of Fourier’s series. Vol. I*, Dover Publications Inc., New York, N.Y., 1958.
- [11] J. MALÝ, *Absolutely continuous functions of several variables*, J. Math. Anal. Appl., 231 (1999), pp. 492–508.
- [12] T. RADO AND P. V. REICHELDERFER, *Continuous transformations in analysis. With an introduction to algebraic topology*, Springer-Verlag, Berlin, 1955.

DEPARTMENT OF MATHEMATICS, MIAMI UNIVERSITY, OXFORD, OH 45056, USA

E-mail address: randrib@miamioh.edu

DEPARTMENT OF MATHEMATICS, MIAMI UNIVERSITY, OXFORD, OH 45056, USA

E-mail address: xuhuaqiang1990@gmail.com