

NECESSARY AND SUFFICIENT CONDITIONS FOR THE SOLVABILITY OF INVERSE PROBLEM FOR A CLASS OF DIRAC OPERATORS

KH. R. MAMEDOV^{1,*} AND O. AKCAY²

ABSTRACT. In this paper, we consider a problem for the first order Dirac differential equations system with spectral parameter dependent in boundary condition. The asymptotic behaviors of eigenvalues, eigenfunctions and normalizing numbers of this system are investigated. The expansion formula with respect to eigenfunctions is obtained and Parseval equality is given. The main theorem on necessary and sufficient conditions for the solvability of inverse problem is proved and the algorithm of reconstruction of potential from spectral data (the sets of eigenvalues and normalizing numbers) is given.

1. INTRODUCTION

We consider the boundary value problem generated by one dimensional Dirac differential equations system

$$By' + \Omega(x)y = \lambda y, \quad 0 < x < \pi \quad (1)$$

with boundary condition

$$\begin{aligned} U(y) &:= y_1(0) = 0, \\ V(y) &:= (\lambda + h_1)y_1(\pi) + h_2y_2(\pi) = 0, \end{aligned} \quad (2)$$

where

$$B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \Omega(x) = \begin{pmatrix} p(x) & q(x) \\ q(x) & -p(x) \end{pmatrix}, \quad y = \begin{pmatrix} y_1(x) \\ y_2(x) \end{pmatrix},$$

$p(x)$, $q(x)$ are real valued functions in $L_2(0, \pi)$, λ is a spectral parameter, h_1 and h_2 are real numbers, $h_2 > 0$.

In the finite interval, the inverse problems for Dirac differential equations system by different spectral characteristics (e.g., one spectrum and norming constants or two spectra or spectral function) are solved in [1, 2, 3, 4, 5, 6, 7]. When spectral parameter contained in boundary condition, reconstruction of potential from spectral function is carried out in [7] and the uniqueness of inverse problem for Dirac operator according to Weyl function is worked in [8]. Inverse spectral problem for Dirac operator with potentials belong entrywise to $L_p(0, 1)$; for some $p \in [1, \infty)$, was studied in [3] and in this work, not only Gelfand-Levitan-Marchenko method but also Krein method [9] was used. Using Weyl-Titchmarsh function, direct and inverse problems for Dirac type-system were investigated in [10, 11]. For weighted Dirac systems, inverse spectral problems was examined in [12].

2010 *Mathematics Subject Classification.* 34A55; 34L40.

Key words and phrases. Dirac operator, eigenvalues and normalizing numbers, inverse problem, necessary and sufficient conditions.

*corresponding author.

On a positive half line, inverse scattering problem for a system of Dirac equations of order $2n$ is completely solved in [13] and when boundary condition involving spectral parameter, for Dirac operator, inverse scattering problem worked in [14, 15]. The applications of Dirac differential equations system has been widespread in various areas of physics, such as [17, 18, 19] ever since Dirac equation was discovered to be associated with nonlinear wave equation in [16].

This paper is organized as follows: In section 2, the asymptotic formulas of eigenvalues, eigenfunctions and normalizing numbers of the boundary value problem (1), (2) are investigated. In section 3, by the eigenfunctions, completeness theorem is proved and expansion formula is obtained. Parseval equality is given. In section 4, the main equation namely Gelfand-Levitan-Marchenko type equation is derived. In section 5, we show that the boundary value problem (1),(2) can be uniquely determined from its eigenvalues and normalizing numbers. Finally in section 6, the solution of inverse problem is obtained. Let's express this more clearly. We can state the inverse problem for a system of Dirac equations in the following way: let λ_n and α_n , ($n \in \mathbb{Z}$) are respectively eigenvalues and normalizing numbers of boundary value problem (1), (2) and the quantities $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$) are called spectral data. Knowing the spectral data $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$) to indicate a method of determining the potential $\Omega(x)$ and to find necessary and sufficient conditions for λ_n and α_n , ($n \in \mathbb{Z}$) to be the spectral data of a problem (1),(2), for this, we derive differential equation, Parseval equality and boundary conditions. The main theorem on the necessary and sufficient conditions for the solvability of inverse problem is proved and then algorithm of the construction of the function $\Omega(x)$ by spectral data is given. Note that throughout this paper, we use the following notation: $\tilde{\phi}$ denotes the transposed matrix of ϕ .

2. ASYMPTOTIC FORMULAS OF EIGENVALUES, EIGENFUNCTION AND NORMALIZING NUMBERS

The inner product in Hilbert space $H = L_2(0, \pi; \mathbb{C}^2) \oplus \mathbb{C}$ is defined by

$$\langle Y, Z \rangle := \int_0^\pi [y_1(x)\bar{z}_1(x) + y_2(x)\bar{z}_2(x)] dx + \frac{1}{h_2}y_3\bar{z}_3,$$

where

$$Y = \begin{pmatrix} y_1(x) \\ y_2(x) \\ y_3 \end{pmatrix} \in H, \quad Z = \begin{pmatrix} z_1(x) \\ z_2(x) \\ z_3 \end{pmatrix} \in H.$$

Let us define

$$L(Y) := \begin{pmatrix} l(y) \\ -h_1y_1(\pi) - h_2y_2(\pi) \end{pmatrix}$$

with

$$D(L) = \left\{ Y \mid Y = (y_1(x), y_2(x), y_3)^T \in H, y_1(x), y_2(x) \in AC[0, \pi], \right. \\ \left. y_3 = y_1(\pi), l(y) \in L_2(0, \pi; \mathbb{C}^2), U(y) = 0, V(y) = 0 \right\}$$

where

$$l(y) = \begin{pmatrix} y_2' + p(x)y_1 + q(x)y_2 \\ -y_1' + q(x)y_1 - p(x)y_2 \end{pmatrix}.$$

Let $\varphi(x, \lambda)$ and $\psi(x, \lambda)$ be solutions of the system (1) satisfying the initial conditions

$$\varphi(0, \lambda) = \begin{pmatrix} 0 \\ -1 \end{pmatrix}, \quad \psi(\pi, \lambda) = \begin{pmatrix} h_2 \\ -\lambda - h_1 \end{pmatrix}. \quad (3)$$

The solution $\varphi(x, \lambda)$ has the following representation (see [8])

$$\varphi(x, \lambda) = \varphi_0(x, \lambda) + \int_0^x K(x, t)\varphi_0(t, \lambda)dt, \quad (4)$$

where $\varphi_0(x, \lambda) = \begin{pmatrix} \sin \lambda x \\ -\cos \lambda x \end{pmatrix}$, $K_{ij}(x, \cdot) \in L_2(0, \pi)$, $i, j = 1, 2$ for fixed $x \in [0, \pi]$ and $K(x, t)$ is solution of the problem

$$\begin{aligned} BK'_x(x, t) + K'_t(x, t)B &= -\Omega(x)K(x, t), \\ \Omega(x) &= K(x, x)B - BK(x, x), \\ K_{11}(x, 0) &= K_{21}(x, 0) = 0. \end{aligned} \quad (5)$$

The formula (5) gives the relation between the kernel $K(x, t)$ and the coefficient of $\Omega(x)$ of the equation (1).

The characteristic function $\Delta(\lambda)$ of the problem L is

$$\Delta(\lambda) := W[\varphi(x, \lambda), \psi(x, \lambda)] = \varphi_2(x, \lambda)\psi_1(x, \lambda) - \varphi_1(x, \lambda)\psi_2(x, \lambda), \quad (6)$$

where $W[\varphi(x, \lambda), \psi(x, \lambda)]$ is Wronskian of the solutions $\varphi(x, \lambda)$ and $\psi(x, \lambda)$ and independent of $x \in [0, \pi]$. The zeros of $\Delta(\lambda)$ coincide with the eigenvalues λ_n of problem L . The functions $\varphi(x, \lambda)$ and $\psi(x, \lambda)$ are eigenfunctions and there exists a sequence β_n such that

$$\psi(x, \lambda_n) = \beta_n \varphi(x, \lambda_n), \quad \beta_n \neq 0. \quad (7)$$

Normalizing numbers are

$$\alpha_n := \int_0^\pi \left(|\varphi_1(x, \lambda_n)|^2 + |\varphi_2(x, \lambda_n)|^2 \right) dx + \frac{1}{h_2} |\varphi_1(\pi, \lambda_n)|^2.$$

The following relation holds (see [20]):

$$\dot{\Delta}(\lambda_n) = \beta_n \alpha_n, \quad (8)$$

where $\dot{\Delta}(\lambda) = \frac{d}{d\lambda} \Delta(\lambda)$.

Theorem 1. *i) The eigenvalues λ_n , ($n \in \mathbb{Z}$) of boundary value problem (1), (2) are*

$$\lambda_n = \lambda_n^0 + \epsilon_n, \quad \{\epsilon_n\} \in l_2, \quad (9)$$

where $\lambda_n^0 = n$ are zeros of function $\lambda \sin \lambda \pi$. For the large n , the eigenvalues are simple.

ii) The eigenfunctions of the boundary value problem can be represented in the form

$$\varphi(x, \lambda_n) = \begin{pmatrix} \sin nx \\ -\cos nx \end{pmatrix} + \begin{pmatrix} \zeta_n^{(1)}(x) \\ \zeta_n^{(2)}(x) \end{pmatrix}, \quad (10)$$

where $\sum_{n=-\infty}^{\infty} \left\{ \left| \zeta_n^{(1)}(x) \right|^2 + \left| \zeta_n^{(2)}(x) \right|^2 \right\} \leq C$, in here C is a positive number.

iii) Normalizing numbers of the problem (1), (2) are as follows

$$\alpha_n = \pi + \tau_n, \quad \{\tau_n\} \in l_2. \quad (11)$$

Proof. i) Using (4), as $|\lambda| \rightarrow \infty$ uniformly in $x \in [0, \pi]$ the following asymptotic formulas are obtain:

$$\begin{aligned}\varphi_1(x, \lambda) &= \sin \lambda x + O\left(\frac{1}{|\lambda|}e^{Im\lambda|x}\right), \\ \varphi_2(x, \lambda) &= -\cos \lambda x + O\left(\frac{1}{|\lambda|}e^{Im\lambda|x}\right).\end{aligned}\tag{12}$$

Substituting the asymptotic formulas (12) into

$$\Delta(\lambda) = (\lambda + h_1)\varphi_1(\pi, \lambda) + h_2\varphi_2(\pi, \lambda),$$

we get as $|\lambda| \rightarrow \infty$

$$\Delta(\lambda) = \lambda \sin \lambda \pi + O\left(e^{Im\lambda|\pi}\right).\tag{13}$$

Denote

$$G_\delta := \{\lambda : |\lambda - n| \geq \delta, \quad n = 0, \pm 1, \pm 2, \dots\},$$

where δ is a sufficiently small positive number. There exists a positive number C_δ such that (see [21])

$$|\sin \lambda \pi| \geq C_\delta e^{Im\lambda|\pi}, \quad \lambda \in G_\delta.\tag{14}$$

On the other hand from (13), as $|\lambda| \rightarrow \infty$

$$\Delta(\lambda) - \lambda \sin \lambda \pi = O\left(e^{Im\lambda|\pi}\right).\tag{15}$$

Therefore, on infinitely expanding contours

$$\Gamma_n = \left\{ \lambda : |\lambda| = n + \frac{1}{2} \right\}$$

for sufficiently large n , using (14) and (15) we have

$$|\Delta(\lambda) - \lambda \sin \lambda \pi| < |\lambda| |\sin \lambda \pi|, \quad \lambda \in \Gamma_n.$$

Applying the Rouché theorem, it is obtained that the number of zeros of the function $\{\Delta(\lambda) - \lambda \sin \lambda \pi\} + \lambda \sin \lambda \pi = \Delta(\lambda)$ inside the counter Γ_n coincides with the number of zeros of function $\lambda \sin \lambda \pi$. Moreover, using the Rouché theorem, there exist only one zero λ_n of the function $\Delta(\lambda)$ in the circle $\gamma_n = \{\lambda : |\lambda - n| < \delta\}$ is concluded. Since $\delta > 0$ is arbitrary, we have

$$\lambda_n = n + \epsilon_n, \quad \lim_{n=\pm\infty} \epsilon_n = 0.\tag{16}$$

Substituting (16) into (13), we get $\sin \epsilon_n \pi = O\left(\frac{1}{n}\right)$, i.e. $\epsilon_n = O\left(\frac{1}{n}\right)$. Thus $\{\epsilon_n\} \in l_2$ is found. For the large n , the eigenvalues are simple. In fact, $\alpha_n \beta_n = \dot{\Delta}(\lambda_n)$ and $\alpha_n \neq 0, \beta_n \neq 0$, we get $\dot{\Delta}(\lambda_n) \neq 0$.

ii) Putting (9) into (4), we have

$$\varphi(x, \lambda_n) = \begin{pmatrix} \sin nx \\ -\cos nx \end{pmatrix} + \begin{pmatrix} \zeta_n^{(1)}(x) \\ \zeta_n^{(2)}(x) \end{pmatrix},$$

where

$$\begin{aligned}\zeta_n^{(1)}(x) &= \sin nx [\cos \epsilon_n x - 1] + \cos nx \sin \epsilon_n x + \\ &+ \int_0^x K_{11}(x, t) \sin(n + \epsilon_n)t dt - \int_0^x K_{12}(x, t) \cos(n + \epsilon_n)t dt,\end{aligned}$$

$$\begin{aligned}\zeta_n^{(2)}(x) &= -\cos nx [\cos \epsilon_n x - 1] + \sin nx \sin \epsilon_n x + \\ &\quad + \int_0^x K_{21}(x, t) \sin(n + \epsilon_n)t dt - \int_0^x K_{22}(x, t) \cos(n + \epsilon_n)t dt.\end{aligned}$$

Since $K_{ij}(\pi, \cdot) \in L_2(0, \pi)$, $i, j = 1, 2$, according to (see [21], pp.66) we get

$$\left\{ \int_0^\pi K_{i1}(\pi, t) \sin \lambda_n t dt \right\} \in l_2, \quad \left\{ \int_0^\pi K_{i2}(\pi, t) \cos \lambda_n t dt \right\} \in l_2.$$

Using these relations and $\{\epsilon_n\} \in l_2$, we obtain

$$\sup_{0 \leq x \leq \pi} \sum_{n=-\infty}^{\infty} \left\{ \left| \zeta_n^{(1)}(x) \right|^2 + \left| \zeta_n^{(2)}(x) \right|^2 \right\} < +\infty.$$

iii) Using the definition of normalizing numbers and (10)

$$\begin{aligned}\alpha_n &= \int_0^\pi \left(|\varphi_1(x, \lambda_n)|^2 + |\varphi_2(x, \lambda_n)|^2 \right) dx + \frac{1}{h_2} |\varphi_1(\pi, \lambda_n)|^2 \\ &= \int_0^\pi dx + \tau_n = \pi + \tau_n,\end{aligned}$$

where

$$\begin{aligned}\tau_n &= \int_0^\pi \left(\zeta_n^{(1)}(x) \right)^2 dx + \int_0^\pi \left(\zeta_n^{(2)}(x) \right)^2 dx + 2 \int_0^\pi \sin nx \zeta_n^{(1)}(x) dx - \\ &\quad - 2 \int_0^\pi \cos nx \zeta_n^{(2)}(x) dx + \frac{1}{h_2} \left(\zeta_n^{(1)}(\pi) \right)^2.\end{aligned}$$

Since $\{\zeta_n^{(1)}(x)\} \in l_2$ and $\{\zeta_n^{(2)}(x)\} \in l_2$, we find $\{\tau_n\} \in l_2$. The theorem is proved. \square

Proposition 2. *The specification of the eigenvalues λ_n , ($n \in \mathbb{Z}$) uniquely determines the characteristic function $\Delta(\lambda)$ by formula*

$$\Delta(\lambda) = -\pi(\lambda_0^2 - \lambda^2) \prod_{n=1}^{\infty} \frac{(\lambda_n^2 - \lambda^2)}{n^2}. \quad (17)$$

Proof. Since the function $\Delta(\lambda)$ is entire function of order 1, from Hadamard's theorem (see [22]), using (13) we obtain (17). \square

3. PARSEVAL EQUALITY

Theorem 3. *a) The system of eigenfunctions $\varphi(x, \lambda_n)$, ($n \in \mathbb{Z}$) of boundary value problem (1), (2) is complete in space $L_2(0, \pi; \mathbb{C}^2)$.*

b) Let $f(x) \in D(L)$. Then the expansion formula

$$f(x) = \sum_{n=-\infty}^{\infty} a_n \varphi(x, \lambda_n), \quad (18)$$

$$a_n = \frac{1}{\alpha_n} \langle f(x), \varphi(x, \lambda_n) \rangle$$

is valid and the series converges uniformly with respect to $x \in [0, \pi]$. For $f(x) \in L_2(0, \pi; \mathbb{C}^2)$ series (18) converges in $L_2(0, \pi; \mathbb{C}^2)$, moreover, Parseval equality holds

$$\|f\|^2 = \sum_{n=-\infty}^{\infty} \alpha_n |a_n|^2. \quad (19)$$

Proof. a) Denote

$$G(x, t, \lambda) := -\frac{1}{\Delta(\lambda)} \begin{cases} \psi(x, \lambda) \tilde{\varphi}(t, \lambda), & x \geq t \\ \varphi(x, \lambda) \tilde{\psi}(t, \lambda), & t \geq x. \end{cases}$$

Consider the function

$$Y(x, \lambda) = \int_0^\pi G(x, t, \lambda) f(t) dt \quad (20)$$

which gives solution of boundary value problem

$$\begin{aligned} BY' + \Omega(x)Y &= \lambda Y + f(x), \\ Y_1(0, \lambda) &= 0, \\ (\lambda + h_1) Y_1(\pi, \lambda) + h_2 Y_2(\pi, \lambda) &= 0. \end{aligned} \quad (21)$$

Using (7) and (8), we have

$$\psi(x, \lambda_n) = \frac{\dot{\Delta}(\lambda_n)}{\alpha_n} \varphi(x, \lambda_n).$$

Using this expression, we find

$$\operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = -\frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^\pi \tilde{\varphi}(t, \lambda_n) f(t) dt. \quad (22)$$

Let $f(x) \in L_2(0, \pi; \mathbb{C}^2)$ be such that

$$\langle f(x), \varphi(x, \lambda_n) \rangle = \int_0^\pi \tilde{\varphi}(t, \lambda_n) f(t) dt = 0. \quad (23)$$

Then, it follows from (22) that $\operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = 0$. Consequently $Y(x, \lambda)$ is entire function with respect to λ for each fixed $x \in [0, \pi]$. Taking into account

$$|\Delta(\lambda)| \geq C_\delta |\lambda| e^{|Im\lambda|\pi} \quad (24)$$

and the following equalities being valid according to ([21], Lemma 1.3.1)

$$\lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} \exp(-|Im\lambda|x) \left| \int_0^x \tilde{\varphi}(t, \lambda) f(t) dt \right| = 0, \quad (25)$$

$$\lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} \frac{1}{|\lambda|} \exp(-|Im\lambda|(\pi-x)) \left| \int_x^\pi \tilde{\psi}(t, \lambda) f(t) dt \right| = 0, \quad (26)$$

we find

$$\lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} |Y(x, \lambda)| = 0.$$

Hence $Y(x, \lambda) \equiv 0$ is obtained. From here and (21), $f(x) = 0$ a.e. on $(0, \pi)$.

b) Since $\varphi(x, \lambda)$ and $\psi(x, \lambda)$ are solution of the problem (1), (2),

$$\begin{aligned} Y(x, \lambda) &= -\frac{1}{\lambda \Delta(\lambda)} \psi(x, \lambda) \int_0^x \left\{ -\frac{\partial}{\partial t} \tilde{\varphi}(t, \lambda) B + \tilde{\varphi}(t, \lambda) \Omega(t) \right\} f(t) dt \\ &\quad - \frac{1}{\lambda \Delta(\lambda)} \varphi(x, \lambda) \int_x^\pi \left\{ -\frac{\partial}{\partial t} \tilde{\psi}(t, \lambda) B + \tilde{\psi}(t, \lambda) \Omega(t) \right\} f(t) dt \end{aligned}$$

can be written. Integrating by parts and using (6),

$$Y(x, \lambda) = -\frac{1}{\lambda}f(x) - \frac{1}{\lambda}Z(x, \lambda) \quad (27)$$

is obtained, where

$$Z(x, \lambda) = \frac{1}{\Delta(\lambda)} \left\{ \psi(x, \lambda) \int_0^x \tilde{\varphi}(t, \lambda) B f'(t) dt + \varphi(x, \lambda) \int_x^\pi \tilde{\psi}(t, \lambda) B f'(t) dt \right. \\ \left. + \psi(x, \lambda) \int_0^x \tilde{\varphi}(t, \lambda) \Omega(t) f(t) dt + \varphi(x, \lambda) \int_x^\pi \tilde{\psi}(t, \lambda) \Omega(t) f(t) dt \right\}.$$

By applying (25) and (26), we have

$$\lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} |Z(x, \lambda)| = 0, \quad \lambda \in G_\delta. \quad (28)$$

Now, we integrate of $Y(x, \lambda)$ with respect to λ over the contour Γ_N with oriented counter clockwise as follows:

$$I_n(x) = \frac{1}{2\pi i} \oint_{\Gamma_N} Y(x, \lambda) d\lambda,$$

where $\Gamma_N = \{\lambda : |\lambda| = N + \frac{1}{2}\}$, N is sufficiently large natural number. Using residue theorem, we get

$$I_n(x) = \sum_{n=-N}^N \operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = - \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^\pi \tilde{\varphi}(t, \lambda_n) f(t) dt.$$

On the other hand, considering the equation (27)

$$f(x) = \sum_{n=-N}^N a_n \varphi(x, \lambda_n) + \epsilon_N(x) \quad (29)$$

is found, where

$$\epsilon_N(x) = -\frac{1}{2\pi i} \oint_{\Gamma_N} \frac{1}{\lambda} Z(x, \lambda) d\lambda$$

and

$$a_n = \frac{1}{\alpha_n} \int_0^\pi \tilde{\varphi}(t, \lambda_n) f(t) dt.$$

It follows from (28) that,

$$\lim_{N \rightarrow \infty} \max_{0 \leq x \leq \pi} |\epsilon_N(x)| = 0.$$

Thus, by going over in (29) to the limit as $N \rightarrow \infty$ the expansion formula (18) is obtained. The system $\{\varphi(x, \lambda_n)\}$, ($n \in \mathbb{Z}$) is complete and orthogonal in $L_2(0, \pi; \mathbb{C}^2)$. Therefore, it forms orthogonal basis in $L_2(0, \pi; \mathbb{C}^2)$ and Parseval equality (19) holds. \square

4. MAIN EQUATION

Theorem 4. For each fixed $x \in (0, \pi]$ the kernel $K(x, t)$ satisfies the following equation:

$$F(x, t) + K(x, t) + \int_0^x K(x, \xi)F(\xi, t)d\xi = 0, \quad 0 < t < x, \quad (30)$$

where

$$F(x, t) = \sum_{n=-\infty}^{\infty} \left[\frac{1}{\alpha_n} \varphi_0(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right]. \quad (31)$$

Proof. Using the transformation operators (see [23]), according to (4) we have

$$\varphi_0(x, \lambda) = \varphi(x, \lambda) + \int_0^x H(x, t)\varphi(t, \lambda)dt. \quad (32)$$

It follows from (4) and (32) that

$$\begin{aligned} & \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) = \\ & = \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi_0(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) + \int_0^x K(x, \xi) \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi_0(\xi, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) d\xi \end{aligned}$$

and

$$\begin{aligned} & \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) = \\ & = \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}(t, \lambda_n) + \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^t \tilde{\varphi}(\xi, \lambda_n) \tilde{H}(t, \xi) d\xi. \end{aligned}$$

Using the last two equalities, we obtain

$$\begin{aligned} & \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] = \\ & = \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] + \\ & + \int_0^x K(x, \xi) \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(\xi, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] d\xi + \\ & + \int_0^x K(x, \xi) \sum_{n=-N}^N \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(\xi, \lambda_n^0) d\xi - \\ & - \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^t \tilde{\varphi}(\xi, \lambda_n) \tilde{H}(t, \xi) d\xi \end{aligned}$$

or

$$\Phi_n(x, t) = I_{N1}(x, t) + I_{N2}(x, t) + I_{N3}(x, t) + I_{N4}(x, t), \quad (33)$$

where

$$\begin{aligned}\Phi_n(x, t) &:= \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right], \\ I_{N1}(x, t) &:= \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right], \\ I_{N2}(x, t) &:= \int_0^x K(x, \xi) \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(\xi, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] d\xi, \\ I_{N3}(x, t) &:= \int_0^x K(x, \xi) \sum_{n=-N}^N \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(\xi, \lambda_n^0) d\xi, \\ I_{N4}(x, t) &:= \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^t \tilde{\varphi}(\xi, \lambda_n) \tilde{H}(t, \xi) d\xi.\end{aligned}$$

Let $f(x)$ be an absolutely continuous function. Then using expansion formula (18),

$$\begin{aligned}\lim_{N \rightarrow \infty} \int_0^\pi \Phi_N(x, t) f(t) dt &= \lim_{N \rightarrow \infty} \int_0^\pi \sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \tilde{\varphi}(t, \lambda_n) f(t) dt - \\ &- \lim_{N \rightarrow \infty} \int_0^\pi \sum_{n=-N}^N \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) f(t) dt = 0,\end{aligned}\quad (34)$$

$$\begin{aligned}\lim_{N \rightarrow \infty} \int_0^\pi I_{N1}(x, t) f(t) dt &= \\ &= \lim_{N \rightarrow \infty} \int_0^\pi \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(x, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \frac{1}{\pi} \varphi_0(x, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] f(t) dt \\ &= \int_0^\pi F(x, t) f(t) dt,\end{aligned}\quad (35)$$

$$\begin{aligned}\lim_{N \rightarrow \infty} \int_0^\pi I_{N2}(x, t) f(t) dt &= \lim_{N \rightarrow \infty} \int_0^\pi \int_0^x K(x, \xi) \sum_{n=-N}^N \left[\frac{1}{\alpha_n} \varphi_0(\xi, \lambda_n) \tilde{\varphi}_0(t, \lambda_n) - \right. \\ &\left. - \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(t, \lambda_n^0) \right] f(t) d\xi dt = \int_0^\pi \left(\int_0^x K(x, \xi) F(x, \xi) d\xi \right) f(t) dt,\end{aligned}\quad (36)$$

$$\begin{aligned}\lim_{N \rightarrow \infty} \int_0^\pi I_{N3}(x, t) f(t) dt &= \\ &= \lim_{N \rightarrow \infty} \int_0^\pi \int_0^x K(x, \xi) \sum_{n=-N}^N \frac{1}{\pi} \varphi_0(\xi, \lambda_n^0) \tilde{\varphi}_0(\xi, \lambda_n^0) f(t) d\xi dt \\ &= \int_0^x K(x, \xi) f(\xi) d\xi,\end{aligned}\quad (37)$$

$$\begin{aligned}
& \lim_{N \rightarrow \infty} \int_0^\pi I_{N4}(x, t) f(t) dt = \\
& = \lim_{N \rightarrow \infty} \int_0^\pi \left(\sum_{n=-N}^N \frac{1}{\alpha_n} \varphi(x, \lambda_n) \int_0^t \tilde{\varphi}(\xi, \lambda_n) \tilde{H}(t, \xi) d\xi \right) f(t) dt \\
& = \int_x^\pi H(t, x) f(t) dt. \tag{38}
\end{aligned}$$

We put $K(x, t) = H(x, t) = 0$ for $x < t$. Using (33), (34), (35), (36), (37) and (38) we obtain

$$\begin{aligned}
& \int_0^\pi F(x, t) f(t) dt + \int_0^\pi \left(\int_0^x K(x, \xi) F(\xi, t) d\xi \right) f(t) dt + \\
& + \int_0^x K(x, \xi) f(\xi) d\xi - \int_x^\pi H(t, x) f(t) dt = 0.
\end{aligned}$$

Since $f(x)$ can be chosen arbitrarily,

$$F(x, t) + K(x, t) + \int_0^x K(x, \xi) F(\xi, t) d\xi - H(t, x) = 0.$$

When $t < x$, this equality implies (30). \square

5. THEOREM FOR THE SOLUTION OF THE INVERSE PROBLEM

Lemma 5. *For each fixed $x \in (0, \pi]$ main equation (30) has a unique solution $K(x, t) \in L_2(0, x)$.*

Proof. It suffices to prove that homogeneous equation

$$g(t) + \int_0^x g(s) F(s, t) ds = 0$$

has only trivial solution $g(t) = 0$. Let $g(t) = (g_1(t), g_2(t)) \in L_2(0, x)$ be a solution of the above equation and $g(t) = 0$ for $t \in (x, \pi)$. Then

$$(g(t), g(t)) + \left(\int_0^x g(s) F(s, t) ds, g(t) \right) = 0.$$

Using the expression (31), we get

$$\begin{aligned}
& \int_0^x (g_1^2(t) + g_2^2(t)) dt + \\
& + \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \int_0^x \left[\int_0^x (g_1(s) \sin \lambda_n(s) - g_2(s) \cos \lambda_n(s)) ds \right] g_1(t) \sin \lambda_n t dt \\
& - \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \int_0^x \left[\int_0^x (g_1(s) \sin \lambda_n(s) - g_2(s) \cos \lambda_n(s)) ds \right] g_2(t) \cos \lambda_n t dt \\
& - \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \int_0^x \left[\int_0^x (g_1(s) \sin \lambda_n^0(s) - g_2(s) \cos \lambda_n^0(s)) ds \right] g_1(t) \sin \lambda_n^0 t dt \\
& + \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \int_0^x \left[\int_0^x (g_1(s) \sin \lambda_n^0(s) - g_2(s) \cos \lambda_n^0(s)) ds \right] g_2(t) \cos \lambda_n^0 t dt
\end{aligned}$$

$$\begin{aligned}
&= \int_0^x (g_1^2(t) + g_2^2(t)) dt + \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^x [g_1(t) \sin \lambda_n t - g_2(t) \cos \lambda_n t] dt \right)^2 \\
&\quad - \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \left(\int_0^x [g_1(t) \sin \lambda_n^0 t - g_2(t) \cos \lambda_n^0 t] dt \right)^2 = 0.
\end{aligned}$$

Thus, it follows from the last relation that

$$\begin{aligned}
&\int_0^x (g_1^2(t) + g_2^2(t)) dt + \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^x g(t) \varphi_0(t, \lambda_n) dt \right)^2 - \\
&\quad - \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \left(\int_0^x g(t) \varphi_0(t, \lambda_n^0) dt \right)^2 = 0.
\end{aligned}$$

Using Parseval equality,

$$\|g\|^2 = \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \left(\int_0^x g(t) \varphi_0(t, \lambda_n^0) dt \right)^2$$

we have

$$\sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^x g(t) \varphi_0(t, \lambda_n) dt \right)^2 = 0.$$

Since $\alpha_n > 0$ and from statement a) in Theorem 3, the system $\{\varphi_0(t, \lambda_n)\}$, ($n \in \mathbb{Z}$) is complete in $L_2(0, \pi; \mathbb{C}^2)$, we obtain $g(t) = 0$. \square

Theorem 6. Let $L(\Omega(x), h_1, h_2)$ and $\hat{L}(\hat{\Omega}(x), \hat{h}_1, \hat{h}_2)$ be two boundary value problems and

$$\lambda_n = \hat{\lambda}_n, \quad \alpha_n = \hat{\alpha}_n, \quad (n \in \mathbb{Z}).$$

Then

$$\Omega(x) = \hat{\Omega}(x) \text{ a.e. on } (0, \pi), \quad h_1 = \hat{h}_1, \quad h_2 = \hat{h}_2.$$

Proof. According to (31), $F(x, t) = \hat{F}(x, t)$. Then, from the main equation (30), we have $K(x, t) = \hat{K}(x, t)$. It follows from (5) that $\Omega(x) = \hat{\Omega}(x)$ a.e. on $(0, \pi)$. Taking into account (4), we find $\varphi(x, \lambda_n) = \hat{\varphi}(x, \lambda_n)$. In consideration of (13), we get $\hat{\Delta}(\lambda_n) \equiv \hat{\Delta}(\lambda_n)$ and from (8), $\beta_n = \hat{\beta}_n$. Thus, using (3) and (7), we obtain $h_1 = \hat{h}_1$, $h_2 = \hat{h}_2$. \square

6. SOLUTION OF INVERSE PROBLEM

Let the real numbers $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$) of the form (9) and (11) be given. Using these numbers, we construct the function $F(x, t)$ by the formulas (31) and determine $K(x, t)$ from the main equation (30).

Now, let us construct the functions $\varphi(x, \lambda)$ by the formula (4), $\Omega(x)$ by the formula (5), $\Delta(\lambda)$ by the formula (17) and β_n by the formula (8) respectively, i.e.,

$$\begin{aligned}
\varphi(x, \lambda) &:= \varphi_0(x, \lambda) + \int_0^x K(x, t) \varphi(t, \lambda) dt, \\
\Omega(x) &:= K(x, x) B - BK(x, x), \\
\Delta(\lambda) &:= -\pi(\lambda_0^2 - \lambda^2) \prod_{n=1}^{\infty} \frac{(\lambda_n^2 - \lambda^2)}{n^2},
\end{aligned}$$

$$\beta_n := \frac{\dot{\Delta}(\lambda_n)}{\alpha_n} \neq 0.$$

The function $F(x, t)$ can be rewritten as follows:

$$F(x, t) = \frac{1}{2} [a(x-t) + a(x+t)T],$$

where

$$a(x) = \sum_{n=-\infty}^{\infty} \left[\frac{1}{\alpha_n} \begin{pmatrix} \cos \lambda_n x & -\sin \lambda_n x \\ \sin \lambda_n x & \cos \lambda_n x \end{pmatrix} - \frac{1}{\pi} \begin{pmatrix} \cos nx & -\sin nx \\ \sin nx & \cos nx \end{pmatrix} \right]$$

and $T = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$. It is shown analogously in ([24], Lemma 1.5.4) that the function $a(x) \in W_2^1[0, 2\pi]$.

6.1. Derivation of the Differential Equation.

Theorem 7. *The relations hold:*

$$B\varphi'(x, \lambda) + \Omega(x)\varphi(x, \lambda) = \lambda\varphi(x, \lambda), \quad (39)$$

$$\varphi_1(0, \lambda) = 0, \quad \varphi_2(0, \lambda) = -1. \quad (40)$$

Proof. Differentiating on x and y the main equation (30) respectively, we get

$$F'_x(x, t) + K'_x(x, t) + K(x, x)F(x, t) + \int_0^x K'_x(x, \xi)F(\xi, t)d\xi = 0, \quad (41)$$

$$F'_t(x, t) + K'_t(x, t) + \int_0^x K(x, \xi)F'_t(\xi, t)d\xi = 0. \quad (42)$$

It follows from (31) that

$$F'_t(x, t)B + BF'_x(x, t) = 0. \quad (43)$$

Since $F(x, 0)BS = 0$, where $S = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$, using the main equation (30), we obtain

$$K(x, 0)BS = 0, \quad (44)$$

or

$$K_{11}(x, 0) = K_{21}(x, 0) = 0.$$

Multiplying the equation (41) on the left by B , we get

$$BF'_x(x, t) + BK'_x(x, t) + BK(x, x)F(x, t) + \int_0^x BK'_x(x, \xi)F(\xi, t)d\xi = 0,$$

and multiplying the equation (42) on the right by B , we find

$$F'_t(x, t)B + K'_t(x, t)B + \int_0^x K(x, \xi)F'_t(\xi, t)Bd\xi = 0.$$

Adding the last two equalities and using (43)

$$\begin{aligned} & BK'_x(x, t) + BK(x, x)F(x, t) + \int_0^x BK'_x(x, \xi)F(\xi, t)d\xi = \\ & = -K'_t(x, t)B + \int_0^x K(x, \xi)BF'_\xi(\xi, t)d\xi \equiv I(x, t). \end{aligned} \quad (45)$$

Integrating by parts in (45) and from (44),

$$I(x, t) = -K'_t(x, t)B + K(x, x)BF(x, t) - \int_0^x K'_\xi(x, \xi)BF(\xi, t)d\xi \quad (46)$$

is obtained. Substituting (46) into (45), we have

$$\begin{aligned} & BK'_x(x, t) + BK(x, x)F(x, t) + K'_t(x, t)B - K(x, x)BF(x, t) + \\ & + \int_0^x [BK'_x(x, \xi) + K'_\xi(x, \xi)B] F(\xi, t)d\xi = 0. \end{aligned} \quad (47)$$

Multiplying (30) on the left by $\Omega(x)$ in form of (5) and adding to (47),

$$\begin{aligned} & BK'_x(x, t) + K'_t(x, t)B + \Omega(x)K(x, t) + \\ & + \int_0^x [BK'_x(x, \xi) + K'_\xi(x, \xi)B + \Omega(x)K(x, \xi)] F(\xi, t)d\xi = 0 \end{aligned} \quad (48)$$

is obtained. Setting

$$J(x, t) := BK'_x(x, t) + K'_t(x, t)B + \Omega(x)K(x, t),$$

we can rewrite equation (48) as follows

$$J(x, t) + \int_0^x J(x, \xi)F(\xi, t)d\xi = 0. \quad (49)$$

According to Lemma 5, homogeneous equation (48) has only trivial solution, i.e.,

$$BK'_x(x, t) + K'_t(x, t)B + \Omega(x)K(x, t) = 0, \quad 0 < t < x. \quad (50)$$

Now, differentiating (4) and multiplying on the left by B , we have

$$\begin{aligned} B\varphi'_x(x, \lambda) &= \lambda B \begin{pmatrix} \cos \lambda x \\ \sin \lambda x \end{pmatrix} + BK(x, x) \begin{pmatrix} \sin \lambda x \\ -\cos \lambda x \end{pmatrix} + \\ &+ \int_0^x BK'_x(x, t) \begin{pmatrix} \sin \lambda t \\ -\cos \lambda t \end{pmatrix} dt. \end{aligned} \quad (51)$$

On the other hand, multiplying (4) on the left by λ and integrating by parts and using (44), we find

$$\begin{aligned} \lambda\varphi(x, \lambda) &= \lambda B \begin{pmatrix} \cos \lambda x \\ \sin \lambda x \end{pmatrix} + K(x, x)B \begin{pmatrix} \sin \lambda x \\ -\cos \lambda x \end{pmatrix} - \\ &- \int_0^x K'_t(x, t)B \begin{pmatrix} \sin \lambda t \\ -\cos \lambda t \end{pmatrix} dt. \end{aligned} \quad (52)$$

It follows from (51) and (52) that

$$\begin{aligned} \lambda\varphi(x, \lambda) &= B\varphi'_x(x, \lambda) + [K(x, x)B - BK(x, x)] \begin{pmatrix} \sin \lambda x \\ -\cos \lambda x \end{pmatrix} - \\ &- \int_0^x [K'_t(x, t)B + BK'_x(x, t)] \begin{pmatrix} \sin \lambda t \\ -\cos \lambda t \end{pmatrix} dt. \end{aligned}$$

Taking into account (5) and (50),

$$B\varphi'_x(x, \lambda) + \Omega(x)\varphi(x, \lambda) = \lambda\varphi(x, \lambda)$$

is obtained. For $x = 0$, from (4) we get (40). \square

6.2. Derivation of Parseval Equality.

Theorem 8. For any $g(x) \in L_2(0, \pi; \mathbb{C}^2)$, the following relation holds:

$$\|g\|_{L_2} = \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^\pi \tilde{\varphi}(t, \lambda_n) g(t) dt \right)^2. \quad (53)$$

Proof. Denote

$$Q(\lambda) := \int_0^\pi \tilde{\varphi}(t, \lambda) g(t) dt.$$

It follows from (4) that

$$Q(\lambda) := \int_0^\pi \tilde{\varphi}_0(t, \lambda) h(t) dt,$$

where

$$h(t) = g(t) + \int_t^\pi \tilde{K}(s, t) g(s) ds. \quad (54)$$

Similarly, in view of (32),

$$g(t) = h(t) + \int_x^\pi \tilde{H}(s, t) h(s) ds, \quad (55)$$

where for the kernel $H(x, t)$, we have the identity

$$\tilde{H}(x, t) = F(t, x) + \int_0^x K(x, \xi) F(\xi, t) d\xi. \quad (56)$$

Using (54), we get

$$\begin{aligned} \int_0^\pi F(x, t) h(t) dt &= \int_0^\pi F(x, t) \left[g(t) + \int_t^\pi \tilde{K}(u, t) g(u) du \right] dt = \\ &= \int_0^\pi \left[F(x, t) + \int_0^t F(x, u) \tilde{K}(t, u) du \right] g(t) dt \\ &= \int_0^x \left[F(x, t) + \int_0^t F(x, u) \tilde{K}(t, u) du \right] g(t) dt + \\ &\quad + \int_x^\pi \left[F(x, t) + \int_0^t F(x, u) \tilde{K}(t, u) du \right] g(t) dt. \end{aligned}$$

It follows from (30) and (56) that

$$\int_0^\pi F(x, t) h(t) dt = \int_0^x H(x, t) g(t) dt - \int_x^\pi \tilde{K}(t, x) g(t) dt. \quad (57)$$

From (31) and Parseval equality, we obtain

$$\begin{aligned} \int_0^\pi [h_1^2(t) + h_2^2(t)] dt + \int_0^\pi \int_0^\pi \tilde{h}(x) F(x, t) h(t) dt dx &= \\ &= \int_0^\pi [h_1^2(t) + h_2^2(t)] dt + \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^\pi \tilde{\varphi}_0(t, \lambda_n) h(t) dt \right)^2 - \\ &\quad - \sum_{n=-\infty}^{\infty} \frac{1}{\pi} \left(\int_0^\pi \tilde{\varphi}_0(t, \lambda_n^0) h(t) dt \right)^2 \\ &= \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^\pi \tilde{\varphi}_0(t, \lambda_n) h(t) dt \right)^2 = \sum_{n=-\infty}^{\infty} \frac{Q^2(\lambda_n)}{\alpha_n}. \end{aligned}$$

Taking into account (57), we have

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{Q^2(\lambda_n)}{\alpha_n} &= \int_0^{\pi} [h_1^2(t) + h_2^2(t)] dt + \\ &+ \int_0^{\pi} \tilde{h}(x) \left(\int_0^x H(x,t)g(t)dt \right) dx - \int_0^{\pi} \tilde{h}(x) \left(\int_x^{\pi} \tilde{K}(t,x)g(t)dt \right) dx \\ &= \int_0^{\pi} [h_1^2(t) + h_2^2(t)] dt + \int_0^{\pi} \left(\int_t^{\pi} \tilde{h}(x)H(x,t)dx \right) g(t)dt - \\ &- \int_0^{\pi} \tilde{h}(x) \left(\int_x^{\pi} \tilde{K}(t,x)g(t)dt \right) dx, \end{aligned}$$

whence by formulas (54) and (55),

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{Q^2(\lambda_n)}{\alpha_n} &= \int_0^{\pi} [h_1^2(t) + h_2^2(t)] dt + \\ &+ \int_0^{\pi} (\tilde{g}(t) - \tilde{h}(t)) g(t)dt - \int_0^{\pi} \tilde{h}(x) (h(x) - g(x)) dx \\ &= \int_0^{\pi} (g_1^2(t) + g_2^2(t)) dt \end{aligned}$$

is obtained, i.e., the relation (53) is valid. \square

Corollary 9. For any function $f(x)$ and $g(x) \in L_2(0, \pi; \mathbb{C}^2)$, the relation holds:

$$\int_0^{\pi} \tilde{g}(x)f(x)dx = \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^{\pi} \tilde{g}(t)\varphi(t, \lambda_n)dt \right) \left(\int_0^{\pi} \tilde{\varphi}(t, \lambda_n)f(t)dt \right). \quad (58)$$

Lemma 10. For any $f(x) \in W_2^1[0, \pi]$, the expansion formula

$$f(x) = \sum_{n=-\infty}^{\infty} c_n \varphi(x, \lambda_n) \quad (59)$$

is valid, where

$$c_n = \frac{1}{\alpha_n} \int_0^{\pi} \tilde{\varphi}(x, \lambda_n) f(x) dx.$$

Proof. Consider the series

$$f^*(x) = \sum_{n=-\infty}^{\infty} c_n \varphi(x, \lambda_n), \quad (60)$$

where

$$c_n := \frac{1}{\alpha_n} \int_0^{\pi} \tilde{\varphi}(x, \lambda_n) f(x) dx. \quad (61)$$

Using Theorem 7 and integrating by parts, we get

$$\begin{aligned} c_n &= \frac{1}{\alpha_n \lambda_n} \int_0^{\pi} \left[-\frac{\partial}{\partial x} \tilde{\varphi}(x, \lambda_n) B + \tilde{\varphi}(x, \lambda_n) \Omega(x) \right] f(x) dx \\ &= -\frac{1}{\alpha_n \lambda_n} [\tilde{\varphi}(\pi, \lambda_n) B f(\pi) - \tilde{\varphi}(0, \lambda_n) B f(0)] + \\ &+ \frac{1}{\alpha_n \lambda_n} \int_0^{\pi} \tilde{\varphi}(x, \lambda_n) [B f'(x) + \Omega(x) f(x)] dx. \end{aligned}$$

Applying the asymptotic formulas in Theorem 1, we find $\{c_n\} \in l_2$. Consequently the series (60) converges absolutely and uniformly on $[0, \pi]$. According to (58) and (61), we have

$$\begin{aligned} \int_0^\pi \tilde{g}(x)f(x)dx &= \sum_{n=-\infty}^{\infty} \frac{1}{\alpha_n} \left(\int_0^\pi \tilde{g}(t)\varphi(t, \lambda_n)dt \right) \left(\int_0^\pi \tilde{\varphi}(t, \lambda_n)f(t)dt \right) \\ &= \sum_{n=-\infty}^{\infty} c_n \left(\int_0^\pi \tilde{g}(t)\varphi(t, \lambda_n)dt \right) = \int_0^\pi \tilde{g}(t) \left(\sum_{n=-\infty}^{\infty} c_n\varphi(t, \lambda_n) \right) = \int_0^\pi \tilde{g}(t)f^*(t)dt. \end{aligned}$$

Since $g(x)$ is arbitrary, $f(x) = f^*(x)$ is obtained, i.e., the expansion formula (59) is found. \square

6.3. Derivation of Boundary Condition.

Lemma 11. *The following equality holds:*

$$\sum_{n=-\infty}^{\infty} \frac{\varphi(x, \lambda_n)}{\alpha_n \beta_n} = 0. \quad (62)$$

Proof. Using residue theorem, we get

$$\sum_{n=-\infty}^{\infty} \frac{\varphi(x, \lambda_n)}{\alpha_n \beta_n} = \sum_{n=-\infty}^{\infty} \frac{\varphi(x, \lambda_n)}{\Delta(\lambda_n)} = \sum_{n=-\infty}^{\infty} \operatorname{Res}_{\lambda=\lambda_n} \frac{\varphi(x, \lambda)}{\Delta(\lambda)} = \frac{1}{2\pi i} \int_{\Gamma_N} \frac{\varphi(x, \lambda)}{\Delta(\lambda)} d\lambda, \quad (63)$$

where $\Gamma_N = \{\lambda : |\lambda| = N + \frac{1}{2}\}$. From (17) and ([21], Lemma 3.4.2),

$$\Delta(\lambda) = \lambda \sin \lambda\pi + O(e^{|\operatorname{Im}\lambda|\pi}). \quad (64)$$

We denote $G_\delta = \{\lambda : |\lambda - n| \geq \delta, \ n = 0, \pm 1, \pm 2, \dots\}$ for some small fixed $\delta > 0$ and $|\sin \lambda\pi| \geq C_\delta e^{|\operatorname{Im}\lambda|\pi}$, $\lambda \in G_\delta$, where C_δ positive number. Therefore, we have

$$|\Delta(\lambda)| \geq C_\delta |\lambda| e^{|\operatorname{Im}\lambda|\pi}, \quad \lambda \in G_\delta.$$

Using this inequality and (12), we obtain (62). \square

Theorem 12. *The following relation is valid:*

$$(\lambda_n + h_1)\varphi_1(\pi, \lambda_n) + h_2\varphi_2(\pi, \lambda_n) = 0. \quad (65)$$

Proof. From (62), we can write for any $n_0 \in \mathbb{Z}$

$$\frac{\varphi(x, \lambda_{n_0})}{\alpha_{n_0}} = - \sum_{\substack{n=-\infty \\ n \neq n_0}}^{\infty} \frac{\beta_{n_0}\varphi(x, \lambda_n)}{\alpha_n \beta_n} \quad (66)$$

Let $m \neq n_0$ be any fixed number and $f(x) = \varphi(x, \lambda_k)$. Then substituting (66) in (59)

$$\varphi(x, \lambda_k) = \sum_{\substack{n=-\infty \\ n \neq n_0}}^{\infty} c_{nk}\varphi(x, \lambda_n),$$

where

$$c_{nk} = \frac{1}{\alpha_n} \int_0^\pi \left[\tilde{\varphi}(t, \lambda_n) - \frac{\beta_{n_0}}{\beta_n} \tilde{\varphi}(t, \lambda_{n_0}) \right] \varphi(t, \lambda_k) dt.$$

The system of functions $\{\varphi_0(x, \lambda_n)\}$, $(n \in \mathbb{Z})$ is orthogonal in $L_2(0, \pi; \mathbb{C}^2)$. Then by (4), the system of functions $\{\varphi(x, \lambda_n)\}$, $(n \in \mathbb{Z})$ is orthogonal in $L_2(0, \pi; \mathbb{C}^2)$ as well. Therefore, $c_{nk} = \delta_{nk}$, where δ_{nk} is Kronecker delta. Let us define

$$a_{nk} := \int_0^\pi \tilde{\varphi}(t, \lambda_n) \varphi(t, \lambda_k) dt. \quad (67)$$

Using this expression, we have for $n \neq k$

$$a_{kk} - \frac{\beta_n}{\beta_k} a_{nk} = \alpha_k. \quad (68)$$

It follows from (67) that $a_{nk} = a_{kn}$. Taking into account this equality and (68),

$$\beta_k^2 (\alpha_k - a_{kk}) = \beta_n^2 (\alpha_n - a_{nn}) = H, \quad k \neq n,$$

where H is a constant. Then, we have

$$\int_0^\pi \tilde{\varphi}(t, \lambda_n) \varphi(t, \lambda_n) dt = \alpha_n - \frac{H}{\beta_n^2}$$

and

$$\int_0^\pi \tilde{\varphi}(t, \lambda_k) \varphi(t, \lambda_n) dt = -\frac{H}{\beta_k \beta_n}, \quad k \neq n.$$

It is easily obtained that for $k \neq n$,

$$\begin{aligned} & \int_0^\pi [\varphi_1(x, \lambda_k) \varphi_1(x, \lambda_n) + \varphi_2(x, \lambda_k) \varphi_2(x, \lambda_n)] dx = \\ & = \frac{1}{(\lambda_k - \lambda_n)} [\varphi_2(\pi, \lambda_k) \varphi_1(\pi, \lambda_n) - \varphi_1(\pi, \lambda_k) \varphi_2(\pi, \lambda_n)] = -\frac{H}{\beta_k \beta_n}. \end{aligned}$$

According to the last equation, for $n \neq k$,

$$\beta_k \varphi_2(\pi, \lambda_k) \beta_n \varphi_1(\pi, \lambda_n) - \beta_k \varphi_1(\pi, \lambda_k) \beta_n \varphi_2(\pi, \lambda_n) = -H(\lambda_k - \lambda_n). \quad (69)$$

We denote

$$D_n := \beta_n \varphi_1(\pi, \lambda_n), \quad E_n := \beta_n \varphi_2(\pi, \lambda_n). \quad (70)$$

Then, we can rewrite equation (69) as follows

$$D_k E_n - E_k D_n = H(\lambda_k - \lambda_n), \quad n \neq k. \quad (71)$$

Let i, j, k, n be pairwise distinct integers, then we get

$$\begin{aligned} D_k E_n - E_k D_n &= H(\lambda_k - \lambda_n), \\ D_n E_i - E_n D_i &= H(\lambda_n - \lambda_i), \\ D_i E_k - E_i D_k &= H(\lambda_i - \lambda_k). \end{aligned}$$

Adding them together, we find

$$D_n(E_i - E_k) + E_n(D_k - D_i) = E_i D_k - D_i E_k.$$

In this equation, replacing n by j , we get another equation

$$D_j(E_i - E_k) + E_j(D_k - D_i) = E_i D_k - D_i E_k.$$

Subtracting the last two equation,

$$(D_n - D_j)(E_i - E_k) = (D_i - D_k)(E_n - E_j).$$

In the case of $E_n = E_j$, for some $n, j \in \mathbb{Z}$, then $E_n = \text{const}$. From (71), $D_n = \mu_1 \lambda_n + \mu_2$. In the case of $E_n \neq E_j$, then we obtain $D_n = \mu_1 \lambda_n + \mu_2$ and $E_n =$

$\mu_3\lambda_n + \mu_4$, where in both cases $\mu_1, \mu_2, \mu_3, \mu_4$ are constant. Therefore, using these relation in (70), we find

$$\beta_n\varphi_1(\pi, \lambda_n) = \mu_1\lambda_n + \mu_2, \quad \beta_n\varphi_2(\pi, \lambda_n) = \mu_3\lambda_n + \mu_4.$$

Using

$$\varphi_1(\pi, \lambda_n) = O\left(\frac{1}{n}\right), \quad \varphi_2(\pi, \lambda_n) = (-1)^{n+1} + O\left(\frac{1}{n}\right),$$

$\lambda_n = n + O\left(\frac{1}{n}\right)$ and $\beta_n = n(-1)^n + O(1)$ derived from (8) and (64), we obtain $\mu_1 = 0, \mu_3 = -1$. Denoting $h_2 := \mu_2$ and $h_1 := -\mu_4$,

$$h_2\varphi_2(\pi, \lambda_n) = -(\lambda_n + h_1)\varphi_1(\pi, \lambda_n), \quad n \in \mathbb{Z}$$

is obtained and it follows from (71) that $H = h_2$. \square

Thus, we have proved the following theorem:

Theorem 13. *For the sequences $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$), to be the spectral data for a certain boundary value problem $L(\Omega(x), h_1, h_2)$ of the form (1), (2) with $\Omega(x) \in L_2(0, \pi)$ it is necessary and sufficient that the relations (9) and (11) hold.*

Algorithm of the construction of the function $\Omega(x)$ by spectral data $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$) follows from the proof of the theorem:

- 1) By the given numbers $\{\lambda_n, \alpha_n\}$, ($n \in \mathbb{Z}$) the function $F(x, t)$ is constructed by formula (31),
- 2) The function $K(x, t)$ is found from equation (30),
- 3) $\Omega(x)$ calculated by the formula (5).

Acknowledgement 14. *This work is supported by The Scientific and Technological Research Council of Turkey (TÜBİTAK).*

REFERENCES

- [1] M. G. Gasyimov, T.T. Dzabiev, Solution of the inverse problem by two spectra for the Dirac equation on a finite interval, Dokl. Akad. Nauk Azerbaidzan. SSR 22(7) (1966) 3-6.
- [2] T. T. Dzabiev, The inverse problem for the Dirac equation with a singularity, Dokl. Akad. Nauk Azerbaidzan. SSR 22(11) (1966) 8-12.
- [3] S. Albeverio, R. Hryniv, Ya. Mykytyuk, Inverse spectral problems for Dirac operators with summable potentials. Russ. J. Math. Phys. 12 (2005) 406-423.
- [4] Ya. V. Mykytyuk, D. V. Puyda, Inverse spectral problems for Dirac operators on a finite interval, J. Math. Anal. Appl. 386(1) (2012) 177-194.
- [5] M. Horvath, On the inverse spectral theory of Schrödinger and Dirac operators, Trans. Amer. Math. Soc. 353 (2001) 4155-4171.
- [6] K. Mochizuki, I. Trooshin, Inverse problem for interior spectral data of the Dirac operator on a finite interval, Publ. RIMS, Kyoto Univ. 38 (2002) 387-395.
- [7] S. G. Mamedov, The inverse boundary value problem on a finite interval for Dirac's system of equations, Azerbaidzan. Gos. Univ. Ucen. Zap. Ser. Fiz-Mat. Nauk (5) (1975) 61-67.
- [8] R. Kh. Amirov, B. Keskin, A. S. Ozkan, Direct and inverse problems for the Dirac operator with spectral parameter linearly contained in boundary condition, Ukrainian Math. J. 61(9) (2009) 1365-1379.
- [9] M. G. Krein, On integral equations generating differential equations of the second order, Dokl. Akad. SSSR. 97 (1954) 21-24.
- [10] A. Sakhnovich, Skew-self-adjoint discrete and continuous Dirac-type systems: inverse problems and Borg-Marchenko theorems, Inverse Problems 22(6) (2006) 2083-2101.
- [11] B. Fritzsche, B. Kirstein, I. Ya. Roitberg, A. Sakhnovich, Skew-self-adjoint Dirac system with a rectangular matrix potential: Weyl theory, direct and inverse problems, Integral Equations and Operator Theory 74(2) (2012) 163-187.

- [12] B. A. Watson, Inverse spectral problems for weighted Dirac systems, *Inverse Problems* 15 (1999) 793-805.
- [13] M. G. Gasymov, The inverse scattering problem for a system of Dirac equations of order $2n$, *Trans. Moskow Math. Soc.* 19 (1968) 41-119.
- [14] Kh. R. Mamedov, A. Çöl, On an inverse scattering problem for a class Dirac operator with discontinuous coefficient and nonlinear dependence on the spectral parameter in the boundary condition, *Mathematical Methods in the Applied Sciences* 35(14) (2012) 1712-1720.
- [15] A. Çöl, Kh. R. Mamedov, On an inverse scattering problem for a class of Dirac operators with spectral parameter in the boundary condition, *J. Math. Anal. Appl.*, 393 (2012) 470-478.
- [16] M. J. Ablowitz, D. J. Kaup, A. C. Newell, H. Segur, Nonlinear-evolution equations of physical significance, *Phys. Rev. Lett.* 31 (1973) 125-127.
- [17] D. G. Shepelsky, An inverse spectral problem for a Dirac-type operator with sewing, in: *Dynamical Systems and Complex Analysis*, Naukova Dumka, Kiev, 1992, pp.104-112 (Russian).
- [18] A. B. de Monvel, D. Shepelsky, Initial boundary value problem for the mKdV equation on a finite interval, *Ann. Inst. Fourier, Grenoble*, 54(5) (2004) 1477-1495.
- [19] B. Thaller, *The Dirac Equation*, Springer, Berlin, 1992.
- [20] Kh. R. Mamedov, Ö. Akçay, Inverse problem for a class of Dirac operator, *Taiwanese Journal of Mathematics*, (2013) DOI: 10.11650/tjm.17.2013.2768
- [21] V. A. Marchenko, *Sturm-Liouville Operators and Applications*, AMS Chelsea Publishing, Providence, Rhode Island, 2011.
- [22] Ya. B. Levin, *Lectures on Entire Functions*, Providence, American Mathematical Society, 1996.
- [23] B. M. Levitan, I. S. Sargsjan. *Strum-Liouville and Dirac Operator*, Kluwer Academic, Publisher, Dordrecht, Boston, London, 1991.
- [24] G. Freiling, V. Yurko, *Inverse Sturm-Liouville Problems and Their Applications*, Nova Science Publisher, Inc., Huntington, New York, 2008.

^{1,2}SCIENCE AND LETTERS FACULTY, MATHEMATICS DEPARTMENT, MERSIN UNIVERSITY, 33343, TURKEY

E-mail address: ¹hanlar@mersin.edu.tr and ²ozge.akcy@gmail.com