

# CORRECTIONS TO "ON CONTINUED FRACTIONS OF GIVEN PERIOD"

VLADIMIR PLETSER

ABSTRACT. Corrections are brought to an article of Friesen on continued fractions of a given period.

Friesen has proven [1] that, for any  $k \in \mathbb{Z}^+$ , there are infinitely many squarefree integers  $N$ , where the continued fraction expansion of  $\sqrt{N}$  has period equal to  $k$ . This was demonstrated in a Corollary following a Theorem stating that

**Theorem 1.** *Let  $\lfloor \sqrt{N} \rfloor$  denote the greatest integer  $\leq \sqrt{N}$ . Then the equation*

$$\sqrt{N} = \left[ \lfloor \sqrt{N} \rfloor ; \overline{a_1, a_2, \dots, a_{k-2} = a_2, a_{k-1} = a_1, a_k = 2 \lfloor \sqrt{N} \rfloor} \right]$$

*has, for any symmetric set of positive integers  $\{a_1, \dots, a_{k-1}\}$ , infinitely many squarefree solutions  $N$  whenever either  $Q_{k-2}$  or  $(Q_{k-2}^2 - (-1)^k) / Q_{k-1}$  is even. If both quantities are odd, then there are no solutions  $N$  even if the squarefree condition is dropped.*

where  $Q_k$  are convergents of the of the continued fraction of  $\sqrt{N}$ . This theorem is demonstrated using three Lemmas.

Unfortunately, this paper contains two mistakes, that we want to correct here.

First, in the demonstration of the Lemma 1, (see [1], p. 12, line 19), the expression of  $N$  should be  $N = N(b) = \alpha b^2 + \beta b + \gamma$  (instead of  $\alpha b^2 + \beta b^2 + \gamma$ , which would not make sense). However this does not change the final result of this Lemma 1.

Second, Friesen proved the Corollary with a sufficient condition that for each  $k \in \mathbb{Z}^+$ , a symmetric set of positive integers  $\{a_1, \dots, a_{k-1}\}$  exists such that  $Q_{k-1}$  is odd. The case  $k = 1$  is direct as it yields  $N(b) = b^2 + 1$  giving an infinity of squarefree  $N$ .

However, for the case  $k > 1$ , the demonstration (see [1], p. 13, lines 19-25) is wrong as it contradicts the final statement made in the conclusion (see [1], p. 13, lines 35-36).

I have attempted to correct his demonstration herebelow.

"Assume  $k > 1$ . If  $k \equiv 0 \pmod{3}$  [instead of  $k \not\equiv 0 \pmod{3}$ ] then set  $a_1 = a_{k-1} = 2$  and  $a_i = 1$  for  $i = 2, \dots, k-2$ . If  $k \not\equiv 0 \pmod{3}$  [instead of  $k \equiv 0 \pmod{3}$ ] set  $a_i = 1$  for  $i = 1, \dots, k-1$ . In both cases we have the recursion formula for  $Q_n$  giving us copies of the Fibonacci sequence (...). In the first instance [i.e.  $k \equiv 0 \pmod{3}$ ] we have  $Q_n = F_{n+2}$  for  $n = 2, \dots, k-2$  [instead of  $i = 1, \dots, k-2$ ] and  $Q_{k-1} = 2Q_{k-2} + Q_{k-3} = 2F_k + F_{k-1}$  [instead of  $2F_k + F_{k-1} + F_{k+2}$ ]. If  $k \not\equiv 0 \pmod{3}$  we have  $Q_n = F_{n+1}$  for  $n = 1, \dots, k-1$ , hence  $Q_{k-1} = F_k$ . But, as  $F_k$  is even only

---

2000 *Mathematics Subject Classification.* Primary: 11A55, Secondary: 11J70.

*Key words and phrases.* Continued Fractions.

when  $k \equiv 0 \pmod{3}$ , we see that  $Q_{k-1}$  is odd in either situation. Therefore, either  $Q_{k-2}$  or  $(Q_{k-2}^2 - (-1)^k)/Q_{k-1}$  is even and we have satisfied the conditions of the Theorem, thus proving the Corollary."

With these corrections, these statements agree with those of the conclusion namely "By setting  $a_i = 1$  for  $i = 1, \dots, k-1$  if  $k \not\equiv 0 \pmod{3}$  (and  $a_1 = a_{k-1} = 2$ ,  $a_i = 1$  for  $i = 2, \dots, k-2$  if  $k \equiv 0 \pmod{3}$ ) it was shown that the conditions of the Theorem are met."

Despite these mistakes, the final result still stands.

#### REFERENCES

- [1] C. Friesen, On continued fractions of given period, Proc. Amer. Math. Soc. 103 (1988), no. 1, 9-14.

EUROPEAN SPACE RESEARCH AND TECHNOLOGY CENTRE, ESA-ESTEC P.O. Box 299,  
NL-2200 AG NOORDWIJK, THE NETHERLANDS

EUROPEAN SPACE RESEARCH AND TECHNOLOGY CENTRE, ESA-ESTEC P.O. Box 299,  
NL-2200 AG NOORDWIJK, THE NETHERLANDS  
*E-mail address:* `Vladimir.Pletser@esa.int`