A Picard-S Iterative Scheme for Approximating Fixed Point of Weak-Contraction Mappings

Faik GÜRSOY

Department of Mathematics, Yildiz Technical University, Davutpasa Campus, Esenler, 34220 Istanbul, Turkey e-mail: faikgursoy02@hotmail.com

The date of receipt and acceptance will be inserted by the editor

Abstract We study the convergence analysis of a Picard-S iteration method for a particular class of weak-contraction mappings. Furthermore, we prove a data dependence result for fixed point of the class of weak-contraction mappings with the help of the Picard-S iteration methods.

Key words Picard-S iterative scheme – Weak-Contraction mappings – Convergence – Rate of Convergence- Data Dependence

1 Introduction

Most of the problems that arise in different disciplines of science can be formulated by the equations in the form

$$Fx = 0, (1)$$

where F is some function. The equations given by (1) can easily be reformulated as the fixed point equations of type

$$Tx = x. (2)$$

where T is a self-map of an ambient space X and $x \in X$. These equations are often classified as linear or nonlinear, depending on whether the mappings used in the equation is linear with respect to the variables. Over the years, a considerable attention has been paid to solving such equations by using different techniques such as direct and iterative methods. In case of linear equations, both direct and iterative methods are used to obtain solutions of

Send offprint requests to: Offprints Assistant

Correspondence to: Address for offprint requests

the equations. But in case of nonlinear equations, due to various reasons, direct methods can be impractical or fail in solving equations, and thus iterative methods become a viable alternative. Nonlinear problems are of importance interest to mathematicians, physicists and engineers and many other scientists, simply because most systems are intrinsically nonlinear in nature. That is why, researchers in various disciplines of sciences are often faced with the solving such problems. It would be hard to fudge that the role of iterative approximation of fixed points have played in the recent progress of nonlinear science. Indeed, the phrase iterative approximation has been introduced to describe repetition-based researches into nonlinear problems that are inaccessible to analytic methods. For this reason, the iterative approximation of fixed points has become one of the major and basic tools in the theory of equations, and as a result, numerous iterative methods have been introduced or improved and studied for many years in detail from various points of aspects by a wide audience of researchers, see, [1-14, 16-42, 44, 45].

In this paper, we show that a Picard-S iteration method [14] can be used to approximate fixed point of weak-contraction mappings. Also, we show that this iteration method is equivalent and converges faster than CR iteration method [9] for the aforementioned class of mappings. Furthermore, by providing an example, it is shown that the Picard-S iteration method converges faster than CR iteration method and hence also faster than all Picard [31], Mann [25], Ishikawa [17], Noor [26], SP [30], S [2] and some other iteration methods in the existing literature when applied to weakcontraction mappings. Finally, a data dependence result is proven for fixed point of weak-contraction mappings with the help of the Picard-S iteration method.

Throughout this paper the set of all positive integers and zero is shown by N. Let B be a Banach space, D be a nonempty closed convex subset of B and T a self-map of D. An element x_* of D is called a fixed point of T if and only if $Tx_* = x_*$. The set of all fixed point of T denoted by F_T . Let $\{a_n^i\}_{n=0}^{\infty}, i \in \{0, 1, 2\}$ be real sequences in [0, 1] satisfying certain control condition(s).

Renowned Picard iteration method [31] is formulated as follow

$$\begin{cases} p_0 \in D, \\ p_{n+1} = Tp_n, \ n \in \mathbb{N}, \end{cases}$$
(3)

and generally used to approximate fixed points of contraction mappings satisfying: for all $x, y \in B$ there exists a $\delta \in (0, 1)$ such that

$$\|Tx - Ty\| \le \delta \|x - y\|.$$

$$\tag{4}$$

The following iteration methods are known as Noor [26] and SP [30] iterations, respectively:

$$\begin{cases} \omega_0 \in D, \\ \omega_{n+1} = (1 - a_n^0) \,\omega_n + a_n^0 T \varpi_n, \\ \varpi_n = (1 - a_n^1) \,\omega_n + a_n^1 T \rho_n, \\ \rho_n = (1 - a_n^2) \,\omega_n + a_n^2 T \omega_n, \, n \in \mathbb{N}, \end{cases}$$
(5)

$$\begin{cases} q_0 \in D, \\ q_{n+1} = (1 - a_n^0) r_n + a_n^0 T r_n, \\ r_n = (1 - a_n^1) s_n + a_n^1 T s_n, \\ s_n = (1 - a_n^2) q_n + a_n^2 T q_n, n \in \mathbb{N}, \end{cases}$$
(6)

Remark 1 (i) If $a_n^2 = 0$ for each $n \in \mathbb{N}$, then the Noor iteration method reduces to iterative method of Ishikawa [17].

(ii) If $a_n^2 = 0$ for each $n \in \mathbb{N}$, then the SP iteration method reduces to iterative method of Thianwan [42].

(iii) When $a_n^1 = a_n^2 = 0$ for each $n \in \mathbb{N}$, then both Noor and SP iteration methods reduce to an iteration method due to Mann [25].

Recently, Gürsoy and Karakaya [14] introduced a Picard-S iterative scheme as follows:

$$\begin{cases} x_0 \in D, \\ x_{n+1} = Ty_n, \\ y_n = (1 - a_n^1) Tx_n + a_n^1 Tz_n, \\ z_n = (1 - a_n^2) x_n + a_n^2 Tx_n, n \in \mathbb{N}, \end{cases}$$
(7)

The following definitions and lemmas will be needed in obtaining the main results of this article.

Definition 1 [5] Let $\{a_n\}_{n=0}^{\infty}$ and $\{b_n\}_{n=0}^{\infty}$ be two sequences of real numbers with limits a and b, respectively. Assume that there exists

$$\lim_{n \to \infty} \frac{|a_n - a|}{|b_n - b|} = l.$$
(8)

(i) If l = 0, the we say that $\{a_n\}_{n=0}^{\infty}$ converges faster to a than $\{b_n\}_{n=0}^{\infty}$ to b.

(ii) If $0 < l < \infty$, then we say that $\{a_n\}_{n=0}^{\infty}$ and $\{b_n\}_{n=0}^{\infty}$ have the same rate of convergence.

Definition 2 [5] Assume that for two fixed point iteration processes $\{u_n\}_{n=0}^{\infty}$ and $\{v_n\}_{n=0}^{\infty}$ both converging to the same fixed point p, the following error predictions

$$||u_n - p|| \le a_n \text{ for all } n \in \mathbb{N},\tag{9}$$

$$\|v_n - p\| \le b_n \text{ for all } n \in \mathbb{N},\tag{10}$$

are available where $\{a_n\}_{n=0}^{\infty}$ and $\{b_n\}_{n=0}^{\infty}$ are two sequences of positive numbers (converging to zero). If $\{a_n\}_{n=0}^{\infty}$ converges faster than $\{b_n\}_{n=0}^{\infty}$, then $\{u_n\}_{n=0}^{\infty}$ converges faster than $\{v_n\}_{n=0}^{\infty}$ to p.

Definition 3 [4] Let $(B, \|\cdot\|)$ be a Banach space. A map $T : B \to B$ is called weak-contraction if there exist a constant $\delta \in (0, 1)$ and some $L \ge 0$ such that

$$||Tx - Ty|| \le \delta ||x - y|| + L ||y - Ty||, \text{ for all } x, y \in B.$$
(11)

Definition 4 [4] Let $T, \tilde{T} : B \to B$ be two operators. We say that \tilde{T} is an approximate operator of T if for all $x \in B$ and for a fixed $\varepsilon > 0$ we have

$$\left\| Tx - \widetilde{T}x \right\| \le \varepsilon. \tag{12}$$

Lemma 1 [43]Let $\{\beta_n\}_{n=0}^{\infty}$ and $\{\rho_n\}_{n=0}^{\infty}$ be nonnegative real sequences satisfying the following inequality:

$$\beta_{n+1} \le (1 - \lambda_n) \,\beta_n + \rho_n, \tag{13}$$

where $\lambda_n \in (0,1)$, for all $n \ge n_0$, $\sum_{n=1}^{\infty} \lambda_n = \infty$, and $\frac{\rho_n}{\lambda_n} \to 0$ as $n \to \infty$. Then $\lim_{n\to\infty} \beta_n = 0$.

Lemma 2 [39] Let $\{\beta_n\}_{n=0}^{\infty}$ be a nonnegative sequence for which one assumes there exists $n_0 \in \mathbb{N}$, such that for all $n \geq n_0$ one has satisfied the inequality

$$\beta_{n+1} \le (1-\mu_n)\,\beta_n + \mu_n \gamma_n,\tag{14}$$

where $\mu_n \in (0,1)$, for all $n \in \mathbb{N}$, $\sum_{n=0}^{\infty} \mu_n = \infty$ and $\gamma_n \ge 0$, $\forall n \in \mathbb{N}$. Then the following inequality holds

$$0 \le \lim \sup_{n \to \infty} \beta_n \le \lim \sup_{n \to \infty} \gamma_n. \tag{15}$$

2 Main Results

Theorem 1 Let $T: D \to D$ be a weak-contraction map satisfying condition (11) with $F_T \neq \emptyset$ and $\{x_n\}_{n=0}^{\infty}$ an iterative sequence defined by (7) with real sequences $\{a_n^i\}_{n=0}^{\infty}$, $i \in \{1,2\}$ in [0,1] satisfying $\sum_{k=0}^{\infty} a_k^1 a_k^2 = \infty$. Then $\{x_n\}_{n=0}^{\infty}$ converges to a unique fixed point u^* of T.

Proof Uniqueness of u^* comes from condition (11). Using Picard-S iterative scheme (7) and condition (11), we obtain

$$||z_n - u^*|| \le (1 - a_n^2) ||x_n - u^*|| + a_n^2 ||Tx_n - Tu^*||$$

$$\le (1 - a_n^2) ||x_n - u^*|| + a_n^2 \delta ||x_n - u^*|| + a_n^2 L ||u^* - Tu^*||$$

$$= [1 - a_n^2 (1 - \delta)] ||x_n - u^*||, \qquad (16)$$

$$||y_n - u^*|| \le (1 - a_n^1) ||Tx_n - Tu^*|| + a_n^1 ||Tz_n - Tu^*|| \le (1 - a_n^1) \,\delta \,||x_n - u^*|| + a_n^1 \delta \,||z_n - u^*||,$$
(17)

Title Suppressed Due to Excessive Length

$$||x_{n+1} - u^*|| \le \delta ||y_n - u^*||.$$
(18)

Combining (16), (17) and (18)

$$|x_{n+1} - u^*|| \le \delta^2 \left[1 - a_n^1 a_n^2 \left(1 - \delta \right) \right] ||x_n - u^*||.$$
(19)

By induction

$$||x_{n+1} - u^*|| \le \delta^{2(n+1)} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right] ||x_0 - u^*||$$

$$\le \delta^{2(n+1)} ||x_0 - u^*||^{n+1} e^{-(1-\delta) \sum_{k=0}^n a_k^1 a_k^2}.$$
 (20)

Since $\sum_{k=0}^{\infty} a_k^1 a_k^2 = \infty$,

$$e^{-(1-\delta)\sum_{k=0}^{n}a_{k}^{1}a_{k}^{2}} \to 0 \text{ as } n \to \infty,$$

$$(21)$$

which implies $\lim_{n\to\infty} ||x_n - u^*||$.

Theorem 2 Let $T: D \to D$ with fixed point $u^* \in F_T \neq \emptyset$ be as in Theorem 1 and $\{q_n\}_{n=0}^{\infty}$, $\{x_n\}_{n=0}^{\infty}$ two iterative sequences defined by SP (6) and Picard-S (7) iteration methods with real sequences $\{a_n^i\}_{n=0}^{\infty}$, $i \in \{0, 1, 2\}$ in [0,1] satisfying $\sum_{k=0}^{n} a_k^1 a_k^2 = \infty$. Then the following are equivalent: (i) $\lim_{n\to\infty} ||x_n - u^*|| = 0$; (ii) $\lim_{n\to\infty} ||q_n - u^*|| = 0$.

Proof (i) \Rightarrow (ii): It follows from (6), (7), and condition (11) that

$$\|x_{n+1} - q_{n+1}\| = \|(1 - a_n^0) (Ty_n - r_n) + a_n^0 (Ty_n - Tr_n)\|$$

$$\leq (1 - a_n^0) \|Ty_n - r_n\| + a_n^0 \|Ty_n - Tr_n\|$$

$$\leq [1 - a_n^0 (1 - \delta)] \|y_n - r_n\| + [1 - a_n^0 (1 - L)] \|y_n - Ty_n\|,$$
(22)

$$||y_n - r_n|| = ||(1 - a_n^1) (Tx_n - s_n) + a_n^1 (Tz_n - Ts_n)||$$

$$\leq (1 - a_n^1) ||Tx_n - s_n|| + a_n^1 ||Tz_n - Ts_n||$$

$$\leq (1 - a_n^1) ||Tx_n - s_n|| + a_n^1 \delta ||z_n - s_n|| + a_n^1 L ||z_n - Tz_n||,$$
(23)

$$\|Tx_n - s_n\| = \|(1 - a_n^2)(Tx_n - q_n) + a_n^2(Tx_n - Tq_n)\|$$

$$\leq [1 - a_n^2(1 - \delta)] \|x_n - q_n\| + [1 - a_n^2(1 - L)] \|x_n - Tx_n\|,$$
(24)

$$||z_n - s_n|| \le (1 - a_n^2) ||x_n - q_n|| + a_n^2 ||Tx_n - Tq_n||$$

$$\le [1 - a_n^2 (1 - \delta)] ||x_n - q_n|| + a_n^2 L ||x_n - Tx_n||.$$
(25)

Combining (22), (23), (24), and (25)

$$\begin{aligned} \|x_{n+1} - q_{n+1}\| &\leq \left[1 - a_n^0 \left(1 - \delta\right)\right] \left[1 - a_n^1 \left(1 - \delta\right)\right] \left[1 - a_n^2 \left(1 - \delta\right)\right] \|x_n - q_n\| \\ &+ \left[1 - a_n^0 \left(1 - \delta\right)\right] \left\{ \left(1 - a_n^1\right) \left[1 - a_n^2 \left(1 - L\right)\right] + a_n^1 a_n^2 \delta L \right\} \|x_n - T x_n\| \\ &+ \left[1 - a_n^0 \left(1 - \delta\right)\right] a_n^1 L \|z_n - T z_n\| + \left[1 - a_n^0 \left(1 - L\right)\right] \|y_n - T y_n\|. \end{aligned}$$

$$(26)$$

It follows from the facts $\delta \in (0,1)$ and $a_n^i \in [0,1]$, $\forall n \in \mathbb{N}$, $i \in \{0,1,2\}$ that $\left[1 - a_n^0 (1-\delta)\right] \left[1 - a_n^1 (1-\delta)\right] \left[1 - a_n^2 (1-\delta)\right] < 1 - a_n^1 a_n^2 (1-\delta)$. (27)

Hence, inequality (26) becomes

$$\|x_{n+1} - q_{n+1}\| \leq \left[1 - a_n^1 a_n^2 (1 - \delta)\right] \|x_n - q_n\|$$

$$+ \left[1 - a_n^0 (1 - \delta)\right] \left\{ \left(1 - a_n^1\right) \left[1 - a_n^2 (1 - L)\right] + a_n^1 a_n^2 \delta L \right\} \|x_n - T x_n\|$$

$$+ \left[1 - a_n^0 (1 - \delta)\right] a_n^1 L \|z_n - T z_n\| + \left[1 - a_n^0 (1 - L)\right] \|y_n - T y_n\|.$$
(28)

Denote that

$$\begin{aligned} \beta_{n} &:= \|x_{n} - q_{n}\|, \end{aligned} (29) \\ \lambda_{n} &:= a_{n}^{1} a_{n}^{2} \left(1 - \delta\right) \in (0, 1), \\ \rho_{n} &:= \left[1 - a_{n}^{0} \left(1 - \delta\right)\right] \left\{ \left(1 - a_{n}^{1}\right) \left[1 - a_{n}^{2} \left(1 - L\right)\right] + a_{n}^{1} a_{n}^{2} \delta L \right\} \|x_{n} - T x_{n}\| \\ &+ \left[1 - a_{n}^{0} \left(1 - \delta\right)\right] a_{n}^{1} L \|z_{n} - T z_{n}\| + \left[1 - a_{n}^{0} \left(1 - L\right)\right] \|y_{n} - T y_{n}\|. \end{aligned}$$

Since $\lim_{n \to \infty} ||x_n - u^*|| = 0$ and $Tu^* = u^*$

$$\lim_{n \to \infty} \|x_n - Tx_n\| = \lim_{n \to \infty} \|y_n - Ty_n\| = \lim_{n \to \infty} \|z_n - Tz_n\| = 0, \quad (30)$$

which implies $\frac{\rho_n}{\lambda_n} \to 0$ as $n \to \infty$. Therefore, inequality (28) perform all assumptions in Lemma 1 and thus we obtain $\lim_{n\to\infty} ||x_n - q_n|| = 0$. Since

$$||q_n - u^*|| \le ||x_n - q_n|| + ||x_n - u^*|| \to 0 \text{ as } n \to \infty,$$
(31)

 $\lim_{n \to \infty} \|q_n - u^*\| = 0.$

 $(ii) \Rightarrow (i)$: It follows from (6), (7), and condition (11) that

$$||q_{n+1} - x_{n+1}|| = ||r_n - Ty_n + a_n^0 (Tr_n - r_n)||$$

$$\leq \delta ||r_n - y_n|| + (1 + a_n^0 + L) ||r_n - Tr_n||,$$
(32)

$$\|r_{n} - y_{n}\| \leq (1 - a_{n}^{1}) \|s_{n} - Tx_{n}\| + a_{n}^{1} \|Ts_{n} - Tz_{n}\|$$

$$\leq (1 - a_{n}^{1}) \|s_{n} - Tx_{n}\| + a_{n}^{1}\delta \|s_{n} - z_{n}\| + a_{n}^{1}L \|s_{n} - Ts_{n}\|,$$
(33)

$$||s_{n} - Tx_{n}|| \leq ||Ts_{n} - Tx_{n}|| + ||s_{n} - Ts_{n}||$$

$$\leq \delta ||s_{n} - x_{n}|| + (1 + L) ||s_{n} - Ts_{n}||$$

$$\leq \delta ||q_{n} - x_{n}|| + \delta a_{n}^{2} ||Tq_{n} - q_{n}|| + (1 + L) ||s_{n} - Ts_{n}|| ,$$
(34)

$$||s_n - z_n|| \le (1 - a_n^2) ||q_n - x_n|| + a_n^2 ||Tq_n - Tx_n||$$

$$\le [1 - a_n^2 (1 - \delta)] ||q_n - x_n|| + a_n^2 L ||q_n - Tq_n||.$$
(35)

Combining (32), (33), (34), and (35)

$$\|q_{n+1} - x_{n+1}\| \le \delta^2 \left[1 - a_n^1 a_n^2 (1 - \delta) \right] \|q_n - x_n\|$$

$$+ \delta^2 a_n^2 \left[1 - a_n^1 (1 - L) \right] \|q_n - Tq_n\|$$

$$+ \left(1 + a_n^0 + L \right) \|r_n - Tr_n\| + \delta \left(1 - a_n^1 + L \right) \|s_n - Ts_n\| .$$
(36)

Title Suppressed Due to Excessive Length

Since $\delta \in (0,1)$

$$\delta^{2} \left[1 - a_{n}^{1} a_{n}^{2} \left(1 - \delta \right) \right] < 1 - a_{n}^{1} a_{n}^{2} \left(1 - \delta \right).$$
(37)

Hence, inequality (36) becomes

$$\|q_{n+1} - x_{n+1}\| \leq \left[1 - a_n^1 a_n^2 (1 - \delta)\right] \|q_n - x_n\|$$

$$+ \delta^2 a_n^2 \left[1 - a_n^1 (1 - L)\right] \|q_n - Tq_n\|$$

$$+ \left(1 + a_n^0 + L\right) \|r_n - Tr_n\| + \delta \left(1 - a_n^1 + L\right) \|s_n - Ts_n\|.$$
(38)

Denote that

$$\begin{aligned} \beta_{n} &:= \|q_{n} - x_{n}\|, \\ \lambda_{n} &:= a_{n}^{1} a_{n}^{2} \left(1 - \delta\right) \in (0, 1), \\ \rho_{n} &:= \delta^{2} a_{n}^{2} \left[1 - a_{n}^{1} \left(1 - L\right)\right] \|q_{n} - Tq_{n}\| \\ &+ \left(1 + a_{n}^{0} + L\right) \|r_{n} - Tr_{n}\| + \delta \left(1 - a_{n}^{1} + L\right) \|s_{n} - Ts_{n}\|. \end{aligned}$$

$$(39)$$

Since $\lim_{n \to \infty} ||q_n - u^*|| = 0$ and $Tu^* = u^*$

$$\lim_{n \to \infty} \|q_n - Tq_n\| = \lim_{n \to \infty} \|r_n - Tr_n\| = \lim_{n \to \infty} \|s_n - Ts_n\| = 0, \quad (40)$$

which implies $\frac{\rho_n}{\lambda_n} \to 0$ as $n \to \infty$. Therefore, inequality (38) perform all assumptions in Lemma 1 and thus we obtain $\lim_{n\to\infty} ||q_n - x_n|| = 0$. Since

$$||x_n - u^*|| \le ||q_n - x_n|| + ||q_n - u^*|| \to 0 \text{ as } n \to \infty,$$
(41)

 $\lim_{n \to \infty} \|x_n - u^*\| = 0.$

Taking R. Chugh et al.'s result ([9], Corollary 3.2) into account, Theorem 2 leads to the following corollary under weaker assumption:

Corollary 1 Let $T: D \to D$ with fixed point $u^* \in F_T \neq \emptyset$ be as in Theorem 1. Then the followings are equivalent:

- 1) The Picard iteration method (3) converges to u^* ,
- 2) The Mann iteration method [25] converges to u^* ,
- 3) The Ishikawa iteration method [17] converges to u^* ,
- 4) The Noor iteration method (5) converges to u^* ,
- 5) S-iteration method [2] converges to u^* ,

6) The SP-iteration method (6) converges to u^* ,

- 7) CR-iteration method [9] converges to u^* ,
- 8) The Picard-S iteration method (7) converges to u^* .

Theorem 3 Let $T: D \to D$ with fixed point $u^* \in F_T \neq \emptyset$ be as in Theorem 1. Suppose that $\{\omega_n\}_{n=0}^{\infty}$, $\{q_n\}_{n=0}^{\infty}$ and $\{x_n\}_{n=0}^{\infty}$ are iterative sequences, respectively, defined by Noor (5), SP (6) and Picard-S (7) iterative schemes with real sequences $\{a_n^i\}_{n=0}^{\infty} \subset [0,1], i \in \{0,1,2\}$ satisfying

- $(i) \ 0 \le a_n^i < \frac{1}{1+\delta},$
- (*ii*) $\lim_{n \to \infty} a_n^i = 0.$

Then the iterative sequence defined by (7) converges faster than the iterative sequences defined by (5) and (6) to a unique fixed point of T, provided that the initial point is the same for all iterations.

Proof From inequality (20), we have

$$\|x_{n+1} - u^*\| \le \delta^{2(n+1)} \|x_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right]$$
(42)

Using (6) we obtain

$$\begin{aligned} \|q_{n+1} - u^*\| &= \left\| \left(1 - a_n^0\right) r_n + a_n^0 T r_n - u^* \right\| \end{aligned}$$
(43)

$$\geq \left(1 - a_n^0\right) \|r_n - u^*\| - a_n^0 \|T r_n - T u^*\| \end{aligned}$$
(43)

$$\geq \left[1 - a_n^0 (1 + \delta)\right] \|r_n - u^*\| \end{aligned}$$

$$\geq \left[1 - a_n^0 (1 + \delta)\right] \left\{ \left(1 - a_n^1\right) \|s_n - u^*\| - a_n^1 \delta \|s_n - u^*\| \right\} \end{aligned}$$

$$= \left[1 - a_n^0 (1 + \delta)\right] \left[1 - a_n^1 (1 + \delta)\right] \|s_n - u^*\| \end{aligned}$$

$$\geq \left[1 - a_n^0 (1 + \delta)\right] \left[1 - a_n^1 (1 + \delta)\right] \left\{ \left(1 - a_n^2\right) \|q_n - u^*\| - a_n^2 \delta \|q_n - u^*\| \right\} \end{aligned}$$

$$= \left[1 - a_n^0 (1 + \delta)\right] \left[1 - a_n^1 (1 + \delta)\right] \left[1 - a_n^2 (1 + \delta)\right] \|q_n - u^*\| \end{aligned}$$

$$\geq \cdots$$

$$\geq \|q_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^0 (1 + \delta)\right] \left[1 - a_k^1 (1 + \delta)\right] \left[1 - a_k^2 (1 + \delta)\right] .$$

Using now (42) and (43)

$$\frac{\|x_{n+1} - u^*\|}{\|q_{n+1} - u^*\|} \le \frac{\delta^{2(n+1)} \|x_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right]}{\|q_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^0 \left(1 + \delta\right)\right] \left[1 - a_k^1 \left(1 + \delta\right)\right] \left[1 - a_k^2 \left(1 + \delta\right)\right]}.$$
(44)

Define

$$\theta_n = \frac{\delta^{2(n+1)} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right]}{\prod_{k=0}^n \left[1 - a_k^0 \left(1 + \delta\right)\right] \left[1 - a_k^1 \left(1 + \delta\right)\right] \left[1 - a_k^2 \left(1 + \delta\right)\right]}.$$
 (45)

By the assumtion

$$\lim_{n \to \infty} \frac{\theta_{n+1}}{\theta_n}$$
(46)
$$= \lim_{n \to \infty} \frac{\frac{\delta^{2(n+2)} \prod_{k=0}^{n+1} [1-a_k^1 a_k^2 (1-\delta)]}{\prod_{k=0}^{n+1} [1-a_k^0 (1+\delta)] [1-a_k^1 (1+\delta)] [1-a_k^2 (1+\delta)]}}{\frac{\delta^{2(n+1)} \prod_{k=0}^{n} [1-a_k^1 a_k^2 (1-\delta)]}{\prod_{k=0}^{n} [1-a_k^0 (1+\delta)] [1-a_k^1 (1+\delta)] [1-a_k^2 (1+\delta)]}}$$
$$= \lim_{n \to \infty} \frac{\delta^2 [1-a_{n+1}^1 a_{n+1}^2 (1-\delta)]}{[1-a_{n+1}^1 (1+\delta)] [1-a_{n+1}^2 (1+\delta)]} [1-a_{n+1}^2 (1+\delta)]}$$
$$= \delta^2 < 1.$$

Title Suppressed Due to Excessive Length

It thus follows from ratio test that $\sum_{n=0}^{\infty} \theta_n < \infty$. Hence, we have $\lim_{n\to\infty} \theta_n = 0$ which implies that the iterative sequence defined by (7) converges faster than the iterative sequence defined by SP iteration method (6).

Using Noor iteration method (5), we get

$$\begin{aligned} \|\omega_{n+1} - u^*\| &= \left\| \left(1 - a_n^0\right) \omega_n + a_n^0 T \varpi_n - u^* \right\| \end{aligned}$$
(47)

$$\geq (1 - a_n^0) \|\omega_n - u^*\| - a_n^0 \|T \varpi_n - T u^*\|$$

$$\geq (1 - a_n^0) \|\omega_n - u^*\| - a_n^0 \delta \|\varpi_n - u^*\|$$

$$\geq \left[1 - a_n^0 - a_n^0 \delta \left(1 - a_n^1\right)\right] \|\omega_n - u^*\| - a_n^0 a_n^1 \delta^2 \|\rho_n - u^*\|$$

$$\geq \left\{1 - a_n^0 - a_n^0 \delta \left(1 - a_n^1\right) - a_n^0 a_n^1 \delta^2 \left[1 - a_n^2 \left(1 - \delta\right)\right]\right\} \|\omega_n - u^*\|$$

$$\geq \left\{1 - a_n^0 - a_n^0 \delta \left[1 - a_n^1 \left(1 - \delta\right)\right]\right\} \|\omega_n - u^*\|$$

$$\geq \left[1 - a_n^0 \left(1 + \delta\right)\right] \|\omega_n - u^*\|$$

$$\geq \cdots$$

$$\geq \left\|\omega_0 - u^*\right\|^{n+1} \prod_{k=0}^n \left[1 - a_k^0 \left(1 + \delta\right)\right].$$

It follows by (42) and (47) that

$$\frac{\|x_{n+1} - u^*\|}{\|\omega_{n+1} - x_*\|} \le \frac{\delta^{2(n+1)} \|x_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right]}{\|\omega_0 - u^*\|^{n+1} \prod_{k=0}^n \left[1 - a_k^0 \left(1 + \delta\right)\right]}.$$
 (48)

Define

$$\theta_n = \frac{\delta^{2(n+1)} \prod_{k=0}^n \left[1 - a_k^1 a_k^2 \left(1 - \delta\right)\right]}{\prod_{k=0}^n \left[1 - a_k^0 \left(1 + \delta\right)\right]}.$$
(49)

By the assumption

$$\lim_{n \to \infty} \frac{\theta_{n+1}}{\theta_n} = \lim_{n \to \infty} \frac{\delta^2 \left[1 - a_{n+1}^1 a_{n+1}^2 \left(1 - \delta \right) \right]}{\left[1 - a_{n+1}^0 \left(1 + \delta \right) \right]}$$
(50)
= $\delta^2 < 1$.

It thus follows from ratio test that $\sum_{n=0}^{\infty} \theta_n < \infty$. Hence, we have $\lim_{n\to\infty} \theta_n = 0$ which implies that the iterative sequence defined by (7) converges faster than the iterative sequence defined by Noor iteration method (5).

By use of the following example due to [45], it was shown in ([9], Example 4.1) that CR iterative method [9] is faster than all of Picard (3), S [2], Noor (5) and SP (6) iterative methods for a particular class of operators which is included in the class of weak-contraction mappings satisfying (11). In the following, for the sake of consistent comparison, we will use the same example as that of ([9], Example 4.1) in order to compare the rates of convergence between the Picard-S iterative scheme (7) and the CR iteration method [9] for weak-contraction mappings. In the following example, for convenience, we use the notations (PS_n) and (CR_n) for the iterative sequences associated to Picard-S (7) and CR [9] iteration methods, respectively.

Example 1 [45] Define a mapping $T : [0,1] \to [0,1]$ as $Tx = \frac{x}{2}$. Let $a_n^0 = a_n^1 = a_n^2 = 0$, for n = 1, 2, ..., 24 and $a_n^0 = a_n^1 = a_n^2 = \frac{4}{\sqrt{n}}$, for all $n \ge 25$.

It can be seen easily that the mapping T satisfies condition (11) with the unique fixed point $0 \in F_T$. Furthermore, it is easy to see that Example 1 satisfies all the conditions of Theorem 1. Indeed, let $x_0 \neq 0$ be an initial point for the iterative sequences (PS_n) and (CR_n) . Utilizing Picard-S (7) and CR [9] iteration methods we obtain

$$PS_{n} = \frac{1}{2} \left(\frac{1}{2} - \frac{4}{n} \right) x_{n}$$

= ...
= $\prod_{k=25}^{n} \left(\frac{1}{4} - \frac{2}{k} \right) x_{0},$ (51)

$$CR_{n} = \left(\frac{1}{2} - \frac{1}{\sqrt{n}} - \frac{4}{n} + \frac{8}{n\sqrt{n}}\right)x_{n}$$

= ...
= $\prod_{k=25}^{n} \left(\frac{1}{2} - \frac{1}{\sqrt{k}} - \frac{4}{k} + \frac{8}{k\sqrt{k}}\right)x_{0}.$ (52)

It follows from (51) and (52) that

$$\frac{|PS_n - 0|}{|CR_n - 0|} = \frac{\prod_{k=25}^n \left(\frac{1}{4} - \frac{2}{k}\right) x_0}{\prod_{k=25}^n \left(\frac{1}{2} - \frac{1}{\sqrt{k}} - \frac{4}{k} + \frac{8}{k\sqrt{k}}\right) x_0}$$
$$= \prod_{k=25}^n \frac{\frac{1}{2} - \frac{1}{\sqrt{k}} - \frac{4}{k} + \frac{8}{k\sqrt{k}}}{\frac{1}{2} - \frac{1}{\sqrt{k}} - \frac{4}{k} + \frac{8}{k\sqrt{k}}}$$
$$= \prod_{k=25}^n \frac{(k - 8)\sqrt{k}}{2\left(k\sqrt{k} - 2k - 8\sqrt{k} + 16\right)}$$
$$= \prod_{k=25}^n \frac{(k - 8)\sqrt{k}}{2\left(\sqrt{k} - 2\right)(k - 8)}$$
$$= \prod_{k=25}^n \frac{\sqrt{k}}{2\left(\sqrt{k} - 2\right)}.$$
(53)

For all $k \geq 25$, we have

$$\frac{(k-2)\left(\sqrt{k}-4\right)}{4} > 1$$

$$\Rightarrow (k-2)\left(\sqrt{k}-4\right) > 4$$

$$\Rightarrow k\left(\sqrt{k}-4\right) > 2\left(\sqrt{k}-2\right)$$

$$\Rightarrow \frac{\sqrt{k}-4}{2\left(\sqrt{k}-2\right)} > \frac{1}{k}$$

$$\Rightarrow \frac{\sqrt{k}}{2\left(\sqrt{k}-2\right)} < 1 - \frac{1}{k},$$
(54)

which yields

$$\frac{|PS_n - 0|}{|CR_n - 0|} = \prod_{k=25}^n \frac{\sqrt{k}}{2\left(\sqrt{k} - 2\right)} < \prod_{k=25}^n \left(1 - \frac{1}{k}\right) = \frac{24}{n}.$$
 (55)

Therefore, we have

$$\lim_{n \to \infty} \frac{|PS_n - 0|}{|CR_n - 0|} = 0,$$
(56)

which implies that the Picard-S iterative scheme (7) is faster than the CR iteration method [9].

Having regard to R. Chugh et al.'s result ([9], Example 4.1), L.B. Ciric et al.'s results [10] and Example 1 above, we conclude that Picard-S iteration method is faster than all Picard (3), Mann [25], Ishikawa [17], S [2], Noor (5) and SP (6) iterative methods.

We are now able to establish the following data dependence result.

Theorem 4 Let T with fixed point $u^* \in F_T \neq \emptyset$ be as in Theorem 1 and \widetilde{T} an approximate operator of T. Let $\{x_n\}_{n=0}^{\infty}$ be an iterative sequence generated by (7) for T and define an iterative sequence $\{\widetilde{x}_n\}_{n=0}^{\infty}$ as follows

$$\begin{cases} \widetilde{x}_0 \in D, \\ \widetilde{x}_{n+1} = \widetilde{T}\widetilde{y}_n, \\ \widetilde{y}_n = (1 - a_n^1) \widetilde{T}\widetilde{x}_n + a_n^1 \widetilde{T}\widetilde{z}_n, \\ \widetilde{z}_n = (1 - a_n^2) \widetilde{x}_n + a_n^2 \widetilde{T}\widetilde{x}_n, n \in \mathbb{N}, \end{cases}$$
(57)

where $\{a_n^i\}_{n=0}^{\infty}$, $i \in \{1, 2\}$ be real sequences in [0, 1] satisfying $(i) \frac{1}{2} \leq a_n^1 a_n^2$ for all $n \in \mathbb{N}$, and $(ii) \sum_{n=0}^{\infty} a_n^1 a_n^2 = \infty$. If $\widetilde{T}\widetilde{u}^* = \widetilde{u}^*$ such that $\widetilde{x}_n \to \widetilde{u}^*$ as $n \to \infty$, then we have

$$\|u^* - \widetilde{u}^*\| \le \frac{5\varepsilon}{1 - \delta},\tag{58}$$

where $\varepsilon > 0$ is a fixed number.

Proof It follows from (7), (11), (12), and (57) that

$$\begin{aligned} \|z_n - \widetilde{z}_n\| &\leq \left(1 - a_n^2\right) \|x_n - \widetilde{x}_n\| + a_n^2 \left\| Tx_n - \widetilde{T}\widetilde{x}_n \right\| \\ &\leq \left[1 - a_n^2 + a_n^2 \delta\right] \|x_n - \widetilde{x}_n\| + a_n^2 L \|x_n - Tx_n\| + a_n^2 \varepsilon, \tag{59} \end{aligned}$$

$$\|y_{n} - \widetilde{y}_{n}\| \leq (1 - a_{n}^{1}) \,\delta \,\|x_{n} - \widetilde{x}_{n}\| + a_{n}^{1} \delta \,\|z_{n} - \widetilde{z}_{n}\| + (1 - a_{n}^{1}) \,L \,\|x_{n} - Tx_{n}\| + a_{n}^{1} L \,\|z_{n} - Tz_{n}\| + (1 - a_{n}^{1}) \,\varepsilon + a_{n}^{1} \varepsilon,$$
(60)

$$\|x_{n+1} - \widetilde{x}_{n+1}\| \le \delta \|y_n - \widetilde{y}_n\| + L \|y_n - Ty_n\| + \varepsilon.$$
(61)

From the relations (59), (60), and (61)

$$\begin{aligned} \|x_{n+1} - \widetilde{x}_{n+1}\| &\leq \delta^2 \left[1 - a_n^1 a_n^2 \left(1 - \delta \right) \right] \|x_n - \widetilde{x}_n\| \\ &+ \left\{ a_n^1 a_n^2 \delta^2 L + \left(1 - a_n^1 \right) \delta L \right\} \|x_n - T x_n\| \\ &+ L \|y_n - T y_n\| + a_n^1 \delta L \|z_n - T z_n\| \\ &+ a_n^1 a_n^2 \delta^2 \varepsilon + \left(1 - a_n^1 \right) \delta \varepsilon + a_n^1 \delta \varepsilon + \varepsilon. \end{aligned}$$
(62)

Since $a_n^1, \, a_n^2 \in [0,1]$ and $\frac{1}{2} \leq a_n^1 a_n^2$ for all $n \in \mathbb{N}$

$$1 - a_n^1 a_n^2 \le a_n^1 a_n^2, (63)$$

$$1 - a_n^1 \le 1 - a_n^1 a_n^2 \le a_n^1 a_n^2, \tag{64}$$

$$1 \le 2a_n^1 a_n^2. \tag{65}$$

Use of the facts δ , $\delta^2 \in (0, 1)$, (63), (64), and (65) in (62) yields

$$||x_{n+1} - \widetilde{x}_{n+1}|| \leq \left[1 - a_n^1 a_n^2 (1 - \delta)\right] ||x_n - \widetilde{x}_n|| + a_n^1 a_n^2 (1 - \delta) \left\{ \frac{L\delta (1 + \delta) ||x_n - Tx_n||}{1 - \delta} + \frac{2L ||y_n - Ty_n|| + 2\delta L ||z_n - Tz_n|| + 5\varepsilon}{1 - \delta} \right\}.$$
 (66)

Define

$$\begin{aligned} \beta_n &:= \|x_n - \widetilde{x}_n\|, \\ \mu_n &:= a_n^1 a_n^2 \left(1 - \delta\right) \in (0, 1), \\ \gamma_n &:= \frac{L\delta \left(1 + \delta\right) \|x_n - Tx_n\| + 2L \|y_n - Ty_n\| + 2\delta L \|z_n - Tz_n\| + 5\varepsilon}{1 - \delta} \ge 0. \end{aligned}$$
(67)

Hence, the inequality (66) perform all assumptions in Lemma 2 and thus an application of Lemma 2 to (66) yields

$$0 \leq \lim \sup_{n \to \infty} \|x_n - \widetilde{x}_n\|$$

$$\leq \lim \sup_{n \to \infty} \frac{L\delta \left(1 + \delta\right) \|x_n - Tx_n\| + 2L \|y_n - Ty_n\| + 2\delta L \|z_n - Tz_n\| + 5\varepsilon}{1 - \delta}.$$
(68)

We know from Theorem 1 that $\lim_{n\to\infty} x_n = u^*$ and since $Tu^* = u^*$

$$\lim_{n \to \infty} \|x_n - Tx_n\| = \lim_{n \to \infty} \|y_n - Ty_n\| = \lim_{n \to \infty} \|z_n - Tz_n\| = 0.$$
(69)

Therefore the inequality (68) becomes

$$\|u^* - \widetilde{u}^*\| \le \frac{5\varepsilon}{1 - \delta}.\tag{70}$$

References

- 1. Abbas, M. & Nazir, T. 2013. A new faster iteration process applied to constrained minimization and feasibility problems. Matematicki Vesnik, Article.
- Agarwal, R., O Regan, D. & Sahu, D. 2007. Iterative construction of fixed points of nearly asymptotically nonexpansive mappings. Journal of Nonlinear and Convex Analysis 8: 61-79.
- Babu, G. V. R. & Prasad, K. N. V. V. 2006. Mann iteration converges faster than Ishikawa iteration for the class of Zamfirescu operators. Fixed Point Theory and Applications 2006(1).
- 4. Berinde, V. 2007. Iterative Approximation of Fixed Points. Springer, Berlin.
- 5. Berinde, V. 2004. Picard iteration converges faster than the Mann iteration in the class of quasi-contractive operators. Fixed Point Theory Appl. 2: 97-105.
- 6. Berinde, V. 2004. Comparing Krasnoselskij and Mann iterations for Lipschitzian generalized pseudocontractive operators. Proceed. of Int. Conf. On Fixed Point Theory, Univ. of Valencia. Yokohama Publishers.
- Berinde, V. & Berinde, M. 2005. The fastest Krasnoselskij iteration for approximating fixed points of strictly pseudo-contractive mappings. Carpathian J. Math 21(1-2): 13-20.
- Chang, S. S., Cho, Y. J. & Kim, J. K. 2003. The equivalence between the convergence of modified Picard, modified Mann, and modified Ishikawa iterations. Mathematical and Computer Modelling 37(9): 985-991.
- Chugh, R., Kumar, V. & Kumar, S. 2012. Strong Convergence of a new three step iterative scheme in Banach spaces. American Journal of Computational Mathematics 2: 345-357.
- Ciric, L. B., Lee, B. S., & Rafiq, A. 2010. Faster Noor iterations. Indian Journal of Mathematics, 52(3), 429-436.
- 11. Gürsoy, F. & Karakaya, V. 2014. Some Convergence and Stability Results for Two New Kirk Type Hybrid Fixed Point Iterative Algorithms. Journal of Function Spaces doi:10.1155/2014/684191.
- Gürsoy, F., Karakaya, V. & Rhoades, B.E. 2014. Some Convergence and Stability Results for the Kirk Multistep and Kirk-SP Fixed Point Iterative Algorithms. Abstract and Applied Analysis doi:10.1155/2014/806537.
- Gürsoy, F., Karakaya, V. & Rhoades, B.E. 2013. Data dependence results of new multistep and S-iterative schemes for contractive-like operators. Fixed Point Theory and Applications doi:10.1186/1687-1812-2013-76.
- Gürsoy, F., & Karakaya, V. 2014. A Picard-S hybrid type iteration method for solving a differential equation with retarded argument. arXiv preprint arXiv:1403.2546.
- 15. Hämmerlin, G. & Hoffmann, K. H. 1991. Numerical mathematics. Springer.

- Hussain, N., Rafiq, A., Damjanović, B. & Lazović, R. 2011. On rate of convergence of various iterative schemes. Fixed Point Theory and Applications 2011(1): 1-6.
- 17. Ishikawa, S. 1974. Fixed points by a new iteration method. Proceedings of the American Mathematical Society 44: 147-150.
- Karahan, I. & Ozdemir, M. 2013. A general iterative method for approximation of fixed points and their applications. Advances in Fixed Point Theory 3(3).
- Karakaya, V., Gürsoy, F. & Ertürk, M. 2014. Comparison of The Speed of Convergence Among Various Iterative Schemes. arXiv preprint arXiv:1402.6080.
- Karakaya, V., Gürsoy, F., Doğan, K. & Ertürk, M. 2103. Data Dependence Results for Multistep and CR Iterative Schemes in the Class of Contractive-Like Operators. Abstract and Applied Analysis doi:10.1155/2013/381980.
- Karakaya, V., Doğan, K., Gürsoy, F. & Ertürk, M. 2013. Fixed Point of a New Three-Step Iteration Algorithm under Contractive-Like Operators over Normed Spaces. Abstract and Applied Analysis 2013.
- 22. Khan, S. H. 2013. A Picard-Mann hybrid iterative process. Fixed Point Theory and Applications 2013(1): 1-10.
- Kirk, W. 1971. On successive approximations for nonexpansive mappings in Banach spaces. Glasgow Math. J. 12:6-9.
- Krasnosel'skii, M.A. 1955. Two remarks on the method of successive approximations. Uspekhi Matematicheskikh Nauk 10: 123-127.
- Mann, W.R. 1953. Mean value methods in iteration. Proceedings of the American Mathematical Society 4: 506-510.
- Noor, M.A. 2000. New approximation schemes for general variational inequalities. Journal of Mathematical Analysis and Applications 251: 217-229.
- 27. Olatinwo, M. 2009. Some stability results for two hybrid fixed point iterative algorithms in normed linear space. Matematique Vesnik 61: 247-256.
- Olatinwo, M.O. 2008. Some stabilty results for two hybrid fixed point iterative algorithms of Kirk-Ishikawa and Kirk-Mann type. Journal of AdvancedMathematical Studies 1(1-2): 87-96.
- Phuengrattana, W. & Suantai, S. 2012. Comparison of the Rate of Convergence of Various Iterative Methods for the Class of Weak Contractions in Banach Spaces. Thai Journal of Mathematics 11(1): 217-226.
- Phuengrattana, W. & Suantai, S. 2011. On the rate of convergence of Mann, Ishikawa, Noor and SP-iterations for continuous functions on an arbitrary interval. Journal of Computational and Applied Mathematics 235: 3006-3014.
- 31. Picard, E., (1890). "Mémoire sur la théorie des équations aux dérivées partielles et la méthode des approximations successives", Journal de Mathématiques pures et appliquées, 6:145-210.
- 32. Popescu, O. 2007. Picard iteration converges faster than Mann iteration for a class of quasiontractive operators. Math. Commun. 12: 195-202.
- 33. Rhoades, B. E. & Şoltuz, S. M. 2004. The equivalence between the convergences of Ishikawa and Mann iterations for an asymptotically nonexpansive in the intermediate sense and strongly successively pseudocontractive maps. Journal of mathematical analysis and applications 289(1): 266-278.
- Rhoades, B. & Soltuz, S.M. 2004. The equivalence between Mann–Ishikawa iterations and multistep iteration. Nonlinear Analysis: Theory, Methods & Applications 58: 219-228.

- Rhoades, B. E. 1976. Comments on two fixed point iteration methods. J. Math. Anal. Appl. 56(2): 741-750.
- Sahu, D. R. 2011. Applications of the S-iteration process to constrained minimization problems and split feasibility problems. Fixed Point Theory 12(1): 187-204.
- 37. Sahu, D. R. & Petrusel, A. 2011. Strong convergence of iterative methods by strictly pseudocontractive mappings in Banach spaces. NonlinearAnalysis: Theory, Methods & Applications 74(17): 6012-6023.
- Schaefer, H. 1957. Über die Methode sukzessiver Approximationen. Jahresbericht der Deutschen Mathematiker-Vereinigung 59: 131-140.
- Soltuz, S.M. & Grosan, T. 2008. Data dependence for Ishikawa iteration when dealing with contractive like operators. Fixed Point Theory and Applications 2008: 242916(1-7).
- 40. Soltuz, S. M. & Otrocol, D. 2007. Classical results via Mann-Ishikawa iteration. Revue d'Analyse Numerique de l'Approximation 36(2).
- Sun, H. & Xue, Z. 2013. Reconsiderations on the Equivalence of Convergence between Mann and Ishikawa Iterations for Asymptotically Pseudocontractive Mappings. Journal of Applied Mathematics 2013.
- 42. Thianwan, S. 2009. Common fixed points of new iterations for two asymptotically nonexpansive nonself-mappings in a Banach space. Journal of Computational and Applied Mathematics 224: 688-695.
- Weng, X. 1991. Fixed point iteration for local strictly pseudocontractive mapping. Proc. Amer. Math. Soc. 113: 727-731.
- 44. Xue, Z. 2008. The comparison of the convergence speed between Picard, Mann, Krasnoselskij and Ishikawa iterations in Banach spaces. Fixed Point Theory Appl. 2008.
- 45. Yuan, Q. & Rhoades, B.E. 2008. Comments on the rate of convergence between Mann and Ishikawa iterations applied to Zamfirescu operators. Fixed Point Theory and Applications 2008.