Yet another algorithm to compute the nonlinearity of a Boolean function

E. Bellini

Department of Mathematics, University of Trento, Italy eemanuele.bellini@gmail.com

Abstract

We associate to each Boolean function a polynomial whose evaluations represents the distances from all possible Boolean affine functions. Both determining the coefficients of this polynomial from the truth table of the Boolean function and computing its evaluation vector requires a worst-case complexity of $O(n2^n)$ integer operations. This way, with a different approach, we reach the same complexity of established algorithms, such as those based on the fast Walsh transform.

Keywords: Boolean function, nonlinearity, fast Walsh transform

1 Introduction

Any function from $(\mathbb{F}_2)^n$ to \mathbb{F}_2 is called a Boolean function. Boolean functions are important in symmetric cryptography, since they are used in the confusion layer of ciphers. An affine Boolean function does not provide an effective confusion. To overcome this, we need functions which are as far as possible from being an affine function. The effectiveness of these functions is measured by several parameters, one of these is called "nonlinearity".

In this paper, we provide a method to compute the nonlinearity of a Boolean function, estimating its complexity compared to that of the classical methods which uses fast Walsh and fast Mobius transform.

In Section 2 we recall the basic notions and theorems we need. In Section 3 we associate to each Boolean function in n variables a polynomial whose evaluations represent the distance from all possible affine functions, yielding an algorithm to compute the nonlinearity. We also provide a theorem to express the coefficients of this polynomials. Finally, in Section 4 we analyze the complexity of the proposed method, both experimentally and theoretically. In particular, we arrive at a worst-case complexity of $O(n2^n)$ operations over the integers, that is, sums and doublings. This way, with a different approach, we reach the same complexity of established algorithms, such as those based on the fast Walsh transform.

2 Preliminaries and Notation on Boolean functions

In this chapter we summarize some definitions and known results from [Car10] and [MS77], concerning B.f. and the classical techniques to determine their nonlinearity.

Let \mathbb{F} denote the field \mathbb{F}_2 . The set \mathbb{F}^n is the set of all binary vectors of length n, viewed as an \mathbb{F} -vector space.

Let $v \in \mathbb{F}^n$. The Hamming weight w(v) of the vector v is the number of its nonzero coordinates. For any two vectors $v_1, v_2 \in \mathbb{F}^n$, the Hamming distance between v_1 and v_2 , denoted by $d(v_1, v_2)$, is the number of coordinates in which the two vectors differ.

A Boolean function (B.f.) is any function $f : \mathbb{F}^n \to \mathbb{F}$. The set of all B.f.'s from \mathbb{F}^n to \mathbb{F} will be denoted by \mathcal{B}_n .

We assume implicitly to have ordered \mathbb{F}^n , so that $\mathbb{F}^n = \{\mathsf{p}_1, \ldots, \mathsf{p}_{2^n}\}.$

A B.f. f can be specified by a *truth table*, which gives the evaluation of f at all \mathbf{p}_i 's. We consider the evaluation map from \mathcal{B}_n to \mathbb{F}^{2^n} , associating to each B.f. f the vector $\underline{f} = (f(\mathbf{p}_1) \dots, f(\mathbf{p}_{2^n}))$, which is called the *evaluation vector* of f. Once the order on \mathbb{F}^n is chosen, i.e. the \mathbf{p}_i 's are fixed, it is clear that the evaluation vector of f uniquely identifies f.

A B.f. $f \in \mathcal{B}_n$ can be expressed in a unique way as a polynomial in $\mathbb{F}[X] = \mathbb{F}[x_1, \ldots, x_n]$, as $f = \sum_{v \in \mathbb{F}^n} b_v X^v$, where $X^v = x^{v_1} \cdots x^{v_n}$. This representation is called the *Algebraic Normal* Form (ANF).

Let $\mathcal{A}_n = \{ \alpha \in \mathcal{B}_n \mid \alpha(X) = a_0 + \sum_{i=1}^n a_i x_i, (a_0, \dots, a_n) \in \mathbb{F}^{n+1} \}$ denote the set of all affine functions.

In [CG99] a useful representation of B.f. 's is introduced for characterizing several cryptographic criteria.

B.f. 's can be represented as elements of $\mathbb{K}[X]/\langle X^2 - X \rangle$, where $\langle X^2 - X \rangle$ is the ideal generated by the polynomials $x_1^2 - x_1, \ldots, x_n^2 - x_n$, and \mathbb{K} is \mathbb{Z} , \mathbb{Q} , \mathbb{R} , or \mathbb{C} . Let f be a function on \mathbb{F}^n taking values in a field \mathbb{K} . We call the *numerical normal form* (NNF) of f the following expression of f as a polynomial: $f(x_1, \ldots, x_n) = \sum_{u \in \mathbb{F}^n} \lambda_u(\prod_{i=1}^n x_i^{u_i}) = \sum_{u \in \mathbb{F}^n} \lambda_u X^u$, with $\lambda_u \in \mathbb{K}$ and $u = (u_1, \ldots, u_n)$.

It can be proved that any B.f. f admits a unique NNF, consideiring values in $\mathbb K.$

From now on let $\mathbb{K} = \mathbb{Q}$. The truth table of f can be recovered from its NNF by the formula $f(u) = \sum_{a \leq u} \lambda_a, \forall u \in \mathbb{F}^n$, where $a \leq u \iff \forall i \in \{1, \ldots, n\} a_i \leq u_i$. Conversely, it is possible to derive an explicit formula for the coefficients of the NNF by means of the truth table of f.

Proposition 2.1. Let f be any integer-valued function on \mathbb{F}^n . For every $u \in \mathbb{F}^n$, the coefficient λ_u of the monomial X^u in the NNF of f is:

$$\lambda_u = (-1)^{w(u)} \sum_{a \in \mathbb{F}^n | a \preceq u} (-1)^{w(a)} f(a) \,. \tag{1}$$

Let $f, g \in \mathcal{B}_n$. The distance d(f, g) between f and g is the number of $v \in \mathbb{F}^n$ such that $f(v) \neq g(v)$. It is obvious that $d(f, g) = d(\underline{f}, \underline{g}) = w(\underline{f} + \underline{g})$.

Definition 2.2. Let $f \in \mathcal{B}_n$. The nonlinearity of f is the minimum of the distances between f and any affine function, i.e. $N(f) = \min_{\alpha \in \mathcal{A}_n} d(f, \alpha)$.

Using a simple divide-and-conquer butterfly algorithm it is possible to compute the ANF from the truth-table of a B.f., by performing $O(n2^n)$ bit operations, and storing $O(2^n)$ bits. This algorithm is known as the *fast Möbius transform*. From the ANF it is possible to compute the nonlinearity of a B.f. by means of a similar algorithm known as the *fast Walsh transform*, requiring $O(n2^n)$ integer sums and storing $O(2^n)$ integers.

Faster methods are known in particular cases, for example when the ANF is a sparse polynomial [Qal13].

3 Computing the nonlinearity using fast polynomial evaluation

Let A be the variable set $A = \{a_i\}_{0 \le i \le n}$. We denote by $\mathfrak{g}_n \in \mathbb{F}[A, X]$ the following polynomial:

$$\mathfrak{g}_n = a_0 + \sum_{i=1}^n a_i x_i \; .$$

Determining the nonlinearity of $f \in \mathcal{B}_n$ is the same as finding the minimum weight of the vectors in the set $\{\underline{f} + \underline{g} \mid g \in \mathcal{A}_n\} \subset \mathbb{F}^{2^n}$. We can consider the evaluation vector of the polynomial \mathfrak{g}_n as follows (see [SS07]):

$$\underline{\mathfrak{g}}_{\mathfrak{n}} = (\mathfrak{g}_n(A, \mathsf{p}_1), \dots, \mathfrak{g}_n(A, \mathsf{p}_{2^n})) \in (\mathbb{F}[A])^{2^n}$$

From now on we present original results.

For each $0 \le i \le 2^n$, we define the following Boolean affine polynomials:

$$f_i^{(\mathbb{F})}(A) = \mathfrak{g}_n(A, \mathsf{p}_i) + f(\mathsf{p}_i).$$

We also define

$$f_i^{(\mathbb{Z})}(A) = \operatorname{NNF}(f_i^{(\mathbb{F})}(A)).$$

Definition 3.1. We call $\mathfrak{n}_f(A) = f_1^{(\mathbb{Z})}(A) + \cdots + f_n^{(\mathbb{Z})}(A) \in \mathbb{Z}[A]$ the integer nonlinearity polynomial (or simply the nonlinearity polynomial) of the B.f. f. For any $t \in \mathbb{N}$ we define the ideal $\mathcal{N}_f^t \subseteq \mathbb{Q}[A]$ as follows:

$$\mathcal{N}_{f}^{t} = \langle E[A] \bigcup \{ f_{1}^{(\mathbb{Z})} + \dots + f_{2^{n}}^{(\mathbb{Z})} - t \} \rangle = \langle E[A] \bigcup \{ \mathfrak{n}_{f} - t \} \rangle$$
(2)

Notice that the integer evaluation vector \mathbf{n}_f represents all the distances of f from all possible affine functions in n variables, and so the following theorem is straitforward.

Theorem 3.2. The variety of the ideal \mathcal{N}_f^t is non-empty if and only if the Boolean function f has distance t from an affine function. In particular, N(f) = t, where t is the minimum positive integer such that $\mathcal{V}(\mathcal{N}_f^t) \neq \emptyset$.

Thus, to compute the nonlinearity of f we have to find the minimum nonnegative integer t in the set of the evaluations of \mathfrak{n}_f , that is, in $\{\mathfrak{n}_f(\bar{a}) \mid \bar{a} \in \{0,1\}^{n+1} \subset \mathbb{Z}^{n+1}\}$.

Now we claim a theorem to calculate the coefficients of the nonlinearity polynomial. Proof is omitted due to lack of space (see [BSS14]).

Theorem 3.3. Let $v = (v_0, v_1, \ldots, v_n) \in \mathbb{F}^{n+1}$, $\tilde{v} = (v_1, \ldots, v_n) \in \mathbb{F}^n$, $A^v = a_0^{v_0} \cdots a_n^{v_n} \in \mathbb{F}[A]$ and $c_v \in \mathbb{Z}$ be such that $\mathfrak{n}_f = \sum_{v \in \mathbb{F}^{n+1}} c_v A^v$. Then the coefficients of \mathfrak{n}_f can be expressed as:

$$c_v = \sum_{u \in \mathbb{F}^n} f(u) = w(\underline{f}) \quad \text{if } v = 0 \tag{3}$$

$$c_v = (-2)^{\mathsf{w}(v)} \sum_{\substack{u \in \mathbb{F}^n \\ v \leq u}} \left[f(u) - \frac{1}{2} \right] \quad if \ v \neq 0 \tag{4}$$

4 Complexity considerations

We write the algorithm (Algorithm 1) to calculate the nonlinearity polynomial in $O(n2^n)$ integer operations.

In Figure 1 Algorithm 1 is shown for n = 3.

Algorithm 1 Algorithm to calculate the nonlinearity polynomial \mathfrak{n}_f in $O(n2^n)$ integer operations.

Input: The evaluation vector f of a B.f. $f(x_1, \ldots, x_n)$ **Output:** the vector $c = (c_1, \ldots, c_{2^{n+1}})$ of the coefficients of \mathfrak{n}_f Calculation of the coefficients of the monomials *not* containing a_0 1: $(c_1, \ldots, c_{2^n}) = f$ 2: for i = 0, ..., n - 1 do 3: $b \leftarrow 0$ 4: repeat 5: for $x = b, \ldots, b + 2^i - 1$ do 6: $c_{x+1} \leftarrow c_{x+1} + c_{x+2^i+1}$ 7:if x = b then 8: $c_{x+2^{i}+1} \leftarrow 2^{i} - 2c_{x+2^{i}+1}$ 9: else 10: $c_{x+2^i+1} \leftarrow -2c_{x+2^i+1}$ 11: end if end for 12: $b \leftarrow b + 2^{i+1}$ 13:until $b = 2^n$ 14:15: end for Calculation of the coefficients of the monomials containing a_0 16: $c_{1+2^n} \leftarrow 2^n - 2c_1$ 17: **for** $i = 2, ..., 2^n$ **do** 18: $c_{i+2^n} \leftarrow -2c_i$ 19: **end for** 20: return c

 $(x_1, x_2, x_3) = f(x_1, x_2, x_3)$ Step 1 $Step \ 2$ $Step \ 3$ 000 $e_1 + e_2$ $e_1 + e_2 + e_3 + e_4$ $e_1 + e_2 + e_3 + e_4 + e_5 + e_6 + e_7 + e_8$ $1 - 2e_2$ $2 - 2e_2 - 2e_4$ 001 $4 - 2e_2 - 2e_4 - 2e_6 - 2e_8$ $e_{3} + e_{4}$ $2 - 2e_3 - 2e_4$ 010 $4 - 2e_3 - 2e_4 - 2e_7 - 2e_8$ $1 - 2e_4$ 011 $-2 + 4e_4$ $-4 + 4e_4 - 4e_8$ $\frac{1}{2}$ + $e_5 + e_6$ \rightarrow + $e_5 + e_6 + e_7 + e_8$ 100 $4 - 2e_5 - 2e_6 - 2e_7 - 2e_8$ $2 - 2e_6 - 2e_8$ $1 - 2e_6$ $-4 + 4e_6 - 4e_8$ 101 $e_7 + e_8$ $2 - 2e_7 - 2e_8$ 110 $-4 + 4e_7 - 4e_8$ $4 - 8e_8$ 111 $1 - 2e_8$ $-2 + 4e_8$

Figure 1: Butterfly scheme to obtain a fast computation of the nonlinearity polynomial coefficients, where $(e_1, \ldots, e_8) = (f(\mathbf{p}_1), \ldots, f(\mathbf{p}_8))$.

We claim the following theorem without proof (see [BSS14]).

Theorem 4.1. Algorithm 1 requires $O(n2^n)$ integer sums and doublings, in particular circa $n2^{n-1}$ integer sums and circa $n2^{n-1}$ integer doublings, and the storage of $O(2^n)$ integers of size less than or equal to 2^n .

In Table 1 we report the coefficients of growth of the analyzed algorithm and the standard algorithm which uses the fast Walsh transform¹, comparing them with the value $\log_2 \left[\frac{(n+1)2^{n+1}}{n2^n}\right]$. For each algorithm we compute the average time t_n to compute the nonlinearity of a B.f. with n variables and the average time t_{n+1} to compute the nonlinearity of a B.f. with n+1 variables. Then we report in the table the expected theoretical value $\log_2 \left(\frac{t_{n+1}}{t_n}\right)$. Thanks to theorem 4.1

n	4-5	5-6	6-7	7-8	8-9	9-10	10-11
$\frac{\log_2\left[\frac{(n+1)2^{n+1}}{n2^n}\right]}{FWT}$ NLP+FPE	1.22	1.17	1.14	1.12	1.11	1.09	1.09
FWT	0.90	0.98	1.01	1.22	0.95	1.25	1.07
NLP+FPE	1.02	1.09	1.13	1.07	1.17	1.07	1.11

Table 1: Experimental comparisons of the coefficients of growth of the analyzed algorithm.

and known facts on fast polynomial evaluation, we obtain:

Theorem 4.2. Determining the coefficients of the polynomial \mathfrak{n}_f from the truth table of f and then finding $N(f) = \min\{\mathfrak{n}_f(\bar{a}) \mid \bar{a} \in \{0,1\}^{n+1}\}$ requires a total $O(n2^n)$ integers operations (sums and doublings).

5 Acknowledgments

The author would like to thank his supervisor, Massimiliano Sala. An extended version of this note, jointly with M. Sala and I. Simonetti, can be found at [BSS14].

References

- [BSS14] E. Bellini, I. Simonetti, and M. Sala. Nonlinearity of Boolean functions: an algorithmic approach based on multivariate polynomials. Preprint, 2014.
- [Çal13] Çağdaş Çalık. Computing cryptographic properties of boolean functions from the algebraic normal form representation. PhD thesis, Middle East Technical University, 2013.
- [Car10] C. Carlet. Boolean functions for cryptography and error correcting codes. Boolean Models and Methods in Mathematics, Computer Science, and Engineering, pages 257– 397, 2010.
- [CG99] C. Carlet and P. Guillot. A new representation of Boolean functions. In Applied Algebra, Algebraic Algorithms and Error-Correcting Codes, pages 94–103. Springer, 1999.
- [MS77] F. J. MacWilliams and N. J. A. Sloane. The theory of error-correcting codes. I. North-Holland Publishing Co., Amsterdam, 1977. North-Holland Mathematical Library, Vol. 16.
- [SS07] M. Sala and I. Simonetti. An algebraic description of Boolean functions. Proc. of WCC 2007, pages 343–349, 2007.

 $^{^{1}}$ To compute the values in the columns FWT and NLP+FPE we tested 15000 random B.f. 's.