

George William Scott Blair – the pioneer of fractional calculus in rheology¹

Sergei Rogosin^{1,2}, Francesco Mainardi³

¹IMPACS: Institute of Mathematics, Physics and Computer Sciences,
Aberystwyth University, Wales, UK
ser14@aber.ac.uk

²Department of Economics, Belarusian State University, Minsk, Belarus
rogosin@bsu.by

³Department of Physics, University of Bologna, and INFN, Italy
francesco.mainardi@bo.infn.it

Abstract

The article shows the pioneering role of the British scientist, Professor G.W.Scott Blair, in the creation of the application of fractional modelling in rheology. Discussion of his results is presented. His approach is highly recognized by the rheological society and is adopted and generalized by his successors. Further development of this branch of Science is briefly described in this article too.

Keywords : fractional calculus, rheology, Scott Blair’s fractional model, hereditary solid mechanics, relaxation.

2010 MSC: 26A33, 60G22, 74-03, 74D05, 76-03, 76A10.

Foreword to the second extended version.

After appearance of the first version of the paper in arXiv it has got an interest as from Mathematical as from Rheological Society. In March 2014 one of the authors (SR) made a report at the research seminar at Aberystwyth University. The content of this seminar was described later by Prof. Simon Cox in his note published at Rheological Bulletin [12]. For readers’ convenience we present this note in the Appendix to the extended version of this ArXiv paper. We also add some additional references justifying the role of Scott Blair’s results ([23], [24], [30]). On this respect it is interesting to mention the “Bingham Lecture” [32] by G.H. McKinley. We pay attention of the

¹Short version published at *Communications in Applied and Industrial Mathematics*. – 2014, v. 6, no. 1. – e481. ISSN 2038-0909; DOI: 10.1685/journal.caim.481

readers to our recent book [21] devoted to the description of the properties of the Mittag-Leffler function, the Queen Function of the Fractional Calculus.

1 Introduction.

The interest to applications of fractional calculus in modelling of different phenomena in Physics, Chemistry, Biology is rapidly increasing in the recent three decades. First of all we have to point out the constitutive modelling of non-Newtonian fluids. The main reason is that the fractional models give us possibility for simple description of complex behaviour of a viscoelastic material. In pioneering (mainly experimental) works of 1940-1950th, it was discovered, for instance, that the relaxation processes in some materials exhibit an algebraic decay, which cannot be described in the framework of the Maxwell model based on exponential behaviour of the relaxation moduli.

In order to see perspective in the development of fractional models it is important to understand how such models appear, and what was really done by the pioneers. Among the works which are in the core of the first period of the fractional modelling one can single out the series of articles and monographs by G.W. Scott Blair. His role is not overestimated by the rheological society (see, e.g. [5], [13]), but anyway, some details of his work are still of great importance. We propose here an analysis of the results by G.W. Scott Blair along with their influence on the modern development of the fractional modelling in rheology.

1.1 Short biography by G.W. Scott Blair

Dr. George William Scott Blair (1902–1987) was born on 23 July 1902, of Scottish ancestry, in Weybridgein Surrey, England. After graduated a famous public school at Charterhouse he went to Trinity College Oxford in 1920, where he studied Chemistry, with Prof. Sir Cyril Hinshelwood as his tutor. He carried out a one-year research project in colloid chemistry to complete his master thesis with honour degree.

After graduating Scott Blair was employed as a colloid chemist with a Manchester firm of Henry Simon, working there on the viscometry of flour suspensions, publishing his first rheology paper in 1927. In 1926 he was offered a post in the Physics Department of the Rothamsted Experimental Station, where he was working on the flow properties of soils and clays until

1937. It was there, where he made with his colleagues the first quantitative study of so called sigma-phenomenon, which was originally described by Bingham and Green in 1919. Schofield and Scott Blair (see [49]) studied this phenomenon from 1930 at Rothamsted for soil and clay pastes and named it “sigma effect”. These studies were probably unknown to Fåhræus and Lidquist, who first discovered the sigma effect for blood, referred to as the “Fåhræus-Lindquist phenomenon”. At this period some preliminary experiments were provided by Scott Blair which led him later to the necessity to consider anomalous relationship between stress, strain and time (see, e.g., [50]).

In 1929 Scott Blair took up a Rockefeller Fellowship at Cornell University in Ithaca, New York state. He worked there on the flow of potter’s clay and developed a means of measuring its plasticity. He attended the inaugural meeting of Society of Rheology in December 1929 in Washington DC, met there all pioneers of Rheology, Eugene Bingham and Markus Reiner among them, and many became his life-time friends.

After returning to Rothamsted, Scott Blair made rheological research on honey and flour doughs. He also studied together with the well-known psychologist David Katz, psychophysical problems in bread making. His interest in psychology led him, together with F.M.V. Coppen, to initiate a new field, for which he coined the word “psychorheology”. It is considered as one of the fields of biorheology.

In 1936 he submitted his PhD thesis to the University of London, and it was examined by Prof. Freundlich. The same institution later awarded him a D.Sc. for his labours in rheology (probably the first ever rheology D.Sc.) In 1937 he joined National Institute for Research in Dairying, University of Reading as a head of Chemistry but soon took over the newly formed Physics Department and remained in that position until his retirement thirty years later.

In 1940 the British Rheological Society was founded. Scott Blair played a prominent role and took active part in the development of rheology. He was a Founder-Member and, later, president of the British Society of Rheology (1949–1951). He took part in the organization of the First International Congress on Rheology, held at Scheveningen, Holland, in 1948. He was a Secretary of the Second International Congress on Rheology in Oxford in 1953 and a member of Committee on Rheology, set up by the International Council of Scientific Union. Scott Blair was given to flights of fancy into psychorheology, fractional differentiation etc. In 1969 he was awarded the

Poiseuille Gold Medal of the International Society of Haemorheology (now Biorheology) and in 1970 he received the Founders Gold Medal of the British Society of Rheology. Together with J. Burgers he published a monograph on rheological nomenclature [7]. For many years Scott Blair was the Chairman of the British Standard Institute Committee on Rheological Nomenclature.

During almost a half of a century, George W. Scott Blair was one of the leading rheologists. Beginning from 1957 Scott Blair devoted his experimental and theoretical work entirely to hemorheology. Since the foundation of the International Society of Hemorheology in Reykjavik, Iceland in 1966, he was a member of its Council and acted as Chairman of its Committee on Standards and Terminology. After he retired he worked on the flow and coagulation of blood at the Oxford Haemophilia Centre.

Scott Blair was very active in publication and editorial work. He was a co-founder of the Journal “Biorheology” and its Co-Editor-in-Chief from its inception in November 1959 to December 1978 (see [11]). The books and research papers of Scott Blair were donated to the British Society of Rheology and later deposited in the Library of Aberystwyth University in early 1980’s. The collection has over 550 books and its aim is to develop this “into an up-to-date library of rheological literature available to all members of Society”. Rheology Abstracts and the British Society of Rheology Bulletin are two journals published by/for the Society which form an important of the Collection. The books and journals catalogued online (access via <http://primo.aber.ac.uk>).

2 Rheology and Psychophysics

It was Professor Bingham who had chosen the name “Rheology” for this branch of the Science and gave the definition of it: “The Science of Deformation and Flow of Matter” (see [53]) motivated by Heraclitus’ quote “ $\pi\alpha\nu\tau\alpha\ \rho\epsilon\iota$ ” (“everything flows”). Rheology is one of the very few disciplines having exact day of its birth, April 29, 1929, when the preliminary scope of the Society of Rheology was set up by a committee met at Columbus, Ohio. Anyway, the ancient Egyptian scientist Amenemhet (ca 1600 BC), who made the earliest application of the viscosity effect, can be considered as the first rheologist (see, *e.g.*, [52]).

The observables in rheology are deformations or strains, and the changes of strains in time. Changes of strains in time constitute a flow. Thus, these

changes are generally associated with internal flow of certain kind. States of stress are inferred either from the comparative strain behaviour of complex and simple systems in interaction or from the behaviour of a known mass in the gravitational field. In physical testing, stresses (S), strains (σ) or their differentials with respect to time ($\dot{S}, \dot{\sigma}$) are normally held constant, leaving either a length to be measured, or the time (t).

There is a group of fluids which is characterized by a coefficient of viscosity for a specific temperature. These fluids, known as Newtonian fluids, were singled out by Newton who proposed the definition of resistance (or viscosity in modern language) of an ideal fluid. Pioneering work on the laws of motion for real (i.e. non-ideal) fluids with finite viscosities was carried out by Navier [36] and later by Stokes [66]. The Navier-Stokes equation enabled, among other things, prediction of velocity distributions and flow between rotating cylinders and cylindrical tubes (see [13]).

Nowadays rheology generally accounts for the behaviour of non-Newtonian fluids, by characterizing the minimum number of functions that are needed to relate stresses with rate of change of strains or strain rates. This kind of fluids is called Newtonian since Newton's introduction the concept of viscosity.

In practice, rheology is concerned with extending continuum mechanics to characterize flow of materials, that exhibits a combination of elastic, viscous and plastic behaviour by properly combining elasticity and (Newtonian) fluid mechanics. In [13] the main directions in the development of the rheology are described. First of all this is linear *viscoelasticity*. One of the most important early contribution in this area is the work by Maxwell [31]. In order to explain the behaviour of the materials which are neither truly elastic nor viscous he proposed a constant relaxation time (t_r) and justified implicitly the model of a dash-pot and spring in series. Anyway, he realized that for some materials the assumption of constancy of the relaxation time is oversimplification, in these cases t_r has to be a function of stress. Meanwhile, the notion of Maxwell's units (i.e. pieces of a material having constant relaxation time) has been widely explored by rheologists. Later the conception of the "orientation times" τ has been developed (see, *e.g.*, [2]). It is considered unit stress conditions, supposing that the strain is approaching to an equilibrium value. Thus, the immediate Hookean strain is first subtracted and τ is defined as the time taken for the remaining strain, resulting from the orientation of the chains.

Another direction which was singled out in [13] is the study of *generalized Newtonian materials*. This type of fluid behaviour is associated with the work by Bingham [6] who proposed so called yield stress to describe the flow of paints. In [60], it has been pointed out the close similarity between the usual experimental Bingham curve and the curve of a high power-law. Thus, it shows possibility of existence of systems for which the Bingham plot gives a fairer and simpler account of the data.

The study of *non-linear viscoelasticity* started at the beginning of XXs century, when the area of rheology was most rapidly grown (see, *e.g.* [13]). Thus, Poynting [42] in his experiment discovered that loaded wires increased by a length that was proportional to the square of the twist, what did not correspond to the usual expectation of the linear viscoelasticity theory. Probably the first theoretical work on non-linear viscoelasticity was done by Zaremba [73], who extended linear theory to the non-linear regime by introducing corotational derivative in order to incorporate a frame of reference that was translating and rotating with the material. More extended description of the results in non-linear viscoelasticity can be found in [13] (see also [28], [69] and references therein).

Not all properties of flowing matter can be interpreted in term of real rheological sense. In this case psychophysical approach with its psychophysical experiments can be helpful. Psychophysics is defined as the scientific study of the relation between stimulus and sensation (see, *e.g.* [20]). Psychophysicists usually employ experimental stimuli that can be objectively measured. Psychophysical experiments have traditionally used three methods for testing subjects' perception in stimulus detection and difference detection experiments: the method of limits, the method of constant stimuli and the method of adjustment.

G.W. Scott Blair widely used psychophysical experiments in his research (see [54]). Therefore, it is interesting to recall how he described the role of psychophysics in rheology (see [53]): "The complex and commercially important rheological "properties" of many industrial materials are still assessed subjectively by handling in factory and are expressed in terms "body", "firmness", "spring", "deadness", shortness", "nerve", etc. - concepts which cannot be interpreted ... in terms of simple rheological properties at all. In view of this fact ... it is clearly advisable to know something of the accuracy with which these handling judgements can be made and, by bulking sufficiently large numbers of data together so that reproducibility is ensured, to attempt to correlate the entities so derived with manageable functions of

$S : \sigma : t$. A start has been made in this direction and not only have a number of reproducible regularities been observed, but the information obtained has laid the foundations of a theory of “*Quasi-properties*” which it is hoped will facilitate the study of purely “physical” rheology of complex materials.”

This observation is a core of Scott Blair’s method which he used along his career.

3 Scott Blair Fractional Element

3.1 Nutting’s Law

In 1921 Nutting reported ([37]) about his observation that mechanical strains appeared at the deformation of the viscoelastic materials decreasing as power-type functions in time. From a series of experiments, which covered a range of materials from the elastic solid to the viscous fluid, Nutting suggested a general formula relating shear stress, shear strain and time, whenever shear stress remains constant:

$$\sigma(t) \sim C\Delta S \cdot t^{-\alpha}, \quad (1)$$

with constant order $\alpha \in (0, 1)$ which is close to $1/2$ for many materials. This conclusion was in a strong contradiction to the standard exponential law. Later the Nutting’s observation was justified by Gemant who studied the properties of viscoelastic materials under harmonic load. It was shown that the memory function $\eta(t)$ can have power-type relaxation behaviour proportional to $t^{-3/2}$. In 1950 Gemant published a series of 16 articles entitled “Frictional Phenomena” in Journal of Applied Physics since 1941 to 1943, which were collected in a book of the same title [18]. In his eighth chapter-paper [17, p. 220], he referred to his previous articles [15], [16] for justifying the necessity of fractional differential operators to compute the shape of relaxation curves for some elasto-viscous fluids.

Gemant has used half-differential, but in his later papers he says that fractional differential “only occurs as a useful mathematical symbol, whereas the underlying elementary process, whatever it may be, will probably contain differential quotient of an integral order”.

Scott Blair surely knew the attempts by Gemant (see, *e.g.* [63]) to generalize Maxwell’s theory by changing various (but integer) powers in complex modulus of the Maxwell Fluid Model to fractional powers. In fact, Scott Blair (together with Coppen) also came to the form of Nutting equation,

but from another consideration. They argued that the material properties are determined by various states between an elastic solid and a viscous fluid, rather than a combination of an elastic and a viscous element as proposed by Maxwell. In [57] it was pointed out that, since for Hookian solids, strain is proportional to stress and to unit power of time, for intermediate materials, it might be expected to be proportional to stress and to some fractional power of time with exponent α , $0 < \alpha < 1$ and described this relation in the form

$$\psi = S^\beta \sigma^{-1} t^\alpha, \quad (2)$$

where proportionality coefficient ψ is a constant. Derived in this way the equation (2) looks entirely empirical, though the fundamental significance of α (which is called the *dissipation coefficient*) is shown in psychophysical experiments described by Scott Blair and Coppen (see [58], [59], [61]).

A comparison of the partially differentiated Nutting equation and Maxwell's equation may be written (see [51]), namely, for Nutting:²

$$-\left(\frac{\partial S}{\partial t}\right)_\sigma = \frac{\alpha}{\beta} \frac{S}{t}, \quad (3)$$

and for Maxwell:

$$-\left(\frac{\partial S}{\partial t}\right)_\sigma = \frac{S}{t_r}. \quad (4)$$

Since Nutting equation gives $t = \psi^{1/\alpha} S^{-\beta/\alpha} \sigma^{1/\alpha}$, it is apparent that the Nutting treatment postulates a single *relaxation time* proportional to a power of the stress. This is simplest possible way of implementing Maxwell's suggestion that relaxation time t_r may be some function of stress. From the other side it justifies the believe that the use of fractional calculus in description of processes toward equilibrium is necessary if one has to keep the Newtonian time scale.

3.2 Scott Blair's fractional model

It was suggested in [61] that, in considered cases, comparative firmness is judged neither by σ , nor by $\dot{\sigma}$, nor by any mixture of these two magnitude,

²The suffix σ indicates shear strains.

but by some intermediate entity, namely by fractional derivative $\frac{\partial^\nu \sigma}{\partial t^\nu}$.³ More exactly, he wrote: The general constitutive equation “... is applicable to integral values of n but a more general equation may be used even n is a fraction. The numerical coefficient is expressed as a quotient of Γ -functions and may be written

$$\frac{\delta^n \sigma}{\delta t^n} = \frac{\Gamma(k+1)}{\Gamma(k-n+1)} t^{k-n} \Psi^{-1} S. \quad (5)$$

The expression $\Gamma(k+1)$ is given by

$$\int_0^\infty e^{-x} x^k dx$$

and, whatever the value of n , the Γ factor is a number independent of Ψ_m and t_c , so that the validity of the plot is unchallenged.”

In his work Scott Blair did not specify what kind of fractional derivative he used. From the way how he has calculated derivative of any power we can conclude that this is the standard Riemann-Liouville derivative. It is quite instructive to cite some words by Scott-Blair quoted by Stiassnie in their correspondence, see [65]: *I was working on the assessing of firmness of various materials (e.g. cheese and clay by experts handling them) these systems are of course both elastic and viscous but I felt sure that judgements were made not on an addition of elastic and viscous parts but on something in between the two so I introduced fractional differentials of strain with respect to time.* Later, in the same letter Scott-Blair added: *I gave up the work eventually, mainly because I could not find a definition of a fractional differential that would satisfy the mathematicians.*

The above said *Principle of Intermediacy* was discussed in details by Scott Blair in [51] basing on purely physical grounds. The theory of fractional modelling in rheology is developed by Scott Blair, Veinoglou and Caffyn in [63]. In [53, p. 30] it is briefly summarized: “... times are normally defined as equal when “free” Newtonian bodies (or alternatively light) traverse equal (superposable) distances in them. This leads to a definition of velocity as the first differential of length with respect to time which, because of this

³In fact, few misprints had been made in [61] later corrected by Scott Blair. Thus, in the original reprint of [61] in the Scott Blair reading room one can find hand-written corrections made by Scott Blair.

definition of time equality, is constant for Newtonian bodies; and to the second differential, called acceleration.

When bodies are not influenced by other bodies, and their velocities change with time, a force is postulated and defined as rate of change of (velocity \times mass). It is long been realized that the Newtonian time scale arbitrary (see [41, p. 80]) and in the case of a complex plastic being strained, the rheologically active units are certainly not independent Newtonian bodies. It should, therefore, be easy to choose a non-Newtonian time equality definition which would reduce the entities by which firmness is judged to simple whole-number⁴ differential expression. The use of separate time scales for different materials is not convenient, however, so Newtonian time is used, but, as a result of this arbitrary procedure, the derived constants cannot be expected to be built up entirely from whole-number differentials. It is thus apparent that fractional differential is an essential feature of our whole mode of approach.”

In [56] are discussed the circumstances under which it is practicable to express the Nutting equation and its fractional derivatives in a simple dimensional form. Three main principles of a new proposal are formulated: (1) the fact that the treatment does not lead to any understanding of structure of the materials or of their molecular configurations; (2) the only entities are used whose dimensions depend of the nature of of the material; (3) fractional derivatives and corresponding coefficients are understood as something intermediate between zero and first derivatives and corresponding coefficients. Scott Blair highly supported (see, it e.g. [60]) the ideas by Nutting supposing that for that moment it describe a special but very frequently adequate cases. Anyway, he though that the phenomenon dealing with Nutting equation are related to the fundamental structure of materials.

Fractional derivative of order μ , $0 < \mu < 1$, with respect to time t of the Nutting equation (in the form (2) gives (see [61], [63])

$$\frac{\partial^\mu \sigma}{\partial t^\mu} = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - \mu + 1)} t^{\alpha - \mu} \psi^{-1} S. \quad (6)$$

Another way to justify this relation is to introduce quasi-property χ_1 by the *Principle of Intermediacy*

$$\chi_1 = S \div \frac{\partial^\mu \sigma}{\partial t^\mu}, \quad (7)$$

⁴I.e. non-integer order.

since the viscosity can be defined by the relation $\eta = S \div \frac{d\sigma}{dt}$, and shear modulus as $n = S \div \sigma$.

The relation (7) can be integrated to give

$$\sigma = S^\beta \left(At^{\alpha'} + Bt^{\alpha'-1} + Ct^{\alpha'-2} + \dots \right), \quad (8)$$

where A , B , C etc. are constants. Clearly this equation coincides with Nutting equation if $A \gg B, C \dots$

4 Fractional models in rheology

After the first applications of the fractional derivatives in the modelling of the processes in rheology several other fractional models were proposed to describe certain rheological phenomenon. We briefly outline here the most discussed models of such a type.⁵

Gerasimov [19] used similar arguments as Scott Blair (in, e.g. [61], [63]), namely, interpolation between Hook and Newton's law, in order to introduce a rheological constitutive equation in terms of a precise notation of fractional derivative

$$\sigma(t) = \kappa_\alpha D_{0+,t}^\alpha \varepsilon(t). \quad (9)$$

This equation was used in [19] for description of the flow of the viscoelastic between two parallel plates. He obtained an exact solution by using operational method. He has started his consideration by appealing to the Boltzmann equation saying that from experiments follow the importance of a special case of the Boltzmann equation corresponding only to the hereditary part of the stress $\sigma(t)$

$$\sigma(t) = \int_0^t G(t-\tau) \dot{\varepsilon}(\tau) d\tau, \quad (10)$$

or even the processes for which $\sigma(t)$ has a memory on the velocity of all earlier deformations

$$\sigma(t) = \int_0^t K(t-\tau) \dot{\varepsilon}(\tau) d\tau. \quad (11)$$

⁵Here and in what follows we will use modern notations for stress (σ) and strain (ε) that are not be confused with the corresponding notations used by Scott Blair, namely (S) and (σ). Furthermore we will write D_{0+}^α to denote the Riemann-Liouville fractional derivative implicitly adopted by Scott Blair.

For the kernel in this integro-differential relation he claim that for certain materials this kernel (relaxation function) has the a form

$$K(\tau) = \frac{A}{\tau^\alpha}, \quad A > 0, \quad 0 < \alpha < 1.$$

Hence, equation (11) can be written as⁶

$$\sigma(t) = \frac{\kappa_\alpha}{\Gamma(1-\alpha)} \int_0^t \frac{\dot{\varepsilon}(t-\tau)}{\tau^\alpha} d\tau. \quad (12)$$

In particular, $\alpha = 1$ gives us the Newton law, and $\alpha = 0$ corresponds to Hookean law.

Same approach is used in the above article for the study of the rotational viscoelastic flow between two concentric cylinders.

Similar to (9) formulation of the fractional model was proposed by Slonimsky [64]).

Rabotnov (see [43] and more extended description in his monograph [44]) presented a general theory of hereditary solid mechanics using integral equations (see also [25], where the use of integral equations for viscoelasticity was revisited and interjects fractional calculus into Rabotnov's theory by the introduction of the spring-pot was presented).

Rabotnov introduced an hereditary elastic rheological model with constitutive equation in form of Volterra integral equation with weakly singular kernel of special type⁷

$$\sigma(t) = E \left[\varepsilon(t) - \beta \int_0^{t_\alpha} R_\alpha(-\beta, t_\alpha - \tau) \varepsilon(\tau) d\tau \right], \quad (13)$$

where t_α is the aging time, $\alpha \in (-1, 0]$, $\beta \neq 0$, and the kernel R is represented in the form of power series

$$R_\alpha(\beta, x) = x^\alpha \sum_{n=0}^{\infty} \frac{\beta^n x^{n(\alpha+1)}}{\Gamma((n+1)(\alpha+1))}. \quad (14)$$

⁶Up to the constant multiplier the right-hand side coincides with the fractional derivative known as Caputo derivative.

⁷The integral stress - strain relationship by Rabotnov can be re-interpreted in terms of the fractional differential constitutive equation of Zener type, see later.

Rabotnov's kernel function $R_\alpha(\beta, x)$ is related to the well-known Mittag-Leffler function $E_{\alpha,\beta}(z)$ highly explored nowadays in the fractional calculus and its applications, namely

$$R_\alpha(\beta, x) = x^\alpha E_{\alpha+1,\alpha+1}(\beta x^{\alpha+1}). \quad (15)$$

Of course Scott-Blair did not know the Mittag-Leffler function and its asymptotic behaviours (stretched exponential for short times and power law for long times). Presumably that Scott-Blair had guessed the behaviour of the M-L function but he did not have the mathematical background being overall an experimentalist.

Both Scott Blair's model (6) and Gerasimov's model (9) are naturally considered later as special cases of *fractional Maxwell's model* with rheological constitutive equation of the form

$$\sigma(t) + \lambda^\alpha D_{0+,t}^\alpha \sigma(t) = E \lambda^\beta D_{0+,t}^\beta \varepsilon(t), \quad 0 \leq \alpha \leq \beta \leq 1, \quad (16)$$

where E is the shear modulus, and λ is the relaxation time. This equation generalizes celebrating Maxwell equation in which for the first time Newtonian law for viscous fluid and Hook's law for elastic solid are combined to describe the behaviour of visco-elastic media

$$\sigma(t) + \tau D_t \sigma(t) = E \tau D_t \varepsilon(t). \quad (17)$$

Partial case of fractional Maxwell's model is the so-called *three-parametric generalized Maxwell's model* with constitutive equation of the type

$$\sigma(t) + a_1 D_{0+,t}^\alpha \sigma(t) = b_0 \varepsilon(t), \quad 0 < \alpha < 1. \quad (18)$$

Another popular fractional model with three parameters is the *Kelvin-Voigt fractional model* that presumably for the first time was introduced by Caputo [8] in 1967,

$$\sigma(t) = b_0 \varepsilon(t) + b_1 D_{0+,t}^\alpha \varepsilon(t), \quad 0 < \alpha < 1. \quad (19)$$

It is a generalization of the classical Kelvin model having the following constitutive equation

$$\sigma(t) = E [\sigma(t) + \tau D_t \varepsilon(t)]. \quad (20)$$

More general constitutive equation corresponds to the so called *fractional Zener model*:

$$\sigma(t) + a_1 D_{0+,t}^\alpha \sigma(t) = b_0 \varepsilon(t) + b_1 D_{0+,t}^\alpha \varepsilon(t), \quad 0 < \alpha < 1. \quad (21)$$

formerly introduced in 1971 by Caputo and Mainardi [9]. Theoretical background for this was done by Bagley and Torvik, see [3], [4]. It has to be pointed out that the above considered Rabotnov's model (13) is equivalent to the fractional Zener model, see [68].

Sometimes the *Poynting-Thomson fractional model* is discussed with rheological constitutive equation of the type

$$\sigma(t) + \frac{E}{E_0} \lambda^{\alpha-\gamma} D_{0+,t}^{\alpha-\gamma} \sigma(t) + \frac{E}{E_0} \lambda^{\beta-\gamma} D_{0+,t}^{\beta-\gamma} \sigma(t) = E \lambda^\alpha D_{0+,t}^\alpha \varepsilon(t) + E \lambda^\beta D_{0+,t}^\beta \varepsilon(t), \quad (22)$$

$$(0 \leq \gamma \leq \alpha \leq \beta \leq 1.)$$

More extended discussion of the fractional models in rheology can be found in [10], [28], [40], [48], [72].

5 The second generation of fractional modelling in rheology

The development of the fractional differential approach in rheology is associated with such names as Scott-Blair, Bagley and Torvik, Caputo, Gorenflo and Mainardi [29], Friedrich, Schiessel and Blumen [14], [48], Metzler, Nonnenmacher, Glöckle, Klafter and Schlesinger [34], [35], Koeller [27], Podlubny and Heymans [22], Rossikhin and Shitikova [33], [46], [47], and others.

In a series of papers (see, *e.g.* [3], [4]) Bagley and Torvik extended the ideas of Gemant, and of Caputo–Mainardi [10]. They have shown that the complex modulus of many materials can be approximated by fractional powers in the frequency. They proposed a general fractional constitutive relation describing visco-elastic behaviour in its different appearance.

This approach has been successfully applied to describe rheological behaviour of organic glasses, elastomers, polyurethane, polyisobutylene, monodisperse polybutadiene and solid amorphous polymers in a wide temperature range (see for example [1] and references therein). In [71] and [67] have been derived equations governing the time-dependent indentation response for axisymmetric indenters into a fractional viscoelastic half-space and have proposed an original method for the inverse analysis of fractional viscoelastic properties and applied to experimental indentation creep data of polystyrene. The method is based on fitting the time-dependent indentation data (in the Laplace domain) to the fractional viscoelastic model response. It is shown

that the particular time-dependent response of polystyrene is best captured by a bulk-and-deviator fractional viscoelastic model of the Zener type. We shall dwell in details on fractional differential models of viscoelasticity and then consider a few standard hydrodynamic problems in the simplest model of this type.

It is impossible to describe the modern state in the fractional rheology. We refer interested readers to the recent monographs [28], [69], and to the survey paper [70] for some additional comments on pioneering works in applications of fractional calculus.

Acknowledgements.

One of the inspiration of this article was a possibility to work at the G.W. Scott Blair reading room at the Aberystwyth University. The authors are grateful to the official Custodian of the collection, Prof. S. Cox, for supporting the idea of the article and for fruitful discussion of the subject. The work by S.R. is partly supported by PEOPLE IAPP Project PIAP-GA-2009-251475 HYDROFRAC.

References

- [1] Alcoutlabi, M., Martinez-Vega, J.J.: The effect of physical ageing on the time relaxation spectrum of amorphous polymers: the fractional calculus approach. *J. Material Science*, **34**, 2361–2369 (1999)
- [2] Alexandrov, A.P., Lazurkin, J.S.: Strength of amorphous and crystallizing rubber polymers. *Doklady Akad Nauk SSSR*, **45**, 291-294 (1944)
- [3] Bagley, R.L., Torvik, P.J.: A theoretical basis for the application of fractional calculus to viscoelasticity, *J. Rheology*, **27**, 201-210 (1983)
- [4] Bagley, R.L., Torvik, P.J.: On the fractional calculus model of viscoelastic behavior, *J. Rheology*, **30**, 133-155 (1986)
- [5] Barnes, H.: George William Scott Blair MA PhD DSc FRIC FInstP (1902–1987) - the Man and his Work, In: “The Scott Blair Collection”, Aberystwyth, 6 p. (1999)

- [6] Bingham, E.C.: *Fluidity and Plasticity*, McGraw Hill, New York (1922)
- [7] Burgers, J.M., Scott Blair, G.W.: *Report on the Principles of Rheological Nomenclature*, North Holland P.C., Amsterdam (1949)
- [8] Caputo, M.: Linear models of dissipation whose q is almost frequency independent, part II, *Geophys. J. R. Astr. Soc.*, **13**, 529–539 (1967)
- [9] Caputo, M., Mainardi, F.: A new dissipation model based on memory mechanism, *Pure and Applied Geophysics*, **91**, 134–147 (1971)
- [10] Caputo, M., Mainardi, F.: Linear models of dissipation in anelastic solids, *Riv. Nuovo Cimento* (Ser. II), **1**, 161–198 (1971)
- [11] Copley, A.L.: George William Scott Blair (1902–1987): Obituary, *Thrombosis Research*, **51**, No. 4, 339–353 (1988)
- [12] Cox, S.: From mathematical curiosity to practical significance, *Rheology Bulletin*, **55**, No. 2, 45–46 (2014)
- [13] Doraiswamy D.: Origins of Rheology: a Short Historical Excursion, In: [<http://www.rheology.org/sor/publications/Rheology-B/Jan02/Origin-of-Rheology.pdf>] (mentioned in *Rheology Bulletin*, **71**, No. 1 (2002))
- [14] Friedrich, Chr.: Relaxation and retardation functions of the Maxwell model with fractional derivatives, *Rheol. Acta*, **30**, 151–158 (1991)
- [15] Gemant, A.: A method of analyzing experimental results obtained from elastiviscous bodies, *Physics*, **7**, 311–317 (1936)
- [16] Gemant, A.: On fractional differentials, *Phil. Mag.* (Ser. 7) **25**, 540–549 (1938)
- [17] Gemant, A.: Frictional phenomena: VIII, *J. Appl. Phys.* **13**, 210–221 (1942)
- [18] Gemant, A.: *Frictional Phenomena*, Chemical Publ. Co, Brooklyn N.Y. (1950)
- [19] Gerasimov, A.N.: Generalization of linear laws of deformation and its application to problems with internal friction, *Prikl. Mat. Mekh.*, **12**, 251–260 (1948) (in Russian)

- [20] Gescheider, G.: *Psychophysics: The Fundamentals* (3rd ed.), Lawrence Erlbaum Associates (1997)
- [21] Gorenflo, R., Kilbas, A.A., Mainardi, F., Rogosin, S.V.: *Mittag-Leffler Functions, Related Topics and Applications*, Springer Verlag (2014)
- [22] Heymans, N., Podlubny, I.: Physical interpretation of initial conditions for fractional differential equations with Rieman-Liouville fractional derivatives, *Rheol. Acta*, **45**, 765–771 (2006)
- [23] Jaishankar, A.: Fractional constitutive equations and the rheology of multiscale materials, *Rheology Bulletin*, **55**, No. 1, 4–14 (2014)
- [24] Jaishankar, A., McKinley, G.H.: Power-law rheology in the bulk and at the interface: quasi-properties and fractional constitutive equations, *Proc. R. Soc., A Math. Phys. Eng. Sci.* **469**, 2012.0284 (2013)
- [25] Koeller, R.C.: Application of fractional calculus to the theory of viscoelasticity, *ASME J. Appl. Mech.* **51**, 299307 (1984)
- [26] Koeller, R.C.: Polynomial operators, Stieltjes convolution, and fractional calculus in hereditary mechanics, *Acta Mechanica*, **58**, 251–264 (1986)
- [27] Koeller, R.C.: A theory relating creep and relaxation for linear materials with memory, *ASME J. Appl. Mech.*, **77**, 031008/1–9 (2010)
- [28] Mainardi, F.: *Fractional Calculus and Waves in Linear Viscoelasticity*, Imperial College Press, London and World Scientific, Singapore (2010)
- [29] Mainardi, F., Gorenflo, R.: Time-fractional derivatives in relaxation processes: a tutorial survey, *Fract. Calc. Appl. Anal.*, **10**, 269–308 (2007)
- [30] Mainardi, F., Spada, G.: Creep, relaxation and viscosity properties for basic fractional models in rheology, *The European Physical Journal, Special Topics*, **193**, 133–160 (2011)
- [31] Maxwell, J.C.: On the dynamical theory of gases, *Phil. Mag.* **35**, 129–146 (1868); 185–217 (1868)

- [32] McKinley, G.H., Jaishankar, A.: Critical Gels, Scott Blair and the Fractional Calculus of Soft Squishy Materials. *Bingham Lecture*. 85th Annual Meeting of the Society of Rheology, Montréal, Québec, October 2013.
web.mit.edu/nnf/McKinley_BinghamLecture2013_FractionalCalculus.pdf
- [33] Meshkov, S., Pachevskaya, G., Postnikov, V., Rossikhin, Y.: Integral representation of ε_Y -functions and their application to linear viscoelasticity, *Int. J. Eng. Sci.*, **9**, 387–398 (1971)
- [34] Metzler, R., Glöckle, W. G., Nonnenmacher, T. F.: Fractional model equation for anomalous diffusion, *Physica A*, **211**, 13–24 (1994)
- [35] Metzler, R., Klafter, J.: The restaurant at the end of the random walk: Recent developments in the description of anomalous transport by fractional dynamics, *J. Phys. A. Math. Gen.*, **37**, R161–R208 (2004)
- [36] Navier, C.L.M.H.: Sur les lois de l'équilibre et du mouvement des corps solides élastique, *Bull. Soc. Philomath.*, **75**, 177–183 (1823); also in *Memoire Acad. Sci.* **7**, 375–393 (1827), reported in Acad. Sci. at 14 May 1821
- [37] Nutting, P.G.: A new general law of deformation, *J. Frankline Inst.* **191**, 679–685 (1921)
- [38] Nutting, P.G.: A general stress-strain-time formula, *J. Frankline Inst.* **235**, 513–524 (1943)
- [39] Nutting, P.G.: Deformation in relation to time, pressure and temperature. *J. Frankline Inst.* **242**, 449–458 (1946)
- [40] Podlubny, I.: *Fractional Differential Equations*, Academic Press, New York (1999)
- [41] Poincare, H.: *La Valeur de la Science*, Flammarion, Paris (1904)
- [42] Poynting, J. H., Thomson, J. J.: *A Textbook of Physics. I: Properties of Matter*, Griffin, London (1913) [6th edition carefully revised]
- [43] Rabotnov, Yu.N.: Equilibrium of an elastic medium with after effect, *Prikl. Mat. Mekh.*, **12**, No. 1, 53–62 (1948) (in Russian)

- [44] Rabotnov, Yu.N.: *Elements of Hereditary Solid Mechanics*, Mir Publishers, Moscow (1980)
- [45] Rogosin, S., Mainardi, F.: George William Scott Blair – the pioneer of fractional calculus in rheology, *Communications in Applied and Industrial Mathematics*, **6**, no. 1, e481 (2014)
- [46] Rossikhin, Y., Shitikova, M.: Comparative analysis of viscoelastic models involving fractional derivatives of different orders, *Fract. Calc. Appl. Anal.*, **10**, No. 2, 111–121 (2007)
- [47] Rossikhin, Y., Shitikova, M.: Two approaches for studying the impact response of viscoelastic engineering systems: an overview, *Comp. Math. Appl.*, **66**, 755–773 (2013)
- [48] Schissel, H., Friedrich, Chr., Blumen, A.: Applications to problems in polymer physics and rheology. In: *Applications of Fractional Calculus in Physics* (R. Hilfer Ed.), World Scientific, Singapore (2000), pp. 331–376
- [49] Schofield, R.K., Scott Blair, G.W.: The influence of the proximity of a solid wall on the consistency of viscous and plastic materials. *J. Phys. Chem.* **34**, 248–262 (1930); III - **35**, 1212–1215 (1931); IV - **39**, 973–981 (1935)
- [50] Schofield, R.K., Scott Blair, G.W.: The relationship between viscosity, elasticity and plastic strength of a soft materials as illustrated by some mechanical properties of flour dough. - III. *Proc. Roy. Soc. A.* **141**, 72–85 (1933)
- [51] Scott Blair, G.W.: Analytical and integrative aspects of the stress-strain-time problem, *J. Sci. Instruments*, **21**, No. 5, 80–84 (1944)
- [52] Scott Blair, G.W.: *A Survey of General and Applied Rheology*, Pitman & Sons, London (1944)
- [53] Scott Blair, G.W.: The role of psychophysics in rheology, *J. Colloid Sci.* **2**, No. 1, 21–32 (1947)
- [54] Scott Blair, G.W.: Psychoreology: links between the past and present, *Journal of Texture Studies*, **5**, 3–12 (1974)

- [55] Scott Blair, G.W., Caffyn, J.E.: Significance of power-law relations in rheology, *Nature*, **155**, 171–172 (1945)
- [56] Scott Blair, G.W., Caffyn, J.E.: An application of the theory of quasi-properties to the treatment of anomalous strain-stress relations, *Phil. Mag.* (ser. 7), **40**, 80–94 (1949)
- [57] Scott Blair, G.W., Coppen, F.M.V.: The subjective judgement of the elastic and plastic properties of soft bodies: the differential thresholds for viscosities and compression moduli, *Proc. Roy. Soc. B*, **128**, 109–125 (1939)
- [58] Scott Blair, G.W., Coppen, F.M.V.: The subjective judgement of the elastic and plastic properties of the soft bodies, *British J. Psychol.*, **31**, Part 1, 61–79 (1940)
- [59] Scott Blair, G.W., Coppen, F.M.V.: The subjective conception of the firmness of the soft materials, *Amer. J. Psychol.*, **55**, 215–229 (1942)
- [60] Scott Blair, G.W., Coppen, F.M.V.: The classification of rheological properties of industrial materials in the light of power-law relations between stress, strain, and time, *J. Sci. Instr.*, **19**, 88–93 (1942)
- [61] Scott Blair, G.W., Coppen, F.M.V.: The estimation of firmness in soft materials, *Amer. J. Psychol.*, **56**, 234–246 (1943)
- [62] Scott Blair, G.W., Reiner, M.: The rheological law underlying the Nutting equation, *Appl. Sci. Res.*, **A 2**, 225–234 (1950)
- [63] Scott Blair, G.W., Veinoglou, B.C., in collaboration with Caffyn, J.E.: Limitation of the Newtonian time scale in relation to non-equilibrium rheological states and a theory of quasi-properties, *Proc. Roy. Soc. A*, **189**, 69–87 (1947)
- [64] Slonimsky, G.L.: On the law of deformation of highly plastic polymeric bodies, *Dokl. Akad. Nauk SSSR*, **110**, No. 2, 343–346 (1961) (in Russian)
- [65] Stiassnie, M. On the application of fractional calculus on the formulation of viscoelastic models, *Appl. Math. Modelling*, **3**, 300–302 (1979)
- [66] Stokes, G.G.: *Mathematical and physical papers*. Vol. **1-5** Reprint. New York and London: Johnson Reprint Corporation (1966).

- [67] Shahsavari R., Ulm F.-J.: Indentation analysis of fractional viscoelastic solids, *Journal of Mechanics of Materials and Structures*, **4**, 523–550 (2009)
- [68] Uchaikin, V.V.: *Method of Fractional Derivatives*, Artishok, Ul’yanovsk (2008) (in Russian)
- [69] Uchaikin, V.V.: *Fractional Derivatives for Physicists and Engineers*, Vol. II Applications, Springer, Berlin and Higher Education Press, Beijing (2013)
- [70] Valério, D., Machado, J. T. and Kiryakova, V.: Some pioneers of the applications of Fractional Calculus, *Fract. Calc. Appl. Anal.*, **17**, No. 2, 552–578 (2014)
- [71] Vandamme M., Ulm F.-J.: Viscoelastic solutions for conical indentation, *Int. J. Solids Struct.* **43**, 3142–3165 (2006)
- [72] West, B., Bologna, M., Grigolini, P.: *Physics of Fractal Operators*, Springer, New York - Berlin - Heidelberg (2003)
- [73] Zaremba, St.: Sur une forme perfectionnée de la théorie de la relaxation, *Bull. Acad. Sci. Cracow*, 594–614 (1903) (in French)

Appendix

From mathematical curiosity to practical significance

Rheology Bulletin, **55**, No. 2, 45–46 (2014)

In March, mathematicians in Aberystwyth were treated to an historical review of GW Scott Blair’s contributions to both philosophy and rheology, with particular emphasis on his development of fractional calculus from a mathematical curiosity to a tool of practical significance to rheology. The speaker, Professor Sergei Rogosin, is a Senior Marie Curie Fellow in the Department of Mathematics and Physics at Aberystwyth, on leave from Minsk University in Belarus. Professor Rogosin has worked on fractional calculus previously, and serendipitously discovered the BSR’s Scott Blair Collection on arriving in Aberystwyth. He used the collection to discover more about

Scott Blair's work on fractional calculus, and his talk was based on a recent paper with Francesco Mainardi (Bologna) [45].

As readers may know, not only did Scott Blair help found the BSR, and serve as one of its first presidents, he also contributed to the founding of the (American) Society of Rheology twenty years earlier. His interest in fractional calculus was motivated by trying to explain his experimental results on food rheology, trying to quantify effects such as "firmness" and "taste" and the influence of material memory on rheological response [63]. Gareth McKinley spoke about the same topic in Chicheley Hall last year at the INNFM meeting [32] (see also [24]). Professor Rogosin explained how Scott Blair picked up on Nutting's theoretical work, and Gemant's experiments some time later, indicating that fractional exponents in constitutive relations offered a way to interpolate between Newtonian liquids and Hookean solids, and consequently introduced the idea of quasiproperties. The speaker made the link with Volterra integral equations, which had been developed earlier in the 20th century, providing a robust mathematical framework in which to explore these relationships (although Scott Blair himself probably did not recognise this). At the same time as Scott Blair was working in this area, both Gerasimov and Rabotnov were tackling this problem of interpolation in similar ways and publishing in the Russian literature (although Scott Blair spoke Russian, the 1940s were of course a difficult time to exchange scientific information!). In particular, Rabotnov recognised that using fractional exponents offered the possibility of predicting power-law decay in time, rather than exponential decay, allowing much better fitting of the rheological behaviour of complex fluids. He later acknowledged Scott Blair's contributions in a textbook published in the 1980s; indeed, Scott Blair's contributions were widely valued both by experts in fractional modelling [27] and by rheologists [13].

The seminar was followed by a discussion of nonlinear elasticity and integral equations, upon which I shall not report here.

The Scott Blair Collection of books on rheology continues to grow thanks to a financial bequest to the BSR. Donations are also welcome. Further details are available at <https://www.aber.ac.uk/en/is/collections/scottblair/>

Simon Cox, Aberystwyth, March 2014