

(March 2014)

# On the zeros of some families of polynomials satisfying a three-term recurrence associated to Gribov operator

\*\*\*\*\*

Abdelkader Intissar

- *Equipe d'Analyse Spectrale, UMR-CNRS no:6134, University of Corsica Pascal Paoli. Faculty of Science and Technology, Quartier Grossetti, 20250 Corté, France. intissar@univ-corse.fr, Tél-Fax: 00 33 (0) 4 95 45 00 33.*

&amp;

- *Le Prador, 129, rue du Commandant Rolland, 13008 Marseille, France*

## Abstract

We consider families of tridiagonal- matrices with diagonal  $\beta_k = \mu k$  and off-diagonal entries  $\alpha_k = i\lambda k\sqrt{k+1}$ ;  $1 \leq k \leq n$ ,  $n \in \mathbb{N}$  and  $i^2 = -1$  where  $\mu \in \mathbb{C}$  and  $\lambda \in \mathbb{C}$ . In Gribov theory ([7], A reggeon diagram technique, Soviet Phys. JETP 26 (1968), no. 2, 414-423), the parameters  $\mu$  and  $\lambda$  are reals and they are important in the reggeon field theory. In this theory  $\mu$  is the intercept of Pomeron which describes the energy of dependence of total hadronic cross sections in the currently available range of energies and  $\lambda$  is the triple coupling of Pomeron.

The main motive of the paper is the localization of eigenvalues  $z_{k,n}(\mu, \lambda)$  of the above matrices which are the zeros of the polynomials  $P_{n+1}^{\mu, \lambda}(z)$  satisfying a three-term recurrence :

$$\begin{cases} P_0^{\mu, \lambda}(z) = 0 \\ P_1^{\mu, \lambda}(z) = 1 \\ \alpha_{n-1}P_{n-1}^{\mu, \lambda}(z) + \beta_n P_n^{\mu, \lambda}(z) + \alpha_n P_{n+1}^{\mu, \lambda}(z) = z P_n^{\mu, \lambda}(z); \quad n \geq 1 \end{cases}$$

If  $\mu \in \mathbb{R}$  and  $\lambda \in \mathbb{R}$  then the above matrices are complex symmetric, in this case we show existence of complex-valued function  $\xi(z)$  of bounded variation on  $\mathbb{R}$  such that the polynomials  $P_n^{\mu, \lambda}(z)$  are orthogonal with this weight  $\xi(z)$ .

---

**Keywords:** Jacobi-Gribov matrices; Non self-adjoint Gribov operator; Bargmann space; Polynomials satisfying a three-term recurrence; Eigenvalue problem; Reggeon field theory.

**MR(2010) Subject Classification :** 47B36, 47B39, 33C45.

---

## 1. Introduction and some preliminaries results

Usually, quantum Hamiltonians are constructed as selfadjoint operators; for certain situations, however, non-selfadjoint Hamiltonians are also of importance. In particular, the reggeon field theory (as invented by V. Gribov [7]) for the high energy behavior of soft processes is governed in zero transverse dimension by the non-selfadjoint operator:

$$\mathbb{H} = \mu a^* a + i\lambda a^*(a + a^*)a \quad (1.1)$$

Where  $a$  and  $a^*$  are the standard Bose annihilation and creation operators that:

$$[a, a^*] = I \quad (1.2)$$

and

$\mu$  and  $\lambda$  are real numbers ( $\mu$  is Pomeron intercept and  $\lambda$  is the triple coupling of Pomeron see Gribov [7] or the excellent survey of the development of Reggeon theory and its application to hadron interactions at high energies given by Borskov et al. at 2006 in [3].) and  $i^2 = -1$ .

In Bargmann representation [2], the principal spectral properties of  $\mathbb{H}$  have been studied in ref.[8].

The mathematical difficulties of this problem come of course from the non-self-adjointness of  $\mathbb{H}$ . Notice that this non-self-adjointness is a rather wild one; the word "wild" meaning here that the domains of the adjoint and anti-adjoint parts are not included in one another, nor is the domain of their commutator.

We denote the Bargmann space [2] by :

$$\mathbb{B} = \{ \phi : \mathcal{C} \rightarrow \mathcal{C} \text{ entire}; \int_{\mathcal{C}} |\phi(z)|^2 e^{-|z|^2} dx dy < \infty \} \quad (1.3)$$

The scalar product on  $\mathbb{B}$  is defined by

$$\langle \phi, \psi \rangle = \int_{\mathcal{C}} \phi(z) \overline{\psi(z)} e^{-|z|^2} dx dy \quad (1.4)$$

and the associated norm is denoted by  $\| \cdot \|$ .

$\mathbb{B}$  is closed in  $L_2(\mathcal{C}, d\mu(z))$  where the measure  $d\mu(z) = e^{-|z|^2} dx dy$  and it is closed related to  $L_2(\mathbb{R})$  by an unitary transform of  $L_2(\mathbb{R})$  onto  $\mathbb{B}$  given in [2] by the following integral transform

$$\phi(z) = \int_{\mathbb{R}} e^{-\frac{1}{2}(z^2+u^2)+\sqrt{2}uz} f(u) du \quad (1.5)$$

if  $f \in L_2(\mathbb{R})$  the integral converges absolutely.

In  $\mathbb{B}$  representation  $a$  and  $a^*$  are the operators of derivation and of multiplication and Gribov operator  $\mathbb{H} := \mathbb{H}_{\mu,\lambda}$  is defined on its maximal domain by

$$\left\{ \begin{array}{l} \mathbb{H}\phi(z) = i\lambda z\phi''(z) + (i\lambda z^2 + \mu z)\phi'(z) \\ \text{with maximal domain :} \\ D(\mathbb{H}_{max}) = \{\phi \in \mathbb{B} \text{ such that } \mathbb{H}\phi \in \mathbb{B}\} \end{array} \right. \quad (1.6)$$

Now we give some elementary properties of the operator  $\mathbb{H}$  and some remarks.

i)  $\mathbb{H}$  has the form  $p(z)\frac{d^2}{dz^2} + q(z)\frac{d}{dz}$  with  $p(z) = i\lambda z$  and  $q(z) = i\lambda z^2 + \mu z$  of degree 2 then it is of Heun operator type and its eigenvalue problem on  $\mathbb{B}$  ( $\mathbb{B} = \mathbb{B}_0 \oplus \{\text{constants}\}$  where  $\mathbb{B}_0 = \{\phi \in \mathbb{B} \text{ such that } \phi(0) = 0\}$  and zero is eigenvalue of  $\mathbb{H}$  without interest) does not satisfying the classical ordinary differential equation of the form:

$$\sigma(z)\phi''(z) + \tau(z)\phi'(z) + \alpha\phi(z) = 0$$

where  $\sigma(z)$  is a polynomial of degree at most two,  $\tau(z)$  is a polynomial of degree exactly one, and  $\alpha$  is a constant..

ii) An orthonormal basis of  $\mathbb{B}$  is given by  $e_k(z) = \frac{z^k}{\sqrt{k!}}; k = 0, 1, \dots$  (1.7)

iii) The action of the operators  $a, a^*$  and  $\mathbb{H}$  on the basis  $e_k(z) = \frac{z^k}{\sqrt{k!}}; k = 0, 1, \dots$ :

$$a(e_k) = \sqrt{k}e_{k-1} \text{ with the convention } e_{-1} = 0$$

$$a^*(e_k) = \sqrt{k+1}e_{k+1}$$

and

$$\mathbb{H}(e_k) = i\lambda(k-1)\sqrt{k}e_{k-1} + \mu ke_k + i\lambda k\sqrt{k+1}e_{k+1}$$

iv) Let  $\mathcal{P}$  be space of polynomials then it is dense in  $\mathbb{B}$ .

v) Let  $\mathbb{H}|_{\mathcal{P}}$  be the restriction of  $\mathbb{H}$  to polynomials space, we can define  $\mathbb{H}_{min}$  as the closure of operator  $\mathbb{H}|_{\mathcal{P}}$  in Bargmann space:

$$\left\{ \begin{array}{l} \mathbb{H}_{min}\phi = i\lambda z\phi'' + (i\lambda z^2 + \mu z)\phi' \\ \text{with domain :} \\ D(\mathbb{H}_{min}) = \{\phi \in \mathbb{B}; \exists p_n \in \mathcal{P}, \psi \in \mathbb{B}, p_n \rightarrow \phi \text{ and } \mathbb{H}p_n \rightarrow \psi\} \end{array} \right. \quad (1.8)$$

vi)  $\mathbb{H}_{min}^* = \mathbb{H}_{\mu, -\lambda}$  with domain  $D(\mathbb{H}_{min}^*) = D(\mathbb{H}_{max})$  (1.9)

where  $\mathbb{H}_{min}^*$  is the adjoint of  $\mathbb{H}_{min}$ .

Let  $l^2(\mathbb{N}) = \{(\phi_k)_{k=0}^\infty \in \mathbb{C} \text{ such that } \sum_{k=0}^\infty |\phi_k|^2 < +\infty\}$  with the inner product:

$$\langle (\phi_k)_{k=0}^\infty, (\psi_k)_{k=0}^\infty \rangle = \sum_{k=0}^\infty \phi_k \bar{\psi}_k \quad (1.10)$$

In the representation  $l^2(\mathbb{N})$  where the coefficients  $\phi_k$  define an entire function

$$\phi(z) = \sum_{k=0}^{\infty} \phi_k e_k(z) \text{ in Bargmann space we have}$$

$$\left\{ \begin{array}{l} (a\phi)_k = \sqrt{k}\phi_{k-1}, \phi_{-1} = 0 \\ \text{with domain :} \\ D(a) = \{(\phi)_k \in l^2(\mathbb{N}); \sum_{k=0}^{\infty} k |\phi_k|^2 < \infty\} \end{array} \right. \quad (1.11)$$

$$\left\{ \begin{array}{l} (a^*\phi)_k = \sqrt{k+1}\phi_{k+1} \\ \text{with domain :} \\ D(a^*) = \{(\phi)_k \in l^2(\mathbb{N}); \sum_{k=0}^{\infty} k |\phi_k|^2 < \infty\} \end{array} \right. \quad (1.12)$$

$$\left\{ \begin{array}{l} (\mathbb{H}\phi)_k = i\lambda(k-1)\sqrt{k}\phi_{k-1} + \mu k\phi_k + i\lambda k\sqrt{k+1}\phi_{k+1} \\ \text{with domain :} \\ D(\mathbb{H}) = \{\phi \in l^2(\mathbb{N}); H\phi \in l^2(\mathbb{N})\} \end{array} \right. \quad (1.13)$$

We define  $l_0^2(\mathbb{N}) = \{\phi = (\phi_k)_{k=0}^{\infty} \in l^2(\mathbb{N}); \phi_0 = 0\}$  and the operators  $A$  and  $A^*$  by

$$A(\phi) = a(\phi) \text{ with domain } D(A) = D(a) \cap l_0^2(\mathbb{N})$$

$$A^*(\phi) = a^*(\phi) \text{ with domain } D(A^*) = D(a^*) \cap l_0^2(\mathbb{N})$$

and now

$$\mathbb{H} = \mu A^* A + i\lambda A^*(A + A^*)A \text{ with domain } D(\mathbb{H}) = \{\phi \in l_0^2(\mathbb{N}); \mathbb{H}\phi \in l_0^2(\mathbb{N})\}$$

In the representation  $l_0^2(\mathbb{N})$  where the coefficients  $\phi_k$  define an entire function

$\phi(z) = \sum_{k=1}^{\infty} \phi_k e_k(z)$  in Bargmann space  $B_0$ , In [11], we have study the class of Jacobi-Gribov matrices with unbounded entries:

$$(H\phi)_k = \alpha_{k-1}\phi_{k-1} + \beta_k\phi_n + \alpha_k\phi_{k+1}, k \geq 2 \quad (1.14)$$

with the initial condition

$$(H\phi)_1 = \beta_1\phi_1 + \alpha_1\phi_2,$$

where



Now, we describe the contents of this paper, section by section. In Section 2, As in [13] or [14], we use a method of functional analysis which transform the problem of the zeros of Gribov-Intissar polynomials  $P_n^{(\mu,\lambda)}(z)$  to the equivalent problem of the eigenvalues of Gribov operator and for  $\mu \in \mathbb{C}$  and  $\lambda \in \mathbb{C}$ , we locate the position of the zeros of these polynomials. In Section 3, for  $\mu \in \mathbb{R}$  and  $\lambda \in \mathbb{R}$ , we give some common properties for Complex Symmetric tridiagonal matrices associated respectively to  $L_n^\alpha(z)$ ,  $P_n^\lambda(z)$ ,  $P_n^{(\alpha,\beta)}(z)$  and  $P_n^{(\mu,\lambda)}(z)$  and we show the existence of complex-valued function  $\xi(z)$  of bounded variation on  $\mathbb{R}$  such that the polynomials  $P_n^{(\mu,\lambda)}(z)$  are orthogonal with this weight  $\xi(z)$ .

For  $\mu = 0$ , we have applied some results of the excellent article of B. Simon [15](on the classical moment problem as a self-adjoint finite difference operator) to our Gribov operator, see the references [9] and [10] on the complete indeterminacy of this operator acting on Bargmann space.

For a treatment of the spectral analysis of certain Schrödinger operators associated to classical orthogonal polynomials, we refer the interested reader to the excellent article of Ismail-Koelink [12].

## 2. Localization of zeros of polynomials $P_n^{(\mu,\lambda)}$

Let  $\mathbb{B}_0 = \{\phi \in \mathbb{B}; \phi(0) = 0\}$  and  $e_k(z) = \frac{z^k}{\sqrt{k!}}; k = 1, \dots$  an orthonormal basis of  $\mathbb{B}_0$ .

Let  $\mathbb{V}$  be the shift operator defined by  $\mathbb{V}e_k = e_{k+1}, k = 1, \dots$  and its adjoint is the shift operator  $\mathbb{V}^*$  defined by  $\mathbb{V}^*e_k = e_{k-1}, k = 1, \dots$  and  $\mathbb{V}^*e_1 = 0$

Let also  $\mathbb{A}$  and  $\mathbb{B}$  be the diagonal operators defined by  $\mathbb{A}e_k = \alpha_k e_k, k = 1, \dots$  and  $\mathbb{B}e_k = \beta_k e_k, k = 1, \dots$  then

$$\mathbb{H} = \mathbb{A}\mathbb{V}^* + \mathbb{V}\mathbb{A} + \mathbb{B} \tag{2.1}$$

For  $1 \leq k \leq n$  the truncated Gribov operator is complex symmetric tridiagonal matrix defined by:

$$\mathbb{H}_n = \mathbb{A}\mathbb{V}^* + \mathbb{V}\mathbb{A} + \mathbb{B}; \mathbb{V}e_n = 0 \tag{2.2}$$

### Remark 2.1

Let  $\tilde{P}_n^{(\mu,\lambda)}(z) = \det(\mathbb{H}_n - zI)$  then we have:

$$\alpha_1 \alpha_2 \dots \alpha_n P_{n+1}^{(\mu,\lambda)}(z) = (-1)^n \det(\mathbb{H}_n - zI) = (-1)^n \tilde{P}_n^{(\mu,\lambda)}(z)$$

Let  $\mu = \mu_1 + i\mu_2; \mu_1, \mu_2 \in \mathbb{R}, \lambda = \lambda_1 + i\lambda_2; \lambda_1, \lambda_2 \in \mathbb{R}$  and  $\delta_k = k\sqrt{k+1}$

Let  $z_{k,n}(\mu, \lambda) = \mathcal{R}e z_{k,n} + i\mathcal{I}m z_{k,n}$  the zeros of the polynomials  $P_{n+1}^{\mu,\lambda}(z)$  satisfying three-term recurrence (1.16)

Then we have the following proposition:

**Proposition 2.2**

- i)  $|\operatorname{Re} z_{k,n}| \leq 2 |\lambda_2| \operatorname{Sup}_k \delta_k + |\mu_1| \operatorname{Sup}_k k = 2 |\lambda_2| \delta_n + n |\mu_1|$
- ii)  $|\operatorname{Im} z_{k,n}| \leq 2 |\lambda_1| \operatorname{Sup}_k \delta_k + |\mu_2| \operatorname{Sup}_k k = 2 |\lambda_1| \delta_n + n |\mu_2|$

**Proof**

$$\text{Let } \phi_k = \begin{pmatrix} \phi_{k,1} \\ \phi_{k,2} \\ \cdot \\ \cdot \\ \phi_{k,k} \\ \cdot \\ \cdot \\ \phi_{k,n} \end{pmatrix} \in \mathbb{C}^n; 1 \leq k \leq n \text{ and } \mathbb{H}_n \phi_k = z_{k,n} \phi_k; \quad \|\phi_k\| = 1$$

Then we have:

$$z_{k,n} = \langle \mathbb{H}_n \phi_k, \phi_k \rangle \tag{2.3}$$

As  $\mu \in \mathbb{C}$  and  $\lambda \in \mathbb{C}$ , we write  $\mu = \mu_1 + i\mu_2; (\mu_1, \mu_2) \in \mathbb{R}^2$  and  $\lambda = \lambda_1 + i\lambda_2; (\lambda_1, \lambda_2) \in \mathbb{R}^2$  for get:

$$\beta_n = \mu_1 n + i\mu_2 n \text{ and } \alpha_n = -\lambda_2 \delta_n + i\lambda_1 \delta_n; \delta_n = n\sqrt{n+1}$$

$$\text{Let } \mathbb{A}_1 e_k = -\lambda_2 \delta_k e_k, \mathbb{A}_2 e_k = -\lambda_1 \delta_k e_k, \mathbb{B}_1 e_k = \mu_1 k e_k \text{ and } \mathbb{B}_2 e_k = \mu_2 k e_k$$

Then

$$\mathbb{H}_n = \mathbb{H}_n^1 + i\mathbb{H}_n^2 \text{ where}$$

$$\mathbb{H}_n^1 = \mathbb{A}_1 \mathbb{V}^* + \mathbb{V} \mathbb{A}_1 + \mathbb{B}_1 \tag{2.4}$$

$$\mathbb{H}_n^2 = \mathbb{A}_2 \mathbb{V}^* + \mathbb{V} \mathbb{A}_2 + \mathbb{B}_2 \tag{2.5}$$

and the relation (2.3) takes the form:

$$z_{k,n} = \langle \mathbb{H}_n^1 \phi_k, \phi_k \rangle + i \langle \mathbb{H}_n^2 \phi_k, \phi_k \rangle \tag{2.6}$$

Since the operators  $\mathbb{H}_n^1$  and  $\mathbb{H}_n^2$  are selfadjoint, the inner products  $\langle \mathbb{H}_n^1 \phi_k, \phi_k \rangle$  and  $\langle \mathbb{H}_n^2 \phi_k, \phi_k \rangle$  are real and as a consequence,

$\operatorname{Re} z_{k,n} = \langle \mathbb{H}_n^1 \phi_k, \phi_k \rangle$  and  $\operatorname{Im} z_{k,n} = \langle \mathbb{H}_n^2 \phi_k, \phi_k \rangle$  which immediately result into the following inequalities:

- i)  $|\operatorname{Re} z_{k,n}| \leq 2 |\lambda_2| \operatorname{Sup}_k \delta_k + |\mu_1| \operatorname{Sup}_k k = 2 |\lambda_2| \delta_n + n |\mu_1|$
- ii)  $|\operatorname{Im} z_{k,n}| \leq 2 |\lambda_1| \operatorname{Sup}_k \delta_k + |\mu_2| \operatorname{Sup}_k k = 2 |\lambda_1| \delta_n + n |\mu_2|$

**Remark 2.3**

Let  $\mu \in \mathbb{R}$  and  $\lambda \in \mathbb{R}$  then, in Bargmann representation, we have the following properties on Gribov operator:

i) If  $n$  goes to infinity, the matrix  $\mathbb{H}_n^1$  generates a self-adjoint operator (the harmonic oscillator).

ii) If  $n$  goes to infinity, the matrix  $\mathbb{H}_n^2$  generates a symmetric operator (the cubic Heun Operator), but this operator is not self-adjoint because its spectrum is all  $\mathbb{C}$ . This specificity is attached to degree 2 of  $q(z) = i\lambda z^2 + \mu z$  in (1.6). See the reference [11] for a systematic study of this case.

iii) We note that  $\alpha_n$  is purely imaginary and  $\beta_n$  real, then  $\mathbb{H}_n^* = \mathbb{V}\mathbb{A}^* + \mathbb{A}^*\mathbb{V}^* + \mathbb{B}$  and thus  $\mathbb{H}_n$  is obviously not self-adjoint operator. However, it is complex symmetric operator and in the next section, we show that  $\mathbb{H}_n$  (respectively  $\mathbb{H}$ ) belongs to an interesting class of operators.

From the last proposition, we deduce the following corollary:

**Corollary 2.4**

If  $\mu \in \mathbb{R}$  and  $\lambda \in \mathbb{R}$ , we have:

- i)  $|\operatorname{Re} z_{k,n}| \leq |\mu| \quad \operatorname{Sup}_k k = n \quad |\mu|$
- ii)  $|\operatorname{Im} z_{k,n}| \leq 2 \quad |\lambda_1| \quad \operatorname{Sup}_k \delta_k = 2 \quad |\lambda| \quad \delta_n$

For the end of this section, we recall from [11] some properties established on the sequence  $\tilde{P}_n^{\mu,\lambda}(z)$  (under some conditions on the real parameters  $\mu$  and  $\lambda$ ).

Let  $\mu > 0$  then:

- i) Two successive polynomials  $\tilde{P}_{n-1}^{\mu,\lambda}(z)$  and  $\tilde{P}_n^{\mu,\lambda}(z)$  can not vanish simultaneously.
- ii) Let  $x \in \mathbb{R}$  such that  $\tilde{P}_{n-1}^{\mu,\lambda}(x) = 0$  then  $\tilde{P}_n^{\mu,\lambda}(x)\tilde{P}_{n-2}^{\mu,\lambda}(x) > 0 \quad \forall n$
- iii) If  $x < \mu$  then  $\tilde{P}_n^{\mu,\lambda}(x) > 0 \quad \forall n \geq 2$
- iv) If  $x > n\mu$  then  $(-1)^n \tilde{P}_n^{\mu,\lambda}(x) > 0 \quad \forall n \geq 2$
- v) Let  $x \in [\mu, n\mu]$ ;  $n \geq 2$ , if  $\tilde{P}_{n-1}^{\mu,\lambda}(x)$  and  $\tilde{P}_{n-2}^{\mu,\lambda}(x)$  have same sign on  $[\mu, n\mu]$  then on this interval  $\tilde{P}_k^{\mu,\lambda}(x)$  has this same sign  $\forall k \geq n - 2$
- vi) Let  $\lambda < \frac{\mu}{2\sqrt{2}}$  and  $x_{2k+1}$  the smallest zero of  $\tilde{P}_{2k+1}^{\mu,\lambda}(x)$  on  $[\mu, n\mu]$ ;  $n \geq 2$  then the sequence  $x_{2k+1}$  is in  $[\mu, x_2[$  and it is increasing, where  $x_2$  is the smallest zero of  $\tilde{P}_2^{\mu,\lambda}(x)$

From the above properties we have obtained in [11] the next fundamental result:

**Theorem 2.5** [11]

For  $\lambda < \frac{\mu}{2\sqrt{2}}$ , the Gribov operator  $\mathbb{H}$  has a least real eigenvalue.

### 3. On Complex Symmetric tridiagonal matrix associated to $P_n^{(\mu,\lambda)}$ and orthogonality

The Jacobi-Gribov matrix (1.15):

$$\left\{ \begin{array}{l} \mathbb{H} = (h_{j,k})_{j,k=1}^{\infty} \text{ with the elements :} \\ h_{kk} = \mu k = \beta_k, h_{k,k+1} = h_{k+1,k} = i\lambda k \sqrt{k+1} = \alpha_k; k = 1, 2, \dots \\ \text{and} \\ h_{jk} = 0 \text{ for } |j - k| > 1 \end{array} \right.$$

determines two linear operators in  $l_0^2(\mathbb{N})$  by the formal matrix product  $\mathbb{H}\phi$ . the first operator is defined in the linear manifold of vectors in  $l_0^2(\mathbb{N})$  with finite support related to the set  $\mathcal{P}$  of polynomials in Bargmann space, this operator is densely defined and closable. Let  $\mathbb{H}_{min}$  be its closure.

The second operator  $\mathbb{H}_{max}$  has the domain  $D(\mathbb{H}_{max}) = \{\phi = (\phi_n)_{n=0}^{\infty} \in l_0^2(\mathbb{N}); H\phi \in l_0^2(\mathbb{N})\}$

it was showed in [8] or in [11] (by another method) that  $\mathbb{H}_{max} = \mathbb{H}_{min}$

Let  $l_0^2(\mathbb{N})$  with its usual scalar product :

$$\langle \phi, \psi \rangle = \sum_{k=1}^{\infty} \phi_k \bar{\psi}_k$$

where

$$\phi = (\phi_1, \phi_2, \dots, \phi_k, \dots) \in l_0^2(\mathbb{N}) \text{ and } \psi = (\psi_1, \psi_2, \dots, \psi_k, \dots) \in l_0^2(\mathbb{N})$$

$$\text{Let } \mathbb{J} : l_0^2(\mathbb{N}) \rightarrow l_0^2(\mathbb{N}); \mathbb{J}(\phi) = \bar{\phi} = (\bar{\phi}_1, \bar{\phi}_2, \dots, \bar{\phi}_k, \dots) \text{ (conjugaison)}$$

The conjugation  $\mathbb{J}$  is an antilinear operator in  $l_0^2(\mathbb{N})$  such that

$$\mathbb{J}^2 \phi = \phi, \phi \in l_0^2(\mathbb{N}) \tag{3.1}$$

and

$$\langle \mathbb{J}\phi, \mathbb{J}\psi \rangle = \langle \phi, \psi \rangle ; \phi \in l_0^2(\mathbb{N}), \psi \in l_0^2(\mathbb{N}) \tag{3.2}$$

By using the definitions of complex symmetric matrix (complex symmetric operators) given by Garcia-Putinar in [5] or [6], we deduce obvious that  $\mathbb{H}_n$  is  $\mathbb{J}$ -symmetric :

$$[\mathbb{H}\phi, \psi]_{\mathbb{J}} = [\phi, \mathbb{H}\psi]_{\mathbb{J}} \tag{3.3}$$

where

$$[\phi, \psi]_{\mathbb{J}} := \langle \phi, \mathbb{J}\psi \rangle$$

Observe that (3.3) is equivalent to

$$\mathbb{J}\mathbb{H}\mathbb{J} = \mathbb{H}^* \quad (3.4)$$

Now, let  $\mathbb{T}$  be an unbounded operator on a separable infinite dimensional Banach space  $\mathbb{X}$ .

We define the following sets:

$$D(\mathbb{T}) = \{\phi \in \mathbb{X}; \mathbb{T}\phi \in \mathbb{X}\} \quad (3.5)$$

$$D(\mathbb{T}^\infty) = \bigcap_{n=0}^{\infty} D(\mathbb{T}^k) \quad (3.6)$$

**Definition 3.1**

A linear unbounded densely defined operator  $(\mathbb{T}, D(\mathbb{T}))$  on a Banach space  $\mathbb{X}$  is said to have a simple spectrum if the following conditions are met:

- 1)  $\mathbb{T}^k$  is closed for all positive integers  $k$ .
- 2) there exists an element  $\phi \in D(\mathbb{T}^\infty)$  ;  $\overline{Sp\{\phi, \mathbb{T}\phi, \mathbb{T}^2\phi, \dots\}} = \mathbb{X}$

**Lemma 3.2**

In Bargmann representation  $\mathbb{B}_0$ , the Gribov operator have a simple spectrum.

**Proof**

Let  $\{e_k\}_{k=1}^{\infty}(z) = \frac{z^k}{\sqrt{k!}}$  be an orthonormal basis of  $\mathbb{B}_0$  and  $\mathbb{H}$  be the Jacobi-Gribov matrix defined by (1.15):

$$\left\{ \begin{array}{l} \mathbb{H}e_1 = \beta_1 e_1 + \alpha_1 e_2 \\ \mathbb{H}e_k = \alpha_{k-1} e_{k-1} + \beta_k e_k + \alpha_k e_{k+1}; \quad k = 2, \dots \\ \text{where} \\ \beta_k = \mu k; \quad k = 1, 2, \dots \\ \text{and} \\ \alpha_k = i\lambda k \sqrt{k+1}; \quad k = 1, 2, \dots \end{array} \right. \quad (3.7)$$

$$\mathbb{S} = Sp\{\mathbb{H}^{k-1}e_1; \quad 0 \leq k \leq n\}, \quad n = 1, 2, \dots$$

We observe that  $e_1 \in \mathbb{S}$  and suppose that

$$e_m \in Sp\{\mathbb{H}^{k-1}e_1; \quad 1 \leq k \leq m\}, \quad 1 \leq m \leq n.$$

As  $\alpha_n \neq 0$  ;  $n = 1, 2, \dots$ , then by (3.7) we may write :

$$e_{n+1} = \frac{1}{\alpha_n} [\mathbb{H}e_n - \alpha_{n-1}e_{n-1} - \beta_n e_n]$$

then  $e_{n+1} \in Sp\{\mathbb{H}^{k-1}e_1; \quad 1 \leq k \leq n+1\}$ .

By induction we conclude that

$$e_m \in Sp\{ \mathbb{H}^{k-1}e_1; \quad 1 \leq k \leq m \}, \quad m = 1, 2, \dots \quad (3.8)$$

Therefore  $\bar{\mathbb{S}} = \mathbb{B}_0$ , i.e. the operator  $\mathbb{H}$  has a simple spectrum.

**Lemma 3.3**

Let  $\mathbb{G}$  the Gram matrix associated to Gribov operator  $\mathbb{H}$  for the system:

$$\{\hat{e}_1, \hat{e}_1, \dots, \hat{e}_n, \hat{e}_n^*\}; \quad n \in \mathbb{N}.$$

where

$$\hat{e}_n = H^{k-1}e_1 \text{ and } \hat{e}_n = (H^*)^{k-1}e_1; \quad k \geq 1 \quad (3.5)$$

Then

the vectors  $\{\hat{e}_1, \hat{e}_1, \dots, \hat{e}_n, \hat{e}_n^*\}$ , are linearly dependent and their Gram determinant is equal to zero.

**Proof**

$$\text{Set } \mathbb{S}_m := Sp\{ \mathbb{H}^{k-1}e_1; \quad 1 \leq k \leq m \}, \quad m = 1, 2, \dots$$

By (3.8), we see that  $e_1, e_2, \dots, e_m \in \mathbb{S}_m, m = 1, 2, \dots$ , and therefore  $\{e_k\}_{k=1}^m$  is an orthonormal basis in  $\mathbb{S}_m, m = 1, 2, \dots$

Since  $\mathbb{J}e_k = e_k, k = 1, 2, \dots, \mathbb{J}^2 = \mathbb{J}$  and  $\mathbb{J}\mathbb{H}\mathbb{J} = \mathbb{H}^*$  we have

$$\text{i) } \mathbb{J}\mathbb{S}_m \subset \mathbb{S}_m, m = 1, 2, \dots$$

and

$$\text{ii) } (\mathbb{H}^*)^{m-1}e_1 = (\mathbb{J}\mathbb{H}\mathbb{J})^{m-1}e_1 = \mathbb{J}\mathbb{H}^{m-1}\mathbb{J}e_1 = \mathbb{J}\mathbb{H}^{m-1}e_1 \in \mathbb{S}_m, m = 1, 2, \dots$$

Therefore vectors  $e_1, \mathbb{H}e_1, \dots, \mathbb{H}^{m-1}e_1, (\mathbb{H}^*)^{m-1}e_1$ , are linearly dependent and their Gram determinant is equal to zero.

**Theorem 3.4**

Let  $P_n^{\mu, \lambda}$  be the sequence of polynomials satisfying following three-term recurrence associated to gribov operator:

$$\begin{cases} P_0^{\mu, \lambda}(z) = 0 \\ P_1^{\mu, \lambda}(z) = 1 \\ \alpha_{n-1}P_{n-1}^{\mu, \lambda}(z) + \beta_n P_n^{\mu, \lambda}(z) + \alpha_n P_{n+1}^{\mu, \lambda}(z) = zP_n^{\mu, \lambda}(z); \quad n \geq 1 \end{cases}$$

where

$$\alpha_n = i\lambda n\sqrt{n+1}; n = 1, 2, \dots \text{ and } \beta_n = \mu n; n = 1, 2, \dots$$

then

there exists a complex-valued functions  $\xi(z)$  of bounded variation on  $\mathbb{R}$  such that

$$\int_{\mathbb{R}} P_m^{\mu,\lambda}(z)P_n^{\mu,\lambda}(z)d\xi(z) = \delta_{m,n} \quad m, n = 1, 2, \dots \quad (3.9)$$

**Proof**

Let the sequence of polynomials defined recursively by

$$\alpha_{n-1}P_{n-1}^{\mu,\lambda}(z) + \beta_n P_n^{\mu,\lambda}(z) + \alpha_n P_{n+1}^{\mu,\lambda}(z) = zP_n^{\mu,\lambda}(z); \quad n \geq 1 \quad (3.10)$$

where

$$P_1^{\mu,\lambda}(z) = 1 \text{ and } P_2^{\mu,\lambda}(z) = z - \beta_1$$

set

$$P_n^{\mu,\lambda}(z) = a_{n-1}z^{n-1} + a_{n-2}z^{n-2} + \dots + a_j z^j + \dots + a_0; n \geq 2 \text{ and } a_j \in \mathbb{C}, 0 \leq j \leq n-1$$

Comparing coefficients by  $z^n$  in (3.10), we get

$$a_n = \frac{1}{\alpha_n} a_{n-1}, n = 1, 2, \dots \text{ with } a_0 = -\beta_1$$

By induction, we see that

$$a_n = \left( \prod_{k=1}^n \alpha_k \right)^{-1} a_0$$

$$\text{Set } \mathbb{P}_n(z) = \prod_{k=1}^n \alpha_k P_n^{\mu,\lambda}(z)$$

Multiplying the both sides of (3.10) by  $\prod_{k=1}^n \alpha_k$ , we obtain:

$$\alpha_{n-1}^2 \mathbb{P}_{n-1} + \beta_n \mathbb{P}_n + \mathbb{P}_{n+1} = z \mathbb{P}_n$$

By theorem 6.4 in [4] there exists a complex-valued functions  $\xi(z)$  of bounded variation on  $\mathbb{R}$  such that

$$\int_{\mathbb{R}} \mathbb{P}_m(z)\mathbb{P}_n(z)d\xi(z) = \left( \prod_{k=1}^n n\alpha_k \right)^2 \delta_{m,n} \quad m, n = 1, 2, \dots$$

Therefore we get

$$\int_{\mathbb{R}} P_m^{\mu,\lambda}(z)P_n^{\mu,\lambda}(z)d\xi(z) = \delta_{m,n} \quad m, n = 1, 2, \dots$$

**Remark 3.5**

In Bargmann space and for  $\mu > 0$ , it was showed by Aimar and al. in [1], that the inverse of non-self-adjoint Gribov operator associated to above matrices belongs  $\mathcal{C}_{1+\epsilon}$  for all  $\epsilon > 0$  ( $\mathcal{C}_p; p > 0$  is the class of Carleman operators). By applying the results of this article, we show (in another paper) that more precisely, this inverse is not in  $\mathcal{C}_1$  where  $\mathcal{C}_1$  is the class of nuclear operators or trace operators by showing under some conditions on the parameters  $\mu$  and  $\lambda$  that  $z_{k,n}(\mu, \lambda) \rightarrow k\mu$  as  $n \rightarrow +\infty$ .

**References**

- [1] **M.T. Aimar, A. Intissar and J.M. Paoli**, Quelques nouvelles propriétés de régularité de l'opérateur de Gribov, *Comm. Math. Phys.* 172 (1995) 461-466.
- [2] **V. Bargmann, V.**, On a Hilbert space of analytic functions and an associated integral transform I, *Comm. Pure Appl. Math.*, (1962), 14: 187-214.
- [3] **K.G. Boreskov, A. B. Kaidalov, and O.V. Kancheli**, Strong interactions at high energies in the Reggeon approach, *Phys. Atomic Nuclei*, 2006, 69(10): 1765-1780.
- [4] **Chihara T.S.**, An introduction to orthogonal polynomials, *Mathematics and its Applications*, Vol. 13, Gordon and Breach Science Publishers, New York London Paris, 1978
- [5] **Garcia S.R.**, Putinar M., Complex symmetric operators and applications, *Trans. Amer. Math. Soc.* 358 (2006), 12851315.
- [6] **Garcia S.R.**, Putinar M., Complex symmetric operators and applications. II, *Trans. Amer. Math. Soc.* 359 (2007), 39133931.
- [7] **N. Gribov**, A reggeon diagram technique, *Soviet Phys. JETP* 26 (1968), no. 2, 414-423
- [8] **A. Intissar**, Etude spectrale d'une famille d'opérateurs non-symétriques intervenant dans la théorie des champs de Reggeons, *Comm. Math. Phys.*, 1987, 113(2): 263-297 (in French).
- [9] **A. Intissar**, Analyse de scattering d'un opérateur cubique de Heun dans l'espace de Bargmann, *Comm. Math.Phys.* 199, (1998), 243-256.
- [10] **A. Intissar**, On the complete indeterminacy and the chaoticity of generalized Heun's operator in Bargmann space, arXiv:1401.6560v1 [math.SP] 25 Jan (2014)
- [11] **A. Intissar**, Spectral Analysis of Non-self-adjoint Jacobi-Gribov Operator and Asymptotic Analysis of Its Generalized Eigenvectors, *Advances in Mathematics (China)*, Vol.43, No.x, (2014) doi: 10.11845/sxjz.2013117b
- [12] M.E.H Ismail, E. Koelink, Spectral Analysis of Certain Schrödinger Operators, arXiv:1205.0821v2 [math.CA] 15 Sep 2012
- [13] **C. G. Kokologiannaki, D. Rizos**, On the zeros of polynomials satisfying a three-term recurrence relation with complex coefficients, *Journal of Inequalities and Special Functions*, Volume 3, Issue 3 (2012), Pages 29-33

- [14] **E.N. Petropoulou**, On the Complex Zeros of Some Families of Orthogonal Polynomials, *Abstract and Applied Analysis*, Volume 2010, Article ID 263860, doi:10.1155/2010/263860
- [15] **B. Simon**, The Classical Moment Problem as a Self-Adjoint Finite Difference Operator, *Advances in Mathematics* 137, 82-203 (1998)