

The sine-Gordon equation on time scales

Jan L. Cieśliński*, Tomasz Nikiciuk†

Uniwersytet w Białymstoku, Wydział Fizyki,
ul. Lipowa 41, 15-424 Białystok, Poland

Kamil Waśkiewicz‡

Instytut Geofizyki PAN,
ul. Ks. Janusza 64, 01-452 Warszawa, Poland

Abstract

We formulate and discuss integrable analogue of the sine-Gordon equation on arbitrary time scales. This unification contains the sine-Gordon equation, discrete sine-Gordon equation and the Hirota equation (doubly discrete sine-Gordon equation) as special cases. We present the Lax pair, check compatibility conditions and construct the Darboux-Bäcklund transformation. Finally, we obtain a soliton solution on arbitrary time scale. The solution is expressed by the so called Cayley exponential function.

Keywords: integrable systems, solitons, time scales, discretization

PACS numbers: 02.30.Ik, 05.45.Yv

MSC 2010: 34N05, 35Q51, 39A12

*e-mail: janek@alpha.uwb.edu.pl

†e-mail: niki@alpha.uwb.edu.pl

‡e-mail: kamilwas@igf.edu.pl

1 Introduction

The sine-Gordon equation $\phi_{,xy} = \sin \phi$ is one of the classical soliton equations with numerous applications in many fields [2, 14, 23], from differential geometry [4, 7, 25, 28] to applied physics, including relativistic field theory [29], Josephson junctions [30, 33], propagation of deformations along DNA double helix [36], displacements in crystals [16], domain walls in ferroelectric and ferromagnetic materials [22], mechanical transmission lines [30, 32], and many others. In spite of the long history the sine-Gordon equation and its numerous extensions and generalizations still attracts attention of researchers [13, 15, 24, 31, 35]. In this paper we formulate and study an integrable extension of the sine-Gordon equation on arbitrary time scales, which includes, as special cases, the discrete case [27] and doubly discrete case (the Hirota equation) [20].

A time scale \mathbb{T} is just any (non-empty) closed subset of \mathbb{R} [18], including special cases like $\mathbb{T} = \mathbb{R}$, $\mathbb{T} = \mathbb{Z}$, $\mathbb{T} = h\mathbb{Z}$ and q -calculus. Time scales (or measure chains) were introduced in order to unify continuous and discrete calculus [19]. We recall several notions which will be used throughout the paper (here we assume $t \in \mathbb{T}$, note that in the rest of the paper we have two time scales with elements denoted respectively by x, y). The forward jump operator is defined as

$$t^\sigma := \inf\{s \in \mathbb{T} : s > t\}. \quad (1.1)$$

we denote also

$$\phi^\sigma(t) := \phi(t^\sigma). \quad (1.2)$$

In the case $\mathbb{T} = h\mathbb{Z}$ the forward jump is a shift. Graininess is defined by

$$\mu(t) := t^\sigma - t. \quad (1.3)$$

In the case $\mathbb{T} = h\mathbb{Z}$ we have $\mu = h$ (time step). In the case $\mathbb{T} = \mathbb{R}$, we have, obviously, $\mu = 0$. The delta derivative is defined by

$$f^\Delta(t) := \lim_{s \rightarrow t} \frac{f(t^\sigma) - f(s)}{t^\sigma - s}. \quad (1.4)$$

Note that

$$f + \mu f^\Delta = f^\sigma. \quad (1.5)$$

Delta exponential function $e_a(t)$, see [6, 18], satisfies initial value problem

$$e_a^\Delta = ae_a, \quad e_a(0) = 1, \quad (1.6)$$

where, in general, $a = a(t)$. We also have:

$$e_a^\sigma = (1 + a\mu)e_a \quad (1.7)$$

In some applications (e.g., in trigonometry) another definition of the exponential function is much more convenient [11, 12]. This is the Cayley exponential function which satisfies

$$E_a^\Delta = \frac{1}{2}a(E_a + E_a^\sigma), \quad E_a(0) = 1, \quad E_a^\sigma = \frac{1 + \frac{a\mu}{2}}{1 - \frac{a\mu}{2}}E_a. \quad (1.8)$$

In the continuous case ($\mathbb{T} = \mathbb{R}$) exponential functions become identical and for $a = \text{const}$ we have: $e_a(t) = E_a(t) = e^{at}$.

Dynamic equations on time scales are counterparts of differential equations. They unify continuous and discrete dynamical systems [1, 5, 21]. Some soliton equations on time scales were already studied within the Hamiltonian framework [3, 17, 34]. This paper is a continuation of [8], where the Darboux-Bäcklund transformation for a class of 2×2 linear problems was constructed. We will construct this transformation for the integrable analogue of the sine-Gordon equation on time scales and compute a soliton solution expressed by the Cayley exponential function.

2 Lax pair

A standard Lax pair for the sine-Gordon equation $\phi_{,xy} = \sin \phi$:

$$\begin{aligned} \Psi_{,x} &= \begin{pmatrix} i\zeta & -\frac{1}{2}\phi_{,x} \\ \frac{1}{2}\phi_{,x} & -i\zeta \end{pmatrix} \equiv i\zeta\sigma_3 - \frac{1}{2}i\phi_{,x}\sigma_2, \\ \Psi_{,y} &= \frac{1}{4i\zeta} \begin{pmatrix} \cos \phi & \sin \phi \\ \sin \phi & -\cos \phi \end{pmatrix} \equiv \frac{1}{4i\zeta} (\cos \phi\sigma_3 + \sin \phi\sigma_1), \end{aligned} \quad (2.1)$$

where $\sigma_1, \sigma_2, \sigma_3$ are Pauli matrices. We postulate the following time scale analogue for this Lax pair:

$$\begin{aligned} \Psi^{\Delta x} &= U\Psi, \quad U = i\zeta\sigma_3 - \frac{\sin \frac{\mu_x \phi^{\Delta x}}{2}}{\mu_x} i\sigma_2 - \frac{1 - \cos \frac{\mu_x \phi^{\Delta x}}{2}}{\mu_x} \\ \Psi^{\Delta y} &= V\Psi, \quad V = \frac{1}{4i\zeta} \left(\cos \frac{\phi^{\sigma_y} + \phi}{2} \sigma_3 + \sin \frac{\phi^{\sigma_y} + \phi}{2} \sigma_1 \right) \end{aligned} \quad (2.2)$$

We make important assumption throughout this paper:

$$\mu_x = \mu_x(x) , \quad \mu_y = \mu_y(y) , \quad (2.3)$$

i.e., we consider time scales $\mathbb{T} = \mathbb{T}_1 \times \mathbb{T}_2$ with graininess $\mu = (\mu_x(x), \mu_y(y))$. If $\mu \rightarrow (0, 0)$, then the Lax pair (2.2) becomes (2.1). It is convenient to denote (compare [4]):

$$\delta := \frac{1}{2} \phi^{\Delta_x} , \quad \alpha := \frac{1}{2} (\phi + \phi^{\sigma_y}) . \quad (2.4)$$

Then, the Lax pair (2.2) takes the form

$$\begin{aligned} \Psi^{\Delta_x} &= \left(-\zeta \mathbf{e}_3 + \frac{\sin \mu_x \delta}{\mu_x} \mathbf{e}_2 - \frac{1 - \cos \mu_x \delta}{\mu_x} \right) \Psi , \\ \Psi^{\Delta_y} &= \frac{1}{4\zeta} (\mathbf{e}_3 \cos \alpha + \mathbf{e}_1 \sin \alpha) \Psi , \end{aligned} \quad (2.5)$$

where $\mathbf{e}_1 = -i\sigma_1$, $\mathbf{e}_2 = -i\sigma_2$, $\mathbf{e}_3 = -i\sigma_3$. We have

$$\begin{aligned} \mathbf{e}_1 \mathbf{e}_2 &= -\mathbf{e}_2 \mathbf{e}_1 = \mathbf{e}_3, & \mathbf{e}_2 \mathbf{e}_3 &= -\mathbf{e}_3 \mathbf{e}_2 = \mathbf{e}_1, \\ \mathbf{e}_3 \mathbf{e}_1 &= -\mathbf{e}_1 \mathbf{e}_3 = \mathbf{e}_2, & \mathbf{e}_1^2 &= \mathbf{e}_2^2 = \mathbf{e}_3^2 = -1. \end{aligned} \quad (2.6)$$

Lemma 2.1

$$\alpha^{\Delta_x} = \delta^{\sigma_y} + \delta , \quad \mu_x (\delta + \delta^{\sigma_y}) = \alpha^{\sigma_x} - \alpha. \quad (2.7)$$

Proof: Using (2.4) we get

$$\alpha^{\Delta_x} = \frac{1}{2} (\phi^{\Delta_x} + \phi^{\sigma_2 \Delta_x}) = \delta + \delta^{\sigma_y} . \quad (2.8)$$

The second equality follows by (1.5). \square

Proposition 2.2 *Compatibility conditions, $U^{\Delta_y} - V^{\Delta_x} + U^{\sigma_y} V - V^{\sigma_x} U = 0$, for (2.2) are given by*

$$\frac{4}{\mu_x \mu_y} \sin \frac{\mu_x \mu_y \phi^{\Delta_x \Delta_y}}{4} = \sin \frac{\phi^{\sigma_x \sigma_y} + \phi^{\sigma_x} + \phi^{\sigma_y} + \phi}{4} , \quad (2.9)$$

where for $\mu_x = 0$ or $\mu_y = 0$ the left-hand side becomes $\phi^{\Delta_x \Delta_y}$ (as limits $\mu_x \rightarrow 0$ or $\mu_y \rightarrow 0$ suggest).

Proof: Compatibility conditions yield a system of four scalar equations given by the decomposition in the basis $1, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$. Coefficients by $\mathbf{e}_1, \mathbf{e}_3$ yield

$$\begin{aligned} (\sin \alpha)^{\Delta_x} &= \frac{\sin \mu_x \delta^{\sigma_y} \cos \alpha - (1 - \cos \mu_x \delta^{\sigma_y}) \sin \alpha + \cos \alpha^{\sigma_x} \sin \mu_x \delta + (1 - \cos \mu_x \delta) \sin \alpha^{\sigma_x}}{\mu_x}, \\ (\cos \alpha)^{\Delta_x} &= \frac{-\sin \mu_x \delta^{\sigma_y} \sin \alpha - (1 - \cos \mu_x \delta^{\sigma_y}) \cos \alpha - \sin \alpha^{\sigma_x} \sin \mu_x \delta + (1 - \cos \mu_x \delta) \cos \alpha^{\sigma_x}}{\mu_x}. \end{aligned} \quad (2.10)$$

If $\mu_x \neq 0$, then we use (1.5) to transform these equations into

$$\begin{aligned} \sin \alpha^{\sigma_x} \cos \mu_x \delta - \cos \alpha^{\sigma_x} \sin \mu_x \delta &= \sin \alpha \cos \mu_x \delta^{\sigma_y} + \cos \alpha \sin \mu_x \delta^{\sigma_y}, \\ \cos \alpha^{\sigma_x} \cos \mu_x \delta + \sin \alpha^{\sigma_x} \sin \mu_x \delta &= \cos \alpha \cos \mu_x \delta^{\sigma_y} - \sin \alpha \sin \mu_x \delta^{\sigma_y}. \end{aligned} \quad (2.11)$$

Using well known trigonometric identities, we get

$$\begin{aligned} \sin(\alpha^{\sigma_x} - \mu_x \delta) &= \sin(\mu_x \delta^{\sigma_y} + \alpha), \\ \cos(\alpha^{\sigma_x} - \mu_x \delta) &= \cos(\mu_x \delta^{\sigma_y} + \alpha), \end{aligned} \quad (2.12)$$

which is identically satisfied by (1.5) and Lemma 2.1. If $\mu_x = 0$, then (2.10) reduce to

$$\alpha^{\Delta_x} \cos \alpha = (\delta^{\sigma_y} + \delta) \cos \alpha, \quad \alpha^{\Delta_x} \sin \alpha = (\delta^{\sigma_y} + \delta) \sin \alpha, \quad (2.13)$$

which is satisfied by Lemma 2.1, as well.

Dealing with coefficients by 1 and \mathbf{e}_2 we will consider three cases separately. In the first case we assume $\mu_x \neq 0, \mu_y \neq 0$. Then coefficients by 1 and \mathbf{e}_2 yield

$$\begin{aligned} \cos \mu_x \delta^{\sigma_y} &= \cos \mu_x \delta + \frac{1}{4} \mu_x \mu_y (\cos \alpha^{\sigma_x} - \cos \alpha), \\ \sin \mu_x \delta^{\sigma_y} &= \sin \mu_x \delta + \frac{1}{4} \mu_x \mu_y (\sin \alpha^{\sigma_x} + \sin \alpha), \end{aligned} \quad (2.14)$$

Hence, using trigonometric identities, we get

$$\begin{aligned} \sin \frac{\mu_x(\delta^{\sigma_y} + \delta)}{2} \sin \frac{\mu_x(\delta^{\sigma_y} - \delta)}{2} &= \frac{1}{4} \mu_x \mu_y \sin \frac{\alpha^{\sigma_x} - \alpha}{2} \sin \frac{\alpha^{\sigma_x} + \alpha}{2}, \\ \cos \frac{\mu_x(\delta^{\sigma_y} + \delta)}{2} \sin \frac{\mu_x(\delta^{\sigma_y} - \delta)}{2} &= \frac{1}{4} \mu_x \mu_y \cos \frac{\alpha^{\sigma_x} - \alpha}{2} \sin \frac{\alpha^{\sigma_x} + \alpha}{2}. \end{aligned} \quad (2.15)$$

By virtue of Lemma 2.1 these equations reduce to the following single equation

$$\sin \frac{\mu_x(\delta^{\sigma_y} - \delta)}{2} = \frac{1}{4} \mu_x \mu_y \sin \frac{\alpha^{\sigma_x} + \alpha}{2}. \quad (2.16)$$

Using (2.4) and taking into account that $\delta^{\sigma_y} = \delta + \mu_y \delta^{\Delta_y}$ we get (2.9).

In the second case we assume $\mu_x \neq 0$ and $\mu_y = 0$ (hence, in particular, $\delta^{\sigma_y} = \delta$ and $\alpha = \phi$). Then coefficients by 1 and \mathbf{e}_2 yield

$$\begin{aligned} -\delta^{\Delta_y} \sin \mu_x \delta &= \frac{1}{4}(\cos \alpha^{\sigma_x} - \cos \alpha), \\ \delta^{\Delta_y} \cos \mu_x \delta &= \frac{1}{4}(\sin \alpha^{\sigma_x} + \sin \alpha), \end{aligned} \quad (2.17)$$

In this case Lemma 2.1 implies $\mu_x \delta = \frac{1}{2}(\alpha^{\sigma_x} - \alpha)$. Therefore

$$\begin{aligned} \sin \frac{\alpha^{\sigma_x} - \alpha}{2} (\delta^{\Delta_y} - \frac{1}{2} \sin \frac{\alpha^{\sigma_x} + \alpha}{2}) &= 0, \\ \cos \frac{\alpha^{\sigma_x} - \alpha}{2} (\delta^{\Delta_y} - \frac{1}{2} \sin \frac{\alpha^{\sigma_x} + \alpha}{2}) &= 0, \end{aligned} \quad (2.18)$$

hence

$$\delta^{\Delta_y} = \frac{1}{2} \sin \frac{\alpha^{\sigma_x} + \alpha}{2}, \quad (2.19)$$

and, substituting $\delta = \frac{1}{2}\alpha^{\Delta_x}$, $\alpha = \phi$, we obtain

$$\alpha^{\Delta_x \Delta_y} = \sin \frac{\alpha^{\sigma_x} + \alpha}{2} \quad \longrightarrow \quad \phi^{\Delta_x \Delta_y} = \sin \frac{\phi^{\sigma_x} + \phi}{2}, \quad (2.20)$$

which can be considered as a special case of (2.9).

In the third case we assume $\mu_x = 0$. Now the Lax pair essentially simplifies

$$\Psi^{\Delta_x} = (i\zeta\sigma_3 - i\sigma_2\delta)\Psi, \quad \Psi^{\Delta_y} = \frac{1}{4i\zeta}(\sigma_3 \cos \alpha + \sigma_1 \sin \alpha)\Psi \quad (2.21)$$

Compatibility conditions yield three equations (the term proportional to the unit matrix vanishes):

$$\begin{aligned} \delta^{\Delta_y} &= \frac{1}{2} \sin \alpha, \\ (-\alpha^{\Delta_x} + \delta^{\sigma_y} + \delta) \cos \alpha &= 0, \\ (\alpha^{\Delta_x} - \delta^{\sigma_y} - \delta) \sin \alpha &= 0, \end{aligned} \quad (2.22)$$

The last two equations are identically satisfied by virtue of Lemma 2.1. Then, using (2.4), we obtain

$$\phi^{\Delta_x \Delta_y} = \sin \frac{\phi^{\sigma_y} + \phi}{2}, \quad (2.23)$$

which can be considered as a special case of (2.9). \square

Corollary 2.3 *The equation (2.9) will be called the sine-Gordon equation on time scales. For $\mu_x \neq 0, \mu_y \neq 0$ it coincides with the Hirota equation (doubly discrete sine-Gordon equation) [20, 26] and for $\mu_y = 0$ it coincides with the discrete sine-Gordon equation [27].*

The equation (2.9) can be rewritten as

$$\phi^{\Delta_1 \Delta_2} = F \left(\frac{1}{4} \mu_x \mu_y \sin \langle \langle \phi \rangle \rangle \right) \sin \langle \langle \phi \rangle \rangle, \quad (2.24)$$

where

$$\langle \langle \phi \rangle \rangle = \frac{1}{4} (\phi^{\sigma_x \sigma_y} + \phi^{\sigma_x} + \phi^{\sigma_y} + \phi), \quad (2.25)$$

and $F(x) := x^{-1} \arcsin x$, $F(0) = 1$.

3 Darboux-Bäcklund transformation

The Darboux-Bäcklund transformation on time scales was formulated and derived in [8]. This is a gauge-like transformation preserving the structure of the Lax pair, see [9]. Let $\tilde{\Psi} = D\Psi$ (D is called “Darboux matrix”). Then matrices U, V of the Lax pair (2.2) transform as follows

$$\begin{aligned} \tilde{U} &= D^{\Delta_1} D^{-1} + D^{\sigma_1} U D^{-1}, \\ \tilde{V} &= D^{\Delta_2} D^{-1} + D^{\sigma_2} V D^{-1}. \end{aligned} \quad (3.1)$$

In this paper we confine ourselves to the binary Darboux matrix which adds (as a kind of nonlinear superposition) a soliton onto a given background. Fortunately, the Darboux matrix has exactly the same form in all cases (continuous, discrete and time scales) [8]:

$$D = \frac{\lambda - \kappa_1 A}{\lambda - i\kappa_1}. \quad (3.2)$$

where $A = i(I - 2P)$ and P is a Hermitean projector ($P^2 = P$, $P^\dagger = P$) defined by

$$\ker P = \Psi(i\kappa_1) \vec{c}_1, \quad \text{Im } P = \Psi(-i\kappa_1) \vec{c}_1^\perp, \quad (3.3)$$

where \vec{c}_1 is a constant vector. Moreover, the Lax pair (2.2) satisfies:

$$U(-\zeta) = \mathbf{e}_2^{-1}U(\zeta)\mathbf{e}_2, \quad V(-\zeta) = \mathbf{e}_2^{-1}V(\zeta)\mathbf{e}_2, \quad (3.4)$$

which implies the following constraint on the projector P [8]:

$$\mathbf{e}_2 P = (I - P)\mathbf{e}_2. \quad (3.5)$$

Hence [8]:

$$P = \frac{1}{2}(I + \sigma_1 \sin \theta + \sigma_3 \cos \theta) = \frac{1}{2} \begin{pmatrix} 1 + \cos \theta & \sin \theta \\ \sin \theta & 1 - \cos \theta \end{pmatrix}. \quad (3.6)$$

Taking into account well known trigonometric identities, we can represent P as

$$P = \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} \end{pmatrix} \left(\cos \frac{\theta}{2}, \sin \frac{\theta}{2} \right). \quad (3.7)$$

Likewise, we have

$$I - P = \frac{1}{2} \begin{pmatrix} 1 - \cos \theta & -\sin \theta \\ -\sin \theta & 1 + \cos \theta \end{pmatrix}, \quad (3.8)$$

and

$$I - P = \begin{pmatrix} -\sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} \end{pmatrix} \left(-\sin \frac{\theta}{2}, \cos \frac{\theta}{2} \right). \quad (3.9)$$

Therefore,

$$\text{im} P = f \begin{pmatrix} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} \end{pmatrix}, \quad \text{ker} P = g \begin{pmatrix} -\sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} \end{pmatrix}, \quad (3.10)$$

where f, g are arbitrary functions. The function θ , parameterizing the projector, can be uniquely derived from (3.3), provided that $\Psi(\zeta)$ is known. Alternatively, we can solve dynamic equations for θ .

Proposition 3.1 *Dynamic equations for θ parameterizing the projector P read*

$$\begin{aligned}
\frac{1}{\mu_x} \sin\left(\mu_x \delta - \frac{\theta^{\sigma_x} - \theta}{2}\right) &= \kappa_1 \sin\frac{\theta^{\sigma_x} + \theta}{2}, & (\text{for } \mu_x \neq 0), \\
\frac{1}{\mu_y} \sin\frac{\theta^{\sigma_y} - \theta}{2} &= \frac{1}{4\kappa_1} \sin\left(\alpha - \frac{\theta + \theta^{\sigma_y}}{2}\right) & (\text{for } \mu_y \neq 0), \\
\delta - \frac{1}{2}\theta_{,x} &= \kappa_1 \sin\frac{\theta^{\sigma_x} + \theta}{2}, & (\text{for } \mu_x = 0), \\
\theta_{,y} &= \frac{1}{2\kappa_1} \sin(\alpha - \theta), & (\text{for } \mu_y = 0).
\end{aligned} \tag{3.11}$$

Proof: We confine ourselves to the case $\mu_x \neq 0, \mu_y \neq 0$. Other cases can be considered in exactly the same way (all intermediate steps can be obtained as a formal limit). From (3.3) it follows that

$$(\ker P)^{\Delta_x} = U(i\kappa_1)\ker P, \quad (\ker P)^{\Delta_y} = V(i\kappa_1)\text{im}P. \tag{3.12}$$

Analogical equations for $\text{im}P$ can be omitted because in this case $\text{im}P$ is uniquely determined by $\ker P$. Indeed, $\text{im}P = (\ker P)^\perp$ and $\text{im}P = \mathbf{e}_2\ker P$. Note that $U(i\kappa_1)$ and $V(i\kappa_1)$ are real matrices, provided that all coefficients and fields are real:

$$\begin{aligned}
U(i\kappa_1) &= \kappa_1 \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{\sin\mu_x\delta}{\mu_x} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} - \frac{1-\cos\mu_x\delta}{\mu_x} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \\
V(i\kappa_1) &= -\frac{1}{4\kappa_1} \begin{pmatrix} \cos\alpha & \sin\alpha \\ \sin\alpha & -\cos\alpha \end{pmatrix}.
\end{aligned} \tag{3.13}$$

Substituting (3.10) into (3.12) and dividing by, respectively, f and g , we obtain:

$$\begin{aligned}
(\sin\frac{\theta}{2})^{\Delta_x} + \frac{f^{\Delta_x}}{f} \sin\frac{\theta^{\sigma_x}}{2} &= -\kappa_1 \sin\frac{\theta}{2} + \frac{\sin\mu_x\delta}{\mu_x} \cos\frac{\theta}{2} - \frac{1-\cos\mu_x\delta}{\mu_x} \sin\frac{\theta}{2}, \\
(\cos\frac{\theta}{2})^{\Delta_x} + \frac{f^{\Delta_x}}{f} \cos\frac{\theta^{\sigma_x}}{2} &= \kappa_1 \cos\frac{\theta}{2} - \frac{\sin\mu_x\delta}{\mu_x} \sin\frac{\theta}{2} - \frac{1-\cos\mu_x\delta}{\mu_x} \cos\frac{\theta}{2}, \\
(\sin\frac{\theta}{2})^{\Delta_y} + \frac{g^{\Delta_y}}{g} \sin\frac{\theta^{\sigma_y}}{2} &= -\frac{1}{4\kappa_1} \left(\cos\alpha \cos\frac{\theta}{2} - \sin\alpha \cos\frac{\theta}{2}\right), \\
(\cos\frac{\theta}{2})^{\Delta_y} + \frac{g^{\Delta_y}}{g} \cos\frac{\theta^{\sigma_y}}{2} &= \frac{1}{4\kappa_1} \left(\sin\alpha \sin\frac{\theta}{2} + \cos\alpha \cos\frac{\theta}{2}\right),
\end{aligned} \tag{3.14}$$

Eliminating f and g from (3.14) we get two equations

$$\begin{aligned}
\frac{1}{\mu_x} \sin\frac{\theta^{\sigma_x} - \theta}{2} &= -\kappa_1 \sin\frac{\theta^{\sigma_x} + \theta}{2} + \frac{\sin\mu_x\delta}{\mu_x} \cos\frac{\theta^{\sigma_x} - \theta}{2} + \frac{1-\cos\mu_x\delta}{\mu_x} \sin\frac{\theta^{\sigma_x} - \theta}{2}, \\
\frac{1}{\mu_y} \sin\frac{\theta - \theta^{\sigma_y}}{2} &= \frac{1}{4\kappa_1} \sin\left(\frac{\theta + \theta^{\sigma_y}}{2} - \alpha\right)
\end{aligned}$$

and, after elementary calculation, we get (3.11). \square

Proposition 3.2 ([8]) *If D is given by (3.2), (3.3) and the Lax pair is of the form*

$$U = u_0 + \zeta u_1, \quad V = v_0 + \frac{1}{\zeta} v_1, \quad (3.15)$$

where u_0, u_1, v_0, v_1 are linear combinations of $I, \mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ with real coefficients and (3.4) is satisfied, then the Darboux-Bäcklund transformation is given by

$$\begin{aligned} \tilde{u}_1 &= u_1, & \tilde{u}_0 &= u_0 + \kappa_1(u_1 A - A^{\sigma_x} u_1), \\ \tilde{v}_0 &= v_0, & \tilde{v}_1 &= -A^{\sigma_y} v_1 A^{-1}, \end{aligned} \quad (3.16)$$

where $A = i(I - 2P) = \mathbf{e}_3 \cos \theta + \mathbf{e}_1 \sin \theta$.

Proposition 3.2 can be applied directly to the Lax pair (2.2). Two equations are trivially satisfied ($\tilde{u}_1 = u_1 = -\mathbf{e}_3$ and $\tilde{v}_0 = v_0 = 0$). The equation for \tilde{u}_0 splits into two equations:

$$\begin{aligned} \frac{\sin \mu_x \tilde{\delta}}{\mu_x} &= \frac{\sin \mu_x \delta}{\mu_x} - \kappa_1(\sin \theta^{\sigma_x} + \sin \theta), \\ \frac{1 - \cos \mu_x \tilde{\delta}}{\mu_x} &= \frac{1 - \cos \mu_x \delta}{\mu_x} + \kappa_1(\cos \theta^{\sigma_x} - \cos \theta), \end{aligned} \quad (3.17)$$

for $\mu_x \neq 0$. In the case $\mu_x = 0$ we have, instead,

$$\begin{aligned} \tilde{\delta} &= \delta - 2\kappa_1 \sin \theta, \\ \frac{1}{2} \tilde{\delta}^2 &= \frac{1}{2} \delta^2 - \kappa_1 \theta_{,x} \sin \theta. \end{aligned} \quad (3.18)$$

The last equation of (3.16) will be simplified using an elegant approach of Clifford numbers instead of cumbersome straightforward calculations.

Lemma 3.3 *For any scalar function $\beta = \beta(x, y)$ we have*

$$\begin{aligned} \cos \beta + \mathbf{e}_1 \mathbf{e}_3 \sin \beta &= \exp(\beta \mathbf{e}_1 \mathbf{e}_3), \\ \mathbf{e}_3 \cos \beta + \mathbf{e}_1 \sin \beta &= \mathbf{e}_3 (\cos \beta + \mathbf{e}_1 \mathbf{e}_3 \sin \beta) = \mathbf{e}_3 \exp(\beta \mathbf{e}_1 \mathbf{e}_3), \\ \mathbf{e}_3 \exp(\beta \mathbf{e}_1 \mathbf{e}_3) &= \exp(-\beta \mathbf{e}_1 \mathbf{e}_3) \mathbf{e}_3 \end{aligned} \quad (3.19)$$

Proof: We use (2.6) and observe that $(\mathbf{e}_1\mathbf{e}_3)^2 = -1$. The first equation can be proved using power series, like the famous formula $\cos \beta + i \sin \beta = \exp(i\beta)$. \square

Taking into account properties listed in Lemma 3.3 we can easily transform the last equation of (3.16) into

$$e_3 \exp(\tilde{\alpha}\mathbf{e}_1\mathbf{e}_3) = -e_3^3 \exp(\mathbf{e}_1\mathbf{e}_3\theta^{\sigma_x}) \exp(-\mathbf{e}_1\mathbf{e}_3\alpha) \exp(\mathbf{e}_1\mathbf{e}_3\theta). \quad (3.20)$$

Because $\mathbf{e}_3^3 = -\mathbf{e}_3$ and exponents of all exponential functions commute, we have

$$\mathbf{e}_3 \exp(\tilde{\alpha}\mathbf{e}_1\mathbf{e}_3) = \exp \mathbf{e}_1\mathbf{e}_3(\theta^{\sigma_x} + \theta - \alpha), \quad (3.21)$$

hence we finally get: $\tilde{\alpha} = -\alpha + \theta^{\sigma_x} + \theta$.

Proposition 3.4 *The Darboux-Bäcklund transformation (3.16) for the Lax pair (2.2) is equivalent to the system:*

$$\begin{aligned} \tilde{\alpha} &= -\alpha + \theta^{\sigma_y} + \theta, \\ \tilde{\delta} + \delta &= \theta^{\Delta_x}, \end{aligned} \quad (3.22)$$

which can be reduced to one equation for ϕ :

$$\tilde{\phi} = 2\theta - \phi, \quad (3.23)$$

where θ is computed either from (3.3) or from Proposition 3.1.

Proof: The first equation of (3.22) was already obtained from (3.21). Considering the second equation, we first assume $\mu_x \neq 0$. The system (3.17) can be rewritten as

$$\begin{aligned} \frac{1}{\mu_x} \sin \frac{\mu_x(\tilde{\delta}-\delta)}{2} \cos \frac{\mu_x(\tilde{\delta}+\delta)}{2} &= -\kappa_1 \sin \frac{\theta^{\sigma_x}+\theta}{2} \cos \frac{\theta^{\sigma_x}-\theta}{2}, \\ \frac{1}{\mu_x} \sin \frac{\mu_x(\tilde{\delta}-\delta)}{2} \sin \frac{\mu_x(\tilde{\delta}+\delta)}{2} &= -\kappa_1 \sin \frac{\theta^{\sigma_x}+\theta}{2} \sin \frac{\theta^{\sigma_x}-\theta}{2}, \end{aligned} \quad (3.24)$$

Dividing these equations side by side, we get

$$\frac{1}{\mu_x} \tan \frac{\mu_x(\tilde{\delta}+\delta)}{2} = \frac{1}{\mu_x} \tan \frac{\theta^{\sigma_x}-\theta}{2}, \quad (3.25)$$

which implies the first equation of (3.22). The remaining equation,

$$\frac{1}{\mu_x} \sin \frac{\mu_x(\tilde{\delta}-\delta)}{2} = -\kappa_1 \sin \frac{\theta^{\sigma_x}+\theta}{2}, \quad (3.26)$$

after substituting $\tilde{\delta} = \theta^{\Delta_x} - \delta$, reduces to the first equation of (3.11). The case $\mu_x = 0$ is simpler. Substituting $\tilde{\delta} - \delta$ from the first equation of (3.18) into the second equation of (3.18) we get the second equation of (3.22).

Finally, substituting (2.4) into (3.22), we get

$$\begin{aligned} (\tilde{\phi} + \phi - 2\theta)^{\Delta_x} &= 0, \\ (\tilde{\phi} + \phi - 2\theta)^{\sigma_y} &= -(\tilde{\phi} + \phi - 2\theta). \end{aligned} \tag{3.27}$$

Hence, $\tilde{\phi} + \phi - 2\theta = \chi$, where χ is a function of y such that $\chi^{\sigma_y} = -\chi$. Without loss of generality we can take $\chi = 0$ because transformation $\phi \rightarrow \phi + \chi$ does not change the Lax pair (2.2). \square

4 Solitons on an arbitrary time scale

Pure soliton solutions are obtained, as usual, by applying the Darboux-Bäcklund transformation to the simplest (“vacuum”) background $\phi = 0$. In this case the Lax pair reduces to

$$\Psi^{\Delta_x} = i\zeta\sigma_3\Psi, \quad \Psi^{\Delta_y} = \frac{1}{4i\zeta}\sigma_3\Psi, \tag{4.1}$$

which can be easily solved (assuming initial condition $\Psi(0, 0) = I$):

$$\Psi = \begin{pmatrix} e_{i\zeta}(x)e_{\frac{1}{4i\zeta}}(y) & 0 \\ 0 & e_{-i\zeta}(x)e_{-\frac{1}{4i\zeta}}(y) \end{pmatrix}, \tag{4.2}$$

where e denotes the delta exponential function (1.6). From the previous section we know that $\tilde{\phi} = 2\theta - \phi$. Therefore one-soliton solution is given by

$$\phi_{sol} = 2\theta, \tag{4.3}$$

where θ is associated with the simplified linear system (4.2).

Proposition 4.1 *One-soliton solution to the sine-Gordon equation on an arbitrary time scale (equation (2.9)) is given by*

$$\phi_{sol} = 4 \arctan \left(c E_{-2\kappa_1}(x) E_{-\frac{1}{2\kappa_1}}(y) \right), \tag{4.4}$$

where E denotes the Cayley-exponential function (1.8) and c, κ_1 are real parameters.

Proof: In the case $\alpha = \delta = 0$ equations (3.11) read

$$\begin{aligned}
\frac{1}{\mu_x} \sin \frac{\theta^{\sigma_x} - \theta}{2} &= -\kappa_1 \sin \frac{\theta^{\sigma_x} + \theta}{2}, & (\text{for } \mu_x \neq 0) \\
\frac{1}{\mu_y} \sin \frac{\theta^{\sigma_y} - \theta}{2} &= -\frac{1}{4\kappa_1} \sin \frac{\theta^{\sigma_y} + \theta}{2}, & (\text{for } \mu_y \neq 0) \\
\theta_{,x} &= -2\kappa_1 \sin \theta, & (\text{for } \mu_x = 0), \\
\theta_{,y} &= -\frac{1}{2\kappa_1} \sin \theta & (\text{for } \mu_y = 0).
\end{aligned} \tag{4.5}$$

Equations (4.5) can be easily transformed into:

$$\begin{aligned}
\tan \frac{\theta^{\sigma_x}}{2} &= \frac{1 - \mu_x \kappa_1}{1 + \mu_x \kappa_1} \tan \frac{\theta}{2}, & (\text{for } \mu_x \neq 0) \\
\tan \frac{\theta^{\sigma_y}}{2} &= \frac{1 - \frac{\mu_y}{4\kappa_1}}{1 + \frac{\mu_y}{4\kappa_1}} \tan \frac{\theta}{2}, & (\text{for } \mu_y \neq 0) \\
(\tan \frac{\theta}{2})_{,x} &= -2\kappa_1 \tan \frac{\theta}{2}, & (\text{for } \mu_x = 0), \\
(\tan \frac{\theta}{2})_{,y} &= -\frac{1}{2\kappa_1} \tan \frac{\theta}{2}, & (\text{for } \mu_y = 0)
\end{aligned} \tag{4.6}$$

which yields (4.4) by virtue of (1.8) (the continuous case agrees with this solution, as well). Another method to get (4.4), even more straightforward, consists in using (3.3) and (3.10). \square

In the continuous and discrete case the formula (4.4) reduces to known expressions:

$$\begin{aligned}
\phi_{sol-cont} &= 4 \arctan \exp(-2\kappa_1 x - \frac{1}{2\kappa_1} y + c_0), \\
\phi_{sol-discr} &= 4 \arctan \left(c \left(\frac{1 - \kappa_1 h_x}{1 + \kappa_1 h_x} \right)^{\frac{x}{h_x}} \left(\frac{1 - \frac{h_y}{4\kappa_1}}{1 + \frac{h_y}{4\kappa_1}} \right)^{\frac{y}{h_y}} \right),
\end{aligned} \tag{4.7}$$

where $h_x = \mu_x$ and $h_y = \mu_y$. The discrete soliton has, in fact, a similar shape as the smooth soliton. Indeed,

$$\phi_{sol-discr} = 4 \arctan \exp(-2\tilde{\kappa}_1 x - \frac{1}{2\hat{\kappa}_1} y + c_0), \tag{4.8}$$

where $c = \exp c_0$ and

$$\tilde{\kappa}_1 = \frac{1}{2h_x} \ln \left(\frac{1 + \kappa_1 h_x}{1 - \kappa_1 h_x} \right), \quad \hat{\kappa}_1^{-1} = \frac{1}{2h_y} \ln \left(\frac{1 + \frac{h_y}{4\kappa_1}}{1 - \frac{h_y}{4\kappa_1}} \right). \tag{4.9}$$

Therefore, on two most popular time scales ($\mathbb{T} = \mathbb{R}$, $\mathbb{T} = h\mathbb{Z}$) solitons are solitary waves. In the discrete case velocity and shape depend on h_x and h_y . However, in general, one-soliton solution (4.4) is not a solitary wave.

5 Summary

We formulated and studied an integrable analogue of the sine-Gordon equation on an arbitrary time scale. In particular, we provided parallel computations for the discrete and continuous case. The Darboux-Bäcklund transformation has the same form on any time scale. Soliton solutions of the sine-Gordon equation on time scales are expressed by the Cayley exponential function. Therefore, like in the continuous case, soliton solutions of the nonlinear equation are expressed in terms of exponential functions which were introduced as solutions to linear equations.

Acknowledgement. The first author (J.L.C.) is partly supported by the National Science Centre (NCN) grant no. 2011/01/B/ST1/05137.

References

- [1] C.D.Ahlbrandt, C.Morian: “Partial differential equations on time scales”, *J. Comp. Appl. Math.* 141 (2002) 35-55.
- [2] A.Barone, F. Esposito, C.J.Magee, A.C.Scott: “Theory and applications of the sine-Gordon equation”, *Rev. Nuovo Cim.* 1 (2) (1971) 227-267.
- [3] M.Błaszak, B.Silindir, B.M.Szablikowski: “The R-matrix approach to integrable systems on time scales”, *J. Phys. A: Math. Theor.* 41 (2008) 385203.
- [4] A.I.Bobenko, U.Pinkall: “Discrete surfaces with constant negative Gaussian curvature and the Hirota equation”, *J. Diff. Geom.* 43 (1996) 527-611.
- [5] M.Bohner, A.Peterson: *Dynamic equations on time scales. An introduction with applications*, Birkhäuser, Boston 2001.
- [6] M.Bohner, A.Peterson: “A survey of exponential functions on time scales”, *Rev. Cubo Mat. Edu.* 3 (2) (2001) 285-301.
- [7] E.Bour: ”Théorie de la déformation des surfaces”, *J. Ecole Imperiale Polytechnique* 19 (1862) 148.
- [8] J.L.Cieśliński: “Pseudospherical surfaces on time scales: a geometric definition and the spectral approach”, *J. Phys. A: Math. Theor.* 40 (2007) 12525-12538.
- [9] J.Cieśliński: “An algebraic method to construct the Darboux matrix”, *J. Math. Phys.* 36 (1995) 5670-5706.
- [10] J.L.Cieśliński: “The structure of spectral problems and geometry: hyperbolic surfaces in E^3 ”, *J. Phys. A: Math. Gen.* 36 (2003) 6423-6440.

- [11] J.L.Cieřliński: “Some implications of a new approach to exponential functions on time scales”, *Discr. Contin. Dyn. Syst.* , supplement 2011, 302-311.
- [12] J.L.Cieřliński: “New definitions of exponential, hyperbolic and trigonometric functions on time scales”, *J. Math. Anal. Appl.* 388 (2012) 8-22.
- [13] C.Q.Dai, X.Cen, S.S.Wu: “Exact travelling wave solutions of the discrete sine-Gordon equation obtained via the exp-function method”, *Nonlinear Analysis: Theory, Methods & Applications* 70 (1) (2009) 58-63.
- [14] A.S.Davydov: *Solitons in Molecular Systems*, Reidel, Dordrecht 1985.
- [15] L.A.Ferreira, W.J.Zakrzewski: “Numerical and analytical tests of quasi-integrability in modified sine-Gordon models”, *J. High Energy Phys.* 2014 (1) (2014) 058.
- [16] J.Frenkel, T.Kontorova: “On the theory of plastic deformation and twinning”, *Izv. Akad. Nauk Ser. Fiz.* 1 (1939) 137-149.
- [17] M.Gürses, G.Sh.Guseinov, B.Silindir: “Integrable equations on time scales”, *J. Math. Phys.* 46 (2005) 113510 (22 pp).
- [18] S.Hilger: “Analysis on measure chains – a unified approach to continuous and discrete calculus”, *Results Math.* 18 (1990) 19-56.
- [19] S.Hilger: “Differential and difference calculus – unified!”, *Nonl. Anal. Theory, Meth. Appl.* 30 (1997) 2683-2694.
- [20] R.Hirota: “Nonlinear partial difference equations III: discrete sine-Gordon equation”, *J. Phys. Soc. Japan* 43 (6) (1977) 279-286.
- [21] B.Jackson: “Partial dynamic equations on time scales”, *J. Comp. Appl. Math.* 186 (2006) 391-415.
- [22] A.M.Kosevich, B.A.Ivanov, A.S.Kovalev: “Magnetic solitons”, *Phys. Reports* 194 (3-4) (1990) 117-238.
- [23] G. L. Lamb, Jr: *Elements of Soliton Theory*, Wiley, New York 1980.
- [24] J.Lenells, A.S.Fokas: “On a novel integrable generalization of the sine-Gordon equation”, *J. Math. Phys.* 51 (2) (2010) 055912JMP.
- [25] M.Melko, I.Sterling: “Application of soliton theory to the construction of pseudo-spherical surfaces in \mathbb{R}^3 ”, *Ann. Glob. Anal. Geom.* 11 (1993) 65-107.
- [26] S.J. Orfanidis: “Sine-Gordon equation and nonlinear σ model on a lattice”, *Phys. Rev. D* 18 (1978) 38283832.
- [27] L. Pilloni, D. Levi: “The inverse scattering transform for solving the discrete sine-Gordon equation”, *Phys. Lett. A* 92 (1982) 58.
- [28] A.G.Popov, E.V.Maevskii: “Analytical approaches to the study of the sine-Gordon equation and pseudospherical surfaces”, *J. Math. Sci.* 142 (2007) 2377-2418.
- [29] R.Rajaraman: *Solitons and Instantons: An Introduction to Solitons and Instantons in Quantum Field Theory*, North-Holland 1989.
- [30] M. Remoissenet: *Waves Called Solitons. Concepts and Experiments*, Springer, Berlin 2003.
- [31] A.Sakhnovich: “Sine-Gordon theory in a semi-strip”, *Nonlinear Analysis: Theory, Methods & Applications*, 75 (2) (2012) 964-974.
- [32] A. C. Scott: “A nonlinear Klein-Gordon equation”, *Am. J. Phys.* 37 (1969) 5261.
- [33] A.C.Scott: “Propagation of magnetic flux on a long Josephson tunnel junction”, *Nuovo Cim. B* 69 (1970) 241-261.

- [34] B.M.Szablikowski, M.Błaszak, B.Silindir: “Bi-Hamiltonian structures for integrable systems on regular time scales”, *J. Math. Phys.* 50 (2009) 073502 (17 pp).
- [35] R.Wilcox, A.Ramani, B.Grammaticos: “Solutions of the lattice sine-Gordon equation and the solitons of its cellular automaton”, *J. Phys. A: Math. Theor.* 47 (2014) 125202.
- [36] L.V.Yakushevich: *Nonlinear physics of DNA*, second edition, Wiley, Chichester 2004.