

Criterion for stability of Goldstone Modes and Fermi Liquid behavior in a metal with broken symmetry

Haruki Watanabe¹ and Ashvin Vishwanath^{2,3}

¹*Department of Physics, University of California, Berkeley, California 94720, USA*

²*Department of Physics, University of California, Berkeley, CA 94720, USA*

³*Materials Science Division, Lawrence Berkeley National Laboratories, Berkeley, CA 94720*

There are few general physical principles that protect the low energy excitations of a quantum phase. Of these, Goldstone's theorem and Landau Fermi liquid theory are the most relevant to solids. We investigate the stability of the resulting gapless excitations - Nambu Goldstone bosons (NGBs) and Landau quasiparticles - when coupled to one another, which is of direct relevance to metals with a broken continuous symmetry. Typically, the coupling between NGBs and Landau quasiparticles vanishes at low energies leaving the gapless modes unaffected. If however the low energy coupling is non-vanishing, non-Fermi liquid behavior and overdamped bosons are expected. Here we prove a general criterion which specifies when the coupling is non-vanishing. It is satisfied by the case of a nematic Fermi fluid, consistent with earlier microscopic calculations. In addition, the criterion identifies a new kind of symmetry breaking - of magnetic translations - where non-vanishing couplings should arise, opening a new route to realizing non-Fermi liquid phases.

According to the Goldstone theorem, spontaneous breaking of a continuous symmetry leads to gapless Nambu-Goldstone bosons (NGBs). In a Lorentz invariant theory, these bosons are expected to be well defined excitations, even in the presence of other gapless fields, such as massless Dirac fermions, providing a powerful general mechanism for low energy excitation [1]. A key ingredient leading to their stability is the fact that interactions with NGBs are strongly constrained by symmetry, leading to suppressed couplings at small momentum transfer.

In non-relativistic systems though, such general results are not applicable. A particularly important scenario is spontaneous symmetry breaking in a metallic environment, of which there are numerous examples such as magnetic order in a metal. Do the NGBs then survive as well defined low energy modes? Or does coupling to the high density of gapless fermionic excitations of the metal lead to overdamped excitations? In this work we will establish a general criterion to answer this question based on the pattern of symmetry breaking.

A closely related question has to do with the stability of the Fermi liquid (FL) when coupled to gapless bosonic modes. Besides NGBs, gauge bosons can be gapless over an entire phase, i.e. photons of the electromagnetic field or emergent gauge bosons of spin liquids or quantum Hall states. Alternately, one can tune to a quantum critical point where gapless critical modes centered at wave vector $q = 0$ interact with the FL. The latter two cases, of gauge bosons or $q = 0$ quantum critical bosons coupled to a Fermi sea have been studied in many works [2–17]. These studies conclude that, for example in $d = 2 + 1$ dimensions the lifetime of excitations near the Fermi surface is significantly reduced, leading to an absence of well defined quasiparticles and a breakdown of FL theory. Similarly, the bosonic modes get overdamped and can no longer be observed as well-defined particle-like

excitations. In some cases however, superconductivity intervenes at low energies [18].

In contrast, coupling electrons in a metal to NGBs typically leads to a much more benign outcome. We know from examples of magnons in ferromagnets and phonons in crystals, that NGBs are typically underdamped even in a metallic environment, and the FL theory remains valid. In other words, in these cases the coupling between NGBs and FLs is irrelevant, leading to effectively independent fermionic excitations and NGBs at low energies. This is because interactions involving NGBs are very strongly restricted by both broken and unbroken symmetries. In particular, for these cases the scattering amplitude of electrons off NGBs in the limit of small energy-momentum transfer must vanish. In contrast, quantum critical modes and gauge bosons couple directly to fermions, without the derivative coupling.

However, there is one known exception to this rule. When the continuous spatial rotation in $d = 2 + 1$ dimensions is spontaneously broken by a Fermi surface distortion [19–21], the resulting orientational NGB strongly couples to electrons; i.e., their coupling does *not* vanish in the limit of small energy-momentum transfer. We refer to this type of couplings as *nonvanishing couplings*. In this context, Oganesyan, Fradkin and Kivelson [19] discussed non-Fermi liquid (NFL) behavior and Landau damping of the NGBs, in close analogy with the case of critical bosons or gauge bosons coupled to a FL. However, the deeper reason why this example violates the standard rule of vanishing NGB-electron couplings in the infrared, has been left unclear. Also, whether this is the only pattern of symmetry breaking with nonvanishing coupling, remains an open question.

In this Letter we formulate a simple criterion that allows one to diagnose the nature of the NGB-electron coupling. If the *broken symmetry generator fails to commute with translations*, the coupling is anomalous and is non-

vanishing in the infrared. Furthermore, armed with this criterion we are able to identify a new physical setting, distinct from the spontaneous breaking of rotation symmetry, that also leads to nonvanishing couplings, and thus, if we follow standard arguments, to a NFL and overdamped NGBs.

THE GENERAL CRITERION FOR NONVANISHING COUPLINGS

Let us assume that we are at zero temperature and we make no assumption about spatial dimensionality except that it allows for spontaneous symmetry breaking. NGBs can be associated with symmetry generators that are spontaneously broken, which we label Q_a . Furthermore, to sharply define a Fermi surface we assume the existence of a conserved momentum $\vec{P}$. This could be either the conserved momentum of continuous translation symmetry, or crystal momentum (of discrete translation symmetry). Let

$$[Q_a, P_i] = i\Lambda_{ai}. \quad (1)$$

We now state the general criterion. If $\Lambda_{ai} = 0$, this is the usual situation where the coupling *does* vanish. However, if $\Lambda_{ai} \neq 0$ then the coupling between the NGB and electrons does *not* vanish in the limit of small energy-momentum transfer. Note, this criterion is very general and only involves the pattern of symmetry breaking. For any internal symmetry (for example spin rotation or number conservation), the commutator is zero. Thus, for nonvanishing couplings one must consider a space dependent symmetry. The simple case of broken space translation symmetry has $Q_a = P_a$ and the commutator is again zero, which implies that the corresponding Goldstone modes, the phonons, have vanishing coupling to electrons at small momentum transfer, as is well known.

However, for the case of rotational symmetry breaking, $Q_a = L_z$, which satisfies $[L_z, P_i] = i\epsilon_{ij}P_j \neq 0$ where $i, j \in \{x, y\}$. Thus, nonvanishing couplings are expected in this case, consistent with the results of Oganesyan-Fradkin-Kivelson in the context of nematic order in a 2D Fermi fluid [19–21].

This general criterion allows us to identify an entirely new example of nonvanishing coupling. The criterion for nonvanishing coupling is also fulfilled by spontaneous breaking of *magnetic* translations. That is, begin with charged particles in a uniform magnetic field, with magnetic translation symmetry. Spontaneous formation of a crystal breaks this symmetry, resulting in phonons. Now, the magnetic translation operator $\vec{\mathcal{P}}$ generates NGBs (phonons) and satisfies the non-abelian algebra, $[\mathcal{P}_x, \mathcal{P}_y] = -ieBN$. Thus electron-phonon interactions under a uniform magnetic field are predicted to have nonvanishing coupling as we verify by explicit calculation. This surprising conclusion may be rationalized




FIG. 1. A new route to strongly coupling NGBs and quasi-particles - electron-phonon interaction in a uniform magnetic field. A lattice distortion (phonon fluctuation) with $\vec{\nabla} \cdot \vec{u} \neq 0$ changes the local flux threading each unit cell (darker blue indicates a larger flux), inducing fluctuations of the phase of the hopping matrix t_{ij} . The phonon therefore couples like a gauge field to the fermions, without spatial derivatives.

by imagining the fermions to hop between sites of the corresponding tight-binding model. The external magnetic field affects the phase of the hopping matrix as $t_{ij} \exp(i \int_{\vec{x}_i}^{\vec{x}_j} \vec{A}(\vec{x}', t) \cdot d\vec{x}')$. However, a phonon fluctuation $\vec{u}$ which changes the local flux per a unit cell and produces a fluctuation of t_{ij} , as illustrated in Fig. 1. One can imagine this as resulting from a fluctuating gauge field $\Delta B = B \nabla \cdot \vec{u} = \nabla \times \delta A$. Therefore, for electrons, some part of phonon fluctuation under a magnetic field is equivalent to that of a vector potential $\delta A = B \hat{z} \times \vec{u}$ and the problem resembles that of NFL behavior arising from minimal coupling to a fluctuating gauge field.

PROOF OF THE GENERAL CRITERION

The total Hamiltonian of the system can be split into three pieces, $\mathcal{H}_{\text{tot}} = \mathcal{H}_{\text{el}} + \mathcal{H}_{\text{NGB}} + \mathcal{H}_{\text{int}}$, and each of these terms commutes with symmetry generators. We will mainly be concerned with $\mathcal{H}_{\text{int}}$ which we expand as a series in the NGB fields π^a , $\mathcal{H}_{\text{int}} = \mathcal{H}_{\text{int}}^{(0)} + \mathcal{H}_{\text{int}}^{(1)} + \dots$. Note, $\mathcal{H}_0 \equiv \mathcal{H}_{\text{el}}(\bar{\psi}, \psi) + \mathcal{H}_{\text{int}}^{(0)}(\bar{\psi}, \psi)$ is the mean field Hamiltonian, that defines a one electron problem by picking a symmetry broken ground state. The interaction with NGBs is then determined by symmetry, e.g. the linear coupling for a constant π^a is simply obtained by rotating the mean field Hamiltonian by the corresponding symmetry generator Q_a (see [22] for details).

$$\mathcal{H}_{\text{int}}^{(1)} = -[i\pi^a Q_a, \mathcal{H}_0]. \quad (2)$$

To setup the perturbation theory, we first solve the single-particle electron problem described by $\mathcal{H}_0 \equiv \mathcal{H}_{\text{el}}(\bar{\psi}, \psi) + \mathcal{H}_{\text{int}}^{(0)}(\bar{\psi}, \psi)$ and obtain simultaneous eigenstates $|n\vec{k}\rangle$ of $\mathcal{H}_0$ and the momentum $\vec{P}$,

$$\mathcal{H}_0|n\vec{k}\rangle = \epsilon_{n\vec{k}}|n\vec{k}\rangle, \quad \vec{P}|n\vec{k}\rangle = \vec{k}|n\vec{k}\rangle, \quad (3)$$

where n is the band index. When the translation symmetry is discrete, we replace the second relation with

$$T_i|n\vec{k}\rangle = e^{i\vec{k}\cdot\vec{a}_i}|n\vec{k}\rangle, \quad T_i = e^{i\vec{P}\cdot\vec{a}_i}. \quad (4)$$

where $\{\vec{a}_i\}_{i=1,\dots,d}$ are primitive lattice vectors. The interaction of electrons and NGBs to lowest order can then be written as (see Fig. 2 (1)):

$$\mathcal{H}_{\text{int}}^{(1)} = \sum_{n', n, a} \int \frac{d^d k d^d k'}{(2\pi)^{2d}} v_{n'\vec{k}', n\vec{k}}^a c_{n'\vec{k}'}^\dagger c_{n\vec{k}} \pi_{\vec{q}}^a, \quad (5)$$

where $\vec{q} = \vec{k} - \vec{k}'$ and $v_{n'\vec{k}', n\vec{k}}^a$ is the (bare) vertex function, which is the matrix element of $\mathcal{H}_{\text{int}}^{(1)}$. This can be written via Eqn. 2 as:

$$\begin{aligned} \pi_{\vec{q}}^a v_{n'\vec{k}', n\vec{k}}^a &= -i\pi_{\vec{q}}^a \langle n'\vec{k}' | [Q_a, \mathcal{H}_0] | n\vec{k} \rangle \\ &= i\pi_{\vec{q}}^a \langle n'\vec{k}' | Q_a | n\vec{k} \rangle (\epsilon_{n'\vec{k}'} - \epsilon_{n\vec{k}}), \end{aligned} \quad (6)$$

which, for low energy scattering: $n = n'$ and $\vec{q} \rightarrow 0$ is

$$v_{n\vec{k}', n\vec{k}}^a \approx i \langle n\vec{k}' | Q_a | n\vec{k} \rangle \vec{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}. \quad (7)$$

Clearly, as long as $\langle n\vec{k}' | Q_a | n\vec{k} \rangle$ is *finite*, the vertex vanishes as $\vec{q} \rightarrow 0$. This is why scatterings of electrons off NGBs usually vanish at $\vec{q} = 0$, leaving behind well-defined NGBs and Fermi liquid quasiparticles.

However, we can evade this conclusion if, and only if the matrix element $\langle n\vec{k}' | Q_a | n\vec{k} \rangle$ *diverges* as $\vec{k}' \rightarrow \vec{k}$. To ensure an appropriate divergence, we will need $[Q_a, P_i] \neq 0$ as we now explain. If $[Q_a, P_i] = i\Lambda_{ai} \neq 0$, then the matrix element: $\langle n\vec{k}' | Q_a | n\vec{k} \rangle = -i \frac{\langle n\vec{k}' | \Lambda_{ai} | n\vec{k} \rangle}{k'_i - k_i}$. Substituting this in Eqn. 7, and setting $\vec{q} \rightarrow 0$ we have the coupling:

$$v_{n\vec{k}', n\vec{k}}^a = \sum_i \langle n\vec{k}' | \Lambda_{ai} | n\vec{k} \rangle \partial_{k_i} \epsilon_{n\vec{k}}, \quad (8)$$

which is generically nonvanishing. This proves our claim that when the symmetry generator corresponding to the NGBs fails to commute with translations, nonvanishing couplings result. If the translation symmetry is discrete rather than continuous, we simply replace P^i by the discrete translation operator T_i , and require $[Q_a, T_i] \neq 0$. Then the matrix element

$$\langle n\vec{k}' | Q_a | n\vec{k} \rangle = - \frac{\langle n\vec{k}' | [Q_a, T_i] | n\vec{k} \rangle}{e^{i\vec{k}'\cdot\vec{a}_i} - e^{i\vec{k}\cdot\vec{a}_i}} \quad (9)$$

is inversely proportional to $(\vec{k}' - \vec{k}) \cdot \vec{a}_i$ leading again to a non vanishing coupling in Eqn. 7 as $\vec{k}' \rightarrow \vec{k}$. Note that generically $[Q_a, T_i] \neq 0$ follows from $[Q_a, P_i] \neq 0$. Before turning to specific examples we discuss consequences of the non vanishing couplings.




FIG. 2. (1): The bare vertex with one NG line. (2a) and (2b): 1-loop diagrams for the self-energy of boson. (3): the same for electrons.

NON-FERMI LIQUID AND OVERDAMPED NAMBU-GOLDSTONE BOSONS

A non-vanishing coupling connects our problem to well-studied problems of a Fermi surface interacting with gauge or critical bosons. The vertex $\vec{v}_{\vec{k}', \vec{k}} = -e(\vec{k}' + \vec{k})/2m$ of the gauge coupling $-e\vec{A} \cdot \vec{j}$ does not vanish at $\vec{k}' = \vec{k}$. Similarly, the Yukawa interaction between $q = 0$ critical bosons and electrons are not severely restricted by symmetries and nonvanishing couplings are expected (e.g., Yukawa couplings). We can readily argue, via the one-loop calculation below, that nonvanishing couplings destabilize the fixed point of free NGBs and a decoupled Fermi liquid. The actual fate of this strongly coupled problem - a non-Fermi liquid, a superconductor or some other state - requires a case by case analysis and is currently under active investigation.

The boson self-energy correction $\Pi_{ab}(\nu, \vec{q})$ from the diagrams (2a) and (2b) of Fig. 2 is dominated by [22]

$$\Pi_{ab}(\nu, \vec{q}) = -i\pi \frac{\nu}{|\vec{q}|} \gamma^{ab}(\hat{q}) \quad (10)$$

$$\gamma^{ab}(\hat{q}) = \int \frac{d^d k}{(2\pi)^d} v_{\vec{k}, \vec{k}}^a v_{\vec{k}, \vec{k}}^b \delta(\epsilon_{\vec{k}}) \delta(\hat{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{\vec{k}}).$$

The first delta function puts the electron momentum $\vec{k}$ on the Fermi surface and the second one further restricts $\vec{k}$ into a subspace where $\vec{q}$ is tangential to the Fermi surface. Note that the correction in Eqn. 10 vanishes if $v_{\vec{k}, \vec{k}}^a = 0$ (we have suppressed the band index n). The 1-loop corrected boson propagator $D^{-1} = D_0^{-1} - \Pi$ has over-damped poles $\nu \propto -iq^3$ due to the singularity in Eqn. 10. Thus the NGBs are destroyed (overdamped) by interaction with the Fermi surface at this order.

Now, one can study the lifetime of fermionic quasiparticles by evaluating the diagram (3) of Fig. 2 with the corrected propagator D ,

$$\tau^{-1} \equiv -2\text{Im}\Sigma \propto \omega^{d/3}. \quad (11)$$

Therefore, Landau's criterion $\omega\tau \rightarrow \infty$ as $\omega \rightarrow 0$ does

not hold when $d \leq 3$, implying the breakdown of the FL theory.

Thus, this one-loop treatment at least shows the instability of FLs and NGBs against infinitesimal couplings with $\vec{v}_{\vec{k},\vec{k}} \neq 0$. The ultimate fate of these interacting systems continues to be an active area of research [10, 12–14] and we do not expand further on that aspect here. We merely establish the condition when interactions with NGBs are relevant and render the decoupled fixed point unstable, similar to other well-studied cases.

Below, we demonstrate our general criterion through examples.

EXAMPLE 1: INTERNAL SYMMETRIES - CONVENTIONAL COUPLING

Let us first discuss interactions between electrons and magnons in ferromagnets (in the absence of spin-orbit interactions). The coupling between the ferromagnetic order parameter $\vec{m}$ and the electron spin $\vec{s} = \psi^\dagger \frac{\vec{\sigma}}{2} \psi$ ($\vec{\sigma}$ is the Pauli matrix) may not contain any derivatives, e.g.,

$$\mathcal{H}_{\text{el-magnon}} = J\vec{m} \cdot \vec{s}. \quad (12)$$

Hence it is not obvious that the electron-magnon vertex vanishes in the limit of small momentum transfer. However, we know it must from our general criterion, as the spin $\vec{S}$ and the momentum $\vec{P}$ commute.

To see this explicitly, we perform a local SU(2) rotation $U(\vec{x}, t)$ defined by $U^\dagger(\vec{x}, t)\hat{m}(\vec{x}, t) \cdot \vec{\sigma}U(\vec{x}, t) = \sigma_z$. Now, the spin-spin interaction becomes a site dependent Zeeman field along s_z , while electron-magnon interactions are included in derivatives of the rotated electron field $\partial_\mu \psi = U(\partial_\mu + i\mathcal{A}_\mu)\psi'$ through $\mathcal{A}_\mu \equiv -iU^\dagger \partial_\mu U$. If we expand $\mathcal{A}_\mu$ in series of NGB fields, each term contains one derivative acting on them. Therefore, electron-magnon interactions vanish in the limit of small energy-momentum transfer (see [22] for details).

In general, generators Q_a of internal symmetries commute with $\vec{P}$ and therefore we always obtain *vanishing* couplings.

EXAMPLE 2: CONTINUOUS SPACE ROTATION - NONVANISHING COUPLING

Our first nontrivial example is the spontaneous breaking of continuous spatial rotation symmetry. For concreteness, consider nematic order in 2 + 1 dimensions, in which a circular Fermi surface is distorted into an ellipse in the ordered phase. The generator of SO(2) spatial rotations is L_z , which does not commute with the momentum operator, $[L_z, P_i] = i\epsilon_{ij}P_j \neq 0$ (where $i, j \in \{x, y\}$). Hence, from Eqn. 8, we expect a nonvanishing coupling

$$v_{\vec{k}',\vec{k}} = \epsilon_{ij}\langle \vec{k}'|p_j|\vec{k} \rangle \partial_{k_i} \epsilon_{\vec{k}}. \quad (13)$$

where we assume a single band, and for simplicity, ignore spin.

To see this from an explicit calculation, suppose that the spatial SO(2) rotation is spontaneously broken by the order parameter $\langle \vec{n} \rangle = (1, 0)^T$. The Goldstone fluctuation θ of the order parameter $\vec{n} = (\cos \theta, \sin \theta)^T$ can couple to the spinless electron field ψ via, e.g., $\mathcal{H}_{\text{int}} = (\chi/2m)|\vec{n} \cdot \vec{\nabla} \psi|^2$ as both $\nabla \psi$ and $\vec{n}$ are vector. Expanding the interaction to first order in θ ($\vec{n} \sim (1, \theta)^T$), we have

$$\mathcal{H}_{\text{int}}^{(0)} = \chi \frac{\nabla_x \psi^\dagger \nabla_x \psi}{2m}, \quad (14)$$

$$\mathcal{H}_{\text{int}}^{(1)} = \chi \theta \frac{\nabla_x \psi^\dagger \nabla_y \psi + \nabla_y \psi^\dagger \nabla_x \psi}{2m}. \quad (15)$$

Hence, the single-particle electron Hamiltonian is given by

$$\mathcal{H}_0 = \frac{p^2}{2m} + \chi \frac{p_x^2}{2m} \quad (16)$$

leading to a single particle dispersion $\epsilon_{\vec{k}} = [(1 + \chi)k_x^2 + k_y^2]/2m$ for plane waves with wave vector k .

By directly evaluating the matrix element of Eqn. 15 for plane waves, we get

$$v_{\vec{k}',\vec{k}} = \frac{\chi}{2m}(k'_x k_y + k_x k'_y). \quad (17)$$

This is consistent with our criterion Eqn. 13. Indeed, using the above dispersion $\epsilon_{\vec{k}}$ and $\langle n\vec{k}|p_j|n\vec{k} \rangle = k_j$ in Eqn. 13, one gets $v_{\vec{k}',\vec{k}} = \chi k_x k_y / m$, which agrees with Eqn. 17 in the limit $\vec{k}' \rightarrow \vec{k}$.

Note, the vertex does not vanish at $\vec{k}' = \vec{k}$ for generic points on the Fermi surface, except for few high-symmetry points with $k_x = 0$ or $k_y = 0$. Therefore, at the most of the part of the Fermi surface, the quasi-particle lifetime is heavily suppressed by the interaction with the NGB θ originated from spontaneously broken continuous rotation. A nematic order of a elliptically distorted Fermi surface [19, 20] and a ferromagnetic order in the presence of a Rashba interaction [21, 23] are known examples of this mechanism.

Finally, let us remark on a subtlety regarding space-time symmetries. In certain cases, even if the spatial rotation is spontaneously broken, NGBs associated with the broken rotation may not appear. Suppose translations $p_{x,y}$ are spontaneously broken in 2 + 1 dimensions. Although rotation symmetry is also broken it does not lead to independent NGBs. Phonons originating from $p_{x,y}$ play the role of the NGB of ℓ_z as well, and the fluctuation θ associated with ℓ_z is related to displacement fields by $\theta = \partial_x u_y - \partial_y u_x$. Although the field θ can couple strongly to electrons, these additional derivatives annihilate the scattering in the limit of small energy-momentum transfer. Even when only p_x or p_y is broken, ℓ_z cannot produce an independent NGB. For example, helimagnets in 3 + 1 dimensions with the spiral vector along the z -axis




FIG. 3. (a) Electron band structure under a uniform magnetic field (Landau levels). (b) A spontaneously generated periodic lattice potential $V(\vec{x})$ produces dispersing bands. The quasi-particle excitation of the partially-filled band (filled states shaded in blue) has a reduced life time due to the non vanishing electron-phonon interaction.

breaks $p_z - \ell_z$ and $\ell_{x,y}$ but the phonon associated with p_z plays the role of NGBs of $\ell_{x,y}$ and orientational NGBs are absent [24–27].

EXAMPLE 3: MAGNETIC TRANSLATION - NONVANISHING COUPLING

As a new example of nonvanishing couplings, we discuss continuous translation under a uniform magnetic field in 2+1 dimensions. Suppose that a crystalline order with lattice vectors $\{\vec{a}_i\}_{i=1,2}$ is spontaneously formed, breaking the magnetic translation and giving birth to phonons (NGBs). We assume an integer flux quantum per a unit cell for the commutativity of the lattice translations $T_i \equiv e^{i\vec{p}^B \cdot \vec{a}_i}$, where (p_x^B, p_y^B) is given by $(-i\partial_x, -i\partial_y + eBx)$ in the Landau gauge $\vec{A} = -By\hat{x}$. Thanks to an effective periodic potential, the electron band structure becomes dispersive (Fig. 3). We are interested in coupling the NGBs (phonons) to quasiparticle excitations near the Fermi surface of a partially filled band.

In this case, the conserved (magnetic) momenta $\vec{p}^B$ also plays the role of broken generators $Q_a = p_a^B$ ($a = x, y$) that produce phonons. Hence, we should look at the commutation relation $[p_a^B, p_b^B] = -i\epsilon_{ab}eB \neq 0$. For discrete translations, we have $[\vec{p}^B, T_i] = -eB\hat{z} \times \vec{a}_i T_i$ (no sum over i), and we expect nonvanishing coupling from Eqns. 7 and 9:

$$\vec{v}_{n\vec{k}, n\vec{k}} = eB\hat{z} \times \vec{a}_i (\vec{b}_i \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}). \quad (18)$$

Here $\{\vec{b}_i\}_{i=1,2}$ are reciprocal lattice vectors $\vec{a}_i \cdot \vec{b}_j = \delta_{ij}$. Therefore, electrons may show NFL behaviors as a result of the nonvanishing interaction with phonons. As we know, in the absence of the magnetic field $B = 0$, the electron-phonon coupling is conventional as one can see from Eqn. 18.

Let us confirm the nonvanishing coupling in Eqn. 18 from a direct calculation. For simplicity, we assume one

flux quantum per square lattice unit cell and assume the mean-field lattice potential experienced by electrons is:

$$V(\vec{x}) = -V_0 [\cos(2\pi x/a) + \cos(2\pi y/a)]. \quad (19)$$

The electrons and phonons interact with each other through the potential $\mathcal{H}_{\text{int}} = V(\vec{x} - \vec{u})\psi^\dagger\psi$. Expanding $\mathcal{H}_{\text{int}}$ in series of $\vec{u}$, we have

$$\mathcal{H}_0 = \frac{(\vec{p} - e\vec{A})^2}{2m} + V(\vec{x}), \quad (20)$$

$$\mathcal{H}_{\text{int}}^{(1)} = -\vec{u} \cdot \vec{\nabla} V. \quad (21)$$

We diagonalize $\mathcal{H}_0$ in the strong magnetic field limit, perturbatively taking into account the lattice potential to the lowest order in mV_0/eB . In the Landau gauge, the lowest Landau level wave functions that simultaneously diagonalize T_i are given by [22, 28]

$$\Psi_{\vec{k}} \propto \sum_{m \in \mathbb{Z}} e^{-\frac{1}{2}(\frac{y}{\ell} + k_x \ell + \frac{2\pi \ell}{a} m)^2 + i(k_x + \frac{2\pi}{a} m)x - ik_y a m}. \quad (22)$$

Therefore, the lowest electron band to first order in perturbation theory is

$$\begin{aligned} \epsilon_{\vec{k}} &= E_0 + \langle \vec{k} | V | \vec{k} \rangle \\ &= \frac{eB}{2m} - \tilde{V} [\cos(k_y a) + \cos(k_x a)] \end{aligned} \quad (23)$$

with $\tilde{V} \equiv V_0 \exp[-(\pi \ell/a)^2]$. Utilizing our criterion 18, and the dispersion above we predict the non vanishing coupling $\vec{v}_{\vec{k}', \vec{k}} = eB\tilde{V}a (-\sin k_y a, \sin k_x a)$

Now we directly computing the electron-phonon vertex by evaluating matrix elements of Eqn. 21 with the zeroth-order wave function (22) and $u = u_{\vec{q}} e^{i\vec{q} \cdot \vec{x}}$, which gives:

$$\vec{v}_{\vec{k}', \vec{k}} = eB\tilde{V}a e^{-\frac{(q\ell)^2}{4} - iq_y \frac{k'_x + k_x}{2} \ell^2} \begin{pmatrix} -\sin \frac{(k'_y + k_y + iq_x)a}{2} \\ \sin \frac{(k'_x + k_x - iq_y)a}{2} \end{pmatrix}, \quad (24)$$

which in fact agrees with our formula when $\vec{k}' \rightarrow \vec{k}$.

Let us now note some important physical consequences. For our results, it is important that spontaneous breaking of translation symmetry occurs in a system with a uniform magnetic field. On the other hand, if the underlying symmetry is regular translation, and magnetic flux is spontaneously generated in the symmetry breaking process (as in a skyrmion lattice) this does *not* lead to non-vanishing coupling [29], and a Fermi liquid results. A different but equally valid viewpoint on our result is to consider the magnetic field being applied after breaking the translation symmetry (as in a crystal), which should modify the electron-phonon coupling. Therefore, in a clean metal, a magnetic field should induce a nonvanishing coupling between phonons and electrons. Although in principle this would have important consequences, in a typical solid, even at the highest available magnetic fields there is a wide separation $\ell \gg a$ between the magnetic

length $\ell = (eB)^{-1/2}$ and lattice spacing a . The typical dispersion of the Landau levels induced by the lattice, and hence the coupling constant is: $e^{-C(\ell/a)^2} \ll 1$ [22]. Thus, although a non vanishing coupling is expected, its absolute magnitude is extremely small. A more promising physical scenario is the quantum Hall regime, where $\ell \sim a$, and where translation symmetry breaking into stripe and bubble phases are predicted and may have been observed in higher Landau levels [30–33]. We leave an analysis of this interesting possibility to future work.

We are grateful to Andrew Potter, Siddharth Parameswaran, Yasaman Bahri, Philipp Dumitrescu, and Tomáš Brauner for discussions and especially thank Max Metlitski for stimulating discussions at the early stages of this work and for useful comments on the draft. H. W. was supported by the Honjo International Scholarship Foundation and A. V. is supported by NSF - DMR 1206728.

-
- [1] Weinberg, S. *The Quantum Theory of Fields, Vol. II* (Cambridge University Press, 1996).
 - [2] Halperin, B. I., Lee, P. A. & Read, N. Theory of the half-filled Landau level. *Phys. Rev. B* **47**, 7312–7343 (1993).
 - [3] Stern, A. & Halperin, B. I. Singularities in the fermi-liquid description of a partially filled Landau level and the energy gaps of fractional quantum Hall states. *Phys. Rev. B* **52**, 5890 (1995).
 - [4] Altshuler, B. L., Ioffe, L. B. & Millis, A. J. Low-energy properties of fermions with singular interactions. *Phys. Rev. B* **50**, 14048–14064 (1994).
 - [5] Nayak, C. & Wilczek, F. Non-fermi liquid fixed point in $2 + 1$ dimensions. *Nuclear Physics B* **417**, 359 – 373 (1994).
 - [6] Nayak, C. & Wilczek, F. Renormalization group approach to low temperature properties of a non-fermi liquid metal. *Nuclear Physics B* **430**, 534 – 562 (1994).
 - [7] Chakravarty, S., Norton, R. E. & Syljuåsen, O. F. Transverse gauge interactions and the vanquished fermi liquid. *Phys. Rev. Lett.* **74**, 1423–1426 (1995).
 - [8] Motrunich, O. I. Variational study of triangular lattice spin-1/2 model with ring exchanges and spin liquid state in κ -(ET)₂Cu₂(CN)₃. *Phys. Rev. B* **72**, 045105 (2005).
 - [9] Lee, S.-S. & Lee, P. A. U(1) gauge theory of the Hubbard model: Spin liquid states and possible application to κ (BEDT-TTF)₂Cu₂(CN)₃. *Phys. Rev. Lett.* **95**, 036403 (2005).
 - [10] Lee, S.-S. Low-energy effective theory of fermi surface coupled with u(1) gauge field in $2 + 1$ dimensions. *Phys. Rev. B* **80**, 165102 (2009).
 - [11] Rech, J., Pépin, C. & Chubukov, A. V. Quantum critical behavior in itinerant electron systems: Eliashberg theory and instability of a ferromagnetic quantum critical point. *Phys. Rev. B* **74**, 195126 (2006).
 - [12] Metlitski, M. A. & Sachdev, S. Quantum phase transitions of metals in two spatial dimensions. i. Ising-nematic order. *Phys. Rev. B* **82**, 075127 (2010).
 - [13] Metlitski, M. A. & Sachdev, S. Quantum phase transitions of metals in two spatial dimensions. ii. spin density wave order. *Phys. Rev. B* **82**, 075128 (2010).
 - [14] Mross, D. F., McGreevy, J., Liu, H. & Senthil, T. Controlled expansion for certain non-fermi-liquid metals. *Phys. Rev. B* **82**, 045121 (2010).
 - [15] Mahajan, R., Ramirez, D. M., Kachru, S. & Raghu, S. Quantum critical metals in $d = 3 + 1$ dimensions. *Phys. Rev. B* **88**, 115116 (2013).
 - [16] Fitzpatrick, A. L., Kachru, S., Kaplan, J. & Raghu, S. Non-fermi-liquid fixed point in a Wilsonian theory of quantum critical metals. *Phys. Rev. B* **88**, 125116 (2013).
 - [17] Sun, K., Fregoso, B. M., Lawler, M. J. & Fradkin, E. Fluctuating stripes in strongly correlated electron systems and the nematic-smectic quantum phase transition. *Phys. Rev. B* **78**, 085124 (2008).
 - [18] Metlitski, M. A., Mross, D. F., Sachdev, S. & Senthil, T. Are non-fermi-liquids stable to Cooper pairing? *arXiv:1403.3694* (2014).
 - [19] Oganesyan, V., Kivelson, S. A. & Fradkin, E. Quantum theory of a nematic fermi fluid. *Phys. Rev. B* **64**, 195109 (2001).
 - [20] Lawler, M. J., Barci, D. G., Fernández, V., Fradkin, E. & Oxman, L. Nonperturbative behavior of the quantum phase transition to a nematic fermi fluid. *Phys. Rev. B* **73**, 085101 (2006).
 - [21] Xu, C. Quantum critical points of helical fermi liquids. *Phys. Rev. B* **81**, 054403 (2010).
 - [22] Watanabe, H. & Vishwanath, A. *Supplemental Material*.
 - [23] Bahri, Y. & Potter, A. *in preparation* (2014).
 - [24] Radzihovsky, L. & Lubensky, T. C. Nonlinear smectic elasticity of helical state in cholesteric liquid crystals and helimagnets. *Phys. Rev. E* **83**, 051701 (2011).
 - [25] Low, I. & Manohar, A. V. Spontaneously broken spacetime symmetries and Goldstone's theorem. *Phys. Rev. Lett.* **88**, 101602 (2002).
 - [26] Watanabe, H. & Murayama, H. Redundancies in Nambu-Goldstone bosons. *Phys. Rev. Lett.* **110**, 181601 (2013).
 - [27] Hayata, T. & Hidaka, Y. Broken spacetime symmetries and elastic variables. *arXiv:1312.0008* (2013).
 - [28] Haldane, F. D. M. & Rezayi, E. H. Periodic Laughlin-Jastrow wave functions for the fractional quantized Hall effect. *Phys. Rev. B* **31**, 2529–2531 (1985).
 - [29] Watanabe, H., Parameswaran, S. A., Raghu, S. & Vishwanath, A. Anomalous fermi liquid phase in metallic skyrmion crystals. *arXiv:1309.7047* (2013).
 - [30] Koulakov, A. A., Fogler, M. M. & Shklovskii, B. I. Charge density wave in two-dimensional electron liquid in weak magnetic field. *Phys. Rev. Lett.* **76**, 499–502 (1996).
 - [31] Fogler, M. M., Koulakov, A. A. & Shklovskii, B. I. Ground state of a two-dimensional electron liquid in a weak magnetic field. *Phys. Rev. B* **54**, 1853–1871 (1996).
 - [32] Moessner, R. & Chalker, J. T. Exact results for interacting electrons in high Landau levels. *Phys. Rev. B* **54**, 5006–5015 (1996).
 - [33] Eisenstein, J. Two-dimensional electrons in excited Landau levels: evidence for new collective states. *Solid State Communications* **117**, 123–131 (2001).

SUPPLEMENTAL MATERIAL

for “Criterion for stability of Goldstone Modes and Fermi Liquid behavior in a metal with broken symmetry”

1. Interaction with constant NGB fields

In this section, we prove the formula $\mathcal{H}_{\text{int}}^{(1)} = -[i\pi^a Q_a, \mathcal{H}_0]$ for constant NG fields π^a . More precisely, we prove the Lagrangian version, $\mathcal{L}_{\text{int}}^{(1)} = -[i\pi^a Q_a, \mathcal{L}_0]$. (Here, commutation relations with Q_a and the Lagrangian density means the symmetry transformation of the fields contained in the Lagrangian, as we explain below.) These two statements are equivalent as long as symmetry generators Q_i commute with the total Hamiltonian of the system. (We discuss few exceptions in the next section.)

In general, we can decompose the total Lagrangian density into three pieces,

$$\mathcal{L}_{\text{tot}} = \mathcal{L}_{\text{el}}(\bar{\psi}, \partial_\mu \bar{\psi}, \psi, \partial_\mu \psi) + \mathcal{L}_{\text{NG}}(\pi^a, \partial_\mu \pi^a) + \mathcal{L}_{\text{int}}(\bar{\psi}, \partial_\mu \bar{\psi}, \psi, \partial_\mu \psi, \pi^a, \partial_\mu \pi^a). \quad (25)$$

We define $\mathcal{L}_{\text{int}}^{(1)}$ and $\mathcal{L}_0$ by

$$\mathcal{L}_{\text{int}}^{(1)} = \left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \pi^a} \right|_{\pi=0} \pi^a + \left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \pi^a} \right|_{\pi=0} \partial_\mu \pi^a, \quad (26)$$

$$\mathcal{L}_0 = \mathcal{L}_{\text{el}}(\bar{\psi}, \partial_\mu \bar{\psi}, \psi, \partial_\mu \psi) + \mathcal{L}_{\text{int}}(\bar{\psi}, \partial_\mu \bar{\psi}, \psi, \partial_\mu \psi, 0, 0). \quad (27)$$

For constant π^a , we can drop the second term of $\mathcal{L}_{\text{int}}^{(1)}$ as $\partial_\mu \pi^a = 0$.

Internal symmetries

Let us start with a general symmetry breaking pattern $G \rightarrow H$ of internal symmetries. We introduce a NG field π^a for each broken generator Q_a ($a = 1, 2, \dots, \dim G/H$) to describe low-energy fluctuations of the order parameter. Under the symmetry transformation $U = e^{i\epsilon^i Q_i}$, NG fields transforms as

$$(\pi^a)' \equiv U_\epsilon \pi^a U_\epsilon^\dagger = \pi^a + \epsilon^i h_i^a(\pi) + O(\epsilon^2), \quad (28)$$

and its infinitesimal form is

$$\delta_i \pi^a \equiv (\pi^a)' - \pi^a = [iQ_i, \pi^a] = h_i^a(\pi). \quad (29)$$

In the standard parametrization introduced by Refs. [S. Coleman *et al*, Phys. Rev. 177, 2239 (1969), C. G. Callan *et al*, Phys. Rev. 177, 2247 (1969)], $h_b^a(\pi) = \delta_b^a + O(\pi)$ for broken generators Q_b and $h_\rho^a(\pi) = O(\pi)$ for unbroken generators Q_ρ . Namely, a broken generator Q_a shifts π^a by a constant and an unbroken generator Q_ρ does not shift any NG fields by a constant amount.

Each term of the Lagrangian density $\mathcal{L}_A$ ($A = \text{el, NG, int}$) is invariant under the symmetry transformation (up to total derivatives). Namely, $U_\epsilon \mathcal{L}_A U_\epsilon^\dagger = \mathcal{L}_A$. Hence,

$$\begin{aligned} 0 &= [iQ_a, \mathcal{L}_{\text{int}}] \\ &= [iQ_a, \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \bar{\psi}} + [iQ_a, \partial_\mu \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \bar{\psi}} + [iQ_a, \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \psi} + [iQ_a, \partial_\mu \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \psi} + [iQ_a, \pi^b] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \pi^b} + [iQ_a, \partial_\mu \pi^b] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \pi^b}. \end{aligned} \quad (30)$$

We set $\pi^a = 0$ after substituting the relation $[iQ_b, \pi^a] = \delta_b^a + O(\pi)$:

$$\left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \pi^a} \right|_{\pi=0} = - \left([iQ_a, \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \bar{\psi}} + [iQ_a, \partial_\mu \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \bar{\psi}} + [iQ_a, \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \psi} + [iQ_a, \partial_\mu \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \psi} \right) \Big|_{\pi=0} \quad (31)$$

The right hand side is nothing but $-[iQ_a, \mathcal{L}_{\text{int}}|_{\pi=0}]$. Hence, by multiplying π^a to both hand side, we get

$$\left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \pi^a} \right|_{\pi=0} \pi^a = -\pi^a [iQ_a, \mathcal{L}_{\text{int}}|_{\pi=0}]. \quad (32)$$

Since $\mathcal{L}_{\text{el}}$ commutes with Q_a , we can add it to the inside of the commutator. Therefore, for constant π^a ,

$$\mathcal{L}_{\text{int}}^{(1)} = -[i\pi^a Q_a, \mathcal{L}_0]. \quad (33)$$

As the simplest example, let us discuss the spin-spin interaction in ferromagnets.

$$\mathcal{L}_{\text{el}} = i\psi^\dagger \partial_t \psi - \frac{|\vec{\nabla}\psi|^2}{2m}, \quad \mathcal{L}_{\text{int}} = J\vec{n} \cdot \vec{s}, \quad (34)$$

where $\vec{n}$ is the normalized ferromagnetic order parameter, ψ is an electron field with the spin degree of freedom, $\vec{s} \equiv \psi^\dagger \vec{\sigma} \psi$ is the electron spin, and $\vec{\sigma}$ is the Pauli matrix. The electron spin satisfies the commutation relation $[s_i(\vec{x}, t), s_j(\vec{x}', t)] = i\epsilon_{ijk} s_k(\vec{x}, t) \delta^d(\vec{x} - \vec{x}')$ and $[s_i(\vec{x}, t), n_j(\vec{x}', t)] = 0$.

We introduce fluctuation $\pi_{x,y}(\vec{x}, t)$ as $\vec{n} = (\pi_y, -\pi_x, 1)^T + O(\pi_{x,y}^2)$. By expanding the interaction to the linear order in fluctuation, we find

$$\mathcal{L}_0 = i\psi^\dagger \partial_t \psi - \frac{|\vec{\nabla}\psi|^2}{2m} + \frac{J}{2} s_z, \quad (35)$$

$$\mathcal{L}_{\text{int}}^{(1)} = J(\pi_y s_x - \pi_x s_y), \quad (36)$$

Then it can be readily shown that

$$\mathcal{L}_{\text{int}}^{(1)} = -\pi^x [iQ_x, \mathcal{L}_0] - \pi^y [iQ_y, \mathcal{L}_0] \quad (37)$$

for $\vec{Q} = \int d^d x (\vec{s} + m\vec{n})$.

Equivalently, in terms of the Hamiltonian,

$$\mathcal{H}_0 = \frac{\vec{p}^2}{2m} - J s_z, \quad \mathcal{H}_{\text{int}}^{(1)} = -J(\pi_y s_x - \pi_x s_y), \quad (38)$$

and it is straightforward to check

$$\mathcal{H}_{\text{int}}^{(1)} = -\pi_x [iQ_x, \mathcal{H}_0] - \pi_y [iQ_y, \mathcal{H}_0]. \quad (39)$$

Translation

Now we move on to spacetime symmetries. As we will see, the above derivation applies with only some minor changes.

Let us discuss translation $\vec{x}' = \vec{x} + \vec{a}$ as the easiest example. The displacement field $\vec{u}(\vec{x}, t)$ obeys the transformation rule,

$$\vec{u}'(\vec{x}, t) \equiv e^{i\vec{a} \cdot \vec{P}} \vec{u}(\vec{x}, t) e^{-i\vec{a} \cdot \vec{P}} = \vec{u}(\vec{x} - \vec{a}, t) + \vec{a}, \quad (40)$$

$$\delta_j u^i(\vec{x}, t) \equiv u'^i(\vec{x}, t) - u^i(\vec{x}, t) = [iP^j, u^i(\vec{x}, t)] = \delta_j^i - \partial_j u^i. \quad (41)$$

Each component $\mathcal{L}_A$ is assumed to transform as a scalar,

$$\mathcal{L}'_A(\vec{x}, t) = e^{i\vec{a} \cdot \vec{P}} \mathcal{L}_A(\vec{x}, t) e^{-i\vec{a} \cdot \vec{P}} = \mathcal{L}_A(\vec{x} - \vec{a}, t), \quad (42)$$

$$\delta \mathcal{L}_A(\vec{x}, t) = [i\vec{P}, \mathcal{L}_A(\vec{x}, t)] = -\vec{\nabla} \mathcal{L}_A(\vec{x}, t), \quad (43)$$

so that $L_A = \int d^d x \mathcal{L}_A$ commutes with $\vec{P}$. On the other hand, by explicitly computing $\delta \mathcal{L}_{\text{int}} = [i\vec{P}, \mathcal{L}_{\text{int}}(\vec{x}, t)]$, we have

$$\delta \mathcal{L}_{\text{int}} = [i\vec{P}, \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \bar{\psi}} + [i\vec{P}, \partial_\mu \bar{\psi}] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \bar{\psi}} + [i\vec{P}, \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \psi} + [i\vec{P}, \partial_\mu \psi] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu \psi} + [i\vec{P}, u^i] \frac{\partial \mathcal{L}_{\text{int}}}{\partial u^i} + [i\vec{P}, \partial_\mu u^i] \frac{\partial \mathcal{L}_{\text{int}}}{\partial \partial_\mu u^i}. \quad (44)$$

Therefore, using Eq. (41) first, then setting $\vec{u} = 0$ and multiplying $\vec{u}$, we get

$$\left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \vec{u}} \right|_{\vec{u}=0} \cdot \vec{u} = -\vec{u} \cdot [i\vec{P}, \mathcal{L}_{\text{int}}|_{\vec{u}=0}] - \vec{u} \cdot \vec{\nabla} \mathcal{L}_{\text{int}}|_{\vec{u}=0}. \quad (45)$$

Adding $0 = -\vec{u} \cdot [i\vec{P}, \mathcal{L}_{\text{el}}] - \vec{u} \cdot \vec{\nabla} \mathcal{L}_{\text{el}}$ to the right hand side, we get

$$\mathcal{L}_{\text{int}}^{(1)} = \left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \vec{u}} \right|_{\vec{u}=0} \cdot \vec{u} = -[i\pi^a Q_a, \mathcal{L}_0] - \vec{\nabla} \cdot (\vec{u} \mathcal{L}_0) \quad (46)$$

for a constant NG field $\vec{u}$. The last term is just a total derivative and can be dropped.

Even for the magnetic translation, this derivation does not change at all, since the displacement field is real and its transformation rule does not involve phase rotation. All characteristic features of the magnetic transformation are hidden in the commutation relation $[i\vec{P}_B, \psi]$.

Rotation

In the case of the spatial rotation $\vec{x}' = R_\epsilon \vec{x}$ with

$$R_\epsilon = \begin{pmatrix} \cos \epsilon & -\sin \epsilon \\ \sin \epsilon & \cos \epsilon \end{pmatrix}, \quad (47)$$

the NG field originating from the rotational symmetry breaking transforms as

$$\theta'(\vec{x}, t) \equiv e^{i\epsilon L_z} \theta(\vec{x}, t) e^{-i\epsilon L_z} = \theta(R_{-\epsilon} \vec{x}, t) + \epsilon, \quad (48)$$

$$\delta\theta(\vec{x}, t) \equiv \theta'(\vec{x}, t) - \theta(\vec{x}, t) = [iL_z, \theta(\vec{x}, t)] = 1 - (x\partial_y - y\partial_x)\theta(\vec{x}, t) \quad (49)$$

and the Lagrangian density transforms as

$$\mathcal{L}'_A(\vec{x}, t) = e^{i\epsilon L_z} \mathcal{L}_A(\vec{x}, t) e^{-i\epsilon L_z} = \mathcal{L}_A(R_{-\epsilon} \vec{x}, t), \quad (50)$$

$$\delta\mathcal{L}_A(\vec{x}, t) = [iL_z, \mathcal{L}_A(\vec{x}, t)] = -(x\partial_y - y\partial_x)\mathcal{L}_A(\vec{x}, t) = -\partial_y(x\mathcal{L}_A) + \partial_x(y\mathcal{L}_A), \quad (51)$$

In exactly the same way as above, we can show

$$\left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \theta} \right|_{\theta=0} \theta = -\theta [iL_z, \mathcal{L}_0] - \theta \partial_y(x\mathcal{L}_0) + \theta \partial_x(y\mathcal{L}_0). \quad (52)$$

Again, when θ is constant, the last two terms are total derivatives and can be dropped.

General spacetime symmetries

It should be now clear how to extend the above derivation to general spacetime symmetries. NG fields π^a corresponding to broken generators Q_a obey

$$\delta\pi^a(\vec{x}, t) = [iQ_a, \pi^b(\vec{x}, t)] = \delta_a^b + f_a^\mu(\vec{x}, t) \partial_\mu \pi^b(\vec{x}, t) \quad (53)$$

with some functions $f_a^\mu(\vec{x}, t)$. For those symmetries such that $\delta\mathcal{L}_A(\vec{x}, t) = [iQ_a, \mathcal{L}_A(\vec{x}, t)]$ is a total derivative, we have

$$\left. \frac{\partial \mathcal{L}_{\text{int}}}{\partial \pi^a} \right|_{\pi=0} \pi^a = -\pi^a [iQ_a, \mathcal{L}_0] - \pi^a \partial_\mu F_a^\mu. \quad (54)$$

Hence, for constant π^a s, $\mathcal{L}_{\text{int}}^{(1)} = -[i\pi^a Q_a, \mathcal{L}_0]$ up to total derivatives.

2. Photons as NGBs

In this section, we discuss the electron-photon interaction $\mathcal{L}_{\text{int}} = -A^\mu j_\mu$ in the framework developed above. To that end, let us first review how to understand photons as NGBs.

The Lagrangian of the electron-photon interacting system is given by

$$\mathcal{L} = i\bar{\psi}(\partial_t + ieA_t)\psi - \frac{|(\vec{\nabla} - ie\vec{A})\psi|^2}{2m} - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\xi}(\partial_\mu A^\mu)^2. \quad (55)$$

Here, we added the term $-(1/2\xi)(\partial_\mu A^\mu)^2$ to fix the gauge (the R_ξ gauge). Then, the local gauge symmetry is reduced to the residual *global* symmetry specified by

$$\psi = \psi e^{-ie\epsilon(x)}, \quad A'_\mu = A_\mu + \partial_\mu \epsilon(x), \quad \epsilon(x) = a + b_\mu x^\mu, \quad a, b_\mu \in \mathbb{R}. \quad (56)$$

Let Q and Q_μ be the corresponding Noether charges and $P^\mu = (H, \vec{P})$ be the (four) momentum operator. Their expressions are given by

$$Q = \int d^d x e\bar{\psi}\psi, \quad (57)$$

$$Q_\mu = \int d^d x (\pi_\mu + ex_\mu \bar{\psi}\psi), \quad (58)$$

$$P_\mu = \int d^d x [\pi_\nu \partial_\mu A^\nu + (i/2)(\bar{\psi}\partial_\mu \psi - \partial_\mu \bar{\psi}\psi) - \delta_{\mu 0}\mathcal{L}], \quad (59)$$

where $[A^\mu(\vec{x}, t), \pi_\nu(\vec{y}, t)] = i\delta_\nu^\mu \delta^d(\vec{x} - \vec{y})$ and $[\psi(\vec{x}, t), \bar{\psi}(\vec{y}, t)] = \delta^d(\vec{x} - \vec{y})$. Using these commutation relations, It is straightforward to verify that $[Q_\mu, A^\nu(\vec{x}, t)] = -i\delta_\mu^\nu$ and $[Q_\mu, P^\nu] = -i\delta_\mu^\nu Q$.

The relation $[Q_\mu, A^\nu(\vec{x}, t)] = -i\delta_\mu^\nu$ indicates that Q_μ 's are always broken. As long as Q is unbroken, they produce gapless bosons, which can be identified as photons. $3-1=2$ transverse components are physical, while the longitudinal and the temporal component are unphysical.

When the U(1) symmetry Q is spontaneously broken, the NGB originating from Q plays the role of those of Q_μ . Hence, photons do not show up. This is one way of understanding the Higgs phenomenon. The U(1) NGB does not appear in the physical spectrum either, since it belongs to the unphysical sector of the Hilbert space.

Let us now discuss the electron-photon scattering from this point of view. The total Lagrangian density in Eq. (55) suggests that

$$\mathcal{L}_{\text{el}} = 0, \quad \mathcal{L}_{\text{NG}} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\xi}(\partial_\mu A^\mu)^2, \quad \mathcal{L}_{\text{int}} = i\bar{\psi}(\partial_t + ieA_t)\psi - \frac{|(\vec{\nabla} - ie\vec{A})\psi|^2}{2m}. \quad (60)$$

Hence,

$$\mathcal{L}_0 = \mathcal{L}_{\text{int}}|_{A^\mu=0} = i\bar{\psi}\partial_t\psi - \frac{|\vec{\nabla}\psi|^2}{2m}, \quad \mathcal{L}_{\text{int}}^{(1)} = -A_0j^0 + \vec{A} \cdot \vec{j} = -A_\mu j^\mu, \quad (61)$$

where $j^0 = e\bar{\psi}\psi$ and $\vec{j} = (1/2mi)(\bar{\psi}\vec{\nabla}\psi - \text{c.c.})$. By the straightforward calculation, one can verify that

$$-iA^\mu[Q_\mu, \mathcal{L}_0] = -A_\mu j^\mu = \mathcal{L}_{\text{int}}^{(1)}. \quad (62)$$

Therefore, the electron-photon interaction can be understood as an example of the general class of electron-NGB interactions.

Note, however, that the Hamiltonian version of this relation $\mathcal{H}_{\text{int}}^{(1)} = A_\mu j^\mu = -iA^\mu[Q_\mu, \mathcal{H}_0]$ is not true for the temporal component A_0 . For the spatial component $\vec{A}$, one can check it particularly easily in the single-particle picture, where $\hat{Q}_i = -e\hat{x}^i$, $\hat{H}_0 = \hat{p}^2/2m$, and $-A^i[i\hat{Q}_i, \hat{H}_0] = ieA^i[\hat{x}^i, \hat{p}^2/2m] = A_i\hat{j}^i$ with $\hat{j}^i = e\hat{p}^i/m$. However, $Q_0 = et$ (t is not an operator) commutes with $\hat{H}_0$ and the linear interaction eA_0 cannot be written as $-A_0[i\hat{Q}_0, \hat{H}_0]$. This failure originates from the fact that Q_0 *does not* commute with Hamiltonian $[Q_0, H] = -iQ \neq 0$, even though Q_0 is still a symmetry of the system in the sense that it leaves the Lagrangian invariant.

Another example of this type is the Galilean boost $\vec{B}$, which satisfies $[\vec{B}, H] = -i\vec{P} \neq 0$. So if the Galilean boost was spontaneously broken and if it produced independent NGBs $\vec{v}$, their coupling to electrons would not satisfy the relation $\mathcal{H}_{\text{int}}^{(1)} = -i\vec{v} \cdot [\vec{B}, \mathcal{H}_0]$. However, the Galilean boost does not usually produce NGBs. For example, in superfluids, the Bogoliubov mode originating from spontaneously broken U(1) symmetry plays the role of the NGB of the boost symmetry, again due to the linear dependence of the current operator.

3. Singularities in the matrix element $\langle n\vec{k}'|Q_a|n\vec{k}\rangle$

When an operator Q_a does not commute with the generator of the translation $\vec{P}$, *i.e.*, $[Q_a, \vec{P}] \neq 0$, we have $[Q_a, e^{i\vec{P}\cdot\vec{a}_i}] \neq 0$. By further assuming that $\langle n\vec{k}|[Q_a, e^{i\vec{P}\cdot\vec{a}_i}]|n\vec{k}\rangle \neq 0$, which is generically true except for some high symmetry points in the Brillouin zone, one can prove that the expectation value $\langle n\vec{k}|Q_a|n\vec{k}\rangle$ is not well-defined.

For example, using commutation relations

$$[x_i, p_j] = i\delta_{ij}, \quad (63)$$

$$[\ell_z, p_i] = i\epsilon_{ij}p_j, \quad (64)$$

$$[p_i^B, p_j^B] = -i\epsilon_{ij}eB, \quad (65)$$

one can show

$$[\vec{x}, e^{i\vec{p}\cdot\vec{a}_i}] = -\vec{a}_i e^{i\vec{p}\cdot\vec{a}_i}, \quad (66)$$

$$[\ell_z, e^{i\vec{p}\cdot\vec{a}_i}] = -\hat{z} \cdot \vec{a}_i \times \vec{p} e^{i\vec{p}\cdot\vec{a}_i}, \quad (67)$$

$$[\vec{p}^B, e^{i\vec{p}^B\cdot\vec{a}_i}] = -eB\hat{z} \times \vec{a}_i e^{i\vec{p}^B\cdot\vec{a}_i}, \quad (68)$$

respectively. We now evaluate the matrix element of these commutation relations using the definition $e^{i\vec{p}\cdot\vec{a}_i}|n\vec{k}\rangle = e^{i\vec{k}\cdot\vec{a}_i}|n\vec{k}\rangle$. One then finds

$$\langle n'\vec{k}'|\vec{x}|n\vec{k}\rangle = -\frac{e^{i\vec{k}\cdot\vec{a}_i}}{e^{i\vec{k}\cdot\vec{a}_i} - e^{i\vec{k}'\cdot\vec{a}_i}}\vec{a}_i\delta_{\vec{k}',\vec{k}}\delta_{n',n} \simeq \frac{i\vec{a}_i}{(\vec{k} - \vec{k}')\cdot\vec{a}_i}\delta_{\vec{k}',\vec{k}}\delta_{n',n} + O((\vec{k} - \vec{k}')^0), \quad (69)$$

$$\langle n'\vec{k}'|\ell_z|n\vec{k}\rangle = -\frac{e^{i\vec{k}\cdot\vec{a}_i}}{e^{i\vec{k}\cdot\vec{a}_i} - e^{i\vec{k}'\cdot\vec{a}_i}}\hat{z}\cdot\vec{a}_i \times \vec{k}\delta_{\vec{k}',\vec{k}}\delta_{n',n} = \frac{i\hat{z}\cdot\vec{a}_i \times \vec{k}}{(\vec{k} - \vec{k}')\cdot\vec{a}_i}\delta_{\vec{k}',\vec{k}}\delta_{n',n} + O((\vec{k} - \vec{k}')^0), \quad (70)$$

$$\langle n'\vec{k}'|\vec{p}^B|n\vec{k}\rangle = -\frac{e^{i\vec{k}\cdot\vec{a}_i}}{e^{i\vec{k}\cdot\vec{a}_i} - e^{i\vec{k}'\cdot\vec{a}_i}}eB\hat{z} \times \vec{a}_i\delta_{\vec{k}',\vec{k}}\delta_{n',n} = \frac{ieB\hat{z} \times \vec{a}_i}{(\vec{k} - \vec{k}')\cdot\vec{a}_i}\delta_{\vec{k}',\vec{k}}\delta_{n',n} + O((\vec{k} - \vec{k}')^0). \quad (71)$$

This is how one usually derives $\langle x|\hat{p}|x'\rangle = -i\hbar\delta(x - x')\partial_{x'}$ in the single-particle quantum mechanics.

4. Comoving frame of NGBs

In the main text, we discuss the property of electron-NGB vertices using commutation relations. In this section we discuss them from an alternative approach.

a. Magnons in ferromagnets

The spin-spin interaction in ferromagnetic metals reads

$$\mathcal{H}_{\text{int}} = -\frac{J}{2}\vec{n}\cdot\vec{\psi}\vec{\sigma}\psi. \quad (72)$$

It is not obvious from this representation that the electron-magnon vertex vanishes in the limit of small momentum transfer, since the NGB fields in this interaction does not contain derivatives acting on them. However, there is a useful trick to convert these non-derivative interactions into those with at least one derivative acting on NGB fields. Namely, we perform a local SU(2) rotation $U(\vec{x}, t)$ defined by $U^\dagger(\vec{x}, t)\vec{n}(\vec{x}, t)\cdot\vec{\sigma}U(\vec{x}, t) = \sigma_z$. In other words, we take the quantization axis of the electron spin in the comoving frame of the ferromagnetic order parameter. The spin-spin interaction in terms of the new field $\psi' = U^{-1}\psi$ becomes a constant spin-dependent chemical potential $J\psi'^\dagger\sigma_z\psi'/2$. Electron-magnon interactions are instead included in derivatives of the electron field $\partial_\mu\psi = U(\partial_\mu + i\mathcal{A}_\mu)\psi'$ through fluctuations of the Berry phase $\mathcal{A}_\mu \equiv -iU^\dagger\partial_\mu U$. If we expand $\mathcal{A}_\mu$ in series of NGB fields, each term contains one derivative acting on them. Therefore, electron-magnon interactions in $i\psi'^\dagger(\partial_t - i\mathcal{A}_0)\psi'$ and $[(\vec{\nabla} - i\vec{\mathcal{A}})\psi']^2$ vanish in the limit of small energy-momentum transfer.

b. Phonons in crystals

Similarly, the electron-phonon interaction in

$$\mathcal{H}_{\text{int}} = V(\vec{x} - \vec{u})\bar{\psi}(\vec{x}, t)\psi(\vec{x}, t) \quad (73)$$

does not contain derivatives acting on the displacement field $\vec{u}(\vec{x}, t)$, but the electron-phonon scattering vanishes in the limit of small energy-momentum transfer as discussed by using commutation relations.

To see the vanishing scattering more clearly, we can convert the non-derivative coupling $V(\vec{x} - \vec{u})$ into derivative ones by going to the comoving frame of the crystal lattice. That is, we change the integration variable of the Lagrangian from $\vec{x}$ to $\vec{x}' = \vec{x} - \vec{u}$ and redefine the electron field $\psi'(\vec{x}', t) = \psi(\vec{x}, t)$. Then the potential $V(\vec{x} - \vec{u}) = V(\vec{x}')$ can no longer fluctuate, analogously to the above spin-spin interaction after the SU(2) rotation. Instead, all the electron-phonon interactions come from rewriting the volume element and derivatives:

$$d^d x dt = d^d x' dt' (1 + \vec{\nabla}' \cdot \vec{u}) + O((\partial\vec{u})^2), \quad (74)$$

$$\partial_\mu = \partial'_\mu - (\partial'_\mu u^i)\partial'_i + O((\partial\vec{u})^2). \quad (75)$$

It is now clear in this representation that all electron-phonon interactions vanish for a constant $\vec{u}$.

c. Orientational NGBs in phases with rotational symmetry breaking

If we can eliminate all non-derivative couplings by going to the comoving frame of NGBs, there is no hope to get non-vanishing couplings, as derivatives on NGBs vanish in the limit of small energy-momentum transfer. Here we discuss why this comoving frame argument fails in the case of spital rotation and magnetic translation. (More generally, spacetime symmetries except for the ordinary translation.)

If possible, we would like to eliminate all non-derivative couplings in the interacting Lagrangian,

$$\int d^d x dt |\vec{n} \cdot \vec{\nabla} \psi|^2 = \int d^d x dt \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \vec{\nabla} \bar{\psi} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \cdot \vec{\nabla} \psi. \quad (76)$$

If we change the integration variable from $\vec{x}$ to $\vec{x}' = R_\epsilon \vec{x}$, we get

$$\int d^d x' dt \begin{pmatrix} \cos(\theta - \epsilon) \\ \sin(\theta - \epsilon) \end{pmatrix} \cdot \vec{\nabla}' \bar{\psi} \begin{pmatrix} \cos(\theta - \epsilon) \\ \sin(\theta - \epsilon) \end{pmatrix} \cdot \vec{\nabla}' \psi, \quad (77)$$

where

$$R_\epsilon = \begin{pmatrix} \cos \epsilon & -\sin \epsilon \\ \sin \epsilon & \cos \epsilon \end{pmatrix} \quad (78)$$

is the orthogonal matrix for the rotation by a *constant* angle ϵ . Therefore, changing the integration variable effectively shifts θ by $-\epsilon$. Thus one may expect that setting $\epsilon(\vec{x}, t) = \theta(\vec{x}, t)$ locally eliminates all θ dependence without derivatives. However, it does not work for the following reason. If we define $\vec{x}' = R_{\theta(\vec{x}, t)} \vec{x}$ and rewrite derivative $\vec{\nabla}$ in terms of $\vec{\nabla}'$, we find

$$\partial_i = (\partial_i x'^j) \partial'_j = \partial_i [(R_\theta)^j_k x^k] \partial'_j = (R_\theta)^j_i \partial'_j + (\partial_i R_\theta)^j_k x^k \partial'_j. \quad (79)$$

Due to the second term of the last expression, the Lagrangian now explicitly depends on the coordinate. This makes the Lagrangian after the rotation completely useless for any realistic calculations. Especially, we cannot use the Fourier transformation (despite the fact that the translation is not actually broken). Therefore, we cannot discuss the behavior of couplings in the limit of the small momentum transfer.

d. Magnetic translation

We now discuss the magnetic translation. We would like to remove $\vec{u}$ without derivatives in the Lagrangian,

$$\mathcal{L}_{\text{el+int}} = i \bar{\psi} \partial_t \psi - \frac{|(\vec{\nabla} - ie \vec{A}) \psi|^2}{2m} - \bar{\psi} \psi V(\vec{x} - \vec{u}). \quad (80)$$

If we just change the integration variable to $\vec{x}' = \vec{x} - \vec{u}(\vec{x}, t)$, then $\vec{u}$ without derivatives appears from the vector potential,

$$\vec{A} = B \begin{pmatrix} -y \\ 0 \\ 0 \end{pmatrix} = B \begin{pmatrix} -y' - u_y \\ 0 \\ 0 \end{pmatrix}. \quad (81)$$

In order to absorb this new $\vec{u}$ dependence, one can further perform a local gauge transformation,

$$\psi' = e^{-ie B x' u_y} \psi. \quad (82)$$

When u_y is a constant, this combination of the translation and the gauge transformation successfully removes all u_y 's from the Lagrangian. However, for a general $u_y(\vec{x}, t)$, we have

$$\vec{\nabla}' \psi' = e^{-ie B x' u_y} \left(\vec{\nabla}' \psi - ie B \hat{x} u_y \psi - ie B x' \psi \vec{\nabla}' u_y \right). \quad (83)$$

Again the last term introduces an undesirable coordinate dependence to the Lagrangian.

5. Landau levels with lattice momentum

In this section, we summarize the wave function of Landau levels (following Ref. [F. D. M. Haldane and E. H. Rezayi, Phys. Rev. B 31, 2529 (1985)]) that simultaneously diagonalize Hamiltonian and lattice translations,

$$H = \frac{(p_x + eBy)^2 + p_y^2}{2m}, \quad T_x = e^{ip_x a_x}, \quad T_y = e^{i(p_y + eBx)a_y}. \quad (84)$$

We assume a rectangular lattice with primitive lattice vectors $\vec{a}_x = a_x \hat{x}$ and $\vec{a}_y = a_y \hat{y}$ and a flux quantum per a unit cell $eBa_x a_y = 2\pi$. We work in a torus $a_x N_x \times a_y N_y$ ($N_x, N_y \in \mathbb{Z}$) and impose the periodic boundary condition $T_x^{N_x} = T_y^{N_y} = 1$. The number of degeneracy is precisely the number of lattice points,

$$\frac{a_x a_y N_x N_y}{2\pi \ell^2} = N_x N_y. \quad \ell \equiv \sqrt{\frac{1}{eB}}. \quad (85)$$

For each $k = \frac{2\pi}{a_x N_x} i$ ($i = 1, 2, \dots, N_x N_y$), the function

$$\psi_{nk}(\vec{x}) = \sum_{j \in \mathbb{Z}} \frac{H_n \left(\frac{y}{\ell} + k\ell + \frac{2\pi\ell}{a_x} j N_y \right) e^{-\frac{1}{2} \left(\frac{y}{\ell} + k\ell + \frac{2\pi\ell}{a_x} j N_y \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k + \frac{2\pi}{a_x} j N_y \right) x}}{\sqrt{a_x N_x}} \quad (86)$$

represents an simultaneous eigenfunction of the Hamiltonian with the eigenvalue $(eB/m)(n + 1/2)$ and the lattice translation T_x :

$$T_x \psi_{nk}(\vec{x}) = \sum_{j \in \mathbb{Z}} \frac{H_n \left(\frac{y}{\ell} + k\ell + \frac{2\pi\ell}{a_x} j N_y \right) e^{-\frac{1}{2} \left(\frac{y}{\ell} + k\ell + \frac{2\pi\ell}{a_x} j N_y \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k + \frac{2\pi}{a_x} j N_y \right) (x + a_x)}}{\sqrt{a_x N_x}} = e^{ika_x} \psi_{nk}(\vec{x}). \quad (87)$$

In order to make it a simultaneous eigenfunction of T_y as well, we take a superposition

$$\begin{aligned} \Psi_{n\vec{k}}(\vec{x}) &\equiv \sum_{m=1}^{N_y} \frac{e^{-ik_y a_y m}}{\sqrt{N_y}} \psi_{n, k_x + \frac{2\pi}{a_x} m}(\vec{x}) \\ &= \sum_{m=1}^{N_y} \sum_{j \in \mathbb{Z}} \frac{e^{-ik_y a_y (m + j N_y)}}{\sqrt{N_y}} \frac{H_n \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} (m + j N_y) \right) e^{-\frac{1}{2} \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} (m + j N_y) \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k_x + \frac{2\pi}{a_x} (m + j N_y) \right) x}}{\sqrt{a_x N_x}} \\ &= \sum_{m \in \mathbb{Z}} \frac{e^{-ik_y a_y m}}{\sqrt{N_y}} \frac{H_n \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} m \right) e^{-\frac{1}{2} \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} m \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k_x + \frac{2\pi}{a_x} m \right) x}}{\sqrt{a_x N_x}}, \end{aligned} \quad (88)$$

where

$$k_x = \frac{2\pi}{a_x} i_x \quad i_x = 1, 2, \dots, N_x, \quad (89)$$

$$k_y = \frac{2\pi}{a_y} i_y \quad i_y = 1, 2, \dots, N_y. \quad (90)$$

Now $\Psi_{n\vec{k}}(\vec{x})$'s are simultaneous eigenstates of T_y as well:

$$\begin{aligned} T_y \Psi_{n\vec{k}}(\vec{x}) &= e^{ieBx a_y} \sum_{m \in \mathbb{Z}} \frac{e^{-ik_y a_y m}}{\sqrt{N_y}} \frac{H_n \left(\frac{y + a_y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} m \right) e^{-\frac{1}{2} \left(\frac{y + a_y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} m \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k_x + \frac{2\pi}{a_x} m \right) x}}{\sqrt{a_x N_x}} \\ &= \sum_{m \in \mathbb{Z}} \frac{e^{-ik_y a_y m}}{\sqrt{N_y}} \frac{H_n \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} (m + 1) \right) e^{-\frac{1}{2} \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} (m + 1) \right)^2}}{\sqrt{2^n n!} \sqrt{\pi} \ell} \frac{e^{i \left(k_x + \frac{2\pi}{a_x} (m + 1) \right) x}}{\sqrt{a_x N_x}} \\ &= e^{ik_y a_y} \Psi_{n\vec{k}}(\vec{x}). \end{aligned} \quad (91)$$

For the lowest Landau levels, we have

$$\Psi_{\vec{k}}(\vec{x}) \equiv \Psi_{0\vec{k}}(\vec{x}) = \sum_{m \in \mathbb{Z}} \frac{e^{-\frac{1}{2} \left(\frac{y}{\ell} + k_x \ell + \frac{2\pi\ell}{a_x} m \right)^2 + i \left(k_x + \frac{2\pi}{a_x} m \right) x - ik_y a_y m}}{\sqrt{\sqrt{\pi} \ell a_x N_x N_y}}. \quad (92)$$

6. The electron Green function under magnetic field

Here we summarize the free electron Green function under the magnetic field. We expand the electron field operator as

$$\psi(\vec{x}, t) = \sum_{n\vec{k}} \psi_{n\vec{k}}(\vec{x}) c_{n\vec{k}}(t), \quad (93)$$

where $c_{n\vec{k}}(t)$ is the annihilation operator of electrons in the Bloch eigenstate $\psi_{n\vec{k}}(\vec{x})$, either with or without an external magnetic field. $c_{n\vec{k}}(t)$'s satisfy the equal-time anti-commutation relation

$$\{c_{n\vec{k}}(t), c_{n'\vec{k}'}^\dagger(t)\} = \delta_{nn'} \delta_{\vec{k}, \vec{k}'}. \quad (94)$$

The free Hamiltonian can be expressed as

$$H_0 = \sum_{n\vec{k}} \epsilon_{n\vec{k}} c_{n\vec{k}}^\dagger c_{n\vec{k}}. \quad (95)$$

Thus the time-evolution of the annihilation operator under H_0 is $c_{n\vec{k}}(t) = c_{n\vec{k}} e^{-i\epsilon_{n\vec{k}}t}$. The free Green function is then given by

$$\begin{aligned} G_n(\vec{k}, t) &\equiv -i \langle T c_{n\vec{k}}(t) c_{n\vec{k}}^\dagger(0) \rangle \\ &= -i \langle c_{n\vec{k}} c_{n\vec{k}}^\dagger \rangle e^{-i\epsilon_{n\vec{k}}t} \theta(t) + i \langle c_{n\vec{k}}^\dagger c_{n\vec{k}} \rangle e^{-i\epsilon_{n\vec{k}}t} \theta(-t) \\ &= \int \frac{d\omega}{2\pi} e^{-i\omega t} \left[\frac{\theta(\epsilon_{n\vec{k}})}{\omega - \epsilon_{n\vec{k}} + i\delta} + \frac{\theta(-\epsilon_{n\vec{k}})}{\omega - \epsilon_{n\vec{k}} - i\delta} \right] \\ &\equiv \int \frac{d\omega}{2\pi} e^{-i\omega t} G_n(\vec{k}, \omega). \end{aligned} \quad (96)$$

In the derivation, we assumed that single-particle states with $\epsilon_{n\vec{k}} < 0$ are filled and otherwise unfilled. Therefore, the electron Green function in the momentum space takes the same form regardless of the presence or absence of the external magnetic field.

7. Cancellation of the induced mass of NGBs

For completeness, here we check the absence of a mass of NGBs generated by integrating out electrons.

a. Rotation

Let us start with the example of the spatial rotation. For a constant θ , we have

$$\begin{aligned} H_{\text{int}} &= \frac{\chi}{2m} [(k_x \cos \theta + k_y \sin \theta)^2 - k_x^2] \psi_k^\dagger \psi_k \\ &= \frac{\chi}{m} \left[\theta k_x k_y + \frac{1}{2} \theta^2 (k_y^2 - k_x^2) + O(\theta^3) \right] \psi_k^\dagger \psi_k \\ &= \left[-\theta \partial_{\phi_{\vec{k}}} \epsilon_{\vec{k}} + \frac{1}{2} \theta^2 \partial_{\phi_{\vec{k}}}^2 \epsilon_{\vec{k}} + O(\theta^3) \right] \psi_k^\dagger \psi_k, \end{aligned} \quad (97)$$




FIG. 4. 1-loop diagrams for boson self-energy corrections.

where $ke^{i\phi_{\vec{k}}} = k_x + ik_y$ and $\epsilon_{\vec{k}}$ is the electron dispersion,

$$\epsilon_{\vec{k}} = \frac{(1 + \chi)k_x^2 + k_y^2}{2m} - \mu. \quad (98)$$

The boson self-energy Π at $\vec{q} = 0$ and $\nu = 0$ has two contributions at the 1-loop level,

$$\Pi(0) = \int \frac{d^2k d\omega}{(2\pi)^3} \left[\left(\partial_{\phi_{\vec{k}}} \epsilon_{\vec{k}} G(\vec{k}, \omega) \right)^2 + \partial_{\phi_{\vec{k}}}^2 \epsilon_{\vec{k}} G(\vec{k}, \omega) \right]. \quad (99)$$

Here the first (second) term represents the left (right) diagram in Fig. 4. To show their cancelation, we use the relation of the electron Green function $G^{-1}(\vec{k}, \omega) = \omega - \epsilon_{\vec{k}}$:

$$\vec{\nabla}_{\vec{k}} G(\vec{k}, \omega) = [G(\vec{k}, \omega)]^2 \vec{\nabla}_{\vec{k}} \epsilon_{\vec{k}}. \quad (100)$$

Then,

$$\begin{aligned} \Pi(0) &= \int \frac{d^2k d\omega}{(2\pi)^3} \left[\partial_{\phi_{\vec{k}}} \epsilon_{\vec{k}} \partial_{\phi_{\vec{k}}} G(\vec{k}, \omega) + \partial_{\phi_{\vec{k}}}^2 \epsilon_{\vec{k}} G(\vec{k}, \omega) \right] \\ &= \int \frac{d^2k d\omega}{(2\pi)^3} \left[-\partial_{\phi_{\vec{k}}}^2 \epsilon_{\vec{k}} G(\vec{k}, \omega) + \partial_{\phi_{\vec{k}}}^2 \epsilon_{\vec{k}} G(\vec{k}, \omega) \right] = 0. \end{aligned} \quad (101)$$

b. Magnetic translation

Next, for the electron-phonon problem under a magnetic field, we have

$$\begin{aligned} H_{\text{int}} &= -\tilde{V}_x \left[\cos(k_y a_y - \frac{2\pi}{a_x} u_x) - \cos(k_y a_y) \right] - \tilde{V}_y \left[\cos(k_x a_x + \frac{2\pi}{a_y} u_y) - \cos(k_x a_x) \right] \\ &= \left[\left(\frac{2\pi u_y}{a_y} \right) \partial_{k_x a_x} \epsilon_{\vec{k}} - \left(\frac{2\pi u_x}{a_x} \right) \partial_{k_y a_y} \epsilon_{\vec{k}} \right] + \frac{1}{2} \left[\left(\frac{2\pi u_y}{a_y} \right)^2 \partial_{k_x a_x}^2 \epsilon_{\vec{k}} + \left(\frac{2\pi u_x}{a_x} \right)^2 \partial_{k_y a_y}^2 \epsilon_{\vec{k}} \right] + O(u^3). \end{aligned} \quad (102)$$

Therefore, again by using Eq. (100),

$$\begin{aligned} \Pi_{xx}(0) &= \left(\frac{2\pi}{a_x} \right)^2 \int \frac{d^2k d\omega}{(2\pi)^3} \left[\left((\partial_{k_y a_y} \epsilon_{\vec{k}}) G(\vec{k}, \omega) \right)^2 + (\partial_{k_y a_y}^2 \epsilon_{\vec{k}}) G(\vec{k}, \omega) \right] \\ &= \left(\frac{2\pi}{a_x} \right)^2 \int \frac{d^2k d\omega}{(2\pi)^3} \left[(\partial_{k_y a_y} \epsilon_{\vec{k}}) \partial_{k_y a_y} G(\vec{k}, \omega) + (\partial_{k_y a_y}^2 \epsilon_{\vec{k}}) G(\vec{k}, \omega) \right] \\ &= \left(\frac{2\pi}{a_x} \right)^2 \int \frac{d^2k d\omega}{(2\pi)^3} \left[-(\partial_{k_y a_y}^2 \epsilon_{\vec{k}}) G(\vec{k}, \omega) + (\partial_{k_y a_y}^2 \epsilon_{\vec{k}}) G(\vec{k}, \omega) \right] = 0. \end{aligned} \quad (103)$$

The same derivation applies to $\Pi_{xy}(0)$, $\Pi_{yx}(0)$, and $\Pi_{yy}(0)$.

8. The dominant self-energy correction of bosons

In this section, we discuss the boson self-energy correction for a general $\vec{q}$ and ν . To the leading order in q , the contribution of the left diagram in Fig. 4 is given by

$$\begin{aligned} \Pi_{ab}(\nu, \vec{q}) &= \int \frac{d^d k d\omega}{(2\pi)^{d+1}} v_{n\vec{k}, n(\vec{k}+\vec{q})}^a v_{n(\vec{k}+\vec{q}), n\vec{k}}^b G_n(\vec{k}, \omega) G_n(\vec{k} + \vec{q}, \omega + \nu) \\ &= \int \frac{d^d k}{(2\pi)^d} v_{n\vec{k}, n(\vec{k}+\vec{q})}^a v_{n(\vec{k}+\vec{q}), n\vec{k}}^b \frac{f(\epsilon_{n\vec{k}}) - f(\epsilon_{n(\vec{k}+\vec{q})})}{\nu + i\delta - (\epsilon_{n(\vec{k}+\vec{q})} - \epsilon_{n\vec{k}})} \\ &\simeq \int \frac{d^d k}{(2\pi)^d} \delta(\epsilon_{n\vec{k}}) v_{n\vec{k}, n\vec{k}}^a v_{n\vec{k}, n\vec{k}}^b \frac{\hat{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}}{\nu/q + i\delta - \hat{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}}. \end{aligned} \quad (104)$$

As discussed in the previous section, the constant term

$$\Pi_{ab}(0) = - \int \frac{d^d k}{(2\pi)^d} \delta(\epsilon_{n\vec{k}}) v_{n\vec{k},n\vec{k}}^a v_{n\vec{k},n\vec{k}}^b \quad (105)$$

is exactly cancelled by the diamagnetic term (the right diagram in Fig. 4). The imaginary part is given by

$$\begin{aligned} \text{Im}\Pi_{ab}(\nu, \vec{q}) &= -\pi \frac{\nu}{q} \int \frac{d^d k}{(2\pi)^d} \delta(\epsilon_{n\vec{k}}) v_{n\vec{k},n\vec{k}}^a v_{n\vec{k},n\vec{k}}^b \delta(\nu/q - \hat{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}) \\ &\simeq -\pi \frac{\nu}{q} \int \frac{d^d k}{(2\pi)^d} \delta(\epsilon_{n\vec{k}}) v_{n\vec{k},n\vec{k}}^a v_{n\vec{k},n\vec{k}}^b \delta(\hat{q} \cdot \vec{\nabla}_{\vec{k}} \epsilon_{n\vec{k}}). \end{aligned} \quad (106)$$

9. The band width of the electron band under magnetic field

Here we show a simple numerical result on the band width of the electron band structure under a uniform magnetic field, in order to support the claim

$$(\text{band width}) \propto e^{-C\ell^2/a^2}. \quad (107)$$

Here $\ell = (eB)^{-1/2}$ is the magnetic length and a is the lattice constant of the tight binding model. In the continuum limit $a \rightarrow 0$, the Landau levels are flat. For a finite a , the lattice periodic potential produces nonzero dispersions.

By denoting the number of the flux per a unit cell ϕ

$$\frac{\ell^2}{a^2} = \frac{1}{eBa^2} = \frac{1}{\phi} \quad (108)$$

Eq. (107) suggests that

$$\log(\text{band width}) = (\text{const.}) - C\phi^{-1}. \quad (109)$$

In Fig. 5 we show the numerical result for the lowest Landau levels in the square lattice tight-binding model with the nearest neighbor hopping. The logarithm of the band width is indeed proportional to ϕ^{-1} . This result holds for other Landau levels as well, as long as the van-Hove singularity energy is avoided.




FIG. 5. The band width of the lowest landau level in the tight binding model as a function of the inverse flux ϕ^{-1} .