

FINITE-DIMENSIONAL REPRESENTATIONS FOR A CLASS OF GENERALIZED INTERSECTION MATRIX LIE ALGEBRAS

YUN GAO¹ AND LI-MENG XIA^{2,†}

¹Department of Mathematics and Statistics, York University, Canada

²Faculty of Science, Jiangsu University, P.R. China

ABSTRACT. In this paper, we study a class of generalized intersection matrix Lie algebras $\mathbf{gim}(M_n)$, and prove that its every finite-dimensional semi-simple quotient is of type $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$. Particularly, any finite dimensional irreducible $\mathbf{gim}(M_n)$ module must be an irreducible module of $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ and any finite dimensional irreducible $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ module must be an irreducible module of $\mathbf{gim}(M_n)$.

Key Words: intersection matrix algebras; irreducible modules; quotient algebras; affine Lie algebras.

1. INTRODUCTION

In the early to mid-1980s, Peter Slodowy discovered that matrices like

$$M = \begin{bmatrix} 2 & -1 & 0 & 1 \\ -1 & 2 & -1 & 1 \\ 0 & -2 & 2 & -2 \\ 1 & 1 & -1 & 2 \end{bmatrix}$$

were encoding the intersection form on the second homology group of Milnor fibres for germs of holomorphic maps with an isolated singularity at the origin [S1], [S2]. These matrices were like the generalized Cartan matrices of Kac-Moody theory in that they had integer entries, 2's along the diagonal, and $m_{i,j}$ was negative if and only if $m_{j,i}$ was negative. What was new, however, was the presence of positive entries off the diagonal. Slodowy called such matrices generalized intersection matrices:

Definition 1.1. ([S1]) *An $n \times n$ integer-valued matrix $M = (m_{i,j})_{1 \leq i,j \leq n}$ is called a generalized intersection matrix (\mathbf{gim}) if*

$$m_{i,i} = 2,$$

$$m_{i,j} < 0 \text{ if and only if } m_{j,i} < 0, \text{ and}$$

$$m_{i,j} > 0 \text{ if and only if } m_{j,i} > 0$$

for $1 \leq i, j \leq n$ with $i \neq j$.

Slodowy used these matrices to define a class of Lie algebras that encompassed all the Kac-Moody Lie algebras:

[†]the corresponding author's email: xialimeng@ujs.edu.cn.

Definition 1.2 (see [BrM], [S1]). Given an $n \times n$ generalized intersection matrix $M = (m_{i,j})$, define a Lie algebra over \mathbb{C} , called a generalized intersection matrix (GIM) algebra and denoted by $\mathbf{gim}(M)$, with:

generators: $e_1, \dots, e_n, f_1, \dots, f_n, h_1, \dots, h_n$,

relations: (R1) for $1 \leq i, j \leq n$,

$$[h_i, e_j] = m_{i,j}e_j, \quad [h_i, f_j] = -m_{i,j}f_j, \quad [e_i, f_i] = h_i,$$

(R2) for $m_{i,j} \leq 0$,

$$[e_i, f_j] = 0 = [f_i, e_j], \quad (\text{ad } e_i)^{-m_{i,j}+1}e_j = 0 = (\text{ad } f_i)^{-m_{i,j}+1}f_j,$$

(R3) for $m_{i,j} > 0, i \neq j$,

$$[e_i, e_j] = 0 = [f_i, f_j], \quad (\text{ad } e_i)^{m_{i,j}+1}f_j = 0 = (\text{ad } f_i)^{m_{i,j}+1}e_j.$$

If the M that we begin with is a generalized Cartan matrix, then the $3n$ generators and the first two groups of axioms, (R1) and (R2), provide a presentation of the Kac-Moody Lie algebras [GbK], [C], [K].

Slodowy and, later, Berman showed that the GIM algebras are also isomorphic to fixed point subalgebras of involutions on larger Kac-Moody algebras [S1], [Br]. So, in their words, the GIM algebras lie both "beyond and inside" Kac-Moody algebras.

Further progress came in the 1990s as a byproduct of the work of Berman and Moody, Benkart and Zelmanov, and Neher on the classification of root-graded Lie algebras [BrM], [BnZ], [N]. Their work revealed that some families of intersection matrix (**im**) algebras, were universal covering algebras of well understood Lie algebras. An **im** algebra generally is a quotient algebra of a GIM algebra associated to the ideal generated by homogeneous vectors those have long roots (i.e., $(\alpha, \alpha) > 2$).


A handful of other researchers also began engaging these new algebras. For example, Eswara Rao, Moody, and Yokonuma used vertex operator representations to show that **im** algebras were nontrivial [EMY]. Analogous compact forms of **im** algebras arising from conjugations over the complex field were considered in [G]. Peng found relations between **im** algebras and the representations of tilted algebras via Ringel-Hall algebras [P]. Berman, Jurisich, and Tan showed that the presentation of GIM algebras could be put into a broader framework that incorporated Borcherds algebras [BrJT].

In present paper, we study the GIM algebra $\mathbf{gim}(M_n)$ constructed through intersection matrix

$$M_n = (m_{i,j})_{n \times n} = \begin{bmatrix} 2 & -1 & 0 & \cdots & 0 & 1 \\ -1 & 2 & -1 & \cdots & 0 & 0 \\ 0 & -1 & 2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 2 & -1 \\ 1 & 0 & 0 & \cdots & -1 & 2 \end{bmatrix}_{n \times n}$$

where $n \geq 3$, and we build the representation theory of finite-dimensional modules for $\mathbf{gim}(M_n)$.

Generally, $\mathbf{gim}(M_n)$ can be illustrated by the following diagram (also called Dynkin diagram):




where the numbered circles present the indices of e_i, f_i ($1 \leq i \leq n$), the solid line between two circles $i, i+1$ means that $m_{i,i+1} = m_{i+1,i} = -1$ and the unique dotted line means that $m_{1,n} = m_{n,1} = 1$. There is no line between any other pair (i, j) , it means that $m_{i,j} = m_{j,i} = 0$.

2. CONSTRUCTION OF EPIMORPHISMS

Definition 2.1. Suppose that \mathcal{L} is a non-trivial semi-simple Lie algebra and $\mathbf{a} \in \mathbb{Z}_{\geq 0}, \mathbf{c}, \mathbf{d} \in \{0, 1\}$. If $n \geq 3$ and \mathcal{L} is isomorphic to a direct sum of \mathbf{a} copies of sl_{2n} , \mathbf{c} copies of sp_{2n} and \mathbf{d} copies of so_{2n} , then we say that \mathcal{L} is of $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ type.

Let $\mathcal{L} = \bigoplus_{k=1}^K \mathcal{L}_k$ be of type $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$, in this section we construct an epimorphism from $\mathbf{gim}(M_n)$ to \mathcal{L} .

For convenience, we fix a Chevalley generators $\{e_{\alpha_i}, f_{\alpha_i} | 1 \leq i \leq n\}$ for simple Lie algebra of type C_n or D_n , and $\{e_{\alpha_i}, f_{\alpha_i} | 1 \leq i \leq 2n-1\}$ for A_{2n-1} . The associative Dynkin diagrams are



Lemma 2.2. Let $a \in \mathbb{C}^\times, a \neq \pm 1$. If $\{e_{\alpha_i}, f_{\alpha_i} | 1 \leq i \leq 2n-1\}$ is the Chevalley generators of A_{2n-1} , let

$$\begin{aligned} e_{\alpha_{2n}} &= a[f_{\alpha_{2n-1}}, \dots [f_{\alpha_2}, f_{\alpha_1}] \dots], \\ f_{\alpha_{2n}} &= a^{-1}[\dots [e_{\alpha_1}, e_{\alpha_2}] \dots, e_{\alpha_{2n-1}}], \end{aligned}$$

then

$$e_i \mapsto e_{\alpha_i} - f_{\alpha_{n+i}}, \quad f_i \mapsto f_{\alpha_i} - e_{\alpha_{n+i}},$$

defines a Lie algebra homomorphism from $\mathbf{gim}(M_n)$ to A_{2n-1} , where $1 \leq i \leq n$.

Proof. Let

$$\begin{aligned} X_i &= e_{\alpha_i} - f_{\alpha_{n+i}}, \\ Y_i &= f_{\alpha_i} - e_{\alpha_{n+i}}, \\ H_i &= [X_i, Y_i], \end{aligned}$$

for all $1 \leq i \leq n$.

Note that the subalgebra generated by $e_{\alpha_i}, f_{\alpha_i}$ ($1 \leq i \leq 2n-1, i \neq n$) is isomorphic to $sl_n \oplus sl_n$. So the map

$$e_i \mapsto e_{\alpha_i} - f_{\alpha_{n+i}}, \quad f_i \mapsto f_{\alpha_i} - e_{\alpha_{n+i}},$$

restricted to $1 \leq i \leq n-1$ induces a diagonal injective map $\Phi_1 \oplus \Phi_2$, where

$$\begin{aligned}\Phi_1 : \quad e_i &\mapsto e_{\alpha_i}, & f_i &\mapsto f_{\alpha_i}, \\ \Phi_2 : \quad e_i &\mapsto -f_{\alpha_{n+i}}, & f_i &\mapsto -e_{\alpha_{n+i}}.\end{aligned}$$

Particularly, Φ_1 can be viewed as an inclusion and Φ_2 can be viewed the composition of Chevalley involution and inclusion.

It is sufficient to check the relations involving elements X_n, Y_n, H_n .

$$\begin{aligned}H_i &= [e_{\alpha_i} - f_{\alpha_{n+i}}, f_{\alpha_i} - e_{\alpha_{n+i}}] = h_{\alpha_i} - h_{\alpha_{n+i}}, \quad \forall 1 \leq i \leq n-1, \\ H_n &= [e_{\alpha_n} - f_{\alpha_{2n}}, f_{\alpha_n} - e_{\alpha_{2n}}] = h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}),\end{aligned}$$

where $h_{\alpha_j} = [e_{\alpha_j}, f_{\alpha_j}]$ for all $1 \leq j \leq 2n-1$. (For the computation of H_n , see Remark 1.) Then

$$\begin{aligned}[H_n, X_n] &= [h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}), e_{\alpha_n} - f_{\alpha_{2n}}] \\ &= \alpha_n(h_{\alpha_{n-1}} + 2h_{\alpha_n} + h_{\alpha_{n+1}})e_{\alpha_n} - (\alpha_1 + \cdots + \alpha_{2n-1})(h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}))f_{\alpha_{2n}} \\ &= \alpha_n(h_{\alpha_n})e_{\alpha_n} - (\alpha_1 + \cdots + \alpha_{2n-1})(h_{\alpha_1} + h_{\alpha_{2n-1}})f_{\alpha_{2n}} \\ &= 2X_n, \\ [H_n, Y_n] &= -\alpha_n(h_{\alpha_n})f_{\alpha_n} + (\alpha_1 + \cdots + \alpha_{2n-1})(h_{\alpha_1} + h_{\alpha_{2n-1}})e_{\alpha_{2n}} \\ &= -2Y_n,\end{aligned}$$

and

$$\begin{aligned}[H_n, X_1] &= \alpha_1(h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}))e_{\alpha_1} - (-\alpha_{n+1})(h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}))f_{\alpha_{n+1}} \\ &= \alpha_1(h_{\alpha_1} + h_{\alpha_2})e_{\alpha_1} + \alpha_{n+1}(2h_{\alpha_n} + h_{\alpha_{n+1}} + h_{\alpha_{n+2}})f_{\alpha_{n+1}} \\ &= X_1, \\ [H_n, Y_1] &= -\alpha_1(h_{\alpha_1} + h_{\alpha_2})f_{\alpha_1} - \alpha_{n+1}(2h_{\alpha_n} + h_{\alpha_{n+1}} + h_{\alpha_{n+2}})e_{\alpha_{n+1}} \\ &= -Y_1, \\ [H_n, X_{n-1}] &= \alpha_{n-1}(h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}))e_{\alpha_{n-1}} - (-\alpha_{2n-1})(h_{\alpha_n} + (h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}))f_{\alpha_{2n-1}} \\ &= \alpha_{n-1}(h_{\alpha_{n-2}} + h_{\alpha_{n-1}} + 2h_{\alpha_n})e_{\alpha_{n-1}} + \alpha_{2n-1}(h_{\alpha_{2n-2}} + h_{\alpha_{2n-1}})f_{\alpha_{2n-1}} \\ &= -X_{n-1}, \\ [H_n, Y_{n-1}] &= -\alpha_{n-1}(h_{\alpha_{n-2}} + h_{\alpha_{n-1}} + 2h_{\alpha_n})f_{\alpha_{n-1}} - \alpha_{2n-1}(h_{\alpha_{2n-2}} + h_{\alpha_{2n-1}})e_{\alpha_{2n-1}} \\ &= Y_{n-1}.\end{aligned}$$

Similar argument implies that

$$\begin{aligned}[H_1, X_n] &= X_n, [H_1, Y_n] = -Y_n, [H_{n-1}, X_n] = -X_n, [H_{n-1}, Y_n] = Y_n, \\ [X_1, X_n] &= [Y_1, Y_n] = [Y_{n-1}, X_n] = [X_{n-1}, Y_n] = 0, \\ [X_i, [X_i, Y_j]] &= [Y_i, [Y_i, X_j]] = 0, \text{ for } \{i, j\} = \{1, n\}, \\ [X_i, [X_i, X_j]] &= [Y_i, [Y_i, Y_j]] = 0, \text{ for } \{i, j\} = \{n-1, n\}.\end{aligned}$$

Finally, the following relation is clear:

$$[X_n, X_i] = [X_n, Y_i] = [Y_n, X_i] = [Y_n, Y_i] = 0$$

for all $2 \leq i \leq n-2$. □

Remark 1. Suppose that $\alpha, \beta, \alpha + \beta \in \Delta^+$ in A_{2n-1} and $[e_\alpha, f_\alpha] = \alpha^\vee, [e_\beta, f_\beta] = \beta^\vee$. Then it holds that $[[e_\alpha, e_\beta], [f_\beta, f_\alpha]] = \alpha^\vee + \beta^\vee$. Repeatedly using this formula, we infer that

$$[f_{\alpha_{2n}}, e_{\alpha_{2n}}] = h_{\alpha_1} + \cdots + h_{\alpha_{2n-1}}.$$

Together with that $\alpha_1 + \cdots + \alpha_{2n-1} \pm \alpha_n$ is not a root, we obtain the computation of H_n .

One can also understand it in an easy way: Let A_{2n-1} be the matrix Lie algebra sl_{2n} and $e_{\alpha_i} = E_{i,i+1}, f_{\alpha_i} = E_{i+1,i}$ for $i < 2n$. Then $e_{\alpha_{2n}} = aE_{2n,1}, f_{\alpha_{2n}} = a^{-1}E_{1,2n}$, which implies that

$$[f_{\alpha_{2n}}, e_{\alpha_{2n}}] = E_{1,1} - E_{2n,2n} = \sum_{i=1}^{2n-1} (E_{i,i} - E_{i+1,i+1}) = \sum_{i=1}^{2n-1} h_{\alpha_i}.$$

Lemma 2.3. If $\{e_{\alpha_i}, f_{\alpha_i} | 1 \leq i \leq n\}$ is the Chevalley generators of C_n , let

$$\begin{aligned} E &= [f_{\alpha_{n-1}}, \cdots [f_{\alpha_1}, f_{\alpha_n}] \cdots], \\ F &= [\cdots [e_{\alpha_n}, e_{\alpha_1}] \cdots, e_{\alpha_{n-1}}], \end{aligned}$$

then

$$\begin{aligned} e_i &\mapsto e_{\alpha_i}, & f_i &\mapsto f_{\alpha_i}, & 1 \leq i \leq n-1, \\ e_n &\mapsto E, & f_n &\mapsto F, \end{aligned}$$

defines a Lie algebra homomorphism from $\mathfrak{gim}(M_n)$ to C_n .

Proof. It is sufficient to check the relation involving elements E, F . Actually, F (respectively E) is a root vector of highest short root (respectively lowest short root). Then

$$\begin{aligned} [F, e_{\alpha_i}] &= [E, f_{\alpha_i}] = 0 & \forall 2 \leq i \leq n-1, \\ [F, f_{\alpha_i}] &= [E, e_{\alpha_i}] = 0 & \forall 1 \leq i \leq n-2, \end{aligned}$$

and

$$\begin{aligned} [F, [F, e_{\alpha_1}]] &= [e_{\alpha_1}, [e_{\alpha_1}, F]] = 0, & [E, [E, f_{\alpha_1}]] &= [f_{\alpha_1}, [f_{\alpha_1}, E]] = 0, \\ [F, [F, f_{\alpha_{n-1}}]] &= [f_{\alpha_{n-1}}, [f_{\alpha_{n-1}}, F]] = 0, & [E, [E, e_{\alpha_{n-1}}]] &= [e_{\alpha_{n-1}}, [e_{\alpha_{n-1}}, E]] = 0. \end{aligned}$$

Moreover, $H := [E, F] = -(2h_{\alpha_n} + h_{\alpha_1} + \cdots + h_{\alpha_{n-1}})$, where $h_{\alpha_i} = [e_{\alpha_i}, f_{\alpha_i}]$ for all $1 \leq i \leq n$. Hence, we have

$$\begin{aligned} [H, E] &= (-\alpha_1 - \cdots - \alpha_n)(-(2h_{\alpha_n} + h_{\alpha_1} + \cdots + h_{\alpha_{n-1}}))E = 2E, \\ [H, F] &= (\alpha_1 + \cdots + \alpha_n)(-(2h_{\alpha_n} + h_{\alpha_1} + \cdots + h_{\alpha_{n-1}}))F = -2F, \end{aligned}$$

and

$$\begin{aligned} [H, e_{\alpha_1}] &= e_{\alpha_1}, & [H, f_{\alpha_1}] &= -f_{\alpha_1}, \\ [H, e_{\alpha_{n-1}}] &= -e_{\alpha_{n-1}}, & [H, f_{\alpha_{n-1}}] &= f_{\alpha_{n-1}}, \\ [h_{\alpha_1}, E] &= E, & [h_{\alpha_{n-1}}, E] &= -E, \\ [h_{\alpha_1}, F] &= -F, & [h_{\alpha_{n-1}}, F] &= F. \end{aligned}$$

The above calculation implies our statement and the proof is completed. \square

Lemma 2.4. *If $\{e_{\alpha_i}, f_{\alpha_i} | 1 \leq i \leq n\}$ is the Chevalley generators of D_n , let*

$$\begin{aligned} E &= [f_{\alpha_{n-1}}, \dots [f_{\alpha_2}, f_{\alpha_n}] \dots], \\ F &= [\dots [e_{\alpha_n}, e_{\alpha_2}] \dots, e_{\alpha_{n-1}}], \end{aligned}$$

then

$$\begin{aligned} e_i &\mapsto e_{\alpha_i}, & f_i &\mapsto f_{\alpha_i}, & 1 \leq i \leq n-1, \\ e_n &\mapsto E, & f_n &\mapsto F, \end{aligned}$$

defines a Lie algebra homomorphism from $\mathfrak{gim}(M_n)$ to D_n .

Proof. It is sufficient to check the relation involving elements E, F . Actually, F (respectively E) is a root vector of highest root (respectively lowest root) of subalgebra generated by $e_{\alpha_i}, f_{\alpha_i}$ ($2 \leq i \leq n$). Then

$$\begin{aligned} [F, e_{\alpha_i}] &= [E, f_{\alpha_i}] = 0 & \forall 2 \leq i \leq n-1, \\ [F, f_{\alpha_i}] &= [E, e_{\alpha_i}] = 0 & \forall 2 \leq i \leq n-2, \end{aligned}$$

and

$$[F, [F, f_{\alpha_{n-1}}]] = [f_{\alpha_{n-1}}, [f_{\alpha_{n-1}}, F]] = 0, \quad [E, [E, e_{\alpha_{n-1}}]] = [e_{\alpha_{n-1}}, [e_{\alpha_{n-1}}, E]] = 0.$$

Notice that $\alpha_1 + \dots + \alpha_n$ is a root, and neither of $-\alpha_1 + \alpha_2 + \dots + \alpha_n$, $2\alpha_1 + \dots + \alpha_n$ and $\alpha_1 + 2(\alpha_2 + \dots + \alpha_n)$ is a root, hence we have

$$\begin{aligned} [F, [F, e_{\alpha_1}]] &= [e_{\alpha_1}, [e_{\alpha_1}, F]] = 0, & [E, [E, f_{\alpha_1}]] &= [f_{\alpha_1}, [f_{\alpha_1}, E]] = 0, \\ [F, f_{\alpha_1}] &= 0, & [E, e_{\alpha_1}] &= 0. \end{aligned}$$

Moreover, $H := [E, F] = -(h_{\alpha_2} + \dots + h_{\alpha_n})$, where $h_{\alpha_i} = [e_{\alpha_i}, f_{\alpha_i}]$ for all $1 \leq i \leq n$. Hence, we have

$$\begin{aligned} [H, E] &= (-\alpha_2 - \dots - \alpha_n)(-(h_{\alpha_2} + \dots + h_{\alpha_n}))E = 2E, \\ [H, F] &= (\alpha_2 + \dots + \alpha_n)(-(h_{\alpha_2} + \dots + h_{\alpha_n}))F = -2F, \end{aligned}$$

and

$$\begin{aligned} [H, e_{\alpha_1}] &= e_{\alpha_1}, & [H, f_{\alpha_1}] &= -f_{\alpha_1}, \\ [H, e_{\alpha_{n-1}}] &= -e_{\alpha_{n-1}}, & [H, f_{\alpha_{n-1}}] &= f_{\alpha_{n-1}}, \\ [h_{\alpha_1}, E] &= E, & [h_{\alpha_{n-1}}, E] &= -E, \\ [h_{\alpha_1}, F] &= -F, & [h_{\alpha_{n-1}}, F] &= F. \end{aligned}$$

The above calculation implies our statement and the proof is completed. \square

Next we begin to explicitly construct an epimorphism from $\mathfrak{gim}(M_n)$ to \mathcal{L} . The construction is divided into four distinguish cases.

Case 1. $\mathcal{L}_k \cong sl_{2n}$ for all $1 \leq k \leq K$.

Let $\{e_{\alpha_i}^{[k]}, f_{\alpha_i}^{[k]} | 1 \leq i \leq 2n-1\}$ be the analogue of Chevalley generators of A_{2n-1} in \mathcal{L}_k for all k . Choose a K -tuple $\underline{a} := (a_1, \dots, a_K) \in (\mathbb{C}^\times)^K$ such that $a_k \neq \pm 1$ and $a_k \neq a_j^{\pm 1}$ for all $k \neq j$.

Set

$$\begin{aligned} e_{\alpha_{2n}}^{[k]} &= a_k [f_{\alpha_{2n-1}}^{[k]}, \dots [f_{\alpha_2}^{[k]}, f_{\alpha_1}^{[k]}] \dots], \\ f_{\alpha_{2n}}^{[k]} &= a_k^{-1} [\dots [e_{\alpha_1}^{[k]}, e_{\alpha_2}^{[k]}] \dots, e_{\alpha_{2n-1}}^{[k]}], \end{aligned}$$

for all $1 \leq k \leq K$, where $e_{\alpha_{2n}}^{[k]}$ and $f_{\alpha_{2n}}^{[k]}$ are root vectors of lowest root and highest root, respectively.

Case 2. $\mathcal{L}_1 \cong sp_{2n}$ and $\mathcal{L}_k \cong sl_{2n}$ for all $2 \leq k \leq K$.

Let $\{e_{\alpha_i}^{[1]}, f_{\alpha_i}^{[1]} | 1 \leq i \leq n\}$ be the analogue of Chevalley generators of C_n in \mathcal{L}_1 . and $\{e_{\alpha_i}^{[k]}, f_{\alpha_i}^{[k]} | 1 \leq i \leq 2n-1\}$ be the analogue of Chevalley generators of A_{2n-1} in \mathcal{L}_k for $k \geq 2$. Choose a K -tuple $\underline{a} := (1, a_2, \dots, a_K) \in (\mathbb{C}^\times)^K$ such that $a_k \neq \pm 1$ and $a_k \neq a_j^{\pm 1}$ for all $k \neq j$. Set

$$\begin{aligned} e_{\alpha_{2n}}^{[k]} &= a_k [f_{\alpha_{2n-1}}^{[k]}, \dots [f_{\alpha_2}^{[k]}, f_{\alpha_1}^{[k]}] \dots], \\ f_{\alpha_{2n}}^{[k]} &= a_k^{-1} [\dots [e_{\alpha_1}^{[k]}, e_{\alpha_2}^{[k]}] \dots, e_{\alpha_{2n-1}}^{[k]}], \end{aligned}$$

for all $2 \leq k \leq K$, and

$$\begin{aligned} e_{\alpha_{2n}}^{[1]} &= f_{\alpha_n}^{[1]} - [\dots [[e_{\alpha_n}^{[1]}, e_{\alpha_1}^{[1]}], e_{\alpha_2}^{[1]}] \dots, e_{\alpha_{n-1}}^{[1]}], \\ f_{\alpha_{2n}}^{[1]} &= e_{\alpha_n}^{[1]} - [f_{\alpha_{n-1}}^{[1]}, \dots [f_{\alpha_2}^{[1]}, [f_{\alpha_1}^{[1]}, f_{\alpha_n}^{[1]}] \dots], \\ e_{\alpha_{n+1}}^{[1]} &= \dots = e_{\alpha_{2n-1}}^{[1]} = 0, \\ f_{\alpha_{n+1}}^{[1]} &= \dots = f_{\alpha_{2n-1}}^{[1]} = 0, \end{aligned}$$

where $f_{\alpha_n}^{[1]} - e_{\alpha_{2n}}^{[1]}$ and $e_{\alpha_n}^{[1]} - f_{\alpha_{2n}}^{[1]}$ are root vectors of highest short root and lowest short root, respectively.

Case 3. $\mathcal{L}_1 \cong so_{2n}$ and $\mathcal{L}_k \cong sl_{2n}$ for all $2 \leq k \leq K$.

Let $\{e_{\alpha_i}^{[1]}, f_{\alpha_i}^{[1]} | 1 \leq i \leq n\}$ be the analogue of Chevalley generators of C_n in \mathcal{L}_1 . and $\{e_{\alpha_i}^{[k]}, f_{\alpha_i}^{[k]} | 1 \leq i \leq 2n-1\}$ be the analogue of Chevalley generators of A_{2n-1} in \mathcal{L}_k for $k \geq 2$. Choose a K -tuple $\underline{a} := (-1, a_2, \dots, a_K) \in (\mathbb{C}^\times)^K$ such that $a_k \neq \pm 1$ and $a_k \neq a_j^{\pm 1}$ for all $k \neq j$. Set

$$\begin{aligned} e_{\alpha_{2n}}^{[k]} &= a_k [f_{\alpha_{2n-1}}^{[k]}, \dots [f_{\alpha_2}^{[k]}, f_{\alpha_1}^{[k]}] \dots], \\ f_{\alpha_{2n}}^{[k]} &= a_k^{-1} [\dots [e_{\alpha_1}^{[k]}, e_{\alpha_2}^{[k]}] \dots, e_{\alpha_{2n-1}}^{[k]}], \end{aligned}$$

for all $2 \leq k \leq K$, and

$$\begin{aligned} e_{\alpha_{2n}}^{[1]} &= f_{\alpha_n}^{[1]} - [\dots [e_{\alpha_n}^{[1]}, e_{\alpha_2}^{[1]}] \dots, e_{\alpha_{n-1}}^{[1]}], \\ f_{\alpha_{2n}}^{[1]} &= e_{\alpha_n}^{[1]} - [f_{\alpha_{n-1}}^{[1]}, \dots [f_{\alpha_2}^{[1]}, f_{\alpha_n}^{[1]}] \dots], \\ e_{\alpha_{n+1}}^{[1]} &= \dots = e_{\alpha_{2n-1}}^{[1]} = 0, \\ f_{\alpha_{n+1}}^{[1]} &= \dots = f_{\alpha_{2n-1}}^{[1]} = 0. \end{aligned}$$

Case 4. $\mathcal{L}_1 \cong so_{2n}$, $\mathcal{L}_2 \cong sp_{2n}$ and $\mathcal{L}_k \cong sl_{2n}$ for all $3 \leq k \leq K$.

Let $\{e_{\alpha_i}^{[1]}, f_{\alpha_i}^{[1]} | 1 \leq i \leq n\}$ be the analogue of Chevalley generators of D_n in \mathcal{L}_1 , $\{e_{\alpha_i}^{[2]}, f_{\alpha_i}^{[2]} | 1 \leq i \leq n\}$ be the analogue of Chevalley generators of C_n in \mathcal{L}_2 , and $\{e_{\alpha_i}^{[k]}, f_{\alpha_i}^{[k]} | 1 \leq i \leq 2n-1\}$ be the analogue of Chevalley generators of A_{2n-1} in \mathcal{L}_k for $k \geq 3$. Choose a K -tuple $\underline{a} := (-1, 1, a_3, \dots, a_K) \in (\mathbb{C}^\times)^K$ such that $a_k \neq \pm 1$ and $a_k \neq a_j^{\pm 1}$ for all $k \neq j$. Set

$$\begin{aligned} e_{\alpha_{2n}}^{[k]} &= a_k [f_{\alpha_{2n-1}}^{[k]}, \dots [f_{\alpha_2}^{[k]}, f_{\alpha_1}^{[k]}] \dots], \\ f_{\alpha_{2n}}^{[k]} &= a_k^{-1} [\dots [e_{\alpha_1}^{[k]}, e_{\alpha_2}^{[k]}] \dots, e_{\alpha_{2n-1}}^{[k]}], \end{aligned}$$

for all $3 \leq k \leq K$, and

$$\begin{aligned}
e_{\alpha_{2n}}^{[1]} &= f_{\alpha_n}^{[1]} - [\cdots [e_{\alpha_n}^{[1]}, e_{\alpha_2}^{[1]}] \cdots, e_{\alpha_{n-1}}^{[1]}], \\
f_{\alpha_{2n}}^{[1]} &= e_{\alpha_n}^{[1]} - [f_{\alpha_{n-1}}^{[1]}, \cdots [f_{\alpha_2}^{[1]}, f_{\alpha_n}^{[1]}] \cdots], \\
e_{\alpha_{n+1}}^{[1]} &= \cdots = e_{\alpha_{2n-1}}^{[1]} = 0, \\
f_{\alpha_{n+1}}^{[1]} &= \cdots = f_{\alpha_{2n-1}}^{[1]} = 0, \\
e_{\alpha_{2n}}^{[2]} &= f_{\alpha_n}^{[2]} - [\cdots [[e_{\alpha_n}^{[2]}, e_{\alpha_1}^{[2]}], e_{\alpha_2}^{[2]}] \cdots, e_{\alpha_{n-1}}^{[2]}], \\
f_{\alpha_{2n}}^{[2]} &= e_{\alpha_n}^{[2]} - [f_{\alpha_{n-1}}^{[2]}, \cdots [f_{\alpha_2}^{[2]}, [f_{\alpha_1}^{[2]}, f_{\alpha_n}^{[2]}] \cdots], \\
e_{\alpha_{n+1}}^{[2]} &= \cdots = e_{\alpha_{2n-1}}^{[2]} = 0, \\
f_{\alpha_{n+1}}^{[2]} &= \cdots = f_{\alpha_{2n-1}}^{[2]} = 0.
\end{aligned}$$

In all above four cases, we define a homomorphism from $\mathbf{gim}(M_n) \rightarrow \mathcal{L}$ via

$$\Psi_{\underline{a}}: \quad e_i \mapsto \sum_{k=1}^K (e_{\alpha_i}^{[k]} - f_{\alpha_{n+i}}^{[k]}), \quad f_i \mapsto \sum_{k=1}^K (f_{\alpha_i}^{[k]} - e_{\alpha_{n+i}}^{[k]}),$$

for all $1 \leq i \leq n$.

Proposition 2.5. $\Psi_{\underline{a}}$ is a Lie algebra epimorphism.

Proof. Let P_k be the projection from \mathcal{L} to \mathcal{L}_k . By Lemmas 2.2-2.4, we infer that $P_k \circ \Psi_{\underline{a}}$ is a homomorphism for all $1 \leq k \leq K$, then $\Psi_{\underline{a}} = \oplus_{k=1}^K P_k \circ \Psi_{\underline{a}}$ is a homomorphism of Lie algebras. The detailed proof for that $\Psi_{\underline{a}}$ is an epimorphism, is very similar to the proof for Lemmas 5.1-5.4 in Section 5. \square

3. MAIN RESULTS

Our main result can be stated by the following three theorems.

Theorem 3.1. *If \mathcal{L} is of type $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$, then there exists an ideal I of $\mathbf{gim}(M_n)$, such that*

$$\mathbf{gim}(M_n)/I \cong \mathcal{L}.$$

Theorem 3.2. *If there exists an ideal I of $\mathbf{gim}(M_n)$, such that $\mathbf{gim}(M_n)/I$ is finite-dimensional and semi-simple, then $\mathbf{gim}(M_n)/I$ is of type $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$.*

The above two theorems will be proved in below sections.

Theorem 3.3. (1) *If V is an irreducible finite-dimensional module of an $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ type Lie algebra \mathcal{L} , then V is an irreducible $\mathbf{gim}(M_n)$ -module.*

(2) *If V is an irreducible $\mathbf{gim}(M_n)$ -module with finite dimension, then V is an irreducible module of some $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ type Lie algebra.*

Proof. (1) Suppose that V is an irreducible finite-dimensional module of $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ type Lie algebra \mathcal{L} , and that $\zeta : \mathcal{L} \rightarrow \text{End}(V)$ is the representation map. By Proposition 2.5, V is an irreducible $\mathbf{gim}(M_n)$ -module with representation map $\zeta \circ \Psi_{\underline{a}}$.

(2) Suppose that V is non-trivial and the representation is define by $\kappa : \mathbf{gim}(M_n) \rightarrow \text{End}(V)$ and the image of κ has Levi decomposition

$$\text{Im}(\kappa) = S \dot{+} H \dot{+} W,$$

where S is semi-simple, H is diagonal and W is nilpotent. Undoubtedly, it holds that $[S, H] = 0$.

Because V is non-trivial and there exists a positive integer k such that W^k trivially acts on V , we infer that $W \cdot V$ is a proper submodule and we may assume that $W = 0$.

Suppose that \mathfrak{h} is a Cartan subalgebra of S . Then V has a basis such that both $\kappa(\mathfrak{h})$ and $\kappa(H)$ are diagonal, then any S -module must be an H -module. So V is irreducible as $\mathbf{gim}(M_n)$ -module if and only if V is irreducible as S -module. By Theorem 3.2, S has to be isomorphic to some $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$ type Lie algebra, and thus the statement holds. \square

4. EVALUATION REPRESENTATIONS OF $A_{2n-1}^{(1)}$

Let \mathfrak{g} be the simple Lie algebra $sl_{2n}(\mathbb{C})$. Then the affine Lie algebra of type $A_{2n-1}^{(1)}$ has a realization:

$$\hat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c,$$

with bracket:

$$[c, \hat{\mathfrak{g}}] = 0, \quad [x \otimes t^m, y \otimes t^k] = [x, y] \otimes t^{m+k} + m\delta_{m,-k}(x, y)c,$$

where $x, y \in \mathfrak{g}, m, k \in \mathbb{Z}$ and $(,)$ is a non-degenerate invariant form on \mathfrak{g} , which is a scalar of the Killing form.

The following is the Dynkin diagram of \mathfrak{g} :

$$1 \circ \text{---} 2 \circ \text{---} 3 \circ \text{---} \cdots \text{---} \circ \text{---} \circ^{2n-1}$$

Let $\Pi = \{\alpha_1, \dots, \alpha_{2n-1}\}$ be the associated prime root system, and $\dot{\Delta}$ be its root system. Set $\{E_\alpha, H_i | \alpha \in \dot{\Delta}, 1 \leq i \leq 2n-1\}$ be the Chevalley basis of \mathfrak{g} . Then elements $E_{\pm\alpha_i} \otimes 1 (1 \leq i \leq 2n-1)$ and $E_{\pm\alpha^b} \otimes t^\mp$ are Chevalley generators of $\hat{\mathfrak{g}}$, where $\alpha^b = \alpha_1 + \dots + \alpha_{2n-1}$.

Set

$$\begin{aligned} \hat{e}_1 &= E_{\alpha_1} \otimes 1 - E_{-\alpha_{n+1}} \otimes 1, & \hat{f}_1 &= E_{-\alpha_1} \otimes 1 - E_{\alpha_{n+1}} \otimes 1, & \hat{h}_1 &= H_1 \otimes 1 - H_{n+1} \otimes 1, \\ \hat{e}_2 &= E_{\alpha_2} \otimes 1 - E_{-\alpha_{n+2}} \otimes 1, & \hat{f}_2 &= E_{-\alpha_2} \otimes 1 - E_{\alpha_{n+2}} \otimes 1, & \hat{h}_2 &= H_2 \otimes 1 - H_{n+2} \otimes 1, \\ & \dots \end{aligned}$$

$$\hat{e}_n = E_{\alpha_n} \otimes 1 + E_{\alpha^b} \otimes t^{-1}, \quad \hat{f}_n = E_{-\alpha_n} \otimes 1 + E_{-\alpha^b} \otimes t, \quad \hat{h}_n = H_n \otimes 1 + (H_1 + \dots + H_{2n-1}) \otimes 1 - c,$$

and let $\hat{\mathfrak{g}}_{fp}$ be the subalgebra generated by $\{\hat{e}_i, \hat{f}_i | 1 \leq i \leq n\}$. $\hat{\mathfrak{g}}_{fp}$ is called a fixed point subalgebra of $\hat{\mathfrak{g}}$.

Proposition 4.1. *There exists an algebra homomorphism $\phi : \mathcal{U}(\mathbf{gim}(M_n)) \rightarrow \mathcal{U}(\hat{\mathfrak{g}}_{fp})$ via the action on generators:*

$$\phi(e_i) = \hat{e}_i, \quad \phi(f_i) = \hat{f}_i, \quad 1 \leq i \leq n.$$

More over, it induces a Lie algebra homomorphism (also denoted by ϕ) from $\mathbf{gim}(M_n)$ to $\hat{\mathfrak{g}}_{fp}$.

Proof. This is a consequence of the following equations:

$$\begin{aligned}
[\hat{e}_i, \hat{f}_i] &= \hat{h}_i, & i &= 1, \dots, n, \\
[\hat{h}_i, \hat{e}_j] &= m_{i,j} \hat{e}_j, [\hat{h}_i, \hat{f}_j] = -m_{i,j} \hat{f}_j, & 1 \leq i, j \leq n, \\
[\hat{e}_i, \hat{f}_j] &= 0, & i \neq j, \{i, j\} \neq \{1, n\}, \\
[\hat{e}_i, [\hat{e}_i, \hat{e}_j]] &= 0, [\hat{f}_i, [\hat{f}_i, \hat{f}_j]] = 0, & |i - j| = 1, \\
[\hat{e}_i, \hat{e}_j] &= 0, [\hat{f}_i, \hat{f}_j] = 0, & |i - j| > 1, \\
[\hat{e}_i, [\hat{e}_i, \hat{f}_j]] &= [\hat{f}_i, [\hat{f}_i, \hat{e}_j]] = 0, & \{i, j\} = \{1, n\}.
\end{aligned}$$

□

Remark 2. In fact, the map ϕ is an isomorphism (see [Br]). More over, if we set

$$\hat{e}_n = E_{\alpha_n} \otimes 1 - E_{\alpha^\flat} \otimes t^{-1}, \quad \hat{f}_n = E_{-\alpha_n} \otimes 1 - E_{-\alpha^\flat} \otimes t,$$

then Proposition 4.1 still holds. However, these two maps are equivalent under the automorphism of \hat{sl}_{2n} induced by $x \otimes t^m \mapsto x \otimes (-t)^m$.

Let $a \in \mathbb{C}^\times$, an irreducible evaluation $\hat{\mathfrak{g}}$ -module $V_{\lambda,a}$ is an irreducible highest weight \mathfrak{g} -module with highest weight λ , and the action is defined by:

$$(x \otimes t^m) \cdot v = a^m x \cdot v, \quad c \cdot v = 0.$$

for all $x \in \mathfrak{g}, m \in \mathbb{Z}$ and $v \in V_{\lambda,a}$.

For any $V_{\lambda,a}$, the map ϕ induces a representation for $\mathbf{gim}(M_n)$. Particularly, for any dominate weight λ , $V_{\lambda,a}$ is finite-dimensional.

Let $V_{\lambda,a}$ be the $2n$ -dimensional irreducible module, i.e., the natural representation of \mathfrak{g} . Particularly, we may assume that $\mathfrak{g} = sl_{2n}$. The following proposition is an important tool of the proof for our main results.

Proposition 4.2. The following map induces a representation for $\mathbf{gim}(M_n)$:

$$\begin{aligned}
\psi_a : \mathbf{gim}(M_n) &\rightarrow sl_{2n} \\
\psi_a(e_1) &= E_{1,2} - E_{n+2,n+1}, \quad \psi_a(f_1) = E_{2,1} - E_{n+1,n+2} \\
\psi_a(e_2) &= E_{2,3} - E_{n+3,n+2}, \quad \psi_a(f_2) = E_{3,2} - E_{n+2,n+3} \\
&\dots \\
\psi_a(e_{n-1}) &= E_{n-1,n} - E_{2n,2n-1}, \quad \psi_a(f_{n-1}) = E_{n,n-1} - E_{2n-1,2n} \\
\psi_a(e_n) &= E_{n,n+1} + a^{-1} E_{1,2n}, \quad \psi_a(f_n) = E_{n+1,n} + a E_{2n,1}.
\end{aligned}$$

Particularly, if $a \neq \pm 1$, the image of ψ_a is sl_{2n} , if $a = 1$, the image of ψ_a is sp_{2n} , and if $a = -1$, then the image of ψ_a is so_{2n} .

Remark 3. In case of assuming that

$$\hat{e}_n = E_{\alpha_n} \otimes 1 - E_{\alpha^\flat} \otimes t^{-1}, \quad \hat{f}_n = E_{-\alpha_n} \otimes 1 - E_{-\alpha^\flat} \otimes t,$$

Proposition 4.2 still holds after exchanging the positions of results when $a = 1$ and $a = -1$, respectively.

Proof. **Proof for Proposition 4.2.**

Set $x_i = \psi_a(e_i)$, $y_i = \psi_a(f_i)$ for $1 \leq i \leq n$, and $h_i = E_{i,i} - E_{i+1,i+1} - E_{i+n,i+n} + E_{i+1+n,i+1+n}$ for $1 \leq i \leq n-1$. Then it is easy to know that the elements $x_i, y_i (1 \leq i \leq n-1)$ generate the Lie algebra

$$S = \left\{ X = \begin{bmatrix} A & 0 \\ 0 & -A^t \end{bmatrix} \mid A \in sl_n \right\},$$

which is isomorphic to sl_n .

More over,

$$\begin{aligned} [x_{n-1}, x_n] &= E_{n-1,n+1} + a^{-1}E_{1,2n-1}, \\ [x_{n-2}, [x_{n-1}, x_n]] &= E_{n-2,n+1} + a^{-1}E_{1,2n-2}, \\ [x_{n-3}, [x_{n-2}, [x_{n-1}, x_n]]] &= E_{n-3,n+1} + a^{-1}E_{1,2n-3}, \\ &\dots \quad \dots \quad \dots \\ [x_2 \cdots, [x_{n-1}, x_n] \cdots] &= E_{2,n+1} + a^{-1}E_{1,n+2}, \\ [x_1, [x_2 \cdots, [x_{n-1}, x_n] \cdots]] &= (1 + a^{-1})E_{1,n+1}, \\ \\ [y_n, y_{n-1}] &= E_{n+1,n-1} + aE_{2n-1,1}, \\ [[y_n, y_{n-1}], y_{n-2}] &= E_{n+1,n-2} + aE_{2n-2,1}, \\ [[[y_n, y_{n-1}], y_{n-2}], y_{n-3}] &= E_{n+1,n-3} + aE_{2n-3,1}, \\ &\dots \quad \dots \quad \dots \\ [\cdots [y_n, y_{n-1}], \cdots y_2] &= E_{n+1,2} + aE_{n+2,1}, \\ [[\cdots [y_n, y_{n-1}], \cdots y_2], y_1] &= (1 + a)E_{n+1,1}, \end{aligned}$$

then we divide the proof into three cases according to the value of a .

(1) *Case 1.* $a \neq \pm 1$.

By equations

$$\begin{aligned} [x_1, [x_2 \cdots, [x_{n-1}, x_n] \cdots]] &= (1 + a^{-1})E_{1,n+1}, \\ [[\cdots [y_n, y_{n-1}], \cdots y_2], y_1] &= (1 + a)E_{n+1,1}, \end{aligned}$$

we have $E_{1,n+1}, E_{n+1,1} \in Im(\psi_a)$, and hence, for every $1 \leq i \leq n-1$,

$$\begin{aligned} 2E_{i+1,n+i+1} &= [y_i, [y_i, E_{i,n+i}]], \\ 2E_{n+i+1,i+1} &= [x_i, [x_i, E_{n+i,i}]], \end{aligned}$$

thus it holds that

$$E_{i,n+i}, E_{n+i,i} \in Im(\psi_a), \quad \forall 1 \leq i \leq n.$$

Set

$$\mathbf{x}_k = E_{n+k,k} \quad \mathbf{y}_k = E_{k,n+k},$$

then

$$[x_n, \mathbf{x}_n] = E_{2n,n+1} - a^{-1}E_{1,n}, \quad [y_n, \mathbf{y}_n] = E_{n+1,2n} - aE_{n,1}$$

this implies that $E_{2n,n+1}, E_{n+1,2n} \in Im(\psi_a)$, and hence, $E_{1,n}, E_{n,1} \in Im(\psi_a)$.

Further, the iteration formula

$$[y_i, E_{i,n}] = E_{i+1,n}, \quad [E_{n,i}, x_i] = E_{n,i+1},$$

implies that $E_{n,i}, E_{i,n} \in \text{Im}(\psi_a)$ for all $i \leq n$, and thus $E_{n+i,2n}, E_{2n,n+i} \in \text{Im}(\psi_a)$ for all $i \leq n$.

We also have iteration formula

$$\begin{aligned} [E_{i,n}, y_{n-1}] &= E_{i,n-1}, [x_{n-1}, E_{n,i}] = E_{n-1,i}, \\ [E_{i,k+1}, y_k] &= E_{i,k}, [x_k, E_{k+1,i}] = E_{k,i}, \quad \text{if } i < k, \end{aligned}$$

and this implies that $E_{i,j}, E_{j,i} \in \text{Im}(\psi_a)$ for all $i < j \leq n$. Finally, we infer that $E_{i,j}, E_{i+n,j+n} \in \text{Im}(\psi_a)$ for all $1 \leq i, j \leq n$ and $i \neq j$. Notice that

$$[[E_{2,n}, E_{n,2}], x_n] = -E_{n,n+1}, \quad [[E_{2,n}, E_{n,2}], y_n] = E_{n+1,n},$$

so we have $E_{i,i+1}, E_{i+1,i} \in \text{Im}(\psi_a)$ for all $1 \leq i \leq 2n-1$, this is a set of Chevalley generators of sl_{2n} , and then $\text{Im}(\psi_a) = sl_{2n}$.

(2) *Case 2.* $a = 1$.

It is easy to check that

$$\begin{aligned} [\mathbf{x}_1, y_i] &= [\mathbf{y}_1, x_i] = 0 & i = 1, \dots, n-1, \\ [\mathbf{x}_1, x_i] &= [\mathbf{y}_1, y_i] = 0 & i = 2, \dots, n-1, \\ [[\mathbf{x}_1, x_1], x_1] &= 2E_{n+2,2} = 2\mathbf{x}_2, & [[\mathbf{y}_1, y_1], y_1] = 2E_{2,n+2} = 2\mathbf{y}_2, \\ [\mathbf{x}_2, x_1] &= [\mathbf{y}_2, y_1] = 0, & [\mathbf{x}_1, [\mathbf{x}_1, x_1]] = [\mathbf{y}_1, [\mathbf{y}_1, y_1]] = 0, \end{aligned}$$

then the Lie algebra generated by S and $\mathbf{x}_1, \mathbf{y}_1$ is isomorphic to sp_{2n} . In particular, $\{x_i, y_i, \mathbf{x}_1, \mathbf{y}_1 | 1 \leq i \leq n-1\}$ is a set of Chevalley generators. The associated Dynkin diagram is

$$n \circ \Longrightarrow 1 \circ \text{ --- } 2 \circ \text{ --- } 3 \circ \text{ --- } \dots \text{ --- } \circ \text{ --- } \circ^{n-1}$$

Notice that $x_i, y_i \in sp_{2n}$ for all $1 \leq i \leq n$, we have

$$\langle S, \mathbf{x}_1, \mathbf{y}_1 \rangle \subset \text{Im}(\psi_a) = \langle x_i, y_i | 1 \leq i \leq n \rangle \subset sp_{2n},$$

the dimension relation $\dim \langle S, \mathbf{x}_1, \mathbf{y}_1 \rangle = \dim sp_{2n}$ implies that $\text{Im}(\psi_a) = sp_{2n}$.

(3) *Case 3.* $a = -1$. Then x_n, y_n have the form $\begin{bmatrix} 0 & B \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & 0 \\ B & 0 \end{bmatrix}$, respectively, and

where $B = -B^t \in gl_n$ is anti-symmetric.

Set

$$\begin{aligned} x^\sharp &= [\dots [y_n, y_{n-1}], \dots y_2] = E_{n+1,2} - E_{n+2,1}, \\ y^\sharp &= [x_2, \dots [x_{n-1}, x_n] \dots] = E_{2,n+1} - E_{1,n+2}, \end{aligned}$$

we have

$$\begin{aligned} [\mathbf{x}^\sharp, y_i] &= [\mathbf{y}^\sharp, x_i] = 0, & i = 1, \dots, n-1, \\ [\mathbf{x}^\sharp, x_i] &= [\mathbf{y}^\sharp, y_i] = 0, & i = 1, 3, \dots, n-1, \end{aligned}$$

and the set $\{\mathbf{x}^\sharp, \mathbf{y}^\sharp, x_2, y_2\}$ generates a simple Lie algebra of type A_2 .

Then the Lie algebra generated by S and $\mathbf{x}^\sharp, \mathbf{y}^\sharp$ is isomorphic to so_{2n} . In particular, $\{x_i, y_i, \mathbf{x}^\sharp, \mathbf{y}^\sharp | 1 \leq i \leq n-1\}$ is a set of Chevalley generators. Notice that $x_i, y_i \in so_{2n}$ for all $1 \leq i \leq n$, we have

$$\langle S, \mathbf{x}^\sharp, \mathbf{y}^\sharp \rangle \subset \text{Im}(\psi_a) = \langle x_i, y_i | 1 \leq i \leq n \rangle \subset so_{2n},$$

the dimension relation $\dim\langle S, \mathbf{x}^\sharp, \mathbf{y}^\sharp \rangle = \dim so_{2n}$ implies that $Im(\psi_a) = so_{2n}$. \square

5. PROOF FOR THEOREM 3.1

Let $\mathcal{A} = \bigoplus_{k=1}^K \mathfrak{g}_i = \mathfrak{g}^{\oplus n}$ be the direct sum of K copies of $\mathfrak{g} = sl_{2n}$. Fix a K -tuple $\underline{a} = (a_1, \dots, a_K) \in (\mathbb{C}^\times)^K$ such that $a_k \neq a_j^{\pm 1}$ for all $k \neq j$. Define the map

$$\psi_{\underline{a}} = \bigoplus_{k=1}^K \psi_{a_k} : \quad \mathbf{gim}(M_n) \rightarrow \mathcal{A},$$

where $\psi_{a_k} : \mathbf{gim}(M_n) \rightarrow \mathfrak{g}_k$ is an analogue of the evaluation map defined in Proposition 4.2..

Theorem 3.1 can be proved by the following Lemmas 5.1-5.4.

Lemma 5.1. *If $a_k \neq \pm 1$ for every k , then $\psi_{\underline{a}}$ is an epimorphism.*

Proof. For convenience, we let $\psi(x)^{[k]}$ denote the image of x under ψ_{a_k} . By the definition of ψ , we have that

$$\begin{aligned} \psi_{a_k}(e_1) &= (E_{1,2} - E_{n+2,n+1})^{[k]}, & \psi_{a_k}(f_1) &= (E_{2,1} - E_{n+1,n+2})^{[k]} \\ \psi_{a_k}(e_2) &= (E_{2,3} - E_{n+3,n+2})^{[k]}, & \psi_{a_k}(f_2) &= (E_{3,2} - E_{n+2,n+3})^{[k]} \\ &\dots & & \\ \psi_{a_k}(e_{n-1}) &= (E_{n-1,n} - E_{2n,2n-1})^{[k]}, & \psi_{a_k}(f_{n-1}) &= (E_{n,n-1} - E_{2n-1,2n})^{[k]} \\ \psi_{a_k}(e_n) &= (E_{n,n+1} + a_k^{-1} E_{1,2n})^{[k]}, & \psi_{a_k}(f_n) &= (E_{n+1,n} + a_k E_{2n,1})^{[k]}. \end{aligned}$$

Set

$$x_i^{[k]} = \psi_{a_k}(e_i), \quad y_i^{[k]} = \psi_{a_k}(f_i).$$

Then

$$\begin{aligned} [x_{n-1}^{[k]}, x_n^{[k]}] &= (E_{n-1,n+1} + a_k^{-1} E_{1,2n-1})^{[k]}, \\ [x_{n-2}^{[k]}, [x_{n-1}^{[k]}, x_n^{[k]}]] &= (E_{n-2,n+1} + a_k^{-1} E_{1,2n-2})^{[k]}, \\ &\dots \quad \dots \quad \dots \\ [x_1^{[k]}, [x_2^{[k]} \dots [x_{n-1}^{[k]}, x_n^{[k]}] \dots]] &= (1 + a_k^{-1})(E_{1,n+1})^{[k]}, \\ [y_n^{[k]}, y_{n-1}^{[k]}] &= (E_{n+1,n-1} + a_k E_{2n-1,1})^{[k]}, \\ [[y_n^{[k]}, y_{n-1}^{[k]}], y_{n-2}^{[k]}] &= (E_{n+1,n-2} + a_k E_{2n-2,1})^{[k]}, \\ &\dots \quad \dots \quad \dots \\ [[\dots [y_n^{[k]}, y_{n-1}^{[k]}], \dots y_2^{[k]}], y_1^{[k]}] &= (1 + a_k)(E_{n+1,1})^{[k]}, \end{aligned}$$

we infer that

$$\begin{aligned} H_{1,n+1} &:= \left[\sum_{k=1}^K (1 + a_k^{-1})(E_{1,1+n})^{[k]}, \sum_{k=1}^K (1 + a_k)(E_{n+1,1})^{[k]} \right] \\ &= \sum_{k=1}^K [(1 + a_k^{-1})(E_{1,1+n})^{[k]}, (1 + a_k)(E_{n+1,1})^{[k]}] \\ &= \sum_{k=1}^K (2 + a_k + a_k^{-1})(E_{1,1} - E_{n+1,n+1})^{[k]}. \end{aligned}$$

Consider its action on $\psi_{\underline{a}}(e_1) = \sum_{k=1}^K x_1^{[k]}$ and $\psi_{\underline{a}}(f_1) = \sum_{k=1}^K y_1^{[k]}$, we have that

$$\begin{aligned}
(\text{ad}H_{1,n})(\psi_{\underline{a}}(e_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1}) x_1^{[k]}, \\
(-\text{ad}H_{1,n})(\psi_{\underline{a}}(f_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1}) y_1^{[k]}, \\
(\text{ad}H_{1,n})^2(\psi_{\underline{a}}(e_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1})^2 x_1^{[k]}, \\
(-\text{ad}H_{1,n})^2(\psi_{\underline{a}}(f_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1})^2 y_1^{[k]}, \\
&\dots \dots \dots \\
(\text{ad}H_{1,n})^K(\psi_{\underline{a}}(e_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1})^K x_1^{[k]}, \\
(-\text{ad}H_{1,n})^K(\psi_{\underline{a}}(f_1)) &= \sum_{k=1}^K (2 + a_k + a_k^{-1})^K y_1^{[k]}.
\end{aligned}$$

If $2 + a_k + a_k^{-1} = 2 + a_j + a_j^{-1}$, then $(a_k a_j - 1)(a_k - a_j) = 0$. By our assumption, every $2 + a_k + a_k^{-1}$ is non-zero and $2 + a_k + a_k^{-1} \neq 2 + a_j + a_j^{-1}$ for all $k \neq j$. By the invertibility of Vandermonde matrix, we infer that

$$x_1^{[1]}, y_1^{[1]}, x_1^{[2]}, y_1^{[2]}, \dots, x_1^{[K]}, y_1^{[K]} \in \text{Im}(\psi_{\underline{a}}).$$

Let $H_{1,2}^{[k]} = [x_1^{[k]}, y_1^{[k]}] = (E_{1,1} - E_{2,2})^{[k]}$, and consider its action on $\psi_{\underline{a}}(e_2), \psi_{\underline{a}}(f_2)$, we infer that

$$x_2^{[1]}, y_2^{[1]}, x_2^{[2]}, y_2^{[2]}, \dots, x_2^{[K]}, y_2^{[K]} \in \text{Im}(\psi_{\underline{a}}).$$

Repeat this process, similar argument implies that

$$x_i^{[k]}, y_i^{[k]} \in \text{Im}(\psi_{\underline{a}})$$

for every k . By Proposition 4.2, we can infer that $\mathbf{g}_k \in \text{Im}(\psi_{\underline{a}})$ for all k , and thus the map $\psi_{\underline{a}}$ is surjective. \square

Lemma 5.2. *If $a_1 = 1$ and $a_k \neq -1$ for every $k \geq 2$, then $\text{Im}(\psi_{\underline{a}})$ is a direct sum of one copy of sp_{2n} and $K - 1$ copies of sl_{2n} .*

Proof. We may repeat the proof for Lemma 5.1 until we obtain the result

$$x_i^{[k]}, y_i^{[k]} \in \text{Im}(\psi_{\underline{a}}), \forall 1 \leq k \leq K, 1 \leq i \leq n,$$

then $\text{Im}(\psi_{\underline{a}}) = \bigoplus_{k=1}^K \text{Im}(\psi_{a_k})$, by Proposition 4.2, $\text{Im}(\psi_{a_1})$ is isomorphic to sp_{2n} , and $\text{Im}(\psi_{a_k})$ is isomorphic to sl_{2n} for $k \geq 2$. \square

Lemma 5.3. *If $a_1 = -1, a_2 = 1$, then $\text{Im}(\psi_{\underline{a}})$ is a direct sum of one copy of so_{2n} , one copy of sp_{2n} and $K - 2$ copies of sl_{2n} .*

Proof. We may repeat the proof for Lemma 5.1 until we obtain $H_{1,n+1}$. Notice that, in this case

$$H_{1,n+1} := \sum_{k=2}^K (2 + a_k + a_k^{-1})(E_{1,1} - E_{n+1,n+1})^{[k]}.$$

Then go on with the process in Lemma 5.2, we get that

$$x_i^{[k]}, y_i^{[k]} \in \text{Im}(\psi_{\underline{a}}), \forall 2 \leq k \leq K, 1 \leq i \leq n,$$

then

$$x_i^{[1]} = \psi_{\underline{a}}(e_i) - \left(\sum_{k=2}^K x_i^{[k]} \right) \in \text{Im}(\psi_{\underline{a}}), \quad y_i^{[1]} = \psi_{\underline{a}}(f_i) - \left(\sum_{k=2}^K y_i^{[k]} \right) \in \text{Im}(\psi_{\underline{a}}),$$

it also holds that $\text{Im}(\psi_{\underline{a}}) = \bigoplus_{k=1}^K \text{Im}(\psi_{a_k})$. By Proposition 4.2, $\text{Im}(\psi_{a_1})$ is isomorphic to so_{2n} , $\text{Im}(\psi_{a_2})$ is isomorphic to sp_{2n} , and $\text{Im}(\psi_{a_k})$ is isomorphic to sl_{2n} for $k \geq 3$. \square

Lemma 5.4. *If $a_1 = -1$ and $a_k \neq 1$ for $k \geq 2$, then $\text{Im}(\psi_{\underline{a}})$ is a direct sum of one copy of so_{2n} and $K - 1$ copies of sl_{2n} .*

Proof. The proof is very similar to that for Lemma 5.3. \square

6. PROOF FOR THEOREM 3.2

If $G(t)$ is a polynomial with non-zero constant term, we may define the quotient algebra

$$\hat{\mathfrak{g}} / \langle \mathfrak{g} \otimes G(t) \mathbb{C}[t, t^{-1}] \rangle,$$

and denoted by $\hat{\mathfrak{g}}/G(t)$. Note that, we may regard the central element c as 0 in the quotient algebra $\hat{\mathfrak{g}}/G(t)$ for all non-trivial polynomial $G(t)$.

Lemma 6.1. *If $G(t) | F(t)$, then $\hat{\mathfrak{g}}/G(t)$ is a quotient of $\hat{\mathfrak{g}}/F(t)$. Particularly,*

$$\hat{\mathfrak{g}}/G(t) \cong \frac{\hat{\mathfrak{g}}/F(t)}{\mathfrak{g} \otimes G(t) \mathbb{C}[t, t^{-1}] / F(t)}.$$

Proof. This result follows from the homomorphism fundamental theorem. \square

Lemma 6.2. *If $(x - a)^r | G(t)$ for some $r > 1$, then $\hat{\mathfrak{g}}/G(t)$ has a nilpotent ideal*

$$I = \mathfrak{g} \otimes \frac{G(t)}{(t - a)} \mathbb{C}[t, t^{-1}].$$

Proof. Obviously, for any $x \in \hat{\mathfrak{g}}$ and $y \in I$, we have $[x, y] \in I$. Suppose that $r^* \geq r$ is the multiple of prime factor $t - a$, then for any $u \otimes f(t), v \otimes g(t) \in I$, we have $(t - a)^2 | G(t)$, and

$$G(t) \mid \frac{G(t)^2}{(t - a)^2}, \quad \frac{G(t)^2}{(t - a)^2} \mid f(t)g(t),$$

and thus $[I, I] = 0$. \square

If $G(t)$ has degree K , then it is well-known that $\hat{\mathfrak{g}}/G(t)$ is the direct sum of K copies of \mathfrak{g} if $G(t)$ has K many different factors $t - a_i$, where $a_i \neq 0 (1 \leq i \leq K)$.

Let $G(t) = \prod_{i=1}^k (t - a_i)$, then there exists $c_i \neq 0$, such that

$$\sum_{i=1}^k c_i \frac{G(t)}{t - a_i} = 1,$$

then

$$\hat{\mathfrak{g}}/G(t) = \bigoplus_{i=1}^k \hat{\mathfrak{g}} \otimes \frac{G(t)}{t - a_i} / G(t) \cong \bigoplus_{i=1}^k \frac{\hat{\mathfrak{g}}}{t - a_i}.$$

Let ϕ be as defined in Section 4. Then

$$\begin{aligned} [\hat{e}_1, \dots [\hat{e}_{n-1}, \hat{e}_n] \dots] &= E_{\alpha_1 + \dots + \alpha_n} \otimes (1 + t^{-1}), \\ [\dots [\hat{f}_{n-1}, \hat{f}_{n-2}], \dots \hat{f}_1] &= E_{-\alpha_1 - \dots - \alpha_{n-1}} \otimes 1 - E_{\alpha_{n+1} + \dots + \alpha_{2n-1}} \otimes 1, \\ [[\dots [\hat{f}_{n-1}, \hat{f}_{n-2}], \dots \hat{f}_1], E_{\alpha_1 + \dots + \alpha_n} \otimes (1 + t^{-1})] &= (E_{\alpha_n} - E_{\alpha_1 + \dots + \alpha_{2n-1}}) \otimes (1 + t^{-1}), \\ [(E_{\alpha_n} - E_{\alpha_1 + \dots + \alpha_{2n-1}}) \otimes (1 + t^{-1}), \hat{f}_n] &= H_n \otimes (1 + t^{-1}) + (H_1 + \dots + H_{2n-1}) \otimes (1 + t) - c, \end{aligned}$$

this implies that

$$\Xi := H_n \otimes t^{-1} + (H_1 + \dots + H_{2n-1}) \otimes t \in \phi(\mathbf{gim}(M_n)),$$

then

$$\begin{aligned} [(\frac{1}{2}\text{ad}\Xi)^m(\hat{e}_n), \hat{f}_n] &= H_n \otimes t^{-m} + (H_1 + \dots + H_{2n-1}) \otimes t^m, \\ [(\text{ad}\Xi)^m(\hat{e}_1), \hat{f}_1] &= H_1 \otimes t^m - H_{n+1} \otimes t^{-m}, \end{aligned}$$

for all $m > 0$. Similarly, we infer that

$$H_i \otimes t^{-m} - H_{n+i} \otimes t^m \in \phi(\mathbf{gim}(M_n)),$$

for all $1 \leq i \leq n$ and $m \in \mathbb{Z} \setminus \{0\}$, where $H_{2n} = -H_1 - \dots - H_{2n-1}$.

Moreover,

$$(H_i \otimes t^{-m} - H_{n+i} \otimes t^m) \pm (H_i \otimes t^m - H_{n+i} \otimes t^{-m}) = (H_i \mp H_{n+i}) \otimes (t^m \pm t^{-m}).$$

Suppose that I is an ideal of $\mathbf{gim}(M_n)$ such that the quotient algebra $\mathbf{gim}(M_n)/I$ is finite-dimensional and semi-simple. Since ϕ is an isomorphism, we have that $\phi(I)$ is an ideal of $\hat{\mathbf{g}}_{fp} = \phi(\mathbf{gim}(M_n)) \cong \mathbf{gim}(M_n)$ and $\hat{\mathbf{g}}_{fp}/\phi(I) \cong \mathbf{gim}(M_n)/I$. Note that every polynomial in variable $t + t^{-1}$ is a linear combination of $t^m + t^{-m}$ ($m \geq 0$). Then there must hold that

$$(H_i - H_{n+i}) \otimes \theta(t) \in \phi(I), \quad 1 \leq i \leq n,$$

where $\theta(t) = \eta(t + t^{-1})$ for some polynomial η . Otherwise, the elements

$$\left\{ H \otimes (t^j + t^{-j}) \mid j \in \mathbb{Z}_{\geq 0} \right\}$$

are linearly independent in $\hat{\mathbf{g}}_{fp}/\phi(I)$ for some $H \in \text{span}\{H_i - H_{n+i} \mid 1 \leq i \leq n\}$, which contradicts to the assumption of the finite-dimension.

Actually, $\hat{h}_i = H_i - H_{n+i}$ for all $1 \leq i \leq n$. Let $\mathfrak{h}_{fp} = \text{span}\{\hat{h}_i \mid 1 \leq i \leq n\}$. Then $\hat{\mathbf{g}}_{fp}$ has a root decomposition

$$\hat{\mathbf{g}}_{fp} = (\hat{\mathbf{g}}_{fp})_0 + \sum_{\alpha} (\hat{\mathbf{g}}_{fp})_{\alpha},$$

where $(\hat{\mathbf{g}}_{fp})_0 = \{x \in \hat{\mathbf{g}}_{fp} \mid [\mathfrak{h}_{fp}, x] = 0\}$ and $(\hat{\mathbf{g}}_{fp})_{\alpha} = \{x \in \hat{\mathbf{g}}_{fp} \mid [h, x] = \alpha(h)x, \forall h \in \mathfrak{h}_{fp}\}$.

By Berman's result ([Br], Proposition 1.12), it holds that $\hat{\mathbf{g}} = \hat{\mathbf{g}}_{fp} \oplus M$, where M is a $\hat{\mathbf{g}}_{fp}$ -module and $[M, M] \subseteq \hat{\mathbf{g}}_{fp}$. If $x = \sum_{j=1}^d x_j \otimes p_j(t)\theta(t) \in (\hat{\mathbf{g}}_{fp})_{\alpha}$, $\alpha \neq 0$, then there exists $H \in \mathfrak{h}_{fp}$ such that $\alpha(H) \neq 0$ and

$$x = \frac{1}{\alpha(H)} \sum_{j=1}^d [H \otimes \theta(t), x_j \otimes p_j(t)] \in \phi(I).$$

Otherwise, we infer that $\sum_{j=1}^d x_j \otimes p_j(t) \notin \hat{\mathbf{g}}_{fp}$ and thus $x \notin \hat{\mathbf{g}}_{fp}$, which is a contradiction.

Moreover, $(\hat{\mathfrak{g}}_{fp})_0 \subseteq \mathfrak{h} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}c$ and

$$(\mathfrak{h} \otimes \mathbb{C}[t + t^{-1}] \oplus \mathbb{C}c) \cap \hat{\mathfrak{g}}_{fp} = \sum_{m=0}^{\infty} \sum_{i=1}^n \mathbb{C}((H_i - H_{n+i}) \otimes (t^m + t^{-m}) - \delta_{i,n}c),$$

then we have the following result:

Lemma 6.3. *We let J be the ideal of $\hat{\mathfrak{g}}$ generated by $\{H_i \otimes \theta(t) | 1 \leq i \leq 2n - 1\}$, then*

$$\phi(\mathbf{gim}(M_n)) \cap J \subseteq \phi(I),$$

and naturally there exists an epimorphism

$$\frac{\phi(\mathbf{gim}(M_n))}{\phi(\mathbf{gim}(M_n)) \cap J} \rightarrow \mathbf{gim}(M_n)/I.$$

For convenience, we let $\theta^*(t) = t^r \eta(t + t^{-1})$, where $r = \deg \eta$, then θ^* is a polynomial with $\theta^*(0) \neq 0$ and it holds that

$$\hat{\mathfrak{g}}/J = \hat{\mathfrak{g}}/\theta^*(t).$$

Moreover, $\theta^*(t)$ and $\eta(t + t^{-1})$ have same roots.

Suppose that $\hat{\mathfrak{g}}/\theta^*(t)$ has a Levi decomposition

$$\hat{\mathfrak{g}}/\theta^*(t) = S + U,$$

where S is semi-simple and U is the solvable radical. Let $\mathbf{p} : S + U \rightarrow (S + U)/U$ be the canonical map. If $\mathbf{i} : \mathbf{gim}(M_n)/I \rightarrow \hat{\mathfrak{g}}/\theta^*(t) \subseteq S$ is the injective map, then $\mathbf{p} \circ \mathbf{i}$ is injective. Hence we may assume that

$$\theta^*(t) = \prod_{i=1}^K (t - a_i),$$

where $a_i \neq a_j$ if $i \neq j$.

Out of questions, $\mathbf{gim}(M_n)/I$ is a subalgebra of $\hat{\mathfrak{g}}/\theta^*(t)$ through isomorphism ϕ .

We may assume that $c_i, d_i \in \mathbb{C}^\times$ such that

$$\sum_{i=1}^K c_i \frac{\theta^*(t)}{t - a_i} = 1, \quad \frac{\theta^*(t)}{t - a_i} \equiv d_i \pmod{t - a_i},$$

we also assume that $\hat{\mathfrak{g}}/\theta^*(t) = \bigoplus_{i=1}^K \mathfrak{g}_i$, and ψ^\clubsuit is the canonical map from $\hat{\mathfrak{g}}$ to $\hat{\mathfrak{g}}/\theta^*(t)$.

Let

$$\psi_{a_i} : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}_i = \frac{\mathfrak{g} \otimes \theta^*(t)/(t - a_i)\mathbb{C}[t, t^{-1}]}{(t - a_i)} \cong \hat{\mathfrak{g}}/(t - a_i)$$

be the evaluation map, where $\psi_{a_i}(x \otimes t^m \frac{\theta^*(t)}{(t - a_i)}) = d_i a_i^m x^{[i]}$.

Lemma 6.4. *For any $x \otimes t^m$, we have*

$$\psi^\clubsuit(x \otimes t^m) = \sum_{i=1}^K c_i d_i \psi_{a_i}(x \otimes t^m) = \sum_{i=1}^K c_i d_i a_i^m x^{[i]}.$$

Particularly, $c_i d_i = 1$ for any $1 \leq i \leq K$, then $\psi^\clubsuit = \psi_{\underline{a}}$, where $\underline{a} = (a_1, \dots, a_K)$.

Proof. Since $\sum_{i=1}^K c_i \frac{\theta^*(t)}{t-a_i} = 1$, evaluate $t = a_i$ in both sides, we have $c_i d_i = 1$. For any $x \otimes t^m$, we have

$$\begin{aligned} \psi^\clubsuit(x \otimes t^m) &= \sum_{i=1}^K c_i x \otimes t^m \frac{\theta^*(t)}{t-a_i} \\ &= \sum_{i=1}^K c_i d_i a_i^m x^{[i]} = \sum_{i=1}^K a_i^m x^{[i]}. \end{aligned}$$

□

Lemma 6.5. *Let $a \neq 0, \pm 1$, then associated to the map*

$$\psi_a \oplus \psi_{a^{-1}} : \mathbf{gim}(M_n)/I \rightarrow \hat{\mathbf{g}}/(t-a) \oplus \hat{\mathbf{g}}/(t-a^{-1}),$$

we have that

$$Im(\psi_a \oplus \psi_{a^{-1}}) \cong Im(\psi_a).$$

Proof. As we know that

$$\begin{aligned} \sigma : \quad \mathbf{g} \otimes \mathbb{C}[t, t^{-1}] &\rightarrow \mathbf{g} \otimes \mathbb{C}[t, t^{-1}] \\ x \otimes t^m &\mapsto x \otimes t^{-m}, \end{aligned}$$

is an isomorphism. Moreover, $\psi_a \circ \sigma = \psi_{a^{-1}}$, hence we have that

$$\psi_a \oplus \psi_{a^{-1}} = \psi_a \circ (1 \oplus \sigma),$$

notice that $1 \oplus \sigma$ is diagonally injective, then the image of $\psi_a \oplus \psi_{a^{-1}}$ is isomorphic to $Im(\psi_a)$. □

Proof for Theorem 2.3: Undoubtedly, since η is a polynomial in variable $t+t^{-1}$ and $\theta^*(0) \neq 0$, we have that $(t-a)|\theta^*(t)$ if and only if $(t-a^{-1})|\theta^*(t)$. By Lemma 6.4, $\psi^\clubsuit = \psi_{\underline{a}}$. Here, $a_k \in \underline{a}$ if and only if $a_k^{-1} \in \underline{a}$. However, the approach of proof for Lemma 5.1 still play a major role in this case. Follow the process of the proof for Lemma 5.1 and Lemma 5.3, we can successfully reach that:

$$\begin{aligned} x_i^{[k]}, y_i^{[k]} &\in Im(\psi_{\underline{a}}), & a_k = \pm 1, 1 \leq i \leq n \\ x_i^{[k]} + x_i^{[j]}, y_i^{[k]} + y_i^{[j]} &\in Im(\psi_{\underline{a}}), & a_k \neq \pm 1, a_k a_j = 1, 1 \leq i \leq n. \end{aligned}$$

By Proposition 4.2 and Lemma 6.5, the subalgebra generated by $x_i^{[k]}, y_i^{[k]} (a_k = \pm 1, 1 \leq i \leq n)$ is one of so_{2n}, sp_{2n} , and the subalgebra generated by $x_i^{[k]} + x_i^{[j]}, y_i^{[k]} + y_i^{[j]} (a_k a_j = 1, 1 \leq i \leq n)$ is sl_{2n} . Since $t-1$ or $t+1$ appears in $\theta(t)$ at most one time, each of so_{2n} and sp_{2n} appears at most one time. Then $\mathbf{gim}(M_n)/I$ is of type $M(n, \mathbf{a}, \mathbf{c}, \mathbf{d})$.

The proof is completed.

Acknowledgements.

The first author is partially supported by NSERC of Canada. The second author is supported by the NNSF of China (Grant No. 11001110, 11271131) and *Jiangsu Government Scholarship for Overseas Studies*.

REFERENCES

- [ABnG] B. Allison, G. Benkart and Y. Gao, *Lie algebras graded by the root system BC_r , $r \geq 2$* , Mem. A.M.S.(751)158(2002)
- [BG] S. Bhargava and Y. Gao, *Realizations of BC_r -graded intersection matrix algebras with grading subalgebras of type B_r , $r \geq 3$* , Pacific Journal of Mathematics, Vol. 263 (2013), No. 2, 257-281.
- [BnZ] G. Benkart and E. Zelmanov, *Lie algebras graded by finite root systems and intersection matrix algebras*, Invent. Math. 126 (1996), 1-45.
- [Br] S. Berman, *On generators and relations for certain involutory subalgebras of Kac-Moody Lie Algebras*, Comm. Alg. 17 (1989), 3165-3185.
- [BrJT] S. Berman, E. Jurisich and S. Tan, *Beyond Borcherds Lie Algebras and Inside*, Trans. AMS 353 (2001), 1183-1219.
- [BrM] S. Berman and R.V. Moody, *Lie algebras graded by finite root systems and the intersection matrix algebras of Slowdow*, Invent. Math. 108 (1992), 323-347.
- [C] R. Carter, *Lie Algebras of Finite and Affine Type*, Cambridge Univ. Press, Cambridge, 2005.
- [CM] V. Chari, A. A. Moura, *Spectral characters of finite-dimensional representations of affine algebras*, Journal of Algebra 279 (2004) 820-839.
- [EMY] S. Eswara Rao, R.V. Moody and T. Yokonuma, *Lie algebras and Weyl groups arising from vertex operator representations*, Nova J. of Algebra and Geometry 1 (1992), 15-57.
- [G] Y. Gao, *Involutive Lie algebras graded by finite root systems and compact forms of IM algebras*, Math. Zeitschrift 223 (1996), 651- 672.
- [GbK] O. Gabber and V.G. Kac, *On Defining Relations of Certain Infinite-Dimensional Lie Algebras*, Bull. Amer. Math. Soc. (N.S.) 5 (1981), 185-189.
- [H] J.E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer, New York, 1972.
- [J] N. Jacobson, *Lie Algebras*, Interscience, New York, 1962.
- [K] V.G. Kac, *Infinite Dimensional Lie Algebras*, 3rd edition, Cambridge Univ. Press, Cambridge, 1990.
- [M1] R. V. Moody, *Euclidean Lie algebras*, Canad. J. Math. 21 (1969), 1432C1454.
- [M2] R. V. Moody, *Simple quotients of Euclidean Lie algebras*, Canad. J. Math. 22 (1970), 839C846.
- [N] E. Neher, *Lie algebras graded by 3-graded root systems and Jordan pairs covered by grids*, Amer. J. Math. 118 (1996), no. 2, 439-491.
- [P] L. Peng, *Intersection matrix Lie algebras and Ringel-Hall Lie algebras of tilted algebras*, Representations of Algebra Vol. I, II, 98-108, Beijing Normal Univ. Press, Beijing, 2002.
- [S1] P. Slodowy, *Beyond Kac-Moody algebras and inside*, Can. Math. Soc. Conf. Proc. 5 (1986), 361-371.
- [S2] P. Slodowy, *Singularitäten, Kac-Moody Lie-Algebren, assoziierte Gruppen und Verallgemeinerungen*, Habilitationsschrift, Universität Bonn, March 1984.