

Tight Bounds on ℓ_1 Approximation and Learning of Self-Bounding Functions

Vitaly Feldman*
Google Brain

Pravesh Kothari*
Carnegie Mellon University

Jan Vondrák
Stanford University

Abstract

We study the complexity of learning and approximation of self-bounding functions over the uniform distribution on the Boolean hypercube $\{0, 1\}^n$. Informally, a function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ is self-bounding if for every $x \in \{0, 1\}^n$, $f(x)$ upper bounds the sum of all the n marginal decreases in the value of the function at x . Self-bounding functions include such well-known classes of functions as submodular and fractionally-subadditive (XOS) functions. They were introduced by Boucheron *et al.* (2000) in the context of concentration of measure inequalities. Our main result is a nearly tight ℓ_1 -approximation of self-bounding functions by low-degree juntas. Specifically, all self-bounding functions can be ϵ -approximated in ℓ_1 by a polynomial of degree $\tilde{O}(1/\epsilon)$ over $2^{\tilde{O}(1/\epsilon)}$ variables. We show that both the degree and junta-size are optimal up to logarithmic terms. Previous techniques considered stronger ℓ_2 approximation and proved nearly tight bounds of $\Theta(1/\epsilon^2)$ on the degree and $2^{\Theta(1/\epsilon^2)}$ on the number of variables. Our bounds rely on the analysis of noise stability of self-bounding functions together with a stronger connection between noise stability and ℓ_1 approximation by low-degree polynomials. This technique can also be used to get tighter bounds on ℓ_1 approximation by low-degree polynomials and a faster learning algorithm for halfspaces.

These results lead to improved and in several cases almost tight bounds for PAC and agnostic learning of self-bounding functions relative to the uniform distribution. In particular, assuming hardness of learning juntas, we show that PAC and agnostic learning of self-bounding functions have complexity of $n^{\tilde{\Theta}(1/\epsilon)}$.

1 Introduction

We consider learning and approximation of several classes of real-valued functions over the uniform distribution on the Boolean hypercube $\{0, 1\}^n$. The most well-studied class of functions that we consider is the class of submodular functions. A related class of functions is that of *fractional subadditive functions*, equivalently known as XOS functions, which generalize monotone submodular functions and have been introduced in the context of combinatorial auctions [LLN06]. XOS functions are also known to have an equivalent definition as Rademacher complexity of a subset of data points for some class of functions [FV15]. It turns out that these classes are all contained in a broader class, that of *self-bounding functions*, introduced in the context of concentration of measure inequalities [BLM00]. Informally, a function f over $\{0, 1\}^n$ is a -self-bounding if for every $x \in \{0, 1\}^n$, $a \cdot f(x)$ upper bounds the sum of all the n marginal decreases in the value of the

*Work done while the author was at IBM Research - Almaden.

function at x . For XOS functions $a = 1$ and for submodular¹ $a = 2$ (a is omitted when it equals 1). See Sec. 2 for formal definitions and examples of self-bounding functions.

Wide-spread applications of submodular functions have recently inspired the question of whether and how such functions can be learned from random examples (of an unknown submodular function). The question was first formally considered by Balcan and Harvey [BH12] who motivate it by learning of valuation functions. Reconstruction of such functions up to some multiplicative factor from value queries (which allow the learner to ask for the value of the function at any point) was also considered by Goemans et al. [GHIM09]. In this work we consider the setting in which the learner gets random and uniform examples of an unknown function f and its goal is to find a hypothesis function h that ϵ -approximates the unknown function for a given $\epsilon > 0$. The measure of the approximation error we use is the standard absolute error or ℓ_1 -distance, which equals $\mathbf{E}_{x \sim D}[|f(x) - h(x)|]$. While other measures of error, such as ℓ_2 , are often studied in machine learning, there is a large number of scenarios where the expected absolute error is used. For example, if the unknown function is Boolean then learning with ℓ_1 error is equivalent to learning with Boolean disagreement error [KKMS08]. In fact, it is known that the complexity of agnostic learning over product distributions in the statistical query model is characterized by how well the Boolean functions can be approximated in ℓ_1 by low-degree polynomials [DSFTWW15]. Applications of learning algorithms for submodular functions to differentially-private data release require ℓ_1 error [GHRU11; CKKL12; FK14] as does learning of probabilistic concepts (which are concepts expressing the probability of an event) [KS94].

Motivated by applications to learning, prior works have also studied a number of natural questions on approximation of submodular and related classes of functions by concisely represented functions. For example, linear functions [BH12], low-degree polynomials [CKKL12; FV15], DNF formulas [RY13], decision trees [FKV13] and functions of few variables (referred to as *juntas*) [FKV13; BOSY13; FV15; FV16]. We survey the prior work in more detail in Section 1.2.

1.1 Our results

In this work, we provide nearly tight bounds on approximation of self-bounding functions by low-degree polynomials and juntas in the ℓ_1 -norm. The results are obtained via the noise-stability analysis of self-bounding functions. Previous approximation bounds for the uniform distribution relied on bounding ℓ_2 error that is more convenient to analyze using Fourier techniques. However this approach has so far led to weaker bounds on ℓ_1 approximation error. Further the known bounds on ℓ_2 approximation are known to be optimal [FV15]. The dependence of the degree and junta size on the error parameter ϵ in our bounds is quadratically better (up to a logarithmic term) than bounds which are known for ℓ_2 error.

Structural results: Our two key structural results can be summarized as follows.

Theorem 1.1. *Let $f : \{0,1\}^n \rightarrow [0,1]$ be an a -self-bounding function and $\epsilon > 0$. For $d = O(a/\epsilon \cdot \log(1/\epsilon))$ there exists a set of indices I of size $2^{O(d)}$ and a polynomial p of degree d over variables in I such that $\|f - p\|_1 \leq \epsilon$.*

¹Technically, self-bounding functions are always non-negative and hence capture only non-negative submodular functions. Submodularity is preserved under shifting of the function and therefore it is sufficient to consider non-negative submodular functions.

This result itself is based on a combination of two structural results. The first one gives a degree bound of $O(\frac{a}{\epsilon} \log \frac{1}{\epsilon})$. Previously, it was known that submodular functions with range $[0, 1]$ can be ϵ -approximated by polynomials of degree $O(1/\epsilon^2)$ [CKKL12; FKV13]. Feldman and Vondrák [FV16] showed that the same upper bound applies all self-bounding functions and, more generally, all functions of low total influence. More recently, it was shown that this upper bound is tight [FV15]. For comparison, as follows from the results in [FV15], for XOS functions there is no significant difference in between ℓ_1 and ℓ_2 approximation. In both cases degree $\Theta(1/\epsilon)$ and junta of size $2^{\Theta(1/\epsilon)}$ are needed. One natural open problem that is left open is the degree of polynomial necessary to approximate a submodular function in ℓ_1 norm.

Our proof is based on a new and simple connection between (the appropriately generalized notion of) noise sensitivity of a real-valued function and its approximability by a low-degree polynomial. The key observation here is that the application of the noise operator to a function f that has low noise sensitivity gives a function that is close to f in ℓ_1 norm. The obtained *smoothed* function is much easier to approximate by a low-degree polynomial since its Fourier spectrum decays rapidly with the growth of the degree. This technique is general and also gives a sharper bound for ℓ_1 approximation of halfspaces by low-degree polynomials (see Cor. 3.6). To apply this technique to self-bounding functions we show that noise-sensitivity can be upper bounded using a bound on the total ℓ_1 influence of all the coordinates on the function. It is known that a -self-bounding functions have total influence of at most a [FV16] and thus we obtain that any a -self-bounding function has bounded noise sensitivity and can be approximated by a degree $O(\frac{a}{\epsilon} \log \frac{1}{\epsilon})$ polynomial.

The second component of this result builds on the work of [FV16], where it was shown that a classic theorem of Friedgut [Fri98], on approximation of Boolean functions by juntas, generalizes to the setting of real-valued functions by including a dependence on ℓ_1 as well as ℓ_2 -influences of the function. We show that by applying the analysis from [FV16] to the smoothed version of f (for which we have better degree bounds) we can obtain approximation by a junta of size $2^{O(a/\epsilon \cdot \log(1/\epsilon))}$. This improves on $2^{O(a/\epsilon^2)}$ bound in [FV16] (that holds also for ℓ_2 error). We note that both of the components also apply to the more general class of functions with low total ℓ_1 influence.

We then study the effect of the noise operator on self-bounding functions in more detail. We demonstrate that the smoothed version is noise stable even in the stronger point-wise sense: for every x , the smoothed function at x cannot be much smaller than $f(x)$. This result generalizes a similar result from [CKKL12] for submodular functions. Such stability implies that for every non-negative a -self-bounding function f , $\|f\|_1 \geq \frac{1}{3a} \|f\|_\infty$ (see Lemma 3.9). This has been known for submodular [FMV07] and XOS [Fei06] functions (with a constant a) and, together with approximation by a junta, can be used to obtain a learning algorithm with multiplicative approximation guarantees for all a -self-bounding functions [FV16].

Algorithmic applications: It is easy to exploit our structural results in existing learning algorithms to obtain better running time and sample complexity bounds. We describe one of these results here and some additional ones in Section 4. Specifically, we give an algorithm for learning all a -self-bounding functions relative to the uniform distribution in the challenging agnostic framework. An agnostic learning algorithm for a class of functions \mathcal{C} is an algorithm that given random examples of *any* function f finds a hypothesis h whose error is at most ϵ -greater than the error of the best hypothesis in \mathcal{C} (see [KSS94] for the Boolean case).

Theorem 1.2. *Let \mathcal{C}_a be the class of all a -self-bounding functions from $\{0, 1\}^n$ to $[0, 1]$. There exists an algorithm \mathcal{A} that given $\epsilon > 0$ and access to random uniform examples of any real-valued*

f , with probability at least $2/3$, outputs a function h , such that $\|f - h\|_1 \leq \Delta + \epsilon$, where $\Delta = \min_{g \in \mathcal{C}_a} \{\|f - g\|_1\}$. Further, \mathcal{A} runs in time $n^{\tilde{O}(a/\epsilon)}$ and uses $2^{\tilde{O}(a^2/\epsilon^2)} \log n$ examples.

This algorithm is based on polynomial ℓ_1 regression with an additional constraint on the spectral norm of the solution to obtain a stronger sample complexity bound [FV16]. The best previous bound of $n^{O(a/\epsilon^2)}$ time and $2^{O(a^2/\epsilon^4)} \log n$ examples follows from the results in [FV16] for function of low total influence.

Lower bounds: We prove that a -self-bounding functions require degree $\Omega(a/\epsilon)$ to ϵ -approximate in ℓ_1 distance (see Cor. 5.7). A construction of a parity function correlated with a submodular function in [FKV13] also implies that even submodular functions require polynomials of degree $\Omega(\epsilon^{-2/3})$ to ϵ -approximate in ℓ_1 .

In [FV16] it is shown that XOS functions require a junta of size $2^{\Omega(1/\epsilon)}$ to ϵ -approximate (however submodular functions admit approximation by exponentially smaller juntas [FV16]). This also implies $2^{\Omega(a/\epsilon)}$ lower bound on junta size for a -self-bounding functions (see Lem. 5.3). Therefore our structural results are essentially tight for self-bounding functions.

We then show that our agnostic learning algorithm for a -self-bounding function is nearly optimal. In fact, even PAC learning of non-monotone a -self-bounding functions requires time $n^{\Omega(a/\epsilon)}$ assuming hardness of learning k -term DNF to accuracy $1/4$ in time $n^{\Omega(k)}$. This is in contrast to the submodular [FKV13; FV16] and monotone self-bounding cases (Thm. 4.3).

Theorem 1.3. *For every $a \geq 1$, if there exists an algorithm that PAC learns a -self-bounding functions with range $[0, 1]$ to ℓ_1 error of $\epsilon > 0$ in time $T(n, 1/\epsilon)$ then there exists an algorithm that PAC learns k -DNF formulas to accuracy ϵ' in time $T(n, k/(a \cdot \epsilon'))$ for some fixed constant c .*

To prove this hardness results we show that a k -DNF formula (of any size) is a k -self-bounding function. Using an additional “lifting” trick we can also embed k -DNF formulas into a -self-bounding functions for any $a \geq 1$. Note that any k -junta can be computed by a k -DNF formula. Learning of DNF expressions is a well-studied problem in learning theory but there are no algorithms for this problem better than the trivial $O(n^k)$ algorithm, even for a constant $\epsilon' = 1/4$. The (potentially simpler) problem of learning k -juntas is also considered very hard [BL97; Blu03]. Until recently, the only non-trivial algorithm for the problem was the $O(n^{0.7k})$ -time algorithm by Mossel et al. [MOS04]. The best known upper bound is $O(n^{0.6k})$ and was given in the recent breakthrough result of Valiant [Val12]. Learning of k -juntas is also known to have complexity of $n^{\Omega(k)}$ for all statistical query algorithms [BFJKMR94]. Theorem 1.3 implies that PAC learning of a -self-bounding functions in time $n^{o(a/\epsilon)}$ would lead to a $n^{o(k)}$ algorithm for learning k -DNF to any constant accuracy and, in particular, an algorithm for PAC learning k -juntas in time $n^{o(k)}$. We note that the dependence on a/ϵ in our lower bound matches our upper bound up to a logarithmic factor.

Finally, we remark that our reduction to learning of k -DNF also implies that PAC learning of a -self-bounding functions requires at least $2^{\Omega(a/\epsilon)}$ random examples or even stronger value queries (see Cor. 5.6). Therefore sample complexity bounds we give are also close to optimal. Further details of lower bounds are given in Section 5.

1.2 Related work

Below we briefly mention some of the other related work. We direct the reader to [BH12] and [FV15] for more detailed surveys. Balcan and Harvey study learning of submodular functions

without assumptions on the distribution and also require that the algorithm output a value which is within a multiplicative approximation factor of the true value with probability $\geq 1 - \epsilon$ (the model is referred to as *PMAC learning*). This is a very demanding setting and indeed one of the main results in [BH12] is a factor- $\sqrt[3]{n}$ inapproximability bound for submodular functions. This notion of approximation is also considered in subsequent works of Badanidiyuru et al. [BDFKNR12] and Balcan et al. [BCIW12] where upper and lower approximation bounds are given for other related classes of functions such as XOS and subadditive. We emphasize that these strong lower bounds rely on a very specific distribution concentrated on a sparse set of points, and show that this setting is very different from uniform/product distributions which are the focus of this paper.

Gupta et al. [GHRU11] motivate learning of submodular functions over the uniform distribution by problems in differentially-private data release. They show that submodular functions with range $[0, 1]$ are ϵ -approximated by a collection of $n^{O(1/\epsilon^2)}$ ϵ^2 -Lipschitz submodular functions. Each ϵ^2 -Lipschitz submodular function can be ϵ -approximated by a constant. This leads to a learning algorithm running in time $n^{O(1/\epsilon^2)}$, which however requires value oracle access to the target function, in order to build the collection.

The work of Cheraghchi et al. [CKKL12] studies approximations of submodular functions by low-degree polynomials. They prove that any submodular function (of unit norm) can be ϵ -approximated in ℓ_1 by a polynomial of degree $O(1/\epsilon^2)$. This leads again to an $n^{O(1/\epsilon^2)}$ -time algorithm, but one which requires only random examples and works even in the agnostic setting. The main tool used in this work is the notion of noise stability. Feldman and Vondrák [FV16] studied approximation of submodular, XOS and self-bounding functions by juntas. Their main result shows that submodular functions can be approximated in ℓ_2 by a junta of size $\tilde{O}(1/\epsilon^2)$ and further that all self-bounding functions can be approximated by a junta of size $2^{O(1/\epsilon^2)}$.

Subsequently, Feldman and Vondrák [FV15] have obtained tight bounds on the degree of a polynomial that is sufficient to approximate any function in each of these function classes in ℓ_2 norm. Specifically, they showed $\tilde{O}(\epsilon^{-4/5})$ bound for submodular functions, $\Theta(1/\epsilon)$ bound for XOS functions and a matching lower bound of $\Omega(1/\epsilon^2)$ for self-bounding functions. The degree bound for XOS functions also implies an upper bound of $2^{O(1/\epsilon)}$ on the size of the junta sufficient to approximate (in ℓ_2) any XOS function.

Raskhodnikova and Yaroslavtsev [RY13] consider learning and testing of submodular functions taking values in the range $\{0, 1, \dots, k\}$ (referred to as *pseudo-Boolean*). The error of a hypothesis in their framework is the probability that the hypothesis disagrees with the unknown function. They build on the approach from [GHRU11] to show that pseudo-Boolean submodular functions can be expressed as $2k$ -DNF and then give a $\text{poly}(n) \cdot k^{O(k \log k/\epsilon)}$ -time PAC learning algorithm using value queries. Blais et al. [BOSY13] proved existence of a junta of size $(k \log(1/\epsilon))^{O(k)}$ and used it to give an algorithm for testing submodularity using $(k \log(1/\epsilon))^{\tilde{O}(k)}$ value queries. Feldman and Vondrák [FV16] and, more recently, Blais and Bommireddi [BB17] have studied testing of various type of valuation functions showing that approximation by a junta can be exploited to get efficient testing algorithms.

2 Preliminaries

2.1 Submodular, subadditive and self-bounding functions

In this section, we define the relevant classes of functions. We refer the reader to [Von10; FV16] for more details.

Definition 2.1. A set function $f : 2^N \rightarrow \mathbb{R}$ is

- *monotone*, if $f(A) \leq f(B)$ for all $A \subseteq B \subseteq N$.
- *submodular*, if $f(A \cup B) + f(A \cap B) \leq f(A) + f(B)$ for all $A, B \subseteq N$.
- *fractionally subadditive*, if $f(A) \leq \sum \beta_i f(B_i)$ whenever $\beta_i \geq 0$ and $\sum_{i:a \in B_i} \beta_i \geq 1 \forall a \in A$.

Submodular functions are not necessarily nonnegative, but in many applications (especially when considering multiplicative approximations), this is a natural assumption. Fractionally subadditive functions are nonnegative by definition (by considering $A = B_1, \beta_1 > 1$). In this paper we work exclusively with functions $f : 2^N \rightarrow \mathbb{R}_+$.

Next, we introduce *a-self-bounding functions*. Self-bounding functions were defined by Boucheron et al. [BLM00] as a unifying class of functions that enjoy strong “dimension-free” concentration properties. Currently this is the most general class of functions known to satisfy such concentration bounds. Self-bounding functions are defined generally on product spaces X^n ; here we restrict our attention to the hypercube, so the reader can assume that $X = \{0, 1\}$. We identify functions on $\{0, 1\}^n$ with set functions on $N = [n]$ in a natural way. Here we define a somewhat more general class of *a-self-bounding functions*, following [MR06].

Definition 2.2. A function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ is *a-self-bounding*, if for all $x \in \{0, 1\}^n$ and $i \in [n]$,

$$f(x) - \min_{x_i} f(x) \leq 1$$

and

$$\sum_{i=1}^n (f(x) - \min_{x_i} f(x)) \leq a f(x).$$

Useful properties of *a-self-bounding functions* that are easy to verify is that they are closed under taking max operation and closed under taking convex combinations. A particular example of a self-bounding function (related to applications of Talagrand’s inequality) is a function with the property of *small certificates*: $f : X^n \rightarrow \mathbb{Z}_+$ has small certificates, if it is 1-Lipschitz and whenever $f(x) \geq k$, there is a set of coordinates $S \subseteq [n]$, $|S| = k$, such that if $y|_S = x|_S$, then $f(y) \geq k$. Such functions often arise in combinatorics, by defining $f(x)$ to equal the maximum size of a certain structure appearing in x . In Section 5 we also show that k -DNF formulas are k -self-bounding.

2.2 Fourier analysis on the Boolean cube

The ℓ_1 and ℓ_2 -norms of a $f : \{0, 1\}^n \rightarrow \mathbb{R}$ are defined by $\|f\|_1 = \mathbf{E}_{x \sim \mathcal{U}}[|f(x)|]$ and $\|f\|_2 = (\mathbf{E}_{x \sim \mathcal{U}}[f(x)^2])^{1/2}$, respectively, where \mathcal{U} is the uniform distribution over $\{0, 1\}^n$. In what follows all probabilities and expectations are relative to \mathcal{U} unless explicitly specified otherwise.

We rely on the standard Fourier transform representation of real-valued functions over $\{0, 1\}^n$ as linear combinations of parity functions. For $S \subseteq [n]$, the parity function $\chi_S : \{0, 1\}^n \rightarrow \{-1, 1\}$ is defined by $\chi_S(x) = (-1)^{\sum_{i \in S} x_i}$. The Fourier expansion of f is given by $f(x) = \sum_{S \subseteq [n]} \hat{f}(S) \chi_S(x)$.

The degree of the highest degree non-zero Fourier coefficient of f is referred to as the *Fourier degree* of f . Note that Fourier degree of f is exactly the polynomial degree of f when viewed over $\{-1, 1\}^n$ instead of $\{0, 1\}^n$ and therefore it is also equal to the polynomial degree of f over $\{0, 1\}^n$. Let $f : \{0, 1\}^n \rightarrow \mathbb{R}$ and $\hat{f} : 2^{[n]} \rightarrow \mathbb{R}$ be its Fourier transform.

Definition 2.3 (The noise operator). *For $\rho \in [-1, +1]$, $x \in \{0, 1\}^n$, we define a distribution $N_\rho(x)$ over $y \in \{0, 1\}^n$ by letting $y_i = x_i$ with probability $\frac{1+\rho}{2}$ and $y_i = 1 - x_i$ with probability $\frac{1-\rho}{2}$, independently for each i . The noise operator T_ρ acts on functions $f : \{0, 1\}^n \rightarrow \mathbb{R}$, and is defined by*

$$(T_\rho f)(x) = \mathbf{E}_{y \sim N_\rho(x)} [f(y)].$$

The noise stability of f at noise rate ρ is

$$\mathbb{S}_\rho(f) = \langle f, T_\rho f \rangle = \mathbf{E}[f(x)T_\rho f(x)].$$

In terms of Fourier coefficients, the noise operator acts as $\widehat{T_\rho f}(S) = \rho^{|S|} \hat{f}(S)$. Therefore, noise stability can be written as $\mathbb{S}_\rho(f) = \sum_{S \subseteq [n]} \rho^{|S|} \hat{f}^2(S)$. Finally, we define noise sensitivity that generalizes the notion of noise sensitivity for Boolean functions.

Definition 2.4 (Noise sensitivity). *For $\delta \in [0, 1]$ and a function $f : \{0, 1\}^n \rightarrow \mathbb{R}$, the noise sensitivity of f at δ is $\text{NS}_\delta(f) = \frac{1}{2} \|f - T_{1-2\delta} f\|_1 = \frac{1}{2} \mathbf{E}[|f(x) - T_{1-2\delta} f(x)|]$.*

We keep the factor $1/2$ in the definition for consistency with the Boolean case. In the Boolean case noise sensitivity has the following relationship to noise stability (e.g. [O'D13]):

$$\text{NS}_\delta(f) = \frac{1}{2} (1 - \mathbb{S}_{1-2\delta}(f)).$$

Definition 2.5 (Discrete derivatives). *For $x \in \{0, 1\}^n$, $b \in \{0, 1\}$ and $i \in [n]$ let $x_{i \leftarrow b}$ denote the vector in $\{0, 1\}^n$ that equals to x with i -th coordinate set to b . For a real-valued $f : \{0, 1\}^n \rightarrow \mathbb{R}$ and indices $i, j \in [n]$ we define, $\partial_i f(x) = \frac{1}{2}(f(x_{i \leftarrow 1}) - f(x_{i \leftarrow 0}))$. We also define $\partial_{i,j} f(x) = \partial_i \partial_j f(x)$.*

Observe that $\partial_i f(x) = \sum_{S \ni i} \hat{f}(S) \chi_{S \setminus \{i\}}(x)$, and $\partial_{i,j} f(x) = \sum_{S \ni i,j} \hat{f}(S) \chi_{S \setminus \{i,j\}}(x)$.

We use several notions of *influence* of a variable on a real-valued function which are based on the standard notion of influence for Boolean functions (e.g. [BOL85; KKL88]).

Definition 2.6 (Influences). *For a real-valued $f : \{0, 1\}^n \rightarrow \mathbb{R}$, $i \in [n]$, and $\kappa \geq 0$ we define the ℓ_κ^κ -influence of variable i as $\text{Inf}_i^\kappa(f) = \|\partial_i f\|_\kappa^\kappa = \mathbf{E}[|\partial_i f|^\kappa]$. We define $\text{Inf}^\kappa(f) = \sum_{i \in [n]} \text{Inf}_i^\kappa(f)$ and refer to it as the total ℓ_κ^κ -influence of f .*

3 Structural results

3.1 Approximation of low-sensitivity functions by low-degree polynomials

In this section we demonstrate a simple approach that allows to approximate low noise-sensitive functions in ℓ_1 norm and also show that noise sensitivity of a function can be upper-bounded by its ℓ_1 influence.

Our approach is based on an observation that if a function is close to its noisy version in ℓ_1 norm then it is well-approximated by a low-degree polynomial.

Lemma 3.1. For every function $f : \{0, 1\}^n \rightarrow \mathbb{R}$, every $\epsilon > 0$ and $\delta \in (0, 1]$ there exists a multilinear polynomial p of degree $d = \lceil \frac{1}{2\delta} \log \frac{1}{\epsilon} \rceil$ such that

$$\|f - p\|_1 \leq \epsilon \|f\|_2 + 2 \cdot \text{NS}_\delta(f).$$

In particular, the polynomial can be chosen as $p(x) = \sum_{|S| < d} (1 - 2\delta)^{|S|} \hat{f}(S) \chi_S(x)$.

Proof. Let $\rho = 1 - 2\delta$. We can estimate the tail of the Fourier expansion as follows: For any d , define $f_{<d}(x) = \sum_{S: |S| < d} \hat{f}(S) \chi_S(x)$, a polynomial of degree at most d . Then, since $T_\rho f(x) = \sum_{S \subseteq [n]} \rho^{|S|} \hat{f}(S) \chi_S(x)$, we get

$$\|T_\rho f_{<d} - T_\rho f\|_1 = \left\| \sum_{S: |S| \geq d} \rho^{|S|} \hat{f}(S) \chi_S \right\|_1 \leq \left\| \sum_{S: |S| \geq d} \rho^{|S|} \hat{f}(S) \chi_S \right\|_2 \leq \rho^d \|f\|_2. \quad (1)$$

Taking $d = \lceil \frac{1}{2\delta} \log \frac{1}{\epsilon} \rceil$ we get that such that $\|T_\rho f_{<d} - T_\rho f\|_1 \leq (1 - 2\delta)^d \cdot \|f\|_2 \leq \epsilon \|f\|_2$. Now, by Definition 2.4, we have that $\|f - T_{1-2\delta} f\|_1 = 2 \cdot \text{NS}_\delta(f)$. The lemma now follows by the triangle inequality. \square

Next we observe that the total ℓ_1 influence of a function can be used to derive an upper-bound on its noise sensitivity.

Lemma 3.2. For every function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ and $\delta \in [0, 1]$,

$$\text{NS}_\delta(f) \leq \delta \cdot \text{Inf}^1(f).$$

Proof. For every $t = 0, 1, \dots, n$ we define a distribution $N_{1-2\delta}^{1:t}(x)$ over $y \in \{0, 1\}^n$ by letting $y_i = x_i$ with probability $1 - \delta$ and $y_i = 1 - x_i$ with probability δ , independently for each $i \leq t$, while for all $i > t$, $y_i = x_i$. Note that $N_{1-2\delta}^{1:0}(x)$ is always equal to x and $N_{1-2\delta}^{1:n}(x)$ is exactly $N_{1-2\delta}(x)$. We also define a distribution $N_{1-2\delta}^t(x)$ over $y \in \{0, 1\}^n$ by letting $y_y = x_t$ with probability $1 - \delta$ and $y_t = 1 - x_t$ with probability δ , while for all $i \neq t$, $y_i = x_i$.

Now,

$$\begin{aligned} \text{NS}_\delta(f) &= \frac{1}{2} \cdot \mathbf{E}[|f(x) - T_{1-2\delta} f(x)|] = \frac{1}{2} \cdot \mathbf{E} \left[\left| f(x) - \mathbf{E}_{y \sim N_{1-2\delta}(x)} [f(y)] \right| \right] \\ &\leq \frac{1}{2} \sum_{t=1}^n \mathbf{E} \left[\left| \mathbf{E}_{y \sim N_{1-2\delta}^{1:t-1}(x)} [f(y)] - \mathbf{E}_{y \sim N_{1-2\delta}^{1:t}(x)} [f(y)] \right| \right] \\ &\leq \frac{1}{2} \sum_{t=1}^n \mathbf{E}_{y \sim N_{1-2\delta}^{1:t-1}(x)} \left[\left| f(y) - \mathbf{E}_{z \sim N_{1-2\delta}^t(y)} [f(z)] \right| \right] \\ &= \frac{1}{2} \sum_{t=1}^n \mathbf{E}_{y \sim \mathcal{U}} \left[\left| f(y) - \mathbf{E}_{z \sim N_{1-2\delta}^t(y)} [f(z)] \right| \right] \\ &= \frac{1}{2} \sum_{t=1}^n \mathbf{E}_{y \sim \mathcal{U}} [2\delta |\partial_t f(y)|] = \delta \sum_{t=1}^n \text{Inf}_t^1(f) = \delta \cdot \text{Inf}^1(f). \end{aligned}$$

\square

An immediate corollary of Lemmas 3.1 and 3.2 is that any function of low total ℓ_1 influence can be well-approximated by a low-degree polynomial:

Corollary 3.3. *For every function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ such that $\|f\|_2 \leq 1$ and every $\epsilon > 0$ there exists a multilinear polynomial p of degree $d = \lceil \frac{2 \cdot \text{Inf}^1(f)}{\epsilon} \log \frac{2}{\epsilon} \rceil$ such that $\|f - p\|_1 \leq \epsilon$.*

It follows easily from the definition of self-bounding functions that they have low total ℓ_1 -influence.

Lemma 3.4 ([FV16], Lemma 4.2). *Let $f : \{0, 1\}^n \rightarrow \mathbb{R}_+$ be an a -self-bounding function. Then $\text{Inf}^1(f) \leq a \cdot \|f\|_1$. In particular, for $f : \{0, 1\}^n \rightarrow [0, 1]$, $\text{Inf}^1(f) \leq a$.*

Therefore we obtain that self-bounding functions are well-approximated by low-degree polynomials.

Theorem 3.5. *For every a -self-bounding function $f : \{0, 1\}^n \rightarrow \mathbb{R}_+$ and every $\epsilon > 0$, there exists a multilinear polynomial p of degree $d = \lceil \frac{2a}{\epsilon} \log \frac{2}{\epsilon} \rceil$ such that*

$$\|f - p\|_1 \leq \epsilon \|f\|_2.$$

In particular, the polynomial can be chosen as $p(x) = \sum_{|S| < d} \rho^{|S|} \hat{f}(S) \chi_S(x)$, for $\rho = 1 - \frac{\epsilon}{2a}$.

Application to approximation and learning of halfspaces. We now briefly show that our approach can also be used to obtain sharper bounds on ℓ_1 -approximation of halfspaces by low-degree polynomials. Recall that a halfspace is a Boolean function expressible as $\text{sign}(\sum_{i \in [n]} a_i x_i - a_0)$ for some real values a_0, a_1, \dots, a_n . Halfspaces are known to be noise-stable. Specifically, Kalai et al. [KKMS08] proved that for every halfspace f and $\delta > 0$, $\text{NS}_\delta(f) \leq 8.8 \cdot \sqrt{\delta}$. Using this fact they showed that any halfspace can be ϵ -approximated in ℓ_2 norm by a polynomial of degree $O(1/\epsilon^4)$ and gave an agnostic learning algorithm for learning halfspaces over the uniform distribution that runs in time $n^{O(1/\epsilon^4)}$. For ℓ_1 norm approximation the best previously known bound is $O(\log^2(1/\epsilon)/\epsilon^2)$ and was given by Diakonikolas et al. [DGJSV10] (note however that their result is substantially more involved and gives a stronger notion of approximation that is necessary for fooling halfspaces). By plugging the upper bound on noise sensitivity into our Lemma 3.1 with $\delta = (4 \cdot 8.8)^{-2} \cdot \epsilon^2$ we obtain the following corollary:

Corollary 3.6. *For every halfspace f and every $\epsilon > 0$, there exists a multilinear polynomial p of degree $d = O(\log(1/\epsilon)/\epsilon^2)$ such that $\|f - p\|_1 \leq \epsilon$.*

We note that the agnostic learning algorithm for halfspaces in [KKMS08] requires only ℓ_1 approximation. Therefore our result implies that halfspaces are agnostically learnable over the uniform distribution in time $n^{O(\log(1/\epsilon)/\epsilon^2)}$.

3.2 Noise stability of self-bounding functions

In this section, we study the action of the noise operator on a self-bounding function in more detail. Specifically, we show that self-bounding functions are noise-stable point-wise. This result strengthens and generalizes a similar one proved for submodular functions in [CKKL12]. It allows us to derive additional properties of self-bounding functions useful for their approximation and learning.

Lemma 3.7. *For any a -self-bounding function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ under the uniform distribution, $a \geq 1$, and any $\rho \in [-1, +1]$, $x \in \{0, 1\}^n$,*

$$T_\rho f(x) \geq \left(\left(1 - \frac{1 - \rho}{2(1 - \frac{a-1}{n})} \right)_+ \right)^a f(x).$$

(We denote by $(z)_+ = \max\{z, 0\}$ the positive part of a real number.)

Proof. First, let us observe that the statement of the lemma is invariant under flipping the hypercube $\{0, 1\}^n$ along any coordinate: the notion of a -self-bounding functions does not change, the action of the noise operator does not change, and the conclusion of the lemma does not change either. So we can assume without loss of generality that $x = (0, 0, \dots, 0)$. We also identify points in $\{0, 1\}^n$ with sets $S \subseteq [n]$ by considering $S = \{i : x_i = 1\}$.

Let us average the values of f over levels of sets of constant $|S|$, and define

$$\phi(t) = \mathbf{E}_{|S|=t} [f(S)] = \frac{1}{\binom{n}{t}} \sum_{S: |S|=t} f(S).$$

In particular, $\phi(0) = f(\emptyset) = f(x)$. We claim the following: for every $t = 0, 1, \dots, n$,

$$\phi(t) \geq \left(\left(1 - \frac{t}{n - a + 1} \right)_+ \right)^a \phi(0). \quad (2)$$

Intuitively, if $f(x)$ is a point of high value, the value cannot drop off too quickly as we move away from x . If we prove (2), then we are done: for $x = (0, 0, \dots, 0)$, $T_\rho f(x)$ is an expectation of $f(S)$ over a distribution where the sets on each level appear with the same probability, namely

$$T_\rho f(x) = \sum_{i=0,1,\dots,n} \left(\frac{1 - \rho}{2} \right)^i \cdot \left(\frac{1 + \rho}{2} \right)^{n-i} \cdot \binom{n}{i} \cdot \phi(i).$$

The expected cardinality of a set sampled from this distribution is $\mathbf{E}[|S|] = \frac{1-\rho}{2}n$. By convexity of the bound in (2) and Jensen's inequality, we obtain

$$T_\rho f(x) \geq \left(\left(1 - \frac{\frac{1-\rho}{2}n}{n - a + 1} \right)_+ \right)^a f(x) = \left(\left(1 - \frac{1 - \rho}{2(1 - \frac{a-1}{n})} \right)_+ \right)^a f(x).$$

So it remains to prove (2).

We proceed by induction. For $t = 0$, the claim is trivial. Let us assume it holds for t , and consider a set S , $|S| = t$. We also assume that $t < n - a + 1$, because otherwise the claim is trivial (recall that f and hence ϕ is nonnegative). By the a -self-bounding property, we have

$$af(S) \geq \sum_{i=1}^n (f(S) - \min\{f(S+i), f(S-i)\}) \geq \sum_{i \in [n] \setminus S} (f(S) - f(S+i)).$$

Note that $|[n] \setminus S| = n - t$. By rearranging this inequality, we get

$$(n - t - a)f(S) \leq \sum_{i \in [n] \setminus S} f(S+i).$$

Now let us add up this inequality over all S of size $|S| = t$:

$$(n - t - a) \sum_{|S|=t} f(S) \leq \sum_{|S|=t, i \notin S} f(S + i) = (t + 1) \sum_{|S'|=t+1} f(S')$$

because every set S' of size $t + 1$ appears $t + 1$ times in the penultimate summation. Expressing this inequality in terms of $\phi(t)$, we get

$$(n - t - a) \binom{n}{t} \phi(t) \leq (t + 1) \binom{n}{t + 1} \phi(t + 1),$$

or equivalently (for $t < n - a$)

$$\phi(t) \leq \frac{n - t}{n - t - a} \phi(t + 1).$$

We replace this by a slightly weaker bound: $\phi(t) \leq \left(\frac{n-t-a+1}{n-t-a}\right)^a \phi(t + 1)$. To see why this holds, consider $\left(\frac{n-t-a+1}{n-t-a}\right)^a = \left(1 + \frac{1}{n-t-a}\right)^a \geq 1 + \frac{a}{n-t-a} = \frac{n-t}{n-t-a}$.

By the inductive hypothesis (2), we assume $\phi(t) \geq \left(\frac{n-a+1-t}{n-a+1}\right)^a \phi(0)$. So we obtain

$$\left(\frac{n - a + 1 - t}{n - a + 1}\right)^a \phi(0) \leq \left(\frac{n - t - a + 1}{n - t - a}\right)^a \phi(t + 1).$$

This holds for $t < n - a$, hence proving the bound in (2) for $t < n - a + 1$:

$$\phi(t) \geq \left(\frac{n - a - t + 1}{n - a + 1}\right)^a \phi(0) = \left(1 - \frac{t}{n - a + 1}\right)^a \phi(0).$$

As we mentioned, (2) is trivially true for $t \geq n - a + 1$. □

Corollary 3.8. *For any a -self-bounding function $f : \{0, 1\}^n \rightarrow \mathbb{R}$ under the uniform distribution, the noise stability with noise parameter ρ is*

$$\mathbb{S}_\rho(f) \geq \left(1 - \frac{1 - \rho}{2(1 - \frac{a-1}{n})}\right)^a \|f\|_2^2.$$

In particular, for $a = 1$ (self-bounding functions), we obtain $\mathbb{S}_\rho(f) \geq \frac{1+\rho}{2} \|f\|_2^2$. In [CKKL12], an analogous bound on noise stability is used to derive an agnostic learning algorithm (over the uniform distribution) with excess ℓ_1 -error ϵ in time $n^{O(1/\epsilon^2)}$.

Comparison of norms for self-bounding functions. Our bound on the noise operator implies a bound on the ℓ_1 norm of a self-bounding function, relative to its ℓ_∞ norm. This has been first shown for submodular functions by Feige et al. [FMV07] and for XOS functions by Feige [Fei06] (with a constant a). Feldman and Vondrák [FV16] show how this property together with approximation by a junta can be used to obtain a learning algorithm with multiplicative approximation guarantees that are required in the PMAC model of Balcan and Harvey [BH12]. Hence our results can be used to extend the PMAC learning algorithm in [FV16] from XOS functions to all a -self-bounding functions. The full details of this construction are relatively involved and therefore we direct the interested reader to the discussion in [FV16](Sec. 6) for additional details.

Lemma 3.9. *For any a -self-bounding function $f : \{0, 1\}^n \rightarrow \mathbb{R}_+$ under the uniform distribution, with $n \geq 4a$,*

$$\|f\|_1 \leq \|f\|_\infty \leq 3^a \|f\|_1.$$

Proof. Let $\|f\|_\infty = f(x^*)$. Since f is nonnegative and $n \geq 4a$, we have by Lemma 3.7

$$\|f\|_1 = \mathbf{E}[f(x)] = T_0 f(x^*) \geq \left(1 - \frac{1}{2(1 - \frac{a-1}{n})}\right)^a f(x^*) \geq \left(1 - \frac{1}{2 \cdot 3/4}\right)^a f(x^*) = \frac{1}{3^a} f(x^*).$$

□

We remark that a factor exponential in a is necessary here. Consider the conjunction function on a variables, $f(x) = x_1 x_2 \cdots x_a$. This is an a -self-bounding function with values in $\{0, 1\}$. We have

$$\|f\|_p = (\Pr[f(x) = 1])^{1/p} = 2^{-a/p}.$$

In particular, $\|f\|_1 = 2^{-a}$, $\|f\|_2 = 2^{-a/2}$ and $\|f\|_\infty = 1$; i.e., the ℓ_1 , ℓ_2 and ℓ_∞ norms can differ by factors exponential in a .

Relative error vs. additive error. In our results, we typically assume that the values of $f(x)$ are in a bounded interval $[0, 1]$ or that $\|f\|_1 \leq 1$ and our goal is to approximate f with an additive error of ϵ . As Lemma 3.9 shows, for a -self-bounding functions (with constant a) the ℓ_1 and ℓ_∞ norms are within a bounded factor, so this does not make much difference.

This means that if we scale $f(x)$ by $1/(3^a \|f\|_1)$, we obtain a function with values in $[0, 1]$. Approximating this function within an additive error of ϵ is equivalent to approximating the original function within an error of $\epsilon 3^a \|f\|_1$. In particular, for submodular functions we have $a = 2$. Hence, the two settings are equivalent up to a constant factor in the error and we state our results for submodular functions in the interval $[0, 1]$.

3.3 Friedgut's theorem for ℓ_1 -approximation

As we have mentioned in Lemma 3.4, self-bounding functions have low total sensitivity. A celebrated result of Friedgut [Fri98] shows that any Boolean function on $\{0, 1\}^n$ of low average sensitivity is close to a function that depends on few variables. His result was extended to ℓ_2 approximation of real-valued functions in [FV16]. We now show that for self-bounding functions a tighter bounds can be achieved for ℓ_1 approximation. Our proof is based on the use of ℓ_1 approximation by polynomials proved in Theorem 3.5 together with the analysis from [FV16] to obtain a smaller ℓ_1 approximating junta.

We now state the main result in more detail.

Theorem 3.10. *Let $f : \{0, 1\}^n \rightarrow [0, 1]$ be a function and $a = \text{Inf}^1(f)$. For every $\epsilon > 0$, let $d = \lceil \frac{4a}{\epsilon} \log \frac{4}{\epsilon} \rceil$ and $I = \{i \in [n] \mid \text{Inf}_i^{4/3}(f) \geq \alpha\}$ for $\alpha = 3^{-2d-1} \epsilon^4 / a^2$. Then $|I| \leq a/\alpha$ and there exists a polynomial p of degree d over variables in I such that $\|f - p\|_1 \leq \epsilon$.*

To prove the theorem we will need the following bound on the sum of squares of all low-degree Fourier coefficients that include a variable of low influence from [FV16].

Lemma 3.11 ([FV16], Lemma 4.7). *Let $f : \{0, 1\}^n \rightarrow \mathbb{R}$, $\kappa \in (1, 2)$, $\alpha > 0$ and d be an integer ≥ 1 . Let $I = \{i \in [n] \mid \text{Inf}_i^\kappa(f) \geq \alpha\}$. Then*

$$\sum_{S \not\subseteq I, |S| \leq d} \hat{f}(S)^2 \leq (\kappa - 1)^{1-d} \cdot \alpha^{2/\kappa-1} \cdot \text{Inf}^\kappa(f) .$$

We can now complete the proof of Thm. 3.10.

Proof. Theorem 3.5 proves that for $d \leq \lceil \frac{4a}{\epsilon} \log \frac{4}{\epsilon} \rceil$ and $\rho = 1 - \frac{\epsilon}{2a}$, the function $T_\rho f_{<d}$ satisfies

$$\|f - T_\rho f_{<d}\|_1 \leq \epsilon \|f\|_2 / 2 \leq \epsilon / 2. \quad (3)$$

We can also apply Lemma 3.11 with $\kappa = 4/3$ and $\alpha = 3^{-2d-1} \epsilon^4 / a^2$ to obtain that

$$\sum_{S \not\subseteq I, |S| \leq d} \hat{f}(S)^2 \leq 3^{d-1} \cdot \alpha^{1/2} \cdot \text{Inf}^{4/3}(f) = 3^{d-1} \cdot \left(3^{-d-1/2} \cdot \frac{\epsilon^2}{a} \right) \cdot \text{Inf}^{4/3}(f) \leq \frac{\epsilon^2}{4}, \quad (4)$$

where the last inequality uses $\text{Inf}^{4/3}(f) \leq \text{Inf}^1(f) \leq a$ which follows from the fact that $\partial_i f$'s have range $[-1/2, 1/2]$ when f has range $[0, 1]$.

For every S , $|\widehat{T_\rho f}(S)| = |\rho^{|S|} \hat{f}(S)| \leq |\hat{f}(S)|$. Therefore eq. (4) implies that

$$\sum_{S \not\subseteq I, |S| \leq d} \widehat{T_\rho f}(S)^2 \leq \sum_{S \not\subseteq I, |S| \leq d} \hat{f}(S)^2 \leq \frac{\epsilon^2}{4}. \quad (5)$$

Now let $p = \sum_{S \subseteq I, |S| \leq d} \widehat{T_\rho f}(S) \chi_S$ be the restriction of $T_\rho f_{<d}$ to variables in I . Equation (5) gives a bound on the sum of squares of all the coefficients that we removed from $T_\rho f_{<d}$ and implies that $\|p - T_\rho f_{<d}\|_1 \leq \|p - T_\rho f_{<d}\|_2 \leq \epsilon / 2$. Together with eq. (3), we get $\|f - p\|_1 \leq \epsilon$. Finally, $|I| \leq \text{Inf}^{4/3}(f) / \alpha \leq \text{Inf}^1(f) / \alpha \leq a / \alpha$. \square

By Lemma 3.4, every a -self-bounding function $f : \{0, 1\}^n \rightarrow [0, 1]$ satisfies, $\text{Inf}^1(f) \leq a$. Hence as an immediate corollary we obtain Thm. 1.1. Another immediate corollary of Thm. 3.10 is that for every a -self-bounding function there exists a polynomial of low total ℓ_1 -spectral norm that approximates it.

Corollary 3.12. *Let $f : \{0, 1\}^n \rightarrow [0, 1]$ be an a -self-bounding function and $\epsilon > 0$. There exist $d = O(a/\epsilon \cdot \log(1/\epsilon))$ and a polynomial p of degree d such that $\|f - p\|_1 \leq \epsilon$ and $\|\hat{p}\|_1 = 2^{O(d^2)}$, where $\|\hat{p}\|_1 = \sum_{S \subseteq [n]} |\hat{p}(S)|$.*

4 Algorithmic applications

We now outline the applications of our structural results. They are based on using our stronger bounds in existing learning algorithms for submodular, XOS and self-bounding functions. We omit the details of the algorithms and their analysis since they follow closely those of the corresponding results in [FV16].

4.1 Learning Models

Our learning algorithms are in one of two standard models of learning. The first one assumes that the learner has access to random examples of an unknown function from a known set of functions. This model can be seen as a generalization of Valiant’s PAC learning model to real-valued functions [Val84]. While in general Valiant’s model does not make assumptions on the distribution \mathcal{D} , here we only consider the *distribution-specific* version of the model in which the distribution is fixed and is uniform over $\{0, 1\}^n$.

Definition 4.1 (Distribution-specific ℓ_1 PAC learning). *Let \mathcal{F} be a class of real-valued functions on $\{0, 1\}^n$ and let \mathcal{D} be a distribution on $\{0, 1\}^n$. An algorithm \mathcal{A} PAC learns \mathcal{F} on \mathcal{D} , if for every $\epsilon > 0$ and any target function $f \in \mathcal{F}$, given access to random independent samples from \mathcal{D} labeled by f , with probability at least $\frac{2}{3}$, \mathcal{A} returns a hypothesis h such that $\mathbf{E}_{x \sim \mathcal{D}}[|f(x) - h(x)|] \leq \epsilon$.*

Agnostic learning generalizes the definition of PAC learning to scenarios where one cannot assume that the input labels are consistent with a function from a given class [Hau92; KSS94] (for example as a result of noise in the labels).

Definition 4.2 (Distribution-specific ℓ_1 agnostic learning). *Let \mathcal{F} be a class of real-valued functions on $\{0, 1\}^n$ and let \mathcal{D} be any fixed distribution on $\{0, 1\}^n$. For any distribution \mathcal{D}' , let $\text{opt}(\mathcal{D}', \mathcal{F})$ be defined as:*

$$\text{opt}(\mathcal{D}', \mathcal{F}) = \inf_{f \in \mathcal{F}} \mathbf{E}_{(x,y) \sim \mathcal{D}'} [|y - f(x)|].$$

An algorithm \mathcal{A} , is said to agnostically learn \mathcal{F} on \mathcal{D} if for every excess error $\epsilon > 0$ and any distribution \mathcal{D}' on $\{0, 1\}^n \times \mathbb{R}$ such that the marginal of \mathcal{D}' on $\{0, 1\}^n$ is \mathcal{D} , given access to random independent examples drawn from \mathcal{D}' , with probability at least $\frac{2}{3}$, \mathcal{A} outputs a hypothesis h such that $\mathbf{E}_{(x,y) \sim \mathcal{D}'} [|h(x) - y|] \leq \text{opt}(\mathcal{D}') + \epsilon$.

The first corollary of our structural results is for PAC learning of monotone self-bounding functions (the results also apply to *unate* functions which are either monotone or anti-monotone in each variable). Note that this class of functions includes XOS functions.

Theorem 4.3. *Let \mathcal{C}_a^+ be the set of all monotone a -self-bounding functions on from $\{0, 1\}^n$ to $[0, 1]$. There exists an algorithm that PAC learns \mathcal{C}_a^+ over the uniform distribution, runs in time $\tilde{O}(n) \cdot 2^{\tilde{O}(a^2/\epsilon^2)}$ and uses $2^{\tilde{O}(a^2/\epsilon^2)} \log n$ examples, where ϵ is the error parameter.*

The proof of this result follows from substituting our bounds in Theorems 1.1 and 3.12 into the simple analysis from [FV16].

Our main application to agnostic learning is the algorithm for learning self-bounding functions from random examples described in Theorem 1.2. The algorithm used to prove this result is again polynomial ℓ_1 regression over all monomials of degree $\tilde{O}(a/\epsilon)$. In addition, we can rely on the existence of a polynomial of low spectral norm to obtain substantially tighter bounds on sample complexity. Namely, as in [FV16], we use the uniform convergence bounds for linear combinations of functions with ℓ_1 constraint on the sum of coefficients [KST08]. Our structural results also have immediate implications for learning with value queries, that is oracle access to the value of the unknown function at any point x . Following the approach from [FKV13], we can use the algorithm of Gopalan et al. [GKK08] together with our bounds on the spectral norm of the approximating polynomial in Cor. 3.12. This leads to the following algorithm.

Theorem 4.4. *Let \mathcal{C}_a be the class of all a -self-bounding functions from $\{0,1\}^n$ to $[0,1]$. There exists an agnostic learning algorithm that for any $\epsilon > 0$, given access to value queries learns \mathcal{C}_a with excess error $\epsilon > 0$ over the uniform distribution in time $\text{poly}(n) \cdot 2^{\tilde{O}(a^2/\epsilon^2)}$.*

5 Lower bounds for learning self-bounding functions

In this section, we show that learning a -self bounding functions within an error of at most ϵ , is at least as hard as learning the class of all DNFs (of any size) of width at most $\lfloor \frac{a}{4\epsilon} \rfloor$ to an accuracy of $\frac{1}{4}$. Our reduction to learning width k -DNFs (also referred to as k -DNFs) is based on the simple observation that k -DNFs are k -self bounding functions combined with a simple linear transformation that reduces approximation and learning of $(a \cdot r)$ -self bounding functions for $r \geq 1$ to that of a -self-bounding functions.

Lemma 5.1. *A function $f : \{0,1\}^n \rightarrow \{0,1\}$ computed by a k -DNF formula is a k -self bounding function.*

Proof. Since f is $\{0,1\}$ -valued, clearly, $f(x) - \min_{x_i} f(x) \leq 1$ for any $i \in [n]$. If $f(x) = 0$, then, $\sum_{i=1}^n (f(x) - \min_{x_i} f(x)) = 0 \leq k \cdot f(x)$. Now suppose $f(x) = 1$. Then, there exists at least one term, say T , of the DNF that is satisfied by the assignment x . Observe that if we flip a literal outside of T , then, the value of f remains unchanged. Thus, if the term indexed by j in $\sum_{i=1}^n (f(x) - \min_{x_i} f(x))$ contributes the value 1, then either $x_j \in T$ or $\bar{x}_j \in T$. In particular, at most k terms in the sum contribute 1 and the rest contribute 0. Thus, $\sum_{i=1}^n (f(x) - \min_{x_i} f(x)) \leq k = k \cdot f(x)$. \square

Remark 5.2. *In light of Lemma 5.1 it is natural to ask whether all Boolean k -self-bounding functions are k -DNF. It is easy to see that for Boolean functions being k -self-bounding can be equivalently stated as having 1-sensitivity of k . The smallest k for which f can be represented by a k -DNF is referred to as 1-certificate complexity of f . It has long been observed that for monotone functions 1-certificate complexity equals 1-sensitivity [Nis89] and therefore all monotone k -self-bounding functions are k -DNF. However this is no longer true for non-monotone functions. A simple example in [Nis89] gives a function with a factor two gap between these two measures. Quadratic gap for every k up to $\Theta(n^{1/3})$ is also known [Cha05].*

Next, we observe that for any a -self-bounding function, the function g defined by $g(x) = 1 - \frac{1}{r} + \frac{f(x)}{r}$ is $\frac{a}{r}$ -self-bounding whenever $r \geq 1$. This “lifting” transforms an a -self-bounding functions into an $\frac{a}{r}$ -self-bounding functions.

Lemma 5.3. *Let $f : \{0,1\}^n \rightarrow [0,1]$ be an a -self-bounding function. Then for any $r \geq 1$, $g(x) = 1 - \frac{1}{r} + \frac{f(x)}{r}$ has range $[0,1]$ and is $\frac{a}{r}$ -self-bounding.*

Proof. Clearly, the $1 - 1/r + f(x)/r$ transformation maps $[0,1]$ to $[1 - 1/r, 1] \subseteq [0,1]$. Observe that for any x and $i \in [n]$, $g(x) - \min_{x_i} g(x) = \frac{1}{r} \cdot (f(x) - \min_{x_i} f(x))$ and also that $g(x) \geq f(x)$. By the definition of a -self-boundedness we obtain that g is a/r -self bounding. \square

Observe that given random examples labeled by f , it is easy to simulate random examples labeled by g . Further, ℓ_1 -approximation of f within ϵ can be translated (via the same “lifting”) to ϵ/r -approximation of g and vice versa. An immediate corollary of this is that one can use a learning algorithm for a/r -self-bounding functions to learn a -self bounding functions. We use \mathcal{C}_a^n to denote the class of all a -self-bounding functions from $\{0,1\}^n$ to $[0,1]$.

Lemma 5.4. *Let $a \geq 1$ and $a \geq r \geq 1$. Suppose there is an algorithm that PAC (or agnostically) learns $\mathcal{C}_{a/r}^n$ over a distribution D with ℓ_1 error of ϵ in time $T(n, 1/\epsilon)$. Then, there is an algorithm that PAC (or, respectively, agnostically) learns \mathcal{C}_a^n over D with ℓ_1 error of ϵ in time $T(n, 1/(r\epsilon))$.*

The simple structural observations above give us our lower bounds for learning and approximation of a -self-bounding functions. Using Lemmas 5.1 and 5.4, we have the the following lower bound on the time required to PAC learn a -self-bounding functions.

Theorem 5.5 (Thm. 1.3 restated). *Suppose there exists an algorithm that PAC learns \mathcal{C}_a^n with ℓ_1 error of $\epsilon > 0$ with respect to the uniform distribution in time $T(n, 1/\epsilon)$. Then, for any $k \geq a$, there exists an algorithm that PAC learns k -DNF formulas with disagreement error of at most ϵ' with respect to the uniform distribution in time $T(n, \frac{k}{a\epsilon'})$. Consequently, there exists an algorithm for learning k -juntas on the uniform distribution to an error of at most $1/4$ in time $T(n, \frac{k}{4a})$ for any $k \geq a$.*

Now, k -juntas contain the set of all Boolean functions on any fixed subset of k variables. A standard information-theoretic lower bound implies that any algorithm that PAC learn k -juntas to an accuracy of $1/4$ on the uniform distribution needs $\Omega(2^k)$ random examples or even value queries. This translates into the following unconditional lower bound for learning a -self-bounding functions.

Corollary 5.6. *Any algorithm that PAC learns \mathcal{C}_a over the uniform distribution needs $\Omega(2^{a/\epsilon})$ random examples or value queries.*

Finally, observe that the $\{0, 1\}$ -valued parity function on k bits is computed by a k -DNF formula and any polynomial that $1/4$ -approximates in ℓ_1 distance on the uniform distribution must have degree at least k . Thus, we have the following degree lower bound for polynomials that ℓ_1 approximate a -self-bounding functions on the uniform distribution on $\{0, 1\}^n$.

Corollary 5.7. *Fix an $a \geq 1$ and $\epsilon \in (0, 1/4]$. There exists an a -self-bounding function $f : \{0, 1\}^n \rightarrow [0, 1]$, such that every polynomial p that ϵ -approximates f in ℓ_1 norm with respect to the uniform distribution has degree $d \geq a/(4\epsilon)$.*

Proof. Let $k = \frac{a}{4\epsilon}$ (ignoring rounding issues for simplicity) and f be a $\{0, 1\}$ -valued parity on some set of k variables. By Lemma 5.1 f is k -self-bounding. Then, as in the proof of Lemma 5.4, for $r = \frac{1}{4\epsilon} \geq 1$, g defined by $g(x) = 1 - \frac{1-f(x)}{r}$ is an a -self-bounding function. Let p be a polynomial of degree d that approximates g within an ℓ_1 error of ϵ with respect to the uniform distribution on $\{0, 1\}^n$. Then, as in the proof of Lemma 5.4, $p' = 1 - r(1 - p)$ is a polynomial of degree d and approximates f within an ℓ_1 error of at most $\frac{1}{4\epsilon} \cdot \epsilon = 1/4$.

For the $\{-1, 1\}$ -valued parity $\chi = 2f(x) - 1$ and any polynomial p' of degree less than k , $\mathbf{E}[\chi \cdot p'] = 0$. Further, $\mathbf{E}[|\chi - p'|] \geq 1 - \mathbf{E}[\chi \cdot p'] = 1$. This implies that for f the ℓ_1 error of any polynomial of degree at most $k - 1$ is at least $1/2$. In particular, $d \geq a/(4\epsilon)$. \square

We remark that slightly weaker versions of Cor. 5.6 and Cor. 5.7 are known for monotone submodular functions. Specifically, they require $2^{\Omega(\epsilon^{-2/3})}$ random examples or value queries to PAC learn and also degree $\Omega(\epsilon^{-2/3})$ to approximate [FKV13].

References

- [BB17] E. Blais and A. Bommireddi. “Testing submodularity and other properties of valuation functions”. In: *Innovations in Theoretical Computer Science (ITCS)*. 2017.
- [BCIW12] M. Balcan, F. Constantin, S. Iwata, and L. Wang. “Learning Valuation Functions”. In: *Journal of Machine Learning Research - COLT Proceedings* 23 (2012), pp. 4.1–4.24.
- [BDFKNR12] A. Badanidiyuru, S. Dobzinski, H. Fu, R. Kleinberg, N. Nisan, and T. Roughgarden. “Sketching valuation functions”. In: *SODA*. 2012, pp. 1025–1035.
- [BFJKMR94] A. Blum, M. Furst, J. Jackson, M. Kearns, Y. Mansour, and S. Rudich. “Weakly learning DNF and characterizing statistical query learning using Fourier analysis”. In: *STOC*. Montreal, Quebec, Canada, 1994, pp. 253–262.
- [BH12] M. Balcan and N. Harvey. “Submodular Functions: Learnability, Structure, and Optimization”. In: *CoRR* abs/1008.2159 (2012). Earlier version in proceedings of STOC 2011.
- [BL97] A. Blum and P. Langley. “Selection of Relevant Features and Examples in Machine Learning”. In: *Artificial Intelligence* 97.1-2 (1997), pp. 245–271.
- [BLM00] S. Boucheron, G. Lugosi, and P. Massart. “A sharp concentration inequality with applications”. In: *Random Struct. Algorithms* 16.3 (2000), pp. 277–292.
- [Blu03] A. Blum. “Open Problem: Learning a Function of r Relevant Variables.” In: *COLT*. 2003, pp. 731–733.
- [BOL85] M. Ben-Or and N. Linial. “Collective Coin Flipping, Robust Voting Schemes and Minima of Banzhaf Values”. In: *FOCS*. 1985, pp. 408–416.
- [BOSY13] E. Blais, K. Onak, R. Servedio, and G. Yaroslavtsev. *Concise representations of discrete submodular functions*. Personal communication. 2013.
- [Cha05] S. Chakraborty. *Sensitivity, Block Sensitivity and Certificate Complexity of Boolean Functions*. 2005.
- [CKKL12] M. Cheraghchi, A. Klivans, P. Kothari, and H. Lee. “Submodular functions are noise stable”. In: *SODA*. 2012, pp. 1586–1592.
- [DGJSV10] I. Diakonikolas, P. Gopalan, R. Jaiswal, R. A. Servedio, and E. Viola. “Bounded Independence Fools Halfspaces”. In: *SIAM J. Comput.* 39.8 (2010), pp. 3441–3462.
- [DSFTWW15] D. Dachman-Soled, V. Feldman, L.-Y. Tan, A. Wan, and K. Wimmer. “Approximate resilience, monotonicity, and the complexity of agnostic learning”. In: *SODA*. 2015.
- [Fei06] U. Feige. “On maximizing welfare when utility functions are subadditive”. In: *STOC*. 2006, pp. 41–50.
- [FK14] V. Feldman and P. Kothari. “Learning Coverage Functions and Private Release of Marginals”. In: *COLT*. 2014, pp. 679–702.
- [FKV13] V. Feldman, P. Kothari, and J. Vondrák. “Representation, Approximation and Learning of Submodular Functions Using Low-rank Decision Trees”. In: *COLT*. 2013, 30:711–740.

- [FMV07] U. Feige, V. Mirrokni, and J. Vondrák. “Maximizing Non-Monotone Submodular Functions”. In: 2007, pp. 461–471.
- [Fri98] E. Friedgut. “Boolean Functions With Low Average Sensitivity Depend On Few Coordinates”. In: *Combinatorica* 18.1 (1998), pp. 27–35.
- [FV15] V. Feldman and J. Vondrák. “Tight Bounds on Low-Degree Spectral Concentration of Submodular and XOS Functions”. In: *FOCS*. 2015, pp. 923–942.
- [FV16] V. Feldman and J. Vondrák. “Optimal Bounds on Approximation of Submodular and XOS Functions by Juntas”. In: *SIAM J. Comput.* 45.3 (2016), pp. 1129–1170.
- [GHIM09] M. X. Goemans, N. J. A. Harvey, S. Iwata, and V. S. Mirrokni. “Approximating submodular functions everywhere”. In: *SODA*. 2009, pp. 535–544.
- [GHRU11] A. Gupta, M. Hardt, A. Roth, and J. Ullman. “Privately releasing conjunctions and the statistical query barrier”. In: *STOC*. 2011, pp. 803–812.
- [GKK08] P. Gopalan, A. Kalai, and A. Klivans. “Agnostically Learning Decision Trees”. In: *STOC*. 2008, pp. 527–536.
- [Hau92] D. Haussler. “Decision theoretic generalizations of the PAC model for neural net and other learning applications”. In: *Information and Computation* 100.1 (1992), pp. 78–150. ISSN: 0890-5401.
- [KKL88] J. Kahn, G. Kalai, and N. Linial. “The influence of variables on Boolean functions”. In: *FOCS*. 1988, pp. 68–80.
- [KKMS08] A. Kalai, A. Klivans, Y. Mansour, and R. Servedio. “Agnostically Learning Half-spaces”. In: *SIAM Journal on Computing* 37.6 (2008), pp. 1777–1805.
- [KS94] M. Kearns and R. Schapire. “Efficient distribution-free learning of probabilistic concepts”. In: *Journal of Computer and System Sciences* 48 (1994), pp. 464–497.
- [KSS94] M. Kearns, R. Schapire, and L. Sellie. “Toward Efficient Agnostic Learning.” In: *Machine Learning* 17.2-3 (1994), pp. 115–141.
- [KST08] S. Kakade, K. Sridharan, and A. Tewari. “On the Complexity of Linear Prediction: Risk Bounds, Margin Bounds, and Regularization”. In: *NIPS*. 2008, pp. 793–800.
- [LLN06] B. Lehmann, D. J. Lehmann, and N. Nisan. “Combinatorial auctions with decreasing marginal utilities”. In: *Games and Economic Behavior* 55 (2006), pp. 1884–1899.
- [MOS04] E. Mossel, R. O’Donnell, and R. Servedio. “Learning functions of k relevant variables”. In: *Journal of Computer and System Sciences* 69.3 (2004), pp. 421–434.
- [MR06] C. McDiarmid and B. Reed. “Concentration for self-bounding functions and an inequality of Talagrand”. In: *Random structures and algorithms* 29 (2006), pp. 549–557.
- [Nis89] N. Nisan. “CREW PRAMS and Decision Trees”. In: *STOC*. 1989, pp. 327–335.
- [O’D13] R. O’Donnell. *Analysis of boolean functions*. <http://analysisofbooleanfunctions.org>, 2013.
- [RY13] S. Raskhodnikova and G. Yaroslavtsev. “Learning pseudo-Boolean k -DNF and Submodular Functions”. In: *SODA*. 2013.

- [Val12] G. Valiant. “Finding correlations in subquadratic time, with applications to learning parities and juntas”. In: *The 53rd Annual IEEE Symposium on the Foundations of Computer Science (FOCS)*. 2012.
- [Val84] L. G. Valiant. “A theory of the learnable”. In: *Communications of the ACM* 27.11 (1984), pp. 1134–1142.
- [Von10] J. Vondrák. *A note on concentration of submodular functions*. arXiv:1005.2791v1. 2010.