On function field Mordell-Lang and Manin-Mumford

Franck Benoist*

Elisabeth Bouscaren^{*}

Anand Pillav[†]

March 1, 2022

Abstract

We give a reduction of the function field Mordell-Lang conjecture to the function field Manin-Mumford conjecture, for abelian varieties, in all characteristics, via model theory, but avoiding recourse to the dichotomy theorems for (generalized) Zariski geometries. Additional ingredients include the "Theorem of the kernel", and a result of Wagner on commutative groups of finite Morley rank without proper infinite definable subgroups. In positive characteristic, where the main interest lies, there is one more crucial ingredient: "quantifier-elimination" for the corresponding $A^{\sharp} = p^{\infty}A(\mathcal{U})$ where \mathcal{U} is a saturated separably closed field.

1 Introduction

This paper concerns relationships between "known" results. The original motivation was to supply a transparent account of the function field Mordell-Lang conjecture in positive characteristic. The possibility of reducing Mordell-Lang to Manin-Mumford, in the function field case, was initiated in a talk by the third author in Paris, in December 2010. Damian Rössler picked

^{*}Partially supported by ANR MODIG (ANR-09-BLAN-0047) Model theory and Interactions with Geometry and ANR ValCoMo (ANR-13-BS01-0006) Valuations, Combinatorics and Model Theory

 $^{^\}dagger \mathrm{Partially}$ supported by grants from the EPSRC and NSF

up the theme, and eventually with Corpet, produced a successful algebraicgeometric account of such a reduction, in positive characteristic ([30], [10]). Our original strategy (quite different from what Rössler did) presented in the 2010 talk referred to above, involved some soft stable-group theory results, together with, in positive characteristic, a conjectural "quantifier-elimination" for certain type-definable groups. In the current paper we prove that this strategy works, supplying the so far missing ingredient in characteristic p, the "quantifier elimination" result for abelian varieties.

The subtext is Hrushovski's proof of function field Mordell-Lang [15], which depends on a dichotomy theorem for (generalized) Zariski geometries. In the characteristic 0 case, it is classical (strongly minimal) Zariski geometries which are relevant and the dichotomy theorem is proved in [17] and in [35], although all proofs are complicated, to say the least. But in the positive characteristic case, *type-definable* Zariski geometries are the relevant objects. The needed dichotomy in this case is a combination of a complicated axiomatic account of a field construction in [17], together with arguments in [15] showing that the axioms are satisfied for the particular minimal types in separably closed fields that we are interested in. In both cases, the dichotomy theorem is difficult and its proof impenetrable for non model theory experts (as well as for many model-theorists). The current authors have been pre-occupied for some years about seeing what is really going on, in particular avoiding the recourse to (generalized) Zariski geometries, and/or recovering the results by more direct arguments.

In [26] this issue was taken up, and an approach using differential jet spaces was developed. This succeeded in characteristic 0, but not entirely in positive characteristic due to inseparability issues, although the approach recovered some cases due to Abramovich and Voloch [1]. It is still open whether the approach can be tweaked so as to work in general in the characteristic p case.

In [27] Pink and Rössler gave a reasonably transparent algebraic-geometric proof of function field Manin-Mumford in positive characteristic with *all* torsion points in place of prime-to-p torsion points. This suggested to us to try to reduce function field Mordell-Lang to function field Manin-Mumford, but still using model-theoretic methods. In the current paper we succeed in doing this, producing, so to speak, a "second generation" model-theoretic proof of the Mordell-Lang conjecture for abelian varieties over function fields. Exploring connections between the methods of Rössler and Corpet and either the jet space ideas in [26], or the current paper, would be interesting. We should also mention the interesting paper [34] in which another new algebraic-geometric account of Mordell-Lang in positive characteristic is given, but where only finitely generated groups Γ are considered.

Of course we could also consider the absolute Mordell-Lang conjecture in characteristic 0 (proved by Faltings, McQuillan,..) and ask whether there is a "soft" reduction to absolute Manin-Mumford. We doubt that this is the case as these two theorems seem to us (maybe incorrectly) to be of different orders of difficulty. That such a reduction is possible in the function field case has additional interest.

Acknowledgements. We would like to thank both Université Paris-Sud, Orsay, where the third author was a Professeur Invité in March-April 2010 and the Mathematical Sciences Research Institute, Berkeley, where the authors participated in the Spring 2014 model theory program. The first author would like to thank Françoise Delon for very useful discussions about the quantifier elimination question. Thanks also to the referee of a first version of this paper for helpful comments, resulting in a reorganization of the paper.

2 Preliminaries

The statements of the function field Mordell-Lang which we prove in this paper, as well as the main arguments we use, are specific to abelian varieties. They are also restricted to the function field case in one variable ($\mathbb{C}(t)^{\text{alg}}$ in characteristic 0, and $\mathbb{F}_p(t)^{\text{sep}}$ in characteristic p) as the "Theorem of the Kernel" has been proved only in these cases, in [7] and in [29]. Basic background, as well as references, on abelian varieties, Mordell-Lang and Manin Mumford can be for example found in [14].

Statement of function field Mordell-Lang in characteristic 0. Let $K = \mathbb{C}(t)^{\text{alg}}$, the algebraic closure of $\mathbb{C}(t)$. Let A be an abelian variety over K with \mathbb{C} -trace 0. Let X be an irreducible subvariety of A (defined over K), and let Γ be a "finite-rank" subgroup of A(K), namely Γ is contained in the division points of a finitely generated subgroup of A(K). Suppose $X \cap \Gamma$ is Zariski-dense in X. Then X is a translate of an abelian subvariety of A.

Statement of function field Manin-Mumford in characteristic 0. As above, except that the hypothesis on Γ is strengthened to: Γ is contained in

the group of torsion points of A.

Statement of function field Mordell-Lang in characteristic p > 0. Let K be the separable closure of $\mathbb{F}_p^{\mathrm{alg}}(t)$. Let A be an abelian variety over K with $\mathbb{F}_p^{\mathrm{alg}}$ -trace 0. Let X be an irreducible subvariety of A, defined over K. And let Γ be a subgroup of A(K) contained in the prime-to-p-division points of a finitely generated subgroup. Suppose that $X \cap \Gamma$ is Zariski-dense in X. Then X is a translate of an abelian subvariety of A.

The formulation involving prime-to-p division points is due to Abramovich and Voloch [1]. One could also ask what happens when K is algebraically closed and $\Gamma < A(K)$ is the group of *all* division points of some finitely generated subgroup. No obstacle is currently known to this.

Statement of function field Manin-Mumford in characteristic p. As above, except that Γ is assumed to be contained in the group of *all torsion points* of A.

We will write MM for Manin-Mumford and ML for Mordell-Lang. And from now on, k will denote \mathbb{C} in characteristic zero and $\mathbb{F}_p^{\text{alg}}$ in characteristic $p, K = k(t)^{\text{alg}}$ in characteristic zero, and $K = k(t)^{sep}$ in characteristic p.

In characteristic 0, function field MM as stated is clearly a special case of function field ML. And it follows from the absolute case of MM, of which there are many proofs more generally for all commutative algebraic groups (for example see [13] or, for a model theoretic proof, [16]). In positive characteristic MM (with all torsion points) is proved by Pink and Rössler [27]. The proof uses a variety of methods, including Dieudonné modules, but is accessible. A proof was also given by Scanlon [31] using the dichotomy theorem in $ACFA_p$ (algebraically closed fields of characteristic p with a generic automorphism) which itself depends on an even more generalized notion of Zariski geometries than used in separably closed fields.

As explained in the introduction, *in all characteristics*, assuming function field Manin-Mumford, we give a proof of function field Mordell-Lang without appealing to the dichotomy theorems for Zariski geometries.

The beginning of our proof follows the first steps of Hrushovski's original proof in [15], and has the uniform strategy in both characteristic zero and

characteristic p of "embedding" the algebraic-geometric set-up in a differential algebraic environment. But then we replace the use of the dichotomy theorems by some other ingredients which appeal to "softer" model theory. Two of these ingredients are common to all characteristics, as we will explain below: the first one, of an algebraic nature, we refer to as the "Theorem of the Kernel"; the second one, pertaining to model theory, is the structure of g-minimal groups of finite Morley rank.

Then, in characteristic zero, like Hrushovski himself in his proof, we use the "weak socle theorem" for groups of finite Morley rank (see Section 4.1).

In characteristic p, we do not need this socle theorem for groups, but we need a new result, which forms the core of the paper: in section 3, we prove quantifier elimination for the induced structure on the subgroup of infinitely p-divisible K-rational points of the abelian variety A, and this of course is accomplished without appealing to the dichotomy theorems.

We assume familiarity with model theory, basic stability, as well as differentially and separably closed fields. The book [19] is a reasonable reference, as well as [23] for more on stability theory. Definability means with parameters unless we say otherwise.

2.1 Passing to the differential framework, the group A^{\sharp}

As in Hrushovski's approach, (as well as in Buium's in [8] for the characteristic 0 case), we pass to a differential algebraic framework: differentially closed fields of characteristic zero, or separably closed fields of degree of imperfection one, in characteristic p, which can be viewed as the existentially closed fields with Hasse-Schmidt derivations. In these enriched frameworks, one can "replace" the group Γ by some (infinitely) definable subgroup of the rational points of A.

We fix some more notation.

In characteristic 0, K has a unique derivation ∂ extending d/dt on k(t) and K^{diff} denotes a differential closure of (K, ∂) . We work in the language of differential fields, with the first order theory of K^{diff} , the theory of differentially closed fields of characteristic zero, DCF_0 . It will be convenient sometimes to work in a saturated elementary extension \mathcal{U} of K^{diff} . Recall that DCF_0 is ω -stable with quantifier-elimination and elimination of imaginaries.

In characteristic p, \mathcal{U} will be a saturated elementary extension of K in the language of fields. In contradistinction to the characteristic 0 case, passing to \mathcal{U} will be a crucial part of the proof, rather than just a convenience. The

first order theory of K (or \mathcal{U}) in the language of fields is known as or denoted by $SCF_{p,1}$, the theory of separably closed fields of characteristic p and degree of imperfection 1. It is stable, but not superstable, and has quantifierelimination and elimination of imaginaries after either adding symbols for a p-basis and the so-called λ -functions, or just symbols for a strict iterative Hasse-Schmidt derivation.

From now on, A is an abelian variety over K.

Definition 2.1. The group A^{\sharp} .

(i) In characteristic 0, A^{\sharp} denotes the "Kolchin closure of the torsion", namely the smallest definable (in the sense of differentially closed fields) subgroup of $A(\mathcal{U})$ which contains the torsion subgroup (so note that A^{\sharp} is definable over K.)

(ii) In positive characteristic, A^{\sharp} denotes $p^{\infty}(A(\mathcal{U})) =_{def} \bigcap_{n} p^{n}(A(\mathcal{U}))$ (an infinitely definable subgroup over K.

Remark 2.2. In characteristic 0, the smallest definable subgroup containing the torsion subgroup exists by ω -stability of the theory DCF_0 . As A^{\sharp} is definable, we can consider not only $A^{\sharp}(\mathcal{U})$, but also $A^{\sharp}(K^{\text{diff}})$. In positive characteristic, A^{\sharp} is only infinitely definable, it is the maximal divisible subgroup of $A(\mathcal{U})$ and it is also the smallest infinitely definable subgroup of $A(\mathcal{U})$ which contains the prime-to-p torsion of A. Moreover $A^{\sharp}(K) = \bigcap_n p^n(A(K))$ is infinite and is also the maximal divisible subgroup of A(K).

An exposition of further properties of A^{\sharp} in both characteristics, as well as precise references can be found in [2].

For example, the following basic facts hold in all characteristics and were mostly originally proved in [15]:

Fact 2.3. (i) A^{\sharp} is also the smallest Zariski dense (infinitely) definable subgroup of $A(\mathcal{U})$.

(ii) A^{\sharp} is connected (no relatively definable subgroup of finite index), and of finite U-rank in char. p, and finite Morley rank in char. 0.

(iii) If A is a simple abelian variety, A^{\sharp} has no proper infinite type-definable subgroup.

(iv) If A is the sum of simple A_i then A^{\sharp} is the sum of the A_i^{\sharp} .

We can now state:

The Theorem of the Kernel. In all characteristics: Let A be an abelian variety over K with k-trace 0, then $A^{\sharp}(K)$ is contained in the group of torsion points of A.

This is a differential algebraic or model-theoretic theorem of the kernel. In Corollary K3 of [7], the characteristic 0 case is given, where it is deduced from Chai's strengthening of a theorem of Manin. In [29], the positive characteristic case is proved. Note that in the positive characteristic, the hypothesis of k-trace zero is not needed.

We now pass to the purely model-theoretic ingredients.

2.2 Groups of finite Morley rank and induced structures

Definition 2.4. Let G be a group (with additional structure), which has finite Morley rank and is commutative and connected. We say that G is g-minimal if it has no proper nontrivial connected definable subgroup (equivalently, no proper infinite definable subgroup).

Let us remark that in this context, g-minimality of G passes to saturated elementary extensions: In groups of finite Morley rank, there is a bound on the cardinality of uniformly definable families of finite subgroups, as they do not have the "finite cover property" (see for example [28]).

The following appears in [32] (as a direct consequence of Corollary 6):

Theorem 2.5. Suppose that G is g-minimal. Then any infinite algebraically closed subset of G is (the universe of) an elementary substructure of G.

Remark: So a g-minimal group behaves like a strongly minimal set, where it is well known that infinite algebraically closed subsets are elementary substructures. By Zilber's indecomposability theorem a g-minimal group is almost strongly minimal, namely in the algebraic closure of a definable strongly minimal subset D together with a finite set F of parameters. Then it is also well-known that any algebraically closed (over F) subset whose intersection with D is infinite is an elementary substructure. But here we only suppose that X is algebraically closed (over \emptyset) and infinite, so one cannot directly deduce the result from the almost strongly minimal case. We now fix an ambient saturated stable structure \mathcal{U} . A subset X of \mathcal{U}^n is called *type-definable* (or infinitely definable) if it is the intersection of a small collection of definable sets, namely defined by a small partial type. In the case of interest X will be a countable intersection of definable sets.

Suppose X is type-definable over the small set of parameters A. By a *relatively definable* subset Y of X^m we mean the intersection of some definable (with parameters) subset Z of a suitable Cartesian power of \mathcal{U} with X^m . We will say that Y is relatively A-definable, or relatively definable over A, if Z can be chosen to be A-definable.

Let us fix a set X, type-definable over some small set of parameters A.

Lemma 2.6. Let Y be a relatively definable subset of X (or some Cartesian power of X) which is invariant under automorphisms of \mathcal{U} fixing A pointwise. Then Y is relatively A-definable.

Proof. Suppose Y is relatively definable by formula $\phi(x, b)$ (namely Y is the set of solutions in X of $\phi(x, b)$), where we are exhibiting the required parameters b. By our assumptions, we have that if $tp(b_1/A) = tp(b/A)$ then $\phi(x, b_1)$ relatively defines the same subset of X as does $\phi(x, b)$. We can apply compactness to find a formula $\psi(y) \in tp(b/A)$ such that Y is relatively defined by the formula $\exists y(\psi(y) \land \phi(x, y))$.

Let us denote by \mathcal{X}_A the structure with universe X and predicates for all relatively definable over A subsets of X^n , for all n. With this notation:

Definition 2.7. We will say that X_A has quantifier elimination or QE, if $Th(\mathcal{X}_A)$ has quantifier elimination in the language above.

It is clear from this definition that:

Remark 2.8. X_A has QE if and only if whenever Y is a relatively A-definable subset of X^{n+1} then the projection of Y to X^n is relatively A-definable.

Let us recall a few basic facts about induced structures:

Lemma 2.9. (i) Suppose $A \subseteq B$. Then X_A has QE iff X_B has QE (so we just say that X has QE).

(ii) X has QE just if the projection of any relatively definable subset of any X^{n+1} to X^n is relatively definable (noting that in general it is only type-definable).

(iii) X having QE is equivalent to \mathcal{X}_A being a saturated structure.

Proof. (i) Right implies left follows from Lemma 2.6 as a projection of an A-invariant set is also A-invariant. For left to right: Suppose $Y \subset X^{n+1}$ is relatively definable over B, by $\phi(x_1, ..., x_{n+1}, b)$ where we witness the parameters $b \in B$, which may live outside X. Let Y_1 be the projection of Y to X^n . By Lemma 2.6 it suffices to prove that Y_1 is relatively definable (as clearly it is invariant under automorphisms of \mathcal{U} fixing b pointwise). Now by definability of types we may find some L-formula $\psi(x_1, ..., x_{n+1}, z)$ and c from X such that Y is relatively definable by $\psi(x_1, ..., x_{n+1}, c)$. Let z be an m-tuple of variables. Let Y' be the subset of X^{n+1+m} relatively defined by ψ . By our assumption that X_A has QE, the projection Y'' of Y obtained by existentially quantifying out x_{n+1} , is a relatively A-definable subset of X^{n+m} , (relatively) defined by a formula $\chi(x_1, ..., x_n, z)$ say. So $\chi(x_1, ..., x_n, c)$ relatively defines Y_1 , as required.

(ii) By (i) and Lemma 2.6

(iii) The point is that to say that X_A has QE means precisely that if b_1, b_2 are *n*-tuples from X, then $tp(b_1/A) = tp(b_2/A)$ in the sense of \mathcal{U} iff $tp(b_1) = tp(b_2)$ in the sense of \mathcal{X}_A .

Note that in general \mathcal{X}_A is only quantifier-free saturated (and homogeneous) and is sometimes referred to as a Robinson structure.

Of course when X is definable (rather than type-definable) over A, then X_A always has QE and is referred to as "X with its induced structure".

Using Theorem 2.5 applied to the G_i 's with their induced structure, we deduce easily:

Corollary 2.10. Suppose G is a (saturated) commutative, connected, group of finite Morley rank (with additional structure) which is a sum of finitely many g-minimal \emptyset -definable subgroups G_i . Then any algebraically closed subset of G which meets each G_i in an infinite set, is an elementary substructure of G.

In positive characteristic, in order to apply 2.5 and Corollary 2.10, to A^{\sharp} , we will need to show that A^{\sharp} has quantifier elimination, in the sense above. For example, we know (2.3) that if A is a simple abelian variety, then A^{\sharp} has no proper relatively definable subgroup, but we will need to know that

the group A^{\sharp} , equipped with predicates for relatively definable sets, is a *g*-minimal group, as a first order structure in its own right; QE says that every definable subgroup of this first order structure A^{\sharp} is relatively definable (in \mathcal{U}), which is exactly what we need.

Now the (generalized) Zariski geometry arguments from [15] give the dichotomy theorem for minimal "thin" types in separably closed fields, implying that if A is simple with k-trace 0, then A^{\sharp} is minimal (i.e. U-rank 1), connected, and 1-based, from which it easily follows that A^{\sharp} has QE. For arbitrary traceless abelian varieties A, A^{\sharp} will be a sum of such minimals, hence also 1-based and so we also have QE. So the QE hypothesis is true, after the fact so to speak, once one knows A^{\sharp} is one-based.

Our method of proving QE is related to proofs that minimal thin types in SCF are Zariski, but does not use the dichotomy theorem for (type-definable) Zariski geometries (nor in fact even the so called dimension theorem). In fact we prove QE for A^{\sharp} directly, for any abelian variety over a separably closed field of imperfection degree 1, without any assumption on the trace. The fact that A is an abelian variety, and not a semiabelian variety plays an essential role, though. Indeed, we can define G^{\sharp} in the same way for semiabelian varieties G. But building on work in [2], Alexandra Omar Aziz, in [21] gave semiabelian examples G for which G^{\sharp} does not have QE.

3 Quantifier elimination for A^{\sharp} in characteristic p

Here we prove the quantifier elimination result in full generality. As this is the case we need, and it simplifies notation, we keep the assumption that we are working with a separably closed field of degree of imperfection one, but the same proof will work for any finite non zero degree of imperfection.

Theorem 3.1. Let A be an abelian variety over any separably closed field K of degree of imperfection 1, and \mathcal{U} be a saturated extension of K. We consider $A^{\sharp} = p^{\infty}A(\mathcal{U})$, and we denote by \mathcal{A} the structure A^{\sharp} with relatively definable sets (with parameters from K). Then \mathcal{A} has quantifier elimination.

Note that by Lemma 2.9, it will follow that $\mathcal{A}_{\mathcal{U}}$ has quantifier elimination as well. But, conversely, Lemma 2.9 says that, in order to prove the theorem for \mathcal{A} , we can suppose, for this proof only, that K itself is sufficiently saturated.

This will be used in Claim 3.5.

We will work in the language of rings augmented by λ -functions and a constant t for the p-basis (see [12] as reference). It means that for each a in the field, $a = \sum_{i \in p^n} \lambda_{n,i}(a)^{p^n} M_i$, where elements of p^n are seen as sequences of length n with values between 0 and p-1, and for $i \in p^n$ and $j \in p$, $M_j = t^j$, $M_{i^{\lambda}j} = M_i M_j^{p^n}$ and $\lambda_{n+1,i^{\lambda}j} = \lambda_{1,j} \circ \lambda_{n,i}$. The tuple $a_{=n} := (\lambda_{n,i}(a))$ is called the tuple of λ -components of level n, and we denote $a_{\leq n} = (a_{=m})_{0 \leq m \leq n}$, $a_{\infty} = (a_{=n})_{n \geq 0}$.

Similarly, for X a multivariable, $X_{\infty} = (X_i)_{i \in p^n, n \ge 0}$. Let $I^0(X)$ be the ideal of $K[X_{\infty}]$ generated by the polynomials $X_i - \sum_{j \in p} X_{i \land j}^p M_j$, for all $i \in p^n$, $n \ge 0$. A complete type over K corresponds bijectively to a prime separable ideal of $K[X_{\infty}]$ containing $I^0(X)$. An ideal I of $K[X_{\infty}]$ is separable if and only if whenever $\sum_{j \in p} P_j^p M_j$ is in I, each P_j is in I.

We will use the Λ_n functors, from the category of algebraic varieties (over K) into itself, as defined in [5]. For each algebraic variety V, we have a natural definable map $\lambda_n : V(\mathcal{U}) \to \Lambda_n V(\mathcal{U})$ and a natural morphism $\rho_n : \Lambda_n V \to V$ which define reciprocal bijections between $V(\mathcal{U})$ and $\Lambda_n V(\mathcal{U})$ (the notation λ_n is coherent with the previous one for the affine line). Note that Λ_n , λ_n and ρ_n are obtained from Λ_1 , λ_1 and ρ_1 by the suitable composition. We equip each $V(\mathcal{U})$ with the λ -topology, whose basic closed sets are $\lambda_m^{-1}(V_m)$ for any $m \geq 0$ and V_m any algebraic subvariety of $\Lambda_m V$. Note that for any $n \geq 0$, $\Lambda_n V_m$ is an algebraic subvariety of $\Lambda_{m+n}V$, and $\lambda_m^{-1}(V_m) = \lambda_{m+n}^{-1}(\Lambda_n V_m)$. It follows that finite unions of basic closed sets are still basic closed; we obtain arbitrary closed sets by possibly infinite intersection. It follows from quantifier elimination in the theory of separably closed fields in this language that definable subsets of $V(\mathcal{U})$ are boolean combinations of definable (i.e. basic) closed sets.

If A is an abelian variety, $\Lambda_m A$ is an algebraic group, but is not an abelian variety. But from [3], we know that for each $m \geq 0$, there is an algebraic subgroup $A_m \subset \Lambda_m A$, which is isogenous to A, such that $p^m A(\mathcal{U}) = \lambda_m^{-1}(A_m)$ (it was shown there using the formalism of Weil's restriction of the scalars $\Pi_{K/K^{p^m}}A$, which is known to coincide with $F^{(m)}\Lambda_m A$, $F^{(m)}$ being the *m*-th power of the absolute Frobenius).

Recall that A is a sum of simple abelian varieties, $A = \sum_{1 \le i \le n} A_i$. We will use the following consequence of the Zilber indecomposability theorem (see Fact 3.8 in [2]): as A_i is simple, A_i^{\sharp} has no proper infinite type-definable

subgroups (Fact 2.3) and hence, there exists a minimal type in A_i^{\sharp} whose set of realizations, which will be denoted by Q_i in the following, is such that $A_i^{\sharp} = Q_i + \ldots + Q_i$ (m_i times for some m_i , that is, any element of A_i^{\sharp} is the sum of m_i elements from Q_i).

First we recall some basic facts about the group A^{\sharp} in characteristic p, which will be used here and in Section 4.2

We refer the reader to [2] where we give a precise account of relative Morley rank for type-definable sets in a stable structure (called "internal Morley dimension" in [15]).

Fact 3.2. Both A^{\sharp} and the A_i^{\sharp} are connected groups, with finite relative Morley rank. Moreover, A_i^{\sharp} is the connected component of $A^{\sharp} \cap A_i$ and is relatively definable in A^{\sharp} .

Comments. The fact that A^{\sharp} has finite relative Morley rank is claimed in Remark 2.19 of [15]. However it is also implicitly claimed there that $G^{\sharp} = p^{\infty}(G(\mathcal{U}))$ also has finite relative Morley rank, whenever G is semiabelian, and this is actually wrong, as pointed out in [2]. So we refer the reader rather to the proof of Fact 3.8 from [2]. It is worth remarking that it is this semiabelian counterexample without relative Morley rank which is shown, in [21], not to have QE. But in general there is no reason why a type-definable group of finite relative Morley rank should have QE. Once we know that A^{\sharp} has relative finite Morley rank, as, for each i, A_i^{\sharp} is the connected component of $A_i \cap A^{\sharp}$, it follows that A_i^{\sharp} is relatively definable in A^{\sharp} , but again this is no longer true in the semiabelian counterexample.

Proposition 3.3. For every $n \ge 1$, the λ -topology on $(A^{\sharp})^n$ is Noetherian of finite dimension.

Note that, in the examples of semiabelian varieties which do not have finite relative finite Morley rank given in [2], the topology is not Noetherian.

It was shown in [15] that the trace of the λ -topology on each Q^k is Noetherian and has finite dimension, if Q is a *thin* minimal type (see below), which is the case here. Now the Noetherianity for A_i^{\sharp} follows as we have a continuous relatively definable surjective map from Q_i^{md} onto $A_i^{\sharp d}$.

Passing to A^{\sharp} itself will be a little more complicated.

Let us first note that, once we know that the topology is Noetherian, it is easy to see that it is finite dimensional, that is, that every closed set has finite topological dimension. Indeed, by Noetherianity, every closed set is a finite union of irreducible closed sets, and in $(A^{\sharp})^n$, there is a finite bound on the length of strictly decreasing sequences of *irreducible* closed sets. This follows from "thinness" of the types involved. Recall that if $a \in (A^{\sharp})^n$ then the type of a over K is *thin*, that is, the field generated over K by a and its images under the λ -functions has finite transcendence degree over K ([15] or [6]). More precisely, the transcendence rank of the prime separable ideal I(a/K), of all λ -polynomials vanishing at a, has finite transcendence rank smaller than $(dimA)^n$. Now if $F \subsetneq G \subseteq (A^{\sharp})^n$ are two irreducible closed sets and $I(G) \subsetneq I(F)$ are the associated two prime separable ideals then, the transcendence rank of both are finite and bounded by $(dimA)^n$ and the transcendence rank of I(F) must be strictly smaller than the transcendence rank of I(G).

So, in order to prove Proposition 3.3, we just need to show Noetherianity.

Lemma 3.4. Let p_1, \ldots, p_k be thin minimal types over K and for each i, let P_i denote the set of realizations of p_i in \mathcal{U} . Then for every n_1, \ldots, n_k the topology on $P_1^{n_1} \times \ldots \times P_k^{n_k}$ is Noetherian of finite dimension. In particular every closed subset is relatively definable.

We proceed in two steps.

Claim 3.5. Suppose q_1, \ldots, q_n are pairwise non orthogonal minimal thin types (not necessarily distinct) and Q_j denotes the set of realizations of q_j in \mathcal{U} . Then there is a minimal thin type over K, r, such that, if R denotes the set of realizations of r in \mathcal{U} , then there is a continuous map f from R^n onto $Q_1 \times \ldots \times Q_n$. It follows that the topology on $Q_1 \times \ldots \times Q_n$ is Noetherian.

Proof. Once we have shown that such a type r and the required continuous maps exist, the Noetherianity follows directly from that of Cartesian products of R, which was shown in [15].

Let (a_1, a_2, \ldots, a_n) , realizing $q_1 \times q_2 \times \ldots \times q_n$, be such that $r := tp(a_1, a_2, \ldots, a_n)$ has U-rank equal to one. This exists by the assumption of non pairwise orthogonality and the saturation assumption on K. So the type r is minimal and is easily seen to be also thin (the transcendence degree of the field generated by the a_i and their images by the λ -functions will be finite as each q_i has this property).

For each *i*, the *i*-th component map, π_i from *R* to Q_i is surjective, as all elements of Q_i realize the same type over *K*.

Now consider the Cartesian product \mathbb{R}^n . We claim that there is a surjective definable map f from \mathbb{R}^n onto $Q_1 \times \ldots \times Q_n$ and that it is continuous. Let $(b_1, \ldots, b_n) \in \mathbb{R}^n$, and let $f(b_1, \ldots, b_n) := (\pi_1(b_1), \ldots, \pi_n(b_n)) \in Q_1 \times \ldots \times Q_n$. Let (a_1, \ldots, a_n) be any tuple from $Q_1 \times \ldots \times Q_n$. For each i there is some element b_i from \mathbb{R} such that $\pi_i(b_i) = a_i$, so f is surjective.

Now f is the restriction to \mathbb{R}^n of a projection map $(Q_1 \times \ldots \times Q_n)^n \to Q_1 \times \ldots \times Q_n$, so it is continuous.

Note that as we have not supposed that the non-orthogonal types in Claim 3.5 were distinct, this gives the Noetherianity for any Cartesian product of finitely many non pairwise orthogonal minimal thin types.

Claim 3.6. Suppose q_1, \ldots, q_m are pairwise orthogonal minimal thin types, and for each $i, 1 \leq i \leq m, \{p_{(i,1)}, \ldots, p_{(i,n_i)}\}$ is a set of minimal (thin) types non orthogonal to q_i . Let P_k denote the set of realizations of p_k in \mathcal{U} , and let Q_i denote the cartesian product $P_{(i,1)} \times \ldots \times P_{(i,n_i)}$. Then the topology on $Q_1 \times \ldots \times Q_m$ is Noetherian, and every closed set C is a finite union of closed sets of the form $W_1 \times \ldots \times W_m$, where W_i is a closed subset of Q_i .

Proof. First we note that by pairwise orthogonality and minimality of the types q_j and p_k , if \bar{a}_j is a tuple of elements from Q_j , then $\{\bar{a}_1, \ldots, \bar{a}_m\}$, is an independent set of tuples over K: indeed by minimality, each \bar{a}_j is contained in the algebraic closure (over K) of a K-independent subtuple B_j , each element in B_j realizing one of the types $\{p_{(j,1)}, \ldots, p_{(j,n_j)}\}$, which are all non-orthogonal to q_j . By orthogonality of the q_j , the set $B_1 \cup \ldots \cup B_m$ form an independent set over K and the rest follows.

A first observation: Let $Z = Z_b \subseteq Q_2 \times \ldots \times Q_m$ be a closed set (hence infinitely definable) defined over $K \cup b$ where b is a tuple of elements in Q_1 . Then Z is in fact defined over K. Indeed Z is K-invariant: let b' realize the same type as b over K. By the remark above b and b' must be independent from $Q_2 \cup \ldots \cup Q_m$ over K, it follows that b and b' also realize the same type over $K \cup Q_2 \cup \ldots \cup Q_m$, hence $Z_b = Z_{b'}$.

Now we show by induction on $m \ge 1$ that the topology on $Q_1 \times \ldots \times Q_m$ is Noetherian, and that every closed set is of the required form.

The case m = 1, that is, Cartesian products of a finite set of pairwise nonorthogonal thin minimal types, is Claim 3.5, which says that the topology is Noetherian.

Now, for m > 1, let $C \subseteq Q_1 \times \ldots \times Q_m$ be any closed subset, and let $a := (a_1, a_2, \ldots, a_m) \in C$. We show that

(*) a is contained in a set of the form $W \times Z \subseteq C$ where W is a closed subset of Q_1 defined over K and Z is a closed subset of $Q_2 \times \ldots \times Q_m$, also defined over K.

The Noetherianity and form of the closed sets will follow: by the induction assumption, W is a relatively definable closed subset in Q_1 and Z is a relatively definable closed subset in $Q_2 \times \ldots \times Q_m$ of the right form. By (*), C which is infinitely definable (over K) is covered by a union of such relatively definable sets each of the form $W \times Z$ each defined over K. By (model theoretic) compactness, it follows that C is a finite union of such sets. And as Q_1 and $Q_2 \times \ldots \times Q_m$ are both Noetherian, the result follows.

It remains only to check condition (*).

Let $Z := Z_{a_1} = \{(a_2, \ldots, a_m) \in Q_2 \times \ldots \times Q_m : (a_1, a_2, \ldots, a_m) \in C\}$. The set Z is closed and defined over $K \cup a_1$. By the first observation made above, Z is in fact defined over K. Now let $W = \{x \in Q_1 : \text{ for all } y \in Z, (x, y) \in C\}$. Then W is closed (it is the intersection of the C_y , for $y \in Z$) and clearly defined over K.

And of course $a \in W \times Z$ and $W \times Z \subseteq C$.

We can now conclude the proof of Lemma 3.4: For each non orthogonality class represented amongst the p_i , choose a representative q and apply Claim 3.6.

We now go back to the proof of Proposition 3.3: Let A be any abelian variety over K, $A = \sum_{1 \le i \le n} A_i$, where each A_i is a simple abelian variety over K. It follows that $A^{\sharp} = \sum_{0 \le i \le n} A_i^{\sharp}$. For each *i* there is a minimal thin type q_i such that $A_i^{\sharp} = Q_i + \ldots + Q_i$ (m_i times). So $A^{\sharp} = Q_1 + \ldots + Q_1 + Q_2 + \ldots + Q_2 + \ldots + Q_n + \ldots + Q_n$.

So we have a continuous relatively definable surjective map, for every d from the Cartesian product $(Q_1^{m_1} \times \ldots \times Q_n^{m_n})^d$ onto A^{\sharp^d} . And Noetherianity now follows from Lemma 3.4, and this concludes the proof of Proposition 3.3.

Note, as a corollary, that each closed set of $(A^{\sharp})^n$ is actually relatively definable.

If $\mathcal{U}_0 \prec \mathcal{U}$, and $a \in \mathcal{U}$, $\mathcal{U}_0[a_\infty]$ denotes the ring generated by a and its images by the λ -functions over \mathcal{U}_l and $\mathcal{U}_0(a_\infty)$, the fraction field of $\mathcal{U}_0[a_\infty]$.

Lemma 3.7. Let X and Y be closed sets in $(A^{\sharp})^d$, with X irreducible, and $pr: X \times \underline{Y} \to X$ the projection. Let $G \subsetneq F$ be closed subsets of $X \times Y$, such that $\overline{pr(F)} = X$ and F is irreducible. Let a be a topological generic of X over some small model \mathcal{U}_0 of definition for F and G, and we denote $F(a) = \{y \in Y \mid (a, y) \in F\}$, and similarly for G. Then $G(a) \subsetneq F(a)$. Moreover F(a) is irreducible as a closed set over $\mathcal{U}_0(a_{\infty})$ (note however that it may be reducible as a closed set over \mathcal{U}).

Proof. We denote by $\mathcal{U}_0[X]_{\infty} := \mathcal{U}_0[T_{\infty}]/I(X)$ the ring of λ -coordinates of X over \mathcal{U}_0 , and $\mathcal{U}_0[Y]_{\infty}$ in a similar way. By irreducibility, $\mathcal{U}_0[X]_{\infty}$ is an integral domain, and by choice of a, $\mathcal{U}_0[X]_{\infty} \simeq \mathcal{U}_0[a_{\infty}]$.

We denote by I(F) and I(G) the separable ideals in $\mathcal{U}_0[a_\infty][Y]_\infty \simeq \mathcal{U}_0[X \times Y]_\infty$ corresponding to F and G. Since $\overline{pr(F)} = X$, $I(F) \cap \mathcal{U}_0[a_\infty] = 0$. Now I(F(a)) is the ideal generated by I(F) in $\mathcal{U}_0(a_\infty)[Y]_\infty$, and similarly for I(G(a)). We claim that $I(F) \subsetneq I(G)$ implies that $I(F(a)) \subsetneq I(G(a))$. If I(F(a)) = I(G(a)), then for every $P \in I(G)$, there is some non zero $d \in \mathcal{U}_0[a_\infty]$ such that $dP \in I(F)$, which implies that $P \in I(F)$ since $d \notin I(F)$, which is prime. That contradicts $I(F) \neq I(G)$. We get that I(F(a)) is prime by the same kind of argument. This means that F(a) is irreducible as a closed set over $\mathcal{U}_0(a_\infty)$.

We now prove that A^{\sharp} is complete in the category of λ -closed varieties, in the following sense:

Proposition 3.8. Let F be a definable λ -closed subset of $(X \times A)(\mathcal{U})$, where X is an algebraic variety over K, and $pr : X \times A \to X$ the projection. Then $pr((X(\mathcal{U}) \times A^{\sharp}) \cap F)$ is λ -closed.

We first need the following lemmas.

Lemma 3.9. Let V, W be two affine varieties defined over K. Let g be a morphism from V to W defined over K.

Let $b \in W(\mathcal{U})$. Then $b \in g(V(\mathcal{U}))$ if and only if, for every $n, \lambda_n(b) \in (\Lambda_n g)(\Lambda_n V)$.

Proof. If $V \subset \mathbb{A}^r$ and $W \subset \mathbb{A}^k$, then $g = (g_1, \ldots, g_k)$, where each g_j is a polynomial map from V to \mathbb{A}^1 , and $\Lambda_m g_j$ is a p^m -tuple of elements in $K[X_{=m}]$ for X a multivariable of length r.

One direction is clear: if $b = g(a) \in g(V(\mathcal{U}))$, this follows from the fact that $\lambda_n \circ g(a) = \Lambda_n g \circ \lambda_n(a)$.

Now for the other direction, let $b \in W(\mathcal{U})$ be such that for every n, $\lambda_n(b) \in (\Lambda_n g)(\Lambda_n V)$.

Let H be the separable closure of $K(b_{\infty})$, $K \leq H \leq \mathcal{U}$, and consider the following set R of polynomials:

$$R := \{ (\Lambda_m g(X_{=m}))_{\sigma} - b_{\sigma}; \sigma \in p^m, m \ge 0 \}.$$

Let J_0 denote the ideal of the variety V over H, and I in $H[X_\infty]$, be the ideal generated by I^0 , J_0 and R.

Claim 1 $I \neq H[X_{\infty}]$

Proof of Claim 1: It suffices to find an infinite sequence e_{∞} in $\overline{\mathcal{U}}$ such that $I(e_{\infty}) \supset I$. By saturation of \mathcal{U}^{alg} as an algebraically closed field, it suffices to find, for each n, a sequence $e_{\leq n}$ such that $I(e_{\leq n}) \supset I_{=n} := I \cap H[X_{=n}]$. From the hypothesis on b, there is $a_n \in (\Lambda_n V)(\overline{\mathcal{U}})$ such that $(\Lambda_n g)(a_n) = \lambda_n(b)$. Now for $0 \leq m \leq n$, set $e_{=m} = \rho_{n-m}(a_{=n})$, and it gives the required tuple $e_{\leq n}$.

Claim 2 The ideal I is separable.

Proof of Claim 2: Using the fact that $I^0 \subset I$, it can be easily shown that it suffices to show that for each $h \in I_{=n} := I \cap H[X_{=n}]$, and each j < p, $(\Lambda_1 h)_j \in I$ (where $(\Lambda_1 h)_j$ is the j^{th} component of $\Lambda_1 h$ viewed as a tuple of elements in $H[X_{=n+1}]$). Since such h can be written as $h = h_0 + h_1 + \sum_k P_k Q_k$ with $h_0 \in I^0$, $h_1 \in J_0$ (which are separable), and $P_k \in H[X_{=n}]$, $Q_k \in$ $R \cap H[X_{=n}]$, and since we have the relations $(\Lambda_1(f_1+f_2))_j = (\Lambda_1 f_1)_j + (\Lambda_1 f_2)_j$ and $(\Lambda_1 f_1 f_2)_j = \sum_{k+l-m=j} (\Lambda_1 f_1)_k (\Lambda_1 f_2)_l M_m$, it suffices to consider the case of $Q \in R \cap H[X_{=n}]$. But this is clear by the definition of R: indeed, if $Q = (\Lambda_n g(X_{=n}))_\sigma - b_\sigma$ for some $\sigma \in p^n$, $(\Lambda_1 Q)_j = (\Lambda_{n+1} g(X_{=n+1}))_{\sigma^{\wedge} j} - b_{\sigma^{\wedge} j}$.

Now since I is a proper separable ideal containing I^0 , there is $a \in \mathcal{U}$ such that $I(a_{\infty}) \supset I$, which implies that $a \in V(\mathcal{U})$ and g(a) = b.

The following result, a direct corollary of the previous Lemma, is certainly well-known, probably in the formalism of Weil restrictions.

Lemma 3.10. Let V, W be two irreducible varieties defined over K. Let g be a morphism from V to W defined over K. Then there is some N such that

$$x \in g(V(\mathcal{U}))$$
 iff $\lambda_N(x) \in ((\Lambda_N g)(\Lambda_N V))(\mathcal{U})).$

Proof. Going from affine varieties to arbitrary algebraic varieties comes from the fact that the functors Λ_n preserve open affine coverings. Now we know that $g(\mathcal{U}(\mathcal{U}))$ is a definable set in the separably closed field \mathcal{U} . By the previous Lemma it is also infinitely definable as the infinite conjunction of the formulas $\lambda_k(x) \in (\Lambda_k g)(\Lambda_k V)$. The result then follows by compactness.

Proof of Proposition 3.8. For some m and some subvariety F_m of $\Lambda_m(X \times A)$, $F = \lambda_m^{-1}(F_m)$. We also have for any $n \ge m$ that $F = \lambda_n^{-1}(F_n)$, where $F_n = \Lambda_{n-m} F_m \subset \Lambda_n(X \times A)$. Let x be in $X(\mathcal{U})$. By saturation, $x \in \mathcal{I}$ $pr((X \times A^{\sharp}) \cap F)$ iff for every $n \geq m$, there is some $z_n \in p^n A(\mathcal{U})$ such that $(x, z_n) \in F$, which in turn is equivalent to $(*)_n$: there is some $y_n \in A_n(\mathcal{U})$ such that $(\lambda_n(x), y_n) \in F_n$. We now use Corollary 3.10: there is some $l = l_n$ (which depends on n but not on x) such that $(*)_n$ is equivalent to $\lambda_l \circ \lambda_n(x) \in (\Lambda_l pr)(\Lambda_l F_n)$, where $\Lambda_l pr : \Lambda_{l+n}(X \times A) \simeq \Lambda_{l+n}X \times \Lambda_{l+n}A \to$ $\Lambda_{l+n}X$ is still the projection on the first factor. Now look at the condition $(**)_n: \lambda_l \circ \lambda_n(x) \in pr(\Lambda_l F_n \cap (\Lambda_{l+n}X \times A_{l+n})).$ Recall that $A_{l+n} \subset \Lambda_{l+n}A$ characterizes p^{l+n} -divisible points. Since A_{l+n} is complete, $(**)_n$ is a λ closed condition. If x satisfies $(**)_n$, it satisfies $(*)_n$ a fortiori. Conversely, if $x \in pr((X \times A^{\sharp}) \cap F)$, the z_n 's appearing in the previous discussion can be chosen to be p^{∞} -divisible, which in particular implies that we can find y_n such that $\lambda_l(y_n) = \lambda_l \circ \lambda_n(z_n) \in A_{l+n}$, hence that $(**)_n$ is satisfied. We conclude that $pr((X \times A^{\sharp}) \cap F)$ is given by the conjunction of the con-

We conclude that $pr((X \times A^*) \cap F)$ is given by the conjunction of the conditions $(**)_n$, hence is λ -closed.

From Proposition 3.8 and Proposition 3.3, we get:

Corollary 3.11. Let F be a definable closed subset of A^d and $pr : A^d \to A^{d-1}$ the projection on the first coordinates. Then $pr(F(\mathcal{U}) \cap (A^{\sharp})^d)$ is closed and relatively definable.

In order to go from the case of closed sets to the case of constructible sets, we really follow the lines of the proof of quantifier elimination for onedimensional Zariski geometries given in [35] or [17] (note that QE for one dimensional Zariski geometries is a basic consequence of the axioms and does not involve the deep dichotomy result). We know that $A^{\sharp} = Q_1 + \ldots + Q_1 +$ $Q_2 + \ldots + Q_2 + \ldots + Q_n + \ldots + Q_n$, where Q_i is the set of realizations of a thin minimal type. It would be convenient to work with the relatively definable closed sets \overline{Q}_i , the closure of Q_i in the sense of the λ -topology, but it is not clear a priori why it should be of topological dimension 1. A more general fact would be that the U-rank of a type t coincides with the topological dimension of \overline{t} (the closure of the set of its realizations, or equivalently, the closed set given by the type ideal corresponding to t). It is true for types in A^{\sharp} , actually, but we know it only a posteriori, via the dichotomy theorem, which we do not want to use. We know however that U-rank $(t) \leq dim(\overline{t})$: because of the correspondence between closed type-definable sets and prime separable ideals in the suitable polynomials algebra, $dim(\overline{t})$ is given by the separable depth of the corresponding ideal I_t , and the separable depth is a stability rank, hence greater than or equal to the U-rank (see [11]).

It follows that if $F \subseteq A^{\sharp}$ is irreducible closed of topological dimension one, then F has a unique type of U-rank one, which is the type of maximal rank and is also its topological generic.

So, rather than Q_i , we will consider suitable relatively definable irreducible closed sets H_i of dimension 1. We proceed as follows. For each *i*, let H_i be a relatively definable irreducible closed subset of A_i^{\sharp} of dimension 1 (it exists since we know that the topology is Noetherian, and since translations are homeomorphisms, we are allowed to replace H_i by any of its translates in the following). By the comparison between the *U*-rank and the topological dimension above, it is clear that the generic type (in the topological sense) of H_i is a minimal type, hence, as A_i^{\sharp} has no proper infinite type-definable subgroups, we can apply Zilber's indecomposability theorem to get $A_i^{\sharp} = H_i + \ldots + H_i$ (m_i times for some m_i).

Hence $A^{\sharp} = \sum_{m_1} H_1 + \ldots + \sum_{m_n} H_n$, where the H_i are relatively definable closed subsets of A^{\sharp} of (topological) dimension 1.

Now for any formula $\phi(x, \overline{a})$ with parameters in A^{\sharp} , $\mathcal{A} \models \exists x \phi(x, \overline{a})$ if and only if

$$\mathcal{A} \models \exists x_{1,1} \in H_1 \dots \exists x_{1,m_1} \in H_1 \dots \exists x_{n,1} \in H_n \dots \exists x_{n,m_n} \in H_n \phi(\sum_{i,j} x_{i,j}, \overline{a}).$$

Hence it is sufficient to consider projections of the form $pr : H \times (A^{\sharp})^d \to (A^{\sharp})^d$, where H is one of the H_i 's. From Corollary 3.11 and the fact that H is closed, we get that pr takes closed sets to closed sets. From quantifier elimination in the separably closed field \mathcal{U} and Noetherianity in \mathcal{A} , we just

have to consider projections of definable sets $F \setminus G$, where $G \subsetneq F \subseteq H \times (A^{\sharp})^d$ are closed relatively definable sets, with F irreducible.

Proposition 3.12. The projection $pr(F \setminus G)$ is constructible in \mathcal{A} .

Proof. We proceed by induction on dim(F). The case dim(F) = 0 is obvious since it implies that F is a singleton.

Now $\dim(F) = k + 1$. We consider the closed sets $F_1 = pr(F)$, $G_1 = pr(G)$, $F_0 = \{\overline{y} \in (A^{\sharp})^d \mid \forall x \in H, (x, \overline{y}) \in F\} = \bigcap_{x \in H} F_x$, where $F_x = \{\overline{y} \in (A^{\sharp})^d \mid (x, \overline{y}) \in F\}$ is closed (note that we allow parameters in A^{\sharp} in the definition of the topology), and $G_0 = \{\overline{y} \in (A^{\sharp})^d \mid \forall x \in H, (x, \overline{y}) \in G\}$. There are three cases:

- 1. if $F_0 = F_1$, we see easily that $pr(F \setminus G) = F_0 \setminus G_0$, hence is constructible.
- 2. if $G_1 \subsetneq F_1$, we have a proper closed subset $(H \times G_1) \cap F \subsetneq F$, hence $dim((H \times G_1) \cap F) < dim(F)$ since F is irreducible. But we can write $pr(F \setminus G) = F_1 \setminus G_1 \cup pr(((H \times G_1) \cap F) \setminus G)$, and the result comes from the induction hypothesis.
- 3. if $F_0 \subsetneq F_1 = G_1$, we consider a generic point a of F_1 over \mathcal{U}_0 , a model of definition for F and G (note that F_1 is irreducible since F is). In particular $a \not\in F_0$, which implies that the fibre F(a) is finite, as it is a proper closed set of irreducible dimension one H. Furthermore, by Lemma 3.7, $G(a) \subsetneq F(a)$, and F(a) is irreducible as a closed set over $\mathcal{U}_0(a_\infty)$. It follows that F(a) is the orbit of any of its points under $\operatorname{Aut}(\mathcal{U}/\mathcal{U}_0(a_\infty))$ (note that $\mathcal{U}_0(a_\infty)$) is definably closed in \mathcal{U} , see [12]), and a fortiori, it is the orbit of G(a). Hence F(a) = G(a), a contradiction.

So we have proved Theorem 3.1.

4 Deriving function field ML from function field MM

4.1 The characteristic 0 case

We first deal with the characteristic 0 case.

We work with the notation introduced in Section 2: $k = \mathbb{C}, K = k(t)^{\text{alg}}, K^{\text{diff}}$ is a differential closure of the differential field (K, d/dt) and \mathcal{U} a saturated elementary extension of K^{diff} . In fact because the theory DCF_0 is ω -stable, we will mostly be working in K^{diff} , without needing to go up to the big model \mathcal{U} , contrary to the characteristic p case where it is essential, in the course of the proof, to go up to a sufficiently saturated model.

As explained before, we want to deduce Mordell-Lang from Manin-Mumford and the Kernel Theorem, avoiding reference to the dichotomy theorem for strongly minimal sets in differentially closed fields and/or the local modularity and strong minimality of A^{\sharp} when A is simple with k-trace 0. But we will use relatively softer ingredients of Hrushovski's proof in [15], among them the weak socle theorem, which we recall below.

Definitions 4.1. 1. Let G be a commutative connected group of finite Morley rank, definable in some ambient stable structure M. We say that G is generated (abstractly) by some sets X_1, \ldots, X_n if $G \subset acl(F \cup X_1 \cup \ldots \cup X_n)$, where F is a finite set. If $G \subset acl(F \cup X)$, where X is a strongly minimal set, G is said to be almost strongly minimal.

2. The model-theoretic (or stability-theoretic) socle S(G) of G, is the greatest connected definable subgroup of G which is generated by strongly minimal definable subsets of G.

3. For X a definable subset of G with Morley degree 1, define the model theoretic stabilizer of X in G, $Stab_G(X)$, to be $\{g \in G : MR(X \cap (X+g)) = MR(X)\}$ (so the stabilizer of the generic type of X over M).

4. A definable subgroup of G, H, defined over some B, is said to be rigid if, passing to a saturated model, all connected definable subgroups of H are defined over acl(B).

We will need to know some properties of socles:

Fact 4.2. (Lemma 4.6 in [15], or Section 7 in [18]) The group S(G) is an almost direct sum of pairwise orthogonal definable groups G_i , where each G_i is almost strongly minimal.

In this context, Hrushovski's so called "weak socle theorem" can be stated as:

Proposition 4.3. ((Prop. 4.3 in [15] or Prop; 2.10 in [4]) Let G be a commutative connected group of finite Morley rank. Suppose that the socle of G, S(G), is rigid. Let X be a definable subset of G of Morley degree one, such that $Stab_G(X)$ is finite. Then, up to subtracting a definable subset of X of strictly smaller Morley rank than that of X, some translate of X is contained in S(G).

The reason we call the above Proposition the weak socle theorem, is because a stronger statement is proved in [25] (see Theorem 2.1 there) in the special case of algebraic D-groups, and we have sometimes referred to the latter as "the socle theorem".

Let A be an abelian variety over K and $A^{\sharp} := A^{\sharp}(\mathcal{U})$, the smallest definable subgroup of $A(\mathcal{U})$ which contains the torsion of A (see Section 2.1).

We have already seen some basic properties of A^{\sharp} (Fact 2.3). Recall that A^{\sharp} is definable over K, connected with finite Morley rank and that if A is simple, A^{\sharp} is a g-minimal group.

We are going to use the following now classical fact (see for example 6.5 in [33]):

Fact 4.4. Let H be an almost strongly minimal definable group in \mathcal{U} , non orthogonal to the field of constants \mathcal{C} . Then there is an algebraic group R over \mathcal{C} and a definable isogeny from H onto $R(\mathcal{C})$.

The following, true in characteristic 0, are also well known, except maybe (iv).

Lemma 4.5. (i) $A(\mathcal{U})/A^{\sharp}$ embeds definably (in the sense of DCF_0) in (the group of \mathcal{U} -points of) a unipotent algebraic group over \mathcal{U} . (ii) A^{\sharp} is rigid.

(iii) If A is simple, then A has k-trace 0 if and only if it is not isogenous to an abelian variety over k if and only if A^{\sharp} is orthogonal to the field of constants C.

(iv) If H is a connected finite Morley rank definable subgroup of $A = A(\mathcal{U})$ containing A^{\sharp} , then $S(H) = A^{\sharp}$.

Remark: We will only use (iv) with the extra assumption that A has k-trace 0, but we give below the proof for the general case.

Proof. The following fundamental fact, due to Buium ([9]), and also proved in [22], is essential:

(*) Suppose $G = G(\mathcal{U})$ is a commutative connected algebraic group over \mathcal{U} and H is a Zariski-dense definable subgroup. Then G/H definably embeds

in a unipotent algebraic group, namely $(\mathcal{U}, +)^d$ for some d. (i) is given by (*).

(ii) It follows for example from [7], section 3.1, that any definable connected subgroup H of A^{\sharp} is of the form B^{\sharp} , for some abelian subvariety B of A. Hence it is the Kolchin closure of the torsion of B which is contained in the torsion of A, hence in A(K). This shows that H is itself also defined over K. (iii) The main point is to show that if A^{\sharp} is nonorthogonal to the field of constants, then A is isomorphic to an abelian variety over k. (The rest follows from the definition of trace 0 and simplicity of A.) This is already contained in [15] but we recall the proof. Applying Fact 4.4 and the dual isogeny we obtain a definable isogeny f from $R(\mathcal{C})$ to A^{\sharp} . By QE in DCF_0 , f is given by a rational function, and hence extends to a a surjective homomorphism of algebraic groups from R to A, which we again call f. Now R is a connected commutative algebraic group over \mathcal{C} , so as \mathcal{C} is algebraically closed, and Ker(f) contains the unique maximal connected linear algebraic subgroup of R, R/Ker(f) is again defined over \mathcal{C} , and isomorphic as an algebraic group to A.

(iv) This reduces to the case where A is simple. Now A^{\sharp} is g-minimal, hence, by Zilber's indecomposability theorem, generated by some strongly minimal subset, so $A^{\sharp} = S(A^{\sharp})$ and is contained in S(H).

Suppose by way of contradiction that S(H) properly contains A^{\sharp} . Then by (*) $S(H)/A^{\sharp}$ is an infinite finite-dimensional vector space over the constants C of U.

If A^{\sharp} has k-trace 0, then by (iii) A^{\sharp} is orthogonal to $S(H)/A^{\sharp}$, and so, by 4.2, $S(H) = A^{\sharp} + V$ where V is infinite, orthogonal to A^{\sharp} and has finite intersection with A^{\sharp} . As A is simple, V is Zariski dense in A, which contradicts Fact 2.3 (i) which says that A^{\sharp} is the smallest Zariski dense definable subgroup of A.

If A has non zero k-trace, as k is algebraically closed and we are in characteristic zero, A is isomorphic to an abelian variety over k, so we can suppose that A itself is defined over k. It follows that $A^{\sharp} = A(\mathcal{C})$. Now as both A^{\sharp} and $S(H)/A^{\sharp}$ are non orthogonal to the strongly minimal field of constants, S(H) is almost strongly minimal, $S(H) \subset acl(F \cup \mathcal{C})$. It follows by 4.4 and arguments which are now standard that S(H) is definably isomorphic to $R(\mathcal{C})$ for R a connected algebraic group over \mathcal{C} (see for example in [15] or [4] the proof of Proposition 2.1). Now let N be the biggest connected linear subgroup of R, N is defined over \mathcal{C} also, and we know that R/N is an abelian variety. Pulling back the situation by the definable isomorphism it follows that there is some infinite definable group V such that $V \subset S(H)$ and $V \cap A^{\sharp}$ is finite. As A is simple, V must be Zariski dense in A, but this contradicts Fact 2.3 (i) again.

Remark about (ii): In [15] it is shown that for any H, definable connected subgroup of $A(\mathcal{U})$ of finite Morley rank, S(H) is rigid. We do know that $A^{\sharp} = S(A^{\sharp})$ because $A^{\sharp} = \sum_{0 \leq i \leq n} A_i^{\sharp}$, where each A_i^{\sharp} are g-minimal, hence generated by a strongly minimal set. But we do not want to quote this result from [15] because its proof uses the full dichotomy theorem for strongly minimal sets in DCF_0 . In fact, the easy proof we use here is really the same as the one given in [15] to show that any abelian variety is rigid.

Remark on stabilizers. In the proof below, when we speak of stabilizers of (differential) algebraic subvarieties of a (differential) algebraic group we mean set-theoretic stabilizers. But in the contexts we consider this coincides with the model-theoretic stabilizer defined above. Likewise, if X is an irreducible differential algebraic subvariety of a finite Morley rank differential algebraic group G, then the conclusion of 4.3 says that X is, up to translation, contained in S(G).

Proving Mordell-Lang We now suppose that A has k-trace 0. The first steps are exactly the same as the ones in [15], which were inspired by Buium in [8]. First, quotienting by the connected component of $Stab_A(X)$ we obtain another abelian variety over K with k-trace 0. So, we will assume that $Stab_A(X)$ is finite (note that in the end when we get to the conclusion that X is the translate of a subabelian variety of A, the fact that the stabilizer is finite means that X is just one point).

By (*) in the proof of Lemma 2.2, A/A^{\sharp} definably embeds via some μ in a vector group. So $\mu(\Gamma)$ is contained in a finite-dimensional vector space over \mathcal{C} , the preimage of which we call H: a connected definable finite Morley rank subgroup of A containing both A^{\sharp} and Γ , and defined over K. As $X \cap \Gamma$ is Zariski dense in X, so is $X \cap H$. One reduces to the case when $X \cap H$ is irreducible as a differential algebraic variety. It follows that $Stab_H(X \cap H)$ is finite. So, by Lemma 4.5 (ii) and (iv), we can apply the weak socle theorem, Proposition 4.3, and, after replacing $X \cap H$ (and so X) by a suitable translate (which will be defined over K^{diff} , $X \cap H$ is contained in $A^{\sharp} = S(H)$.

From then on, we diverge from the proof in [15] which at this point appealed to the dichotomy.

We now consider \mathcal{A} , the K-definable group $A^{\sharp}(\mathcal{U})$ with all its induced

structure over K, namely equipped with predicates for K-definable subsets of Cartesian powers. As A^{\sharp} is definable, we automatically have quantifier elimination for \mathcal{A} and of course $\mathcal{A}(K^{\text{diff}})$ is an elementary substructure of \mathcal{A} . Furthermore, as A^{\sharp} is a definable group of finite Morley Rank, so is \mathcal{A} . Now consider the subset $\mathcal{A}(K)$ of \mathcal{A} . One can check easily that it must be algebraically closed in \mathcal{A} . Now A^{\sharp} is a finite sum of K-definable g-minimal subgroups A_i^{\sharp} . And we likewise have the \mathcal{A}_i , the A_i^{\sharp} with the induced structure. Each $A_i^{\sharp}(K)$ is clearly infinite as it contains the torsion of A_i , so by Corollary 2.10, $\mathcal{A}(K)$ is an elementary substructure of \mathcal{A} .

The following was proved in [20], with the extra assumption that A^{\sharp} was strongly minimal:

Claim. $A^{\sharp}(K) = A^{\sharp}(K^{\text{diff}}).$

Proof of claim. If $b \in A^{\sharp}(K^{\text{diff}})$ then tp(b/K) is isolated in DCF_0 . But then $tp(b/\mathcal{A}(K))$ is isolated in the structure \mathcal{A} . As $\mathcal{A}(K)$ is an elementary substructure of $\mathcal{A}, b \in A^{\sharp}(K)$.

Note that X is now defined over K^{diff} , and $X \cap A^{\sharp}(K^{\text{diff}})$ is Zariski-dense in X. By the Claim, $A^{\sharp}(K^{\text{diff}}) = A^{\sharp}(K)$ (so in fact X is defined over K), so, by the Theorem of the Kernel, $A^{\sharp}(K^{\text{diff}}) = A_{torsion}$, hence by Manin-Mumford, X is a translate of an abelian subvariety of A.

4.2 The characteristic *p* case

Once we have quantifier elimination by Theorem 3.1, the proof in the characteristic p case is substantially simpler than the characteristic 0 case, avoiding in particular any recourse to the socle theorem. We work with the notation introduced in Section 2: $k = \mathbb{F}_p^{\text{alg}}$, $K = k(t)^{\text{sep}}$, A, X are over K, Γ is contained in the prime-to-p division points of a finitely generated subgroup of A(K) (so $\Gamma < A(K)$ too), $X \cap \Gamma$ is Zariski-dense in X, and we take \mathcal{U} to be a saturated elementary extension of K. As before, A^{\sharp} denotes $A^{\sharp}(\mathcal{U})$ and A is a sum of simple abelian subvarieties $A_1, ..., A_n$, all defined over K. We know that each A_i^{\sharp} is a g-minimal subgroup of A(K) (Fact 2.3(iii)). Moreover A is assumed to have k-trace 0.

We now consider $\mathcal{A} := A^{\sharp}$ (respectively $\mathcal{A}_i := A_i^{\sharp}$) with their induced structure over K.

Lemma 4.6. The groups \mathcal{A} and \mathcal{A}_i are connected groups of finite Morley rank, the \mathcal{A}_i are g-minimal and $\mathcal{A}(K)$ is an elementary substructure of \mathcal{A} .

Proof. We know by Fact 3.2 that A^{\sharp} and the A_i^{\sharp} are connected groups, with relative Morley rank and that the A_i^{\sharp} are g-minimal subgroups of A^{\sharp} . By quantifier elimination of the induced structures (Theorem 3.1), it follows that \mathcal{A} and the \mathcal{A}_i are connected groups of finite Morley rank, that each of the \mathcal{A}_i remains a g-minimal subgroup of \mathcal{A} , and that \mathcal{A} is saturated. Consider the subset $\mathcal{A}(K)$ of \mathcal{A} . One can check easily that as K is an elementary substructure of \mathcal{U} , quantifier elimination implies that $\mathcal{A}(K)$ is an algebraically closed subset of \mathcal{A} . Furthermore, each $\mathcal{A}_i(K)$ is infinite (it contains all the prime-to-p torsion of A_i) and we can apply Corollary 2.10 to conclude.

We now complete the proof of Mordell-Lang:

First here we again follow Hrushovski's proof to see that $\Gamma \cap X$ can be "replaced" by $E \cap X$, for some E, translate of A^{\sharp} , without losing the Zariski denseness in X. Note that $p^{n+1}\Gamma$ has finite index in $p^n\Gamma$ for all n. As Γ meets X in a Zariski-dense set, we find cosets D_i of $p^i\Gamma$ in Γ such that i < jimplies $D_j \subseteq D_i$, and each D_i meets X in a Zariski-dense set. Hence we obtain a descending chain of cosets E_i of $p^iA(K)$ in A(K), each meeting X in a Zariski-dense set. Passing to the saturated model \mathcal{U} , let E be the intersection of the E_i , E is a translate of A^{\sharp} and by compactness $X \cap E$ is Zariski-dense in X.

At this point, Hrushovski appeals to the dichotomy theorem for thin minimal types in separably closed fields. We replace this by Manin-Mumford and the Theorem of the Kernel, but in order to do this, we need to go down to the smaller field K and we would need to know that $E(K) \cap \Gamma$ is dense in X. But K is no longer saturated, and in fact, although E is type definable over K, E(K) might be empty. So we need to work with a slightly bigger translate than E(K).

We consider the two-sorted structure $M = (\mathcal{A}, \mathcal{E})$, that is $(A^{\sharp}(\mathcal{U}), E(\mathcal{U}))$ with all the K-induced structure. It easily follows from Lemma 4.6 that Th(M) has finite Morley rank and moreover that the sort \mathcal{A} with induced structure has $\mathcal{A}(K)$ as an elementary substructure. Let $M_0 \prec M$ be prime (so atomic) over $\mathcal{A}(K)$. It follows that M_0 is of the form $(\mathcal{A}(K), \mathcal{E}_0)$ for some elementary substructure \mathcal{E}_0 of \mathcal{E} . Note that \mathcal{E}_0 is definably (without parameters) a principal homogeneous space for $\mathcal{A}(K)$.

Claim. $X \cap \mathcal{E}_0$ is Zariski-dense in X.

Proof of claim. Suppose not. Then there is a proper subvariety Z of X (defined over some field) such that $X \cap \mathcal{E}_0 \subseteq Z$. We may replace Z by the

Zariski closure of $X \cap \mathcal{E}_0$, and so we may assume that Z is defined over \mathcal{E}_0 . We now have that $X \cap \mathcal{E}_0 = Z \cap \mathcal{E}_0$. Now $Z \cap \mathcal{E}$ viewed as a set definable in the structure \mathcal{E} (or M) is defined over \mathcal{E}_0 , so as $\mathcal{E}_0 \prec \mathcal{E}$ we easily conclude that $X \cap \mathcal{E} = Z \cap \mathcal{E}$, contradicting Zariski-denseness of $X \cap \mathcal{E}$ in X. This completes the proof of the claim.

Let $a \in X \cap \mathcal{E}_0$. Let $X_1 = X - a$. Then $X_1 \cap A^{\sharp}(K)$ is Zariski-dense in X_1 . In particular X_1 is defined over K. Moreover using the Theorem of the Kernel, $X_1 \cap A_{torsion}$ is Zariski-dense in X_1 , so by MM, X_1 is a translate of an abelian subvariety of A. This completes the proof.

References

- D. Abramovich and F. Voloch, Towards a proof of the Mordell-Lang conjecture in positive characteristic, International Math. Research Notices 5 (1992), 103-115.
- [2] F. Benoist, E. Bouscaren, and A. Pillay, Semiabelian varieties over separably closed fields, maximal divisible subgroups, and exact sequences, Journal of the Institute of Mathematics of Jussieu DOI: 10.1017/S147474801400022X.
- [3] F. Benoist and F. Delon, Questions de corps de définition pour les variétés abéliennes en caractéristique positive, Journal de l'Institut de Mathématiques de Jussieu, vol. 7 (2008), 623–639.
- [4] E. Bouscaren, Proof of the Mordell-Lang Conjecture for Function fields, in *Model theory and algebraic geometry*, edited by E. Bouscaren, Lecture Notes in Mathematics 1696, 2nd edition, Springer, 1999.
- [5] E. Bouscaren and F. Delon, Groups definable in separably closed fields, Transactions of the AMS 354 (2002), no. 3, 945-966.
- [6] E. Bouscaren and F. Delon, Minimal groups in separably closed fields, J. Symbolic Logic 67 (2002), no. 1, 239259.
- [7] D. Bertrand and A. Pillay, A Lindemann-Weierstrass theorem for semiabelian varieties over function fields, Journal AMS, 23 (2010), 491-533.

- [8] A. Buium, Intersections in jet spaces and a conjecture of S. Lang, Annals of Math., 136 (1996), 557-567.
- [9] A. Buium, Differential algebraic groups of finite dimension, LNM 1506, Springer, 1992.
- [10] C. Corpet, Around the Mordell-Lang and Manin-Mumford conjectures in positive characteristic, preprint 2012.
- [11] F. Delon, Idéaux et types sur les corps séparablement clos, Supplément au Bulletin de la SMF, T116 (1988).
- [12] F. Delon, Separably closed fields, in *Model theory and algebraic geom*etry, edited by E. Bouscaren, Lecture Notes in Mathematics 1696, 2nd edition, Springer, 1999.
- [13] M. Hindry, Autour d'une conjecture de Serge Lang, Inventiones Math. 94 (1988), 567-603.
- [14] M. Hindry, Introduction to abelian varieties and the Mordell-Lang conjecture, in *Model Theory and Algebraic Geometry*, E. Bouscaren (Ed.), Lecture Notes in Mathematics 1696, 2nd edition, Springer, 1999.
- [15] E. Hrushovski, The Mordell-Lang conjecture for function fields, Journal AMS 9(1996), 667-690.
- [16] E. Hrushovski, The Manin-Mumford conjecture and the model theory of difference fields, Annals of Pure and Applied Logic, 112 (2001), 43-115.
- [17] E. Hrushovski and B. Zilber, Zariski geometries, Journal AMS 9(1996), 1-56.
- [18] D. Lascar, ω -stable groups, in *Model theory and algebraic geometry*, edited by E. Bouscaren, Lecture Notes in Mathematics 1696, 2nd edition, Springer, 1999.
- [19] D. Marker, M. Messmer and A. Pillay, Model theory of Fields, Lecture Notes in Logic 5, second edition, ASL-AK Peters, 2003.
- [20] D. Marker and A. Pillay, Differential Galois theory III: Some inverse problems, Illinois Journal of Mathematics, Vol. 41, Number 3 (1997), 453-461.

- [21] A. Omar Aziz, Type definable stable groups and separably closed fields, Ph. D. thesis, Leeds, 2012.
- [22] A. Pillay, Differential algebraic groups and the number of countable models, in *Model theory of Fields* above.
- [23] A. Pillay, *Geometric Stability Theory*, Oxford University Press, 1996.
- [24] A. Pillay, The model-theoretic content of Lang's conjecture, Model theory and algebraic geometry, edited by E. Bouscaren, Lecture Notes in Mathematics 1696, Springer, 1996.
- [25] A. Pillay, Mordell-Lang conjecture for function fields in characteristic 0, revisited, Compositio Math. 140 (2004) 64-68.
- [26] A. Pillay and M. Ziegler, Jet spaces of varieties over differential and difference fields, Selecta Math. 9 (2003), 579-599.
- [27] R. Pink and D. Rössler, On ψ -invariant subvarieties of semiabelian varieties and the Manin-Mumford conjecture, J. Algebraic Geometry 13 (2004), 771-798.
- [28] B. Poizat, *Stable groups*, Mathematical Surveys and Monographs, Vol. 87, American Mathematical Society.
- [29] D. Rössler, Infinitely *p*-divisible points on abelian varieties defined over function fields of characteristic p > 0, Notre Dame Journal of Formal Logic 54 (2013), no. 3-4, 579-589
- [30] D. Rössler, On the Manin-Mumford and Mordell-Lang conjectures in positive characteristic, Algebra and Number Theory 7 (2013), 2039-2057.
- [31] T. Scanlon, A positive characteristic Manin-Mumford theorem, Compositio Math. 141 (2005),1351-1364.
- [32] F. Wagner, Fields of finite Morley Rank, Journal of Symbolic Logic, 66(2001), 703-706
- [33] M. Ziegler, Introduction to stability theory and Morley rank, in *Model theory and algebraic geometry*, edited by E. Bouscaren, Lecture Notes in Mathematics 1696, Springer, 1996.

- [34] P. Ziegler, Mordell-Lang in positive characteristic, Rend. Sem. Mat. Univ. Padova, vol 134 (2015), 93-131.
- [35] B. Zilber, Zariski geometries, LMS Lecture series, CUP, 2011.

Franck Benoist Laboratoire de Mathématiques d'Orsay Univ. Paris-Sud, CNRS Université Paris-Saclay 91405 Orsay, France. franck.benoist@math.u-psud.fr

Elisabeth Bouscaren Laboratoire de Mathématiques d'Orsay Univ. Paris-Sud, CNRS Université Paris-Saclay 91405 Orsay, France. elisabeth.bouscaren@math.u-psud.fr

Anand Pillay Department of Mathematics University of Notre Dame 281 Hurley Hall Notre Dame, IN 46556, USA. apillay@nd.edu