LOCAL RAMSEY SPACES IN MATET FORCING EXTENSIONS AND FINITELY MANY NEAR-COHERENCE CLASSES

HEIKE MILDENBERGER

ABSTRACT. We introduce Gowers–Matet forcing with a finite sequence of pairwise non-isomorphic Ramsey ultrafilters over ω , and with this forcing we settle the long-standing problem about the spectrum of numbers of near-coherence classes. We prove that for any finite $n \geq 1$, there is a forcing extension with exactly n near-coherence classes of non-principal ultrafilters.

For evaluating the new forcing, we prove a strengthening of Gowers' theorem on colourings of $\mathrm{Fin}_k.$

Contents

1.	Introduction	1
2.	Matet forcing with Milliken–Taylor ultrafilters	ϵ
3.	Ramsey-theoretic computations in $\mathbb{M}(\mathcal{U})$ -extensions	11
4.	A name for a Matet-adequate family at limit stages	19
5.	Generalisation to Fin_k	23
6.	A Ramsey subspace of the space of Fin_k -sequences	28
7.	Localised Gowers–Matet Forcing	43
References		49

1. Introduction

We connect two lines of research: The topic of whether special kinds of ultrafilters from the ground model have extensions of the same kind in a forcing extension by a Ramsey-theoretic forcing and the investigation of the possible numbers of near-coherence classes. We answer Banakh's and Blass' question [1, Question 31] on the finite part of near-coherence spectrum.

On the existence of special ultrafilters: For ${}^{\omega}\omega$ -bounding forcings and P-points or even Ramsey ultrafilters the following is known: Kunen [27] proved that no Ramsey ultrafilter can be extended to a P-point after addition of any number of random reals at once, Shelah [37] constructed a model with no P-points, recently Chodounsky and Guzmán [16] proved that there are no P-points in the Silver model and that no P-point from the ground model can be extended in a Silver extension. On the other hand [41] and [20] proved that there are Milliken–Taylor ultrafilters

1

Date: July 25, 2019.

²⁰¹⁰ Mathematics Subject Classification. 03E05, 03E35, 05C55, 05D10.

Key words and phrases. Iterated proper forcing, combinatorics with block-sequences, P-point, Milliken–Taylor ultrafilters, near coherence of ultrafilters, selective coideals, preservation theorems, Gowers' Fin_k theorem.

in the Sacks model, indeed, any Milliken–Taylor ultrafilter from the ground model is preserved. Blass [9] proved that the minimum and the maximum projection of a Milliken–Taylor ultrafilter are Ramsey ultrafilters.

In the case of forcings adding an unbounded real, Ketonen proved that $\mathfrak{d}=\mathfrak{c}$ implies the existence of a P-point, Canjar [15] proved the generic existence of Ramsey ultrafilters under $\operatorname{cov}(\mathcal{M})=\mathfrak{c}$ and Eisworth [17] proved the generic existence of a Milliken–Taylor ultrafilter with the Galvin–Glazer technique (see, e.g. [26]) under the same condition. We refer the reader to [10] for the definitions of the cardinal characteristics $\operatorname{cov}(\mathcal{M})$, \mathfrak{u} , \mathfrak{d} , etc. We write \mathfrak{c} for 2^{\aleph_0} .

Here we work with variants of Matet forcing [29] that come from various constraints on the reservoir of the pure components of the conditions. The full Matet forcing preserves any P-point from the ground model [17, Theorem 4] and destroys any Ramsey ultrafilter, since it adds an unbounded real. The (non-complete) subforcings with pure parts from a Milliken–Taylor ultrafilter have specific preservation properties; they destroy some P-points and preserve others, see [17, Theorem 2.5]. We show that the reservoir Milliken–Taylor ultrafilter $\mathcal U$ can be extended to a new Milliken–Taylor ultrafilter after forcing. A new technical ingredient is the work with names for diagonal constructions.

We prove a preservation theorem for countable support iterations and show that there is a model of $\aleph_1 = \mathfrak{u} < \mathfrak{d} = \mathfrak{c} = \aleph_2$ with at least three names of different near coherence classes of ultrafilters and a Milliken–Taylor ultrafilter of character \mathfrak{c} .

We fix some $k \in \omega \setminus \{0\}$. We introduce Milliken–Taylor ultrafilters that are related to Hindman's theorem for colourings of Fin_k as Milliken–Taylor ultrafilters are related to Hindman's theorem, and investigate Matet forcing with Milliken–Taylor ultrafilters. We show that this generalisation gives nothing new in point of near coherence classes.

Then we change the condensation order into the Tetris condensation order $\sqsubseteq_k^{\text{Tetr}}$ (see Def. 6.1) and introduce a variant of Matet forcing in which particular projections of pure conditions are taken from prescribed pairwise nnc Ramsey ultrafilters $\mathcal{R}_{i,x}$, $i=1,\ldots,k,\ x=\min,\max$. To get the pure decision property and hence properness, we introduce a space

$$(\operatorname{Fin}_{k})^{\omega}(\bar{\mathcal{R}}) = \{ \bar{a} \in (\operatorname{Fin}_{k})^{\omega} : \forall (X_{i} \in \mathcal{R}_{i,\min}, Y_{i} \in \mathcal{R}_{i,\max})_{1 \leq i \leq k} \}$$

$$(1.1) \qquad (\exists^{\min-\mathrm{unb}} s \in \operatorname{TFU}_{k}(\bar{a}))$$

$$(\forall i \in \{1, \dots, k\}) (\min_{i}(s) \in X_{i} \wedge \max_{i}(s) \in Y_{i}) \}$$

and investigate $((\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}}), \sqsubseteq_k^{\operatorname{Tetr}})$ (see Def. 6.8) and prove a new generalisation of Gowers' theorem and of Blass' theorem [7, Theorem 2.3].

Theorem 6.16. Let $k \geq 1$ and let $\bar{\mathcal{R}} = \langle \mathcal{R}_{i,\min}, \mathcal{R}_{i,\max} : i = 1, ... k \rangle$ be a sequence of pairwise non nearly coherent Ramsey ultrafilters.

- (1) Any $\sqsubseteq_k^{\text{Tetr}}$ -descending ω -sequence of elements of $(\text{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ has a $\sqsubseteq_k^{\text{Tetr},*}$ -lower bound in $(\text{Fin}_k)^{\omega}(\bar{\mathcal{R}})$.
- (2) Let $n \in \omega \setminus \{0\}$ and $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ and let c be a colouring of $[\operatorname{TFU}_k(\bar{a})]_{<}^n$ into finitely many colours. Then there is a $\bar{b} \sqsubseteq_k \bar{a}$, $\bar{b} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ such that $[\operatorname{TFU}_k(\bar{b})]_{>}^n$ is c-monochromatic.

As a corollary we get the following theorems about existence and additional structure:

Theorem 6.33. For any pairwise nnc Ramsey ultrafilters $\mathcal{R}_{i,x}$, i = 1, ..., k, $x = \min$, max under CH or MA(σ -centred) there is a Gowers-Milliken-Taylor ultrafilter \mathcal{U} such that for i = 1, ..., k, $\min_i(\mathcal{U}) = \mathcal{R}_{i,\min}$ and $\max_i(\mathcal{U}) = \mathcal{R}_{i,\max}$.

Pairwise non-near-coherence and Ramseyness is necessary, however, more structure emerges in the cases $k \geq 2$: The statement about the higher core filters $\Phi_{\geq i+1}$ is novel. For the definition of $\Phi_{\geq i+1}$ see Def. 6.34.

Theorem 6.35. For any Gowers-Milliken-Taylor ultrafilter \mathcal{U} over Fin_k the following holds:

- (1) For each $1 \leq i \leq k$, $\mathcal{R}_{i,\min}$ and $\mathcal{R}_{i,\max}$ are nnc and for i < k, $\mathcal{R}_{i,\min}$ and $\mathcal{R}_{i,\max}$ are nnc $\Phi_{\geq i+1}(\mathcal{U})$.
- (2) The projections $\min_1(\mathcal{U}), \ldots, \min_k(\mathcal{U}), \max_k(\mathcal{U}), \ldots, \max_1(\mathcal{U})$ are pairwise non-nearly coherent Ramsey ultrafilters over ω .
- (3) All cores $\Phi_i(\mathcal{U})$ are nearly coherent.

Now we turn to the second line of research: the number of near-coherence classes of (non-principal) ultrafilters over ω , see Def. 1.1. In [11] a model with one near-coherence class is given. Blass [5] showed that under $\mathfrak{d} \leq \mathfrak{u}$ there are $2^{\mathfrak{u}}$ near-coherence classes. Banakh and Blass [1] showed: If the number of near-coherence classes is infinite then it is $2^{(2^{\omega})}$.

One of our main results about one iterand is:

Theorem 3.3. Let \mathcal{E} be a P-point and \mathcal{U} be a Milliken-Taylor ultrafilter with $\Phi(\mathcal{U}) \not\leq_{RB} \mathcal{E}$. Then in the forcing extension by $\mathbb{M}(\mathcal{U})$ the Milliken-Taylor ultrafilter \mathcal{U} is destroyed and can be completed to a Milliken-Taylor ultrafilter $\mathcal{U}^{\mathrm{ext}} \supseteq \mathcal{U}$ with $\Phi(\mathcal{U}^{\mathrm{ext}}) \not\leq_{RB} \mathcal{E}$.

The notion of a P-point will be explained in this section, the core $\Phi(\mathcal{U})$ is defined in Def. 2.11, the Rudin–Blass order \leq_{RB} is defined in Def. 2.12. Milliken–Taylor ultrafilters are defined in Def. 2.6(6), the forcing $\mathbb{M}(\mathcal{U})$ is defined in Def. 2.14. By [17, Theorem 2.5], \mathcal{E} generates a P-point in the extension.

Theorem 3.3 serves as a successor step in the forcing that is used in the following theorem:

Theorem 4.9. Assume CH. Then there is a countable support iteration of proper iterands $\mathbb{P} = \langle \mathbb{P}_{\alpha}, \mathbb{M}(\mathcal{U}_{\beta}) : \beta < \omega_{2}, \alpha \leq \omega_{2} \rangle$ such that in the extension there at least three near-coherence classes of ultrafilters and there is a Milliken-Taylor ultrafilter of character \mathfrak{c} .

We generalise from the set of blocks to Fin_k for some $k \geq 1$ and show that this does hardly change the situation.

We use the new Ramsey space $((\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}}), \sqsubseteq_k^{\operatorname{Tetr}})$ to define Gowers–Matet forcing $\mathbb{GM}_k(\bar{\mathcal{R}})$ in a localisation $\bar{\mathcal{R}}$ to a 2k-sequence $\bar{\mathcal{R}}$ of non-isomorphic Ramsey ultrafilters and prove our main result:

Theorem 7.18. Assume CH and let $k \geq 1$ and fix 2k pairwise nnc Ramsey ultrafilters $\bar{\mathcal{R}}_0 = \langle \mathcal{R}_{i,x,0} : i = 1, \ldots, k, x = \min, \max \rangle$. Then there is a countable support iteration of proper iterands $\mathbb{P} = \langle \mathbb{P}_{\alpha}, \mathbb{GM}_k(\bar{\mathcal{R}}_{\beta}) : \beta < \omega_2, \alpha \leq \omega_2 \rangle$ such that in the extension there exactly 2k+1 near-coherence classes of ultrafilters. Namely,

¹The last index in the ultrafilters indicates the iteration stage $\mathbf{V}^{\mathbb{P}_{\alpha}}$. The ultrafilters grow, i.e., $\mathcal{R}_{i,x,\alpha} = \mathcal{R}_{i,x,\beta} \cap \mathbf{V}^{\mathbb{P}_{\alpha}}$ for $\alpha < \beta \leq \omega_2$.

one class is represented by a P-point of character ω_1 and 2k classes represented by Ramsey ultrafilters $\mathcal{R}_{i,x,\omega_2}$, $i=1,\ldots,k$, $x=\min,\max$.

A slight variation of the model gives exactly 2k near coherence classes. We thus get the full finite near-coherence spectrum.

Corollary 7.20. For any $n \in \omega$, the statement "there are exactly n+1 near-coherence classes of ultrafilters" is consistent relative to ZFC.

By work of Mioduszewski our result has applications to analysis, namely the number of composants of $\beta(\mathbb{R}^+) - \mathbb{R}^+$ corresponds by [33, 34] to the number of near-coherence classes of ultrafilters. Blass [5] gives applications to cofinality classes of short non standard models of arithmetic, and to the decomposition of the ideal of compact linear operators on a Hilbert space into proper subideals. His results on equivalent characterisations of indecomposability can be translated to: For $n \geq 1$, There is a decomposition of the ideal of compact operators into n proper subideals such that the union of any two different of them is the whole ideal, if and only if there are exactly n near-coherence classes. The correspondence is defined in [13].

In the remainder of the introduction we recall some definitions from the realm of near coherence and special ultrafilters over ω . For the cardinal invariants \mathfrak{d} and $cov(\mathcal{M})$ we refer the reader to [10].

Let S be a countable set. By a *filter over* S we mean a non-empty subset of $\mathcal{P}(S)$ that is closed under supersets and under finite intersections and that does not contain the empty set. We call a filter over S non-principal if it contains all cofinite subsets of S. A \subseteq -maximal filter is an ultrafilter.

For $B \subseteq \omega$ and $f : \omega \to \omega$, we let $f[B] = \{f(b) : b \in B\}$ and $f^{-1}[B] = \{n : f(n) \in B\}$. The set of all infinite subsets of ω is denoted by $[\omega]^{\omega}$. For $\mathcal{B} \subseteq \mathcal{P}(\omega)$ we let $f(\mathcal{B}) = \{X \subseteq \omega : f^{-1}[X] \in \mathcal{B}\}$. This double lifting is an important function from $\mathcal{PP}(\omega)$ into itself. In analysis the special case of f being finite-to-one (that means that the preimage of each natural number is finite) is particularly useful, see e.g., [6].

From now on all filters over ω will be non-principal filters over ω , though we write only "filter" over ω . If $f: \omega \to \omega$ is finite-to-one, then also $f(\mathcal{F})$ is a non-principal filter. It is the filter generated by $\{f[X]: X \in \mathcal{F}\}$.

Definition 1.1. (1) A non-empty family $\mathcal{G} \subseteq [\omega]^{\omega}$ is called a *filter subbase (over* ω) if any intersection of finitely many elements of \mathcal{G} is infinite. We write

$$fil(\mathcal{G}) = \{ X \in [\omega]^{\omega} : (\exists n \in \omega)(\exists G_0 \dots \exists G_n \in \mathcal{G})(X \supseteq G_0 \cap \dots \cap G_n) \}$$

for the filter generated by \mathcal{G} . The *character* of a filter \mathcal{F} is the smallest size of a generating subbase.

- (2) Two filters $\mathcal{F}, \mathcal{G} \subseteq [\omega]^{\omega}$ are nearly coherent, if there is some finite-to-one $f: \omega \to \omega$ such that $f(\mathcal{F}) \cup f(\mathcal{G})$ generates a filter.
- (3) On the set of non-principal ultrafilters near-coherence is an equivalence relation (for a proof see [5], e.g.) whose equivalence classes are called *near-coherence classes*.
- (4) Two subsets \mathcal{H}_1 , \mathcal{H}_2 of $[\omega]^{\omega}$ are called nnc, if for any $X_i \in \mathcal{H}_i$, i = 1, 2 and any finite-to-one h there is $Y_i \subseteq X_i$, $Y_i \in \mathcal{H}_i$, i = 1, 2 such that $h[Y_1] \cap h[Y_2] = \emptyset$.

²If f is surjective, then $f(\mathcal{B})$ can be written as $\{f[X]: X \in \mathcal{B}\}$. In any case, $f(\mathcal{B})$ is contained in the set of supersets of members of the latter set.

(5) $\mathcal{H} \subseteq [\omega]^{\omega}$ is called *nowhere almost a filter* if for any $X \in \mathcal{H}$, and any finite-to-one h the set $\{h[Y]: Y \in \mathcal{H} \land Y \subseteq X\}$ does not have the finite intersection property.

For filters, nnc is the negation of near coherence. Near coherence is witnessed by a weakly increasing surjective finite-to-one function. f is weakly increasing if $x < y \to f(x) \le f(y)$. A coideal \mathcal{H} is nowhere almost a filter if and only if it is nowhere almost an ultrafilter. We will use the property for coideals.

We say "A is almost a subset of B" and write $A \subseteq^* B$ if $A \setminus B$ is finite. Similarly, the symbol $=^*$ denotes equality up to finitely many exceptions between elements of $[S]^{\omega}$ for a set S.

Let κ be a regular uncountable cardinal. An ultrafilter \mathcal{W} is called a P_{κ} -point if for every $\gamma < \kappa$, for every $A_i \in \mathcal{U}$, $i < \gamma$, there is some $A \in \mathcal{W}$ such that for any $i < \gamma$, $A \subseteq^* A_i$; such an A is called a *pseudo-intersection* of $\{A_i : i < \gamma\}$. A P_{\aleph_1} -point is called a P-point.

Let \mathbb{P} be a notion of forcing. We say that \mathbb{P} preserves an ultrafilter \mathcal{W} over I if

$$\Vdash_{\mathbb{P}}$$
 " $(\forall X \subseteq I)(\exists Y \in \mathcal{W})(Y \subseteq X \lor Y \subseteq I \setminus X)$ "

and in the contrary case we say " \mathbb{P} destroys \mathcal{W} ". In the first case $\{X \in [\omega]^{\omega} \cap \mathbf{V}[G] : (\exists Y \in \mathcal{W})X \supseteq Y\}$ is an ultrafilter in $\mathbf{V}[G]$ and \mathcal{W} generates an ultrafilter in $\mathbf{V}[G]$. We just say: \mathcal{W} is an ultrafilter in $\mathbf{V}[G]$. If \mathbb{P} is proper and preserves \mathcal{W} and \mathcal{W} is a P-point in the ground model, then \mathcal{W} stays a P-point in the forcing extension by [11, Lemma 3.2].

An ultrafilter W over ω is called a Q-point if for every strictly increasing sequence $\langle n_i : i \in \omega \rangle$ of natural numbers there is $X \in W$ such that for every $i, |X \cap [n_i, n_{i+1})| \leq 1$. Any Q-point from the ground model ceases to be a Q-point after adding an unbounded real.

An ultrafilter \mathcal{R} is called *selective* (or *Ramsey ultrafilter*) if it is a P-point and a Q-point. We use the von Neumann natural numbers $n = \{0, \dots, n-1\}$. We often use the following, equivalent (see [14, Theorem 4.9] characterisation of selectivity:

(*) For any \subseteq -descending sequence $\langle A_n : n \in \omega \rangle$ of sets $A_n \in \mathcal{R}$ there is $A \in \mathcal{R}$ such that $A \subseteq A_0$ and $(\forall n \in A)(A \setminus (n+1) \subseteq A_n)$.

Kunen [27] contains more information on Ramsey ultrafilters. Two Ramsey ultrafilters are nnc iff they are not isomorphic [4].

In forcing, we follow the Kunen style that the stronger condition is the *smaller* one. This corresponds to the close relationship between the \leq -relation in Matet forcing and the condensation relation \sqsubseteq on the second components, the so-called pure parts, of a condition in Matet forcing.

Readers who want to focus on our extension of Gowers' theorem in Theorem 6.16, that does not contain any forcing, can just read the notational part from Def. 5.1 to Def. 5.5 and then the new Ramsey-theoretic work from the beginning of Section 6 through Question 6.31.

Biggest thanks go to the referee for careful reading and detecting some gaps and errors in an earlier version. I am grateful for Andreas Blass for explaining me his proof of [7, Theorem 2.1].

2. Matet forcing with Milliken-Taylor ultrafilters

In this section we review results of Blass, Eisworth, and Hindman and carry it a bit further. Our nomenclature follows Blass [7], Eisworth [17] and Todorčević [40].

Definition 2.1.

- (1) We let Fin denote the set of finite non-empty subsets of ω .
- (2) An element $a \in \text{Fin is called a block.}$
- (3) For $a, b \in \text{Fin}$ we write a < b if $(\forall n \in a)(\forall m \in b)(n < m)$.
- (4) We define a well-order (of type ω) $\leq_{\text{lex,Fin}}$ on the set Fin via $a <_{\text{lex,Fin}} b$ if $\max(a) < \max(b)$ or $(\max(a) = \max(b))$ and $\min(a \triangle b) \in a$.
- (5) A sequence $\bar{a} = \langle a_n : n \in \omega \rangle$ of members of Fin is called *unmeshed* if for all $n, a_n < a_{n+1}$.
- (6) By $(\text{Fin})^{\omega}$ we denote the set of unmeshed sequences of members in Fin.
- (7) Let a, b be blocks. We let $a \cup b$ be undefined unless a < b. Otherwise, $a \cup b$ is defined as the union.
- (8) A set $X \subseteq \text{Fin}$ is called *min-unbounded* if for any $n \in \omega$ there is some $x \in X$ with $\min(x) \ge n$.
- (9) (Fin, \cup) is a partial semigroup. The associative partial binary operation \cup lifts to $\gamma(\text{Fin})$, the space of min-unbounded ultrafilters over Fin, as follows (and we write \cup for the lifted operation):

$$\mathcal{U}_1 \cup \mathcal{U}_2 = \{ X \subseteq \text{Fin} : \{ s : \{ t : s \cup t \in X \} \in \mathcal{U}_2 \} \in \mathcal{U}_1 \}.$$

For details and history see [26, Section 4.1].

- (10) If X is a subset of Fin, we write FU(X) for the set of all unions of finitely many members of X. We write $FU(\bar{a})$ instead of $FU(\{a_n : n \in \omega\})$. We call X an FU-set if there is some unmeshed \bar{a} such that $X = FU(\bar{a})$.
- (11) For $\emptyset \neq X \subseteq \text{Fin we let } \min_{\leq_{\text{lex},\text{Fin}}}(X) \text{ be the } \leq_{\text{lex},\text{Fin}}\text{-element of } X.$ For $\bar{a} \in (\text{Fin})^{\omega}$ we let $\min_{\text{Fin}}(\bar{a}) = \min_{\leq_{\text{lex},\text{Fin}}}\{a_n : n < \omega\}$, which is a_0 .
- (12) A filter over Fin is a non-empty subset of $\mathcal{P}(\text{Fin})$ that is closed under intersections and supersets and does not contain the empty set.
- (13) For $X \subseteq \text{Fin}$, the set $(\text{FU}(X))^{\omega}$ denotes the collection of all infinite unmeshed sequences in FU(X). For $\bar{a} \in (\text{Fin})^{\omega}$, the set $(\text{FU}(\bar{a}))^{\omega}$ denotes the collection of all infinite unmeshed sequences in $\text{FU}(\bar{a})$ (recall item (9)).
- (14) For $Y \subseteq \text{Fin and } s \in \text{Fin we write } (Y ; \text{past } s) \text{ for } \{u \in Y : \max(s) < \min(u)\}.$ For $\bar{a} \in (\text{Fin})^{\omega}$ and $s \in \text{Fin we write } (\bar{a} ; \text{past } s) \text{ for } \langle a_n : n \geq n_0 \rangle, \text{ where } n_0 = \min\{n : \max(s) < \min(a_n)\}.$

Now the set of min-unbounded elements $\mathcal{P}(\text{Fin})$ is equipped with a partial order \sqsubseteq that makes it into a topological Ramsey space in the sense of [40]. We will work with the (closed) subspace of sets of the form FU(\bar{a}).

³Note that this is not the usual lexicographic well-order; e.g., $\{0,1\} <_{\text{lex,Fin}} \{1\}$. The aim of this well-order is to define the \sqsubseteq -largest common lower bound of two \sqsubseteq -compatible elements of $(\text{Fin})^{\omega}$ by induction on the blocks. The relation \sqsubseteq is defined in Def. 2.2.

⁴Differences in behaviour of union ultrafilters that contain only min-unbounded sets, and ordered-union ultrafilters that have a bases of unmeshed sets, are not yet known. Any known proof of Hindman's theorem results in an unmeshed sequence.

Definition 2.2. Given X and $Y \subseteq \operatorname{Fin}$, we say that Y is a condensation of X and we write $Y \subseteq X$ if $Y \subseteq \operatorname{FU}(X)$. We say Y is almost a condensation of X and we write $Y \subseteq^* X$ if there is an $n \in \omega$ such that $(Y ; \operatorname{past} \{n\})$ is a condensation of X.

The "blurred" order \sqsubseteq^* is an $< \omega_1$ -complete preorder.

We use the relation \sqsubseteq mainly for $Y = \operatorname{range}(\bar{b})$ with $\bar{b} \in (\operatorname{Fin})^{\omega}$, $X = \operatorname{range}(\bar{a})$ for $\bar{a} \in (\operatorname{Fin})^{\omega}$, and then we write $\bar{b} \sqsubseteq \bar{a}$ for $\operatorname{range}(\bar{b}) \sqsubseteq \operatorname{range}(\bar{a})$, and analogously for \sqsubseteq^* .

Definition 2.3. Let $\bar{a}, \bar{b} \in (\operatorname{Fin})^{\omega}$. We say \bar{a} and \bar{b} are compatible, if there is a $\bar{c} \sqsubseteq \bar{a}, \bar{b}$. In the contrary case we write $\bar{a} \perp \bar{b}$.

Lemma 2.4. If \bar{a} and \bar{b} are compatible, there is a weakest $\bar{c} \in (\operatorname{Fin})^{\omega}$ such that $\bar{c} = \langle c_n : n < \omega \rangle \sqsubseteq \bar{b}, \bar{a}$. Moreover, \bar{c} is given by the following procedure. By induction on n we define c_n as follows $c_0 = \min_{\langle \operatorname{lex}, \operatorname{Fin} \rangle} (\operatorname{FU}(\bar{a}) \cap \operatorname{FU}(\bar{b})), c_{n+1} = \min_{\langle \operatorname{lex}, \operatorname{Fin} \rangle} \{s \in \operatorname{FU}(\bar{a}) \cap \operatorname{FU}(\bar{b}) : c_n < s \}.$

Proof. By definition $\bar{c} \sqsubseteq \bar{a}, \bar{b}$. To see that \bar{c} is the largest witness, for a contradiction, we suppose that there is $\bar{d} \sqsubseteq \bar{a}, \bar{b}$ such that $\bar{d} \not\sqsubseteq \bar{c}$. We take the first n such that $d_n \neq c_n$.

Since \bar{a} and \bar{b} are unmeshed sequences, there are $i_1 < \cdots < i_k$, and $j_1 < \cdots < j_\ell$ such that

$$(2.1) c_n = a_{i_1} \cup \cdots \cup a_{i_k} = b_{j_1} \cup \cdots \cup b_{j_\ell}$$

and there are no common maxima of support in the sequence $(a_{i_1}, a_{i_1+1}, \ldots, a_{i_k-1})$ and in the sequence $(b_{j_1}, b_{j_1+1}, \ldots, b_{j_\ell-1})$ in at any pair $(a_{\varrho}, b_{\sigma})$ with $\varrho \in [j_1, \ldots, j_\ell)$ and $\sigma \in [j_1, \ldots, j_\ell)$ and there are no $i < i_1$. $j < j_1$, and a_i , b_j past $\max(\sup(c_{n-1})(-1)$ in the case n = 1) such that $\min(\sup(a_i)) = \min(\sup(b_i))$.

First case: $\max(\operatorname{supp}(d_n)) > \max(\operatorname{supp}(c_n))$. However, then there is a block d'_n with $\max(\operatorname{supp}(d'_n)) = \max(\operatorname{supp}(c_n))$ and there is $d''_n \in \operatorname{FU}(\bar{a}; \operatorname{past} c_{n-1}) \cap \operatorname{FU}(\bar{b}; \operatorname{past} c_n)$ such that $d_n = d'_n + d''_n$ and \bar{d} could be replaced the strictly larger upper bound

$$\bar{d}' = \langle d_0, \dots, d_{n-1}, d'_n, d''_n, d_{n+1}, \dots \rangle \supset \bar{d}.$$

In the second and the third case, applied to d'_n , we show that such a $\bar{d}' \not\sqsubseteq \bar{c}$ cannot exist and thus we get a contradiction.

Second case: $\max(\operatorname{supp}(d_n)) < \max(\operatorname{supp}(c_n))$. Since \bar{a} and \bar{b} are unmeshed sequences, then \bar{c} could be broken up, similar to \bar{d} is the previous case. This possibility to break up contradicts the $<_{\operatorname{lex,Fin}}$ -minimality of c_n .

Third case: $\max(\operatorname{supp}(d_n)) = \max(\operatorname{supp}(c_n))$. First subcase $d_n <_{\operatorname{lex},\operatorname{Fin}} c_n$. This contradicts the choice of c_n .

Second subcase: $d_n >_{\text{lex,Fin}} c_n$. Then d_n uses in its sum not all the summands $a_{i_1}, \ldots, a_{i_\ell}$ that combine to c_n and hence $d_n = 0$, because leaving off summands on one side or on both sides of (2.1) is not possible by the choice of c_n and since $\max(\text{supp}(d_n)) = \max(\text{supp}(c_n))$. However, d_n is a block and hence not zero. Thus also this case ends in a contradiction.

We write $\bar{c} = \bar{a} \wedge \bar{b}$.

Definition 2.5. A set non-empty subset $C \subseteq (\operatorname{Fin})^{\omega}$ is called *centred*, if for any finite $C \subseteq C$ there is $\bar{a} \in C / \bar{a} \in (\operatorname{Fin})^{\omega}$ that is a condensation of any $\bar{c} \in C$ and if C is closed under finite alterations i.e., if $\bar{d} \in C$ and there are $n, m \in \omega$ such that $\langle d_m : m \geq m_0 \rangle = \langle e_n : n \geq n_0 \rangle$ then $\bar{e} \in C$.

We specialise \mathcal{C} further.

Definition 2.6.

- (1) Let \mathcal{F} be a filter. A basis \mathcal{B} of \mathcal{F} is a subset of \mathcal{F} such that $(\forall F \in \mathcal{F})(\exists B \in \mathcal{B})(B \subseteq F)$.
- (2) A non-principal filter \mathcal{F} over Fin is said to be a *union* filter if it has a basis of sets of the form $\mathrm{FU}(X)$ for $X\subseteq\mathrm{Fin}$ such that the elements of X are pairwise disjoint. Note $\mathrm{FU}(X)$ need not be an FU-set.
- (3) A non-principal filter \mathcal{F} over Fin is said to be an *min-unbounded* filter if $(\forall n \in \omega)(\exists X \in \mathcal{F})(\forall s \in X)(\min(s) > n)$.
- (4) A non-principal filter \mathcal{F} over Fin is said to be an *ordered-union* filter if it has a basis of sets of the form $\mathrm{FU}(\bar{d})$ for $\bar{d} \in (\mathrm{Fin})^{\omega}$.
- (5) Let μ be an uncountable cardinal. A union filter is said to be $(<\mu)$ -stable if, whenever it contains $\mathrm{FU}(X_\alpha)$ for $X_\alpha\subseteq\mathrm{Fin},\ \alpha<\kappa$, for some $\kappa<\mu$, then it also contains some $\mathrm{FU}(Y)$ for some Y such that for each $\alpha<\kappa$ there is $n_\alpha\in\omega$ with $(Y\ ;\mathrm{past}\ \{n_\alpha\})\subseteq\mathrm{FU}(X_\alpha)$. Such an Y is called a lower bound of $\{X_\alpha:\alpha<\kappa\}$. For " $<\omega_1$ -stable" we say "stable".
- (6) A stable ordered-union ultrafilter is also called a Milliken-Taylor ultrafilter.
- (7) An ultrafilter is called *idempotent* if $\mathcal{U} \cup \mathcal{U} = \mathcal{U}$.

Ordered-union ultrafilters need not exist, as their existence implies the existence of Q-points [7] and there are models without Q-points [32]. Even union ultrafilters need not exist: Blass [9, Theorem 38] showed that the existence of a union ultrafilter implies the existence of at least two near-coherence classes of ultrafilters. In [11] Blass and Shelah show that it is consistent relative to ZFC to have exactly one near-coherence class of non-principal ultrafilters. Union ultrafilters are idempotent. Idempotent ultrafilters exist by the Ellis–Numakura Lemma [18, 35]. With the help of Hindman's theorem one shows that CH or Martin's Axiom for σ -centred posets and $< 2^{\omega}$ dense sets implies that (even $< 2^{\omega}$ -) stable Milliken–Taylor ultrafilters exist [7]. We recall Hindman's theorem:

Theorem 2.7. (Hindman, [24, Corollary 3.3]) If the set Fin is partitioned into finitely many pieces then there is a set $\bar{d} \in (\text{Fin})^{\omega}$ such that $\text{FU}(\bar{d})$ is included in one piece.

The theorem also holds if instead of Fin we partition some $\mathrm{FU}(\bar{c})$ for a $\bar{c} \in (\mathrm{Fin})^{\omega}$ and search for a homogeneous sequence $\bar{d} \sqsubseteq \bar{c}$, see [7, p. 92].

Corollary 2.8. (See [7, p. 93].) Under CH or $MA_{<2^{\omega}}(\sigma\text{-centred})$, for every $\bar{a} \in (\text{Fin})^{\omega}$ there is a Milliken-Taylor ultrafilter \mathcal{U} such that $FU(\bar{a}) \in \mathcal{U}$.

We let for $X \subseteq \text{Fin}$, $[X]_{<}^n$ be the set of increasing unmeshed *n*-sequences of members of X. For the evaluation of our forcings, Taylor's theorem [39] is utilised:

Theorem 2.9. (Taylor [39].) Let \mathcal{U} be a Milliken-Taylor ultrafilter, $n \in \omega$. Let $[\operatorname{Fin}]^n_{\leq}$ be partitioned into finitely many sets. Then there is $A \in \mathcal{U}$ such that $[\operatorname{FU}(A)]^n_{\leq}$ is monochromatic.

Corollary 2.10. ([17, Cor. 1.3]) Existence of diagonal lower bounds in $((\operatorname{Fin})^{\omega}, \sqsubseteq)$. Let \mathcal{U} be a Milliken-Taylor ultrafilter, and let $\langle A_n : n \in \omega \rangle$ be a

 \sqsubseteq -descending sequence of members of \mathcal{U} . Then there is a $B \in \mathcal{U}$ such $B \sqsubseteq A_0$ and

(2.2)
$$(\forall s \in B)((B ; past s) \text{ is a condensation of } A_{\max(s)+1}).$$

Such a B is called a diagonal lower bound of $\langle A_n : n \in \omega \rangle$.

In Equation (2.2) we can equivalently let s range over FU(B). If B is a diagonal lower bound and $B' \subseteq B$ then B' is a diagonal lower bound as well.

Definition 2.11. Let \mathcal{H} be a subset of $\mathcal{P}(Fin)$.

(1) The core of \mathcal{H} is the set $\Phi(\mathcal{H}) \subseteq [\omega]^{\omega}$ such that

$$X \in \Phi(\mathcal{H}) \text{ iff } (\exists Y \in \mathcal{H})(\bigcup Y \subseteq X).$$

(2) The minimum projection of \mathcal{H} is the set

$$\min(\mathcal{H}) = \{\min[Y] : Y \in \mathcal{H}\},\$$

where

$$\min[Y] = \{\min(y) : y \in Y\},\$$

and analogously we define the maximum projection $\max(\mathcal{H})$.

- (3) For $\bar{a} \in (\operatorname{Fin})^{\omega}$ we write $\min[\bar{a}] = \{\min(a_i) : i \in \omega\}$.
- (4) For $\bar{a} \in (\text{Fin})^{\omega}$ we write $\text{set}(\bar{a}) = \bigcup \{a_i : i \in \omega\}.$
- (5) For $\mathcal{H} \subseteq (\operatorname{Fin})^{\omega}$ we let $\Phi(\mathcal{H}) = \Phi(\{\operatorname{range}(\bar{a}) : \bar{a} \in \mathcal{H}\})$ and $\min(\mathcal{H}) = \{\min[\bar{a}] : \bar{a} \in \mathcal{H}\}.$

Note that $\min[\bar{a}] = \min[\mathrm{FU}(\bar{a})]$ and $\bigcup \{a_n : n \in \omega\} = \bigcup \mathrm{FU}(\bar{a}).$

If $\mathcal{C} \subseteq (\operatorname{Fin})^{\omega}$ is centred, then the core of \mathcal{C} is a filter over ω . For centred $\mathcal{C} \subseteq (\operatorname{Fin})^{\omega}$ and for filters \mathcal{C} over Fin the projection $\min(\mathcal{C})$ is a filter and $\min(\mathcal{C})$, $\max(\mathcal{C}) \supseteq \Phi(\mathcal{C})$. Blass ([7, 3.6–3.9] together with [9, Theorem 38]) showed that for a Milliken–Taylor ultrafilter \mathcal{U} , $\min(\mathcal{U})$ and $\max(\mathcal{U})$ are nnc Ramsey ultrafilters.

The cores of centred systems are just filters over ω . Even if \mathcal{U} is an ultrafilter over Fin, for any finite-to-one f, $f(\Phi(\mathcal{U}))$ need not be an ultrafilter over ω . Blass [9, Theorem 38]) showed that for union-ultrafilters \mathcal{U} , for any finite-to-one f, $f(\Phi(\mathcal{U}))$ is not an ultrafilter because among its supersets there two nnc ultrafilters generated by the f-images of the minimum projection and the maximum projection.

If \mathcal{U} is a Milliken–Taylor ultrafilter, then $\Phi(\mathcal{U})$ does not have a pseudointersection (see [17, Prop. 2.3]) and also any finite-to-one image of $\Phi(\mathcal{U})$ does not have a pseudointersection by the same proof. Hence, by Talagrand [38] $\Phi(\mathcal{U})$ is not meagre. Thus the filter dichotomy principle (see, e.g., [10]) precludes the existence of a Milliken–Taylor ultrafilter.

Definition 2.12. The *Rudin–Blass ordering* for filters over ω is defined as follows: Let $\mathcal{F} \leq_{RB} \mathcal{G}$ if there is a finite-to-one f such that $f(\mathcal{F}) \subseteq f(\mathcal{G})$.⁵

For filters \mathcal{F} , \mathcal{G} , the relation $\mathcal{F} \leq_{RB} \mathcal{G}$ implies that \mathcal{F} is nearly coherent to \mathcal{G} . If \mathcal{G} is an ultrafilter, also the converse holds.

Now we turn to forcing:

⁵Also the definition $f(\mathcal{F}) \subseteq \mathcal{G}$ is used in the literature. If \mathcal{G} is a P-point both definitions are closely related.

Definition 2.13. Conditions in *Matet forcing*, \mathbb{M} , are pairs (s, \bar{c}) such that $s \in \text{Fin}$ and $\bar{c} \in (\text{Fin})^{\omega}$ and $s < c_0$. The forcing order is $(t, \bar{d}) \leq (s, \bar{c})$ (recall the stronger condition is the smaller one) if $s \subseteq t$ and t is the union of s and finitely many of the c_n and \bar{d} is a condensation of \bar{c} .

Definition 2.14. For a family $\mathcal{H} \subseteq (\operatorname{Fin})^{\omega}$, the notion of forcing $\mathbb{M}(\mathcal{H})$ consists of all pairs (s, \bar{a}) such that $\bar{a} \in \mathcal{H}$. The forcing order is the same as in the Matet forcing.

We write $set(\bar{a}) = \bigcup \{a_n : n \in \omega\}.$

For a centred system $\mathcal{C} \subseteq (\operatorname{Fin}_k)^{\omega}$, the set $\Phi(\mathcal{C})$ is the filter fil($\{\operatorname{set}(\bar{a}) : \bar{a} \in \mathcal{C}\}$). The forcing $\mathbb{M}(\mathcal{C})$ diagonalises $\Phi(\mathcal{C})$. Let G be a $\mathbb{M}(\mathcal{C})$ -generic filter over \mathbf{V} . Then the generic real

$$\mu := \bigcup \{ s : \exists \bar{c} : (s, \bar{c}) \in G \}$$

is a pseudointersection of $\Phi(\mathcal{C})$.

The following property of Milliken–Taylor ultrafilters $\mathcal U$ will be important for our proof:

Theorem 2.15. (Eisworth [17, " \rightarrow " Cor. 2.5, " \leftarrow " Theorem 4, this direction works also with non-P ultrafilters]) Let \mathcal{U} be a Milliken-Taylor ultrafilter over Fin and let \mathcal{W} be a P-point. Then $\mathcal{W} \ngeq_{RB} \Phi(\mathcal{U})$ if and only if \mathcal{W} continues to generate an ultrafilter after we force with $\mathbb{M}(\mathcal{U})$.

We remark:

Proposition 2.16. Suppose that \mathcal{U} is a Milliken-Taylor ultrafilter and \mathcal{F} is a filter over ω and $\Phi(\mathcal{U}) \leq_{RB} \mathcal{F}$. Then $\mathbb{M}(\mathcal{U})$ forces that for any finite-to-one function from the ground model, $f(\mathcal{F})$ is not an ultrafilter.

Proof. Let $f \in \mathbf{V}$ be finite-to-one such that $f(\Phi(\mathcal{U})) \subseteq f(\mathcal{F})$. Let G be $\mathbb{M}(\mathcal{U})$ -generic over \mathbf{V} , and let μ be a name for the generic real μ . We show:

$$(\forall (s,\bar{a}) \, \in \, \mathbb{M}(\mathcal{U}))(\forall Y \, \in \, \mathcal{F})(\exists (t,\bar{b}) \, \leq_{\mathbb{M}(\mathcal{U})} \, (s,\bar{a}))(t,\bar{b}) \, \Vdash_{\mathbb{M}(\mathcal{U})} \, f[Y] \, \cap \, f[\underline{\check{\mu}}] \, \neq \, \emptyset$$

$$(\forall (s,\bar{a}) \in \mathbb{M}(\mathcal{U}))(\forall Y \in \mathcal{F})(\exists (t,\bar{b}) \leq_{\mathbb{M}(\mathcal{U})} (s,\bar{a}))(t,\bar{b}) \Vdash_{\mathbb{M}(\mathcal{U})} f[Y] \cap (\omega \smallsetminus f[\mu]) \neq \emptyset.$$

Let (s, \bar{a}) and Y be given. Since $f[\operatorname{set}(\bar{a})] \in f(\Phi(\mathcal{U})) \subseteq f(\mathcal{F})$, we have $f[\operatorname{set}(\bar{a})] \cap f[Y]$ is infinite. So there is $t \in \operatorname{FU}(\bar{a})$ such that $f[t] \cap f[Y] \neq \emptyset$. It follows that $(s \cup t, \bar{a} : \operatorname{past} t) \Vdash_{\mathbb{M}(\mathcal{U})} f[Y] \cap f[\mu] \neq \emptyset$.

Now for the second property:

Next we define a colouring h of $[FU(\bar{a}; past s)]^2 < by$

$$h(u < v) = \begin{cases} 1 & \text{if } f[u] < f[v] \land \\ & \left(\max(f[u]), \min(f[v]) \right) \cap f[Y] \neq \emptyset, \\ 0 & \text{else.} \end{cases}$$

Since \mathcal{U} is a Milliken–Taylor ultrafilter by Theorem [39] there is a monochromatic $\bar{b} \sqsubseteq (\bar{a} ; \text{past } s), \ \bar{b} \in \mathcal{U}$. Since f is finite-to-one, the colour is 1. So we have $(s, \bar{b}) \Vdash_{\mathbb{M}(\mathcal{U})} f[Y] \cap (\omega \setminus f[\mu]) \neq \emptyset$.

Now let f be a name for the function $n \mapsto |\mu \cap n|$. Then

$$\mathbb{M}(\mathcal{U}) \Vdash f$$
 is finite-to-one and $f[\mu] =^* \omega$

and the above proof breaks down. Information on $g(\mathcal{F})$ for particular filters \mathcal{F} with $\Phi(\mathcal{U}) \leq_{\mathrm{RB}} \mathcal{F}$ and any name g for a finite-to-one function is contained in Theorem 3.25.

Definition 2.17. Let \mathcal{F} be a filter. $\mathcal{F}^+ = \{X \in [\omega]^\omega : (\forall Y \in \mathcal{F})(Y \cap X \neq \emptyset)\}.$

For a non-principal \mathcal{F} , \mathcal{F}^+ coincides with $\{X \in [\omega]^\omega : (\forall Y \in \mathcal{F})(Y \cap X \in [\omega]^\omega)\}.$

3. Ramsey-theoretic computations in $\mathbb{M}(\mathcal{U})$ -extensions

Now we consider the Ramsey space $((\operatorname{Fin})^{\omega}, \sqsubseteq)$ in an $\mathbb{M}(\mathcal{U})$ -extension. We use the notation $\mathbf{V}^{\mathbb{P}}$ for any $\mathbf{V}[G]$, with a \mathbb{P} -generic filter G over \mathbf{V} . All computations are about names and we use the forcing theorem freely and identify names often with their evaluations. The first aim is to examine \mathcal{U} in the forcing extension $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$. Is there an extension of the destroyed Milliken–Taylor ultrafilter \mathcal{U} to a new Milliken–Taylor ultrafilter? We show that under CH there are even 2^{ω_1} possibilities with pairwise nnc cores. This will be the successor step of an iteration of iterands of type $\mathbb{M}(\mathcal{U}_{\alpha})$. We introduce Ramsey-theoretic computations with names for elements of $(\operatorname{Fin})^{\omega}$ in order to establish the existence of names for Milliken–Taylor ultrafilters. Although in the end, like in Cor. 2.10, min-unbounded FU-subsets of Fin are the elements of Milliken–Taylor ultrafilters, intermediate work is better carried out with sequences $\bar{a} \in (\operatorname{Fin})^{\omega}$. We use letters $\bar{a}, \bar{b}, \bar{a}, \ldots, A, B, \ldots$ for elements of $(\operatorname{Fin})^{\omega}$, where capital letters are in particular used in work with sequences of sequences. Capital letters are also used for subsets of Fin.

- **Definition 3.1.** (1) Let $\bar{a} \in (\operatorname{Fin})^{\omega}$ and $X \in [\omega]^{\omega}$. We let $\bar{a} \upharpoonright X = \langle a_n : n \in \omega, a_n \subseteq X \rangle$. Note, we do not take those a_n with $a_n \cap X \neq \emptyset$ that are not subsets of X.
- (2) Let $\mathcal{U} \subseteq (\operatorname{Fin})^{\omega}$ and $X \in [\omega]^{\omega}$. We use the restriction symbol also for subsets of $(\operatorname{Fin})^{\omega}$ and let $\mathcal{U} \upharpoonright X = \{\bar{a} \upharpoonright X : \bar{a} \in \mathcal{U}\}$).

Lemma 3.2. Now let μ be a name for the Matet generic real $\bigcup \{s : \exists \bar{a}(s,\bar{a}) \in G\}$.

- (1) $\mathbb{M}(\mathcal{U}) \Vdash \forall \bar{a} \in \mathcal{U}\bar{a} \upharpoonright \mu \in (\mathrm{Fin})^{\omega}$.
- (2) $\mathbb{M}(\mathcal{U}) \Vdash \mathcal{U} \upharpoonright \mu$ is an ordered-union filter.

Proof. (1) is an easy density argument. For (2), we use

$$\mathbb{M}(\mathcal{U}) \Vdash (\bar{a} \wedge \bar{b}) \upharpoonright \mu = (\bar{a} \upharpoonright \mu) \wedge (\bar{b} \upharpoonright \mu) \wedge \mathrm{FU}(\bar{a} \upharpoonright \mu) = \mathrm{FU}(\bar{a}) \upharpoonright \mu.$$

Now we restate and prove

Theorem 3.3. Let \mathcal{E} be a \mathbb{P} -point and let \mathcal{U} be a Milliken-Taylor ultrafilter such that $\Phi(\mathcal{U}) \not\leq_{RK} \mathcal{E}$. In $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$ there is a Milliken-Taylor ultrafilter $\mathcal{U}^{\mathrm{ext}} \supseteq \mathcal{U} \upharpoonright \mu$ such that $\Phi(\mathcal{U}^{\mathrm{ext}}) \not\leq_{RK} \mathcal{E}$.

In the remainder of this section we prove this theorem in an iterable form, Theorem 3.9.

Definition 3.4. (See [17, Def. 3.1]) A set $\mathcal{H} \subseteq (\operatorname{Fin})^{\omega}$ is called a *Matet-adequate family* if the following hold:

(i) \mathcal{H} is closed \sqsubseteq^* -upwards.

- (ii) \mathcal{H} is countably closed, i.e., any \sqsubseteq -descending ω -sequence of members of \mathcal{H} has a \sqsubseteq^* -lower bound in \mathcal{H} .
- (iii) \mathcal{H} has the *Hindman property*: If $\bar{a} \in \mathcal{H}$ and $\mathrm{FU}(\bar{a})$ is partitioned into two pieces then there is some $\bar{b} \sqsubseteq \bar{a}$, $\bar{b} \in \mathcal{H}$ such that $\mathrm{FU}(\bar{b})$ is a subset of a single piece of the partition.

Definition 3.5. Let $\langle \bar{a}_n : n < \omega \rangle$ be \sqsubseteq -descending of elements $\bar{a}_n \in (\text{Fin})^{\omega}$. A sequence $\bar{b} \in (\text{Fin})^{\omega}$ is a diagonal lower bound of $\langle \bar{a}_n : n < \omega \rangle$ if

(3.1)
$$(\forall s \in \mathrm{FU}(\bar{b}))((\bar{b} ; \mathrm{past} \; s) \sqsubseteq \bar{a}_{\max(s)+1}).$$

It is equivalent to say $(\forall s \in \bar{b}))((\bar{b} ; \text{past } s) \sqsubseteq \bar{a}_{\max(s)+1})$ in (3.1).

The Hindman property of \mathcal{H} together with the countable closure of \mathcal{H} implies the existence of diagonal lower bounds of sequences in \mathcal{H} , see [17, Cor.1.3], which is based on the deep theorem [7, 4.2]. We give an alternative proof in Lemma 3.6.

Matet-adequate families have better properties than stated in the definition; this is similar to [7, Theorem 4.2].

Lemma 3.6. Let \mathcal{H} be a Matet-adequate family.

- (a) Let $n \geq 1$. If $A \in \mathcal{H}$ and $[FU(\bar{a})]_{<}^n$ is partitioned into two pieces then there is some $\bar{b} \sqsubseteq \bar{a}$, $\bar{b} \in \mathcal{H}$ such that $[FU_k(\bar{b})]_{<}^n$ is a subset of a single piece of the partition.
- (b) H contains for each descending sequence a diagonal lower bound.

Proof. (a) is proved in [7, Theorem 4.2]. We write an alternative proof that also serves to show that ultraness is not used. For simplicity we write the step from 1 to 2. We let $c : [\mathrm{FU}_k(\bar{a})]_<^2 \to r$ for some finite $r \geq 1$. We enumerate $\mathrm{FU}_k(\bar{a})$ as $\langle s_\ell : \ell \in \omega \rangle$ such that for any ℓ , all the s_i with $\max(\mathrm{supp}(s_i)) < \max(\mathrm{supp}(s_\ell))$ have $i < \ell$. Let $\bar{c}_{-1} = \bar{a}$. For each $s_\ell \in \mathrm{FU}_k(\bar{a})$ by induction hypothesis we may take a $\bar{c}_\ell \sqsubseteq_k (\bar{c}_{\ell-1}; \mathrm{past}\ s_\ell)$ such that

$$f_{\ell} \colon \mathrm{FU}_k(\bar{c}_{\ell-1} ; \mathrm{past} \ s_{\ell}) \to m$$

 $t \mapsto c(s_{\ell}, t)$

is monochromatic on $\mathrm{FU}_k(\bar{c}_\ell)$. Let $\bar{b} \sqsubseteq_k^* \bar{c}_\ell$ for any $\ell < \omega$. Now we take $g \colon \omega \to \omega$ with g(0) = 0 and $g(\ell+1) > g(\ell)$ so large that

$$(3.2) (\forall r \leq g(\ell))((\bar{b} ; past \{g(\ell+1)\}) \sqsubseteq_k \bar{c}_r).$$

Next we let for $s \in \mathrm{FU}_k(\bar{b})$, say $s = s_\ell$, f(s) = c(s,t) for any $t \in \mathrm{FU}_k(\bar{c}_\ell) \cap \mathrm{FU}(\bar{b})$. By the induction hypothesis there is $\bar{c} \sqsubseteq_k \bar{b}$, $\bar{c} \in \mathcal{H}$, such that $\mathrm{FU}_k(\bar{c})$ is monochromatic under the colouring f. Then we colour $s \in \mathrm{FU}_k(\bar{c})$ with colour $j \in 2$ if $\max(\sup p(s))$ is in $\bigcup \{ [g(2r+j), g(2r+j+1)) : r \in \omega \}$, and take $\bar{d} \sqsubseteq_k \bar{c}$ such that $\mathrm{FU}_k(\bar{d})$ is monochromatic for the latter colouring. Finally we take $\bar{e} \sqsubseteq_k \bar{d}$ such that for each r there is at most one ℓ such that $\sup (e_\ell) \cap [g(2r+j), g(2r+j+1)) \neq \emptyset$. The existence of such an $\bar{e} \in \mathcal{H}$ follows by [7, Theorem 3.9]. Then $[\mathrm{FU}_k(\bar{e})]_{<}^2$ is c-monochromatic.

(b) Let $\langle \bar{a}_n : n \in \omega \rangle$ be a \sqsubseteq_k -descending sequence of element of \mathcal{H} and let $\bar{c} \in \mathcal{H}$ be a \sqsubseteq_k^* -lower bound such that $\bar{c} \sqsubseteq_k \bar{a}_0$. We colour $\mathrm{FU}_k(\bar{c})]_<^2$ via

$$c(u,v) = \begin{cases} 1 & \text{if } v \in \mathrm{FU}_k(\bar{a}_{\max(\mathrm{supp}(u))+1}); \\ 0 & \text{else.} \end{cases}$$

Since \bar{c} is a \sqsubseteq_k^* -lower bound of $\langle \bar{a}_n : n < \omega \rangle$, any $\bar{d} \sqsubseteq_k \bar{c}$ such that $[\mathrm{FU}_k(\bar{d})]_<^2$ is c-monochromatic has colour 1 and hence is a diagonal lower bound.

Definition 3.7. Let $\mathcal{C} \subseteq (\operatorname{Fin})^{\omega}$ be centred. $\mathcal{C}^+ = \{\bar{a} \in (\operatorname{Fin})^{\omega} : \forall \bar{c} \in \mathcal{C}, \bar{c} \not\perp \bar{a}\}$

We introduce an abbreviation:

Definition 3.8. Let $\mathcal{H} \subseteq (\operatorname{Fin})^{\omega}$ and let \mathcal{E} be a P-point. We say \mathcal{H} avoids \mathcal{E} if $\{\operatorname{set}(\bar{a}) : \bar{a} \in \mathcal{H}\}$ is nnc to \mathcal{E} . For nnc see Definition 1.1(4).

The technical core of the proof of Theorem 3.3 is:

Theorem 3.9. After forcing with $\mathbb{M}(\mathcal{U})$, $(\mathcal{U} \upharpoonright \mu)^+$ is a Matet-adequate family that for any finite-to-one h, for any $\bar{a} \in (\mathcal{U} \upharpoonright \mu)^+$,

$$(\exists \bar{b}^1, \bar{b}^2 \sqsubseteq \bar{a})(\bar{b}^0 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge \bar{b}^1 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge h[\operatorname{set}(\bar{b}^1)] \cap h[(\operatorname{set}(\bar{b}^2)] = \emptyset).$$

By the latter property, $(\mathcal{U} \upharpoonright \mu)^+$ avoids any ultrafilter from the ground model, in particular \mathcal{E} . Once Theorem 3.9 is proved, a routine downwards construction along ω_1 (see e.g., [7, Theorem 2.4]) completes the proof of Theorem 3.3: Under CH in $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$ there is a Milliken–Taylor ultrafilter $\mathcal{U}^{\mathrm{ext}} \supseteq \mathcal{U}$ that avoids \mathcal{E} .

Showing adequacy requires some technical work to evaluate the forcing.

Definition 3.10. $(t, \bar{b}) \leq (s, \bar{a})$ is called a pure extension of (s, \bar{a}) if s = t.

Lemma 3.11. ([17, Lemma 2.6]) $\mathbb{M}(\mathcal{U})$ has the pure decision property, that is, for any φ in the forcing language for any $(s, \bar{a}) \in \mathbb{M}(\mathcal{U})$, there is $\bar{b} \in \mathcal{U}$, $\bar{b} \sqsubseteq \bar{a}$ such that (s, \bar{b}) decides φ .

Eisworth introduced the notion of a neat condition for a name of a subset of ω . We extend the notion of neatness for our purposes. ⁶

Definition 3.12. (1) Let \underline{A} be a name for an infinite subset of Fin (that means, the weakest condition forces this). We say (s, \overline{b}) is neat for \underline{A} if

$$(\forall t \in \mathrm{FU}(\bar{b}))(\forall r \in \mathrm{FU}(\bar{b} ; \mathrm{past} \; t))(\forall u \subseteq \mathrm{max}(r))$$
$$(\forall r' \in \mathrm{FU}(\bar{b} ; \mathrm{past} \; r))\big((s \cup t \cup r', (\bar{b} ; \mathrm{past} \; r'))$$
 decides $u \in A$ and the decision does not depend on r').

(2) Let \underline{h} be a name for a finite-to-one function such that $h(i) \leq i$. We say (s, \overline{b}) is neat for h if

$$(\forall t \in \mathrm{FU}(\bar{b}))(\forall r \in \mathrm{FU}(\bar{b} ; \mathrm{past} \; t))(\forall i \leq \mathrm{max}(r))$$

$$(\forall r' \in \mathrm{FU}(\bar{b} ; \mathrm{past} \; r))\big((s \cup t \cup r', (\bar{b} ; \mathrm{past} \; r'))$$
 decides $\underline{h}(i)$ and the decision does not depend on r').

(3) Let \underline{c} be a name for a function \underline{c} : Fin $\to \{0,1\}$. We say (s, \overline{b}) is neat for \underline{c} if $(\forall t \in \mathrm{FU}(\overline{b}))(\forall r \in \mathrm{FU}(\overline{b}))(\forall u \subseteq \mathrm{max}(r))$

$$(\forall r' \in \mathrm{FU}(\bar{b} ; \mathrm{past} \; r))((s \cup t \cup r', (\bar{b} ; \mathrm{past} \; r')))$$

decides g(u) and the decision does not depend on r').

⁶It can be used for names of subsets of $H(\omega)$ and names for any ω -hierarchy of hereditary finite sets whose union is a subset $H(\omega)$. We define neatness by tailoring initial segments towards computations with diagonal lower bounds.

(4) Let $\langle A_j : j < \omega \rangle$ be a sequence of names for elements of $(\text{Fin})^{\omega}$. We say (s, \bar{b}) is neat for $\langle A_j : j < \omega \rangle$ if

$$(\forall t \in \mathrm{FU}(\bar{b}))(\forall r \in \mathrm{FU}(\bar{b} \; ; \mathrm{past} \; t))(\forall j \leq \mathrm{max}(r))(\forall u \subseteq \mathrm{max}(r)) \\ (\forall r' \in \mathrm{FU}(\bar{b} \; ; \mathrm{past} \; r))\big((s \cup t \cup r', (\bar{b} \; ; \mathrm{past} \; r')) \\ \mathrm{decides} \; u \in A_j \; \mathrm{and} \; \mathrm{the} \; \mathrm{decision} \; \mathrm{does} \; \mathrm{not} \; \mathrm{depend} \; \mathrm{on} \; r'\big).$$

Lemma 3.13. ([17, Lemma 2.7, Lemma 2.8]) $(s, \bar{a}) \in \mathbb{M}(\mathcal{U})$. Let $(X, \underline{h}, \underline{c}, \langle A_j : j < \omega \rangle)$ given such that the weakest condition forces: X is a min-unbounded subset of Fin, \underline{h} is a surjective weakly increasing finite-to-one function, \underline{c} is a name for a colouring, $\langle A_j : j < \omega \rangle$ is a sequence of members of $(\text{Fin})^{\omega}$. Then there is $\overline{b} \sqsubseteq \overline{a}$ such that $(s, \overline{b}) \in M(\mathcal{U})$ is neat for $X, \underline{h}, \underline{c}, \langle A_j : j < \omega \rangle$.

Remark 3.14. Since the proof does not use the fact that \mathcal{U} is a filter, Lemma 3.13 also holds for $\mathbb{M}(\mathcal{H})$ for any Matet-adequate family \mathcal{H} .

Now we prove Theorem 3.9. It is obvious that the set of positive sets $(\mathcal{U} \upharpoonright \mu)^+$ is upwards closed in the \sqsubseteq^* -order. Now we show that any \sqsubseteq -descending ω -sequence has a \sqsubseteq^* -lower bound. It is not harder to directly show that there is a diagonal lower bound.

We recall that $\min_{\langle,\text{lex},\text{Fin}}$ was defined in Definition 2.1(11). A \mathbb{Q} -name is a set of the form $\tau = \{\langle \sigma, q \rangle : \langle \sigma, q \rangle \in \tau\}$ with names σ of lower rank. For $x \in \mathbf{V}$ we have the \mathbb{Q} -name $\check{x} = \{\langle \check{y}, q \rangle : y \in x, q \in \mathbb{Q}\}$. We drop the \check{x} -sign.

The following technique is one of the cornerstones of our forcing constructions and interesting for itself:

Lemma 3.15. (Existence of positive diagonal lower bounds) Let \mathcal{U} be a Milliken-Taylor ultrafilter, \mathcal{E} be a P-point, $\Phi(\mathcal{U}) \not\leq_{RB} \mathcal{E}$. Let $\mathbb{Q} = \mathbb{M}(\mathcal{U})$ and let μ be the name for the generic real. Let $\bar{X} = \langle \bar{X}_n : n \in \omega \rangle$ be a sequence of \mathbb{Q} -names for elements of $(\operatorname{Fin})^{\omega}$ such that

$$\mathbb{Q} \Vdash (\forall n \in \omega) (X_n \in (\mathcal{U} \upharpoonright \mu)^+ \wedge X_{n+1} \sqsubseteq X_n).$$

Then

fulfils

$$(3.4) \qquad \mathbb{Q} \Vdash \underline{\mathcal{D}} \in (\mathcal{U} \upharpoonright \mu)^+ \wedge \underline{\mathcal{D}} \sqsubseteq \underline{X}_0 \wedge (\forall t \in \underline{\mathcal{D}})((\underline{\mathcal{D}} ; \mathrm{past} \; t) \sqsubseteq X_{\max(t)+1}).$$

Proof. Let (s, \bar{a}) be given such that (s, \bar{a}) is neat for \bar{D} and for $\langle X_i \upharpoonright \mu : i < \omega \rangle$. We show that there is $(s, \bar{b}) \leq_{\mathbb{Q}} (s, \bar{a})$ that forces that $\bar{D} \upharpoonright \mu \in (\mathcal{U} \upharpoonright \mu)^+$.

Let $k \in \omega$ and $u \in \mathrm{FU}(\bar{a})$. We say that u is good for (s, \bar{a}) , $\bar{t} = (t_0, \ldots, t_{k-1}) \in [\mathrm{Fin}]_{<}^k$, \bar{X} , if

$$(s \cup u, \bar{a}; \operatorname{past} u) \Vdash t_0 = \min_{\leq_{\operatorname{lex}, \operatorname{Fin}}} (X_0 \upharpoonright \mu) \land \bigwedge_{i < k-1} (t_{i+1} = \min_{\leq_{\operatorname{lex}, \operatorname{Fin}}} ((X_{\max(t_i)+1} \upharpoonright \mu); \operatorname{past} t_i)).$$

Note that goodness requires $t_i \in \text{Fin}$ and not just names for elements of Fin. We define a colouring of $[\text{FU}(\bar{a})]^2_{<}$ as follows

$$(3.5) \qquad F(u < v) = \begin{cases} 1 & \text{if for any } \bar{t} \text{ such that} \\ u \text{ is good for } (s, \bar{a}), \bar{t}, \bar{X}, \\ & \text{there is a proper end extension } \bar{t'} \text{ of } \bar{t} \\ & \text{such that } u \cup v \text{ is good for } (s, \bar{a}), \bar{t'}, \bar{X}; \\ 0 & \text{else.} \end{cases}$$

By Taylor's theorem 2.9 there is $(s, \bar{b}) \leq_{\mathbb{Q}} (s, \bar{a})$ such that F is monochromatic on $[\operatorname{FU}(\bar{b})]^2_{<}$. We argue that the monochromatic colour can only be 1: It suffices to find $u < v \in \operatorname{FU}(\bar{b})$ such that F(u, v) = 1. Suppose that $u \in \operatorname{FU}(\bar{b})$ is good for $(s, \bar{a}), \bar{t} = (t_0, \ldots, t_{k-1}), \bar{X}$. If k = 0, we let $X_{t_{k-1}+1} = X_0$. Then $(s \cup u, \bar{b}) \Vdash (X_{\max(t_{k-1})+1} \upharpoonright \mu)$; past $t_{k-1}) \in (\mathcal{U} \upharpoonright \mu)^+$. So $(s \cup u, \bar{b}) \Vdash \exists t \in (X_{\max(t_{k-1})+1} \upharpoonright \mu)$; past t_{k-1}).

Now we show

$$(3.6) (s, \bar{b}) \Vdash \bar{\mathcal{D}} \in (\mathcal{U} \upharpoonright \mu)^+ \land \bar{\mathcal{D}} \sqsubseteq X_0 \land (\forall t \in \mathrm{FU}(\bar{\mathcal{D}}))((\bar{\mathcal{D}}; \mathrm{past}\ t) \sqsubseteq X_{\max(t)+1}).$$

The proof comes in four parts. First we show that \tilde{D} is a min-unbounded subset of Fin. Given any $k \in \omega$, the procedure above for finding v, t' is iterated k times, starting at stage k = 0 with u = s. Thus we find v_i and t_i such that

$$(s \cup v_0, \bar{b} ; \operatorname{past} v_0) \Vdash t_0 = \min_{<_{\operatorname{lex}, \operatorname{Fin}}} (X_0 \upharpoonright \mu),$$

$$(s \cup v_0 \cup v_1, \bar{b} ; \operatorname{past} v_1) \Vdash t_1 = \min_{<_{\operatorname{lex}, \operatorname{Fin}}} (X_{\max(t_0)+1} \upharpoonright \mu),$$

$$\vdots$$

$$(s \cup v_0 \cup \dots \cup v_k, \bar{b} ; \operatorname{past} v_k) \Vdash t_k = \min_{<_{\operatorname{lex}, \operatorname{Fin}}} (X_{\max(t_{k-1})+1} \upharpoonright \mu), \text{ and thus }$$

$$(s \cup v_1 \cup \dots \cup v_k, \bar{b} ; \operatorname{past} v_k) \Vdash t_0 < \dots < t_k \in \bar{\mathcal{D}}.$$

Next we show $(s, \bar{b}) \Vdash D \in (\mathcal{U} \upharpoonright \mu)^+$. Suppose for a contradiction, that $\bar{c} \in \mathcal{U}$ and $(s', \bar{b}') \leq (s, \bar{b})$ and $(s', \bar{b}') \Vdash \mathrm{FU}(\bar{c}) \cap \mathrm{FU}(D) = \emptyset$. Then we take $\bar{d} \sqsubseteq \bar{b}', (\bar{c} ; \mathrm{past} \; s'),$ such that $\bar{d} \in \mathcal{U}$ and see that $(s', \bar{d}) \Vdash (D; \mathrm{past} \; s') \subseteq \mathrm{FU}(\bar{c} \upharpoonright \mu)$. Contradiction.

Since
$$\mathbb{Q} \Vdash X_{n+1} \sqsubseteq X_n$$
, by definition of D , $(s, \bar{b}) \Vdash D \subseteq X_0$.

For the last conjunctive clause in Equation (3.6), we work with the characterisation of diagonal lower bound that is given immediately after Def. 3.5. We suppose that $(s \cup v, \bar{b} ; \text{past } v) \Vdash t < t' \in \bar{\mathcal{D}}$. Then by the definition of $\bar{\mathcal{D}}(s \cup v, \bar{b} ; \text{past } v) \Vdash t' \in \mathrm{FU}(\bar{\mathcal{X}}_{\max(t)+1} \upharpoonright \mu; ; \text{past } t)$.

Recall set
$$(\langle a_n : n < \omega \rangle) = \bigcup \{a_n : n < \omega \}.$$

Lemma 3.16. Let $(s, \bar{a}) \Vdash_{\mathbb{M}(\mathcal{U})} \bar{c} \in (\mathcal{U} \upharpoonright \mu)^+ \land h$ is finite-to-one, onto and monotone. Then there are $E \in \mathcal{E}$ and $(s, \bar{b}) \leq (s, \bar{a})$ and \bar{d}^j , j = 0, 1, such that

$$(3.7) \qquad (s,\bar{b}) \Vdash \bigwedge_{j=0,1} (\bar{d}^{j} \in (\mathcal{U} \upharpoonright \mu)^{+} \wedge \bar{d}^{j} \sqsubseteq \bar{c}) \wedge \\ \underline{h}[\operatorname{set}(\bar{d}^{0})] \cap \underline{h}[\operatorname{set}(\bar{d}^{1}] = \emptyset \wedge \bigvee_{j=0,1} \underline{h}[E] \cap \underline{h}[\operatorname{set}(\bar{d}^{j}) = \emptyset.$$

Proof. We assume w.l.o.g. that (s, \bar{a}) is neat for \bar{c} and h. Every name for a positive set \bar{c} and every condition p let us define names \bar{d}^j , j=0,1, such that densely many q below p force $p[\text{set}(\bar{d}^0)] \cap p[\text{set}(\bar{d}^1)] = \emptyset \wedge \bigwedge_{j=0,1} (\bar{d}^j \sqsubseteq \bar{c})$. Again this is proved with a colouring.

Let $k < \omega$ and $u \in FU(\bar{a})$. We say that u is good for (s, \bar{a}) , $\bar{d} = (d_0, \ldots, d_{k-1})$, \bar{c} , and h if

$$(s \cup u, \bar{a} ; \operatorname{past} u) \Vdash d_0 = \min_{<_{\operatorname{lex}, \operatorname{Fin}}} (\bar{c} \upharpoonright \mu) \land \\ \bigwedge_{i < k - 1} (d_{i + 1} = \min_{<_{\operatorname{lex}, \operatorname{Fin}}} \{ d \in \operatorname{FU}(\bar{c} \upharpoonright \mu) : \underline{h}[d_i] \cap \underline{h}[d] = \emptyset \}).$$

Again the d_i are in the ground model, not names. This, though, is not important, since we do not use them as indices. We define a colouring of $[FU(\bar{a})]^2$ as follows

$$(3.8) \qquad F(u < v) = \begin{cases} 1 & \text{if for any \bar{d} such that} \\ & u \text{ is good for } (s, \bar{a}), \bar{d}, \bar{c}, \bar{h} \\ & \text{there is a proper end extension \bar{d}' of \bar{d}} \\ & \text{such that } u \cup v \text{ is good for } (s, \bar{a}), \bar{d}', \bar{c}, \bar{h}; \\ 0 & \text{else.} \end{cases}$$

By Taylor's theorem 2.9 there is $(s, \bar{b}) \leq_{\mathbb{M}(\mathcal{U})} (s, \bar{a})$ such that F is monochromatic on $[\mathrm{FU}(\bar{b})]^2_{\leq}$. Since for any k, $((\bar{c} \upharpoonright \mu) ; \mathrm{past} \ k)$ is forced to be in $(\mathcal{U} \upharpoonright \mu)^+$, the monochromatic colour can only be 1. Now there are names d_i such that

$$(s, \bar{b}) \Vdash \langle d_{2\underline{k}+1} : k \in \omega \rangle \in (\mathcal{U} \upharpoonright \mu)^{+} \wedge$$

$$\langle d_{2\underline{k}} : k \in \omega \rangle \in (\mathcal{U} \upharpoonright \mu)^{+} \wedge$$

$$\langle d_{2} : k \in \omega \rangle \sqsubseteq \bar{c} \wedge$$

$$h[\bigcup \{d_{2\underline{k}} : k \in \omega\}] \cap h[\bigcup \{d_{2\underline{k}+1} : k \in \omega\}] = \emptyset.$$

The first two conjunctive clauses are shown as in the proof of Equation (3.6). The last conjunctive clause follows from the new definition of goodness. We let $(s, \bar{b}) \Vdash \bar{d}^j = \langle d_{2k+j} : k < \omega \rangle$ for j = 0, 1. Since $\mathbb{M}(\mathcal{U}) \Vdash h(\mathcal{E})$ is an ultrafilter, (s, \bar{b}) forces there are a j = 0, 1 and an $E \in \mathcal{E}$ such that $h[E] \cap h[\operatorname{set}(\bar{d}^j)] = \emptyset$.

The next lemma is the most important step in the proof of Theorem 3.9. Indeed, it includes again a proof that positive diagonal lower bounds exist.

Lemma 3.17. In $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$, $(\mathcal{U} \upharpoonright \mu)^+$ has the Hindman property.

For the proof of this lemma, we adapt a theorem of Eisworth.

Theorem 3.18. [17, Theorem 5] Let \mathcal{F} be an ordered-union filter generated by $\langle \cos(\mathcal{B}) \rangle$ sets and let c be a partition of Fin into finite sets. Then there is an $\bar{a} \in \mathcal{F}^+$ such that $\mathrm{FU}(\bar{a})$ is included in one piece of the partition.

At a crucial point in Eisworth's proof a Cohen real over an elementary submodel provides a name in a Galvin–Glazer framework. We show that also a Matet-real and even an $\mathbb{M}(\mathcal{U})$ -generic real can be used. We recall the Galvin–Glazer [25] technique.

Definition 3.19. We denote by $\gamma(\text{Fin})$ the set of min-unbounded (see Def. 2.6(3)) ultrafilters over Fin. This set is endowed with the topology generated by

$$\{\{\mathcal{U} \in \gamma(\operatorname{Fin}) : A \in \mathcal{U}\} : A \subseteq \operatorname{Fin}\}.$$

The space $\gamma(\text{Fin})$ is a compact zero-dimensional Hausdorff space. With the named topology and the semigroup operation \cup from Def. 2.1(9), the semigroup $(\gamma(\text{Fin}), \cup)$ is a right-topological semigroup. Details can be found in [26].

Lemma 3.20. (Ellis, [18]) Each compact subsemigroup of $(\gamma(\text{Fin}), \cup)$ contains an idempotent ultrafilter.

Now we apply Ellis' lemma to $\{\mathcal{U} \in \gamma(\mathrm{Fin}) : \mathcal{U} \supseteq \mathcal{F}\}$ for a min-unbounded filter \mathcal{F} .

Lemma 3.21. ([17, Prop. 4.2]) Let \mathcal{F} be a min-unbounded filter. There is an idempotent ultrafilter $\mathcal{U} \in \gamma(\text{Fin})$ that extends \mathcal{F} .

Now we prove Lemma 3.17. Let p force that \underline{c} is a partition of $\underline{b}_0 \in (\mathcal{U} \upharpoonright \mu)^+$ into finitely many pieces and $\langle \underline{b}_n : n < \omega \rangle$ is a \sqsubseteq -descending sequence of elements $\overline{b}_n \in (\mathcal{U} \upharpoonright \mu)^+$. By Lemma 3.21 there is an $\mathbb{M}(\mathcal{U})$ -name \mathcal{U}^i such that

$$\mathbb{M}(\mathcal{U}) \Vdash \mathcal{U}^i \supseteq \mathrm{fil}((\mathcal{U} \upharpoonright \mu) \cup \{\{b_{n,m} : m \in \omega\} : n \in \omega\}) \wedge \mathcal{U}^i \cup \mathcal{U}^i = \mathcal{U}^i.$$

For $X \subseteq \text{Fin and } t \in \text{Fin we set}$

$$X\ominus t=\{s\,:\,s\cup t\in X\}.$$

Since \mathcal{U}^i is forced to be idempotent,

$$\mathbb{M}(\mathcal{U}) \Vdash (\forall X \in \mathcal{U}^i)(\{t : X \ominus t \in \mathcal{U}^i\} \in \mathcal{U}^i).$$

Now we use again the Milliken–Taylor trick. We assume that (s, \bar{a}) is neat for \underline{c} , $\langle b_n : n < \omega \rangle$.

Let $n \geq 1$. We call $u \in \mathrm{FU}(\bar{a})$ good for $(s,\bar{a}), (X_m,d_m:m < n)$ if $(s \cup u,\bar{a};\mathrm{past}\,u)$ forces the following statements:

- (1) X_0 is the piece of the partition c of $\mathrm{FU}(\bar{b})$ that is in \mathcal{U}^i .
- (2) We let $d_{-1} = \{-1\}$. For any $0 \le m < n$ d_m is the $\le_{\text{lex},\text{Fin}}$ -least element of

$$(3.10) \qquad \{d \in X_m \cap \mathrm{FU}(\{a_k : k \in \omega\} \upharpoonright \mu) \cap \mathrm{FU}(\bar{b}_{\max(d_{m-1})+1}) : X_m \ominus d \in \mathcal{U}^i \text{ and } \min(d) > \max(d_{m-1})\}$$

(3) For any $0 \le m < n-1$, $X_{m+1} = X_m \cap (X_m \ominus d_m)$.

Here we allow names. Only the natural numbers are meant to be pinned down. We colour $[FU(\bar{a}; past s)]^2$ as follows:

$$(3.11) F(u < v) = \begin{cases} 1 & \text{if for any } (X_m, d_m : m < n) \text{ such that} \\ u \text{ is good for } (s, \bar{a}), (X_m, d_m : m < n), \\ \text{there is a proper end extension } (X_m, d_m : m < n') \\ \text{such that } v \text{ is good for } (s, \bar{a}), (X_m, d_m : m < n'); \\ 0 & \text{else.} \end{cases}$$

Then we find a monochromatic $\bar{b} \in \mathcal{U}$ with $(s, \bar{b}) \leq (s, \bar{a})$. Since \mathcal{U}^i is idempotent and $\bar{b} \in \mathcal{U} \subseteq \mathcal{U}^i$, the set in (3.10) is in \mathcal{U}^i . Hence the monochromatic colour can only be 1. We let \bar{e} be a name such that $(s, \bar{b}) \Vdash (\forall n) \langle e_0, \dots, e_{n-1} \rangle = \langle d_0, \dots, d_{n-1} \rangle$.

The monochromaticity statement

$$\mathbb{M}(\mathcal{U}) \Vdash \mathrm{FU}(\bar{e}) \subseteq X_0$$

is proved literally as in Eisworth [17, page 460]. By item (2) in the current definition of "good", the sequence \bar{e} is a diagonal lower bound of $\langle \bar{b}_n : n < \omega \rangle$. Now we show that \bar{e} is positive. For this we use the conditions on $\mathrm{FU}(\bar{a})$ in the goodness clause (2). Suppose for a contradiction that \bar{e} is not forced to be $(\mathcal{U} \upharpoonright \mu)$ -positive. Hence there is $q \in \mathbb{M}(\mathcal{U})$, $\bar{c} \in \mathcal{U}$ such that q is neat for \bar{e} and μ , $q \leq (s, \bar{b})$, and

$$q \Vdash_{\mathbb{M}(\mathcal{U})} \mathrm{FU}(\bar{e}) \cap \mathrm{FU}(\bar{c} \upharpoonright \mu) = \emptyset.$$

Since \mathcal{U} is a filter, we can assume $q=(t,\bar{c})$. We produce an extension r of q that forces the contrary.

There is a minimal m such that q does not determine e_m . So q determines $e_0 = d_0, \ldots, e_{m-1} = d_{m-1}$.

Since $[FU(\bar{c}; past t)]^2$ has colour 1,

$$(t,\bar{c}) \Vdash Y = \{d \in X_m \cap \mathrm{FU}(\{c_k : k \in \omega\} \upharpoonright \mu) \cap \mathrm{FU}(\bar{b}_{\max(d_{m-1})+1}) : X_m \ominus d \in \mathcal{U}^i \text{ and } \min(d) > \max(d_{m-1})\} \in \mathcal{U}^i.$$

Since $[\mathrm{FU}(\bar{c})]^2_{<}$ has colour 1 and since q is neat for \bar{e} and μ , there is $r \leq q$ of the form $(t \cup u, \bar{c}; \mathrm{past}\; u)$ and there is $d \in \mathrm{Fin}$

$$r \Vdash d \in \mathrm{FU}(\bar{e}) \cap \mathrm{FU}(\bar{c} \upharpoonright \mu),$$

in contradiction to the assumption on q.

 $\square_{3.17,3.9,3.3}$

Henceforth we drop the tildes underneath the names.

Now we return to filters over ω and answer some instances of the question left open in the previous section: What happens to filters with $\Phi(\mathcal{U}) \leq_{RB} \mathcal{F}$?

Mathias introduced the following notion under the name "happy family" [30, Def. 0.1.]. Louveau studied it in the special case of ultrafilters [28]. Todorcevic [40, Chapter 7] uses the name "selective coideal" for a happy family.

Definition 3.22. (See [30, Def. 0.1.], [40, Def. 7.3]) A set $\mathcal{H} \subseteq [\omega]^{\omega}$ is called a *selective coideal/happy family* if the following hold:

- (i) $\mathcal{I}_{\mathcal{H}} := \mathcal{P}(\omega) \setminus \mathcal{H}$ is an ideal that contains all singletons.
- (ii) If $\langle A_i : i \in \omega \rangle$ is a \subseteq -descending sequence of elements $A_i \in \mathcal{H}$, then there is $B \in \mathcal{H}$ such that $B \subseteq A_0$ and $(\forall i \in B)B \setminus (i+1) \subseteq A_i$. We call such a B a diagonal lower bound of $\langle A_i : i \in \omega \rangle$.

We write $\mathcal{F}_{\mathcal{H}} = \{ \omega \setminus X : X \in \mathcal{I}_{\mathcal{H}} \}$ for the filter that is dual to $\mathcal{I}_{\mathcal{H}}$. Then \mathcal{H} coincides with the $\mathcal{F}_{\mathcal{H}}$ -positive sets, i.e.,

$$\mathcal{H} = \mathcal{F}_{\mathcal{H}}^+ := \{ X \in [\omega]^\omega : (\forall Y \in \mathcal{F}_{\mathcal{H}})(X \cap Y \neq \emptyset) \}.$$

Lemma 3.23. Let \mathcal{F} be a filter over ω and let \mathcal{R} be an ultrafilter over ω . $\mathcal{R} \subseteq \mathcal{F}^+$ iff $\mathcal{R} \supset \mathcal{F}$.

So the forward implication, which will be invoked many times, uses that \mathcal{R} is ultra.

Remark 3.24. Let $\mathcal{H} \subseteq [\omega]^{\omega}$ and let \mathcal{E} be a filter. \mathcal{H} and \mathcal{E} are nnc iff for any finite-to-one function f and $X \in \mathcal{H}$ there are a $E \in \mathcal{E}$ and a $Y \subseteq X$, such that $Y \in \mathcal{H}$ and such that $f[E] \cap f[H] = \emptyset$.

The following theorem provides information on $\min(\mathcal{U})$ and $\max(\mathcal{U})$.

Theorem 3.25. Assume CH and that we force with $\mathbb{M}(\mathcal{U})$ for a Milliken-Taylor ultrafilter \mathcal{U} , $\mathcal{R} \in \{\min(\mathcal{U}), \max(\mathcal{U})\}$ and μ is the generic real. After forcing with $\mathbb{M}(\mathcal{U})$, the set of positive sets

$$(\operatorname{fil}(\mathcal{R} \cup \{\mu\}))^+ = \{X \in ([\omega]^\omega)^{\mathbf{V}^{\mathbb{M}(u)}} : (\forall Y \in \mathcal{R})|X \cap Y \cap \mu| = \omega\}$$

is a happy family that is nowhere almost a filter in $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$ (and hence it is nnc to \mathcal{E}) and by [30, Prop. 011] is a Ramsey ultrafilter $\mathcal{R}^{\mathrm{ext}} \supseteq \mathcal{R} \cup \{\mu\}$ that is nnc to \mathcal{E} ,

Proof. The theorem is proved like Theorem 3.3, however, it is much easier. Lemma 3.15, giving diagonal lower bounds, and Lemma 3.16, showing the nnc-part, are adapted to names for $(\mathcal{R} \cup \{\mu\})$ -positive subsets of ω . There are no new ideas. \square

Remark 3.26. We remark that by an analogous proof to Mathias' [30, Prop. 011], under CH any happy family that is nnc to \mathcal{E} contains a Ramsey ultrafilter as a subset that is nnc to \mathcal{E} . So we see that instead of heading for a Milliken–Taylor ultrafilter $\mathcal{U}^{\text{ext}} \supseteq \mathcal{U} \upharpoonright \mu$ that yields of course $\min(\mathcal{U}^{\text{ext}}) \supseteq \min(\mathcal{U}) \cup \{\mu\}$ and the same for the maximum projection, we could proceed into a different direction: By Theorem 3.25, we can extend the minimum and maximum projections $\min(\mathcal{U})$, $\max(\mathcal{U})$ to new Ramsey ultrafilters in $\mathbf{V}^{\mathbb{M}(\mathcal{U})}$ and not care whether these extensions are the minimum and the maximum of a Milliken–Taylor ultrafilter extending \mathcal{U} . This direction will be important in Section 6.

4. A NAME FOR A MATET-ADEQUATE FAMILY AT LIMIT STAGES

We define by induction on $\alpha \leq \omega_2$ a countable support iteration $\mathbb{P}_{\alpha} = \langle \mathbb{P}_{\beta}, \mathbb{M}(\mathcal{U}_{\gamma}) : \beta \leq \alpha, \gamma < \alpha \rangle$ such that for any $\gamma < \alpha$,

$$\mathbb{P}_{\gamma+1} = \mathbb{P}_{\gamma} * \mathbb{M}(\mathcal{U}_{\gamma}) \text{ and}$$

$$\mathbb{P}_{\gamma+1} \Vdash \mu_{\gamma} = \bigcup \{s : (s, \bar{a}) \in G_{\mathbb{M}(\mathcal{U}_{\gamma})}\}$$

$$\mathbb{P}_{\gamma} \Vdash \mathcal{U}_{\gamma} \supseteq (\bigcup \{\mathcal{U}_{\delta} \upharpoonright \mu_{\delta}) : \delta < \gamma)$$
is a Milliken–Taylor ultrafilter that avoids \mathcal{E} .

In Theorem 3.3 we proved that there are extension of Milliken–Taylor ultrafilters in the successor steps

$$\mathbb{P}_{\gamma} \Vdash_{\mathbb{P}_{\gamma}} \Big(\mathbb{M}(\mathcal{U}_{\gamma}) \Vdash_{\mathbb{M}(\mathcal{U}_{\gamma})} \mu_{\gamma} = \bigcup \{ s : (s, \bar{a}) \in G_{\mathbb{M}(\mathcal{U}_{\gamma})} \}$$

$$\wedge \big(\exists \mathcal{U}_{\gamma+1} \supseteq \mathcal{U}_{\gamma} \big)$$

$$\big(\mathcal{U}_{\gamma+1} \text{ is a Milliken-Taylor ultrafilter and } \mathcal{U}_{\gamma+1} \supseteq \mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma} \big) \Big).$$

This guarantees the continuation of our construction in the successor steps, via $\mathcal{U}_{\gamma+1} \supseteq \mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma}$.

Now we consider limit steps α . If $\operatorname{cf}(\alpha) > \omega$, we can just take $\mathbb{P}_{\alpha} \vdash \mathcal{U}_{\alpha} = \bigcup_{\gamma < \alpha} \mathcal{U}_{\gamma}$ and the inductive hypotheses will be carried on, since in proper forcing every real appears at a step of at most countable cofinality, with the only exception that for $\alpha = \aleph_2$ the CH gets lost. So we concentrate on the hard case, $\operatorname{cf}(\alpha) = \omega$.

Theorem 4.1. Suppose that \mathbb{P}_{β} , \mathcal{U}_{β} are as in Equation (4.1), $\alpha < \omega$, $\operatorname{cf}(\alpha) = \omega$, \mathbb{P}_{α} is the countable support limit of $\langle \mathbb{P}_{\beta}, \mathbb{M}(\mathcal{U}_{\beta}) : \beta < \alpha \rangle$. In $\mathbf{V}^{\mathbb{P}_{\alpha}}$, the filter \mathcal{E} still generates a P-point and the set of positive sets

$$\left(\bigcup_{\gamma<\alpha}(\mathcal{U}_{\gamma}\restriction\mu_{\gamma})\right)^{+}$$

forms a Matet-adequate family such that for any $\bar{a} \in (\mathcal{U} \upharpoonright \mu)^+$,

$$(\exists \bar{b}^1, \bar{b}^2 \sqsubseteq \bar{a})(\bar{b}^0 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge \bar{b}^1 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge h[\operatorname{set}(\bar{b}^1)] \cap h[(\operatorname{set}(\bar{b}^2)] = \emptyset).$$

As in Theorem 3.9, the latter implies avoidance of \mathcal{E} . Again CH and a routine enumeration along ω_1 gives the following.

Corollary 4.2. Suppose that \mathbb{P}_{β} , \mathcal{U}_{β} are as in Equation (4.1) and \mathbb{P}_{α} is the countable support limit of $\langle \mathbb{P}_{\beta}, \mathbb{M}(\mathcal{U}_{\beta}) : \beta < \alpha \rangle$ and that CH holds in $\mathbf{V}^{\mathbb{P}_{\alpha}}$. Then

 $\mathbb{P}_{\alpha} \Vdash \exists \mathcal{U}_{\alpha}(\mathcal{U}_{\alpha} \text{ is a Milliken-Taylor ultrafilter that avoids } \mathcal{E}, \text{ and}$

$$\mathcal{U}_{\alpha} \supseteq \bigcup_{\gamma < \alpha} (\mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma}) \big).$$

Now we prove Theorem 4.1. Blass and Shelah [11, Theorem 4.1] showed that in $\mathbf{V}^{\mathbb{P}_{\alpha}}$ the closure of \mathcal{E} under almost supersets is a P-point.

For the new part, by induction we define an increasing sequence $\bar{R} = \langle R_{\gamma} : \gamma < \alpha \rangle$ of relations R_{γ} in $\mathbf{V}^{\mathbb{P}_{\gamma}}$ such that a property called

(4.2) "
$$\mathbb{P}_{\gamma}$$
 is R_{γ} -preserving"

is a notion we want to carry from $\gamma < \alpha$, $cf(\gamma) < \omega_1$, to α in addition to the property (4.1) and properness in the inductive choice of the iteration.

Now we define the relation R_{α} for which we want to preserve statements of the form $(\forall f)(\exists \bar{g})(fR_{\alpha}\bar{g})$. The relation R_{α} will be a Borel relation on the Baire space in $\mathbf{V}^{\mathbb{P}_{\alpha}}$ that contain complex parameters from the ground model, e.g. Milliken–Taylor ultrafilters and names for Milliken–Taylor ultrafilters.

Definition 4.3. By induction on $\alpha \leq \omega_2$ we define the following relations.

- (1) We say that a \mathbb{P}_{α} -name \bar{a} for an element of $(\operatorname{Fin})^{\omega}$ is α -positive if $1 \Vdash_{\mathbb{P}_{\alpha}} \bar{a} \in (\bigcup \{\mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma} : \gamma < \alpha\})^{+}$.
- (2) Assume that $\langle \mathcal{U}_{\gamma} : \gamma < \alpha \rangle$ is an ascending sequence of Milliken–Taylor ultrafilters $\mathcal{U}_{\gamma} \in \mathbf{V}^{\mathbb{P}_{\gamma}}$, such that $\mathbb{P}_{\gamma} \Vdash \Phi(\mathcal{U}_{\gamma}) \not\leq_{\mathrm{RB}} \mathcal{E}$ and $\forall \gamma < \delta < \alpha$, $\mathbb{P}_{\delta} \Vdash \mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma} \subseteq \mathcal{U}_{\delta}$. We say $fR_{\alpha}\bar{g}$ if the following holds in $\mathbf{V}^{\mathbb{P}_{\alpha}}$:
 - (a) $f = (\bar{A}, h, c),$
 - (b) $\bar{A} = \langle A_{\ell} : \ell \in \omega \rangle$ is a \sqsubseteq -descending sequence of α -positive sequences $A_{\ell} \in (\operatorname{Fin})^{\omega}$,
 - (c) h is finite-to-one,
 - (d) c is a partition of $FU(A_0)$.
 - (e) For j = 0, 1 we let $\bar{g}^j := \langle g_{2n+j} : n \in \omega \rangle$. Then
 - (i) For $j = 0, 1, \bar{q}^j$ is an α -positive diagonal lower bound of \bar{A} .
 - (ii) For j = 0, 1, $FU(\bar{g}^j)$ is in one piece of the partition c.
 - (iii) $h[\operatorname{set}(\bar{g}^0)] \cap h[\operatorname{set}(\bar{g}^1)] = \emptyset.$

So R_{α} is a \mathbb{P}_{α} -name for a relation.

Definition 4.4. We say \mathbb{P}_{α} is R_{α} -preserving if \mathbb{P}_{α} is proper and

$$\mathbb{P}_{\alpha} \Vdash \forall f \in \text{dom}(R_{\alpha}) \exists \bar{g}(fR_{\alpha}\bar{g}).$$

There are two main differences to the known "Case A" of iteration theorem [37, Ch XVIII], [21]: For our R_{α} , it is not the case that for countably many tasks f_n , $n \in \omega$ there is one answer \bar{g} , E such that $\forall n f R_{\alpha,n}(\bar{g},E)$ where $R_{\alpha,n}$ is R_{α} up to mistakes before n. There are \sqsubseteq^* -incompatible positive elements \bar{g} . Secondly, not only the quests f but also the answers \bar{g} are now from the forcing extension. This differs from the traditional applications in the preservation of cardinal invariants, see e.g. [2]. We do not write tildes below the R_{α} 's.

Lemma 4.5. Assume CH, $\alpha < \omega_2$, $cf(\alpha) < \omega_1$. If \mathbb{P}_{α} is R_{α} -preserving then

$$\mathbb{P}_{\alpha} \Vdash (\bigcup \{ \mathcal{U}_{\beta} \upharpoonright \mu_{\beta} : \beta < \alpha \})^{+} \text{ is a Matet-adequate family such that}$$

$$(\forall \bar{a} \in (\mathcal{U} \upharpoonright \mu)^{+})(\exists \bar{b}^{1}, \bar{b}^{2} \sqsubseteq \bar{a})$$

$$(\bar{b}^{0} \in (\mathcal{U} \upharpoonright \mu)^{+} \wedge \bar{b}^{1} \in (\mathcal{U} \upharpoonright \mu)^{+} \wedge h[\operatorname{set}(\bar{b}^{1})] \cap h[(\operatorname{set}(\bar{b}^{2})] = \emptyset).$$

Proof. This follows from Definition 4.4.

Now we carry the preservation property upwards by induction.

Lemma 4.6. Assume CH, $\alpha < \omega_2$, and $\operatorname{cf}(\alpha) = \omega$. Let \mathbb{P}_{α} be the countable support limit of \mathbb{P}_{β} , $\beta < \alpha$. If for $\beta < \alpha$ such that $\operatorname{cf}(\beta) < \omega_1$, \mathbb{P}_{β} is R_{β} -preserving and for any $\beta < \alpha$ Equation (4.1) holds then \mathbb{P}_{α} is R_{α} -preserving.

This lemma will proved with Lemma 4.8. For definiteness, we can take $\chi = (2^{|\mathbb{P}_{\alpha}|})^+$. Under CH, for $\alpha < \omega_2$, $|\mathbb{P}_{\alpha}| \leq \aleph_1$ by [36, page 96]. So $\chi = (2^{\aleph_1})^+$ is sufficiently large. The following lemma on the translation to countable elementary submodels is well-known, see [37, Theorem 2.11 and Ch. XVIII].

Lemma 4.7. The following are equivalent.

- (1) \mathbb{P}_{α} is R_{α} -preserving.
- (2) For all $M \prec H(\chi)$ such that \mathbb{P}_{α} , $\langle \mathcal{U}_{\beta} : \beta < \alpha \rangle$, $\mathcal{E} \in M$ for all \mathbb{P}_{α} -names $f \in M$ for all $p \in \mathbb{P}_{\alpha} \cap M$, if $p \Vdash f \in \text{dom}(R_{\alpha})$ then there is an (M, \mathbb{P}_{α}) -generic $q \geq p$ and there is a \mathbb{P}_{α} -name $\bar{g} \in M$ such that $q \Vdash fR_{\alpha}\bar{g}$.

The property of Lemma 4.7(2) is carried on by induction in the following slightly stronger technical form that is suitable for induction.

Lemma 4.8. The induction lemma. Suppose CH, $\xi < \zeta \leq \aleph_2$, and $\operatorname{cf}(\zeta) < \omega_1$. Let $M \prec H(\chi)$, $\zeta \in M$, $\mathbb{P}_{\zeta} \in M$, $p \in \mathbb{P}_{\zeta} \cap M$ and $q_0 \leq p \upharpoonright \xi$ be (M, \mathbb{P}_{ξ}) -generic. Let $f \in M$ be a \mathbb{P}_{ζ} -name such that

$$p \Vdash_{\mathbb{P}_{\zeta}} f = (\langle A_n : n \in \omega \rangle, c, h) \in \text{dom}(R_{\zeta}).$$

Then there is some (M, \mathbb{P}_{ζ}) -generic condition $q \leq p \upharpoonright [\xi, \eta) \cup q_0$ such that $q \Vdash (\exists \bar{g})(fR_{\zeta}\bar{g})$.

Proof. Moreover we get $dom(q_0) \setminus \xi \subseteq \zeta \cap M$. We go by induction on ζ . For $\zeta = 0$ there is nothing to prove, for ζ successor a proof is included in the proof of Theorem 3.3, namely in Lemmata 3.15, 3.16, 3.17. So let ζ be a limit. We fix a strictly increasing sequence $\langle \zeta_{\ell} : \ell < \omega \rangle$ with $\zeta_0 = \xi$, $\zeta_{\ell} \in M$, and $\sup \zeta_{\ell} = \zeta$. We

also fix an enumeration $\langle D_n : n < \omega \rangle$ of the dense subsets of \mathbb{P}_{ζ} that are elements of M. We let \mathcal{U}^i be a \mathbb{P}_{ζ} -name in M such that

(4.3)
$$\mathbb{P}_{\zeta} \Vdash \mathcal{U}^{i} \text{ is idempotent and}$$

$$\mathcal{U}^{i} \supseteq \text{fil}(\bigcup \{(\mathcal{U}_{\gamma} \upharpoonright \mu_{\gamma}) : \gamma < \zeta\} \cup \{A_{n} : n \in \omega\}).$$

By 3.21 such a name \mathcal{U}^i exists. We choose by induction on $n, q_n \in \mathbb{P}_{\zeta_n}$, a \mathbb{P}_{ζ_n} -name for a $\mathbb{P}_{\zeta_n,\zeta}$ -name d_n for an element of Fin, a \mathbb{P}_{ζ_n} -name for a $\mathbb{P}_{\zeta_n,\zeta}$ -name X_n for an element of \mathcal{U}^i , and a \mathbb{P}_{ζ_n} -name of a $p_n \in \mathbb{P}_{\zeta_n,\zeta}$ with the following properties:

- (a) $q_n \in \mathbb{P}_{\zeta_n}$, $dom(q_n) \setminus \xi \subseteq M \cap \zeta_n$, $q_{n+1} \upharpoonright \zeta_n = q_n$,
- (b) q_n is $(M, \mathbb{P}_{\zeta_n})$ -generic,
- (c) $q_{n+1} \Vdash_{\mathbb{P}_{\zeta_n}} p_{n+1} \in D_n \cap M \cap G$,
- (d) $p_0 \upharpoonright \zeta_0 \ge q_0 \text{ in } \mathbb{P}_{\zeta_0}$,
- (e) $q_{n+1} \Vdash_{\mathbb{P}_{\zeta_{n+1}}} p_{n+1} \upharpoonright [\zeta_n, \zeta_{n+1}) \le p_n \upharpoonright [\zeta_n, \zeta_{n+1})$ (in $\mathbb{P}_{\zeta_n, \zeta_{n+1}}$),
- (f) q_1, p_1, X_0, d_0, X_1 is such that

 $q_1 \Vdash_{\mathbb{P}_{\zeta_1}} \left(p_1 \Vdash_{\mathbb{P}_{\zeta_1,\zeta}} \left(X_0 \text{ is the piece of the partition } c \upharpoonright \mathrm{FU}(A_0) \text{ that lies in } \mathcal{U}^i \right) \right)$

Moreover $q_1 \Vdash p_1 \Vdash d_0 \in M$, $q_1 \Vdash p_1 \in M \cap G \cap D_0$.

(g) $q_{n+1}, p_{n+1}, d_n, X_{n+1}$ is such that

$$q_{n+1} \Vdash_{\mathbb{P}_{\zeta_{n+1}}} \left(p_{n+1} \in M \cap G \cap D_n \wedge \right.$$

$$p_{n+1} \Vdash_{\mathbb{P}_{\zeta_{n+1},\zeta}} \left(d_n = \min_{\text{lex,Fin}} \{ d \in X_n \cap \text{FU}(A_n) : X_n \ominus d \in \mathcal{U}^i \text{ and } \right.$$

$$\min(d) > \max(d_{n-1}) \wedge h[d_{n-1}] \cap h[d] = \emptyset \}$$

$$\wedge X_{n+1} = X_n \cap (X_n \ominus d_n) \right).$$

Since the name d_{n+1} is defined from elements in $M, q_{n+1} \Vdash_{\mathbb{P}_{\zeta_{n+1}}} p_{n+1} \Vdash_{\mathbb{P}_{\zeta_{n+1},\zeta}} d_{n+1} \in M$.

Since $M \prec H(\chi)$, such a sequence exists by the induction hypothesis and the maximum principle.

In the end we let \bar{g} such that

$$q_n \Vdash_{\mathbb{P}_{\zeta_n}} p_n \Vdash_{\mathbb{P}_{\zeta_n,\zeta}} \bar{g} \upharpoonright n = \langle d_i : i < n \rangle$$

and $q = \bigcup_{n < \omega} q_n$ and we let \bar{g}^j be a name such that $q \Vdash \bar{g}^j = \langle g_{2n+j} : n < \omega \rangle$.

Now it is easy to see that q, \bar{g}^j are as desired, i.e., $q \leq p$ is (M, \mathbb{P}_{ζ}) generic and

$$(4.4) q \Vdash fR_{\alpha}\bar{g}.$$

Let G be \mathbb{P}_{α} -generic over V with $q \in G$. Since $M[G] \prec H(\chi)[G]$ (again, see [37, Theorem 2.11]) there are also names $\bar{g} \in M$ as in Equation (4.4). $\square_{4.8.4.1}$

Putting Theorem 3.3 and Theorem 4.1 together yields the following.

Theorem 4.9. Let \mathcal{E} be a P-point and assume CH. Then there is a countable support iteration of proper iterands $\mathbb{P} = \langle \mathbb{P}_{\alpha}, \mathbb{M}(\mathcal{U}_{\beta}) : \beta < \omega_2, \alpha \leq \omega_2 \rangle$ such that in the extension \mathcal{E} is a P-point, there at least three near-coherence classes of ultrafilters and there is a Milliken-Taylor ultrafilter of character $\mathfrak{d} = \mathfrak{c} = \aleph_2$.

We note that Fernández-Bréton [19] recently built a model in which, like in ours, $cov(\mathcal{M}) = \aleph_1 < \mathfrak{d} = \aleph_2$ with a Milliken-Taylor ultrafilter.

5. Generalisation to Fin_k

In this section we generalise the results of Sections 2, 3, and 4 from Fin to Fin_k for $k \geq 1$. We introduce Milliken–Taylor ultrafilters over Fin_k and Matet forcing for Fin_k .

Definition 5.1. Let $k \in \omega \setminus \{0\}$ unless stated otherwise.

- (1) For $p \colon \omega \to k+1$ we let $\operatorname{supp}(p) = \{n \in \omega \, : \, p(n) \neq 0\}.$
 - $\operatorname{Fin}_k = \{p \colon \omega \to k+1 \colon \operatorname{supp}(p) \text{ is finite } \wedge k \in \operatorname{range}(p)\}.$
- (2) For $a, b \in \operatorname{Fin}_k$, we let a < b denote $\operatorname{supp}(a) < \operatorname{supp}(b)$, i.e., $(\forall m \in \operatorname{supp}(a))$ $(\forall n \in \operatorname{supp}(b))(m < n)$, see Def. 2.1(3). A finite or infinite sequence $\langle a_i : i < m \leq \omega \rangle$ of elements of Fin_k is in block-position if for any i < j < m, $a_i < a_j$. The set $(\operatorname{Fin}_k)^{\omega}$ is the set of ω -sequences in block-position, also called block sequences. For $n \geq 1$, the set $[\operatorname{Fin}_k]^n_{<}$ is the set of n-sequences in block-position over Fin_k .
- (3) We define a well-order (of type ω) $\leq_{\text{lex},\text{Fin}_k}$ on the set Fin_k via $a <_{\text{lex},\text{Fin}_k} b$ if $\max(\text{supp}(a)) < \max(\text{supp}(b))$ or $(\max(\text{supp}(a)) = \max(\text{supp}(b))$ and there is an m such that $a \upharpoonright m = b \upharpoonright m$ and a(m) > b(m). For a non-empty set $X \subseteq \text{Fin}_k$ we let $\min_{<_{\text{lex},\text{Fin}_k}}(X)$ be the $\leq_{\text{lex},\text{Fin}_k}$ -least element of X, in parallel to Definition 2.1(11).
- (4) For $k \geq 1$, $a, b \in \operatorname{Fin}_{[1,k]}$, we define the partial semigroup operation + as follows: If $\operatorname{supp}(a) < \operatorname{supp}(b)$, then $a+b \in \operatorname{Fin}_k$ is defined. We let (a+b)(n) = a(n) + b(n). Otherwise a+b is undefined. Thus $a+b = a \upharpoonright \operatorname{supp}(a) \cup b \upharpoonright \operatorname{supp}(b) \cup 0 \upharpoonright (\omega \smallsetminus (\operatorname{supp}(a) \cup \operatorname{supp}(b)))$.
- (5) Let B be a min-unbounded subset of Fin_k , i.e., for any $n, B \cap \{s \in \operatorname{Fin}_k : \{n\} < s\} \neq \emptyset$. We let

$$FU_k(B) = \{b_{n_0} + \dots + b_{n_\ell} : \\ \ell \in \omega, b_{n_i} \in B, b_{n_0} < \dots < b_{n_\ell} \}$$

be the partial subsemigroup of Fin_k generated by B. We call B an FU_k -set if there is a sequence $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ such that $B = \operatorname{FU}_k(\bar{a})$.

- (6) The set $\gamma(\operatorname{Fin}_k)$ is the set of ultrafilters over Fin_k that contain all sets of the form $\{p \in \operatorname{Fin}_k : \{n\} < p\}, n \in \omega$.
- (7) We lift + to $(\bigcup_{k\geq 1}\gamma(\mathrm{Fin}_k))^2$ (so also mixed pairs are allowed) via

$$\mathcal{U} + \mathcal{V} = \Big\{ X \subseteq \operatorname{Fin}_k \,:\, \big\{ s \,:\, \{t \,:\, s + t \in X \} \in \mathcal{V} \big\} \in \mathcal{U} \Big\}.$$

(8) For $A \subseteq \operatorname{Fin}_k$, $s \in \operatorname{Fin}_k$ and $n \in \omega$ we let $(A ; \operatorname{past} s) = \{a \in A : \max(\operatorname{supp}(s)) < \min(\operatorname{supp}(a))\}$. For $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$, $s \in \operatorname{Fin}_k$ and $n \in \omega$ we let $(\bar{a} ; \operatorname{past} s) = \langle a_m : m \geq m_0 \rangle$, where m_0 is the minimal m such that $\max(\operatorname{supp}(s)) < \min(\operatorname{supp}(a_m))$.

(9) For min-unbounded sets $A, B \subseteq \operatorname{Fin}_k$ we let $B \sqsubseteq_k A$ if $B \subseteq \operatorname{FU}_k(A)$. We say B is a k-condensation of A. We let $B \sqsubseteq_k^* A$ if there is an $n \in \omega$ such that $(B : \operatorname{past} \{n\}) \subseteq \operatorname{FU}_k(A)$. For $\bar{a}, \bar{b} \in (\operatorname{Fin}_k)^{\omega}$ we let $\bar{b} \sqsubseteq_k \bar{a}$ if $\bar{b} = \langle b_n : n < \omega \rangle$ and $\{b_n : n < \omega\} \subseteq \operatorname{FU}_k(\{a_n : n < \omega\})$, and $\bar{b} \sqsubseteq^* \bar{a}$ has the obvious meaning.

So $(\operatorname{Fin}_1, +) \cong (\operatorname{Fin}, \cup)$ via $p \mapsto \operatorname{supp}(p)$, and $((\operatorname{Fin}_1)^{\omega}, \sqsubseteq_1) \cong ((\operatorname{Fin})^{\omega}, \sqsubseteq)$ via $\langle a_n : m \in \omega \rangle \mapsto \langle \operatorname{supp}(a_n) : n < \omega \rangle$. For checking whether there is a minunbounded $C \sqsubseteq_k A, B$, a piecewise construction recursively along ω with the aid of $<_{\operatorname{lex},\operatorname{Fin}_k}$ as in the proof of Lemma 3.15 yields \sqsubseteq -weakest (i.e., largest) common lower bounds. Since this is not entirely obvious in the Fin_k -setting, we state it explicitly.

Lemma 5.2. Assume that $\bar{a}, \bar{b} \in (\operatorname{Fin}_k)^{\omega}$ and that there is a $\bar{c} \in (\operatorname{Fin}_k)^{\omega}$ such that $\bar{c} \sqsubseteq_k \bar{a}, \bar{b}$. Then there is a \sqsubseteq_k -largest witness $\bar{c} = \langle c_n : n < \omega \rangle$ to this, that can be found in the following recursive way:

$$c_0 = \min_{\leq_{\text{lex,Fin}_k}} (\text{FU}_k(\bar{a}) \cap \text{FU}_k(\bar{b})),$$

$$c_{n+1} = \min_{\leq_{\text{lex,Fin}_k}} (\text{FU}_k(\bar{a}; \text{past } c_n) \cap \text{FU}_k(\bar{b}; \text{past } c_n)).$$

The proof is like Lemma 2.4.

We write $\bar{c} = \bar{a} \wedge \bar{b}$ for the largest $\bar{c} \sqsubseteq_k, \bar{a}, \bar{b}$.

Hindman's theorem is generalised to $(\operatorname{Fin}_k)^{\omega}$:

Theorem 5.3. (Hindman [24, Cor. 3.3]) Let $k \geq 1$ and $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$. For every finite colouring of $\operatorname{FU}_k(\bar{a})$ there is a infinite block sequence $\bar{b} \sqsubseteq_k \bar{a}$ such that the elements of $\operatorname{FU}_k(\bar{b})$ are monochromatic.

Definition 5.4. A set $\mathcal{H} \subseteq (\operatorname{Fin}_k)^{\omega}$ is called a *adequate family* if the following hold:

- (i) \mathcal{H} is closed \sqsubseteq_{k}^{*} -upwards.
- (ii) \mathcal{H} is stable, i.e., any \sqsubseteq_k -descending ω -sequence of members of \mathcal{H} has a \sqsubseteq^* lower bound in \mathcal{H} .
- (iii) \mathcal{H} has the *Hindman property*: If $\bar{a} \in \mathcal{H}$ and $\mathrm{FU}_k(\bar{a})$ is partitioned into finitely many pieces then there is some $\bar{b} \sqsubseteq_k \bar{a}$, $\bar{b} \in \mathcal{H}$ such that $\mathrm{FU}_k(\bar{b})$ is a subset of a single piece of the partition.

Again Lemma 3.6 applies.

Definition 5.5. (1) For $\bar{a}, \bar{b} \in (\operatorname{Fin}_k)^{\omega}$ we write $\bar{a} \not\perp \bar{b}$ and say that \bar{a}, \bar{b} are compatible, if there is $\bar{c} \sqsubseteq_k \bar{a}, \bar{b}$.

- (2) A set $\mathcal{C} \subseteq (\operatorname{Fin}_k)^{\omega}$ is called *centred*, if for any finite $C \subseteq \mathcal{C}$ there is $\bar{a} \in \mathcal{C}$ that is a generalised condensation of any $\bar{c} \in C$ and if \mathcal{C} is closed under finite alterations i.e., if $\bar{d} \in \mathcal{C}$ and $\bar{d} =^* \bar{e}$ then $\bar{e} \in \mathcal{C}$.
- (3) A non-principal filter \mathcal{F} on Fin_k is said to be an *ordered-union* filter if it has a basis of sets of the form $\operatorname{FU}_k(\bar{d})$ for $\bar{d} \in (\operatorname{Fin}_k)^{\omega}$.
- (4) For a \sqsubseteq_k -descending ω -sequence of members of $(\operatorname{Fin}_k)^{\omega}$ $\langle \bar{a}_n : n \in \omega \rangle$ and $\bar{b} \in (\operatorname{Fin}_k)^{\omega}$ we say: \bar{b} is a diagonal lower bound if $\bar{b} \sqsubseteq \bar{a}_0$ and

$$(\forall s \in \mathrm{FU}_k(\bar{b}))(\bar{b} ; \mathrm{past} \ s) \sqsubseteq_k \bar{a}_{\max(\mathrm{supp}(s))+1}.$$

- (5) Let μ be an uncountable cardinal. An ordered-union filter is said to be $(<\mu)$ stable if, whenever it contains $\mathrm{FU}_k(\bar{b}_\alpha)$ for $\bar{a}_\alpha \in (\mathrm{Fin})^\omega$, $\alpha < \kappa$, for some $\kappa < \mu$,
 then it also contains some $\mathrm{FU}_k(\bar{b})$ for some \bar{b} such that for each α there is n_α with $(\bar{b} ; \mathrm{past} \{n_\alpha\}) \sqsubseteq_k \bar{b}_\alpha)$ for $\alpha < \kappa$. Such an \bar{b} is called a lower bound of $\{\bar{b}_\alpha : \alpha < \kappa\}$. For " $<\omega_1$ -stable" we say "stable".
- (6) A stable ordered-union ultrafilter over Fin_k is called a *Milliken-Taylor ultra-filter*(over Fin_k).
- (7) An ultrafilter is called *idempotent* if $\mathcal{U}+\mathcal{U}=\mathcal{U}$.

In particular, any Milliken–Taylor ultrafilter is Matet-adequate and Lemma 3.6 holds.

Definition 5.6. Let $k \geq 1$. Conditions *Matet forcing (over* Fin_k), \mathbb{M}_k , are pairs (s, \bar{c}) such that $s \in \text{Fin}_k$ and $\bar{c} \in (\text{Fin}_k)^{\omega}$ and $\text{supp}(s) < \text{supp}(c_0)$. The forcing order is $(t, \bar{d}) \leq_k (s, \bar{c})$ (recall the stronger condition is the smaller one) if the following holds:

(a) s = t or there are $n, i_0 < \cdots < i_n \in \omega$, and $j_r \in \{0, \dots, k-1\}$ for $r \le n$ with at least one $j_r = 0$ such that t is a sum in the sense of Def. 5.1(5) of the form

$$t = s + c_{i_0} + \dots + c_{i_n}.$$

(b) $\bar{d} \sqsubseteq_k \bar{c}$ (see Def. 5.1(9)).

In the case of s = t we call (t, \bar{b}) a pure extension of $(s, \bar{a}) = p$. s is called the trunk of p and \bar{a} is called the pure part of p.

Since the trunk and the pure part are in block position and since the pure parts are block-sequences, $s \upharpoonright \operatorname{supp}(s) = t \upharpoonright \operatorname{supp}(s)$ is equivalent to $s \upharpoonright (\operatorname{max}(\operatorname{supp}(s)) + 1) = t \upharpoonright (\operatorname{max}(\operatorname{supp}(s)) + 1)$.

Definition 5.7. Given a Matet-adequate family $\mathcal{H} \subseteq (\operatorname{Fin}_k)^{\omega}$, the notion of forcing $\mathbb{M}_k(\mathcal{H})$ consists of all pairs (s,\bar{a}) such that $\bar{a} \in \mathcal{H}$. The forcing order is the same as in the Matet forcing.

Definition 5.8. Let \mathcal{H} be a subset of $\mathscr{P}(\operatorname{Fin}_k)$, $X \subseteq \operatorname{Fin}_k$, $s \in \operatorname{Fin}_k$.

- (1) For $1 \le i \le k$ we let $set_i(s) = s^{-1}[\{i\}]$, $set_i[X] = \bigcup \{set_i(s) : s \in X\}$. $set[X] = \bigcup \{supp(s) : s \in X\}$.
- (2) For $1 \le i \le k$ we let $\min_i(s) = \min(s^{-1}[\{i\}])$, $\min_i[X] = \{\min_i(s) : s \in X\}$. We have the analogous notions for max.
- (3) The core of \mathcal{H} is the set $\Phi(\mathcal{H}) \subseteq [\omega]^{\omega}$ such that

$$X \in \Phi(\mathcal{H}) \text{ iff } (\exists Y \in \mathcal{H})(\text{set}[Y] \subseteq X).$$

- (4) For j = 1, ..., k, the core of \mathcal{H} at colour j is the set $\Phi_j(\mathcal{H}) \subseteq [\omega]^{\omega}$ such that $X \in \Phi_j(\mathcal{H})$ iff $(\exists Y \in \mathcal{H})(\text{set}_j[Y] \subseteq X)$.
- (5) The minimum projection of \mathcal{H} at colour j is the set

$$\min_{j}(\mathcal{H}) = \{\min_{j}[Y] : Y \in \mathcal{H}\},$$

and analogously we define $\max_{j}(\mathcal{H})$, the maximum projection at colour j.

(6) Let \mathcal{E} be an ultrafilter over ω . We say \mathcal{H} avoids \mathcal{E} if $\Phi(\mathcal{H})$ is nnc to \mathcal{E} .

(7) The $\mathbb{M}_k(\mathcal{H})$ -generic function from ω to k+1 is

$$\mu = \bigcup \{ s \upharpoonright (\max(\operatorname{supp}(s)) + 1) : (\exists \bar{a}) ((s, \bar{a}) \in G) \}.$$

A name for μ is

$$\mu = \bigcup \{ \langle (s \upharpoonright (\max(\operatorname{supp}(s)) + 1), (s, \bar{a}) \rangle : (s, \bar{a}) \in \mathbb{M}_k(\mathcal{H}) \}.$$

Usually we do not write the tildes.

(8) In addition we define the generic *i*-fibres for i = 1, ..., k by letting

$$\mu_i := \mu^{-1}[\{i\}].$$

Again we have $\{\min(y^{-1}[\{i\}]): y \in \mathrm{FU}_k(\bar{a})\} = \{\min(a_n^{-1}[\{i\}]): n \in \omega\}$ and $\bigcup\{b^{-1}[\{i\}]: b \in \mathrm{FU}(\bar{a})\} = \bigcup\{a_n^{-1}[\{i\}]: n \in \omega\}$. Sometimes we identify each FU_k -set $\mathrm{FU}_k(\bar{a})$ with its generating $\bar{a} \in (\mathrm{Fin}_k)^\omega$. Especially in the forcing notion $\mathbb{M}(\mathcal{H})$ we think of \mathcal{H} as a subset of $(\mathrm{Fin}_k)^\omega$ even when we insert an Milliken–Taylor ultrafilter \mathcal{U} for \mathcal{H} .

Remark 5.9. In the case of a centred adequate family $C \subseteq (\operatorname{Fin}_k)^{\omega}$, a density argument shows for any $j = 1, \ldots, k$,

$$\mathbb{M}_k(\mathcal{C}) \Vdash \mu_j \in [\omega]^\omega \land (\forall X \in \Phi_j(\mathcal{C}))(\mu_j \subseteq^* X).$$

The proof of the following theorem is carried out as in the original. It makes frequent use of the good properties from Lemma 3.6.

Theorem 5.10. (Generalisation of [17, " \leftarrow " Theorem 4, " \rightarrow " Cor. 2.5, this direction works also with non-P ultrafilters]) Let \mathcal{U} be a Milliken-Taylor ultrafilter over Fin_k and let \mathcal{W} be a P-point. $\mathcal{W} \succeq_{RB} \Phi(\mathcal{U})$ iff \mathcal{W} continues to generate an ultrafilter after we force with $\mathbb{M}_k(\mathcal{U})$.

Now here is the Fin_k -analogon to Theorem 3.3.

Theorem 5.11. Let \mathcal{E} be a P-point and let \mathcal{U} be a Milliken-Taylor ultrafilter over Fin_k such that $\Phi(\mathcal{U}) \not\leq_{\operatorname{RK}} \mathcal{E}$. In $\mathbf{V}^{\mathbb{M}_k(\mathcal{U})}$ there is a Milliken-Taylor ultrafilter $\mathcal{U}^{\operatorname{ext}} \supseteq \mathcal{U} \upharpoonright \mu$ with $\Phi(\mathcal{U}^{\operatorname{ext}}) \not\leq_{\operatorname{RK}} \mathcal{E}$.

We introduce some notions that allow us to adapt the proof of Theorem 3.3 to a proof of Theorem 5.11.

Definition 5.12. Let $\mathcal{C} \subseteq (\operatorname{Fin}_k)^{\omega}$ be centred. $\mathcal{C}^+ = \{\bar{a} \in (\operatorname{Fin}_k)^{\omega} : \forall \bar{c} \in \mathcal{C}, \bar{c} \not\perp \bar{a}\}$

Definition 5.13. Let $f: \omega \to (k+1)$ and let $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ and let $\mathcal{U} \subseteq (\operatorname{Fin}_k)^{\omega}$.

- (1) We let $\bar{a} \upharpoonright f = \langle a_n : a_n = f \upharpoonright \operatorname{supp}(a_n), n \in \omega \rangle$.
- (2) We let $\mathcal{U} \upharpoonright f = \{\bar{a} \upharpoonright f : \bar{a} \in \mathcal{U}\}.$

Density arguments show:

Lemma 5.14. Let \mathcal{U} be a Milliken-Taylor ultrafilter over Fin_k and let μ be the generic function, see Def. 5.8(7). Then

$$1 \Vdash_{\mathbb{M}_k(\mathcal{U})} \mu \colon \omega \to (k+1).$$
$$1 \Vdash_{\mathbb{M}_k(\mathcal{U})} (\forall \bar{a} \in \mathcal{U}) \bar{a} \upharpoonright \mu \in (\operatorname{Fin}_k)^{\omega}.$$

Also Theorem 3.9 is generalised, and the following now is proved literally as there.

Theorem 5.15. After forcing with $\mathbb{M}_k(\mathcal{U})$, $(\mathcal{U} \upharpoonright \mu)^+$ is a Matet-adequate family for any finite-to-one h, for any $\bar{a} \in (\mathcal{U} \upharpoonright \mu)^+$,

$$(\exists \bar{b}^1, \bar{b}^2 \sqsubseteq \bar{a})(\bar{b}^0 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge \bar{b}^1 \in (\mathcal{U} \upharpoonright \mu)^+ \wedge h[\operatorname{set}(\bar{b}^1)] \cap h[(\operatorname{set}(\bar{b}^2)] = \emptyset).$$

Thus Theorem 5.11 is proved. There is no problem in generalising the iteration theory from Section 4 and hence we arrive at the following result:

Theorem 5.16. Assume CH and let $k \geq 1$. Then there is a countable support iteration of proper iterands $\mathbb{P} = \langle \mathbb{P}_{\alpha}, \mathbb{M}_k(\mathcal{U}_{\beta}) : \beta < \omega_2, \alpha \leq \omega_2 \rangle$ such that in the extension \mathcal{E} is a P-point and there is a Milliken-Taylor ultrafilter over Fin_k of character $\mathfrak{c} = \aleph_2 = \mathfrak{d}$.

The rest of the section is not used in the main theorem.

Now we are concerned with the number of near-coherence classes among the ultrafilters that contain $\Phi(\mathcal{U})$ as a subset.

The following generalises [9, Theorem 38] and adds a new aspect:

Theorem 5.17. For any union-ultrafilter \mathcal{U} over Fin_k , all the ultrafilters among the projections

$$\min_i(\mathcal{U}), i = 1, 2 \dots, k$$

are nearly coherent to $\min_k(\mathcal{U})$ and that class is different from the class of $\max_k(\mathcal{U})$, which is the class of any of the ultrafilters among

$$\max_{j}(\mathcal{U}), j = 1, \dots, k.$$

Proof. We let $M \subseteq \{1, \ldots k-1\}$ be maximal such that for any $m \in M$ the set of $X \in \mathcal{U}$ such that $\text{set}_m[X]$ is infinite is \subseteq -dense in \mathcal{U} . Then exactly for $m \in M \cup \{k\}$, $\max_m(\mathcal{U})$ and $\min_m(\mathcal{U})$ are ultrafilters. Recall, ultrafilter means non-principal ultrafilter. by [7, Proposition 3.9].

In [9, Section 6] Blass proves that the minimum and the maximum class are different. \Box

Theorem 5.18. Let \mathcal{U} be a Milliken-Taylor ultrafilter over Fin_k . Let M be as above. The cores $\Phi_i(\mathcal{U})$, $j \in M \cup \{k\}$, are all nearly coherent.

Proof. Let $A = \mathrm{FU}_k(\bar{a}) \in \mathcal{U}$. We define $I_0 = [0, \max(\mathrm{supp}(a_0)))$, $I_{n+1} = [\max(\mathrm{supp}(a_n)), \max(\mathrm{supp}(a_{n+1})))$. Then we take a finite-to-one function $h_{\bar{a}}$ that is constant on I_n for $n \in \omega$. Then for any $i, j \in M$,

$$(\forall n \in \omega)(h_{\bar{a}}[\operatorname{set}_i(a_n)] \cap h_{\bar{a}}[\operatorname{set}_j(a_n)] \neq \emptyset).$$

Since \mathcal{U} is centred, for any $\mathrm{FU}(\bar{b})$, $\mathrm{FU}(\bar{c}) \in \mathcal{U}$ there is $\bar{d} \sqsubseteq_k \bar{a}, \bar{b}, \bar{c}$, $\mathrm{FU}(\bar{d}) \in \mathcal{U}$. Hence we have for $\bar{d} = \langle d_\ell : \ell < \omega \rangle$ for any $\ell \in \omega$ that there are natural numbers $n, m, r \geq 1$ and natural numbers $i_1 < \cdots < i_n, j_1 < \cdots < j_m, k_1 < \cdots < k_r$, such that

$$d_{\ell} = a_{i_1} + \dots + a_{i_n} = b_{j_1} + \dots + b_{j_m} = c_{k_1} + \dots + c_{k_r}.$$

Then

$$h_{\bar{a}}[\operatorname{set}_i(d_{\ell})] \cap h_{\bar{a}}[\operatorname{set}_i(d_{\ell})]$$
 has size at least $\max(n, m, r)$.

Hence

$$h_{\bar{a}}[\bigcup \{ \operatorname{set}_i(d_{\ell}) : \ell \in \omega \}] \cap h_{\bar{a}}[\bigcup \{ \operatorname{set}_j(d_{\ell}) : \ell \in \omega \}]$$
 is infinite.

Thus also the superset

$$h_{\bar{a}}[\bigcup \{ \operatorname{set}_i(b_\ell) \, : \, \ell \in \omega \}] \cap h_{\bar{a}}[\bigcup \{ \operatorname{set}_j(c_\ell) \, : \, \ell \in \omega \}]$$
 is infinite.

Remark 5.19. Theorem 5.18 can also be proved with the help of Theorem 5.10 and the following folklore result.

Proposition 5.20. Any forcing that diagonalises two ncc filters adds a dominating real.

Proof. Let \mathcal{F} and $\mathcal{F}' \in \mathbf{V}$ be nnc filters and let $x, y \in \mathbf{V}[G]$ be such that $(\forall F \in \mathcal{F})(x \subseteq^* F)$, $(\forall F' \in \mathcal{F}')(y \subseteq^* F')$. Then by the next lemma,

$$n \mapsto \max\{\min(x \setminus (n+1)), \min(y \setminus (n+1))\}$$

is a dominating function.

Lemma 5.21. (Proof of [6, Theorem 3.2]) Let V, W be non-principal filters over ω . V is nearly coherent to W iff

$$\{\max(\text{next}(\cdot, X), \text{next}(\cdot, Y)) : X \in \mathcal{V}, Y \in \mathcal{W}\}$$

is not a \leq^* -dominating family.

We give a proof for the direction we use: Assume that \mathcal{V} and \mathcal{W} are not nearly coherent. Let $h \in {}^{\omega}\omega$ be given, w.l.o.g. we assume that h is strictly increasing and h(0) > 0. We let \tilde{h} be the iterate of h: $\tilde{h}(0) = 0$, $\tilde{h}(n+1) = h(\tilde{h}(n))$. We let $f_e(n) = i$ for $n \in [\tilde{h}(2i), \tilde{h}(2i+2))$ and we let $\tilde{h}(-1) = 0$ and $f_o(n) = i$ for $n \in [\tilde{h}(2i-1), \tilde{h}(2i+1))$. Then there are $V_e, V_o \in \mathcal{V}$ and $W_e, W_o \in \mathcal{W}$ such that $f_e[V_e] \cap f_e[W_e] = \emptyset$ and $f_o[V_o] \cap f_o[W_o] = \emptyset$. Since \mathcal{V} and \mathcal{W} are filters, we can assume $V_e = V_o$ and $W_e = W_o$. Then for any $n \in \omega$ we have $\max(\max(n, V_e), \max(n, W_e)) \geq h(n)$. \square

Now we finish the alternative proof of Theorem 5.18: $\mathbb{M}_k(\mathcal{U})$ preserves \mathcal{E} according to Theorem 5.10 and hence does not add a dominating real. However $\mathbb{M}_k(\mathcal{U})$ diagonalises $\Phi_j(\mathcal{U})$ for $j=1,\ldots,k$ by Remark 5.9. By Proposition 5.20 all the $\Phi_j(\mathcal{U})$, $j=1,\ldots,k$, are nearly coherent.

6. A Ramsey subspace of the space of Fin_k -sequences

Now we change from the Hindman Fin_k space to the Gowers Fin_k space, i.e., we incorporate the Tetris operation and use the resulting generalised type of condensation as partial order. Then we localise the space $((\operatorname{Fin}_k)^\omega, \sqsubseteq_k^{\operatorname{Tetr}})$ by choosing 2k pairwise nnc Ramsey ultrafilters $\mathcal{R}_{i,\min}$, $\mathcal{R}_{i,\max}$, $i=1,\ldots,k$, that will serve as representatives of names of near-coherence classes and get a subspace $((\operatorname{Fin}_k)^\omega(\bar{\mathcal{R}}), \sqsubseteq_k^{\operatorname{Tetr}})$, see Def. 6.8. We show that this space has good Ramsey theoretic properties, as in Theorem 6.15, and we introduce Gowers-Milliken-Taylor ultrafilters.

We tell more about our plans: In Section 7 we define a variant of Matet forcing with a non-centred reservoir that is given by the sequences $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. The resulting forcings are in contrast to the partial order $\mathbb{M}_k(\mathcal{U})$ with a Milliken–Taylor ultrafilter \mathcal{U} (or to $\mathbb{GM}_k(\mathcal{U})$ with a Gowers–Milliken–Taylor ultrafilter \mathcal{U}) not σ -centred. Nevertheless they still fulfil Axiom A and hence are proper.

In order to prove that the new forcings are proper, we insert this section whose main result, Theorem 6.16, is a common strengthening of Gowers' theorem and of Blass' theorem [7, Theorem 2.2].

Definition 6.1. Let $k \in \omega \setminus \{0\}$ unless stated otherwise.

(1) For any $j \geq 2$ we define on Fin_j the *Tetris operation*: $T_j \colon \operatorname{Fin}_j \to \operatorname{Fin}_{j-1}$ by $T_j(p)(n) = \max\{p(n) - 1, 0\}$ and let $T = \bigcup \{T_j : 2 \leq j \leq k\}$.

For $\bar{a} \in (\operatorname{Fin}_j)^{\omega}$, we write $T[\bar{a}]$ for $\langle T(a_n) : n < \omega \rangle$. For $X \subseteq \operatorname{Fin}_j$, we write T[X] for $\{T(x) : x \in X\}$.

(2) Let B be a min-unbounded subset of Fin_k . We let

$$TFU_{k}(B) = \{ T^{(j_{0})}(b_{n_{0}}) + \dots + T^{(j_{\ell})}(b_{n_{\ell}}) : \\ \ell \in \omega, b_{n_{i}} \in B, \\ b_{n_{0}} < \dots < b_{n_{\ell}}, j_{i} \in k, (\exists i \leq \ell)(j_{i} = 0) \}$$

be the partial subsemigroup of Fin_k generated by B. Here $T^{(0)}$ is the identity, $T^{(1)}=T,\,T^{(j+1)}=T\circ T^{(j)}$.

- (3) For $\bar{b} \in (\operatorname{Fin}_k)^{\omega}$ we let $\operatorname{TFU}_k(\bar{b}) = \operatorname{TFU}_k(\{b_n : n \in \omega\})$. We call B a TFU_k - set if $B = \operatorname{TFU}_k(\bar{b})$ for some $\bar{b} \in (\operatorname{Fin}_k)^{\omega}$.
- (4) For $j \geq 2$, we define T: $\gamma(\operatorname{Fin}_j) \to \gamma(\operatorname{Fin}_{j-1})$ via

$$T(\mathcal{U}) = \{ X \subseteq \operatorname{Fin}_{j-1} : \{ s : T(s) \in X \} \in \mathcal{U} \}.$$

- (5) For min-unbounded sets $A, B \subseteq \operatorname{Fin}_k$ we let $B \sqsubseteq_k^{\operatorname{Tetr}} A$ if $B \subseteq \operatorname{TFU}_k(A)$. Note that here k is not allowed to drop. We say B is a k-Tetris-condensation of A. We write $B \sqsubseteq_k^{\operatorname{Tetr},*} A$ if there is an $n \in \omega$ such that $(B ; \operatorname{past} n) \subseteq \operatorname{TFU}_k(A)$.
- (6) We introduce the Tetris condensation order $\sqsubseteq_k^{\text{Tetr}}$ on sequences. For $\bar{a}, \bar{b} \in (\text{Fin}_k)^{\omega}$ we let $\bar{b} \sqsubseteq_k^{\text{Tetr}} \bar{a}$ if $\bar{b} = \langle b_n : n < \omega \rangle$ and $\{b_n : n < \omega\} \subseteq \text{TFU}_k(\{a_n : n < \omega\})$, and $\bar{b} \sqsubseteq_k^{\text{Tetr},*} \bar{a}$ has the obvious meaning.

From the following lemma we will use only the part of the statement that the described procedure produces a lower bound.

Lemma 6.2. Assume that $\bar{a}, \bar{b} \in (\operatorname{Fin}_k)^{\omega}$ and that there is a $\bar{c} \in (\operatorname{Fin}_k)^{\omega}$ such that $\bar{c} \sqsubseteq_k^{\operatorname{Tetr}} \bar{a}, \bar{b}$. Then there is a $\sqsubseteq_k^{\operatorname{Tetr}}$ -largest witness \bar{c} to this, that can be gotten in the following recursive way:

$$c_0 = \min_{\leq_{\text{lex,Fin}_k}} (\text{TFU}_k(\bar{a}) \cap \text{TFU}_k(\bar{b})),$$

$$c_{n+1} = \min_{\leq_{\text{lex,Fin}_k}} (\text{TFU}_k(\bar{a}; \text{past } c_n) \cap \text{TFU}_k(\bar{b}; \text{past } c_n)).$$

The proof that \bar{c} is the largest lower bound requires additional analysis in the second subcase of the third case in the proof of Proposition 2.4, due to the Tetris operation. We skip it.

An important and deep property of the space $((\operatorname{Fin}_k)^{\omega}, \sqsubseteq_k^{\operatorname{Tetr}})$ is the following:

Theorem 6.3. (Gowers, see e.g. [22], [40, Theorem 2.22]) Let $k \geq 1$ and $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$. For every finite colouring of $\operatorname{TFU}_k(\bar{a})$ there is a infinite block sequence $\bar{b} \sqsubseteq_k^{\operatorname{Tetr}} \bar{a}$ such that the elements of $\operatorname{TFU}_k(\bar{b})$ are monochromatic.

The following proposition gives a first picture:

Proposition 6.4. For each $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ there is a $\bar{b} \sqsubseteq_k^{\operatorname{Tetr}} \bar{b}$ such that for each $s \in \operatorname{TFU}_k(\bar{b})$ we have

$$(6.1) \quad \min_{1}(s) < \min_{2}(s) < \dots < \min_{k}(s) < \max_{k}(s) < \max_{k-1}(s) < \dots < \max_{1}(s).$$

Proof. For $s \in \mathrm{TFU}_k(\bar{a})$ we let c(s) = 1 if s fulfils (6.1) and c(s) = 0 otherwise. By Gowers' Theorem there is $\bar{b} \sqsubseteq_k^{\mathrm{Tetr}} \bar{a}$, such that $\mathrm{TFU}_k(\bar{b})$ is monochromatic for c. We show that the colour can only be 1. We take $s_i \in \mathrm{TFU}_k(\bar{b})$, $s_1 < s_2 < \cdots < s_{2k-1}$. Then

(6.2)
$$s = T^{(k-1)}(s_1) + T^{(k-2)}(s_2) + \dots + T(s_{k-1}) + s_k + T(s_{k+1}) + \dots + T^{(k-1)}(s_{2k-1}) \in TFU_k(\bar{b})$$
 fulfils $c(s) = 1$.

Definition 6.5.

(1) A non-principal filter \mathcal{F} over Fin_k is said to be an *Tetris-ordered-union* filter if it has a basis of sets of the form $\operatorname{TFU}_k(\bar{d})$ for $\bar{d} \in (\operatorname{Fin}_k)^{\omega}$.

(2) For a $\sqsubseteq_k^{\text{Tetr}}$ -descending ω -sequence of members of $(\text{Fin}_k)^{\omega} \langle \bar{a}_n : n \in \omega \rangle$ and $\bar{b} \in (\text{Fin}_k)^{\omega}$ we say: \bar{b} is a diagonal lower bound if $\bar{b} \sqsubseteq_k^{\text{Tetr}} \bar{a}_0$ and

$$(\forall s \in \mathrm{TFU}_k(\bar{b}))(\bar{b} ; \mathrm{past} \; s) \sqsubseteq_k^{\mathrm{Tetr}} \bar{a}_{\max(\mathrm{supp}(s))+1}.$$

- (3) Let μ be an uncountable cardinal. A Tetris-ordered-union filter is said to be $(<\mu)$ -stable if, whenever it contains $\mathrm{TFU}_k(\bar{b}_\alpha)$ for $\bar{b}_\alpha \in (\mathrm{Fin}_k)^\omega$, $\alpha < \kappa$, for some $\kappa < \mu$, then it also contains some $\mathrm{TFU}_k(\bar{b})$ for some $\bar{b} \in (\mathrm{Fin}_k)^\omega$ such that for each α there is n_α with $(\bar{b}; \mathrm{past} \{n_\alpha\}) \sqsubseteq_k^{\mathrm{Tetr}} \bar{b}_\alpha$ for $\alpha < \kappa$. Such an \bar{b} is called a lower bound of $\{\bar{b}_\alpha : \alpha < \kappa\}$. For " $<\omega_1$ -stable" we say "stable".
- (4) A stable Tetris-ordered-union ultrafilter is called a Gowers-Milliken-Taylor ultrafilter.

Definition 6.6. A set $\mathcal{H} \subseteq (\operatorname{Fin}_k)^{\omega}$ is called a *Gowers-adequate family* if the following hold:

- (i) \mathcal{H} is closed $\sqsubseteq_k^{\text{Tetr},*}$ -upwards.
- (ii) \mathcal{H} is stable, i.e., any $\sqsubseteq_k^{\text{Tetr}}$ -descending ω -sequence of members of \mathcal{H} has a $\sqsubseteq_k^{\text{Tetr},*}$ -lower bound in \mathcal{H} .
- (iii) \mathcal{H} has the Gowers property: If $\bar{a} \in \mathcal{H}$ and $\mathrm{TFU}_k(\bar{a})$ is partitioned into finitely many pieces then there is some $\bar{b} \sqsubseteq_k^{\mathrm{Tetr}} \bar{a}, \ \bar{b} \in \mathcal{H}$, such that $\mathrm{TFU}_k(\bar{b})$ is a subset of a single piece of the partition.

In particular, any Gowers–Milliken–Taylor ultrafilter is Gowers-adequate. Gowers-adequate families have better properties than stated in the definition; the analogue to Lemma 3.6 hold for them.

Definition 6.7. In the general case of $i \in \{1, ..., k\}$, $x \in \{\min, \max\}$, we let the variable x range over $\{\min, \max\}$, and we also write $x_i(a)$ for $\min_i(a)$, $\max_i(a)$. We call $i \in \{1, ..., k\}$ sometimes a coordinate, a value or a level.

We introduce a new space, called $((\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}}), \sqsubseteq_k^{\operatorname{Tetr}})$, the one advertised in the Equation (1.1) in the introduction.

Definition 6.8. Let $k \geq 1$ and let

$$\bar{\mathcal{R}} = \langle \mathcal{R}_{i,x} : i = 1, \dots, k, x = \min, \max \rangle$$

be a $\{1,\ldots,k\} \times \{\min,\max\}$ -sequence of pairwise nnc Ramsey ultrafilters. We define

(6.3)
$$(\operatorname{Fin}_{k})^{\omega}(\bar{\mathcal{R}}) = \{ \bar{a} \in (\operatorname{Fin}_{k})^{\omega} : \forall (X_{i} \in \mathcal{R}_{i,\min}, Y_{i} \in \mathcal{R}_{i,\max})_{1 \leq i \leq k}$$

$$(\exists^{\min-\mathrm{unb}} s \in \operatorname{TFU}_{k}(\bar{a})) \bigwedge_{1 \leq i \leq k} (\min_{i}(s) \in X_{i} \wedge \max_{i}(s) \in Y_{i}) \}.$$

Here we write $(\exists^{\min-\text{unb}}s)$ as a abbreviation for "the set of witnesses s in min-unbounded".

Remark 6.9. We remark that Equation (6.3) is equivalent to

$$(\operatorname{Fin}_{k})^{\omega}(\bar{\mathcal{R}}) = \{ \bar{a} \in (\operatorname{Fin}_{k})^{\omega} : \forall (X_{i} \in \mathcal{R}_{i,\min}, Y_{i} \in \mathcal{R}_{i,\max})_{1 \leq i \leq k}$$

$$(\exists \bar{b} \sqsubseteq_{k}^{\operatorname{Tetr}} \bar{a})(\forall n) \bigwedge_{1 \leq i \leq k} (\min_{i}(b_{n}) \in X_{i} \wedge \max_{i}(b_{n}) \in Y_{i}) \}.$$

However, it speaks about the b_n and not about the elements of $TFU_k(\bar{b})$.

First we recall [7, Lemma 1.1]:

Lemma 6.10. Let $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ be nnc Ramsey ultrafilters, $X_1 \in \mathcal{R}_{1,\min}$, $Y_1 \in \mathcal{R}_{1,\max}$ and let $(B_n)_n$ be a descending sequence of elements of $\mathcal{R}_{1,\min}$, $(D_n)_n$ be a descending sequence of elements of $\mathcal{R}_{1,\max}$. Then there are $X_1^1 \subseteq X_1$, $Y_1^1 \subseteq Y_1$, $X_1^1 \in \mathcal{R}_{1,\min}$, $Y_1^1 \in \mathcal{R}_{1,\max}$ such that

- (1) $(\forall x \in X_1^1)(\forall y \in Y_1^1)(y < x \to x \in B_y).$
- (2) $(\forall y \in Y_1^1)(\forall x \in X_1^1)(x < y \to y \in D_x).$
- (3) In the increasing enumeration of $X_1^1 \cup Y_1^1$ the elements of X_1^1 and Y_1^1 alternate.

Lemma 6.11. There are \sqsubseteq_k^* -incompatible elements in $(\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$.

Proof. For notational simplicity we assume that k=1 and the Ramsey ultrafilters are \mathcal{R}_{\min} \mathcal{R}_{\max} . For transferring the situation to the general case, just replace the value 1 by k and work only with the top ultrafilters $\mathcal{R}_{k,\min}$, $\mathcal{R}_{k,\max}$. Lemma 6.10, applied to $X_1 = Y_1 = \omega$, $A_n = B_n = [2n, \omega)$ yields sets $X \in \mathcal{R}_{\min}$ and $Y \in \mathcal{R}_{\max}$ such that in the increasing enumeration $\langle x_n : n < \omega \rangle$ of $X \cup Y$ the even numbers just give X and the odd number just give Y and that the x_n have distance at least 2. Then we take a natural number, call ℓ_n , such that $x_{2n} < \ell_n < x_{2n+1}$. Then

$$\bar{a} = \langle \{x_{2n}, x_{2n+1}\} : n \in \omega \rangle,$$

$$\bar{b} = \langle \{x_{2n}, \ell_n, x_{2n+1}\} : n \in \omega \rangle$$

have

(6.5)
$$X = \{ \min_{1}(a_n) : n \in \omega \} \land Y = \{ \max_{1}(a_n) : n \in \omega \},$$

and hence $\bar{a} \in (\operatorname{Fin}_1)(\mathcal{R}_{\min}, \mathcal{R}_{\max})$, and the same holds for \bar{b} . Obviously, the conditions $\bar{a}, \bar{b} \in (\operatorname{Fin}_1)^{\omega}(\bar{\mathcal{R}})$ are $\sqsubseteq_1^{\operatorname{Tetr}}$ -incompatible.

By the way, our only possibility to verify that some sequence is in $(\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ is to construct one that has ultrafilter projections as in Equation (6.5).

By applying e.g. the trick in [7, Cor. 2.3] one sees that also below every condition there are incompatible conditions.

Lemma 6.12. For any ultrafilter \mathcal{U} over Fin_k , the projections $\min_k(\mathcal{U})$, $\max_k(\mathcal{U})$ generate (by adding supersets) ultrafilters, and for $1 \leq i < k$, either each of the projections $\min_i(\mathcal{U})$, $\max_i(\mathcal{U})$ generates a filter or each of them contains the empty set.

Proof. They are filters: $\min_i(\mathcal{U}) = \{\min_i[X] : X \in \mathcal{U}\}$. Since \mathcal{U} is a filter, the projection $\min_i(\mathcal{U})$ is closed under finite intersections. For i < k it is possible that there is an $X \in \mathcal{U}$ such that $\min_i[X] = \max_i[X] = \emptyset$. By definition of Fin_k , the top coordinate $\min_k(\mathcal{U})$ does not contain the empty set. The family $\min_k(\mathcal{U})$ generates an ultrafilter: Let $A \subseteq \omega$. Then $\operatorname{Fin}_k = \{a \in \operatorname{Fin}_k : \min_k(a) \in A\} \cup \{a \in \operatorname{Fin}_k : \min_k(a) \notin A\}$ is a partition of Fin_k , and one part lies in \mathcal{U} . The same holds for the maximum projection.

Definition 6.13. For an ultrafilter $\mathcal{U} \in \gamma(\operatorname{Fin}_k)$ we say \mathcal{U} projects to $\overline{\mathcal{R}}$, if

$$(\forall i \in \{1, \dots, k\})(\min_i(\mathcal{U}) = \mathcal{R}_{i,\min} \land \max_i(\mathcal{U}) = \mathcal{R}_{i,\max}).$$

Definition 6.14. Let $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$. $\operatorname{set}(\bar{a}) = \bigcup \{\operatorname{supp}(a_n) : n < \omega\}$.

In the following Theorem 6.15, the case of k=1 is due to Blass' [7, Theorem 2.2], and the unlocalised case of $k \geq 1$ is proved in Gowers' theorem [40, 2.22]. In a sense the result can be seen as an instance of the programme outlined by Todorcevic in [40, Remark 7.27]. The author is thankful to Andreas Blass for his careful explanation of his proof of [7, Theorem 2.1].

Theorem 6.15. Assume CH. For any k, $\bar{\mathcal{R}}$ as in Def. 6.8 the following hold:

- (1) Let $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ and let c be a colouring of $\operatorname{TFU}_k(\bar{a})$ into finitely many colours. Then there is a $\bar{b} \sqsubseteq_k^{\operatorname{Tetr}} \bar{a}$, such that $\bar{b} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ and $\operatorname{TFU}_k(\bar{b})$ is c-monochromatic.
- (2) Let $\bar{a}_n \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$, $n \in \omega$, be $\sqsubseteq_k^{\operatorname{Tetr}}$ -decending. Then there is a $\sqsubseteq_k^{\operatorname{Tetr},*}$ -lower bound in $(\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$.
- (3) If $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ and $\mathcal{R}_{0,\min}$, $\mathcal{R}_{0,\max}$ are two additional nnc ultrafilters, that are nnc to each of the $\mathcal{R}_{i,x}$, $i=1,\ldots,k$, $x=\min,\max$, then there is a Gowers-Milliken-Taylor ultrafilter \mathcal{U}_k over Fin_k such that $\operatorname{TFU}_k(\bar{a}) \in \mathcal{U}_k$ and for $i=1,\ldots,k$, $\min_i(\mathcal{U}) = \mathcal{R}_{i,\min}$, $\max_i(\mathcal{U}) = \mathcal{R}_{i,\max}$ and $\Phi(\mathcal{U}_k)$ is nnc $\mathcal{R}_{0,x}$, $x=\min,\max$.

Since parts (1) and (2) are absolute, we get:

Theorem 6.16. Let $k \geq 1$ and let $\bar{\mathcal{R}} = \langle \mathcal{R}_{i,\min}, \mathcal{R}_{i,\max} : i = 1, ... k \rangle$ be a sequence of pairwise non nearly coherent Ramsey ultrafilters.

- (1) Any $\sqsubseteq_k^{\text{Tetr}}$ -descending ω -sequence of elements of $(\text{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ has a $\sqsubseteq_k^{\text{Tetr},*}$ -lower bound in $(\text{Fin}_k)^{\omega}(\bar{\mathcal{R}})$.
- (2) Let $n \in \omega \setminus \{0\}$ and $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ and let c be a colouring of $[\operatorname{TFU}_k(\bar{a})]_{<}^n$ into finitely many colours. Then there is a $\bar{b} \sqsubseteq_k \bar{a}$, $\bar{b} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ such that $[\operatorname{TFU}_k(\bar{b})]_{<}^n$ is c-monochromatic.

We prove Theorem 6.15.

Proof. We prove (1), (2) and (3) simultaneously by induction on k. The case of k = 1 is given by Blass and will be repeated below for k = 1.

It suffices to prove (1) of the theorem for a partition of Fin_k instead of TFU_k(\bar{a}) since $T^{j_0}(a_{n_0}) + \cdots + T^{j_r}(a_{n_r}) \in \text{TFU}_k(\bar{a})$ corresponds to $(n_0, k - j_0), \ldots, (n_r, k - j_r) \in \text{Fin}_k$.

We collect some technical facts separately, before we prove Theorem 6.15.

First we give a general pattern for proving the existence of a particular ultrafilter over Fin_i via compactness.

Lemma 6.17. Let \mathcal{R}_{\min} , \mathcal{R}_{\max} be two nnc Ramsey ultrafilters. Let $\bar{a} \in (\operatorname{Fin}_1)^{\omega}(\mathcal{R}_{\min}, \mathcal{R}_{\max})$. The compact Hausdorff space

$$\{\mathcal{U} \in \gamma(\operatorname{Fin}_1) : \min(\mathcal{U}) = \mathcal{R}_{\min} \wedge \max(\mathcal{U}) = \mathcal{R}_{\max} \wedge \operatorname{TFU}_1(\bar{a}) \in \mathcal{U}\}$$

is not empty.

Proof. Since min and max are continuous function, the space is compact. We show that there is an ultrafilter \mathcal{U} over Fin_1 in the named space. For this it suffices to show that for any $X \in \mathcal{R}_{\min}$ and any $Y \in \mathcal{R}_{\max}$

R(X,Y) There are min-unboundedly many
$$s \in \mathrm{TFU}_1(\bar{a})$$
 such that $\min_1(s) \in X$ and $\max_1(s) \in Y$.

We argue why this suffices: Then the open and closed set

 $C(X,Y) = \{ \mathcal{U} \in \gamma(\operatorname{Fin}_1) : (\exists A \in \mathcal{U})(A \text{ is a min-unb. set of witnesses to } R(X,Y)) \}$ is not empty.

Since $\mathcal{R}_{i,x}$ are ultrafilters, the intersection of finitely many sets of the form C(X,Y) is again a set of the C(X',Y') with X' being the intersection of the finitely many X's and Y' being the intersection of the finitely many Y's.

Thus any finite intersection of sets of the form C(X,Y) is not empty, and by compactness of $\gamma(\operatorname{Fin}_i)$, there is an ultrafilter

$$\mathcal{U} \in \bigcap \{C(X,Y) : X \in \mathcal{R}_{k,\min}, Y \in \mathcal{R}_{k,\max}\} \subseteq \gamma(\operatorname{Fin}_i),$$

which means

$$\min_{i}(\mathcal{U}) = \mathcal{R}_{k,\min} \wedge \max_{1}(\mathcal{U}) = \mathcal{R}_{k,\max}.$$

To fulfil requirement R(X,Y): We verify that for any $n \in \omega$ there is a block $s \in (\mathrm{TFU}_1(\bar{a}); \mathrm{past}\ \{n\})$ such that $\min_1(s) \in X$ and $\max_1(s) \in Y$. Why? Since $\bar{a} \in (\mathrm{Fin}_k)^{\omega}(\bar{\mathcal{R}})$, and $X \in \mathcal{R}_{\min}$ and $Y \in \mathcal{R}_{\max}$ there are min-unboundedly many $s \in \mathrm{TFU}_1(\bar{a})$ such that $\min_1(s) \in X \wedge \max_1(s) \in Y$.

Starting from now, we just write the requirements and show how to fulfil them. For proving statement (1) at k on the bases of (1) and (3) at k-1, we rework and extend the proof of Gowers' original theorem as given by Todorcevic [40, pp. 35–36].

Lemma 6.18. Let $\mathcal{R}_{i,x}$, $i=1,\ldots,k$, $x=\min,\max$ be pairwise nnc Ramsey ultrafilters. For any $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ the following are equivalent:

- (1) $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}}).$
- (2) There is an ultrafilter $\mathcal{U} \in \gamma(\operatorname{Fin}_k)$ such that

$$\mathrm{TFU}_k(\bar{a}) \in \mathcal{U} \wedge (\forall i \in \{1, \dots k\}) (\min_i(\mathcal{U}) = \mathcal{R}_{i,\min} \wedge \max_i(\mathcal{U}) = \mathcal{R}_{i,\max}).$$

Proof. (1) implies (2): By the compactness method as explained in the previous lemma it suffices to show that for any $(X_i \in \mathcal{R}_{i,\min} Y_i \in \mathcal{R}_{i,\max})$

$$(6.6) \qquad (\exists^{\min-\mathrm{unb}} s \in \mathrm{TFU}_k(\bar{a})) (\forall i \in \{1, \dots, k\}) (\min_i(s) \in X_i \wedge \max_i(s) \in Y_i).$$

However, this is guaranteed by $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\mathcal{R})$.

(2) implies (1). Let \mathcal{U} be as in (2). Given sets $X_i \in \mathcal{R}_{i,\min}$, $Y_i \in \mathcal{R}_{i,\max}$, $i = 1, \ldots, k$, there is a set $S \in \mathcal{U}$ such that

$$S \subseteq \mathrm{TFU}_k(\bar{a}) \wedge (\forall i \in \{1, \dots, k\}) (\min_i [S] \subseteq X_i \wedge \max_i [S] \subseteq Y_i).$$

Since $S \in \mathcal{U}$, S is min-unbounded. Any $s \in S$ fulfils $\min_i(s) \in X_i \wedge \max_i(s) \in Y_i$.

We name the set of ultrafilters with given projections:

Definition 6.19. For $1 \le i \le k$ and a sequence $\langle \mathcal{R}_{j,x} : 1 \le j \le i, x \in \{\min, \max\} \rangle$ or pairwise nnc Ramsey ultrafilters we let

(6.7)
$$\gamma^{\text{good}}(\operatorname{Fin}_{i}, \bar{\mathcal{R}}) = \{ \mathcal{U} \in \gamma(\operatorname{Fin}_{i}) : (\forall j \in \{1, \dots, i\}) \\ (\min_{j}(\mathcal{U}) = \mathcal{R}_{j,\min} \wedge \max_{j}(\mathcal{U}) = \mathcal{R}_{j,\max}) \}.$$

Cave: Sometimes we use ultrafilter sequences that carry shifted indices.

Lemma 6.20. For any i, $\bar{\mathcal{R}}$ as above, the space $\gamma^{\text{good}}(\operatorname{Fin}_i, \bar{\mathcal{R}})$ is closed under +.

Proof. This is proved in [7, page 99, second but last paragraph]. \Box

In Lemma 6.22 we shall look at "mixed additions." First we state some simple observations.

Lemma 6.21.

(1) For any $X \subseteq \operatorname{Fin}_i$, for any $1 \le j < j' \le i$ we have

$$\min_{j}[X] \cap \min_{j'}[X] = \emptyset,$$

since for each $s \in \operatorname{Fin}_i$, $\min_i(s) \neq \min_{i'}(s)$.

- (2) Given $t \in \operatorname{Fin}_j$, there is a particular (in $<_{\operatorname{Fin}_{j+1},lex}$ largest) $s \in \operatorname{Fin}_{j+1}$ with T(s) = t: We let $\operatorname{supp}(t) = \operatorname{supp}(s)$ and t(n) = s(n) + 1 for $n \in \operatorname{supp}(s)$. We write for this t now s^{\sharp} .
- (3) We can lift the sharp operation by letting $X^{\sharp} := \{s^{\sharp} : s \in X\}$. Given $\mathcal{U} \in \gamma(\operatorname{Fin}_{j})$, there is a particular ultrafilter $\mathcal{V} \in \gamma(\operatorname{Fin}_{j+1})$ with $T(\mathcal{V}) = \mathcal{U}$, namely

$$\mathcal{V} = \{ X^{\sharp} : X \in \mathcal{U} \}.$$

We write $V = \text{lift}(\mathcal{U})$. Then $\min_{j'+1}(V) = \min_{j'}(\mathcal{U})$ for $j' = 1, \ldots, j$, and $\min_1(V) = \{\emptyset\}$ and analogously for max.

Now we prove a technical lemma about the sums of good ultrafilters. Prima facie the ultrafilters do not fit, e.g., $\min_k(\mathcal{X}) = \mathcal{R}_{k,\min}$ and $\min_{k-1}(\mathcal{V}) = \mathcal{R}_{k,\min}$. However, this mixed sum is defined on purpose.

Lemma 6.22. Let $\bar{\mathcal{R}} = \langle \mathcal{R}_{i,x} : 1 \leq i \leq k, x = \min, \max \rangle$. Let $\mathcal{X} \in \gamma^{\text{good}}(\operatorname{Fin}_k, \bar{\mathcal{R}})$ and $\mathcal{V} \in \gamma^{\text{good}}(\operatorname{Fin}_{k-1}, \langle \mathcal{R}_{2,x}, \dots, \mathcal{R}_{k,x}, x = \min, \max \rangle)$. Then $\mathcal{W} = \mathcal{X} + \mathcal{V} \in \gamma^{\text{good}}(\operatorname{Fin}_k, \bar{\mathcal{R}})$.

Proof. We have $W \in \gamma(\operatorname{Fin}_k, \overline{\mathcal{R}})$ iff for $v = 1, \dots, k, x = \min, \max, \text{ for any } X \in \mathcal{W},$

(6.8)
$$\{x_v(s) : s \in X\} \in \mathcal{R}_{v,x}.$$

We fix some $X \in \mathcal{W}$. Then $\{s : \{t : s + t \in X\} \in \mathcal{V}\} \in \mathcal{X}$. For any $s \in \text{Fin}_k$, $t \in \text{Fin}_{k-1}$, such that s < t and $s + t \in X$, and for any $v = 1, \ldots, k$, $x = \min, \max$,

 $x_v(s+t)$ is either $x_v(s)$ or $x_v(t)$. Either the set of (s,t) on which x_v is the first is in

$$\mathcal{X} \times \mathcal{V} = \{ Z \subseteq \operatorname{Fin}_k \times \operatorname{Fin}_{k-1} : \{ s : \{ t : (s,t) \in Z \} \in \mathcal{V} \} \in \mathcal{X} \},$$

or its complement is in $\mathcal{X} \times \mathcal{V}$.

We start with the high values of v: For the values v=k, we have for any $s \in \operatorname{Fin}_k$, $t \in \operatorname{Fin}_{k-1}$, $(s+t)^{-1}[\{v\}] = s^{-1}[\{v\}]$ and hence its minimum and/or maximum is from $x_v(s)$. Thus for $X \in \mathcal{X} + \mathcal{V}$, $x = \min$, max we have

(6.9)
$$\{x_v(s+t) : s+t \in X\} \in \mathcal{R}_{v,x},$$

as requested in Equation (6.8).

Now we proceed by downwards induction on v.

The next value is v = k - 1 and $x = \min$ or $x = \max$. For $s \in \operatorname{Fin}_k$, $t \in \operatorname{Fin}_{k-1}$, s < t, $s + t \in X$, we have $x_v(s + t) = x_v(s)$ or $x_v(s + t) = x_v(t)$, and there is a set $Z \in \mathcal{X} \times \mathcal{V}$ such that for any $(s,t) \in Z$ the decision is the same. We claim that this decision can only be for the s-parts and hence $\{x_v(s + t) : s + t \in X\} = \{x_v(s) : s + t \in X\} \in \mathcal{R}_{k-1,x}$. This is seen as follows: Since any $s + t \in X$ is an element of Fin_k , we have

$$(6.10) \{x_{v+1}(s+t) : s+t \in X\} \cap \{x_v(s+t) : s+t \in X\} = \emptyset.$$

For $X \in \mathcal{X}+\mathcal{V}$, the left-hand set on the left-hand side of Equation (6.10), i.e., the set $\{x_k(s+t): s+t \in X\}$, is already in the ultrafilter $\mathcal{R}_{k,x}$, by the analysis of the coordinate k in Equation 6.9. Hence by Equation (6.10),

$$\{x_{k-1}(s+t): s+t\in X\}\not\in\mathcal{R}_{k,x}$$

and so $Z \in \mathcal{X} \times \mathcal{V}$ did not decide for the t-part. Since there must be a decision on Z as to whether $x_{k-1}(s+t) = x_{k-1}(s)$ or $x_{k-1}(s+t) = x_{k-1}(t)$, we have $x_{k-1}(s+t) = x_{k-1}(s)$ for $s \in Z$ and hence

$${x_{k-1}(s+t) : s+t \in X} \in \mathcal{R}_{k-1,x}.$$

In the next step we verify Equation (6.8) for v = k - 2 with the same technique, shifted by -1 in the v-component, and proceed by downwards induction down to v = 1.

Lemma 6.23. (See [40, Lemma 2.23]) Let $k \geq 2$. Let $2 \leq j \leq k$. Then $T: \gamma(\operatorname{Fin}_{j}) \to \gamma(\operatorname{Fin}_{j-1})$ is a continuous onto +-homomorphism.

So we use freely $T(\mathcal{U}+\mathcal{V}) = T(\mathcal{U})+T(\mathcal{V})$.

The following is a strengthening of [40, Lemma 2.24]. It is stated in more generality than we need for the proof of statement (1) at k so that we can use it also for the proof of statement (3) at k.

Lemma 6.24. Let $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. Let \mathcal{U}_{k-1} be a Gowers-Milliken-Taylor ultrafilter over Fin_{k-1} , such that $\operatorname{TFU}_{k-1}(T[\bar{a}]) \in \mathcal{U}_{k-1}$ and $\operatorname{min}_i(\mathcal{U}_{k-1}) = \mathcal{R}_{1+i,\min}$, $i = 1, \ldots, k-1$, and same for \max and such that $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ are Ramsey ultrafilters that are $\operatorname{nnc} \Phi(\mathcal{U}_{k-1})$. In addition we let \mathcal{S}_1 , \mathcal{S}_2 be two ultrafilters that are $\operatorname{nnc} \Phi(\mathcal{U}_{k-1})$ and nnc to $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$. Tet h be a finite-to-one function. The

⁷ The Gowers–Milliken–Taylor ultrafilter \mathcal{U}_{k-1} is given by the by the induction hypothesis, (2) at k-1 with shifted indices in the ultrafilters.

set

$$S_{k} := \{ \mathcal{X} \in \gamma^{\text{good}}(\operatorname{Fin}_{k}, \bar{\mathcal{R}}) : \operatorname{T}(\mathcal{X}) = \mathcal{U}_{k-1} \land$$

$$\bigwedge_{n} \operatorname{TFU}_{k}(\bar{a}) \in \mathcal{X} \land$$

$$\bigwedge_{j=1,2} (\exists E_{j} \in \mathcal{S}_{j})(\exists X \in \mathcal{X})(h[\operatorname{set}(X)] \cap h[E_{j}] = \emptyset) \}$$

is a closed non-empty subset of $\gamma(\operatorname{Fin}_k)$.

Proof. In case k=1, just let \mathcal{U}_0 be the principal ultrafilter $\{f\}$ with f(n)=0 for any n, and we leave out B from the requirements below. Let \mathcal{U}_{k-1} be given by the induction hypothesis, be as in (2) at k-1 for $\mathcal{R}_{1+i,x}$, $i=1,\ldots k-1$, $x=\min,\max$, such that $\mathcal{R}_{1,x}$ is nnc $\Phi(\mathcal{U}_{k-1})$ and such that $\mathrm{TFU}_{k-1}(T[\bar{a}]) \in \mathcal{U}_{k-1}$. Let h be a finite-to-one function.

Let $B \in \mathcal{U}_{k-1}$, let for $1 \leq i \leq k$ sets $X_i \in \mathcal{R}_{i,\min}$, $Y_i \in \mathcal{R}_{i,\max}$ be given. The requirement

$$R(X_1, \dots, X_k, Y_k, \dots, Y_1, B, h)$$
 says
$$(\exists^{\min-\text{unb}} s)(s_n \in \text{TFU}_k(\bar{a}_n))(\forall i = 1, \dots, k)$$

$$(\min_i(s) \in X_i \land \max_i(s) \in Y_i \land T(s) \in B \land$$

$$\bigwedge_{j=1,2} (\exists E_j \in \mathcal{S}_j)(h[\text{set}(s)] \cap h[E_i]) = \emptyset).$$

Again the requirements are closed under finite conjunctions, ⁸hence it suffices that any single requirement can be fulfilled.

We take $\bar{b} \in \mathcal{U}_{k-1} \cap B$ such that $(\forall n)(\forall i = 1, \dots, k-1)(\min_i(b_n) \in X_{i+1} \wedge \max_i(b_n) \in Y_{i+1})$. Then look at where the sets X_1 , Y_1 lie outside set (\bar{b}) . We take $X_1^1 \subseteq X_1$, $X_1^1 \in \mathcal{R}_{1,\min}$ and $Y_1^1 \subseteq Y_1$, $Y_1^1 \in \mathcal{R}_{1,\max}$ and $\bar{c} \sqsubseteq_k^{\text{Tetr}} \bar{b}$, $\bar{c} \in \mathcal{U}_{k-1}$ such that X_1^1 , Y_1^1 , set (\bar{c}) are pairwise disjoint. We can do this since $\mathcal{R}_{i,x}$ nnc $\Phi(\mathcal{U})$. Let $r_0 \in \omega$ be given. We take $x_0 \in X_1^1$, $x_0 > r_0$, $r(0) \ge r_0$ such that $x_0 < \sup(c_{r(0)})$ and we take $y_0 \in Y_1^1$ that lies past $c_{r(0)}$ and let $r_1 = y$ and such that

$$(\exists d = d_{r(0)} \in TFU_k(\bar{a})(\min_1(d) = x_0 \land T(d) = c_{r(0)}) \land \max_1(d) = y_0).$$

Such at d exists since $\bar{a} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. We choose the next triple $(x_1, c_{r(1)}, y_1)$ with r_1 in the place of r_0 , and so on. Since \mathcal{S}_1 , \mathcal{S}_2 are ultrafilters, $\mathcal{S}_1 \cap \mathcal{S}_2$ is not meager and hence contains a set $E_1 = E_2$ such that $h[E_1]$ that is disjoint from infinitely many of the sets $h[(r_m, r_{m+1}]], m \in \omega$. We choose such an m. Then $s = d_{r(m)}$ fulfils the requirement. Thus we showed that S_k is not empty.

Remark 6.25. The version with the starting set $\mathrm{TFU}_k(\bar{a})$ and the extra ultrafilters \mathcal{S}_1 , \mathcal{S}_2 will be used for the proof of (3) at k. For (1) at k, we can drop the \mathcal{S}_j and replace $\mathrm{TFU}_k(\bar{a})$ by the full Fin_k .

Now we are ready to resume the thread of Todorcevic's proof of [40, Lemma 2.24].

⁸Note that h is fixed. In order to look at all h simultaneously, it is best to work with two finite-to-one functions h_1 , h_2 . For any finitely many finite-to-one functions h^j , j < J, there are two finite-to-one functions h_1 , h_2 such that for any $n \in \omega$, for any j < J, $(h^j)^{-1}[\{n\}] \subseteq h_1^{-1}[\{n\}]$ or $\subseteq h_2^{-1}[\{n\}]$. Such a compactness argument is not used here.

Definition 6.26. For $\mathcal{U}, \mathcal{V} \in \bigcup_{j=1}^k \gamma(\operatorname{Fin}_j)$, we let $\mathcal{V} \leq \mathcal{U}$ if $\mathcal{U} + \mathcal{V} = \mathcal{V} + \mathcal{U} = \mathcal{V} = \mathcal{V} + \mathcal{V}$. So \mathcal{V} is idempotent and \mathcal{V} is stronger and richer than \mathcal{U} .

The relation \leq is a preorder on $\bigcup_{j=1}^{k} \gamma(\operatorname{Fin}_{j})$.

Lemma 6.27. Let \mathcal{U}_{k-1} be as in Lemma 6.24. There is an idempotent ultrafilter $\mathcal{U}^{\text{idem}} \in \gamma^{\text{good}}(\operatorname{Fin}_k, \bar{\mathcal{R}})$ such that $\mathcal{U}^{\text{idem}} \leq \mathcal{U}_{k-1}$ and $T(\mathcal{U}^{\text{idem}}) = \mathcal{U}_{k-1}$.

Proof. We let $S_k = \{\mathcal{X} \in \gamma^{\text{good}}(\operatorname{Fin}_k, \bar{\mathcal{R}}) : \operatorname{T}(\mathcal{X}) = \mathcal{U}_{k-1}\}$. By Lemma 6.24, $S_k \neq \emptyset$. By Lemma 6.22 the set $\{\mathcal{X} + \mathcal{U}_{k-1} : \mathcal{X} \in S_k\}$ is a subset of $\gamma^{\text{good}}(\operatorname{Fin}_k, \bar{\mathcal{R}})$. The set $\{\mathcal{X} + \mathcal{U}_{k-1} : \mathcal{X} \in S_k\}$ is a non-empty closed subsemigroup, since the sum $\mathcal{V} + \mathcal{U}_{k-1} + \mathcal{W} + \mathcal{U}_{k-1}$ belongs to $S_k + \mathcal{U}_{k-1}$. This is seen as follows: We have

$$T(\mathcal{V}+\mathcal{U}_{k-1}+\mathcal{W}) = \mathcal{U}_{k-1} + T\mathcal{U}_{k-1} + \mathcal{U}_{k-1} = \mathcal{U}_{k-1},$$

where the latter equation holds by induction hypothesis. Then we pick an idempotent ultrafilter $W \in S_k + \mathcal{U}_{k-1}$, say $W = \mathcal{V} + \mathcal{U}_{k-1}$ with $\mathcal{V} \in S_k$. Let $\mathcal{U}^{\text{idem}} = \mathcal{U}_{k-1} + \mathcal{V} + \mathcal{U}_{k-1}$. Then $T\mathcal{U}^{\text{idem}} = \mathcal{U}_{k-1}$ and $\mathcal{U}^{\text{idem}} = \mathcal{U}_{k-1} + \mathcal{W}$, being a sum of idempotent ultrafilters, is idempotent.

Moreover $\mathcal{U}^{\text{idem}} + \mathcal{U}_{k-1} = \mathcal{U}_{k-1} + \mathcal{U}^{\text{idem}} = \mathcal{U}^{\text{idem}}$, so $\mathcal{U}^{\text{idem}} \leq \mathcal{U}_{k-1}$. By Lemma 6.22 (and a relative thereof for additions from the righthand-side) we have $\mathcal{U}^{\text{idem}} \in \gamma^{\text{good}}(\text{Fin}_k, \bar{\mathcal{R}})$.

Here is the penultimate step of the proof of statement (1) of Theorem 6.15: We pick an idempotent ultrafilter $\mathcal{U}^{\text{idem}}$ as in the previous lemma. Let P be a piece of the given partition of Fin_k such that $P \in \mathcal{U}^{\text{idem}}$. Now by induction on n we build a tree Tr of finite increasing sequences x_0, x_1, \ldots, x_n of elements of Fin_k and branching sets $A_0 \supseteq A_{1,x_0} \supseteq \cdots \supseteq A_{n,x_0,\ldots x_{n-1}} \in \mathcal{U}_k$ with the following properties for any $n \in \omega$,

- (a) $A_0 = P$.
- (b) $x_n \in A_{n,x_0,x_1,...,x_{n-1}}$ and $T^{(k-\ell)}[A_{n,x_0,...,x_{n-1}}] = A_{n,x_0,...,x_{n-1}}^{\ell}$ for $1 \le \ell \le k$. So $A_{n,x_0,...,x_{n-1}} = A_{n,x_0,...,x_{n-1}}^{k}$.
- (c) $(\mathcal{U}^{\text{idem}}x)(T^{(k-i)}(x_n) + T^{(k-j)}(x) \in A_{n,x_0,\dots,x_{n-1}}^{\max\{i,j\}})$ for $1 \le i, j \le k$.
- (d) The tree Tr has branching sets in $\mathcal{U}^{\text{idem}}$, i.e. given $(x_0, \ldots, x_n) \in Tr$ for any n, the set of immediate tree successors of (x_0, \ldots, x_n) fulfils:

(6.12)
$$\{x_{n+1} : x_{n+1} \text{ could serve in the tree } Tr$$
 as a prolongation of $(x_0, \dots, x_n)\} = A_{n+1, x_0, \dots, x_n} \in \mathcal{U}^{\text{idem}}.$

As shown in the proof of [40, Theorem 2.22], any branch \bar{x} of the tree Tr has $\mathrm{TFU}_k(\bar{x}) \subseteq P$. We show that there is a branch \bar{c} of Tr such that $\bar{c} \in (\mathrm{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. We let $\tilde{A}_0 = A_0$ and or $n \in \omega$ we let

(6.13)
$$\tilde{A}_{n+1} = \bigcap \{ A_{m+1,x_0,\dots,x_m} : m \le n, (x_0,\dots,x_m) \in Tr, \\ \max(\sup (x_m)) < n+1 \}.$$

Then $\tilde{A}_n \in \mathcal{U}^{\text{idem}}$, and they are \subseteq -descending.

Now we come to the last step.

Final Claim

There is a branch \bar{c} through the Grigorieff tree Tr such that $\bar{c} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$.

Proof. We recall another lemma.

Lemma 6.28. If U_{k-1} is a Gowers-Milliken-Taylor ultrafilter over Fin_{k-1} with projections and

$$(\forall i \in \{1, \dots, k-1\})(\dot{x}_i(\mathcal{U}_{k-1}) = \mathcal{R}_{i+1,x})$$

and $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ are given and both are nnc to $\Phi(\mathcal{U}_{k-1})$ and $\mathcal{V} \in S_k + \mathcal{U}_{k-1}$, $\mathcal{V} \leq \mathcal{U}_{k-1}$, then $(\min_1, T, \max_1)(\mathcal{V}) = \mathcal{R}_{1,\min} \times \mathcal{U}_{k-1} \times \mathcal{R}_{1,\max}$.

Proof. No other order is possible. Since $\mathcal{V} = \mathcal{V} + \mathcal{U}_{k-1} = \mathcal{U}_{k-1} + \mathcal{V}$, we have $X \in \mathcal{V}$ iff $\{s: t: s+t \in X\} \in \mathcal{U}_{k-1}\} \in \mathcal{V}$ and $X \in \mathcal{V}$ iff $\{s: t: s+t \in X\} \in \mathcal{V}\} \in \mathcal{U}_{k-1}$. However, since \mathcal{U}_{k-1} is a Gowers-Milliken-Taylor ultrafilter, by Lemma 6.4 the high values of $s \in X \in \mathcal{U}_{k-1}$ are always in the middle. So only the order $\mathcal{R}_{1,\min} \times \mathcal{U}_{k-1} \times \mathcal{R}_{1,\max}$ in the triple projection is possible.

Now we return to the proof of the final claim. From \tilde{A}_n , $n \in \omega$, as given in Equation (6.13) we define three families of sets:

$$B_n := \{ \min_{1}(s) : s \in \tilde{A}_n \},$$

(6.14)
$$C_n := \{ T(s) : s \in \tilde{A}_n \land \min_1(s) \in B_n \},$$

$$D_n := \{ \max_{1}(s) : s \in \tilde{A}_n \wedge \min_{1}(s) \in B_n \wedge T(s) \in C_n \}.$$

Since $\tilde{A}_n \in \mathcal{U}^{\text{idem}} \in \gamma^{\text{good}}(\text{Fin}_k, \bar{\mathcal{R}})$, we have $B_n \in \mathcal{R}_{1,\min}$, $C_n \in \mathcal{U}_{k-1}$, $D_n \in \mathcal{R}_{1,\max}$.

We prove a relative of Blass' Lemma 6.10 for triples:

Lemma 6.29. Fix $k \geq 1$. If k = 1, then there is no Gowers-Milliken-Taylor ultrafilter \mathcal{U}_{k-1} , we replace \mathcal{U}_{k-1} by the filter $\{0\}$, where 0 is a "block" with empty support. Suppose we are given

- (1) A Gowers-Milliken-Taylor ultrafilter \mathcal{U} over Fin_{k-1} with projections $\mathcal{R}_{i+1,x}: 1 \leq i \leq k-1 \rangle$.
- (2) $\mathcal{R}_{1,x}$, Ramsey and nnc $\Phi(\mathcal{U}_{k-1})$, $x = \min, \max$, and elements $X_1 \in \mathcal{R}_{1,x}$ and $Y_1 \in \mathcal{R}_{1,\max}$,
- (3) a descending sequence \tilde{A}_n as in Equation (6.13) of subsets of an idempotent ultrafilter $\mathcal{U}^{\text{idem}} \leq \mathcal{U}_{k-1}$, $T(\mathcal{U}^{\text{idem}}) = \mathcal{U}_{k-1}$ and $\mathcal{U}^{\text{idem}} \in \gamma^{\text{good}}(\text{Fin}_k, \bar{\mathcal{R}})$ contains P.
- (4) B_n , C_n , D_n are defined from $\langle \tilde{A}_n : n < \omega \rangle$ as in Equation (6.14).

Then there is some $\mathrm{TFU}_k(\bar{b}) \in \mathcal{U}_{k-1}$ and there is $X \in \mathcal{R}_{1,\min}$, $Y \in \mathcal{R}_{1,\max}$ with the following properties

(a) the blocks of X, range(\bar{b}) and Y lie as follows

$$x_0 < b_0 < y_0 < x_1 < b_1 < y_1 < \dots$$

⁹We remark that in the case of ideals (or coideals) over ω , Grigorieff [23, Section 1] worked with similar trees.

(b) the following three conditions on diagonal lower bounds hold

$$(\forall n) \qquad (x_n \in B_{y_{n-1}}),$$

$$(\forall n) \qquad (b_n \in C_{x_n}),$$

$$(\forall n) \quad (y_n \in D_{\max(\text{supp}(b_n))}).$$

Proof. First step:

We take on each of the three parts diagonal lower bounds. We take $X_{0.5} \in \mathcal{R}_{1,\text{min}}$ such that

$$\forall x, y \in X_{0.5} (y < x \to x \in B_y).$$

We take $Y_{0.5} \in \mathcal{R}_{1,\text{max}}$ such that

$$\forall x, y \in Y_{0.5}(y < x \to x \in D_y).$$

We take $\bar{b}_{0.5} \in \mathcal{U}_{k-1}$ such that

$$\forall n < m(b_{0.5,m} \in C_{\max(\text{supp}(b_n^1))}).$$

Then we take $X_1 \subseteq X_{0.5}$, $Y_1 \subseteq Y_{0.5}$, $\bar{b}_1 \sqsubseteq_k^{\text{Tetr}} \bar{b}_{0.5}$, $X_1 \in \mathcal{R}_{1,\min}$, $Y_1 \in \mathcal{R}_{1,\max}$, $\bar{b}_1 \in \mathcal{U}_{k-1}$ such that X_1 , X_2 and $\text{set}(\bar{b}_1)$ are pairwise disjoint. Since $\mathcal{R}_{1,\min}$ and $\mathcal{R}_{1,\max}$ are nnc and both are nnc to $\Phi(\mathcal{U}_{k-1})$, there are such strengthenings. The diagonal conditions hold also for X_1 , \bar{b}_1 , Y_1 .

Second step:

We start working towards the triple alternation pattern in clause (a). We take $X_2 \subseteq X_1, X_2 \in \mathcal{R}_{1,\min}, \bar{b}_{1.5} \in \mathcal{U}_{k-1}, \bar{b}_{1.5} \sqsubseteq_k^{\text{Tetr}} \bar{b}_1, Y_{1.5} \in \mathcal{R}_{1,\max}, Y_{1.5} \subseteq Y_1$, such that

$$(6.15) (\forall x < x' \in X_2)(\exists n)(\exists y \in Y_{1.5})(x < \operatorname{supp}(b_{1.5,n}) < y < x').$$

Since $\mathcal{R}_{1,\min}$ is a Q-point and nnc $\Phi(\mathcal{U}_{k-1})$ and nnc to $\mathcal{R}_{1,\min}$, we find such a sparse set X_2 and $Y_{1.5}$ and $\bar{b}_{1.5}$.

We take $\bar{b}_2 \sqsubseteq_k^{\text{Tetr}} \bar{b}_{1.5}$, $\bar{b}_2 \in \mathcal{U}_{k-1}$, such that

$$(6.16) \qquad (\forall n < m)(\exists x \in X_2)(\exists y \in Y_{1.5})(\text{supp}(b_{2.n}) < y < x < \text{supp}(b_{2.m}).$$

Such a \bar{b}_2 can be found via the Taylor-colouring law of $[\text{TFU}_{k-1}(\bar{b}_{1.5})]^2_{<}$ that sends s < t to 1 if there are the desired elements of X_2 and of $Y_{1.5}$ between s and t and to 0 otherwise. We find $\bar{b}_2 \sqsubseteq_k^{\text{Tetr}} \bar{b}_{1.5}$, $\text{TFU}_{k-1}(\bar{b}_2) \in \mathcal{U}_{k-1}$ such that the monochromatic colour is 1.

We take $Y_2 \subseteq Y_{1.5}, Y_2 \in \mathcal{R}_{1,\text{max}}$, such that

$$(\forall y < y' \in Y_2)(\exists n)(\exists x \in X_2)(y < x < \text{supp}(b_{2,n}) < y').$$

Third step:

We fill up a tiny bit: If between two adjacent blocks of \bar{b}_2 there is only an element of X_2 but no larger element of Y_2 , we add such a point from $Y_{1.5}$ to Y_2 and thus get, when doing this for all situations a set Y_3 such that $Y_2 \subseteq Y_3 \subseteq Y_{1.5}$. By Equation (6.16) there are such elements of $Y_{1.5}$ such that the alternating pattern is established.

If between two adjacent points of X_2 there is only an element of $Y_{1.5}$ but no element of Y_2 or if there is only an element of $\bar{b}_{1.5}$ but no element of \bar{b}_2 , we add, if applicable, such a point from Y_1 to Y_3 and thus get, when doing this for all situations a set Y_4 such that $Y_3 \subseteq Y_4 \subseteq Y_{1.5}$, and we add, if applicable, such a block from range($\bar{b}_{1.5}$) to \bar{b}_2 and thus get, when doing this for all situations a set

 \bar{b}_3 such that $\bar{b}_2 \sqsubseteq_k^{\text{Tetr}} \bar{b}_3 \sqsubseteq_k^{\text{Tetr}} \bar{b}_{1.5}$, $\bar{b}_2 \in \mathcal{U}_{k-1}$. By Equation (6.15) such a filling up is possible.

We let $X_4 = X_2$, $\bar{b}_4 = \bar{b}_3$. Thus we have an alternating chain of enumerating $X_4 \cup \text{range}(\bar{b}_4) \cup Y_4$ as $x_0 < b_0 < y_0 < x_1 < \dots$

Fourth step:

We "attack the possible exceptions" to the diagonalisation requirements in clause (b).

Exceptions for x: Suppose that in the common increasing enumeration of $X_4 \cup \text{range}(\bar{b}_4) \cup Y_4$ there is an element $x \in X_4$ such that for no $x' \in X_{1.5}$, $y < x' < x < b_{4,k}$. We map the last \bar{b}_4 -block before x to x by a finite-to-one function that sends the whole interval $[\min(b_{4,n}), x]$ to $\min(b_{4,n})$. We do this for all such x. We define f arbitrarily on the other points so that it will be a finite-to-one function. Then we use that $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ nnc $\Phi(\mathcal{U}_{k-1})$ and take $\bar{b}_5 \sqsubseteq_k^{\mathrm{Tetr}} \bar{b}_4$, $\bar{b}_5 \in \mathcal{U}_{k-1}$ and $X_5 \subseteq X_4$, $X_5 \in \mathcal{R}_{1,\min}$ and $Y_5 \subseteq Y_4$, $Y_4 \in \mathcal{R}_{1,\max}$ such that $h[Y_5] \cap h[\mathrm{set}(\bar{b}_5)] = \emptyset$, $h[X_5] \cap h[\mathrm{set}(\bar{b}_5)] = \emptyset$. Then for the common enumeration of X_5 , \bar{b}_5 and Y_5 we have: Any $x \in X_5$ that is preceded by an element of Y_5 or one of \bar{b}_5 is shielded by an $x' \in X_1$ from them and hence fulfils the condition that corresponds to the first equation in (b) for X_5 , \bar{b}_5 , Y_5 . However clause (a) is already ruined.

Exceptions for $b_{5,k}$: Suppose that in the common increasing enumeration of $X_5 \cup \operatorname{range}(\bar{b}_5) \cup Y_5$ there is an element $b_{5,k} \in \operatorname{range}(\bar{b}_5)$ such that for no $b' \in \bar{b}_1$ we have for the largest $x \in X_5$ preceding $b_{5,k}$, $y < x < b' < b_{5,k}$. (There might be more x's in the pattern, since in the first manipulation of the foruth step we may have destroyed the alternating pattern. It is only important to do the procedure with a smaller y.) We take a finite-to-one function that sends the whole interval $[y, \max(b_5, k)]$ to $\min(b_{k,k-1})$. We do this for all such $b_{5,k}$. We define f arbitrarily on the other points so that it will be a finite-to-one function. Then we use that $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ nnc $\Phi(\mathcal{U}_{k-1})$ and take $\bar{b}_6 \sqsubseteq_k^{\mathrm{Tetr}} \bar{b}_6$, $\bar{b}_6 \in \mathcal{U}_{k-1}$ and $X_6 \subseteq X_5$, $X_6 \in \mathcal{R}_{1,\min}$ and $Y_6 \subseteq Y_5$, $Y_6 \in \mathcal{R}_{1,\max}$ such that $h[Y_6] \cap h[\operatorname{set}(\bar{b}_6)] = \emptyset$, $h[X_6] \cap h[\operatorname{set}(\bar{b}_6)] = \emptyset$.

Exceptions for y: Suppose that in the common increasing enumeration of $X_6 \cup \operatorname{range}(\bar{b}_6) \cup Y_6$ there is an element $y \in Y_6$ such for no $y' \in Y_1$ we have that for the latest block $b_{6,n}$ before y the constellation $x < b_{6,n} < y' < y < x', x, x' \in X_6$. Again the remark in parantheses applies. We map the last interval [x,y] before y to x by a finite-to-one function. We do this for all such y. We define f arbitrarily on the other points so that it will be a finite-to-one function. Since $\mathcal{R}_{1,\min}$, $\mathcal{R}_{1,\max}$ are nnc $\Phi(\mathcal{U}_{k-1})$ there are $\bar{b}_7 \sqsubseteq_k^{\operatorname{Tetr}} \bar{b}_6$, $\bar{b}_7 \in \mathcal{U}_{k-1}$ and $X_7 \subseteq X_6$, $X_7 \in \mathcal{R}_{1,\min}$ and $Y_7 \subseteq Y_6$, $Y_7 \in \mathcal{R}_{1,\max}$ such that $h[Y_7] \cap h[\operatorname{set}(\bar{b}_7)] = \emptyset$, $h[X_7] \cap h[\operatorname{set}(\bar{b}_7)] = \emptyset$.

By the thinning out in the fourth step we ensured that X_7 , Y_7 , \bar{b}_7 respect clause (b) in the following form:

$$(\forall x \in X_7)(\forall y \in Y_7)(y < x \to x \in B_y)$$

$$(6.17) \qquad (\forall b \in \operatorname{range}(\bar{b}_7))(\forall x \in X_7)(x < \operatorname{supp}(b) \to b \in C_x)$$

$$(\forall y \in Y_7)(\forall b \in \operatorname{range}(\bar{b}_7))(\operatorname{supp}(b) < y \to y \in D_{\max(\operatorname{supp}(b))}).$$

However, clause (a) is ruined. Note that the diagonal laws from Equation 6.17 are preserved under further thinning out.

Fifth step:

We restore the triple alternation pattern again. We repeat the steps two and three, and thus get $X_8 \subseteq X_7$, $Y_8 \subseteq Y_7$, $\bar{b}_8 \sqsubseteq_k^{\text{Tetr}} \bar{b}_7$ that respect (a) and (b).

We conclude the proof of part (1) of Theorem 6.15: Let X, \bar{b} , Y be as in Lemma 6.29. By Equation (6.14), for each n, there is

$$c_n \in \tilde{A}_{\max(\text{supp}(c_{n-1}))}$$
 s.t. $(\min_1(c_n) = x_n \wedge T(c_n) = b_n \wedge \max_1(c_n) = y_n)$.

By the definition of the \tilde{A}_n we have $\mathrm{TFU}_k(\bar{c}) \subseteq P$. Since

$$\bar{b} \in \mathcal{U}_{k-1} \in \gamma^{\text{good}}(\operatorname{Fin}_{k-1}, \langle \mathcal{R}_{1+i,x} : 1 \leq i \leq k-1, x = \min, \max \rangle),$$

 $\{\min_{1}(c_n) : n < \omega\} = X \in \mathcal{R}_{1,\min},$
 $\{\max_{1}(c_n) : n < \omega\} = Y \in \mathcal{R}_{1,\max}, \text{ and }$
 $\Phi(\mathcal{U}_{k-1}) \text{ is nnc } \mathcal{R}_{1,\min}, \mathcal{R}_{1,\max},$

we have $\bar{c} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. So \bar{c} is branch of the Grigorieff tree with the desired properties. Thus Theorem 6.15 part (1) is proved.

We prove part (2). Again we encounter the phenomenon (as in the proof of the Hindman colouring in the $\mathbb{M}(\mathcal{U})$ -extension in Lemma 3.17) that we almost proved the existence of diagonal lower bounds already during the proof of the colouring theorem.

Lemma 6.30.

(1) $a \sqsubseteq_k^{\text{Tetr}}$ -descending sequence $\bar{a}_n, n \in \omega, \bar{a}_n \in (\text{Fin}_k)^{\omega}(\bar{\mathcal{R}}),$

$$(6.18) B_n := \{ \min_1(s) : s \in \mathrm{TFU}(\bar{a}_n) \}$$

$$C_n := \{ T(s) : s \in \mathrm{TFU}(\bar{a}_n) \wedge \min_1(s) \in B_n \}$$

$$D_n := \{ \max_1(s) : s \in \mathrm{TFU}(\bar{a}_n) \wedge \min_1(s) \in B_n \wedge T(s) \in C_n \}$$

Then there is some $\bar{b} \sqsubseteq_k^{\text{Tetr},*} T(\bar{a}_n)$, $\bar{b} \in (\text{Fin}_{k-1})^{\omega} (\langle \mathcal{R}_{1+i,x} : 1 \leq i \leq k-1, x = \min, \max \rangle)$ and there is $X \in \mathcal{R}_{1,\min}$, $Y \in \mathcal{R}_{1,\max}$ with the following properties the blocks of \bar{b} and X and Y lie as follows

$$x_0 < b_0 < y_0 < x_1 < b_1 < y_1 < \dots$$

and the following three conditions on diagonal lower bounds hold

$$(6.19) \qquad (\forall n) (x_n \in B_{y_{n-1}} \land b_n \in C_{x_n} \land y_n \in D_{\max(\text{supp}(b_n))})$$

Proof. This is proved as in Lemma 6.29. In the first step, we take separate diagonal lower bounds, call them as there. For the middle part $\bar{b}_1 \in (\operatorname{Fin}_k)^{\omega}(\langle \mathcal{R}_{1+i,x} : 1 \leq i \leq k-1, x = \min, \max \rangle)$ we choose by (3) at k-1 a Gowers-Milliken-Taylor ultrafilter $\mathcal{U}_{k-1} \in \gamma^{\operatorname{good}}(\operatorname{Fin}_{k-1}\langle \mathcal{R}_{1+i,x} : 1 \leq i \leq k-1, x = \min, \max \rangle)$ such that $\Phi(\mathcal{U}_{k-1})$ is nnc $\mathcal{R}_{1,x}$ and such that $\operatorname{TFU}_{k-1}(\bar{b}_1) \in \mathcal{U}_{k-1}$. Then we can go on as in the proof of Lemma 6.29.

Equation 6.19 yields a diagonal bound $\bar{c} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ of $\bar{a}_n, n \in \omega$. This finishes the proof of statement (2) at k.

Now we turn to (3) at k:

Let S_1 and S_2 be two new ultrafilters, nnc to any of the $\langle \mathcal{R}_{i,x} : i = 1, \ldots, k, x = \min, \max \rangle$. Using CH, we enumerate via $\langle X_{\alpha}, h_{\alpha} : \alpha < \omega_1 \rangle$ any subset X_{α} of Fin_k and any finite-to-one function h_{α} such that each pair appears cofinally often in the enumeration. By induction on $\alpha < \omega_1$ we choose $\bar{a}_{\alpha} \in (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ such that

- (1) $\bar{a}_{\alpha+1}$ decides X_{α} and
- (2) for $i = 1, 2, \exists E_i \in \mathcal{S}_i, h_{\alpha}[\operatorname{set}(\bar{a}_{\alpha+1})] \cap h_{\alpha}[E_i] = \emptyset.$
- (3) $(\forall \beta < \alpha)(\bar{a}_{\beta} \sqsubseteq_{k}^{\text{Tetr},*} \bar{a}_{\alpha}).$

Then we let \mathcal{U} be generated by $\{TFU_k(\bar{a}_{\alpha}: \alpha < \omega_1)\}.$

Given \bar{a}_{β} , $\beta < \alpha$, we first take a $\sqsubseteq_k^{\text{Tetr},*}$ -lower bound \bar{a}_{α} of them as guaranteed by (2) at k. We let $h = h_{\alpha}$ By induction hypothesis there is a Gowers–Milliken–Taylor ultrafilter \mathcal{U}_{k-1} avoiding \mathcal{S}_i and containing $\text{TFU}_{k-1}(T[\bar{c}_{\alpha}])$. In step 0, just drop all clauses about \bar{a}_{α} . Then we build S_k in Lemma 6.24 (this time in the form with the $\mathcal{R}_{1,x}$ and the \mathcal{S}_j and the starting set $\text{TFU}_k(\bar{a})$, note that this lemma uses only (3) at k-1), i.e., the requirement pertains to h_{α} , and $\text{TFU}_k(\bar{a})$ such that for any $X \in \mathcal{X} \in S_k$, $(\exists E_i \in \mathcal{S}_i)_{i=1,2}(h[\text{set}[X]] \cap h[E_i] = \emptyset)$ and $\text{TFU}_k(\bar{a}_{\alpha}) \in \mathcal{U}^{\text{idem}}$. We use, that h-avoiding \mathcal{S}_i is preserved by + to find an idempotent ultrafilter $\mathcal{U}^{\text{idem}} \leq \mathcal{U}_{k-1}$, $\mathcal{U}^{\text{idem}} \in S_k + \mathcal{U}_{k-1}$ such that $\text{TFU}_k(\bar{a}_{\alpha}) \in \mathcal{U}^{\text{idem}}$ and $\mathcal{U}^{\text{idem}}$ h avoids \mathcal{S}_j . Then we take Y_{α} to be X_{α} , if $X_{\alpha} \in \mathcal{U}^{\text{idem}}$ or $Y = \text{Fin}_k \setminus X_{\alpha}$ otherwise. Now we build all the Grigorieff tree with $\tilde{A}_n \in \mathcal{U}^{\text{idem}}$ and $\tilde{A}_n \subseteq \text{TFU}_k(\bar{a}_{\alpha})$. A branch of it is taken as $\bar{a}_{\alpha+1}$. Then $\text{TFU}_k(\bar{a}_{\alpha+1}) \subseteq Y_{\alpha}$.

Thus we finish the proof of part (3) at k of Theorem 6.15.

Question 6.31. We do not know a proof in the Baumgartner [3] style. The analogous question for Gowers' Fin_k theorem is asked in [40].

Now we show that under CH or under MA or after forcing with the pure part of $\mathbb{GM}(\bar{\mathcal{R}})$ there is a Gowers–Milliken–Taylor ultrafilter.

Corollary 6.32. Let k, $\bar{\mathcal{R}}$ be as in Def. 6.8. The family $(\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ is Gowers-adequate.

Corollary 6.33. Let k, $\bar{\mathcal{R}}$ be as in Def. 6.8. Under CH or under $\mathsf{MA}(\sigma\text{-centred})$ there is an Gowers-Milliken-Taylor ultrafilter.

Proof. We perform a $\sqsubseteq_k^{\text{Tetr},*}$ -downward construction of length $\mathfrak c$ with lower bound in the limit steps that are either given by CH or provided by $\mathsf{MA}(\sigma\text{-centred})$. In the successor we use Theorem 6.15 to decide whether the next subset of Fin_k is in or out with the aid of Theorem 6.15.

Definition 6.34. Let $k \geq 2$ and let $1 \leq i < k$. Let \mathcal{U} be an ultrafilter over Fin_k . We let

$$\Phi_{\geq i+1}(\mathcal{U}) = \{ \{ s^{-1}[\{i+1,\dots,k\}] : s \in X \} : X \in \mathcal{U} \},$$

and call $\Phi_{>i+1}(\mathcal{U})$ the core strictly above i.

The following generalises [9, Theorem 38] and shows that in any Gowers–Milliken–Taylor ultrafilter over Fin_k the 2k pairwise nnc Ramsey projections can be found. The additional information on the higher cores can be seen as a justification of the inductive procedure in the proof 6.15.

Theorem 6.35. For any Gowers-Milliken-Taylor ultrafilter \mathcal{U} over Fin_k the following holds:

- (1) For each $1 \leq i \leq k$, $\mathcal{R}_{i,\min}$ and $\mathcal{R}_{i,\max}$ are nnc and for i < k, $\mathcal{R}_{i,\min}$ and $\mathcal{R}_{i,\max}$ are nnc $\Phi_{\geq i+1}(\mathcal{U})$.
- (2) The projections $\min_1(\mathcal{U}), \ldots, \min_k(\mathcal{U}), \max_k(\mathcal{U}), \ldots, \max_1(\mathcal{U})$ are pairwise non-nearly coherent Ramsey ultrafilters over ω .

(3) All cores $\Phi_i(\mathcal{U})$ are nearly coherent.

Proof. In [7, Proposition 3.9] it is shown that all minima and all maxima are Ramsey ultrafilters. It remains to show that they are pairwise non-nearly coherent.

We prove (1). In the first step we consider the case of $\min_j(\mathcal{U})$, $\max_j(\mathcal{U})$. Almost verbatim as in [9, Section 6] it is shown that for any $j \in \{1, \ldots, k\}$, the two Ramsey ultrafilters $\min_j(\mathcal{U})$ and $\max_j(\mathcal{U})$ are not nearly coherent. The reasoning uses only the closure of $\mathrm{TFU}_k(\bar{b})$ under + from Def. 5.1(7) and does not refer to the Tetris function.

We show that for any $1 \leq i < \leq k$, $\min_i(\mathcal{U})$ and $\Phi_{\geq i+1}(\mathcal{U})$ are not nearly coherent. The crucial step is the Tetris step that does not exist for the Hindman space. For a contradiction, we assume that f is finite-to-one and $f(\min_i(\mathcal{U})) \supseteq f(\Phi_{\geq i+1}(\mathcal{U}))$. W.l.o.g. we assume that f is non-decreasing and surjective. We let $I_n = f^{-1}[\{n\}]$. We colour Fin_k as follows. Let c(s) = 0 if the k such that $\min(s^{-1}[\{i\}]) \in I_k$ is equal to the k' such that $s^{-1}[\{i+1,\ldots k\}]$ meets $I_{k'}$; and 1 otherwise. We choose a c-monochromatic $\operatorname{TFU}_k(\bar{a}) \in \mathcal{U}$. We show that the colour is 1. For any r < t, $r, t \in \operatorname{range}(\bar{a})$ whose supports do not meet the same I-interval we have for any $j \geq i+1$ that $j \notin \operatorname{range}(T^{(k-i)}(r))$ but $i \in \operatorname{range}(T^{(k-i)}(r))$ and hence

$$c(T^{(k-i)}(r) + t) = 1.$$

However, by the assumed near coherence there are infinitely many n such that

$$I_n\cap \{\min(a_m^{-1}[\{i\}])\,:\, m\in\omega\}\cap (\bigcup\{a_\ell^{-1}[\{i+1,\ldots,k\}]\,:\, \ell\in\omega\})\neq\emptyset.$$

We take n, m, ℓ such that

$$\min(a_m^{-1}[\{i\}]) \in I_n, a_\ell^{-1}[\{j\}] \cap I_n \neq \emptyset.$$

Then in case $m < \ell$ we have $c(a_m + a_\ell) = 0$ and in case $m = \ell$ we have $c(a_m) = 0$. Contradiction.

Similarly one shows the statement about $\mathcal{R}_{i,\text{max}}$.

- (2): Statement (1) implies (2), since for j > i, $\min_{i}(\mathcal{U})$, $\max_{i}(\mathcal{U}) \supseteq \Phi_{\geq i+1}(\mathcal{U})$.
- (3) Now we consider the cores. We let $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ such that $\operatorname{TFU}_k(\bar{a}) \in \mathcal{U}$. Then there is some finite-to-one h such that

$$(\forall n)(h[a_n^{-1}[\{1\}]] = \dots = h[a_n^{-1}[\{k\}]]).$$

Any set in \mathcal{U} is of the form $TFU_k(\bar{b})$ for some \bar{b} compatible with \bar{a} , we can assume $\bar{b} \sqsubseteq_k^{\text{Tetr}} \bar{a}$. Then by the definition of the condensation order, in each block of \bar{b} there must be one block of \bar{a} that does not drop by the Tetris function. Hence

$$(\forall n)(\forall i = 1, \dots, k)(h[b_n^{-1}[\{i\}] \cap h[b_n^{-1}[\{j\}] \neq \emptyset).$$

Thus $\Phi_i(\mathcal{U})$ and $\Phi_i(\mathcal{U})$ are nearly coherent non-meagre filters.

7. Localised Gowers–Matet Forcing

Now we harvest a fruit from the Ramsey theoretic work: a new family of forcing orders that allows us to arrange the number of near-coherence classes. Since we use generalised condensations by including the Tetris operation, we call the new forcing Gowers–Matet forcing.

Definition 7.1. Let $k \geq 1$. Conditions in Gowers-Matet forcing, \mathbb{GM}_k , are pairs (s, \bar{a}) such that $s \in \operatorname{Fin}_k$ and $\bar{a} \in (\operatorname{Fin}_k)^{\omega}$ and $\operatorname{supp}(s) < \operatorname{supp}(a_0)$. Let $p = (s, \bar{a}) \in \mathbb{GM}_k$. The component s is called the trunk of p and the component \bar{a} is called the pure part of p.

The forcing order is $(t, \bar{b}) \leq_k (s, \bar{a})$ (recall the stronger condition is the smaller one) if the following holds:

(a) s=t or there are $n, i_0 < \cdots < i_n \in \omega$, and $j_r \in \{0, \ldots, k-1\}$ for $r \leq n$ with at least one $j_r = 0$ such that t is a sum of the form

$$t = s + T^{(j_0)}(a_{i_0}) + \dots + T^{(j_n)}(a_{i_n}).$$

(b) $\bar{b} \sqsubseteq_k^{\text{Tetr}} \bar{a}$ (see Def. 6.1(6)).

In the case of s = t we call (t, \bar{b}) a pure extension of $(s, \bar{a}) = p$.

Definition 7.2. Given a Gowers–Matet-adequate family $\mathcal{H} \subseteq (\operatorname{Fin}_k)^{\omega}$, the notion of forcing $\mathbb{GM}_k(\mathcal{H})$ consists of all pairs (s, \bar{a}) such that $\bar{a} \in \mathcal{H}$. The forcing order is the same as in the Gowers–Matet forcing.

Definition 7.3. For $\mathcal{H} = (\operatorname{Fin}_k)^{\omega}(\bar{\mathcal{R}})$ from the previous section we let $\mathbb{GM}_k(\bar{\mathcal{R}}) = \mathbb{GM}_k(\mathcal{H})$.

Thanks to Theorem 6.15 we have:

Lemma 7.4. $\mathbb{GM}_k(\bar{\mathcal{R}})$ has the pure decision property, i.e., for any $\varphi \in \mathcal{L}(\in)$, $(s, \bar{a}) \in \mathbb{GM}_k(\bar{\mathcal{R}})$ there is $(s, \bar{b}) \leq_0 (s, \bar{a})$ such that $((s, \bar{b}) \Vdash \varphi \lor (s, \bar{b}) \Vdash \neg \varphi)$.

Lemma 7.5. The forcing poset $(\mathbb{GM}_k(\bar{\mathcal{R}}), \leq, (\leq_n)_{n < \omega}))$ fulfils Axiom A and hence is proper.

Proof. A derivation of Axiom A for the relations $(\leq_n)_n$ from the pure decision property for tree forcings can be found, e.g., in [2, Section 7.1]. Properness alone can also be proved as in [11, 2.3, 2.4, 2.5]. The latter proof uses the existence of lower bounds, i.e., statement (2) of Theorem 6.15, but not the pure decision property.

Definition 7.6. Let \mathcal{H} be a subset of $(\operatorname{Fin}_k)^{\omega}$.

(1) The $\mathbb{GM}_k(\mathcal{H})$ -generic function from ω to k+1 is

$$\mu = \bigcup \{s \upharpoonright (\max(\operatorname{supp}(s)) + 1) \, : \, (\exists \bar{a}) ((s, \bar{a}) \in G)\}.$$

(2) Again the generic *i*-fibres for i = 1, ..., k are written as

$$\mu_i := \mu^{-1}[\{i\}].$$

(3) Any Gowers-Milliken-Taylor ultrafilter \mathcal{U} avoids \mathcal{E} iff $\Phi(\mathcal{U}) \not\leq_{RB} \mathcal{E}$.

Now we investigate the number of near-coherence classes in models our our new forcing.

Definition 7.7. Let $X \in [\omega]^{\omega}$. We let $f_X(n) = |X \cap n|$.

We start with a density argument for evaluating our forcings.

Theorem 7.8. Assume CH. Let $\overline{\mathcal{R}}$ be as in Def. 6.8. Let $h: \omega \to \omega$ be a finite-to-one function. Let \mathcal{E} and \mathcal{W} be ultrafilters over ω such that \mathcal{W} , $\mathcal{E} \not\geq_{RB} \mathcal{R}_{i,x}$ for $i = 1, \ldots, k$, $x = \min, \max$. Then

$$\mathbb{GM}_k(\bar{\mathcal{R}}) \Vdash_{\mathbb{GM}_k(\bar{\mathcal{R}})} f_{\operatorname{supp}(\mu)}(\mathcal{W}) = f_{\operatorname{supp}(\mu)}(\mathcal{E}).$$

Proof. Suppose for some $E \in \mathcal{E}$, $W \in \mathcal{W}$, some $(s,\bar{a}) \in \mathbb{GM}_k(\bar{\mathcal{R}})$, $(s,\bar{a}) \vdash f_{\text{supp}(\mu)}[W] \cap f_{\text{supp}(\mu)}[E] \cap [n,\omega) = \emptyset$. By Theorem 6.15(3), there is a Gowers–Milliken–Taylor ultrafilter \mathcal{U}_k such that $\text{TFU}_k(\bar{a}) \in \mathcal{U}$ and such that $\Phi(\mathcal{U}_k)$ avoids \mathcal{E}, \mathcal{W} .

So there is a $\bar{b} \sqsubseteq_k^{\text{Tetr}} \bar{a}$ such that $\text{TFU}(\bar{b}) \in \mathcal{U}_k$ and such that there are infinitely many elements of E outside $\text{set}(\bar{b})$ and infinitely many elements of W outside $\text{set}(\bar{b})$. Since $\text{TFU}_k(\bar{b}) \in \mathcal{U}_k$, Lemma 6.18 implies that $\bar{b} \in (\text{Fin}_k)^{\omega}(\bar{\mathcal{R}})$. We take some $e \in (E \setminus \text{set}(\bar{b}))$ such that $e > \max\{n, \max(\text{supp}(s))\}$. and some $w \in W \setminus \text{set}(\bar{b})$ such that $w > \max\{n, \max(\text{supp}(s))\}$. Then we can lengthen s by taking t > s such that

$$\min(\operatorname{supp}(t)) > w, e \text{ and } (s+t,\bar{b}) \leq_{\mathbb{GM}_k(\bar{\mathcal{R}})} (s,\bar{a}). \text{ Then } (s+t,\bar{b}) \Vdash f_{\operatorname{supp}(\mu)}(e) = f_{\operatorname{supp}(\mu)}(w). \text{ Contradiction.}$$

Theorem 7.9. (Adaption of [17, Theorem 4]) Let $k \geq 1$ and $\bar{\mathcal{R}}$ be as above and assume that \mathcal{E} is a P-point with $\mathcal{E} \not\geq_{RB} \mathcal{R}_{i,\min}, \mathcal{R}_{j,\max}$ for any $i \in \{1, \dots, k\}$. Then \mathcal{E} continues to generate an ultrafilter after we force with $\mathbb{GM}_k(\bar{\mathcal{R}})$.

Proof. By Theorem 6.15(2) the conditions (s, \bar{a}) such that there is a Gowers–Milliken–Taylor ultrafilter $\Phi(\mathcal{U}_k)$ is nnc to \mathcal{E} (which means $\Phi(\mathcal{U}_k) \not\leq_{rmRB} \mathcal{E}$) and such that $\bar{a} \in \mathcal{U}_k$ are dense in $\mathbb{GM}_k(\bar{\mathcal{R}})$. Since $\mathbb{GM}_k(\bar{\mathcal{R}})$ has the pure decision property we can carry out the proof literally as in [17, Lemma 2.3 – Lemma 2.10].

Now we are concerned with the second iterand. The following follows from an easy density argument.

Lemma 7.10. For any
$$i \in \{1, ..., k\}$$
, $x = \min, \max$, we have

$$\mathbb{GM}_k(\bar{\mathcal{R}}) \Vdash \mathcal{R}_{i,x} \cup \{\mu_i\} \text{ is a filter subbase.}$$

Now we come to the k-dimensional version of Theorem 3.25. It is crucial that the Ramsey ultrafilter $\mathcal{R}_{i,x}$ is not diagonalised. Otherwise we would add a dominating real. For each $1 \leq i \leq k$ the two Ramsey ultrafilters $\mathcal{R}_{\min,i}$ and $\mathcal{R}_{\max,i}$ are destroyed by the i-fiber μ_i .

Theorem 7.11. Let k, $\bar{\mathcal{R}}$ be as in Def. 6.8.

(7.1)
$$\mathbb{GM}_{k}(\bar{\mathcal{R}}) \Vdash (\forall i \in \{1, \dots, k\}, x \in \{\min, \max\})$$
$$(\text{fil}((\mathcal{R}_{i,x} \cup \{\mu_{i}\}))^{+} \text{ is a happy family}$$
$$\text{that is nowhere almost a filter}).$$

and hence by Mathias' [30, Prop. 011] combined with [5, proof of Theorem 14],

$$\mathbb{GM}_k(\bar{\mathcal{R}}) \Vdash (\exists \langle \mathcal{R}_{i,x}^{\text{ext}} : i = 1, \dots, k, x = \min, \max \rangle) \Big((\forall (i,x)) \Big(\mathcal{R}_{i,x}^{\text{ext}} \supseteq (\mathcal{R}_{i,x} \cup \{\mu_i\}) \land (\mathcal{R}_{i,x}^{\text{ext}} \supseteq (\mathcal{$$

 $\mathcal{R}_{i,x}^{\mathrm{ext}}$ is a Ramsey ultrafilter that is nnc to $\mathcal{E} \big)$

and the extensions are pairwise nnc).

Proof. Theorem 6.16 allows us to transfer the proof of Theorem 3.25 to $\mathbb{GM}_k(\bar{\mathcal{R}})$.

We rework Section 4, the iteration theory for limit steps, for iterands of the form $\mathbb{GM}_k(\bar{\mathcal{R}})$. We fix k and $\bar{\mathcal{R}} = \bar{\mathcal{R}}_0$ as in Def. 6.8. We define by induction on $\alpha \leq \omega_2$ a countable support iteration (in the sense of [37, Def. III, 3.1]) $\mathbb{P}_{\alpha} = \langle \mathbb{P}_{\beta}, \mathbb{GM}(\bar{\mathcal{R}}_{\gamma}) : \beta \leq \alpha, \gamma < \alpha \rangle$ such that for any $\beta < \alpha$,

(7.2)
$$\mathbb{P}_{\beta} \Vdash (\forall (i,x)) \big(\mathcal{R}_{i,x,\beta} \supseteq \bigcup \{ \mathcal{R}_{i,x,\gamma} \cup \{ \mu_{\gamma,i} \} : \gamma < \beta \}$$

$$\wedge \mathcal{R}_{i,x,\beta} \text{ is a Ramsey ultrafilter that is nnc to } \mathcal{E} \big)$$
and the $\mathcal{R}_{i,x,\beta}$, $i = 1, \dots, k$, $x = \min, \max$, are pairwise nnc.

We use for names the same letters as for the corresponding evaluated names.

Here $\mathbb{P}_{\beta+1}$ forces that μ_{β} is the $\mathbb{GM}_k(\overline{\mathcal{R}}_{\beta})$ -generic real, and $\mu_{\beta,i}$ denotes its *i*-fibre according to Definition 5.8(8). In Theorem 7.11 we proved that there are extensions in the successor steps. This guarantees the existence of $\overline{\mathcal{R}}_{\beta+1}$ with the desired properties.

Now we consider limit steps α . If $\operatorname{cf}(\alpha) > \omega$, we can just take $\mathbb{P}_{\alpha} \Vdash \mathcal{R}_{i,x,\alpha} = \bigcup_{\gamma < \alpha} \mathcal{R}_{i,x,\gamma}$ and the inductive hypotheses will be carried on, since in proper forcing every real appears at a step of at most countable cofinality, with the only exception that for $\alpha = \aleph_2$ the CH gets lost. So we concentrate on the hard case, $\operatorname{cf}(\alpha) = \omega$.

(7.3)
$$\mathbb{P}_{\alpha} \Vdash (\exists \bar{\mathcal{R}}_{\alpha}) (\bigwedge_{i,x} \mathcal{R}_{i,x,\alpha} \supseteq \bigcup \{\mathcal{R}_{i,x,\beta} \cup \{\mu_{\beta,i}\}) : \beta < \alpha\}$$
 and the $\mathcal{R}_{i,x,\alpha}$, $i = 1, \dots, k$, $x = \min$, max, are pairwise nnc Ramsey ultrafilters that are nnc to \mathcal{E}),

will follow from the CH, a routine enumeration according to [30, Prop. 011] and [5, proof of Theorem 14] the following theorem:

Theorem 7.12. Suppose CH, $\alpha < \omega_2$, $\operatorname{cf}(\alpha) = \omega$, and that $\mathbb{P}_{\beta}, \overline{\mathcal{R}}_{\beta}$ are as in Equation (7.2) and \mathbb{P}_{α} is the countable support limit of $\langle \mathbb{P}_{\beta}, \mathbb{GM}_k(\overline{\mathcal{R}}_{\beta}) : \beta < \alpha \rangle$. In $\mathbf{V}^{\mathbb{P}_{\alpha}}$, for any i, x, the set of positive sets

$$\left(\bigcup_{\gamma<\alpha}(\mathcal{R}_{i,x,\gamma}\cup\{\mu_{\gamma,i}\})\right)^+$$

forms a happy family that is nowhere almost a filter.

As in the proofs of Theorem 4.1 we introduce an increasing sequence $\langle R_{\gamma}(\bar{\mathcal{R}}_{<\gamma}) : \gamma < \alpha \rangle$ of relations $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ in $\mathbf{V}^{\mathbb{P}_{\alpha}}$ such that a property called

(7.4) "
$$\mathbb{P}_{\alpha}$$
 is $R_{\alpha}(\bar{\mathcal{R}}_{\leq \alpha})$ -preserving"

is carried in addition to the property (7.2) and properness in the inductive choice of the iteration.

We define a relation $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ and are concerned with statements of the form $(\forall f)(\exists g)(fR_{\alpha}(\bar{\mathcal{R}}_{<\alpha})g)$ for a Borel relation $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ on the Baire space in $\mathbf{V}^{\mathbb{P}_{\alpha}}$. Here the argument $\bar{\mathcal{R}}_{<\alpha}$ indicates the invoked parameters from the ground model: Ramsey ultrafilters in the stage 0 and names for Ramsey ultrafilters in any later stage.

We do not write tildes below the $R_{\alpha}(\bar{\mathcal{R}}_{\leq \alpha})$'s, although these relations are names.

Definition 7.13. By induction on $\alpha \leq \omega_2$ we define the following relations.

- (1) Let $\mathbb{P}_{\alpha} = \langle \mathbb{P}_{\beta}, \mathbb{GM}_{k}(\bar{\mathcal{R}}_{\gamma}) : \beta \leq \alpha, \gamma < \alpha \rangle$ be defined with (7.2). Let $i \in \{1, \ldots, k\}, x \in \{\min, \max\}$. We say that a \mathbb{P}_{α} -name g for an element of $[\omega]^{\omega}$ is $\alpha \cdot (i, x)$ -positive if $1 \Vdash_{\mathbb{P}_{\alpha}} g \in (\bigcup \{\text{fil}(\mathcal{R}_{i,x,\gamma} \cup \{\mu_{\gamma,i}\}) : \gamma < \alpha\})^{+}$.
- (2) Let $\mathbb{P}_{\alpha} = \langle \mathbb{P}_{\beta}, \mathbb{GM}_{k}(\bar{\mathcal{R}}_{\gamma}) : \beta \leq \alpha, \gamma < \alpha \rangle$ be defined with (7.2). We say $fR_{\alpha}(\bar{\mathcal{R}}_{\leq \alpha})g$ if the following holds in $\mathbf{V}^{\mathbb{P}_{\alpha}}$:
 - (a) $f = (\bar{A}, h, (i, x)),$
 - (b) $\bar{A} = \langle A_{\ell} : \ell \in \omega \rangle$ is a \subseteq -descending sequence of α -(i, x)-positive members of $[\omega]^{\omega}$,
 - (c) h is finite-to-one,
 - (d) for j = 0, 1 there are $g^j \in [\omega]^{\omega}$, with the following properties:
 - (i) $g = g^0 \cup g^1$,
 - (ii) g^j is an α -(i, x)-positive diagonal lower bound of \bar{A} and
 - (iii) $h[g^0] \cap h[g^1] = \emptyset$.

Definition 7.14. We say \mathbb{P}_{α} is $R_{\alpha}(\bar{\mathcal{R}}_{\leq \alpha})$ -preserving if \mathbb{P}_{α} is proper and

$$\mathbb{P}_{\alpha} \Vdash (\forall f \in \text{dom}(R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})))(\exists g)(fR_{\alpha}(\bar{\mathcal{R}}_{<\alpha})g).$$

Lemma 7.15. We assume CH, $\alpha < \omega_2$ and $\operatorname{cf}(\alpha) = \omega$. If $\mathbb{P}_{\alpha} = \langle \mathbb{P}_{\beta}, \mathbb{GM}_k(\bar{\mathcal{R}}_{\gamma}) : \beta \leq \alpha, \gamma < \alpha \rangle$ is $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ preserving then

$$\mathbb{P}_{\alpha} \Vdash (\bigcup \{ \text{fil}(\mathcal{R}_{i,x,\beta} \cup \{\mu_{i,\beta}\}) : \beta < \alpha \})^{+}$$

is a happy family that is nowhere almost a filter).

Proof. This follows from Definition 7.14.

As mentioned a routine enumeration gives:

Corollary 7.16. If $\mathbb{P}_{\alpha} \Vdash \mathsf{CH}$ and \mathbb{P}_{α} is $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ -preserving then Equation (7.3) holds.

We carry the preservation property upwards by induction.

Lemma 7.17. Assume CH, $\alpha < \omega_2$, $\operatorname{cf}(\alpha) = \omega$. Let \mathbb{P}_{α} be the countable support limit of \mathbb{P}_{β} , $\beta < \alpha$. If for $\beta < \alpha$ such that $\operatorname{cf}(\beta) < \omega_1$, \mathbb{P}_{β} is $R_{\beta}(\bar{\mathcal{R}}_{<\beta})$ -preserving and for any $\beta < \alpha$, (7.2) holds then \mathbb{P}_{α} is $R_{\alpha}(\bar{\mathcal{R}}_{<\alpha})$ preserving.

Proof. As in Lemma 4.6.

This finishes the proof of Theorem 7.12.

Finally we answer Banakh's and Blass' question on the finite part near-coherence spectrum:

Theorem 7.18. Assume CH and let $k \geq 1$. Then there is a countable support iteration of proper iterands $\mathbb{P} = \langle \mathbb{P}_{\alpha}, \mathbb{GM}_k(\bar{\mathcal{R}}_{\beta}) : \beta < \omega_2, \alpha \leq \omega_2 \rangle$ such that in the extension there exactly 2k+1 near-coherence classes of ultrafilters. Namely, one class is represented by a P-point of character ω_1 and 2k classes represented by the names Ramsey ultrafilters $\mathcal{R}_{i,x,\omega_2}$, $i=1,\ldots,k$, $x=\min,\max$.

Proof. We start with a ground model V that fulfils CH. By CH there are a P-point \mathcal{E} and a $\{1, \ldots, k\} \times \{\min, \max\}$ -sequence of Ramsey ultrafilters $\bar{\mathcal{R}}_0$, such that any two

of them are nnc. A countable support iteration $\langle \mathbb{P}_{\beta}, \mathbb{GM}_{k}(\overline{\mathcal{R}}_{\gamma}) : \beta \leq \omega_{2}, \gamma < \omega_{2} \rangle$ of proper forcings with (7.2) at any stage β forces: There are at least the near-coherence classes of \mathcal{E} , $\mathcal{R}_{i,x,\omega_{2}}$, $i = 1, \ldots, k$, $x = \min, \max$.

Why are there no other near-coherence classes in $\mathbf{V}^{\mathbb{P}_{\omega_2}}$? For a contradiction, we suppose that \mathcal{W} is a non-principal ultrafilter in $\mathbf{V}[G]$ for some \mathbb{P}_{ω_2} -generic filter G and \mathcal{W} is nnc to any $\mathcal{R}_{i,x\omega_2}$, $i=1,\ldots,k$, $x=\min$, max and nnc to \mathcal{E} . Then by [11, Lemma 5.10] there are a $p \in \mathbb{P}_{\omega_2} \cap G$, say $p \in \mathbb{P}_{\alpha_0}$ for an $\alpha_0 \in \omega_2$ (by [11, Lemma 5.6]), and an ω_1 -club $C \subseteq [\alpha_0, \omega_2)$ such that for each $\alpha \in C \cup \{\omega_2\}$, $\mathcal{W} \cap \mathbf{V}[G \cap \mathbb{P}_{\alpha}]$ has a \mathbb{P}_{α} -name (for which we again write $\mathcal{W} \cap \mathbf{V}[G \cap \mathbb{P}_{\alpha}]$) and

 $p \Vdash_{\mathbb{P}_{\alpha}} \mathcal{W} \cap \mathbf{V}[G \cap \mathbb{P}_{\alpha}]$ is an ultrafilter that is nnc to $\mathcal{R}_{i,x,\alpha}$ and nnc to \mathcal{E} .

We fix $\alpha = \min(C)$.

By Theorem 7.8 and Theorem 7.9 we have for any $\alpha \in C$,

 $p \Vdash_{\mathbb{P}_{\alpha+1}} f_{\operatorname{supp}(\mu_{\alpha})}(\mathcal{W} \cap \mathbf{V}[G \cap \mathbb{P}_{\alpha}]) = f_{\operatorname{supp}(\mu_{\alpha})}(\mathcal{E}) \wedge \mathcal{E} \text{ generates an ultrafilter.}$

Since

$$p \Vdash_{\mathbb{P}_{\omega_2}} \mathcal{W}$$
 is a filter and $\mathcal{W} \supseteq \mathcal{W} \cap \mathbf{V}[G \cap \mathbb{P}_{\alpha}]$,

we have $p \Vdash_{\mathbb{P}_{\omega_2}} f_{\operatorname{supp}(\mu_{\alpha})}(\mathcal{W}) = f_{\operatorname{supp}(\mu_{\alpha})}(\mathcal{E}).$

Theorem 7.19. Under CH there is an iteration $\langle \mathbb{P}_{\beta}, \mathbb{Q}_{\alpha} : \beta \leq \omega_2, \alpha < \omega_2 \rangle$ that forces exactly 2k near-coherence classes.

Proof. We fix a $\Diamond(S_{\aleph_1}^{\aleph_2})$ -sequence $\langle D_{\alpha} : \alpha \in S_{\aleph_1}^{\aleph_2} \rangle$ in the ground model or add one by a preliminary forcing that preserves CH. At iterand $\mathbb{Q}_{\alpha} = \mathbb{GM}_k(\langle \mathcal{R}_{i,x,\alpha} : \mathcal{R}_{i,x,\alpha} : \mathcal{R}_{i,x,\alpha})$ $i=1,\ldots,k,x=\min,\max\rangle$, $cf(\alpha)=\omega_1$, we work with only one top ultrafilter $\mathcal{R}_{k,\min,\alpha}$ that extends the ultrafilters $\mathcal{R}_{k,\min,\beta}$, $\beta < \alpha$. The other top ultrafilter $\mathcal{R}_{k,\max,\alpha}$ is varied according to a $\diamondsuit(S_{\aleph_1}^{\aleph_2})$ sequence, namely to be some Ramsey ultrafilter that is (forced to be) nnc to D_{α} that is handed down by the diamond, if the diamond hands down a name of an ultrafilter D_{α} that is forced to be nnc to \mathcal{E} and nnc to any of the $\mathcal{R}_{i,x,\alpha}$, $i=1,\ldots,k-1$, $x=\min,\max$ or $(i,x)=(k,\min)^{10}$. According to Theorem 7.8 in stage $\mathbf{V}^{\mathbb{P}_{\alpha+1}}$ we have that D_{α} is nearly coherent to \mathcal{E} . If $\mathcal{W} \in \mathbf{V}^{\mathbb{P}_{\omega_2}}$ extends D_{α} , then in the final model \mathcal{W} is nearly coherent to \mathcal{E} via $f_{\mu_{\alpha}}$. At stages α of uncountable cofinality and below conditions where the diamond does not hand down an ultrafilter that is nnc to \mathcal{E} and nnc to any of the $\mathcal{R}_{i,x,\alpha}$, $i=1,\ldots,k-1$, $x=\min,\max$ or $(i,x)=(k,\min)$, or at stages of countable cofinality or successor stages we choose for $\mathcal{R}_{k,\max,\alpha}$ any Ramsey ultrafilter that is nnc to any of the $\mathcal{R}_{i,x,\alpha}$, $i=1,\ldots,k-1$, $x=\min,\max$ or $(i,x)=(k,\min)$ and nnc to \mathcal{E} . Since for any \mathcal{P}_{ω_2} -name τ for an ultrafilter, there are stationarily many $\alpha \in S_{\aleph_1}^{\aleph_2}$ such that $\tau \cap \mathbf{V}^{\mathbb{P}_{\alpha}} = D_{\alpha}$, any ultrafilter \mathcal{W} in \mathbb{P}_{ω_2} extension is nearly coherent to \mathcal{E} or nearly coherent to some of the $\mathcal{R}_{i,x,\omega_2}$, $i=1,\ldots,k-1$, $x = \min, \max \text{ or } (i, x) = (k, \min).$

Corollary 7.20. For any $n \in \omega$, the statement "there are exactly n+1 near-coherence classes of ultrafilters" is consistent relative to ZFC.

¹⁰Details about book-keeping functions that are defined from diamond sequences can for example be found in [31, Section 2].

For n = 1 we can take the Matet model without any localisation [8] or the Miller model [12] or the Blass–Shelah model [11].

The following two propositions give more information on the decompositions of the iterands.

Proposition 7.21. Fix k, $\bar{\mathcal{R}}$ as Definition 6.8. We let $\mathbb{Q}_{pure} = (\operatorname{Fin}_k^{\omega}(\bar{\mathcal{R}}), \sqsubseteq_k^{\operatorname{Tetr},*})$ and we let $\mathcal{U} = \{\langle \bar{a}, \check{a} \rangle : \bar{a} \in \mathbb{Q}_{pure} \}$. Then the following holds:

- (1) \mathbb{Q}_{pure} is ω -closed.
- (2) $\mathbb{GM}_k(\bar{\mathcal{R}})$ is densely embedded into $\mathbb{Q}_{pure} * \mathbb{GM}_k(\mathcal{U})$.
- (3) \mathbb{Q}_{pure} forces that \mathcal{U} is a Gowers-Milliken-Taylor ultrafilter with $\min_i(\mathcal{U}) = \mathcal{R}_{i,\min}$ and $\max_i(\mathcal{U}) = \mathcal{R}_{i,\max}$ for $i = 1, \dots, k$.
- (4) \mathbb{Q}_{pure} forces that $\Phi(\mathcal{U})$ (Φ is defined in Def. 5.8(3)) is nnc to any filter from the ground model that is nnc $\mathcal{R}_{i,x}$. $i=1,\ldots,k$, $x=\min,\max$.
- *Proof.* (1) The forcing order \mathbb{Q}_{pure} is ω -closed by (2) of Theorem 6.15.
- (2) We map $(s, \bar{a}) \in \mathbb{GM}_k(\bar{\mathcal{R}})$ to $(\bar{a}, (s, \bar{a}))$. This is a dense embedding. Details can be found in [17, Proposition 3.2].
- (3) By Theorem 6.15 and density arguments, the first forcing, \mathbb{Q}_{pure} , forces that is generic filter is an Gowers–Milliken–Taylor ultrafilter. Since \mathbb{Q}_{pure} does not add reals, only colourings and descending ω -sequences from the ground model have to be considered.

Statement (4) follows from Theorem 7.8, applied to \mathbb{Q}_{pure} .

Remark 7.22. We sketch an alternative way to force at least 2k + 1 nameable near-cohernce classes: Force with an iteration of iterands $\mathbb{GM}_k(\mathcal{U}_{\alpha})$, \mathcal{U}_{α} a Gowers-Milliken-Taylor ultrafilter in $\mathbf{V}^{\mathbb{P}_{\alpha}}$ without localisation to $\overline{\mathcal{R}}$. In the successor steps one needs to prove the analogon to Theorem 3.9 about completing the destroyed Gowers-Milliken-Taylor ultrafilter \mathcal{U}_{α} , in a non-localised form. So here for the existence of $\mathcal{U}_{\alpha+1} \supseteq \mathcal{U}_{\alpha}$ we need Gower's theorem in a computation about names. We expect that this can be done similarly to the proofs in Section 3. However, there is no way to prevent that more nnc Ramsey ultrafilters than the projections are created in the iteration, since the iteration could just be the Fin_k -shadow of an iteration with Gowers-Milliken-Taylor ultrafilters over Fin_{k+1} .

References

- Taras Banakh and Andreas Blass. The number of near-coherence classes of ultrafilters is either finite or 2^c. In Joan Bagaria and Stevo Todorcevic, editors, Set Theory, Trends in Mathematics, pages 257–273. Birkhäuser, 2006.
- [2] Tomek Bartoszyński and Haim Judah. Set Theory, On the Structure of the Real Line. A K Peters, 1995.
- [3] James Baumgartner. A short proof of Hindman's theorem. J. Comb. Theory (A), 17:384–386, 1974
- [4] Andreas Blass. The Rudin-Keisler ordering of P-points. Trans. Amer. Math. Soc., 179:145–166, 1973.
- [5] Andreas Blass. Near coherence of filters, I: Cofinal equivalence of models of arithmetic. Notre Dame J. Formal Logic, 27:579–591, 1986.
- [6] Andreas Blass. Near coherence of filters, II: Applications to operator ideals, the Stone-Čech Remainder of a half-line, order ideals of sequences, and slenderness of groups. Trans. Amer. Math. Soc., 300:557–581, 1987.

- [7] Andreas Blass. Ultrafilters related to Hindman's finite unions theorem and its extensions. In Stephen Simpson, editor, Logic and Combinatorics, Arcata, California, 1985, volume 65 of Contemp. Math., pages 89–124. Amer. Math. Soc., 1987.
- [8] Andreas Blass. Applications of superperfect forcing and its relatives. In Juris Steprāns and Steve Watson, editors, Set Theory and its Applications, volume 1401 of Lecture Notes in Mathematics, pages 18–40, 1989.
- [9] Andreas Blass. Homogeneous sets from several ultrafilters. Topology Appl., 156:2581–2594, 2009.
- [10] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In Matthew Foreman and Akihiro Kanamori, editors, Handbook of set theory. Vols. 1, 2, 3, pages 395–489, Dordrecht, 2010. Springer.
- [11] Andreas Blass and Saharon Shelah. There may be simple P_{\aleph_1} and P_{\aleph_2} -points and the Rudin-Keisler ordering may be downward directed. Annals of Pure and Applied Logic, 33:213–243, 1987.
- [12] Andreas Blass and Saharon Shelah. Near coherence of filters III: A simplified consistency proof. Notre Dame J. Formal Logic, 30:530–538, 1989.
- [13] Andreas Blass and Gary Weiss. A characterization and sum decomposition of operator ideals. Trans. Amer. Math. Soc., 246:407 – 417, 1978.
- [14] David Booth. Ultrafilters on a countable set. Ann. Math. Logic, 2:1-24, 1970.
- [15] R. Michael Canjar. On the generic existence of special ultrafilters. Proc. Amer. Math. Soc., 110:233–241, 1990.
- [16] David Chodounsky and Osvaldo Guzman. There are not p-points in the Silver model. Israel J. Math., 2019, to appear.
- [17] Todd Eisworth. Forcing and stable ordered-union ultrafilters. J. Symbolic Logic, 67:449–464, 2002.
- [18] Robert Ellis. Distal transformation groups. Pacific J. Math., 8:401-405, 1958.
- [19] David Fernández-Bretón. cov(M) and stable ordered union-ultrafilters. J. Symbolic Logic, 2019, to appear.
- [20] David Fernández-Bretón and Michael Hrušák. A parametrized diamond principle and union ultrafilters. Colloq. Math., 153(2):261–271, 2018.
- [21] Martin Goldstern. Tools for your forcing construction. In Set theory of the reals (Ramat Gan, 1991), volume 6 of Israel Math. Conf. Proc., pages 305–360. Bar-Ilan Univ., Ramat Gan, 1993.
- [22] W. T. Gowers. Lipschitz functions on classical spaces. European J. Combin., 13(3):141–151, 1992
- [23] Serge Grigorieff. Combinatorics of ideals and forcing. Ann. Math. Logic, 3:363-394, 1971.
- [24] Neil Hindman. Finite sums from sequences within cells of a partition of N. J. Combin. Theory Ser. A, 17:1–11, 1974.
- [25] Neil Hindman. Ultrafilters and combinatorial number theory. In Melvyn B. Nathanson, editor, Number theory, Carbondale 1979 (Proc. Southern Illinois Conf., Southern Illinois Univ., Carbondale, Ill., 1979), volume 751 of Lecture Notes in Math., pages 119–184. Springer, Berlin, 1979.
- [26] Neil Hindman and Dona Strauss. Algebra in the Stone-Čech Compactification. De Gruyter,
- [27] Kenneth Kunen. Some points in βN . Math. Proc. Cambridge Philos. Soc., 80(3):385–398, 1976.
- [28] Alain Louveau. Une méthode topologique pour l'étude de la propriété de Ramsey. Israel J. Math., 23(2):97-116, 1976.
- [29] Pierre Matet. Partitions and filters. J. Symbolic Logic, 51:12–21, 1986.
- [30] Adrian Mathias. Happy families. Ann, Math. Logic, 12:59–111, 1977.
- [31] Heike Mildenberger, Saharon Shelah, and Boaz Tsaban. Covering the Baire space with meager sets. Ann. Pure Appl. Logic, 140:60–71, 2006.
- [32] Arnold Miller. There are no Q-points in Laver's model for the Borel conjecture. Proc. Amer. Math. Soc., 78:103–106, 1980.
- [33] Jerzy Mioduszewski. On composants of $\beta R R$. In J. Flachsmeyer, Z. Frolík, and F. Terpe, editors, *Proceedings Conf. Topology and Measure (Zinnowitz 1974)*, pages 257–283. Ernst-Moritz-Arndt-Universität zu Greifswald, 1978.

- [34] Jerzy Mioduszewski. An approach to $\beta \mathbf{R} \mathbf{R}$. In Á. Császár, editor, *Topology*, volume 23 of *Colloq. Math. Soc. János Bolyai*, pages 853–854. North-Holland, 1980.
- $[35] \ \ Katsui \ \ Namakura. \ On \ bicompact \ semigroups. \ \textit{Math. J. Okayama Univ.}, \ 1:99-108, \ 1952.$
- [36] Saharon Shelah. *Proper forcing*, volume 940 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, xxix+496 pp, 1982.
- [37] Saharon Shelah. Proper and Improper Forcing, 2nd Edition. Springer, 1998.
- [38] Michel Talagrand. Compacts de fonctions mesurables et filtres non mesurables. Studia Mathematicae, 67:13 43, 1980.
- [39] Alan D. Taylor. A canonical partition relation for finite subsets of ω . J. Combinatorial Theory Ser. A, 21(2):137–146, 1976.
- [40] Stevo Todorcevic. Introduction to Ramsey spaces, volume 174 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2010.
- [41] Yuan Yuan Zheng. Selective ultrafilters on FIN. Proc. Amer. Math. Soc., 145(12):5071–5086, 2017.

HEIKE MILDENBERGER, ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT, ABTEILUNG FÜR MATH. LOGIK, ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU, GERMANY

 $E\text{-}mail\ address: \ \texttt{heike.mildenberger@math.uni-freiburg.de}$