arXiv:1404.7690v3 [math.AT] 11 Nov 2019

TWISTED LEFSCHETZ NUMBERS OF INFRA-SOLVMANIFOLDS AND ALGEBRAIC GROUPS

HISASHI KASUYA

Abstract. Twisted Lefschetz numbers are extensions of the ordinary Lefschetz numbers for cohomologies with values in flat bundles. As a generalization of linearization formula for the ordinary Lefschetz number of a diffeomorphism of a nilmanifold, we show that a twisted Lefschetz number of any diffeomorphism of any infra-solvmanifold is equal to the determinant $\det(I-A)$ for some matrix A by using the cohomology of algebraic groups.

1. Introduction

Let M be a compact manifold and E be a flat vector bundle over M. We consider the de Rham cohomology $H^*(M,E)$ with values in E. Let $\varphi : M \to M$ be a smooth map and a morphism $\Xi : \varphi^*E \to E$ of flat bundles where φ^*E is the pull-back of E. Consider the induced map $\Xi \circ H^*(\varphi) : H^*(M,E) \to H^*(M,E)$. We define the twisted Lefschetz number

$$L(\varphi, E, \Xi) = \sum_i (-1)^i \mathrm{trace}(\Xi \circ H^*(\varphi))_{|H^i(M,E)}.$$

As for the ordinary Lefschetz number, we can show the Lefschetz fixed point formula for the twisted Lefschetz number by the Atiyah-Bott theorem ([1]). If φ is a diffeomorphism so that the graph of φ is transversal to the diagonal in $M \times M$, then we have

$$L(\varphi, E, \Xi) = \sum_{\varphi(x)=x} \mathrm{sign}(\det(I - d\varphi_x)) \cdot \mathrm{trace}(\Xi_x).$$

In general, computations of the twisted Lefschetz numbers are harder than computations of usual Lefschetz numbers. In this paper we give a simple formula for twisted Lefschetz numbers of diffeomorphisms of solvmanifolds and infra-solvmanifolds which is a generalization of the "linearization formula" for ordinary Lefschetz numbers of maps of nilmanifolds.

In this paper solvmanifolds (resp. nilmanifolds) are compact homogeneous spaces of connected solvable (resp. nilpotent) Lie groups. We say that a solvmanifold (resp. nilmanifold) M has a discrete presentation (G,Γ) if there exists a simply connected solvable (resp. nilpotent) Lie group G with a cocompact discrete subgroup Γ so that M is diffeomorphic to G/Γ. It is known that every nilmanifold M has a discrete presentation (G,Γ) (see [11]).

Consider a nilmanifold G/Γ for a discrete presentation (G,Γ). For a smooth map $\varphi : G/\Gamma \to G/\Gamma$, since the fundamental group of G/Γ is Γ, we have the map $f : \Gamma \to \Gamma$ induced by φ. It is known that any homomorphism between cocompact discrete subgroups of simply connected nilpotent Lie groups can be extended to a Lie group homomorphism. Hence we have an extension $T : G \to G$ of f. In [14], Nomizu showed

2010 *Mathematics Subject Classification.* 22E25; 20G20 ; 54H25; 55M20 .

Key words and phrases. Topological fixed point theory, cohomology of algebraic group, twisted lefschetz number, infra-solvmanifold.

that the de Rham cohomology of a nilmanifold G/Γ for a discrete presentation (G,Γ) is isomorphic to the Lie algebra cohomology of the Lie algebra $\mathfrak{g}$ of G. Hence, considering the induced homomorphism $T_* : \mathfrak{g} \to \mathfrak{g}$, taking a matrix presentation A of $T_* : \mathfrak{g} \to \mathfrak{g}$, we can get the "linearization formula"

$$L(\varphi) = \det(I - A)$$

where $L(\varphi)$ is the ordinary Lefschetz number.

We are interested in getting a "linearization formula" for solvmanifolds. But there are many difficulties:

- In general, a solvmanifold M does not have a discrete presentation (G,Γ).
- In general, for a simply connected solvable Lie group with a cocompact discrete subgroup Γ, a homomorphism $\Gamma \to \Gamma$ may not extend to a Lie group homomorphism $G \to G$.
- In general, the de Rham cohomology of a solvmanifold G/Γ for a discrete presentation (G,Γ) is not isomorphic to the Lie algebra cohomology of the Lie algebra $\mathfrak{g}$ of G.

Assuming some conditions, we can get a "linearization formula" (see for examples [12] and [10]). However on general solvmanifolds, it seems to be difficult to get a "linearization formula".

In this paper we consider the more general setting. An infra-solvmanifold is a manifold of the form G/Δ, where G is a simply connected solvable Lie group, and Δ is a torsion-free subgroup of $\mathrm{Aut}(G) \ltimes G$ such that the closure of $h(\Delta)$ in $\mathrm{Aut}(G)$ is compact where $h : \mathrm{Aut}(G) \ltimes G \to \mathrm{Aut}(G)$ is the projection. An infra-solvmanifold is a generalization of a solvmanifold. See [2] and [4] for properties of infra-solvmanifolds. The purpose of this paper is to give a "linearization formula" for any diffeomorphism of any infra-solvmanifold by using the twisted Lefschetz number and algebraic groups.

Let M be an infra-solvmanifold with the fundamental group Γ and $\varphi : M \to M$ a diffeomorphism. Denote by $f : \Gamma \to \Gamma$ the homomorphism induced by φ. Then, by methods of algebraic groups given in [16, 2], we can say that $f : \Gamma \to \Gamma$ induces an explicit isomorphism on a certain Lie algebra $\mathfrak{u}_\Gamma$. For a matrix presentation A of such isomorphism, we will prove that by taking some E and Ξ, the formula

$$L(\varphi, E, \Xi) = \det(I - A)$$

holds.

2. Algebraic groups and representations

An algebraic group $\mathcal{G}$ is a Zariski-closed subgroup of $GL_n(\mathbb{C})$. We denote by $\mathcal{U}(\mathcal{G})$ the unipotent radical of $\mathcal{G}$. If $\mathcal{U}(\mathcal{G})$ is trivial, an algebraic group $\mathcal{G}$ is called reductive.

Denote by $\mathrm{Rep}(\mathcal{G})$ the category of finite-dimensional rational representations V_ρ of $\mathcal{G}$. $V_\rho \in \mathrm{Rep}(\mathcal{G})$ says V_ρ is a $\mathbb{C}$-vector space with a rational representation $\rho : \mathcal{G} \to GL(V_\rho)$. For $V_\rho \in \mathrm{Rep}(\mathcal{G})$, we denote by $\overline{V_\rho}$ the $\mathbb{C}$-vector space V_ρ with the trivial representation.

3. Representations of reductive algebraic groups

Let $\mathcal{G}$ be a reductive algebraic group and $V_\rho \in \mathrm{Rep}(\mathcal{G})$. We assume that we have an algebraic group isomorphism $F : \mathcal{G} \to \mathcal{G}$ and a linear isomorphism $\Phi : V_\rho \to V_\rho$ satisfying

$$\rho(g) \circ \Phi = \Phi \circ \rho(F(g))$$

for every $g \in \mathcal{G}$.

We take a finite set $\Lambda_\rho = \{V_\alpha\}$ of irreducible representations of $\mathcal{G}$ such that we have an irreducible decomposition $V_\rho = \bigoplus_{\Lambda_\rho} E_\alpha$ and E_α is a non-trivial irreducible component of V_ρ corresponding to an irreducible $V_\alpha \in \mathrm{Rep}(\mathcal{G})$. Denote $E_{\alpha\circ F} = \Phi(E_\alpha)$. Then we have the irreducible decomposition $V_\rho = \bigoplus_{\Lambda_\rho} E_{\alpha\circ F}$. Consider $V_{\alpha\circ F} \in \mathrm{Rep}(\mathcal{G})$ as the vector space V_α with the representation $\alpha \circ F$. Then, $E_{\alpha\circ F}$ is a non-trivial irreducible component of V_ρ corresponding to an irreducible $V_{\alpha\circ F} \in \mathrm{Rep}(\mathcal{G})$. Comparing two irreducible decomposition $V_\rho = \bigoplus_{\Lambda_\rho} E_\alpha = \bigoplus_{\Lambda_\rho} E_{\alpha\circ F}$, for each $\alpha \in \Lambda_\rho$, we have a unique $\beta \in \Lambda_\rho$ such that we have an isomorphism $\psi_\alpha : V_{\alpha\circ F} \cong V_\beta$.

Let us consider the linear map $\Phi_\alpha : V_\rho \otimes V_\alpha^* \otimes \overline{V_\alpha} \to V_\rho \otimes V_{\alpha\circ F}^* \otimes \overline{V_{\alpha\circ F}}$ defined by $\Phi_\alpha(f_\alpha \otimes v_\alpha) = \Phi \circ f_\alpha \otimes v_\alpha$ for $f_\alpha \in V_\rho \otimes V_\alpha^*$, $v_\alpha \in \overline{V_\alpha}$ (we should notice $\overline{V_\alpha} = \overline{V_{\alpha\circ F}}$). Then this linear map is $\mathcal{G}$-equivariant and hence we can restrict $\Phi_\alpha : (V_\rho \otimes V_\alpha^*)^{\mathcal{G}} \otimes \overline{V_\alpha} \to (V_\rho \otimes V_{\alpha\circ F}^*)^{\mathcal{G}} \otimes \overline{V_{\alpha\circ F}}$. Consider the isomorphism $(\psi_\alpha^{-1})^* \otimes \psi_\alpha : V_{\alpha\circ F}^* \otimes \overline{V_{\alpha\circ F}} \to V_\beta^* \otimes \overline{V_\beta}$. Then we have the linear map $((\psi_\alpha^{-1})^* \otimes \psi_\alpha) \circ \Phi_\alpha : (V_\rho \otimes V_\alpha^*)^{\mathcal{G}} \otimes \overline{V_\alpha} \to (V_\rho \otimes V_\beta^*)^{\mathcal{G}} \otimes \overline{V_\beta}$.

We consider $V_\phi \in \mathrm{Rep}(\mathcal{G})$ as $V_\phi = \bigoplus_{\Lambda_\rho}(V_\rho \otimes V_\alpha^*)^{\mathcal{G}} \otimes \overline{V_\alpha}$. Then we have the linear map $\bigoplus_{\Lambda_\rho}((\psi_\alpha^{-1})^* \otimes \psi_\alpha) \circ \Phi_\alpha : (V_\rho \otimes V_\phi)^{\mathcal{G}} \to (V_\rho \otimes V_\phi)^{\mathcal{G}}$.

Lemma 3.1. *There exists an linear isomorphism $(V_\rho \otimes V_\phi)^{\mathcal{G}} \cong \overline{V_\rho}$ which identifies the linear map $\bigoplus_{\Lambda_\rho}((\psi_\alpha^{-1})^* \otimes \psi_\alpha) \circ \Phi_\alpha : (V_\rho \otimes V_\phi)^{\mathcal{G}} \to (V_\rho \otimes V_\phi)^{\mathcal{G}}$ with $\Phi : \overline{V_\rho} \to \overline{V_\rho}$.*

Proof. By [17, Theorem 27.3.6], we have the isomorphism

$$(V_\rho \otimes V_\alpha^*)^{\mathcal{G}} \otimes \overline{V_\alpha} \ni f_\alpha \otimes v_\alpha \mapsto f(v_\alpha) \in \overline{E_\alpha}.$$

Thus we have the isomorphism $(V_\rho \otimes V_\phi)^{\mathcal{G}} \cong \bigoplus_{\Lambda_\rho} \overline{E_\alpha} = \overline{V_\rho}$. Since we have

$$((\psi_\alpha^{-1})^* \otimes \psi_\alpha) \circ \Phi_\alpha(f_\alpha \otimes v_\alpha) = \Phi \circ f_\alpha \circ \psi_\alpha^{-1} \otimes \psi_\alpha(v_\alpha),$$

the lemma follows from this isomorphism.

□

4. Rational cohomology of algebraic groups

Let $\mathcal{G}$ be an algebraic group. In this section, we denote $\mathcal{U} = \mathcal{U}(\mathcal{G})$ Let $F : \mathcal{G} \to \mathcal{G}$ be an isomorphism of algebraic groups. Consider the extension

$$1 \longrightarrow \mathcal{U} \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}/\mathcal{U} \longrightarrow 1\ . \tag{1}$$

It is known that this extension splits ([13]). The restriction $F_{\mathcal{U}} : \mathcal{U} \to \mathcal{U}$ is also an isomorphism and F induces an isomorphism $\bar{F} : \mathcal{G}/\mathcal{U} \to \mathcal{G}/\mathcal{U}$.

For $V_\phi \in \mathrm{Rep}(\mathcal{G})$, we define the rational cohomology $H^*(\mathcal{G}, V_\phi) = \mathrm{Ext}^*_{\mathcal{G}}(\mathbb{C}, V_\phi)$ as in [7] and [9]. Consider the induced map $H^*(F) : H^*(\mathcal{G}, V_\phi) \to H^*(\mathcal{G}, V_{\phi\circ F})$. Let $\Psi : V_{\phi\circ F} \to V_\phi$ be an isomorphism in $\mathrm{Rep}(\mathcal{G})$. We consider the linear map $\Psi \circ H^*(F) : H^*(\mathcal{G}, V_\phi) \to H^*(\mathcal{G}, V_\phi)$. It is known that we have a natural isomorphism

$$H^*(\mathcal{G}, V_\phi) \cong H^*(\mathcal{U}, V_\phi)^{\mathcal{G}/\mathcal{U}} \cong H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}}$$

induced by the extension (1) and the exponential map $\mathfrak{u} \to \mathcal{U}$ (see [7],[5]) where $\mathfrak{u}$ is the Lie algebra of $\mathcal{U}$ and $H^*(\mathfrak{u}, V_\phi)$ is the Lie algebra cohomology and we regard $H^*(\mathcal{U}, V_\phi), H^*(\mathfrak{u}, V_\phi) \in \mathrm{Rep}(\mathcal{G}/\mathcal{U})$ via the adjoint action. This isomorphism identifies the linear map $\Psi \circ H^*(F) : H^*(\mathcal{G}, V_\phi) \to H^*(\mathcal{G}, V_\phi)$ with the linear map $\Psi \circ H^*(F_{\mathcal{U}*}) : H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}} \to H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}}$ where $F_{\mathcal{U}*} : \mathfrak{u} \to \mathfrak{u}$ is the derivation of $F_{\mathcal{U}} : \mathcal{U} \to \mathcal{U}$.

Denote by $H^*(\mathfrak{u})$ the Lie algebra cohomology with values in the 1-dimensional trivial representation. We apply the arguments in the last section to the objects:

- The reductive algebraic group $\mathcal{G}/\mathcal{U}$.

- The representation $H^*(\mathfrak{u}) \in \mathrm{Rep}(\mathcal{G}/\mathcal{U})$.
- The isomorphism $\bar{F} : \mathcal{G}/\mathcal{U} \to \mathcal{G}/\mathcal{U}$.
- The linear isomorphism $H^*(F_{\mathcal{U}*}) : H^*(\mathfrak{u}) \to H^*(\mathfrak{u})$.

Then, taking $V_\phi \in \mathrm{Rep}(\mathcal{G}/\mathcal{U})$ and $\Psi = \bigoplus_{\Lambda_\rho}((\psi_\alpha^{-1})^* \otimes \psi_\alpha)$ as in the last section and extending $V_\phi \in \mathrm{Rep}(\mathcal{G})$ by the extension (1), the isomorphism

$$H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}} = (H^*(\mathfrak{u}) \otimes V_\phi)^{\mathcal{G}/\mathcal{U}} \cong \overline{H^*(\mathfrak{u})}$$

in Lemma 3.1 identifies the linear map $\Psi \circ H^*(F_{\mathcal{U}*}) : H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}} \to H^*(\mathfrak{u}, V_\phi)^{\mathcal{G}/\mathcal{U}}$ with the linear map $H^*(F_{\mathcal{U}*}) : \overline{H^*(\mathfrak{u})} \to \overline{H^*(\mathfrak{u})}$. This implies the following:

Proposition 4.1. *Let A be a matrix presentation of the linear map $F_{\mathcal{U}*} : \mathfrak{u} \to \mathfrak{u}$. Then we have*

$$\sum_i (-1)^i \mathrm{trace}(\Psi \circ H^*(F))_{|H^i(\mathcal{G}, V_\phi)} = \det(I - A).$$

5. Group cohomology of torsion-free virtually polycyclic groups

Lemma 5.1. ([16, Lemma 4.36.]) *Let Γ be a torsion-free virtually polycyclic group. For a finite-dimensional representation $\rho : \Gamma \to GL(V_\rho)$ on a complex vector space V_ρ, denote by $\mathcal{G}$ the Zariski-closure of $\rho(\Gamma)$ in $GL(V_\rho)$. Then we have* $\dim \mathcal{U}(\mathcal{G}) \leq \mathrm{rank}\, \Gamma$.

By this lemma, we consider the following definition.

Definition 5.2. Let Γ be a torsion-free virtually polycyclic group. For a real algebraic group $\mathcal{G}$, a representation $\rho : \Gamma \to \mathcal{G}$ is called *full* if the image $\rho(\Gamma)$ is Zariski-dense in $\mathcal{G}$ and we have $\dim \mathcal{U}(\mathcal{G}) = \mathrm{rank}\, \Gamma$.

We can compute the group cohomology of torsion-free virtually polycyclic groups by using rational cohomology of algebraic groups. Let $\mathrm{Rep}(\Gamma)$ be the category of representations V_ρ of Γ. For $V_\rho \in \mathrm{Rep}(\Gamma)$, we define the group cohomology $H^*(\Gamma, V_\rho) = \mathrm{Ext}^*_\Gamma(\mathbb{C}, V_\rho)$ as in [3].

Theorem 5.3 ([8])**.** *Let Γ be a torsion-free virtually polycyclic group. We suppose that we have an algebraic group $\mathcal{G}$ and an injection $\Gamma \subset \mathcal{G}$ which is a full representation. Let V be a rational $\mathcal{G}$-module. Then the inclusion $\Gamma \subset \mathcal{G}$ induces a cohomology isomorphism*

$$H^*(\mathcal{G}, V_\rho) \cong H^*(\Gamma, V_\rho).$$

This isomorphism was discovered by Baues for the case of trivial coefficients see the sentence after [2, Theorem 1.8].

For any torsion-free virtually polycyclic group, we have a good full representation.

Definition 5.4. We call an algebraic group $\mathcal{A}_\Gamma$ an *algebraic hull* of Γ if there exists an injective group homomorphism $\psi : \Gamma \to \mathcal{A}_\Gamma$ so that

- $\psi : \Gamma \to \mathcal{A}_\Gamma$ is a full representation.
- $Z_{\mathcal{A}_\Gamma}(\mathcal{U}_\Gamma) \subset \mathcal{U}_\Gamma$ where $Z_{\mathcal{A}_\Gamma}(\mathcal{U}_\Gamma)$ is the centralizer of the unipotent radical $\mathcal{U}_\Gamma$ of $\mathcal{A}_\Gamma$.

Theorem 5.5. ([2, Theorem A.1]) *There exists an algebraic hull of Γ and an algebraic hull of Γ is unique up to algebraic group isomorphism.*

Let Γ be a torsion-free virtually polycyclic group and $f : \Gamma \to \Gamma$ an isomorphism. Take an algebraic hull $\psi : \Gamma \to \mathcal{A}_\Gamma$. Then, f extends uniquely to an isomorphism $F : \mathcal{A}_\Gamma \to \mathcal{A}_\Gamma$ by [16, Lemma 4.41]. In fact, we obtain F by the following way.

Define $\Phi_f : \Gamma \to \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma$ as $\Phi_f(\gamma) = (\psi(\gamma), \psi \circ f(\gamma))$. Let $\mathcal{G}_f$ be the real Zariski-closure of $\Phi_f(\Gamma)$ in $\mathcal{A}_\Gamma \times \mathcal{A}_\Gamma$. For the projections $p_1 : \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma \ni (a, b) \mapsto (a) \in \mathcal{A}_\Gamma$

and $p_2 : \mathcal{A}_\Gamma \times \mathcal{A}_\Gamma \ni (a,b) \mapsto (b) \in \mathcal{A}_\Gamma$, we consider the restriction $\pi_i : \mathcal{G}_f \to \mathcal{A}_\Gamma$ of p_i on $\mathcal{G}_f$ for $i = 1, 2$. Then π_1 is an isomorphism and desired F is $\pi_2 \circ \pi_1^{-1}$.

For the isomorphism $F : \mathcal{A}_\Gamma \to \mathcal{A}_\Gamma$, by Theorem 5.3, for $V_\phi \in \mathrm{Rep}(\mathcal{A}_\Gamma)$, the induced map on the cohomology

$$H^*(f) : H^*(\Gamma, V_\phi) \to H^*(\Gamma, V_{\phi\circ f})$$

is identified with the map

$$H^*(F) : H^*(\mathcal{A}_\Gamma, V_\phi) \to H^*(\mathcal{A}_\Gamma, V_{\phi\circ F}).$$

We apply the arguments in the last section to the algebraic group $\mathcal{A}_\Gamma$. Take $V_\phi \in \mathrm{Rep}(\mathcal{A}_\Gamma/\mathcal{U}_\Gamma)$ and Ψ for Proposition 4.1. The isomorphisms

$$H^*(\Gamma, V_\phi) \cong H^*(\mathcal{A}_\Gamma, V_\phi) \cong H^*(\mathfrak{u}_\Gamma, V_\phi)^{\mathcal{G}/\mathcal{U}_\Gamma} = (H^*(\mathfrak{u}_\Gamma) \otimes V_\phi)^{\mathcal{G}/\mathcal{U}_\Gamma} \cong \overline{H^*(\mathfrak{u}_\Gamma)}$$

identify the linear map $\Psi \circ H^*(f) : H^*(\Gamma, V_\phi) \to H^*(\Gamma, V_\phi)$ with $H^*(F_{\mathcal{U}_\Gamma *}) : \overline{H^*(\mathfrak{u}_\Gamma)} \to \overline{H^*(\mathfrak{u}_\Gamma)}$ where $\mathfrak{u}_\Gamma$ is the Lie algebra of $\mathcal{U}_\Gamma$.

Proposition 5.6. *Let A be a matrix presentation of the linear map $F_{\mathcal{U}*} : \mathfrak{u}_\Gamma \to \mathfrak{u}_\Gamma$. Then we have*

$$\sum_i (-1)^i \mathrm{trace}(\Psi \circ H^*(f))_{|H^i(\Gamma, V_\phi)} = \det(I - A).$$

6. Main result

We give a "linearization formula" for any diffeomorphism of any infra-solvmanifold by using the twisted Lefschetz number and algebraic groups. It is known that the fundamental group of every infra-solvmanifold is a torsion-free virtually polycyclic group (see [4]).

Let M be an infra-solvmanifold with the fundamental group Γ and $\varphi : M \to M$ a diffeomorphism. Denote by $f : \Gamma \to \Gamma$ the homomorphism induced by φ. For the universal covering $\tilde{M}$, we write $M = \tilde{M}/\Gamma$. Then, φ is described as a diffeomorphim $\tilde{\varphi} : \tilde{M} \to \tilde{M}$ so that for $x \in \tilde{M}$ and $g \in \Gamma$, $\tilde{\varphi}(gx) = f(g)\tilde{\varphi}(x)$. For $V_\phi \in \mathrm{Rep}(\Gamma)$, we define the flat bundle $E_\phi = (\tilde{M} \times V_\phi)/\Gamma$. Then $\varphi^* E_\phi = E_{\phi\circ f}$. By the standard theory (see [16, Lemma 7.4] for instance), the induced map $H^*(\varphi) : H^*(M, E_\phi) \to H^*(M, \varphi^* E_\phi)$ is identified with $H^*(f) : H^*(\Gamma, V_\phi) \to H^*(\Gamma, V_{\phi\circ f})$. Take $V_\phi \in \mathrm{Rep}(\mathcal{A}_\Gamma/\mathcal{U}_\Gamma)$ and an isomorphism $\Psi : V_{\phi\circ F} \to V_\phi$ as in the last section. Considering $\Psi : V_{\phi\circ f} \to V_\phi$ as an isomorphism in $\mathrm{Rep}(\Gamma)$, Ψ corresponds to an isomorphism $\Xi : \varphi^* E_\phi \to E_\phi$ of flat bundles.

Thus, by Proposition 5.6, we obtain the following result.

Theorem 6.1. *Let A be a matrix presentation of $F_{\mathcal{U}_\Gamma *} : \mathfrak{u} \to \mathfrak{u}$. Then we have*

$$L(\varphi, E_\phi, \Xi) = \det(I - A).$$

Remark 1. We have a good presentation of the universal covering $\tilde{M}$ of any infra-solvmanifold M. For a torsion-free virtually polycyclic group Γ with an algebraic hull $\psi : \Gamma \to \mathcal{A}_\Gamma$, since the extension (1) splits, we have a splitting $\mathcal{A}_\Gamma = \mathcal{A}_\Gamma/\mathcal{U}_\Gamma \ltimes \mathcal{U}_\Gamma$. We can take $\mathcal{A}_\Gamma$ defined over $\mathbb{R}$ and $\psi'(\Gamma)$ contained in the $\mathbb{R}$-rational points $\mathcal{A}_\Gamma(\mathbb{R})$ of $\mathcal{A}_\Gamma$. By an injection $\psi : \Gamma \to \mathcal{A}_\Gamma(\mathbb{R})$, we have the Γ-action on $\mathcal{U}_\Gamma(\mathbb{R})$ by the splitting. In [2], Baues proved that the quotient space $\mathcal{U}_\Gamma(\mathbb{R})/\Gamma$ is a compact aspherical manifold with the universal covering $\mathcal{U}_\Gamma(\mathbb{R})$ and every infra-solvmanifold M with the fundamental group Γ is diffeomorphic to $\mathcal{U}_\Gamma(\mathbb{R})/\Gamma$. Thus, we can take $\tilde{M} = \mathcal{U}_\Gamma(\mathbb{R})$. We notice that as a Lie group, $\mathcal{U}_\Gamma(\mathbb{R})$ is simply connected nilpotent Lie group and by the exponential map, $\mathcal{U}_\Gamma(\mathbb{R})$ is diffeomorphic to the Euclidean space of dimension $\dim \mathcal{U}_\Gamma$.

Remark 2. If M is a nilmanifold, then Γ is a nilpotent and we have $\mathcal{A}_\Gamma = \mathcal{U}_\Gamma$ (see [16]) and the above quotient space $\mathcal{U}_\Gamma(\mathbb{R})/\Gamma$ is a quotient of a simply connected nilpotent Lie group $\mathcal{U}_\Gamma(\mathbb{R})$ by a discrete subgroup Γ. In this case, the flat bundle E_ϕ is trivial and so Theorem 6.1 is the usual linearization formula for a nilmanifold as we explained in Introduction.

Example 1. Let $G = \mathbb{R} \ltimes_\lambda \mathbb{R}^2$ such that $\lambda(t) = \begin{pmatrix} \cos \pi t & -\sin \pi t \\ \sin \pi t & \cos \pi t \end{pmatrix}$ for $t \in \mathbb{R}$. Then the subgroup $\Gamma = \mathbb{Z} \ltimes_\lambda \mathbb{Z}^2$ is a cocompact discrete subgroup in G. Consider the solvmanifold $M = G/\Gamma$. We have the explicit algebraic hull $\psi : \Gamma \to \mathcal{A}_\Gamma$ where $\mathcal{A}_\Gamma = \langle \tau \rangle \ltimes \mathbb{C}^3$ such that $\tau = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$ and $\psi(t, u, v) = (\tau^t, t, u, v)$ for $(t, u, v) \in \Gamma$. We consider the automorphism $f : \Gamma \to \Gamma$ defined by the matrix $A = \begin{pmatrix} -1 & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{pmatrix}$ such that $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$. This f can not be extended to an automorphism of the Lie group G. We can extend f to the automorphism $F : \mathcal{A}_\Gamma \to \mathcal{A}_\Gamma$ such that $F(\tau) = \tau$ and F on $\mathcal{U}_\Gamma = \mathbb{C}^3$ is defined by the matrix A. We can compute

$$H^i(M, \mathbb{C}) = H^i(\Gamma, \mathbb{C}) = H^i(\mathbb{C}^3)^{\langle \tau \rangle} = \begin{cases} \mathbb{C} & (0 \leq i \leq 3) \\ 0 & (\text{other } i) \end{cases}$$

and we have $H^*(f)_{|H^i(\Gamma, \mathbb{C})} = (-1)^i$. Consider a diffeomorphism $\varphi : M \to M$ inducing f on the fundamental group Γ (e.g. φ is the map on $M = \mathbb{R}^3/\Gamma$ defined by the matrix A where we consider $G = \mathbb{R}^3$ as a smooth manifold). Then the ordinary Lefschetz number is $L(\varphi) = 4$. In contrast, for the above E_ϕ and Ξ, by Theorem 6.1, we have

$$L(\varphi, E_\phi, \Xi) = \det(I - A) = 4 - 2(a + d).$$

7. Application

We consider a solvmanifold G/Γ for a discrete presentation (G, Γ) and a diffeomorphism $\varphi : G/\Gamma \to G/\Gamma$. Let $\mathfrak{g}$ be the Lie algebra of G. We suppose that the induced map $f : \Gamma \to \Gamma$ can be extended to a Lie group homomorphism $T : G \to G$. Consider the derivation $T_* : \mathfrak{g} \to \mathfrak{g}$. If an isomorphism $H^*(G/\Gamma, \mathbb{R}) \cong H^*(\mathfrak{g}, \mathbb{R})$ holds, then we have the equation

$$L(\varphi) = \det(I - B)$$

where B is a matrix presentation of $T_* : \mathfrak{g} \to \mathfrak{g}$. But in general an isomorphism $H^*(G/\Gamma, \mathbb{R}) \cong H^*(\mathfrak{g}, \mathbb{R})$ does not hold. Thus the equation $L(\varphi) = \det(I - B)$ does not hold. Even if an isomorphism $H^*(G/\Gamma, \mathbb{R}) \cong H^*(\mathfrak{g})$ does not hold, it is expected that the number $\det(I - B)$ gives some information for fixed points. As an application of Theorem 6.1, we will answer such expectation.

A solvmanifold G/Γ is an infra-solvmanifold with the fundamental group Γ. For the same notation as in Theorem 6.1, we have:

Theorem 7.1.

$$L(\varphi, E_\phi, \Xi) = \det(I - B).$$

Proof. Let G be a simply connected solvable Lie group. For a finite dimensional representation $\rho : G \to GL(V_\rho)$, taking the Zariski-closure $\mathcal{G}$ of $\rho(G)$, f we have $\dim \mathcal{U}(\mathcal{G}) \leq \dim G$ ([16, Lemma 4.36.]). For an algebraic group $\mathcal{G}$ and a representation

$\rho : G \to \mathcal{G}$, we say that ρ is a full representation if the image $\rho(G)$ is Zariski-dense in $\mathcal{G}$ and we have $\dim \mathcal{U} = \dim G$.

We call an algebraic group $\mathcal{A}_G$ an algebraic hull of G if there exists an injective Lie group homomorphism $\psi' : G \to \mathcal{A}_G$ so that:

- $\psi' : G \to \mathcal{A}_G$ is a full representation.
- $Z_{\mathcal{A}_G}(\mathcal{U}_G) \subset \mathcal{U}_G$ where $\mathcal{U}_G$ is the unipotent radical of $\mathcal{A}_G$.

As similar to virtually polycyclic case, there exists an algebraic hull $\mathcal{A}_G$ of G and an algebraic hull of G is unique up to algebraic group isomorphism (see [16, Proposition 4.40, Lemma 4.41]). Moreover, we can take $\mathcal{A}_G$ defined over $\mathbb{R}$ and $\psi'(G)$ contained in the $\mathbb{R}$-rational points $\mathcal{A}_G(\mathbb{R})$ of $\mathcal{A}_G$.

Take a real algebraic hull $\psi' : G \to \mathcal{A}_G$ of G such that $\mathcal{A}_G$ is defined over $\mathbb{R}$ and $\psi'(G) \subset \mathcal{A}_G(\mathbb{R})$. Then, by the similar arguments on virtually polycyclic groups, for an isomorphism $T : G \to G$ of Lie groups, T extends uniquely to an isomorphism $F : \mathcal{A}_G \to \mathcal{A}_G$. It is known that the composition $Q = q \circ \psi' : G \to \mathcal{U}_G(\mathbb{R})$ as a smooth map is a diffeomorphism (see [2, Proposition 2.3]). Hence, for the derivation $F_{\mathcal{U}_G *} : \mathcal{U}_G(\mathbb{R}) \to \mathcal{U}_G(\mathbb{R})$, we have $F_* = Q_* T_* Q_*^{-1}$. This implies that for matrix presentations A and B of F_* and T_* respectively, we have

$$\det(I - A) = \det(I - B).$$

We suppose that G has a cocompact discrete subgroup Γ and $T : G \to G$ is an extension of $f : \Gamma \to \Gamma$. It is known that the Zariski-closure of $\psi'(\Gamma)$ in $\mathcal{A}_G$ is the algebraic hull $\mathcal{A}_\Gamma$ of a torsion-free polycyclic group Γ and $\mathcal{U}_\Gamma = \mathcal{U}_G$ (see [16, Proof of Theorem 4.34]). Thus we can say that the restriction $F : \mathcal{A}_\Gamma \to \mathcal{A}_\Gamma$ is a unique extension of $f : \Gamma \to \Gamma$.

Now we assume that $f : \Gamma \to \Gamma$ is induced by a diffeomorphism $\varphi : G/\Gamma \to G/\Gamma$. Then, by Theorem 6.1, we have

$$L(\varphi, E_\phi, \Xi) = \det(I - A).$$

Hence the theorem follows.

□

References

[1] M. F. Atiyah, R. Bott, A Lefschetz fixed point formula for elliptic complexes. II. Applications. Ann. of Math. (2) **88** 1968 451–491.

[2] O. Baues, Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups. Topology **43** (2004), no. 4, 903–924.

[3] K. Brown, Cohomology of groups. Berlin, Heidelberg, New York: Springer 1982.

[4] F. T. Farrell and L. E. Jones, Classical Aspherical Manifolds, Amer. Math. Soc., 1990.

[5] R. Hain, Remarks on non-abelian cohomology of proalgebraic groups. J. Algebraic Geom. **22** (2013), no. 3, 581–598.

[6] A. Hattori, Spectral sequence in the de Rham cohomology of fibre bundles. J. Fac. Sci. Univ. Tokyo Sect. I **8** 1960 289–331 (1960).

[7] G. Hochschild, Cohomology of algebraic linear groups. Illinois J. Math. **5** 1961 492–519.

[8] H. Kasuya, Central theorems for cohomologies of certain solvable groups. Trans. Amer. Math. Soc. **369** (2017), no. 4, 2879–2896.

[9] J. C. Jantzen, Representations of algebraic groups. Pure and Applied Mathematics, **131**. Academic Press, Inc., Boston, MA, 1987.

[10] J. Jezierski, W. Marzantowicz, Homotopy methods in topological fixed and periodic points theory. Topological Fixed Point Theory and Its Applications, **3**. Springer, Dordrecht, 2006.

[11] A. Malcev, On a class of homogeneous spaces. (Russian) Izvestiya Akad. Nauk. SSSR. Ser. Mat. **13**, (1949). 9–32.

[12] C. K. McCord, Nielsen numbers and Lefschetz numbers on solvmanifolds. Pacific J. Math. **147** (1991), no. 1, 153–164.

[13] G. D. Mostow, Fully reducible subgroups of algebraic groups. Amer. J. Math. **78** (1956), 200–221.
[14] K. Nomizu, On the cohomology of compact homogeneous spaces of nilpotent Lie groups. Ann. of Math. (2) **59**, (1954). 531–538.
[15] A. L. Onishchik, E. B. Vinberg (Eds), Lie groups and Lie algebras Ⅱ, Springer (2000).
[16] M.S. Raghunathan, Discrete subgroups of Lie Groups, Springer-Verlag, New York, 1972.
[17] P. Tauvel and R. W. T. Yu, Lie algebras and algebraic groups, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2005.

(Hisashi Kasuya) Department of Mathematics, Graduate School of Science, Osaka University, Osaka, Japan.
E-mail address: kasuya@math.sci.osaka-u.ac.jp