PROOF OF GROTHENDIECK-SERRE CONJECTURE ON PRINCIPAL BUNDLES OVER REGULAR LOCAL RINGS CONTAINING A FINITE FIELD

IVAN PANIN

ABSTRACT. Let R be a regular local ring, containing a finite field. Let \mathbf{G} be a reductive group scheme over R. We prove that a principal \mathbf{G} -bundle over Ris trivial, if it is trivial over the fraction field of R. In other words, if K is the fraction field of R, then the map of non-abelian cohomology pointed sets

 $H^1_{\text{\acute{e}t}}(R, \mathbf{G}) \to H^1_{\text{\acute{e}t}}(K, \mathbf{G}),$

induced by the inclusion of R into K, has a trivial kernel.

Certain arguments used in the present preprint do not work if the ring R contains a characteristic zero field. In that case and, more generally, in the case when the regular local ring R contains **an infinite field** this result is proved in [FP].

1. INTRODUCTION

Assume that U is a regular scheme, \mathbf{G} is a reductive U-group scheme. Recall that a U-scheme \mathcal{G} with an action of \mathbf{G} is called a principal \mathbf{G} -bundle over U, if \mathcal{G} is faithfully flat and quasi-compact over U and the action is simple transitive, that is, the natural morphism $\mathbf{G} \times_U \mathcal{G} \to \mathcal{G} \times_U \mathcal{G}$ is an isomorphism, see [Gro3, Section 6]. It is well known that such a bundle is trivial locally in étale topology but in general not in Zariski topology. Grothendieck and Serre conjectured that \mathcal{G} is trivial locally in Zariski topology, if it is trivial generically. More precisely

Conjecture. Let R be a regular local ring, let K be its field of fractions. Let \mathbf{G} be a reductive group scheme over $U := \operatorname{Spec} R$, let \mathcal{G} be a principal \mathbf{G} -bundle. If \mathcal{G} is trivial over $\operatorname{Spec} K$, then it is trivial. Equivalently, the map of non-abelian cohomology pointed sets

$$H^1_{\acute{e}t}(R,\mathbf{G}) \to H^1_{\acute{e}t}(K,\mathbf{G}),$$

induced by the inclusion of R into K, has a trivial kernel.

The main result of this paper is a proof of this conjecture for regular semi-local domains R, containing **a finite field**. Our proof was inspired by the preprint [FP], where the conjecture is proven for semi-local regular domains containing an **infinite** field. Thus, the conjecture holds for semi-local regular domains containing a field.

The proof in the present preprint uses [Pan1, Thm.1.1], [Pan2, Thm.1.0.1], the key ideas of the paper [FP] and a Bertini type theorem from [Poo].

Our result implies that two principal **G**-bundles over U are isomorphic, if they are isomorphic over Spec K as proved in the next section. This result is new even

Key words and phrases. Reductive group schemes; Principal bundles

The author acknowledges support of the RNF-grant 14-11-00456.

for constant group schemes (that is, for group schemes coming from the ground field).

Recall that a part of the Gersten conjecture asserts that the natural homomorphism of K-groups $K_q(R) \to K_q(K)$ is injective. Very roughly speaking, the Grothendieck–Serre conjecture is a non-abelian version of this part of the Gersten conjecture.

1.1. **History of the topic.** Here is a list of known results in the same vein, corroborating the Grothendieck–Serre conjecture.

• The case, where the group scheme **G** comes from an infinite ground field, is completely solved by J.-L. Colliot-Thélène, M. Ojanguren, and M. S. Raghunatan in [CTO] and [Rag1, Rag2]; O. Gabber announced a proof for group schemes coming from arbitrary ground fields.

• The case of an arbitrary reductive group scheme over a discrete valuation ring or over a henselian ring is completely solved by Y. Nisnevich in [Nis1]. He also proved the conjecture for two-dimensional local rings in the case, when **G** is quasi-split in [Nis2].

• The case, where **G** is an arbitrary reductive group scheme over a regular semilocal domain containing an infinite field, was settled by R. Fedorov and I. Panin in [FP].

• The case, where **G** is an arbitrary torus over a regular local ring, was settled by J.-L. Colliot-Thélène and J.-J. Sansuc in [CTS].

• For some simple group schemes of classical series the conjecture is solved in works of the author, A. Suslin, M. Ojanguren, and K. Zainoulline; see [Oja1], [Oja2], [PS1], [OP], [Zai], [OPZ].

• Under an isotropy condition on **G** and assuming that the ring contains an infinite field the conjecture is proved in a series of preprints [PSV] and [Pa2].

• The case of strongly inner simple adjoint group schemes of the types E_6 and E_7 is done by the second author, V. Petrov, and A. Stavrova in [PPS]. No isotropy condition is imposed there, however it is supposed that the ring contains an infinite field.

• The case, when **G** is of the type F_4 with trivial g_3 -invariant and the field is of characteristic zero, is settled by V. Chernousov in [Che]; the case, when **G** is of the type F_4 with trivial f_3 -invariant and the field is infinite and perfect, is settled by V. Petrov and A. Stavrova in [PS2].

1.2. Acknowledgments. The author thanks A.Suslin for his interest to the topic of the present preprint.

2. Main results

Let R be a commutative unital ring. Recall that an R-group scheme \mathbf{G} is called *reductive*, if it is affine and smooth as an R-scheme and if, moreover, for each algebraically closed field Ω and for each ring homomorphism $R \to \Omega$ the scalar extension \mathbf{G}_{Ω} is a connected reductive algebraic group over Ω . This definition of a reductive R-group scheme coincides with [DG, Exp. XIX, Definition 2.7]. A well-known conjecture due to J.-P. Serre and A. Grothendieck (see [Ser, Remarque, p.31], [Gro1, Remarque 3, p.26-27], and [Gro2, Remarque 1.11.a]) asserts that

 $\mathbf{2}$

given a regular local ring R and its field of fractions K and given a reductive group scheme **G** over R, the map

$$H^1_{\text{\'et}}(R, \mathbf{G}) \to H^1_{\text{\'et}}(K, \mathbf{G}),$$

induced by the inclusion of R into K, has a trivial kernel. The following theorem, which is the main result of the present paper, asserts that this conjecture holds, provided that R contains a **finite field**. If R contains an infinite field, then the conjecture is proved in [FP].

Theorem 1. Let R be a regular semi-local domain containing a finite field, and let K be its field of fractions. Let G be a reductive group scheme over R. Then the map

$$H^1_{\acute{e}t}(R,\mathbf{G}) \to H^1_{\acute{e}t}(K,\mathbf{G}),$$

induced by the inclusion of R into K, has a trivial kernel. In other words, under the above assumptions on R and G, each principal G-bundle over R having a Krational point is trivial.

Theorem 1 has the following

Corollary. Under the hypothesis of Theorem 1, the map

$$H^1_{\acute{e}t}(R,\mathbf{G}) \to H^1_{\acute{e}t}(K,\mathbf{G}),$$

induced by the inclusion of R into K, is injective. Equivalently, if \mathcal{G}_1 and \mathcal{G}_2 are two principal bundles isomorphic over Spec K, then they are isomorphic.

Proof. Let \mathcal{G}_1 and \mathcal{G}_2 be two principal **G**-bundles isomorphic over Spec K. Let $\operatorname{Iso}(\mathcal{G}_1, \mathcal{G}_2)$ be the scheme of isomorphisms. This scheme is a principal $\operatorname{Aut} \mathcal{G}_2$ -bundle. By Theorem 1 it is trivial, and we see that $\mathcal{G}_1 \cong \mathcal{G}_2$.

Note that, while Theorem 1 was previously known for reductive group schemes \mathbf{G} coming from the ground field (an unpublished result due to O.Gabber), in many cases the corollary is a new result even for such group schemes.

For a scheme U we denote by \mathbb{A}^1_U the affine line over U and by \mathbb{P}^1_U the projective line over U. Let T be a U-scheme. By a principal **G**-bundle over T we understand a principal $\mathbf{G} \times_U T$ -bundle.

In Section 3 we deduce Theorem 1 from the following result of independent interest (cf. [PSV, Thm.1.3]).

Theorem 2. Let R be the semi-local ring of finitely many closed points on an irreducible smooth affine variety over a finite field k, set $U = \operatorname{Spec} R$. Let \mathbf{G} be a simple simply-connected group scheme over U (see [DG, Exp. XXIV, Sect. 5.3] for the definition). Let \mathcal{E}_t be a principal \mathbf{G} -bundle over the affine line $\mathbb{A}^1_U = \operatorname{Spec} R[t]$, and let $h(t) \in R[t]$ be a monic polynomial. Denote by $(\mathbb{A}^1_U)_h$ the open subscheme in \mathbb{A}^1_U given by $h(t) \neq 0$ and assume that the restriction of \mathcal{E}_t to $(\mathbb{A}^1_U)_h$ is a trivial principal \mathbf{G} -bundle. Then for each section $s: U \to \mathbb{A}^1_U$ of the projection $\mathbb{A}^1_U \to U$ the \mathbf{G} -bundle $s^* \mathcal{E}_t$ over U is trivial.

The derivation of Theorem 1 from Theorem 2 is based on [Pan2, Thm.1.0.1] and [Pan1, Thm.1.1].

Let Y be a semi-local scheme. We will call a simple Y-group scheme quasi-split if its restriction to each connected component of Y contains a **Borel subgroup** scheme.

Theorem 3. Let R, U, and \mathbf{G} be as in Theorem 2. Let $Z \subset \mathbb{P}^1_U$ be a closed subscheme finite over U. Let $Y \subset \mathbb{P}^1_U$ be a closed subscheme finite and étale over U and such that

(i) $\mathbf{G}_Y := \mathbf{G} \times_U Y$ is quasi-split,

(ii) $Y \cap Z = \emptyset$ and $Y \cap \{\infty\} \times U = \emptyset = Z \cap \{\infty\} \times U$,

(iii) for any closed point $u \in U$ one has $Pic(\mathbb{P}^1_u - Y_u) = 0$, where $Y_u := \mathbb{P}^1_u \cap Y$. Let \mathcal{G} be a principal **G**-bundle over \mathbb{P}^1_U such that its restriction to $\mathbb{P}^1_U - Z$ is trivial. Then the restriction of \mathcal{G} to $\mathbb{P}^1_U - Y$ is also trivial.

In particular, the principal G-bundle \mathcal{G} is trivial locally for the Zarisky topology.

The proof of this result is inspired by [FP, Thm.3].

2.1. Organization of the paper. In Section 3, we reduce Theorem 1 to Theorem 2. In Section 4, we reduce Theorem 2 to Theorem 3. This reduction is based on [Pan2, Thm.1.0.1], [Pan1, Thm.1.1], on a theorem of D. Popescu [Pop] and on Proposition 4.1. The latter proposition is a new ingredient comparing with respecting arguments from [FP, Section 4].

In Section 5 we prove Theorem 3. We give an outline of the proof in Section 5.1. We use the technique of henselization.

In Section 6 we give an application of Theorem 1.

In the Appendix we recall the definition of henselization from [Gab, Section 0].

3. Reducing Theorem 1 to Theorem 2

In what follows "**G**-bundle" always means "principal **G**-bundle". Now we assume that Theorem 2 holds. We start with the following particular case of Theorem 1.

Proposition 3.1. Let R, $U = \operatorname{Spec} R$, and G be as in Theorem 2. Let \mathcal{E} be a principal G-bundle over U, trivial at the generic point of U. Then \mathcal{E} is trivial.

Proof. Under the hypothesis of the proposition, the following data are constructed in [Pan1, Thm.1.1]:

(a) a principal **G**-bundle \mathcal{E}_t over \mathbb{A}^1_U ;

(b) a monic polynomial $h(t) \in R[t]$.

Moreover these data satisfies the following conditions:

(1) the restriction of \mathcal{E}_t to $(\mathbb{A}^1_U)_h$ is a trivial principal **G**-bundle;

(2) there is a section $s: U \to \mathbb{A}^1_U$ such that $s^* \mathcal{E}_t = \mathcal{E}$.

Now it follows from Theorem 2 that \mathcal{E} is trivial.

Proposition 3.2. Let U be as in Theorem 2. Let **G** be a reductive group scheme over U. Let \mathcal{E} be a principal **G**-bundle over U trivial at the generic point of U. Then \mathcal{E} is trivial.

Proof. Firstly, using [Pan2, Thm.1.0.1], we can assume that **G** is semi-simple and simply-connected. Secondly, standard arguments (see for instance [PSV, Section 9]) show that we can assume that **G** is simple and simply-connected. (Note that for this reduction it is necessary to work with semi-local rings.) Now the proposition is reduced to Proposition 3.1.

Proof of Theorem 1. Let us prove a general statement first. Let k' be a finite field, X be a k'-smooth irreducible affine variety, **H** be a reductive group scheme over X. Denote by k'[X] the ring of regular functions on X and by k'(X) the field of rational functions on X. Let \mathcal{H} be a principal **H**-bundle over X trivial over k'(X).

Let $\mathfrak{p}_1, \ldots, \mathfrak{p}_n$ be prime ideals in k'[X], and let $\mathcal{O}_{\mathfrak{p}_1,\ldots,\mathfrak{p}_n}$ be the corresponding semilocal ring.

Lemma 3.3. The principal **H**-bundle \mathcal{H} is trivial over $\mathcal{O}_{\mathfrak{p}_1,\ldots,\mathfrak{p}_n}$.

Proof. For each i = 1, 2, ..., n choose a maximal ideal $\mathfrak{m}_i \subset k'[X]$ containing \mathfrak{p}_i . One has inclusions of k'-algebras

$$\mathcal{O}_{\mathfrak{m}_1,\ldots,\mathfrak{m}_n} \subset \mathcal{O}_{\mathfrak{p}_1,\ldots,\mathfrak{p}_n} \subset k'(X).$$

By Proposition 3.2 the principal **H**-bundle \mathcal{H} is trivial over $\mathcal{O}_{\mathfrak{m}_1,\ldots,\mathfrak{m}_n}$. Thus it is trivial over $\mathcal{O}_{\mathfrak{p}_1,\ldots,\mathfrak{p}_n}$.

Let us return to our situation. Let $\mathfrak{m}_1, \ldots, \mathfrak{m}_n$ be all the maximal ideals of R. Let \mathcal{E} be a **G**-bundle over R trivial over the fraction field of R. Clearly, there is a nonzero $f \in R$ such that \mathcal{E} is trivial over R_f . Let k be the prime field of R. Note that k is perfect. It follows from Popescu's theorem ([Pop, Swa]) that R is a filtered inductive limit of smooth k-algebras R_{α} . Modifying the inductive system R_{α} if necessary, we can assume that each R_{α} is integral. There are an index α , a reductive group scheme \mathbf{G}_{α} over R_{α} , a principal \mathbf{G}_{α} -bundle \mathcal{E}_{α} over R_{α} , and an element $f_{\alpha} \in R_{\alpha}$ such that $\mathbf{G} = \mathbf{G}_{\alpha} \times_{\operatorname{Spec} R_{\alpha}} \operatorname{Spec} R$, \mathcal{E} is isomorphic to $\mathcal{E}_{\alpha} \times_{\operatorname{Spec} R_{\alpha}} \operatorname{Spec} R$ as principal \mathbf{G} -bundle, f is the image of f_{α} under the homomorphism $\varphi_{\alpha} : R_{\alpha} \to R$, \mathcal{E}_{α} is trivial over $(R_{\alpha})_{f_{\alpha}}$.

For each maximal ideal \mathfrak{m}_i in R (i = 1, ..., n) set $\mathfrak{p}_i = \varphi_{\alpha}^{-1}(\mathfrak{m}_i)$. The homomorphism φ_{α} induces a homomorphism of semi-local rings $(R_{\alpha})_{\mathfrak{p}_1,...,\mathfrak{p}_n} \to R$. By Lemma 3.3 the principal \mathbf{G}_{α} -bundle \mathcal{E}_{α} is trivial over $(R_{\alpha})_{\mathfrak{p}_1,...,\mathfrak{p}_n}$. Whence the \mathbf{G} -bundle \mathcal{E} is trivial over R.

4. Reducing Theorem 2 to Theorem 3

Now we assume that Theorem 3 is true. Let k, U and \mathbf{G} be as in Theorem 2. Let u_1, \ldots, u_n be all the closed points of U. Let $k(u_i)$ be the residue field of u_i . Consider the reduced closed subscheme \mathbf{u} of U, whose points are u_1, \ldots, u_n . Thus

$$\mathbf{u} \cong \coprod_i \operatorname{Spec} k(u_i)$$

Set $\mathbf{G}_{\mathbf{u}} = \mathbf{G} \times_U \mathbf{u}$. By \mathbf{G}_{u_i} we denote the fiber of \mathbf{G} over u_i ; it is a simple simply-connected algebraic group over $k(u_i)$.

Proposition 4.1. Let $Z \subset \mathbb{A}_U^1$ be a closed subscheme finite over U. There is a closed subscheme $Y \subset \mathbb{A}_U^1$ which is étale and finite over U and such that (i) $\mathbf{G}_Y := \mathbf{G} \times_U Y$ is quasi-split, (ii) $Y \cap Z = \emptyset$,

(iii) for any closed point $u \in U$ one has $Pic(\mathbb{P}^1_u - Y_u) = 0$, where $Y_u := \mathbb{P}^1_u \cap Y$. (Note that Y and Z are closed in \mathbb{P}^1_U since they are finite over U).

Proof. For every u_i in **u** choose a Borel subgroup \mathbf{B}_{u_i} in \mathbf{G}_{u_i} . The laller is possible since the fields $k(u_i)$ are finite. Let \mathcal{B} be the U-scheme of Borel subgroup schemes of **G**. It is a smooth projective U-scheme (see [DG, Cor. 3.5, Exp. XXVI]). The subgroup \mathbf{B}_{u_i} in \mathbf{G}_{u_i} is a $k(u_i)$ -rational point b_i in the fibre of \mathcal{B} over the point u_i . Using a variant of Bertini theorem (see [Poo, Thm.1.2]), we can find a closed subscheme Y' of \mathcal{B} such that Y' is étale over U and all the b_i 's are in Y (take an embedding of \mathcal{B} into a projective space \mathbb{P}_U^N and intersect \mathcal{B} with appropriately chosen family of hypersurfaces containing the points b_i . Arguing as in the proof of

[OP, Lemma 7.2], we get a scheme Y' finite and étale over U). For any closed point u_i in U the fibre Y'_{u_i} of Y' over u_i contains a $k(u_i)$ -rational point (it is the point b_i).

To continue the proof of the Proposition we need the following

Lemma 4.2. Let U be as in the Proposition. Let $Z \subset \mathbb{A}^1_U$ be a closed subscheme finite over U. Let $Y' \to U$ be a finite étale morphism such that for any closed point u_i in U the fibre Y'_{u_i} of Y' over u_i contains a $k(u_i)$ -rational point. Then there are finite field extensions k_1 and k_2 of the finite field k such that

(i) the degrees $[k_1 : k]$ and $[k_2 : k]$ are coprime,

(ii) $k(u_i) \otimes_k k_r$ is a field for r = 1 and r = 2,

(iii) the degrees $[k_1 : k]$ and $[k_2 : k]$ are strictly greater than any of the degrees [k(z) : k], where z runs over all closed points of Z,

(iv) there is a closed embedding of U-schemes $Y'' = ((Y' \otimes_k k_1) \coprod (Y' \otimes_k k_2)) \xrightarrow{i} \mathbb{A}^1_U$, (v) for Y = i(Y'') one has $Y \cap Z = \emptyset$,

(vi) for any closed point u_i in U one has $Pic(\mathbb{P}^1_{u_i} - Y_{u_i}) = 0$.

To prove this Lemma note that it's easy to find field extensions k_1 and k_2 subjecting (i) to (iii). To satisfy (iv) it suffices to require that for any closed point u_i in U and for r = 1 and r = 2 the number of closed points in $Y'_{u_i} \otimes_k k_r$ is the same as the number of closed points in Y'_{u_i} , and to require that for any integer n > 0 and any closed point u_i in U the number of points $y \in Y''_{u_i}$ with $[k(y) : k(u_i)] = n$ is not more than the number of points $x \in \mathbb{A}^1_{u_i}$ with $[k(x) : k(u_i)] = n$. Clearly, these requirements can be satisfied, which proves the item (iv).

The condition (v) holds for any closed U-embedding $i: Y'' \hookrightarrow \mathbb{A}^1_U$ from item (iv), since the property (iii). The condition (vi) holds since the property (i).

Now complete the proof of Proposition 4.1. Take the *U*-scheme $Y' \subset \mathbf{B}$ as in the beginning of the proof. This *U*-scheme Y' satisfies the assumption of Lemma 4.2. Take the closed subscheme Y of \mathbb{A}^1_U as in the item (v) of the Lemma. For this Y the conditions (ii) and (iii) of the Proposition are obviously satisfied. The condition (i) is satisfied too, since already it is satisfied for the *U*-scheme Y'. The Proposition follows.

Proof of Theorem 2. Set $Z := \{h = 0\} \cup s(U) \subset \mathbb{A}^1_U$. Clearly, Z is finite over U. Since the principal **G**-bundle \mathcal{E}_t is trivial over $(\mathbb{A}^1_U)_h$ it is trivial over $\mathbb{A}^1_U - Z$. Note that $\{h = 0\}$ is closed in \mathbb{P}^1_U and finite over U because h is monic. Further, s(U) is also closed in \mathbb{P}^1_U and finite over U because it is a zero set of a degree one monic polynomial. Thus $Z \subset \mathbb{P}^1_U$ is closed and finite over U.

Since the principal **G**-bundle \mathcal{E}_t is trivial over $(\mathbb{A}^1_U)_h$, and **G**-bundles can be glued in Zariski topology, there exists a principal **G**-bundle \mathcal{G} over \mathbb{P}^1_U such that

(i) its restriction to \mathbb{A}^1_U coincides with \mathcal{E}_t ;

(ii) its restriction to $\mathbb{P}^1_U - Z$ is trivial.

Now choose Y in \mathbb{A}^1_U as in Proposition 4.1. Clearly, Y is finite étale over U and closed in \mathbb{P}^1_U . Moreover, $Y \cap \{\infty\} \times U = \emptyset = Z \cap \{\infty\} \times U$ and $Y \cap Z = \emptyset$. Applying Theorem 3 with this choice of Y and Z, we see that the restriction of \mathcal{G} to $\mathbb{P}^1_U - Y$ is a trivial **G**-bundle. Since s(U) is in $\mathbb{A}^1_U - Y$ and $\mathcal{G}|_{\mathbb{A}^1_U}$ coincides with \mathcal{E}_t , we conclude that $s^*\mathcal{E}_t$ is a trivial principal **G**-bundle over U.

 $\mathbf{6}$

5. Proof of Theorem 3

We will be using notation from Theorem 3. Let \mathbf{u} be as in Section 4. For $u \in \mathbf{u}$ set $\mathbf{G}_u = \mathbf{G}|_u$.

Proposition 5.1. Let \mathcal{E} be a **G**-bundle over \mathbb{P}^1_U such that $\mathcal{E}|_{\mathbb{P}^1_u}$ is a trivial \mathbf{G}_u bundle for all $u \in \mathbf{u}$. Assume that there exists a closed subscheme T of \mathbb{P}^1_U finite over U such that the restriction of \mathcal{E} to $\mathbb{P}^1_U - T$ is trivial and $(\infty \times U) \cap T = \emptyset$. Then \mathcal{E} is trivial.

Proof. This follows from Theorem 9.6 of [PSV], since $\mathcal{E}|_{(\infty \times U)}$ is a trivial **G**-bundle.

5.1. An outline of the proof of Theorem 3. Our proof of this Theorem almost literally coincides with the proof of [FP, Thm.3]. Our arguments are simpler at certain points.

An outline of the proof.

Denote by Y^h the henselization of the pair (\mathbb{A}^1_U, Y) , it is a scheme over \mathbb{A}^1_U . Let $s: Y \to Y^h$ be the canonical closed embedding, see Section 5.2 for more details. Set $\dot{Y}^h := Y^h - s(Y)$. Let \mathcal{G}' be a **G**-bundle over $\mathbb{P}^1_U - Y$. Denote by $\operatorname{Gl}(\mathcal{G}', \varphi)$ the **G**-bundle over \mathbb{P}^1_U obtained by gluing \mathcal{G}' with the trivial **G**-bundle $\mathbf{G} \times_U Y^h$ via a **G**-bundle isomorphism $\varphi: \mathbf{G} \times_U \dot{Y}^h \to \mathcal{G}'|_{\dot{Y}^h}$.

Note that the **G**-bundle \mathcal{G} can be presented in the form $\operatorname{Gl}(\mathcal{G}', \varphi)$, where $\mathcal{G}' = \mathcal{G}|_{\mathbb{P}^1_{t-Y}}$. The idea is to show that

There is $\alpha \in \mathbf{G}(\dot{Y}^h)$ such that the $\mathbf{G}_{\mathbf{u}}$ -bundle $\operatorname{Gl}(\mathcal{G}', \varphi \circ \alpha)|_{\mathbb{P}^1_{\mathbf{u}}}$ is trivial (here α (*) is regarded as an automorphism of the **G**-bundle $\mathbf{G} \times_U \dot{Y}^h$ given by the right translation by the element α).

If we find α satisfying condition (*), then Proposition 5.1, applied to $T = Y \cup Z$, shows that the **G**-bundle $\operatorname{Gl}(\mathcal{G}', \varphi \circ \alpha)$ is trivial over \mathbb{P}^1_U . On the other hand, its restriction to $\mathbb{P}^1_U - Y$ coincides with the **G**-bundle $\mathcal{G}' = \mathcal{G}|_{\mathbb{P}^1_U - Y}$. Thus $\mathcal{G}|_{\mathbb{P}^1_U - Y}$ is a trivial **G**-bundle.

To prove (*) it suffices to show that

(i) the bundle $\mathcal{G}|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}}$ is trivial;

(ii) each element $\gamma_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(\dot{Y}_{\mathbf{u}}^{h})$ can be written in the form

$$\alpha|_{\dot{Y}_{\mathbf{u}}^{h}} \cdot \beta_{\mathbf{u}}|_{\dot{Y}_{\mathbf{u}}^{h}}$$

for certain elements $\alpha \in \mathbf{G}(\dot{Y}^h)$ and $\beta_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(Y^h_{\mathbf{u}})$.

A realization of this plan in details is given below in the paper.

5.2. Henselization of affine pairs. We will use the theory of henselian pairs and, in particular, a notion of a henselization A_I^h of a commutative ring A at an ideal I (see Appendix and [Gab, Section 0]). We refer to [FP, subsection 5.2] for the geometric counterpart. Let S = Spec A be a scheme and T = Spec(A/I) be a closed subscheme. Let $(T^h, \pi : T^h \to S, s : T \to T^h)$ be the henselization of the pair (S,T) (cf. Definition A.3). By definition the scheme T^h is affine and the composite morphism $\pi \circ s : T \to S$ is the closed embedding $T \hookrightarrow S$. Recall that the pair $(T^h, s(T))$ is henselian, which means that for any affine étale morphism $\pi : Z \to T^h$, any section σ of π over s(T) uniquely extends to a section of π over T^h . It is known that $\pi^{-1}(T) = s(T)$.

In the notation of [Gab, Section 0] we have $T^h = \operatorname{Spec} A_I^h, \pi: T^h \to S$ is induced by the structure of A-algebra on A_I^h .

Recall three properties of henselization of affine pairs

(i) Let T be a semi-local scheme. Then the henselization commutes with restriction to closed subschemes. In more details, if $S' \subset S$ is a closed subscheme, then there is a natural morphism $(T \times_S S')^h \to T^h \times_S S'$. This morphism is an isomorphism and the canonical section $s': T \times_S S' \to (T \times_S S')^h$ coincides under this identification with

$$s \times_S \operatorname{Id}_{S'} : T \times_S S' \to T^h \times_S S'.$$

(ii) If $T = \coprod_i T_i$ is a disjoint union, then $T^h = \coprod_i T_i^h$. (iii) If we replace in a pair (S, T) the scheme S by an étale affine neighborhood of T, then the (T^h, π, s) remains the same. In more details, given a pair (S, T) as above we write temporarily $(S_T^{hen}, \pi_{S,T}, s_{S,T})$ for (T^h, π, s) . If $p: W \to S$ is an étale morphism and $t: T \hookrightarrow W$ is such that $p \circ t: T \hookrightarrow S$ coincides with the closed embedding T into S, then there is a canonical isomorphism $\rho: W_T^{hen} \to S_T^{hen}$ of the S-schemes $(W_T^{hen}, \pi_{W,T})$ and $(S_T^{hen}, \pi_{S,T})$ such that $\rho \circ s_{W,T} = s_{S,T}$.

5.3. Gluing principal G-bundles. Recall that $U = \operatorname{Spec} R$, where R is the semilocal ring of finitely many closed points on an irreducible k-smooth affine variety over a finite field k. Also, G is a simple simply-connected group scheme over U, and Y is a closed subscheme of \mathbb{P}^1_U finite and étale over U.

We will assume below in the preprint that $Y \subset \mathbb{A}^1_U$ (as in the hypotheses of Theorem 3). Let (Y^h, π, s) be the henselization of the pair (\mathbb{A}^1_U, Y) and let $\dot{Y}^h =$ $Y^h - s(Y)$ and let $in : \mathbb{A}^1_U \hookrightarrow \mathbb{P}^1_U$ be the open inclusion.

Proposition 5.2. [FP] The schemes Y^h and \dot{Y}^h are affine.

Let us make a general remark. Let \mathcal{F} be a **G**-bundle over a U-scheme T. By definition, a trivialization of \mathcal{F} is a **G**-equivariant isomorphism $\mathbf{G} \times_U T \to \mathcal{F}$. Equivalently, it is a section of the projection $\mathcal{F} \to T$. If φ is such a trivialization and $f: T' \to T$ is a U-morphism, we get a trivialization $f^*\varphi$ of $f^*\mathcal{F}$. Sometimes we denote this trivialization by $\varphi|_{T'}$. We also sometimes call a trivialization of $f^*\mathcal{F}$ a trivialization of \mathcal{F} on T'.

The main cartesian square we will work with is

Let \mathcal{A} be the category of pairs (\mathcal{E}, ψ) , where \mathcal{E} is a **G**-bundle on \mathbb{P}^1_U , ψ is a trivialization of $\mathcal{E}|_{Y^h} := (in \circ \pi)^* \mathcal{E}$. A morphism between (\mathcal{E}, ψ) and (\mathcal{E}', ψ') is an isomorphism $\mathcal{E} \to \mathcal{E}'$ compatible with trivializations.

Similarly, let \mathcal{B} be the category of pairs (\mathcal{E}, ψ) , where \mathcal{E} is a **G**-bundle on $\mathbb{P}^1_U - Y$, ψ is a trivialization of $\mathcal{E}|_{\dot{Y}^h}$.

Consider the restriction functor $\Psi : \mathcal{A} \to \mathcal{B}$.

Proposition 5.3. [FP] The functor Ψ is an equivalence of categories.

Construction 5.4. [FP] By Proposition 5.3 we can choose a functor quasi-inverse to Ψ . Fix such a functor Θ . Let Λ be the forgetful functor from \mathcal{A} to the category of **G**-bundles over \mathbb{P}^1_U . For $(\mathcal{E}, \psi) \in \mathcal{B}$ set

$$\operatorname{Gl}(\mathcal{E}, \psi) = \Lambda(\Theta(\mathcal{E}, \psi)).$$


Note that $\operatorname{Gl}(\mathcal{E}, \psi)$ comes with a canonical trivialization over Y^h .

Conversely, if \mathcal{E} is a principal **G**-bundle over \mathbb{P}^1_U such that its restriction to Y^h is trivial, then \mathcal{E} can be represented as $\operatorname{Gl}(\mathcal{E}', \psi)$, where $\mathcal{E}' = \mathcal{E}|_{\mathbb{P}^1_U - Y}, \psi$ is a trivialization of \mathcal{E}' on \dot{Y}^h .

Let **u** be as in Section 4, $Y_{\mathbf{u}} := Y \times_U \mathbf{u}$. Let $(Y_{\mathbf{u}}^h, \pi_{\mathbf{u}}, s_{\mathbf{u}})$ be the henselization of $(\mathbb{A}_{\mathbf{u}}^1, Y_{\mathbf{u}})$. Using property (i) of henselization, we get $Y_{\mathbf{u}}^h = Y^h \times_U \mathbf{u}$. Thus we have a natural closed embedding $Y_{\mathbf{u}}^h \to Y^h$. Set $\dot{Y}_{\mathbf{u}}^h = Y_{\mathbf{u}}^h - s_{\mathbf{u}}(Y_{\mathbf{u}})$. We get a closed embedding

(2)
$$\dot{Y}^h_{\mathbf{u}} \hookrightarrow \dot{Y}^h$$

Thus the pull-back of the cartesian square (1) by means of the closed embedding $\mathbf{u} \hookrightarrow U$ has the form



where $in_{\mathbf{u}} : \mathbb{A}^1_{\mathbf{u}} \to \mathbb{P}^1_{\mathbf{u}}$.

Similarly to the above, we can define categories $\mathcal{A}_{\mathbf{u}}$ and $\mathcal{B}_{\mathbf{u}}$ and an equivalence of categories $\Psi_{\mathbf{u}} : \mathcal{A}_{\mathbf{u}} \to \mathcal{B}_{\mathbf{u}}$. Let $\Theta_{\mathbf{u}}$ be a functor quasi-inverse to $\Psi_{\mathbf{u}}$ and $\Lambda_{\mathbf{u}}$ be the forgetful functor from $\mathcal{A}_{\mathbf{u}}$ to the category of $\mathbf{G}_{\mathbf{u}}$ -bundles over $\mathbb{P}^{1}_{\mathbf{u}}$. Let $\mathcal{E}_{\mathbf{u}}$ be a principal $\mathbf{G}_{\mathbf{u}}$ -bundle over $\mathbb{P}^{1}_{\mathbf{u}} - Y_{\mathbf{u}}$ and $\psi_{\mathbf{u}}$ be a trivialization of $\mathbf{G}_{\mathbf{u}}$ on $\dot{Y}^{h}_{\mathbf{u}}$. Set

$$\operatorname{Gl}_{\mathbf{u}}(\mathcal{E}_{\mathbf{u}},\psi_{\mathbf{u}}) = \Lambda_{\mathbf{u}}(\Theta_{\mathbf{u}}(\mathcal{E}_{\mathbf{u}},\psi_{\mathbf{u}})).$$

Lemma 5.5. [FP] Let $(\mathcal{E}, \psi) \in \mathcal{B}$, and let $\operatorname{Gl}(\mathcal{E}, \psi)$ be the **G**-bundle obtained by Construction 5.4. Then

$$\operatorname{Gl}_{\mathbf{u}}(\mathcal{E}|_{\mathbb{P}^{1}_{\mathbf{u}}-Y_{\mathbf{u}}},\psi|_{\dot{Y}^{h}_{\mathbf{u}}}) and \operatorname{Gl}(\mathcal{E},\psi)|_{\mathbb{P}^{1}_{\mathbf{u}}}$$

are isomorphic as $\mathbf{G}_{\mathbf{u}}$ -bundles over $\mathbb{P}^{1}_{\mathbf{u}}$.

Lemma 5.6. [FP] For any $(\mathcal{E}_{\mathbf{u}}, \psi_{\mathbf{u}}) \in \mathcal{B}_{\mathbf{u}}$ and any $\beta_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(Y_{\mathbf{u}}^{h})$ the $\mathbf{G}_{\mathbf{u}}$ -bundles $\operatorname{Gl}_{\mathbf{u}}(\mathcal{E}_{\mathbf{u}}, \psi_{\mathbf{u}})$ and $\operatorname{Gl}_{\mathbf{u}}(\mathcal{E}_{\mathbf{u}}, \psi_{\mathbf{u}} \circ \beta_{\mathbf{u}}|_{\dot{Y}_{\mathbf{u}}^{h}})$

are isomorphic (here $\beta_{\mathbf{u}}|_{\dot{Y}_{\mathbf{u}}^{h}}$ is regarded as an automorphism of the $\mathbf{G}_{\mathbf{u}}$ -bundle $\mathbf{G}_{\mathbf{u}} \times_{\mathbf{u}}$ $\dot{Y}_{\mathbf{u}}^{h}$ given by the right translation by $\beta_{\mathbf{u}}|_{\dot{Y}_{\mathbf{u}}^{h}}$).

5.4. Proof of Theorem 3: presentation of \mathcal{G} in the form $\operatorname{Gl}(\mathcal{G}', \varphi)$. Let U, \mathbf{G}, Z, Y and \mathcal{G} be as in Theorem 3.

Proposition 5.7. [FP] The **G**-bundle \mathcal{G} over \mathbb{P}^1_U is of the form $\operatorname{Gl}(\mathcal{G}', \varphi)$ for the **G**-bundle $\mathcal{G}' := \mathcal{G}|_{\mathbb{P}^1_U - Y}$ and a trivialization φ of \mathcal{G}' over \dot{Y}^h .

Proof. In view of Construction 5.4, it is enough to prove that the restriction of the principal **G**-bundle \mathcal{G} to Y^h is trivial. Let us choose a closed subscheme $Z' \subset \mathbb{A}^1_U$ such that Z' contains $Z, Z' \cap Y = \emptyset$, and $\mathbb{A}^1_U - Z'$ is affine. Then $\mathbb{A}^1_U - Z'$ is an affine neighborhood of Y. By the property (iii) from subsection 5.2 the henselization of

the pair $(\mathbb{A}_U^1 - Z', Y)$ coincides with the henselization of the pair (\mathbb{A}_U^1, Y) . Since \mathcal{G} is trivial over $\mathbb{A}_U^1 - Z'$, its pull-back to Y^h is trivial too. The proposition is proved.

Our aim is to modify the trivialization φ via an element

 $\alpha \in \mathbf{G}(\dot{Y}^h)$

so that the **G**-bundle $\operatorname{Gl}(\mathcal{G}', \varphi \circ \alpha)$ becomes trivial over \mathbb{P}^1_U .

5.5. **Proof of Theorem 3: proof of property (i) from the outline.** Now we are able to prove property (i) from the outline of the proof. In fact, we will prove the following modification of [FP, Lemma 5.11].

Lemma 5.8. Let $\operatorname{Gl}(\mathcal{G}', \varphi)$ be the presentation of the **G**-bundle \mathcal{G} over \mathbb{P}^1_U given in Proposition 5.7. Set $\varphi_{\mathbf{u}} := \varphi|_{\dot{Y}^h_{\mathbf{u}}}$. Then there is $\gamma_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(\dot{Y}^h_{\mathbf{u}})$ such that the $\mathbf{G}_{\mathbf{u}}$ -bundle $\operatorname{Gl}_{\mathbf{u}}(\mathcal{G}'|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}}, \varphi_{\mathbf{u}} \circ \gamma_{\mathbf{u}})$ is trivial.

Proof. We show first that $\mathcal{G}|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}}$ is trivial. One has

$$\mathbb{P}^1_{\mathbf{u}} = \coprod_{u \in \mathbf{u}} \mathbb{P}^1_u$$

For $u \in \mathbf{u}$ set $Y_u := Y \times_U u$, $\mathbf{G}_u := \mathbf{G} \times_U u$, and $\mathcal{G}_u := \mathcal{G} \times_U u$.

Take $u \in \mathbf{u}$. By our assumption on Y, $Pic(\mathbb{P}_u^1 - Y_u) = 0$. The \mathbf{G}_u -bundle \mathcal{G}_u is trivial over $\mathbb{A}_u^1 - Z_u$. Thus, by [Gil1, Corollary 3.10(a)], it is trivial over $\mathbb{P}_u^1 - Y_u$.

We see that $\mathcal{G}'|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}} = \mathcal{G}|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}}$ is trivial. Choosing a trivialization, we may identify $\varphi_{\mathbf{u}}$ with an element of $\mathbf{G}_{\mathbf{u}}(\dot{Y}^h_{\mathbf{u}})$. Set $\gamma_{\mathbf{u}} = \varphi_{\mathbf{u}}^{-1}$. By the very choice of $\gamma_{\mathbf{u}}$ the $\mathbf{G}_{\mathbf{u}}$ -bundle $\mathrm{Gl}_{\mathbf{u}}(\mathcal{G}'|_{\mathbb{P}^1_{\mathbf{u}}-Y_{\mathbf{u}}}, \varphi_{\mathbf{u}} \circ \gamma_{\mathbf{u}})$ is trivial.

5.6. **Proof of Theorem 3: reduction to property (ii) from the outline.** The aim of this section is to deduce Theorem 3 from the following

Proposition 5.9. [FP] Each element $\gamma_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(\dot{Y}_{\mathbf{u}}^h)$ can be written in the form

$$\alpha|_{\dot{Y}_{h}^{h}} \cdot \beta_{\mathbf{u}}|_{\dot{Y}_{h}^{h}}$$

for certain elements $\alpha \in \mathbf{G}(\dot{Y}^h)$ and $\beta_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(Y^h_{\mathbf{u}})$.

Deduction of Theorem 3 from Proposition 5.9. [FP] Let $\operatorname{Gl}(\mathcal{G}', \varphi)$ be the presentation of the **G**-bundle \mathcal{G} from Proposition 5.7. Let $\gamma_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(\dot{Y}_{\mathbf{u}}^h)$ be the element from Lemma 5.8. Let $\alpha \in \mathbf{G}(\dot{Y}^h)$ and $\beta_{\mathbf{u}} \in \mathbf{G}_{\mathbf{u}}(Y_{\mathbf{u}}^h)$ be the elements from Proposition 5.9. Set

$$\mathcal{G}^{new} = \mathrm{Gl}(\mathcal{G}', \varphi \circ \alpha).$$

Claim. The **G**-bundle \mathcal{G}^{new} is trivial over \mathbb{P}^1_U . Indeed, by Lemmas 5.5 and 5.6 one has a chain of isomorphisms of **G**_u-bundles

$$\begin{split} \mathcal{G}^{new}|_{\mathbb{P}^{1}_{\mathbf{u}}} &\cong \mathrm{Gl}_{\mathbf{u}}(\mathcal{G}'|_{\mathbb{P}^{1}_{\mathbf{u}}-Y_{\mathbf{u}}}, \varphi_{\mathbf{u}} \circ \alpha|_{\dot{Y}^{h}_{\mathbf{u}}}) \cong \\ & \mathrm{Gl}_{\mathbf{u}}(\mathcal{G}'|_{\mathbb{P}^{1}_{\mathbf{u}}-Y_{\mathbf{u}}}, \varphi_{\mathbf{u}} \circ \alpha|_{\dot{Y}^{h}_{\mathbf{u}}} \circ \beta_{\mathbf{u}}|_{\dot{Y}^{h}_{\mathbf{u}}}) = \mathrm{Gl}_{\mathbf{u}}(\mathcal{G}'|_{\mathbb{P}^{1}_{\mathbf{u}}-Y_{\mathbf{u}}}, \varphi_{\mathbf{u}} \circ \gamma_{\mathbf{u}}), \end{split}$$

which is trivial by the choice of $\gamma_{\mathbf{u}}$. The **G**-bundles $\mathcal{G}|_{\mathbb{P}^1_U - Y}$ and $\mathcal{G}^{new}|_{\mathbb{P}^1_U - Y}$ coincide by the very construction of \mathcal{G}^{new} . By Proposition 5.1, applied to $T = Z \cup Y$, the **G**-bundle \mathcal{G}^{new} is trivial. Whence the claim.

The claim above implies that the **G**-bundle $\mathcal{G}|_{\mathbb{P}^1_U-Y} = \mathcal{G}^{new}|_{\mathbb{P}^1_U-Y}$ is trivial. Theorem 3 is proved.

5.7. End of proof of Theorem 3: proof of property (ii) from the outline. In the remaining part of Section 5 we will prove Proposition 5.9. This will complete the proof of Theorem 3.

By our assumption on Y, the group scheme $\mathbf{G}_Y = \mathbf{G} \times_U Y$ is **quasi-split**. Thus we can and will choose a Borel subgroup scheme \mathbf{B}^+ in \mathbf{G}_Y .

Since Y is an affine scheme, by [DG, Exp. XXVI, Cor. 2.3, Th 4.3.2(a)] there is an opposite to \mathbf{B}^+ Borel subgroup scheme \mathbf{B}^- in \mathbf{G}_Y . Let \mathbf{U}^+ be the unipotent radical of \mathbf{B}^+ , and let \mathbf{U}^- be the unipotent radical of \mathbf{B}^- .

Definition 5.10. We will write **E** for the functor, sending a Y-scheme T to the subgroup $\mathbf{E}(T)$ of the group $\mathbf{G}_Y(T) = \mathbf{G}(T)$ generated by the subgroups $\mathbf{U}^+(T)$ and $\mathbf{U}^-(T)$ of the group $\mathbf{G}_Y(T) = \mathbf{G}(T)$.

Lemma 5.11. The functor \mathbf{E} has the property that for every closed subscheme S in an affine Y-scheme T the induced map $\mathbf{E}(T) \to \mathbf{E}(S)$ is surjective.

Proof. The restriction maps $\mathbf{U}^{\pm}(T) \to \mathbf{U}^{\pm}(S)$ are surjective, since \mathbf{U}^{\pm} are isomorphic to vector bundles as Y-schemes (see [DG, Exp. XXVI, Cor. 2.5]).

Recall that (Y^h, π, s) is the henselization of the pair (\mathbb{A}^1_U, Y) . Also, $in : \mathbb{A}^1_U \to \mathbb{P}^1_U$ is the embedding. Denote the projection $\mathbb{A}^1_U \to U$ by pr and the projection $\mathbb{A}^1_Y \to Y$ by pr_Y .

Lemma 5.12. [FP] There is a morphism $r: Y^h \to Y$ making the following diagram commutative

(3)
$$Y^{h} \xrightarrow{r} Y$$
$$\downarrow pr|_{T}$$
$$\mathbb{P}^{1}_{U} \xrightarrow{pr} U$$

and such that $r \circ s = \mathrm{Id}_Y$.

We view Y^h as a Y-scheme via r. Thus various subschemes of Y^h also become Y-schemes. In particular, \dot{Y}^h and $\dot{Y}^h_{\mathbf{u}}$ are Y-schemes, and we can consider

 $\mathbf{E}(\dot{Y}^h) \subset \mathbf{G}(\dot{Y}^h) \ \text{ and } \ \mathbf{E}(\dot{Y}^h_\mathbf{u}) \subset \mathbf{G}(\dot{Y}^h_\mathbf{u}) = \mathbf{G}_\mathbf{u}(\dot{Y}^h_\mathbf{u}).$

Lemma 5.13.

$$\mathbf{G}_{\mathbf{u}}(\dot{Y}_{\mathbf{u}}^h) = \mathbf{E}(\dot{Y}_{\mathbf{u}}^h)\mathbf{G}_{\mathbf{u}}(Y_{\mathbf{u}}^h).$$

Proof. Firstly, one has $Y_{\mathbf{u}} = \coprod_{u \in \mathbf{u}} \coprod_{y \in Y_u} y$. (Note that Y_u is a finite scheme.) Thus by property (ii) of henselization, we have

$$Y^{h}_{\mathbf{u}} = \coprod_{u \in \mathbf{u}} \coprod_{y \in Y_{u}} y^{h}, \qquad \dot{Y}^{h}_{\mathbf{u}} = \coprod_{u \in \mathbf{u}} \coprod_{y \in Y_{u}} \dot{y}^{h},$$

where (y^h, π_y, s_y) is the henselization of the pair $(\mathbb{A}^1_u, y), \dot{y}^h := y^h - s_y(y)$. We see that y^h and \dot{y}^h are subschemes of Y^h , so we can view them as Y-schemes, and $\mathbf{G}_{y^h} := \mathbf{G}_Y \times_Y y^h$ is **quasi-split**. Also, $\mathbf{E}(\dot{y}^h)$ makes sense as a subgroup of $\mathbf{G}(\dot{y}^h) = \mathbf{G}_u(\dot{y}^h) = \mathbf{G}_{y^h}(\dot{y}^h)$. One has

$$\begin{split} \mathbf{G}_{\mathbf{u}}(\dot{Y}_{\mathbf{u}}^{h}) &= \prod_{u \in \mathbf{u}} \prod_{y \in Y_{u}} \mathbf{G}_{u}(\dot{y}^{h}) = \prod_{u \in \mathbf{u}} \prod_{y \in Y_{u}} \mathbf{G}_{y^{h}}(\dot{y}^{h}), \\ \mathbf{E}(\dot{Y}_{\mathbf{u}}^{h}) &= \prod_{u \in \mathbf{u}} \prod_{y \in Y_{u}} \mathbf{E}(\dot{y}^{h}), \\ \mathbf{G}_{\mathbf{u}}(Y_{\mathbf{u}}^{h}) &= \prod_{u \in \mathbf{u}} \prod_{y \in Y_{u}} \mathbf{G}_{u}(y^{h}) = \prod_{u \in \mathbf{u}} \prod_{y \in Y_{u}} \mathbf{G}_{y^{h}}(y^{h}). \end{split}$$

Thus it suffices for each $u \in \mathbf{u}$ and each $y \in Y_u$ to check the equality

$$\mathbf{G}_{y^h}(\dot{y}^h) = \mathbf{E}(\dot{y}^h)\mathbf{G}_{y^h}(y^h).$$

This equality holds by Fait 4.3 and Lemma 4.5 of [Gil2]. In fact, $y^h = \operatorname{Spec} \mathcal{O}$, where $\mathcal{O} = k(u)[t]^h_{\mathfrak{m}_y}$ is a henselian discrete valuation ring, and $\mathfrak{m}_y \subset k(u)[t]$ is the maximal ideal defining the point $y \in \mathbb{A}^1_u$. Further, $\dot{y}^h = \operatorname{Spec} L$, where L is the fraction field of \mathcal{O} . The lemma is proved.

We have the closed embedding (2) and the scheme \dot{Y}^h is affine by Proposition 5.2. Recall that we regard \dot{Y}^h as a Y-scheme via the morphism $r|_{\dot{Y}^h}$. Thus by Lemma 5.11 the restriction map $\mathbf{E}(\dot{Y}^h) \to \mathbf{E}(\dot{Y}^h_{\mathbf{u}})$ is surjective. Since $\mathbf{E}(\dot{Y}^h) \subset \mathbf{G}(\dot{Y}^h)$, the proposition 5.9 follows. This completes the proof of Theorem 3.

6. AN APPLICATION

The following result is a straightforward consequence of Theorem 1 and an exact sequence for étale cohomology. Recall that by our definition a reductive group scheme has geometrically connected fibres.

Theorem 4. Let R be as in Theorem 1 and **G** be a reductive R-group scheme. Let $\mu : \mathbf{G} \to \mathbf{T}$ be a group scheme morphism to an R-torus **T** such that μ is locally in the étale topology on Spec R surjective. Assume further that the R-group scheme $\mathbf{H} := \text{Ker}(\mu)$ is reductive. Let K be the fraction field of R. Then the group homomorphism

$$\mathbf{T}(R)/\mu(\mathbf{G}(R)) \to \mathbf{T}(K)/\mu(\mathbf{G}(K))$$

is injective.

This theorem extends all the known results of this form proven in [CTO], [PS1], [Zai], [OPZ].

Appendix A. [FP]

For a commutative ring A we denote by $\operatorname{Rad}(A)$ its Jacobson ideal. The following definition one can find in [Gab, Section 0].

Definition A.1. If I is an ideal in a commutative ring A, then the pair (A, I) is called *henselian*, if $I \subset \text{Rad}(A)$ and for every two relatively prime monic polynomials $\overline{g}, \overline{h} \in \overline{A}[t]$, where $\overline{A} = A/I$, and monic lifting $f \in A[t]$ of $\overline{g}\overline{h}$, there exist monic liftings $g, h \in A[t]$ such that f = gh. (Two polynomials are called relatively prime, if they generate the unit ideal.)

Lemma A.2. [FP] Let (A, I) be a henselian pair with a semi-local ring A and $J \subset A$ be an ideal. Then the pair (A/J, (I+J)/J) is henselian.

The following definition one can find in [Gab, Section 0].

Definition A.3. The henselization of any pair (A, I) is the pair (A_I^h, I^h) (over (A, I)) defined as follows

 $(A_I^h, I^h) :=$ the filtered inductive limit over the category \mathcal{N} of $(A', \operatorname{Ker}(\sigma))$,

where \mathcal{N} is the filtered category of pairs (A', σ) such that A' is an étale A-algebra and $\sigma \in \operatorname{Hom}_{A-alg}(A', A/I)$.

References

- [Che] Vladimir Chernousov. Variations on a theme of groups splitting by a quadratic extension and Grothendieck-Serre conjecture for group schemes F_4 with trivial g_3 invariant. *Doc. Math.*, (Extra volume: Andrei A. Suslin sixtieth birthday):147–169, 2010.
- [CTO] Jean-Louis Colliot-Thélène and Manuel Ojanguren. Espaces principaux homogènes localement triviaux. Inst. Hautes Études Sci. Publ. Math., (75):97–122, 1992.
- [CTS] Jean-Louis Colliot-Thélène and Jean-Jacques Sansuc. Principal homogeneous spaces under flasque tori: applications. J. Algebra, 106(1):148–205, 1987.
- [DG] Michel Demazure and Alexander Grothendieck. Schémas en groupes. III: Structure des schémas en groupes réductifs. Séminaire de Géométrie Algébrique du Bois Marie 1962/64 (SGA 3). Dirigé par M. Demazure et A. Grothendieck. Lecture Notes in Mathematics, Vol. 153. Springer-Verlag, Berlin, 1970.
- [FP] Fedorov, R.; Panin, I. A proof of Grothendieck-Serre conjecture on principal bundles over a semilocal regular ring containing an infinite field, Preprint, April 2013, http://www.arxiv.org/abs/1211.2678v2.
- [Gab] Ofer Gabber. Affine analog of the proper base change theorem. Israel J. Math., 87(1-3):325–335, 1994.
- [Gil1] Philippe Gille. Torseurs sur la droite affine. Transform. Groups, 7(3):231–245, 2002.
- [Gil2] Philippe Gille. Le problème de Kneser-Tits. Astérisque, (326):Exp. No. 983, vii, 39–81 (2010), 2009. Séminaire Bourbaki. Vol. 2007/2008.
- [Gro1] Alexander Grothendieck. Torsion homologique et sections rationnelles. In Anneaux de Chow et applications, Séminaire Claude Chevalley, number 3. Paris, 1958.
- [Gro2] Alexander Grothendieck. Le groupe de Brauer. II. Théorie cohomologique. In Dix Exposés sur la Cohomologie des Schémas, pages 67–87. North-Holland, Amsterdam, 1968.
- [Gro3] Alexander Grothendieck. Technique de descente et théorèmes d'existence en géometrie algébrique. I. Généralités. Descente par morphismes fidèlement plats. In Séminaire Bourbaki, Vol. 5, Exp. No. 190., pages 299–327. Soc. Math. France, Paris, 1995.
- [Nis1] Yevsey Nisnevich. Espaces homogènes principaux rationnellement triviaux et arithmétique des schémas en groupes réductifs sur les anneaux de Dedekind. C. R. Acad. Sci. Paris Sér. I Math., 299(1):5–8, 1984.
- [Nis2] Yevsey Nisnevich. Rationally trivial principal homogeneous spaces, purity and arithmetic of reductive group schemes over extensions of two-dimensional regular local rings. C. R. Acad. Sci. Paris Sér. I Math., 309(10):651–655, 1989.
- [Oja1] Manuel Ojanguren. Quadratic forms over regular rings. J. Indian Math. Soc. (N.S.), 44(1-4):109-116 (1982), 1980.
- [Oja2] Manuel Ojanguren. Unités représentées par des formes quadratiques ou par des normes réduites. In Algebraic K-theory, Part II (Oberwolfach, 1980), volume 967 of Lecture Notes in Math., pages 291–299. Springer, Berlin, 1982.
- [OP] Manuel Ojanguren and Ivan Panin. Rationally trivial Hermitian spaces are locally trivial. Math. Z., 237(1):181–198, 2001.
- [OPZ] M. Ojanguren, I. Panin, and K. Zainoulline. On the norm principle for quadratic forms. J. Ramanujan Math. Soc., 19(4):289–300, 2004.
- [Pa1] I. Panin. On Grothendieck–Serre's conjecture concerning principal G-bundles over reductive group schemes containing a finite field: I. ArXiv e-prints, 0905.1418v3, April 2013.
- [Pa2] Ivan Panin. On Grothendieck–Serre's conjecture concerning principal G-bundles over reductive group containing a finite field schemes:II. ArXiv e-prints, 0905.1423v3, April 2013.

- [Pan1] Panin, I. On Grothendieck-Serre conjecture concerning principal G-bundles over regular semi-local domains containing a finite field: I, Preprint, May 2014.
- [Pan2] Panin, I. On Grothendieck-Serre conjecture concerning principal G-bundles over regular semi-local domains containing a finite field: II, Preprint, May 2014.
- [Pop] Dorin Popescu. General Néron desingularization and approximation. Nagoya Math. J., 104:85–115, 1986.
- [Poo] Poonen, B., Bertini theorems over finite fields, Annals of Mathematics, 160 (2004), 1099 -1127.
- [PPS] I. Panin, V. Petrov, and A. Stavrova. Grothendieck-Serre conjecture for adjoint groups of types E_6 and E_7 and for certain classical groups. ArXiv e-prints, 0905.1427, December 2009.
- [PS1] Ivan A. Panin and Andrei A. Suslin. On a conjecture of Grothendieck concerning Azumaya algebras. St. Petersburg Math. J., 9(4):851–858, 1998.
- [PS2] Victor Petrov and Anastasia Stavrova. Grothendieck-Serre conjecture for groups of type F_4 with trivial f_3 invariant. ArXiv e-prints, 0911.3132, November 2009.
- [PSV] I. Panin, A. Stavrova, and N. Vavilov. On Grothendieck–Serre's conjecture concerning principal G-bundles over reductive group schemes: I. ArXiv e-prints, 0905.1418v3, April 2013.
- [Rag1] Madabusi S. Raghunathan. Principal bundles admitting a rational section. Invent. Math., 116(1-3):409–423, 1994.
- [Rag2] Madabusi S. Raghunathan. Erratum: "Principal bundles admitting a rational section" [Invent. Math. 116 (1994), no. 1-3, 409–423; MR1253199 (95f:14093)]. Invent. Math., 121(1):223, 1995.
- [Ser] Jean-Pierre Serre. Espaces fibrés algébrique. In Anneaux de Chow et applications, Séminaire Claude Chevalley, number 3. Paris, 1958.
- [Swa] Richard G. Swan. Néron-Popescu desingularization. In Algebra and geometry (Taipei, 1995), volume 2 of Lect. Algebra Geom., pages 135–192. Int. Press, Cambridge, MA, 1998.
- [Zai] Kirill Zainoulline. On Grothendieck conjecture about principal homogeneous spaces for some classical algebraic groups. St. Petersburg Math. J., 12(1):117–143, 2001.

E-mail address: paniniv@gmail.com

Steklov Institute of Mathematics at St.-Petersburg, Fontanka 27, St.-Petersburg 191023, Russia