

# A NEW PROOF FOR THE BORNOLGICITY OF THE SPACE OF SLOWLY INCREASING FUNCTIONS

JULIAN LARCHER AND JOCHEN WENGENROTH

ABSTRACT. A. Grothendieck proved at the end of his thesis that the space  $\mathcal{O}_M$  of slowly increasing functions and the space  $\mathcal{O}'_C$  of rapidly decreasing distributions are bornological. Grothendieck's proof relies on the isomorphy of these spaces to a sequence space and we present the first proof that does not utilize this fact by using homological methods and, in particular, the derived projective limit functor.

## 1. INTRODUCTION AND NOTATION

In [Sch66, p. 243] L. Schwartz introduced the space of multipliers of temperate distributions, i.e., the space of slowly increasing functions

$$\mathcal{O}_M = \{f \in \mathcal{C}^\infty(\mathbb{R}^d); \forall \alpha \in \mathbb{N}_0^d \exists N \in \mathbb{N} : \langle x \rangle^{-N} \partial^\alpha f \in L^\infty\},$$

where  $\mathcal{C}^\infty(\mathbb{R}^d)$  is the space of complex valued, infinitely differentiable functions on  $\mathbb{R}^d$ ,  $\langle x \rangle = 1 + |x|^2$ ,  $\partial^\alpha$  is the partial derivative, and  $L^\infty$  is the Lebesgue space of bounded functions. The dual  $\mathcal{O}'_M$  of  $\mathcal{O}_M$  is the space of very rapidly decreasing distributions.

Schwartz also introduced the space of convolutors of temperate distributions, i.e., the space  $\mathcal{O}'_C$  of rapidly decreasing distributions, which is the dual of the space

$$\mathcal{O}_C = \{f \in \mathcal{C}^\infty(\mathbb{R}^d); \exists N \forall \alpha \in \mathbb{N}_0^d : \langle x \rangle^{-N} \partial^\alpha f \in L^\infty\}$$

of very slowly increasing functions. These spaces are related as in the diagram

$$\begin{array}{ccc} \mathcal{O}_C & \subseteq & \mathcal{O}_M \\ \Downarrow & & \Downarrow \\ \mathcal{O}'_M & \subseteq & \mathcal{O}'_C \end{array}$$

where in both cases the Fourier transform can be taken as the isomorphism.

It is comparatively easy to see that the four spaces are nuclear and semi-reflexive, that  $\mathcal{O}_M$  and  $\mathcal{O}'_C$  are complete and that  $\mathcal{O}_C$  and  $\mathcal{O}'_M$  are (LF)-spaces and hence bornological. But the completeness of  $\mathcal{O}_C$  and  $\mathcal{O}'_M$  and the bornologicity of  $\mathcal{O}_M$  and  $\mathcal{O}'_C$  are not trivial (which was even asserted by Grothendieck, [Gro55, Chap. II, p. 130]). Since the dual of a bornological space is complete and the dual of a complete nuclear space is bornological, these two problems are equivalent (for the definitions of these topological properties and relations between them see [Itō87, Section 424]).

Grothendieck proved that  $\mathcal{O}_M$  is bornological by showing that it is isomorphic to a complemented subspace of the sequence space  $s \hat{\otimes}_\pi s'$  [Gro55, Chap. II, Lemme 18, p. 132] and verified “directly” that the space  $s \hat{\otimes}_\pi s'$  is bornological [Gro55, Chap. II, Prop. 15, p. 125, Cor. 2, p. 128]. We will find out more about this isomorphy in

---

2000 *Mathematics Subject Classification.* Primary 46M18 · 46F05 ; Secondary 46A08 · 46A45.  
*Key words and phrases.* derived projective limit functor · slowly increasing functions · rapidly decreasing distributions · sequence space representations.

Section 2 and also give a homological proof of the bornologicity of  $s \hat{\otimes}_\pi s'$ .

In [Kuc85], J. Kučera claimed to have presented a new (and simple) proof for the main properties of the space  $\mathcal{O}_M$ . That Kučera's proof contains severe mistakes and that it is based on incorrect propositions is clarified in [Lar12], where also the lack of a proof of the bornologicity of  $\mathcal{O}_M$ , that does not use the isomorphism  $\mathcal{O}_M \cong s \hat{\otimes}_\pi s'$ , is pointed out. In Section 3 we will give such a proof.

## 2. PROJECTIVE LIMITS AND THE SPACE $s \hat{\otimes}_\pi s'$

Since quotients (and, in particular, complemented subspaces) of bornological spaces are bornological, it was sufficient for Grothendieck to prove that  $\mathcal{O}_M$  is isomorphic to a complemented subspace of  $s \hat{\otimes}_\pi s'$ , where  $s$  is the space of rapidly decreasing sequences

$$s = \{(x_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}}; \forall k : \sup_{j \in \mathbb{N}} j^k |x_j| < \infty\}$$

and  $s'$  is its dual, the space of slowly increasing sequences

$$s' = \{(x_j)_{j \in \mathbb{N}} \in \mathbb{C}^{\mathbb{N}}; \exists k : \sup_{j \in \mathbb{N}} j^{-k} |x_j| < \infty\}.$$

By  $s \otimes_\pi s'$  we denote the completed projective tensor product of these spaces. E.g., by [Bar12, Remark 1, p. 321], this space  $s \hat{\otimes}_\pi s'$  is canonically isomorphic to

$$s \hat{\otimes}_\pi s' \cong \{x \in \mathbb{C}^{\mathbb{N} \times \mathbb{N}}; \forall n \exists N : \sup_{i,j} i^n j^{-N} |x_{i,j}| < \infty\}.$$

In [Val81], M. Valdivia proved that  $\mathcal{O}_M$  is even isomorphic to  $s \hat{\otimes}_\pi s'$  itself which answered a question posed in [Gro55, Chap. II, p. 134]. C. Bargetz used this fact, the bornologicity of  $s \hat{\otimes}_\pi s'$ , and methods of the theory of topological tensor products to obtain the isomorphism  $\mathcal{O}_C \cong s \hat{\otimes}_l s'$  [Bar12, Prop. 1, p. 318].

The descriptions of the spaces  $\mathcal{O}_M$  and  $s \hat{\otimes}_\pi s'$  already indicate how they can be written as projective limits of LB-spaces (countable inductive limits of Banach spaces)

$$(1) \quad \mathcal{O}_M = \bigcap_{n \in \mathbb{N}} X_n = \bigcap_{n \in \mathbb{N}} \bigcup_{N \in \mathbb{N}} X_{n,N},$$

$$(2) \quad s \hat{\otimes}_\pi s' = \bigcap_{n \in \mathbb{N}} Y_n = \bigcap_{n \in \mathbb{N}} \bigcup_{N \in \mathbb{N}} Y_{n,N},$$

where  $X_{n,N}$  and  $Y_{n,N}$  are the Banach spaces

$$X_{n,N} = \{f \in \mathcal{C}^n(\mathbb{R}^d); \|f\|_{n,N} = \sup_{x \in \mathbb{R}^d, |\alpha| \leq n} \langle x \rangle^{-N} |\partial^\alpha f(x)| < \infty\},$$

$$Y_{n,N} = \{x \in \mathbb{C}^{\mathbb{N} \times \mathbb{N}}; \|x\|_{n,N} = \sup_{i,j} i^n j^{-N} |x_{i,j}| < \infty\}.$$

These representations as projective limits of LB-spaces are not only natural but also extremely useful since there are very good criteria for checking bornologicity. They are related to the derived projective limit functor  $\text{Proj}^1 \mathcal{X}$  (which can be defined as the cokernel of the map  $\prod X_n \rightarrow \prod X_n, (x_n)_n \mapsto (x_n - \varrho_{n+1}^n(x_{n+1}))_n$  where  $\varrho_m^n$  are the connecting maps of the projective spectrum  $\mathcal{X}$ , in our cases,  $\varrho_m^n$  are just inclusions). Indeed, an unpublished theorem of D. Vogt (his proof reproduced in [Wen03, Th. 3.3.4]) says that  $\text{Proj} \mathcal{X}$  is bornological whenever  $\text{Proj}^1 \mathcal{X} = 0$ . Moreover, there is a variety of evaluable conditions ensuring  $\text{Proj}^1 \mathcal{X} = 0$ . We are going to apply the following results of Palamodov-Retakh [Pal71] and the second named author, respectively:

A specturm  $\mathcal{X}$  of LB-spaces satisfies  $\text{Proj}^1 \mathcal{X} = 0$  if and only if there are Banach discs  $D_n$  in  $X_n$  with  $\varrho_m^n(D_m) \subseteq D_n$  and

$$\forall n \in \mathbb{N} \exists m \geq n \forall k \geq m : \varrho_m^n(X_m) \subseteq \varrho_k^n(X_k) + D_n.$$

The requirement  $\varrho_m^n(D_m) \subseteq D_n$  is sometimes very easy to fulfil but in many cases it is very inconvenient. It can be omitted if either all steps  $X_n$  are LS-spaces (i.e., the inclusions  $X_{n,N} \hookrightarrow X_{n,N+1}$  are compact) or if a slightly stronger condition of Palamodov-Retakh type is required. Denoting by  $\varrho_\infty^n : \text{Proj} \mathcal{X} \rightarrow X_n$  the obvious map we have:

A specturm  $\mathcal{X}$  of LB-spaces satisfies  $\text{Proj}^1 \mathcal{X} = 0$  if and only if, for every  $n \in \mathbb{N}$ , there are a Banach discs  $D_n$  in  $X_n$  and  $m \geq n$  with

$$\varrho_m^n(X_m) \subseteq \varrho_\infty^n(\text{Proj} \mathcal{X}) + D_n.$$

We refer to [Wen03] for the proofs of these characterization and much more information about derived functors. Typically, the decompositions required in conditions of Retakh-Palamodov type are quite easy to produce in the case of spaces of sequences (or matrices) since one can write  $x = \chi x + (1 - \chi)x$  where  $\chi$  is the indicator function of a suitably chosen set. We want to exemplify this by giving a very short proof for the bornologicity of  $s \hat{\otimes}_\pi s'$  (which is similar to Vogt's proof of  $\text{Ext}^1(s, s) = 0$  [Vog84, Lemma 2.1, p. 359]).

**Proposition 1.** *The space  $s \hat{\otimes}_\pi s'$  is bornological.*

*Proof.* We keep the notation  $s \hat{\otimes}_\pi s' \cong \bigcap_{n \in \mathbb{N}} Y_n = \bigcap_{n \in \mathbb{N}} \bigcup_{N \in \mathbb{N}} Y_{n,N}$  from above and we will verify the Palamodov-Retakh condition for the unit balls  $D_n$  of  $Y_{n,0}$  which trivially satisfy  $D_{n+1} \subseteq D_n$ . For  $n \in \mathbb{N}$  we take  $m = n + 1$  and fix  $x \in Y_n$  as well as  $k \geq n + 1$ . Since  $x \in Y_{m,M}$  for some  $M \in \mathbb{N}$  we have

$$\|x\|_{m,M} = \sup_{i,j} i^m j^{-N} |x_{i,j}| = c < \infty.$$

We set  $y_{i,j} = x_{i,j}$  if  $i < cj^M$  and  $y_{i,j} = 0$  else, as well as  $z = x - y$ . For  $i < cj^m$  we have  $z_{i,j} = 0$  and for  $i \geq cj^M$  we estimate

$$i^n j^{-0} |z_{i,j}| = i^m j^{-M} |z_{i,j}| j^M / i \leq \|x\|_{m,M} / c = 1$$

which proves  $z \in D_n$ . It remains to show  $y \in Y_{k,K}$  for  $K$  sufficiently large. Indeed, for  $K = M(k - m + 1)$  we have  $y_{i,j} = 0$  if  $i \geq cj^M$  and if  $i < cj^M$  we estimate

$$i^k j^{-K} |y_{i,j}| = i^m j^{-M} |y_{i,j}| i^{k-m} j^{M-K} \leq \|x\|_{m,M} c^{k-m} j^{(k-m)M+M-K} = c^{k-m+1}.$$

This proves  $\|y\|_{k,K} < \infty$ , as required.  $\square$

### 3. THE NEW PROOF

Now we want to prove  $\text{Proj}^1 \mathcal{X} = 0$  for the spectrum  $\mathcal{X} = (X_n)_{n \in \mathbb{N}}$  in (1) in order to obtain that  $\mathcal{O}_M$  is bornological. Splitting up a given function  $f \in X_m$  as  $f = \chi f + (1 - \chi)f$  with a cut-off function  $\chi$  (as in the proof of Proposition 1) does not work in this case. But we will see how  $f$  can be ‘‘split up’’ in the following proof of Grothendieck's result.

**Proposition 2.** *The space  $\mathcal{O}_M$  is bornological.*

*Proof.* To obtain  $\text{Proj}^1 \mathcal{X} = 0$  we will show

$$(3) \quad \forall n \exists m, N : X_m \subseteq \mathcal{O}_M + B_{n,N}$$

where  $B_{n,N}$  is the unit ball of  $X_{n,N}$ . This condition means that we have to approximate every  $f \in X_m$  with respect to the norm  $\|\cdot\|_{n,N}$  by elements of  $\mathcal{O}_M$ . To achieve such an approximation we use a kernel  $K \in \mathcal{O}_M(\mathbb{R}^d \times \mathbb{R}^d)$  satisfying

$$K \geq 0, \quad \int_{\mathbb{R}^d} K(t, x) dt = 1 \text{ for all } x \in \mathbb{R}^d, \quad \text{and}$$

$$\text{supp } K(\cdot, x) \subseteq \prod_{j=1}^d [x_j, x_j + \varepsilon \langle x \rangle^{-\mu}] =: A_x \text{ for all } x \in \mathbb{R}^d$$

where we will see later how  $\varepsilon$  and  $\mu$  have to be chosen in dependence on  $f \in X_m$ . We can obtain such a kernel by defining

$$K(t, x) = \varepsilon^{-d} \langle x \rangle^{\mu d} \varphi(\varepsilon^{-1} \langle x \rangle^\mu (t - x))$$

for a positive test function  $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$  with support in  $[0, 1]^d$  and  $\int_{\mathbb{R}^d} \varphi(t) dt = 1$  (the conditions above can be checked easily and  $K \in \mathcal{O}_M$  since every derivative of  $K$  can be estimated by a polynomial).

We start with the one-dimensional case  $d = 1$  where we can take  $m = n + 1$  and  $N = 0$ . So let  $f \in X_{n+1, M}$  for some  $M \in \mathbb{N}$ . We want to find  $g \in \mathcal{O}_M$  such that  $f - g \in B_{n,0}$ . At first we set

$$g_n(x) = \int_{\mathbb{R}} f^{(n)}(t) K(t, x) dt$$

and show that this is a good approximation to  $f^{(n)}$ . Since for  $l \in \mathbb{N}_0$

$$\left| g_n^{(l)}(x) \right| = \left| \int_{A_x} f^{(n)}(t) \partial_x^l K(t, x) dt \right| \leq \int_{A_x} |P(t)| |Q(t, x)| dt \leq |R(x)|$$

for some polynomials  $P, Q, R$ , the function  $g_n$  is contained in  $\mathcal{O}_M$ . Furthermore we can estimate in virtue of Taylor's formula

$$\left| f^{(n)}(t) - f^{(n)}(x) \right| \leq |t - x| \langle \xi(t, x) \rangle^M \|f\|_{n+1, M}$$

with a point  $\xi(t, x)$  between  $t$  and  $x$ . For  $\varepsilon$  small enough the inequality  $\langle \xi(t, x) \rangle \leq 2\langle x \rangle$  holds for every  $x \in \mathbb{R}$  and  $t \in A_x$ . We obtain

$$\begin{aligned} |g_n(x) - f^{(n)}(x)| &= \left| \int_{\mathbb{R}} \left( f^{(n)}(t) - f^{(n)}(x) \right) K(t, x) dt \right| \\ &\leq \int_{A_x} \left| f^{(n)}(t) - f^{(n)}(x) \right| K(t, x) dt \\ (4) \quad &\leq \int_{A_x} |t - x| \langle \xi(t, x) \rangle^M \|f\|_{n+1, M} K(t, x) dt \\ &\leq \varepsilon 2^M \langle x \rangle^{M-\mu} \|f\|_{n+1, M} \int_{A_x} K(t, x) dt \\ &= \varepsilon 2^M \langle x \rangle^{M-\mu} \|f\|_{n+1, M}. \end{aligned}$$

Now if

$$T : \mathcal{O}_M(\mathbb{R}) \rightarrow \mathcal{O}_M(\mathbb{R}), h \mapsto \left( x \mapsto \int_0^x h(t) dt \right),$$

we can set

$$g(x) = \sum_{j=0}^{n-1} \frac{f^{(j)}(0)}{j!} x^j + (T^n g_n)(x).$$

Then  $g \in \mathcal{O}_M$  and since

$$(T^n f^{(n)})(x) = f(x) - \sum_{j=0}^{n-1} \frac{f^{(j)}(0)}{j!} x^j,$$

integrating (4) (the integral starting at 0) yields

$$|g^{(l)}(x) - f^{(l)}(x)| \leq 1, \quad x \in \mathbb{R}^d, l \leq n$$

for  $\varepsilon$  small enough and  $\mu$  large enough. Hence  $g - f \in B_{n,0}$  and the proof is complete for the one-dimensional case.

Now we will prove the two-dimensional case  $d = 2$ . We set  $m = 2n + 1$  and  $N = n - 1$  in (3). So let  $f \in X_{2n+1,M}$  for some  $M$ . With the help of a kernel  $K \in \mathcal{O}_M(\mathbb{R}^2 \times \mathbb{R}^2)$  like above, we set

$$g_n(x) = \int_{\mathbb{R}^2} \partial^{(n,n)} f(t) K(t, x) dt$$

in order to approximate  $\partial^{(n,n)} f$  by  $g_n$ . Similar to the one-dimensional case we have

$$\left| \partial^{(n,n)} f(t) - \partial^{(n,n)} f(x) \right| \leq c \cdot |t - x| \langle \xi(t, x) \rangle^M \|f\|_{2n+1, M}$$

and  $\langle \xi(t, x) \rangle \leq 2\langle x \rangle$  for  $t \in A_x$  and  $\varepsilon$  small enough and thus

$$(5) \quad \begin{aligned} |g_n(x) - \partial^{(n,n)} f(x)| &\leq c \int_{A_x} |t - x| \langle \xi(t, x) \rangle^M \|f\|_{2n+1, M} K(t, x) dt \\ &\leq \tilde{c} \varepsilon \langle x \rangle^{M-\mu} \|f\|_{2n+1, M}. \end{aligned}$$

Let us denote  $T_j$  the integral with respect to the  $j$ -th component (the integral starting at 0). Applying  $T_1 \circ T_2$   $n$ -times to  $\partial^{(n,n)} f(x)$  yields

$$(T_1^n T_2^n f)(x) = f(x) + \sum_{\alpha < (n,n)} \partial^\alpha f(0, 0) \frac{x^\alpha}{\alpha!} - \sum_{j=0}^{n-1} \partial^{(j,0)} f(0, x_2) \frac{x_1^j}{j!} - \sum_{j=0}^{n-1} \partial^{(0,j)} f(x_1, 0) \frac{x_2^j}{j!}.$$

As in the one-dimensional case we can choose  $g_0^1, \dots, g_{n-1}^1, g_0^2, \dots, g_{n-1}^2 \in \mathcal{O}_M(\mathbb{R})$  such that  $\|g_j^1 - \partial^{(0,j)} f(\cdot, 0)\|_{n,0} \leq \varepsilon$  and  $\|g_j^2 - \partial^{(j,0)} f(\cdot, 0)\|_{n,0} \leq \varepsilon$ . Defining

$$g(x) = (T_1^n T_2^n) g_n(x) - \sum_{\alpha < (n,n)} \partial^\alpha f(0, 0) \frac{x^\alpha}{\alpha!} + \sum_{j=0}^{n-1} g_j^2(x_2) \frac{x_1^j}{j!} + \sum_{j=0}^{n-1} g_j^1(x_1) \frac{x_2^j}{j!}$$

and applying  $T_1^n T_2^n$  to (5) yields

$$\begin{aligned} |g(x) - f(x)| &\leq \\ &\varepsilon + \sum_{j=0}^{n-1} \left( \left| g_j^1(x_1) - \partial^{(0,j)} f(x_1, 0) \right| \frac{|x_2|^j}{j!} + \left| g_j^2(x_2) - \partial^{(j,0)} f(0, x_2) \right| \frac{|x_1|^j}{j!} \right) \end{aligned}$$

for  $\mu$  large enough which implies

$$|g(x) - f(x)| \leq \varepsilon + \varepsilon \sum_{j=0}^{n-1} \frac{|x_2|^j}{j!} + \varepsilon \sum_{j=0}^{n-1} \frac{|x_1|^j}{j!} \leq \varepsilon c \langle x \rangle^{n-1}$$

for some  $c > 1$ . Since similar estimates also hold for  $|\partial^\alpha g(x) - \partial^\alpha f(x)|$ ,  $|\alpha| \leq n$ , we obtain  $g - f \in B_{n,n-1}$  and the proof is complete for  $d = 2$ .

The general case  $d \in \mathbb{N}$  is very similar. Inductively we want to show

$$X_{dn+1} \subseteq \mathcal{O}_M + B_{n,(d-1)(n-1)}$$

and start by approximating  $\partial^{(n,\dots,n)} f$  by  $g_n(x) := \int_{\mathbb{R}^d} \partial^{(n,\dots,n)} f(t) K(t, x) dt$ . Then we integrate the estimate of  $g_n - \partial^{(n,\dots,n)} f$   $n$ -times with respect to each component. The integral  $T_1^n \dots T_d^n \partial^{(n,\dots,n)} f$  contains  $f$  as a summand and terms that are the product of a derivative of  $f$  that only depends on less than  $d$  components and a polynomial in less than  $d$  components with exponents less than  $n$ . But we can estimate the functions that only depend on less than  $d$  variables by the induction hypothesis and hence we can obtain  $g \in \mathcal{O}_M$  with  $g - f \in B_{n,(d-1)(n-1)}$ .  $\square$

## REFERENCES

- [Bar12] Christian Bargetz, *A sequence space representation of L. Schwartz' space  $\mathcal{O}_C$* , Arch. Math. (Basel) **98** (2012), no. 4, 317–326. MR 2914348
- [Gro55] Alexander Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955). MR 0075539 (17,763c)
- [Itō87] Kiyosi Itō (ed.), *Encyclopedic dictionary of mathematics. Vol. I–IV*, second ed., MIT Press, Cambridge, MA, 1987, Translated from the Japanese. MR 901762 (89b:00033)
- [Kuc85] Jan Kucera, *The spaces  $\mathcal{O}_M$  and  $\mathcal{O}_C$  are ultrabornological. A new proof*, Internat. J. Math. Math. Sci. **8** (1985), no. 4, 813–816. MR 821639 (87e:46054)
- [Lar12] Julian Larcher, *Some remarks concerning the spaces of multipliers and convolutors,  $\mathcal{O}_M$  and  $\mathcal{O}'_C$ , of Laurent Schwartz*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **106** (2012), no. 2, 407–417. MR 2978922
- [Pal71] Victor P. Palamodov, *Homological methods in the theory of locally convex spaces*, Uspehi Mat. Nauk **26** (1971), no. 1(157), 3–65. MR 0293365 (45 #2442)
- [Sch66] Laurent Schwartz, *Théorie des distributions*, Publications de l'Institut de Mathématique de l'Université de Strasbourg, No. IX–X. Nouvelle édition, entièrement corrigée, refondue et augmentée, Hermann, Paris, 1966. MR 0209834 (35 #730)
- [Val81] Manuel Valdivia, *A representation of the space  $\mathcal{O}_M$* , Math. Z. **177** (1981), no. 4, 463–478. MR 624226 (82i:46057)
- [Vog84] Dietmar Vogt, *Some results on continuous linear maps between Fréchet spaces*, Functional analysis: surveys and recent results, III (Paderborn, 1983), North-Holland Math. Stud., vol. 90, North-Holland, Amsterdam, 1984, pp. 349–381. MR 761391 (86i:46075)
- [Wen03] Jochen Wengenroth, *Derived functors in functional analysis*, Lecture Notes in Mathematics, vol. 1810, Springer-Verlag, Berlin, 2003. MR 1977923 (2004d:46090)

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT INNSBRUCK, A-6020 INNSBRUCK, AUSTRIA  
*E-mail address:* julian.larcher@uibk.ac.at

UNIVERSITÄT TRIER, FB IV – MATHEMATIK, 54286 TRIER, GERMANY  
*E-mail address:* wengenroth@uni-trier.de