

Complex geodesics in convex tube domains II

Sylwester Zając

Institute of Mathematics, Faculty of Mathematics and Computer Science,
Jagiellonian University, Łojasiewicza 6, 30-348 Kraków, Poland
sylwester.zajac@im.uj.edu.pl

Abstract

We give a description (direct formulas) of all complex geodesics in a convex tube domain in \mathbb{C}^n containing no complex affine lines, expressed in terms of geometric properties of the domain. We next apply that result to give formulas (a necessary condition) for extremal mappings with respect to the Lempert function and the Kobayashi-Royden metric in a big class of bounded, pseudoconvex, complete Reinhardt domains: for all of them in \mathbb{C}^2 and for those of them in \mathbb{C}^n which logarithmic image is strictly convex in geometric sense.

1 Introduction

A non-empty open set $D \subset \mathbb{C}^n$ is called a *tube domain* if $D = \Omega + i\mathbb{R}^n$ for some domain $\Omega \subset \mathbb{R}^n$. We call Ω the *base* of D and in this paper we denote it by $\operatorname{Re} D$. In the recent paper [Zaj] we investigated convex tube domains from the point of view of theory of holomorphically invariant distances. More precisely, we were interested especially in the notion of complex geodesics. Given a convex domain $D \subset \mathbb{C}^n$, we call a holomorphic map $\varphi : \mathbb{D} \rightarrow D$ a *complex geodesic* for D if there exists a *left inverse* of φ , i.e. a holomorphic function $f : D \rightarrow \mathbb{D}$ such that $f \circ \varphi = \operatorname{id}_{\mathbb{D}}$. Complex geodesics of D are exactly holomorphic isometries between the unit disc $\mathbb{D} \subset \mathbb{C}$ equipped with the Poincaré distance and the domain D equipped with the Carathéodory pseudodistance. It follows from the Lempert theorem (see [Lem] or [Jar-Pfl, Chapter 8]) that if $D \subset \mathbb{C}^n$ is a taut convex domain, then for any pair of points in D there exists a complex geodesic passing through them.

In this paper we restrict our considerations to convex tube domains containing no complex affine lines (equivalently, to convex tubes with no real affine lines contained in base). Such a family of domains is equal to the family of taut convex tube domains (see e.g. [Bra-Sor]). This approach has many advantages, among which it is worth mentioning that every holomorphic map with image lying in such a domain admits a boundary measure ([Zaj, Observation

⁽¹⁾ 2010 *Mathematics Subject Classification*: Primary: 32F45, 32A07.

Key words and phrases: Complex geodesics; tube domains; convex domains.

The author was supported by the NCN grant on the basis of the decision number DEC-2012/05/N/ST1/02911.

2.5]). What is more, from [Zaj, Observation 2.4] it follows that doing such a restriction we lose no generality.

This paper may be treated as a continuation of [Zaj]. In [Zaj] we gave an equivalent condition for a holomorphic map $\varphi : \mathbb{D} \rightarrow D$ to be a complex geodesic in a convex tube domain D containing no complex affine lines. It is stated in language of measure theory and formulated in terms of *boundary n -tuple of measures* of φ (in [Zaj] and here shortly called a *boundary measure*). Boundary measure of φ is a unique n -tuple $\mu = (\mu_1, \dots, \mu_n)$ of real Borel measures on the unit circle $\mathbb{T} \subset \mathbb{C}$ such that

$$\varphi(\lambda) = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{\zeta + \lambda}{\zeta - \lambda} d\mu(\zeta) + i \operatorname{Im} \varphi(0), \quad \lambda \in \mathbb{D}.$$

In the main result of this paper, Theorem 3.1, we present a full description of all complex geodesics for D . We derive it using the equivalent condition from [Zaj] and the following, 'spherical' decomposition of n -tuples of measures (Lemma 2.1): given real Borel measures μ_1, \dots, μ_n on \mathbb{T} , there exist a finite positive Borel measure ν on \mathbb{T} singular to the Lebesgue measure $\mathcal{L}^{\mathbb{T}}$ on \mathbb{T} , a Borel-measurable map ϱ from \mathbb{T} to the unit sphere $\partial \mathbb{B}_n$ and a map $g : \mathbb{T} \rightarrow \mathbb{R}^n$ with components in $L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$ such that

$$(\mu_1, \dots, \mu_n) = g d\mathcal{L}^{\mathbb{T}} + \varrho d\nu.$$

The objects ν , ϱ and g are in some sense unique. Theorem 3.1 states that a holomorphic map with a boundary measure $\mu = (\mu_1, \dots, \mu_n)$ is a complex geodesic for D if and only if the parts ν , ϱ and g of the decomposition of μ satisfy several geometric conditions. So, strictly speaking, in Theorem 3.1 we describe the form of every n -tuple of measures which define a complex geodesic for D . But in fact, the complex geodesic itself can be then easily recovered (up to an imaginary constant) from its boundary measure via the above integral.

Later in the paper we apply Theorem 3.1 to obtain more detailed descriptions of complex geodesics in some special classes of convex tube domains. In Subsection 3.1 we do it for, among others, convex tubes $D \subset \mathbb{C}^n$ with the base being bounded from above on each coordinate and satisfying the equality $\operatorname{Re} D + (-\infty, 0]^n = \operatorname{Re} D$. These domains are very useful in studying extremal mappings with respect to the Lempert function and the Kobayashi-Royden metric in bounded, pseudoconvex, complete Reinhardt domains in \mathbb{C}^n . We deal with this topic in Section 5, achieving formulas for extremal mappings in a big class of such Reinhardt domains: for all of them in \mathbb{C}^2 and for those of them in \mathbb{C}^n which logarithmic image is strictly convex in geometric sense (i.e. it is convex and its boundary contains no non-trivial segments). Besides, in Subsection 3.2 we investigate complex geodesics in convex tube domains in \mathbb{C}^2 . The results obtained there, together with the considerations made in Subsection 3.1, simplify the conditions from Theorem 3.1 in two-dimensional case.

Let us briefly summarize the content of the paper. In Section 2 we present the notation which is used in this paper and we recall some facts about boundary measures of holomorphic maps. There we also prove the lemma on the decomposition of n -tuples of measures, which was mentioned above. At the end of that section we define a few objects describing some geometric properties of a convex tube domain in \mathbb{C}^n . In Section 3 we formulate the main result of this paper, Theorem 3.1, we present its applications in special classes of tube domains

and we give some examples. Section 4 contains the proof of Theorem 3.1 and some additional remarks. In Section 5 we apply results from Section 3 to obtain formulas for extremal holomorphic mappings in some classes of Reinhardt domains in \mathbb{C}^n .

2 Preliminaries

Let us begin with some notation. The symbols \mathbb{D} , \mathbb{T} , \mathbb{C}_* denote respectively the unit disc in \mathbb{C} , the unit circle in \mathbb{C} and the punctured plane, namely the set $\mathbb{C} \setminus \{0\}$. By δ_{λ_0} we mean the Dirac delta at a point $\lambda_0 \in \mathbb{T}$, by χ_A we mean the characteristic function $\chi_A : \mathbb{T} \rightarrow \{0, 1\}$ of a set $A \subset \mathbb{T}$ and by e_1, \dots, e_n we mean the canonical basis of \mathbb{R}^n or \mathbb{C}^n . The Poincaré distance in \mathbb{D} is denoted by ρ . By $\langle x, y \rangle$ we mean the standard inner product of vectors $x, y \in \mathbb{R}^n$, by $\|\cdot\|$ we denote the euclidean norm in \mathbb{R}^n and by \mathbb{B}_n we mean the unit euclidean ball in \mathbb{R}^n . For a set $A \subset \mathbb{R}^n$ the symbol A^\perp denotes the set $\{v \in \mathbb{R}^n : \forall a \in A : \langle v, a \rangle = 0\}$.

We use the symbol $\langle \cdot, \cdot \rangle$ also for measures and functions. For example, if μ is a tuple (μ_1, \dots, μ_n) of real (i.e. complex with real values) Borel measures on \mathbb{T} and $v = (v_1, \dots, v_n)$ is a real vector or a bounded Borel-measurable mapping from \mathbb{T} to \mathbb{R}^n , then $\langle d\mu, v \rangle$ or $\langle v, d\mu \rangle$ is the measure $\sum_{j=1}^n v_j d\mu_j$, etc. The fact that a real measure ν is positive (resp. negative, null) is shortly denoted by $\nu \geq 0$ (resp. $\nu \leq 0$, $\nu = 0$). The variation of a complex measure ν is denoted by $|\nu|$. In this paper we consider mostly Borel measures on \mathbb{T} and we sometimes omit the word 'Borel'.

In what follows we use the following families of mappings:

$$\begin{aligned} \mathcal{H}^n &:= \{h \in \mathcal{O}(\mathbb{C}, \mathbb{C}^n) : \forall \lambda \in \mathbb{T} : \bar{\lambda}h(\lambda) \in \mathbb{R}^n\}, \\ \mathcal{H}_+^n &:= \{h \in \mathcal{O}(\mathbb{C}, \mathbb{C}^n) : \forall \lambda \in \mathbb{T} : \bar{\lambda}h(\lambda) \in [0, \infty)^n\}. \end{aligned}$$

We have

$$\mathcal{H}^n = \{h \in \mathcal{O}(\mathbb{C}, \mathbb{C}^n) : \exists a \in \mathbb{C}^n, b \in \mathbb{R}^n : h(\lambda) = \bar{a}\lambda^2 + b\lambda + a, \lambda \in \mathbb{C}\}.$$

Moreover (see e.g. [Jar-Pfl, Lemma 8.4.6]),

$$\mathcal{H}_+^1 = \{h \in \mathcal{O}(\mathbb{C}) : \exists c \geq 0, d \in \mathbb{D} : h(\lambda) = c(\lambda - d)(1 - \bar{d}\lambda), \lambda \in \mathbb{C}\}.$$

In particular, for $h \in \mathcal{H}_+^1$ we have $\bar{\lambda}h(\lambda) = c|\lambda - d|^2$, $\lambda \in \mathbb{T}$, so such a function h has at most one zero on \mathbb{T} (counting without multiplicities).

In this paper we sometimes consider linear dependence or independence of functions $h_1, \dots, h_m \in \mathcal{H}^1$. Note that here it does not matter whether it is meant over the field \mathbb{R} or \mathbb{C} , because these two properties are equivalent, in view of the fact that $\bar{\lambda}h_j(\lambda) \in \mathbb{R}$ for $\lambda \in \mathbb{T}$, $j = 1, \dots, m$.

Now we recall some facts on boundary measures of holomorphic maps. A real Borel measure μ on \mathbb{T} is called *boundary measure* of a holomorphic function $\varphi : \mathbb{D} \rightarrow \mathbb{C}$, if

$$(1) \quad \varphi(\lambda) = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{\zeta + \lambda}{\zeta - \lambda} d\mu(\zeta) + i\text{Im } \varphi(0), \quad \lambda \in \mathbb{D},$$

or equivalently, taking the real parts in this equality, if

$$(2) \quad \text{Re } f(\lambda) = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{1 - |\lambda|^2}{|\zeta - \lambda|^2} d\mu(\zeta), \quad \lambda \in \mathbb{D}.$$

Such a measure μ is uniquely determined by φ . In the case when φ is a map, namely $\varphi = (\varphi_1, \dots, \varphi_n) \in \mathcal{O}(\mathbb{D}, \mathbb{C}^n)$, by *boundary measure* of φ we mean a unique n -tuple (μ_1, \dots, μ_n) of real Borel measures on \mathbb{T} such that μ_j is the boundary measure for φ_j for every $j = 1, \dots, n$. Then formulas analogous to (1) and (2) hold for φ .

Denote

$$\mathcal{M}^n := \{\varphi \in \mathcal{O}(\mathbb{D}, \mathbb{C}^n) : \varphi \text{ admits a boundary measure}\}.$$

Not every holomorphic function on \mathbb{D} admits a boundary measure and hence $\mathcal{M}^n \subsetneq \mathcal{O}(\mathbb{D}, \mathbb{C}^n)$. It is very important that if $D \subset \mathbb{C}^n$ is a convex tube domain containing no complex affine lines, then every holomorphic map $\varphi : \mathbb{D} \rightarrow D$ belongs to \mathcal{M}^n (see [Zaj, Observation 2.5]). In that case for $\mathcal{L}^\mathbb{T}$ -almost every $\lambda \in \mathbb{T}$ the radial limit $\varphi^*(\lambda) = \lim_{r \rightarrow 1^-} \varphi(r\lambda)$ of φ exists and belongs to \overline{D} .

It is worth to recall that if μ is a boundary measure of a holomorphic function $\varphi \in \mathcal{M}$, then μ is a weak-* limit of measures $\operatorname{Re} \varphi(r\lambda) d\mathcal{L}^\mathbb{T}(\lambda)$, when $r \rightarrow 1^-$ (see e.g. [Koo, p. 10]). Here we treat complex measures as linear functionals on $\mathcal{C}(\mathbb{T})$, the space of all complex-valued continuous functions on \mathbb{T} equipped with the supremum norm. The weak-* convergence which we mentioned means that

$$\int_{\mathbb{T}} u(\lambda) \operatorname{Re} \varphi(r\lambda) d\mathcal{L}^\mathbb{T}(\lambda) \xrightarrow{r \rightarrow 1^-} \int_{\mathbb{T}} u(\lambda) d\mu(\lambda), \quad u \in \mathcal{C}(\mathbb{T}).$$

We need the following fact: if μ is a boundary measure of a function $\varphi \in \mathcal{M}$ and $\mu = g d\mathcal{L}^\mathbb{T} + \mu_s$ is the Lebesgue-Radon-Nikodym decomposition of μ with respect to $\mathcal{L}^\mathbb{T}$, i.e. $g \in L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$ and μ_s is a real Borel measure on \mathbb{T} singular to $\mathcal{L}^\mathbb{T}$, then $\operatorname{Re} \varphi^*(\lambda) = g(\lambda)$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ (see e.g. [Koo, p. 11]). In particular, $\operatorname{Re} \varphi^* \in L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$ and if φ_s is a holomorphic function with boundary measure μ_s , then $\operatorname{Re} \varphi_s^*(\lambda) = 0$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$.

In what follows, given a n -tuple $\mu = (\mu_1, \dots, \mu_n)$ of real Borel measures on \mathbb{T} , by its Lebesgue-Radon-Nikodym decomposition with respect to $\mathcal{L}^\mathbb{T}$ we mean a unique decomposition

$$\mu = g d\mathcal{L}^\mathbb{T} + \mu_s,$$

where $g = (g_1, \dots, g_n) : \mathbb{T} \rightarrow \mathbb{R}^n$ is Borel-measurable, $g_1, \dots, g_n \in L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$ and $\mu_s = (\mu_{s,1}, \dots, \mu_{s,n})$ is a n -tuple of real Borel measures on \mathbb{T} , each of which is singular to $\mathcal{L}^\mathbb{T}$. In other words, for every j ,

$$\mu_j = g_j d\mathcal{L}^\mathbb{T} + \mu_{s,j}$$

is the Lebesgue-Radon-Nikodym decomposition of μ_j with respect to $\mathcal{L}^\mathbb{T}$. We call the n -tuples $g d\mathcal{L}^\mathbb{T}$ and μ_s respectively the absolutely continuous part and the singular part of μ (omitting the phrase 'in its Lebesgue-Radon-Nikodym decomposition with respect to $\mathcal{L}^\mathbb{T}$ ', which is assumed by default). The following lemma is a useful variation on the Lebesgue-Radon-Nikodym decomposition of n -tuples of measures:

Lemma 2.1. *Let μ be a n -tuple of real Borel measures on \mathbb{T} . Then there exist a unique finite, positive Borel measure ν on \mathbb{T} singular to $\mathcal{L}^\mathbb{T}$, a unique (up to a set of ν measure zero) Borel-measurable map $\varrho : \mathbb{T} \rightarrow \partial\mathbb{B}_n$ and a unique (up to a set of $\mathcal{L}^\mathbb{T}$ measure zero) Borel-measurable map $g : \mathbb{T} \rightarrow \mathbb{R}^n$ with components in $L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$ such that*

$$(3) \quad \mu = g d\mathcal{L}^\mathbb{T} + \varrho d\nu.$$

In particular, $g d\mathcal{L}^{\mathbb{T}}$ and $\varrho d\nu$ are (respectively) the absolutely continuous and singular parts of μ in its Lebesgue-Radon-Nikodym decomposition with respect to $\mathcal{L}^{\mathbb{T}}$.

Lemma 2.1 follows directly from the following general fact, applied to the singular part of μ :

Lemma 2.2. *If (X, \mathcal{A}) is a measurable space and $\mu = (\mu_1, \dots, \mu_n)$ is a n -tuple of real measures $\mu_j : \mathcal{A} \rightarrow \mathbb{R}$, then there exists a unique finite, positive measure $\nu : \mathcal{A} \rightarrow [0, \infty)$ and a unique (up to a set of ν measure zero) \mathcal{A} -measurable map $\varrho : X \rightarrow \partial\mathbb{B}_n$ such that $\mu = \varrho d\nu$.*

Proof of Lemma 2.2. Define a finite, positive measure $\tilde{\nu}$ as

$$\tilde{\nu} := |\mu_1| + \dots + |\mu_n|.$$

Since each μ_j is absolutely continuous with respect to $\tilde{\nu}$, from the classical Radon-Nikodym theorem it follows that there exists an \mathcal{A} -measurable map $F = (F_1, \dots, F_n) : X \rightarrow \mathbb{R}^n$ such that $F_1, \dots, F_n \in L^1(X, \tilde{\nu})$ and

$$\mu_j = F_j d\tilde{\nu}, \quad j = 1, \dots, n.$$

We have $|\mu_j| = |F_j| d\tilde{\nu}$, so

$$|F_1(x)| + \dots + |F_n(x)| = 1 \text{ for } \tilde{\nu}\text{-a.e. } x \in X.$$

Let $\varrho : X \rightarrow \partial\mathbb{B}_n$ be an \mathcal{A} -measurable map such that $F(x) = \varrho(x)\|F(x)\|$ for $\tilde{\nu}$ -a.e. $x \in X$. Set $\nu := \|F(x)\| d\tilde{\nu}(x)$. We have

$$\mu = F d\tilde{\nu} = \varrho d\nu,$$

what gives a desired decomposition.

It remains to show uniqueness. Assume that there are ν', ϱ' satisfying the same conditions as ν, ϱ . We have $\varrho d\nu = \varrho' d\nu'$. Set $\omega := \nu + \nu'$ and let $G, G' : X \rightarrow [0, \infty)$ be \mathcal{A} -measurable functions, integrable with respect to ω and such that $\nu = G d\omega$ and $\nu' = G' d\omega$. We have

$$G\varrho d\omega = \varrho d\nu = \varrho' d\nu' = G'\varrho' d\omega.$$

Thus, the maps $G\varrho$ and $G'\varrho'$ are equal ω -a.e. on X . This gives

$$G(x) = \|G(x)\varrho(x)\| = \|G'(x)\varrho'(x)\| = G'(x) \text{ for } \omega\text{-a.e. } x \in X.$$

In consequence, $\nu = \nu'$ and ν -almost everywhere on X there holds the equality $\varrho = \varrho'$, because $\varrho d\nu = \varrho' d\nu'$. \square

Example 2.3. In this example we are going to decompose as in Lemma 2.1 the following n -tuple of measures:

$$\mu = g d\mathcal{L}^{\mathbb{T}} + (\alpha_1 \delta_{\lambda_1}, \dots, \alpha_n \delta_{\lambda_n}),$$

where $g = (g_1, \dots, g_n)$, $g_1, \dots, g_n \in L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$, $\alpha_1, \dots, \alpha_n \in \mathbb{R}$ and $\lambda_1, \dots, \lambda_n \in \mathbb{T}$. As the measure ν is required to be singular with respect to $\mathcal{L}^{\mathbb{T}}$, the first part of desired decomposition is equal to $g d\mathcal{L}^{\mathbb{T}}$ and the second part comes from

Lemma 2.2 applied to the measure $(\alpha_1\delta_{\lambda_1}, \dots, \alpha_n\delta_{\lambda_n})$. To find the latter part, we follow the proof of Lemma 2.2 with $X = \mathbb{T}$ and \mathcal{A} being the σ -field of Borel subsets of \mathbb{T} .

For $j \in \{1, \dots, n\}$ let

$$A_j := \{l \in \{1, \dots, n\} : \lambda_l = \lambda_j\}.$$

We have

$$\tilde{\nu} = |\alpha_1|\delta_{\lambda_1} + \dots + |\alpha_n|\delta_{\lambda_n},$$

so we may set

$$F := \left(\frac{\alpha_1}{\sum_{l \in A_1} |\alpha_l|} \chi_{\{\lambda_1\}}, \dots, \frac{\alpha_n}{\sum_{l \in A_n} |\alpha_l|} \chi_{\{\lambda_n\}} \right)$$

(the mapping F is $\tilde{\nu}$ -almost everywhere well defined, because if $\sum_{l \in A_j} |\alpha_l| = 0$ for some j , then $\chi_{\{\lambda_j\}}$ is $\tilde{\nu}$ -a.e. equal to 0 and so is the j -th component of the right hand side of the above definition). Since $\nu = \|F(\lambda)\| d\tilde{\nu}(\lambda)$, the measure ν is supported on the set $\{\lambda_1, \dots, \lambda_n\}$ and

$$\chi_{\{\lambda_j\}} d\nu = \sqrt{\sum_{l \in A_j} \alpha_l^2} \delta_{\lambda_j}.$$

This gives

$$(4) \quad \nu = \sum_{j=1}^n \frac{\sqrt{\sum_{l \in A_j} \alpha_l^2}}{\#A_j} \delta_{\lambda_j},$$

where $\#A_j$ denotes the number of elements of the set A_j .

A map $\varrho : \mathbb{T} \rightarrow \partial\mathbb{B}_n$ has to be taken such that the equality $F(\lambda) = \varrho(\lambda)\|F(\lambda)\|$ holds for $\tilde{\nu}$ -a.e. $\lambda \in \mathbb{T}$, or equivalently, for ν -a.e. $\lambda \in \mathbb{T}$. It means that

$$(5) \quad \varrho = \left(\frac{\alpha_1}{\sqrt{\sum_{l \in A_1} \alpha_l^2}} \chi_{\{\lambda_1\}}, \dots, \frac{\alpha_n}{\sqrt{\sum_{l \in A_n} \alpha_l^2}} \chi_{\{\lambda_n\}} \right) \quad \nu\text{-a.e. on } \mathbb{T}.$$

Note that the right hand side is ν -almost everywhere well defined and it does not matter what values ϱ takes outside the set $\{\lambda_1, \dots, \lambda_n\}$. The desired decomposition consists of the map g , the measure ν given by (4) and a map ϱ satisfying (5).

The situation becomes simpler in the case when $\lambda_1, \dots, \lambda_n$ are pairwise disjoint. We then have

$$\nu = |\alpha_1|\delta_{\lambda_1} + \dots + |\alpha_n|\delta_{\lambda_n}$$

and

$$\varrho = \left(\frac{\alpha_1}{|\alpha_1|} \chi_{\{\lambda_1\}}, \dots, \frac{\alpha_n}{|\alpha_n|} \chi_{\{\lambda_n\}} \right)$$

$\tilde{\nu}$ -almost everywhere on \mathbb{T} (again, j -th component of ϱ may be anyhow if $\alpha_j = 0$).

For a convex tube domain $D \subset \mathbb{C}^n$ introduce the following sets, describing some geometric properties of its base. Define

$$\begin{aligned} W_D &:= \left\{ v \in \mathbb{R}^n : \sup_{x \in \operatorname{Re} D} \langle x, v \rangle < \infty \right\}, \\ S_D &:= \{ y \in \mathbb{R}^n : \forall v \in W_D : \langle y, v \rangle \leq 0 \} \end{aligned}$$

and for a vector $v \in \mathbb{R}^n$,

$$P_D(v) := \{ p \in \overline{\operatorname{Re} D} : \langle x - p, v \rangle < 0 \text{ for all } x \in \operatorname{Re} D \}.$$

It is clear that all these sets are convex, $P_D(v) \subset \partial \operatorname{Re} D$ and if $v \in S_D$, $w \in W_D$ and $t \geq 0$, then $tv \in S_D$ and $tw \in W_D$, i.e. the sets S_D and W_D are infinite cones. Next observation presents a number of their elementary geometric properties.

Observation 2.4. *Let $D \subset \mathbb{C}^n$ be a convex tube domain and let $v \in \mathbb{R}^n$. Then:*

- (i) *the sets $P_D(v)$ and S_D are closed,*
- (ii) *if $P_D(v) \neq \emptyset$, then $v \in W_D$,*
- (iii) *if $p, q \in P_D(v)$, then the vectors $p - q$ and v are orthogonal,*
- (iv) *if the domain $\operatorname{Re} D$ is strictly convex (in the geometric sense, i.e. it is convex and $\partial \operatorname{Re} D$ does not contain any non-trivial segments), then the set $P_D(v)$ contains at most one element,*
- (v) *$v \in S_D$ iff for all $a \in \operatorname{Re} D$ and $t \geq 0$ there holds $a + tv \in \operatorname{Re} D$,*
- (vi) *if $\operatorname{Re} D$ contains no complex affine lines, then $\operatorname{int} W_D \neq \emptyset$,*
- (vii) *if $\operatorname{Re} D$ is bounded, then $W_D = \mathbb{R}^n$ and $S_D = \{0\}$.*

Proof. (i). If $(p_m)_m \subset P_D(v)$ and $p_m \rightarrow p$, then $\langle x - p, v \rangle \leq 0$ for each $x \in \operatorname{Re} D$. As $P_D(v) \neq \emptyset$, we have $v \neq 0$, so the map $x \mapsto \langle x - p, v \rangle$ is open. It is non-positive on the open set $\operatorname{Re} D$, so it is in fact negative on $\operatorname{Re} D$.

(iii). If $p, q \in P_D(v)$, then $\frac{1}{2}(p + q) \in P_D(v)$. Since $p, q \in \overline{\operatorname{Re} D}$, we have $\langle p - \frac{1}{2}(p + q), v \rangle \leq 0$ and $\langle q - \frac{1}{2}(p + q), v \rangle \leq 0$, what gives $\langle p - q, v \rangle = 0$.

(vi). It follows e.g. from [Zaj, Observation 2.4].

(ii), (iv), (v), (vii). The proofs are immediate. \square

3 Description of complex geodesics in an arbitrary convex tube domain and its applications in special classes of domains

In this section we formulate the main result of this paper, Theorem 3.1. It gives a full description of all complex geodesics for D in terms of its geometric properties, i.e. the sets $P_D(v)$, W_D , S_D . In the latter part of this section we show how it can be applied to obtain formulas for boundary measures of complex geodesics in some special classes of convex tube domains. We also give some examples. The proof of Theorem 3.1 is presented in Section 4.

Theorem 3.1. *Let $D \subset \mathbb{C}^n$ be a convex tube domain containing no complex affine lines and let $\varphi \in \mathcal{M}^n$ be a holomorphic map with boundary measure μ . Consider the decomposition*

$$\mu = g d\mathcal{L}^{\mathbb{T}} + \varrho d\nu,$$

where $g = (g_1, \dots, g_n) : \mathbb{T} \rightarrow \mathbb{R}^n$ and $\varrho : \mathbb{T} \rightarrow \partial\mathbb{B}_n$ are Borel-measurable maps, $g_1, \dots, g_n \in L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$ and ν is a positive, finite, Borel measure on \mathbb{T} singular to $\mathcal{L}^{\mathbb{T}}$.

Then

$$\varphi(\mathbb{D}) \subset D \text{ and } \varphi \text{ is a complex geodesic for } D$$

iff there exists a map $h \in \mathcal{H}^n$, $h \not\equiv 0$, such that the following conditions hold:

- (i) $g(\lambda) \in P_D(\bar{\lambda}h(\lambda))$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$,
- (ii) $\langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle \geq 0$ for ν -a.e. $\lambda \in \mathbb{T}$,
- (iii) $\varrho(\lambda) \in S_D$ for ν -a.e. $\lambda \in \mathbb{T}$,
- (iv) $\operatorname{Re} \varphi(0) \in \operatorname{Re} D$.

Moreover, if $\varphi(\mathbb{D}) \subset D$, φ is a complex geodesic for D and $h \in \mathcal{H}^n$, $h \not\equiv 0$ is a map satisfying the conditions (i) - (iv), then there also hold:

- (v) $\varrho(\lambda) \in S_D \cap \{\bar{\lambda}h(\lambda)\}^\perp$ for ν -a.e. $\lambda \in \mathbb{T}$,
- (vi) $\nu(\{\lambda \in \mathbb{T} : \bar{\lambda}h(\lambda) \in \operatorname{int} W_D\}) = 0$.
- (vii) $\bar{\lambda}h(\lambda) \in \overline{W_D}$ for every $\lambda \in \mathbb{T}$.

Note that from (vi) and (vii) it follows that the measure ν is supported on the set $\{\lambda \in \mathbb{T} : \bar{\lambda}h(\lambda) \in \partial W_D\}$.

Remark 3.2. Theorem 3.1 gives quite separate conditions for both parts $g d\mathcal{L}^{\mathbb{T}}$ and $\varrho d\nu$ of the decomposition of μ , what makes it relatively not difficult to construct a measure which defines a complex geodesic for D . The part $g d\mathcal{L}^{\mathbb{T}}$ must satisfy (i), while the part $\varrho d\nu$ must fulfill (ii) and (iii). Everything is connected 'only' by the map h . To construct a measure μ which defines a complex geodesic for D it suffices to choose a map $h \in \mathcal{H}^n$, $h \not\equiv 0$ such that

$$(6) \quad P_D(\bar{\lambda}h(\lambda)) \neq \emptyset \text{ for } \mathcal{L}^{\mathbb{T}}\text{-a.e. } \lambda \in \mathbb{T}$$

and next:

- take a Borel map g with integrable components satisfying (i) (note that it may happen that it is impossible, even if (6) holds - see Example 3.7),
- take a measure ν singular to $\mathcal{L}^{\mathbb{T}}$ and satisfying (vi),
- take a Borel map $\varrho : \mathbb{T} \rightarrow \partial\mathbb{B}_n$ satisfying (v).

Then, if $\mu = g d\mathcal{L}^{\mathbb{T}} + \varrho d\nu$ and additionally $\frac{1}{2\pi}\mu(\mathbb{T}) \in \operatorname{Re} D$ (i.e. $\operatorname{Re} \varphi(0) \in \operatorname{Re} D$), then μ is a boundary measure of a complex geodesic for the domain D .

Before we proceed to applications of Theorem 3.1, let us make a remark on convex tubes with bounded base:

Remark 3.3. If $D \subset \mathbb{C}^n$ is a convex tube domain containing no complex affine lines and $\operatorname{Re} D$ is bounded, then $W_D = \mathbb{R}^n$ and $S_D = \{0\}$, so from the condition (iii) of Theorem 3.1 it follows that $\varrho(\lambda) = 0$ for ν -a.e. $\lambda \in \mathbb{T}$. Hence ν is a null measure, because the image of ϱ lies in $\partial\mathbb{B}_n$. Then also the condition (ii) is automatically fulfilled. Thus, a holomorphic map φ with boundary measure μ is a complex geodesic for D iff

$$\mu = g d\mathcal{L}^{\mathbb{T}}$$

for some g, h satisfying (i) and (iv).

In general case, i.e. for D with unbounded base, the absolutely continuous part of μ is determined by Theorem 3.1 (i) in the same way. But in that case we have to deal also with the singular part of μ , what is a more interesting issue.

3.1 Convex tube domains with $\overline{W_D} = [0, \infty)^n$

In this section we investigate the family \mathcal{D}_n of all convex tube domains $D \subset \mathbb{C}^n$ such that

$$\overline{W_D} = [0, \infty)^n.$$

A convex tube domain D belongs to \mathcal{D}_n iff $e_1, \dots, e_n \in \overline{W_D}$ and

$$\operatorname{Re} D + (-\infty, 0]^n = \operatorname{Re} D.$$

The base of such a domain D contains no real affine lines and there holds the equality

$$S_D = (-\infty, 0]^n.$$

In Corollary 3.4 we describe all complex geodesics for a domain $D \in \mathcal{D}_n$ and we apply it in Section 5 to describe extremal mappings in some classes of Reinhardt domains in \mathbb{C}^n .

Corollary 3.4. *Let $D \in \mathcal{D}_n$, $n \geq 2$, and let $\varphi \in \mathcal{M}^n$ be a holomorphic map with boundary measure μ . Consider the decomposition*

$$\mu = g d\mathcal{L}^{\mathbb{T}} + \varrho d\nu,$$

where $g = (g_1, \dots, g_n) : \mathbb{T} \rightarrow \mathbb{R}^n$ and $\varrho = (\varrho_1, \dots, \varrho_n) : \mathbb{T} \rightarrow \partial\mathbb{B}_n$ are Borel-measurable maps, $g_1, \dots, g_n \in L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$ and ν is a positive, finite, Borel measure on \mathbb{T} singular to $\mathcal{L}^{\mathbb{T}}$. Then

$\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D

iff there exists a map $h \in \mathcal{H}^n$, $h \not\equiv 0$ such that the following conditions hold:

- (i) $h \in \mathcal{H}_+^n$,
- (ii) $g(\lambda) \in P_D(\bar{\lambda}h(\lambda))$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$.
- (iii) $\varrho(\lambda) \in (-\infty, 0]^n$ for ν -a.e. $\lambda \in \mathbb{T}$,
- (iv) $\operatorname{Re} \varphi(0) \in \operatorname{Re} D$,

(v) if $j \in \{1, \dots, n\}$ is such that $h_j \not\equiv 0$, then

$$\varrho_j d\nu = \alpha_j \delta_{\lambda_j}$$

for some $\lambda_j \in \mathbb{T}$ and $\alpha_j \in (-\infty, 0]$ such that $\alpha_j h_j(\lambda_j) = 0$.

Note that the condition (iii) from the above Corollary means that the singular part of μ , i.e. the measure $\varrho d\nu$, is just a n -tuple of negative measures. Moreover, the condition (v) means that if $h_j \not\equiv 0$, then the j -th component of the singular part of μ is of the form $\alpha_j \delta_{\lambda_j}$ with some appropriate α_j and λ_j .

Proof of Corollary 3.4. Assume that $\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D . Let h be as in Theorem 3.1. The conditions (i) - (iv) follow directly from Theorem 3.1, so it remains to show the condition (v).

Set $\varrho = (\varrho_1, \dots, \varrho_n)$. The expression $\langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle$, which is by Theorem 3.1 (v) ν -almost everywhere equal to zero, is a sum of ν -almost everywhere non-positive terms $\bar{\lambda}h_1(\lambda)\varrho_1(\lambda), \dots, \bar{\lambda}h_n(\lambda)\varrho_n(\lambda)$. Therefore, all this terms are ν -a.e. equal to zero. If j is such that $h_j \not\equiv 0$, then the function $h_j \in \mathcal{H}_+^1$ has at most one root on \mathbb{T} (counting without multiplicities). Hence, up to a set of ν measure zero, $\varrho_j = \beta_j \chi_{\{\lambda_j\}}$ for some $\lambda_j \in \mathbb{T}$ and $\beta_j \in (-\infty, 0]$ such that $\beta_j h_j(\lambda_j) = 0$. This gives the condition (v) with $\alpha_j := \beta_j \nu(\{\lambda_j\})$.

On the other hand, it follows from Theorem 3.1 that if h is such that the conditions (i) - (v) are satisfied, then φ is a complex geodesic for D . Indeed, it is clear that the conditions (i), (iii) and (iv) from Theorem 3.1 hold, so it suffices to show that (ii) is also fulfilled. From the assumption (v) we conclude that if j is such that $h_j \not\equiv 0$, then

$$\bar{\lambda}h_j(\lambda)\varrho_j(\lambda) d\nu(\lambda) = \alpha_j \bar{\lambda}h_j(\lambda) d\delta_{\lambda_j}(\lambda) = \alpha_j \bar{\lambda}_j h_j(\lambda_j) d\delta_{\lambda_j}(\lambda) = 0.$$

This implies that $\langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle d\nu(\lambda)$ is a null measure, what involves the condition (ii). The proof is complete. \square

Remark 3.5. Under the assumptions of Corollary 3.4, if φ is a complex geodesic for D , h is as in the corollary and $h_1 \not\equiv 0, \dots, h_n \not\equiv 0$, then it follows from (v) that

$$\varrho d\nu = (\alpha_1 \delta_{\lambda_1}, \dots, \alpha_n \delta_{\lambda_n})$$

for some $\alpha_1, \dots, \alpha_n \in (-\infty, 0]$ and $\lambda_1, \dots, \lambda_n \in \mathbb{T}$ such that

$$\alpha_1 h_1(\lambda_1) = \dots = \alpha_n h_n(\lambda_n) = 0.$$

Thus, the singular part of μ takes then a very special form.

In the opposite situation, i.e. when the set $A := \{j \in \{1, \dots, n\} : h_j \equiv 0\}$ is non-empty, for every $j \in A$ the j -th component of the singular part of μ may be almost arbitrary. More precisely, if $\omega_1, \dots, \omega_n$ are finite, negative Borel measures on \mathbb{T} , singular to $\mathcal{L}^\mathbb{T}$ and such that $\omega_j = \varrho_j d\nu$ for every $j \in \{1, \dots, n\} \setminus A$, then a holomorphic map ψ with boundary measure $g d\mathcal{L}^\mathbb{T} + (\omega_1, \dots, \omega_n)$ is a complex geodesic for D , provided that $\operatorname{Re} \psi(0) \in \operatorname{Re} D$. This fact follows directly from Corollary 3.4, because ψ satisfies the conditions (i) - (v) with h .

Note that if the domain $\operatorname{Re} D$ is strictly convex (in the geometric sense), then there must hold $h_1 \not\equiv 0, \dots, h_n \not\equiv 0$. Indeed, in view of the condition (ii) from Corollary 3.4, $\mathcal{L}^\mathbb{T}$ -almost all sets $P_D(\bar{\lambda}h(\lambda))$ are non-empty. Our claim is a consequence of strict convexity of D and of the following geometric property

of domains from the family \mathcal{D}_n : if for a vector $v = (v_1, \dots, v_n) \in \mathbb{R}^n$ we have $P_D(v) \neq \emptyset$ and $v_j = 0$ for some j , then $\partial \operatorname{Re} D$ contains a half-line of the form $p + (-\infty, 0] e_j$ for any $p \in P_D(v)$.

Example 3.6. Consider the following domain from the family \mathcal{D}_2 ,

$$D := \{(x_1, x_2) \in \mathbb{R}^2 : (\max\{x_1 + 1, 0\})^2 + (\max\{x_2 + 1, 0\})^2 < 1\} + i\mathbb{R}^2.$$

The base of D is shown on Figure 1. We have




Figure 1: The base of D

$$P_D(v) = \begin{cases} \left\{ \frac{v}{\|v\|} - (1, 1) \right\}, & \text{if } v \in (0, \infty)^2, \\ (-\infty, -1] \times \{0\}, & \text{if } v \in \{0\} \times (0, \infty), \\ \{0\} \times (-\infty, -1], & \text{if } v \in (0, \infty) \times \{0\}, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Take a complex geodesic $\varphi = (\varphi_1, \varphi_2) : \mathbb{D} \rightarrow D$ with boundary measure $\mu = (\mu_1, \mu_2)$ and let $g = (g_1, g_2)$, $\nu, \varrho = (\varrho_1, \varrho_2)$ and $h = (h_1, h_2) \in \mathcal{H}_+^2$ be as in Corollary 3.4.

Assume that the functions h_1, h_2 are linearly independent. By Corollary 3.4 (ii), for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ we have

$$g(\lambda) \in P_D(\bar{\lambda}h(\lambda)) = \left\{ \frac{\bar{\lambda}h(\lambda)}{\|\bar{\lambda}h(\lambda)\|} - (1, 1) \right\}.$$

In view of Remark 3.5, we obtain

$$(7) \quad \mu = \left(\frac{\bar{\lambda}h(\lambda)}{\|\bar{\lambda}h(\lambda)\|} - (1, 1) \right) d\mathcal{L}^\mathbb{T}(\lambda) + (\alpha_1 \delta_{\lambda_1}, \alpha_2 \delta_{\lambda_2})$$

for some $\alpha_1, \alpha_2 \in (-\infty, 0]$ and $\lambda_1, \lambda_2 \in \mathbb{T}$ such that

$$(8) \quad \alpha_1 h_1(\lambda_1) = \alpha_2 h_2(\lambda_2) = 0.$$

On the other hand, from Corollary 3.4 it follows that if a holomorphic map has boundary measure of the form (7) with some linearly independent $h_1, h_2 \in \mathcal{H}_+^2$

and some $\alpha_1, \alpha_2 \leq 0$, $\lambda_1, \lambda_2 \in \mathbb{T}$ satisfying (8), then it is a complex geodesic for D (the condition (iv) from Corollary 3.4 is then a consequence of linear independence of h_1, h_2).

Now consider the case when h_1, h_2 are linearly dependent, but $h_1, h_2 \not\equiv 0$. We have $h_2 = \gamma h_1$ for some $\gamma > 0$. Since $P_D(\bar{\lambda}h(\lambda)) = P_D((1, \gamma))$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$, the map g is $\mathcal{L}^\mathbb{T}$ -almost everywhere constant and equal to $\frac{(1, \gamma)}{\|(1, \gamma)\|} - (1, 1)$. Using Remark 3.5 again we conclude that the singular part of μ is of the form $(\alpha_1 \delta_{\lambda_1}, \alpha_2 \delta_{\lambda_2})$ for some $\alpha_1, \alpha_2 \in (-\infty, 0]$ and $\lambda_1, \lambda_2 \in \mathbb{T}$ such that $\alpha_1 h_1(\lambda_1) = \alpha_2 \gamma h_1(\lambda_2) = 0$. But from Corollary 3.4 (iv) it follows that $(\alpha_1, \alpha_2) \neq (0, 0)$, so h_1, h_2 have a common root $\lambda_0 \in \{\lambda_1, \lambda_2\}$. Thus,

$$(9) \quad \mu = \left(\frac{(1, \gamma)}{\|(1, \gamma)\|} - (1, 1) \right) d\mathcal{L}^\mathbb{T} + (\alpha_1, \alpha_2) \delta_{\lambda_0}.$$

On the other hand, if a holomorphic map has boundary measure of the form (9) with some $\gamma > 0$, $\lambda_0 \in \mathbb{T}$ and $\alpha_1, \alpha_2 \leq 0$ such that $\alpha_1 + \alpha_2 < 0$, then it is a complex geodesic for D . It is a consequence of Corollary 3.4 applied to that map and to $h(\lambda) = (\lambda - \lambda_0)(1 - \bar{\lambda}_0 \lambda) \cdot (1, \gamma)$.

It remains to consider the situation when $h_1 \equiv 0$ or $h_2 \equiv 0$. If $h_1 \equiv 0$, then $g(\lambda) \in (-\infty, -1] \times \{0\}$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$, in view of (ii). Moreover, from (v) it follows that $\varrho_2 d\nu = \alpha_2 \delta_{\lambda_2}$ for some $\alpha_2 \leq 0$ and $\lambda_2 \in \mathbb{T}$ such that $\alpha_2 h_2(\lambda_2) = 0$, and (iv) gives that $\alpha_2 \neq 0$. Therefore

$$(10) \quad \mu = (\mu_1, \alpha_2 \delta_{\lambda_2}).$$

On the other hand, one can check that if a holomorphic map has boundary measure given by (10) with some $\alpha_2 < 0$, $\lambda_2 \in \mathbb{T}$ and some real Borel measure μ_1 on \mathbb{T} such that $\mu_1 \leq -\mathcal{L}^\mathbb{T}$, then it is a complex geodesic for the domain D . If $h_2 \equiv 0$, then arguing similarly as before we obtain that

$$(11) \quad \mu = (\alpha_1 \delta_{\lambda_1}, \mu_2)$$

for some $\alpha_1 < 0$, $\lambda_1 \in \mathbb{T}$ and $\mu_2 \leq -\mathcal{L}^\mathbb{T}$. Any holomorphic map with boundary measure of this form is a complex geodesic for D .

Example 3.7. Consider the domain

$$D := \{(x_1, x_2) \in \mathbb{R}^2 : x_1, x_2 < 0, x_1 x_2 > 1\} + i\mathbb{R}^2.$$

It belongs to the family \mathcal{D}_2 . We have

$$(12) \quad P_D(v) = \left\{ \left(-\sqrt{\frac{v_2}{v_1}}, -\sqrt{\frac{v_1}{v_2}} \right) \right\}$$

when $v = (v_1, v_2) \in (0, \infty)^2$ and $P_D(v) = \emptyset$ otherwise.

Take a complex geodesic $\varphi : \mathbb{D} \rightarrow D$ with boundary measure μ and let g, ν, ϱ and $h = (h_1, h_2) \in \mathcal{H}_+^2$ be as in Corollary 3.4. Assume that h_1 and h_2 are linearly independent. Then $h_j(\lambda) = c_j(\lambda - \lambda_j)(1 - \bar{\lambda}_j \lambda)$ for some $c_1, c_2 > 0$, $\lambda_1, \lambda_2 \in \mathbb{D}$ such that $\lambda_1 \neq \lambda_2$. By Corollary 3.4 (ii), for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ we have

$$g(\lambda) = \left(-\sqrt{\frac{c_2}{c_1}} \frac{|\lambda - \lambda_2|}{|\lambda - \lambda_1|}, -\sqrt{\frac{c_1}{c_2}} \frac{|\lambda - \lambda_1|}{|\lambda - \lambda_2|} \right).$$

Since both components of g belong to $L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$, we have $\lambda_1, \lambda_2 \in \mathbb{D}$. Moreover, $\varrho d\nu = 0$, by Remark 3.5. In summary,

$$(13) \quad \mu = \left(c \frac{|\lambda - \lambda_2|}{|\lambda - \lambda_1|}, \frac{1}{c} \frac{|\lambda - \lambda_1|}{|\lambda - \lambda_2|} \right) d\mathcal{L}^\mathbb{T}(\lambda),$$

where $c = -(\frac{c_2}{c_1})^{\frac{1}{2}}$. Such a map φ extends analytically on a neighbourhood of $\overline{\mathbb{D}}$, because the map g is real analytic on \mathbb{T} . It follows from Corollary 3.4 that any holomorphic map with boundary measure of the form (13) (with some $c > 0$ and $\lambda_1, \lambda_2 \in \mathbb{D}$, $\lambda_1 \neq \lambda_2$) is a complex geodesic for D .

If the functions h_1, h_2 are linearly dependent, then similarly as in Example 3.6 we can show that μ is of the form

$$(14) \quad \mu = (-\gamma^{\frac{1}{2}}, -\gamma^{-\frac{1}{2}}) d\mathcal{L}^\mathbb{T} + (\alpha_1, \alpha_2) \delta_{\lambda_0}$$

for some $\gamma > 0$, $\lambda_0 \in \mathbb{T}$ and $\alpha_1, \alpha_2 \leq 0$ such that $\alpha_1 + \alpha_2 < 0$. And again, any holomorphic map with boundary measure of the above form is a complex geodesic for D .

We see that in this example every complex geodesic which admits a map h with linearly independent components can be extended analytically on a neighbourhood of the closed unit disc $\overline{\mathbb{D}}$. However, even in some 'similar' domains this property do not hold. For example, let

$$D' := \{(x_1, x_2) \in \mathbb{R}^2 : x_1, x_2 < 0, x_2 < -x_1^{-2}\} + i\mathbb{R}^2.$$

For $v = (v_1, v_2) \in (0, \infty)^2$ we have

$$P_{D'}(v) = \left\{ \left(-\left(\frac{2v_2}{v_1} \right)^{\frac{1}{3}}, -\left(\frac{v_1}{2v_2} \right)^{\frac{2}{3}} \right) \right\}.$$

Take $h(\lambda) := ((\lambda + 1)^2, \lambda)$ (it belongs to the family \mathcal{H}_+^2) and g such that $g(\lambda) \in P_{D'}(\bar{\lambda}h(\lambda))$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$, i.e.

$$g(\lambda) = \left(-2^{\frac{1}{3}} |\lambda + 1|^{-\frac{2}{3}}, -2^{-\frac{2}{3}} |\lambda + 1|^{\frac{4}{3}} \right).$$

We see that both components of g belong to $L^1(\mathbb{T})$. From Corollary 3.4 it follows that if $\alpha_1 \leq 0$, then the holomorphic map given by the boundary measure $\mu := g d\mathcal{L}^\mathbb{T} + (\alpha_1 \delta_{-1}, 0)$ is a complex geodesic for D' . But this map does not extend analytically to a neighbourhood of $\overline{\mathbb{D}}$.

In these examples we also see that it is possible that for some h there is no map g with components integrable with respect to $\mathcal{L}^\mathbb{T}$ and satisfying $g(\lambda) \in P_D(\bar{\lambda}h(\lambda))$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$, even if these sets are non-empty for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ (cf. Remark 3.2).

Remark 3.8. Although in examples presented above we considered only tube domains with base contained in $(-\infty, 0)^n$, it is clear that in the family \mathcal{D}_n there are domains with base not contained in any set of the form $a + (-\infty, 0)^n$, $a \in \mathbb{R}^n$. An example of such a domain in \mathbb{C}^2 is $\{(x_1, x_2) \in \mathbb{R}^2 : x_2 < -e^{x_1}\} + i\mathbb{R}^2$, where $W_D = \{(0, 0)\} \cup [0, \infty) \times (0, \infty) \subsetneq \overline{W}_D$. Applying Corollary 3.4 in the same way as previously, we can find formulas for boundary measures of all complex geodesics for such tube domains.

3.2 Domains in \mathbb{C}^2

Let $D \subset \mathbb{C}^2$ be a convex tube domain containing no complex affine lines. From Observation 2.4 it follows that the set $\overline{W_D}$ is a closed, convex, infinite cone with vertex at the origin and with non-empty interior. Thus, $\overline{W_D}$ is any of the whole \mathbb{R}^2 , a half-plane or a convex infinite angle, i.e. the set

$$\{(r \cos \theta, r \sin \theta) : r \geq 0, \theta \in [\theta_1, \theta_2]\}$$

for some $\theta_1 < \theta_2 < \theta_1 + \pi$.

If $\overline{W_D}$ is the whole \mathbb{R}^2 , then $\text{Re } D$ is bounded. Tubes with bounded base were considered in Remark 3.3. If $\overline{W_D}$ is an angle, then D is affinely equivalent to a convex tube domain $D' \subset \mathbb{C}^2$ having $\overline{W_{D'}} = [0, \infty)^2$. These domains are exactly those from the family \mathcal{D}_2 and they were considered in Subsection 3.1. If $\overline{W_D}$ is a half-plane, then we may assume that $\overline{W_D} = \mathbb{R} \times (-\infty, 0]$. This case we consider now, in Corollary 3.9. If $D \subset \mathbb{C}^2$ is a convex tube domain with $\overline{W_D} = \mathbb{R} \times (-\infty, 0]$, then D contains no complex affine lines, there holds the equality

$$S_D = \{0\} \times [0, \infty)$$

and D is of the form

$$D = \{(x_1, x_2) \in (a, b) \times \mathbb{R} : x_2 > f(x_1)\} + i\mathbb{R}^2$$

for some $-\infty \leq a < b \leq \infty$ and a convex function $f : (a, b) \rightarrow \mathbb{R}$ such that:

- if $a = -\infty$, then $f'_-(x), f'_+(x) \rightarrow -\infty$, when $x \rightarrow -\infty$, and
- if $b = \infty$, then $f'_-(x), f'_+(x) \rightarrow \infty$, when $x \rightarrow \infty$.

Here f'_- and f'_+ denotes the one-sided derivatives of f . Depending on a, b and f , the set $\overline{W_D} \setminus W_D$ may be any of the empty set, a horizontal half-line starting at the origin or the horizontal line $\mathbb{R} \times \{0\}$. In Corollary 3.9 all of this cases are treated the same, as there only the set $\overline{W_D}$ is important, not W_D itself.

Corollary 3.9. *Let $D \subset \mathbb{C}^2$ be a convex tube domain such that $\overline{W_D} = \mathbb{R} \times (-\infty, 0]$. Take a map $\varphi \in \mathcal{M}^2$ with boundary measure μ and consider the decomposition*

$$\mu = g d\mathcal{L}^{\mathbb{T}} + \varrho d\nu,$$

where $g = (g_1, g_2) : \mathbb{T} \rightarrow \mathbb{R}^2$ and $\varrho : \mathbb{T} \rightarrow \partial\mathbb{B}_2$ are Borel-measurable maps, $g_1, g_2 \in L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$ and ν is a positive, finite, Borel measure on \mathbb{T} singular to $\mathcal{L}^{\mathbb{T}}$. Then

$$\varphi(\mathbb{D}) \subset D \text{ and } \varphi \text{ is a complex geodesic for } D$$

iff there exists a map $h \in \mathcal{H}^2$, $h \not\equiv 0$ such that the following conditions hold:

- (i) $h_2 \in -\mathcal{H}_+^1$,
- (ii) $g(\lambda) \in P_D(\bar{\lambda}h(\lambda))$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$,
- (iii) $\varrho(\lambda) = e_2$ for ν -a.e. $\lambda \in \mathbb{T}$,
- (iv) $\text{Re } \varphi(0) \in \text{Re } D$,

(v) if $h_2 \not\equiv 0$, then $\nu = \alpha \delta_{\lambda_0}$ for some $\alpha \in [0, \infty)$ and $\lambda_0 \in \mathbb{T}$ such that $\alpha h_2(\lambda_0) = 0$.

The condition (iii) from the above Corollary means that $\varrho d\nu = (0, \nu)$. In particular, ν is equal to the singular part of the second component of μ .

Proof of Corollary 3.9. Assume that $\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D and let h be as in Theorem 3.1. The conditions (ii), (iii) and (iv) follow immediately from Theorem 3.1. Since $\bar{\lambda}h(\lambda) \in \overline{W_D}$ for every $\lambda \in \mathbb{T}$, we have $h_2 \in -\mathcal{H}_+^1$, what gives (i).

If $h_2 \not\equiv 0$, then h_2 has at most one root on \mathbb{T} (counting without multiplicities), so the set $\{\lambda \in \mathbb{T} : \bar{\lambda}h(\lambda) \in \partial W_D\}$ contains at most one element. Hence the condition (v) follows from Theorem 3.1 (vi).

It is a direct consequence of Theorem 3.1 that if h is such that the conditions (i) - (v) are fulfilled, then $\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D . \square

Example 3.10. Consider the domain

$$D := \{(x_1, x_2) \in \mathbb{R}^2 : x_1 > 0, x_2 > x_1^2\} + i\mathbb{R}^2.$$

This domain is of the type considered in Corollary 3.9, because

$$W_D = \{(r \cos \theta, r \sin \theta) : r \geq 0, \theta \in [-\pi, 0)\}.$$

For $v = (v_1, v_2) \in \mathbb{R}^2$ we have

$$P_D(v) = \begin{cases} \left\{ \left(-\frac{v_1}{2v_2}, \frac{v_1^2}{4v_2^2} \right) \right\}, & \text{if } v \in (0, \infty) \times (-\infty, 0), \\ \{(0, 0)\}, & \text{if } v \in (-\infty, 0] \times (-\infty, 0), \\ \{0\} \times [0, \infty), & \text{if } v \in (-\infty, 0) \times \{0\}, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Take a complex geodesic $\varphi : \mathbb{D} \rightarrow D$ with boundary measure $\mu = (\mu_1, \mu_2)$ and let $g = (g_1, g_2)$, ν , $\varrho = (\varrho_1, \varrho_2)$ and $h = (h_1, h_2) \in \mathcal{H}^1 \times (-\mathcal{H}_+^1)$ be as in Corollary 3.9. We have $\mu_1 = g_1 d\mathcal{L}^\mathbb{T}$, so from the conditions (ii) and (iv) of Corollary 3.9 it follows that the sets $\{\lambda \in \mathbb{T} : g_1(\lambda) > 0\}$, $\{\lambda \in \mathbb{T} : \bar{\lambda}h_1(\lambda) > 0\}$ are of positive $\mathcal{L}^\mathbb{T}$ measure and $h_2 \not\equiv 0$. In particular, $h_1 \in \mathcal{H}^1 \setminus (-\mathcal{H}_+^1)$. The condition (v) implies that

$$\varrho d\nu = (0, \alpha \delta_{\lambda_0})$$

for some $\alpha \in [0, \infty)$ and $\lambda_0 \in \mathbb{T}$ such that $\alpha h_2(\lambda_0) = 0$. Moreover, as $h_2 \not\equiv 0$, for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \{\zeta \in \mathbb{T} : \bar{\zeta}h_1(\zeta) \leq 0\}$ there holds $g(\lambda) = (0, 0)$. This gives

$$g(\lambda) = \left(-\frac{h_1(\lambda)}{2h_2(\lambda)}, \frac{h_1(\lambda)^2}{4h_2(\lambda)^2} \right) \chi_{\{\zeta \in \mathbb{T} : \bar{\zeta}h_1(\zeta) > 0\}}(\lambda)$$

for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$.

If the functions h_1, h_2 are linearly independent, then h_2 has no roots on the set $\{\zeta \in \mathbb{T} : \bar{\zeta}h_1(\zeta) \geq 0\}$, because $g_1, g_2 \in L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$. In that case

$$(15) \quad \mu = \left(-\frac{h_1(\lambda)}{2h_2(\lambda)}, \frac{h_1(\lambda)^2}{4h_2(\lambda)^2} \right) \chi_{\{\zeta \in \mathbb{T} : \bar{\zeta}h_1(\zeta) > 0\}}(\lambda) d\mathcal{L}^\mathbb{T}(\lambda) + (0, \alpha \delta_{\lambda_0}).$$

On the other hand, if a holomorphic map has boundary measure of the form (15) with some $h_1 \in \mathcal{H}^1 \setminus (-\mathcal{H}_+^1)$, $h_2 \in -\mathcal{H}_+^1$, $\alpha \in [0, \infty)$ and $\lambda_0 \in \mathbb{T}$ such that

h_1, h_2 are linearly independent, h_2 has no roots on $\{\zeta \in \mathbb{T} : \bar{\zeta}h_1(\zeta) \geq 0\}$ and $\alpha h_2(\lambda_0) = 0$, then it is a complex geodesic for D .

If h_1, h_2 are linearly dependent, then applying Corollary 3.9 (v) and arguing similarly as in previously considered examples we obtain that

$$\mu = (\gamma, \gamma^2) d\mathcal{L}^{\mathbb{T}} + (0, \alpha \delta_{\lambda_0})$$

for some $\alpha < 0$, $\lambda_0 \in \mathbb{T}$ and $\gamma > 0$. Any holomorphic map with boundary measure of the above form is a complex geodesic for D .

4 Proof of Theorem 3.1 and further remarks

The aim of this section is to prove the main result of the paper, Theorem 3.1. We begin with investigating the singular and absolutely continuous parts of the boundary measure of a complex geodesic in its Lebesgue-Radon-Nikodym decomposition with respect to $\mathcal{L}^{\mathbb{T}}$. Next we show the proof of Theorem 3.1 and we give some remarks related to it.

A starting point for our considerations is the following fact (see [Zaj, Theorem 1.2]):

Theorem 4.1. *Let $D \subset \mathbb{C}^n$ be a convex tube domain containing no complex affine lines and let $\varphi : \mathbb{D} \rightarrow D$ be a holomorphic map with boundary measure μ . Then φ is a complex geodesic for D iff there exists a map $h \in \mathcal{H}^n$, $h \not\equiv 0$, such that*

$$\langle \bar{\lambda}h(\lambda), \operatorname{Re} z d\mathcal{L}^{\mathbb{T}}(\lambda) - d\mu(\lambda) \rangle \leq 0$$

for every $z \in D$.

Lemma 4.2. *Let $D \subset \mathbb{C}^n$ be a convex tube domain containing no complex affine lines, $h \in \mathcal{H}^n$, $h \not\equiv 0$ and let $\varphi : \mathbb{D} \rightarrow D$ be a holomorphic map with boundary measure μ . Consider*

$$\mu = \operatorname{Re} \varphi^* d\mathcal{L}^{\mathbb{T}} + \mu_s,$$

the Lebesgue-Radon-Nikodym decomposition of μ with respect to $\mathcal{L}^{\mathbb{T}}$. Then

$$(16) \quad \langle \bar{\lambda}h(\lambda), \operatorname{Re} z d\mathcal{L}^{\mathbb{T}}(\lambda) - d\mu(\lambda) \rangle \leq 0 \text{ for each } z \in D$$

iff the following two conditions hold:

$$(i) \operatorname{Re} \varphi^*(\lambda) \in P_D(\bar{\lambda}h(\lambda)) \text{ for } \mathcal{L}^{\mathbb{T}}\text{-a.e. } \lambda \in \mathbb{T},$$

$$(ii) \langle \bar{\lambda}h(\lambda), d\mu_s(\lambda) \rangle \geq 0.$$

Proof. Let $\mu_s = (\mu_{s,1}, \dots, \mu_{s,n})$. There exists a Borel subset $S \subset \mathbb{T}$ such that

$$\mathcal{L}^{\mathbb{T}}(S) = 0, |\mu_{s,1}|(\mathbb{T} \setminus S) = \dots = |\mu_{s,n}|(\mathbb{T} \setminus S) = 0.$$

There hold the equalities

$$(17) \quad \chi_S d\mathcal{L}^{\mathbb{T}} = 0, \chi_{\mathbb{T} \setminus S} d\mathcal{L}^{\mathbb{T}} = \mathcal{L}^{\mathbb{T}}, \chi_S d\mu = \mu_s, \chi_{\mathbb{T} \setminus S} d\mu = \operatorname{Re} \varphi^* d\mathcal{L}^{\mathbb{T}}.$$

For $z \in D$ set

$$\nu_z := \langle \bar{\lambda}h(\lambda), \operatorname{Re} z d\mathcal{L}^{\mathbb{T}}(\lambda) - d\mu(\lambda) \rangle.$$

We have $\nu_z = \chi_{\mathbb{T} \setminus S} d\nu_z + \chi_S d\nu_z$ and from (17) it follows that

$$(18) \quad \chi_{\mathbb{T} \setminus S} d\nu_z = \langle \bar{\lambda} h(\lambda), \operatorname{Re} z - \operatorname{Re} \varphi^*(\lambda) \rangle d\mathcal{L}^{\mathbb{T}}(\lambda)$$

and

$$(19) \quad \chi_S d\nu_z = -\langle \bar{\lambda} h(\lambda), d\mu_s(\lambda) \rangle.$$

If the condition (16) holds, i.e. $\nu_z \leq 0$ for every $z \in D$, then (i) follows from [Zaj, Lemma 3.7] and (ii) follows from the equality (19). On the other hand, if there hold both (i) and (ii), then (18) and (19) gives that for each $z \in D$ the measures $\chi_{\mathbb{T} \setminus S} d\nu_z$ and $\chi_S d\nu_z$ are negative and hence ν_z is so. \square

Lemma 4.3. *Let $D \subset \mathbb{C}^n$ be a convex tube domain containing no complex affine lines, let $\varphi \in \mathcal{M}^n$ be a holomorphic map with boundary measure μ and let*

$$\mu = \operatorname{Re} \varphi^* d\mathcal{L}^{\mathbb{T}} + \mu_s$$

be the Lebesgue-Radon-Nikodym decomposition of μ with respect to $\mathcal{L}^{\mathbb{T}}$. Then $\varphi(\mathbb{D}) \subset \overline{D}$ iff the following two conditions hold:

- (i) $\operatorname{Re} \varphi^*(\lambda) \in \overline{\operatorname{Re} D}$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$,
- (ii) $\langle \mu_s, w \rangle \leq 0$ for every $w \in \overline{W_D}$.

Proof. Again, let $S \subset \mathbb{T}$ be such that there holds (17). Assume that $\varphi(\mathbb{D}) \subset \overline{D}$. The first condition is clear. If $v \in W_D$, then for some constant $C \in \mathbb{R}$ there is $\langle x, v \rangle < C$ for every $x \in \operatorname{Re} D$. In particular, $\langle \operatorname{Re} \varphi(\lambda), v \rangle < C$ for $\lambda \in \mathbb{D}$, what gives a similar inequality for measures:

$$\langle \operatorname{Re} \varphi(r\lambda) d\mathcal{L}^{\mathbb{T}}(\lambda), v \rangle \leq C d\mathcal{L}^{\mathbb{T}}, \quad r \in (0, 1).$$

Taking limit for r tending to 1 we get

$$\langle d\mu, v \rangle \leq C d\mathcal{L}^{\mathbb{T}}.$$

Hence

$$\langle \chi_S d\mu, v \rangle \leq C \chi_S d\mathcal{L}^{\mathbb{T}},$$

what together with (17) gives

$$\langle \mu_s, v \rangle \leq 0.$$

If $w \in \overline{W_D}$, then there exists a sequence $(v_m)_m \subset W_D$ tending to w . The measure $\langle \mu_s, w \rangle$ is a weak-* limit of the sequence $\langle \mu_s, v_m \rangle$ of negative measures, so it is also negative.

Now assume that both (i) and (ii) hold. It suffices to show that if $p \in \mathbb{R}^n \setminus \overline{\operatorname{Re} D}$ and $v \in \mathbb{R}^n$ are such that $\langle x - p, v \rangle \leq 0$ for every $x \in \operatorname{Re} D$, then $\langle \operatorname{Re} \varphi(\lambda) - p, v \rangle \leq 0$ for every $\lambda \in \mathbb{D}$. Fix p, v and λ . It is clear that $v \in W_D$ and $\langle \operatorname{Re} \varphi^*(\zeta) - p, v \rangle \leq 0$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\zeta \in \mathbb{T}$. We have

$$\begin{aligned} \langle \operatorname{Re} \varphi(\lambda) - p, v \rangle &= \frac{1}{2\pi} \int_{\mathbb{T}} \frac{1 - |\lambda|^2}{|\zeta - \lambda|^2} \langle \operatorname{Re} \varphi^*(\zeta) - p, v \rangle d\mathcal{L}^{\mathbb{T}}(\zeta) \\ &+ \frac{1}{2\pi} \int_{\mathbb{T}} \frac{1 - |\lambda|^2}{|\zeta - \lambda|^2} d(\langle \mu_s(\zeta), v \rangle), \end{aligned}$$

so $\langle \operatorname{Re} \varphi(\lambda) - p, v \rangle \leq 0$ and the proof is complete. \square

Proof of Theorem 3.1. We have

$$\operatorname{Re} \varphi^*(\lambda) = g(\lambda) \text{ for } \mathcal{L}^{\mathbb{T}}\text{-a.e. } \lambda \in \mathbb{T}$$

and

$$\mu_s = \varrho d\nu,$$

where $\mu_s = (\mu_{s,1}, \dots, \mu_{s,n})$ is the singular part of μ in its Lebesgue-Radon-Nikodym decomposition with respect to $\mathcal{L}^{\mathbb{T}}$. The condition (ii) from Lemma 4.2 may be written as

$$(20) \quad \langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle d\nu(\lambda) \geq 0$$

and the condition (ii) from Lemma 4.3 can be written as

$$(21) \quad \langle \varrho(\lambda), w \rangle d\nu(\lambda) \leq 0 \text{ for every } w \in \overline{W_D}.$$

Now it is clear that if for some map $h \in \mathcal{H}^n$, $h \not\equiv 0$ the conditions (i) - (iv) from Theorem 3.1 hold, then from Lemmas 4.2 and 4.3 it follows that $\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D . It remains to prove the opposite implication.

Assume that $\varphi(\mathbb{D}) \subset D$ and φ is a complex geodesic for D . Take $h \in \mathcal{H}^n$ as in Theorem 4.1. The condition (iv) is clear and the conditions (i), (ii) of Theorem 3.1 follow directly from (20) and Lemma 4.2.

Lemma 4.3 and the equality (21) imply that for every $w \in W_D$ and ν -a.e. $\lambda \in \mathbb{T}$ there holds

$$(22) \quad \langle \varrho(\lambda), w \rangle \leq 0.$$

This 'almost every' may a priori depend on w , but we can omit this problem in the following way. Take a dense, countable subset $\{w_j : j = 1, 2, \dots\} \subset W_D$ and for each j let $A_j \subset \mathbb{T}$ be a Borel set such that $\nu(\mathbb{T} \setminus A_j) = 0$ and $\langle \varrho(\lambda), w_j \rangle \leq 0$ for every $\lambda \in A_j$. Put $A := \bigcap_{j=1}^{\infty} A_j$. It is clear that $\nu(\mathbb{T} \setminus A) = 0$ and (22) holds every $w \in W_D$ and every $\lambda \in A$. Thus,

$$\varrho(\lambda) \in S_D \text{ for } \nu\text{-a.e. } \lambda \in \mathbb{T}.$$

This is exactly the condition (iii).

It remains to prove the last part of the theorem, i.e. if $h \in \mathcal{H}^n$, $h \not\equiv 0$ satisfy the conditions (i) - (iv), then it satisfy also (v), (vi) and (vii).

From (i) it follows that for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$ there is $\bar{\lambda}h(\lambda) \in W_D$. Hence, (vii) is a consequence of continuity of h .

We prove (v). Fix $\epsilon > 0$. There exists $\delta > 0$ such that $|\bar{\lambda}h_j(\lambda) - \bar{\zeta}h_j(\zeta)| \leq \epsilon$ for $j = 1, \dots, n$, whenever $\lambda, \zeta \in \mathbb{T}$ and $|\lambda - \zeta| \leq \delta$. Take $\lambda_1, \dots, \lambda_m \in \mathbb{T}$ for which the arcs $L_k := \{\lambda \in \mathbb{T} : |\lambda - \lambda_k| < \delta\}$, $k = 1, \dots, m$, cover the circle \mathbb{T} . For ν -a.e. $\lambda \in L_k$ we have

$$\langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle \leq \langle \bar{\lambda}_k h(\lambda_k), \varrho(\lambda) \rangle + \|\bar{\lambda}h(\lambda) - \bar{\lambda}_k h(\lambda_k)\| \leq \epsilon \sqrt{n}.$$

The last inequality follows from (iii) and (vii). As k and ϵ are arbitrary, the condition (v) follows.

Now we prove (vi). For every $\lambda \in \mathbb{T}$ such that $\varrho(\lambda) \in S_D$ and $\bar{\lambda}h(\lambda) \in \operatorname{int} W_D$ there holds $\langle \bar{\lambda}h(\lambda), \varrho(\lambda) \rangle < 0$, because the map $w \mapsto \langle \varrho(\lambda), w \rangle$ is open and non-positive on W_D , so it must be negative on $\operatorname{int} W_D$. Hence, in view of (v), $\bar{\lambda}h(\lambda) \in \operatorname{int} W_D$ holds ν -almost nowhere on \mathbb{T} . The proof is complete. \square

Remark 4.4. It follows from the proof, that if φ is a complex geodesic for D and h is as in Theorem 4.1, then all of the conditions from Theorem 3.1 are satisfied with this h . And vice versa, if the conditions (i) - (iv) from Theorem 3.1 hold for $h \in \mathcal{H}^n$, $h \not\equiv 0$, then h satisfy the condition from Theorem 4.1.

Example 4.5. In most of previously considered domains the singular part of boundary measure of a complex geodesic took a very special form (it was expressed by Dirac deltas), provided that the components of corresponding map h were linearly independent. In this example we will see that in some domains even for such h the singular part may be almost arbitrary. Consider tube domain in \mathbb{C}^2 with the base being a 'half-cone', namely

$$D := \left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : x_2 > 0, x_3 > \sqrt{x_1^2 + x_2^2} \right\} + i\mathbb{R}^3.$$

One can check that

$$W_D = \left\{ x_1 \in \mathbb{R}, x_2 \geq 0, x_3 \leq -\sqrt{x_1^2 + x_2^2} \right\} \cup \{x_1 \in \mathbb{R}, x_2 \leq 0, x_3 \leq -|x_1|\}$$

and

$$S_D = \overline{\text{Re } D}.$$

Let $h \in \mathcal{H}^2$ be such that

$$\bar{\lambda}h(\lambda) = (\text{Re } \lambda, \text{Im } \lambda, -1), \quad \lambda \in \mathbb{T},$$

i.e. $h(\lambda) = \frac{1}{2}(\lambda^2 + 1, -i\lambda^2 + i, -2\lambda)$. Set $g := 0$ and

$$\varrho(\lambda) := 2^{-\frac{1}{2}}(\text{Re } \lambda, \text{Im } \lambda, 1), \quad \lambda \in \mathbb{T}.$$

Note that for $\lambda \in \mathbb{T}$ there holds $\bar{\lambda}h(\lambda) \in \partial W_D$ iff $\text{Im } \lambda \geq 0$. Let ν be an arbitrary finite positive Borel measure on \mathbb{T} singular to $\mathcal{L}^\mathbb{T}$ and such that

$$\nu(\{\lambda \in \mathbb{T} : \text{Im } \lambda < 0\}) = 0.$$

Set $\mu := g d\mathcal{L}^\mathbb{T} + \varrho d\nu = \varrho d\nu$ and let φ be a holomorphic map given by the boundary measure μ . One can see that the conditions (i), (ii) and (iii) from Theorem 3.1 are fulfilled.

Now if we choose ν such that $\frac{1}{2\pi}\mu(\mathbb{T}) \in \text{Re } D$, then in view of Theorem 3.1 the map φ is a complex geodesic for D . To do so, we can e.g. take an arbitrary finite positive Borel measure ω singular to $\mathcal{L}^\mathbb{T}$ and supported on the set $\{\lambda \in \mathbb{T} : \text{Im } \lambda \geq 0\}$, and put $\nu := \omega + \delta_1 + \delta_i$.

Remark 4.6. Let $D \subset \mathbb{C}^n$ be a convex tube domain containing no complex affine lines. Then a map $\varphi \in \mathcal{O}(\mathbb{D}, D)$ is a complex geodesic for D iff there exists a number $m \in \{1, 2, 3\}$ and a real $m \times n$ matrix V with linearly independent rows such that the domain $D' := \{V \cdot z : z \in D\} \subset \mathbb{C}^m$ is a convex tube containing no complex affine lines and $V \cdot \varphi$ is a complex geodesic for D' . This claim follows from [Zaj, Lemma 4.3]: if φ is a complex geodesic for D and $h(\lambda) = \bar{a}\lambda^2 + b\lambda + a$ ($a \in \mathbb{C}^n$, $b \in \mathbb{R}^n$) is as in Theorem 3.1, then V may be chosen such that its rows form a basis of the space $X_h := \text{span}_\mathbb{R}\{\text{Re } a, \text{Im } a, b\}$. Moreover, if in this situation we do an affine change of coordinates so that $X_h = \mathbb{R}^m \times \{0\}^{n-m}$, then we conclude that the map $(\varphi_1, \dots, \varphi_m)$ has to be a complex geodesic for D' and the other components $\varphi_{m+1}, \dots, \varphi_n$ can be arbitrary, provided that $\varphi(\mathbb{D}) \subset D$.

5 Applications of Theorem 3.1 in Reinhardt domains in \mathbb{C}^n

In this section we apply results obtained in Subsection 3.1 to give formulas (more precisely, a necessary condition) for extremal mappings with respect to the Lempert function and the Kobayashi-Royden pseudometric in some classes of complete Reinhardt domains in \mathbb{C}^n . Recall that a non-empty open set $G \subset \mathbb{C}^n$ is called a *complete Reinhardt domain* if for every $(z_1, \dots, z_n) \in G$ and $\lambda_1, \dots, \lambda_n \in \mathbb{D}$ there holds $(\lambda_1 z_1, \dots, \lambda_n z_n) \in G$. To such a domain $G \subset \mathbb{C}^n$ we associate its *logarithmic image*

$$\log G := \{(\log |z_1|, \dots, \log |z_n|) \in \mathbb{R}^n : (z_1, \dots, z_n) \in G \cap (\mathbb{C}_*)^n\}$$

and the tube domain

$$D_G := \log G + i\mathbb{R}^n$$

(we have $\operatorname{Re} D_G = \log G$). The map

$$\exp : D_G \ni (z_1, \dots, z_n) \mapsto (e^{z_1}, \dots, e^{z_n}) \in G \cap (\mathbb{C}_*)^n$$

is then a holomorphic covering. If the domain G is bounded and pseudoconvex, then D_G belongs to the family \mathcal{D}_n . Using an argument from [Edi-Zwo] we obtain a relation between extremal mappings in G and complex geodesics in D_G . It allows us to apply our description of complex geodesics for domain $D_G \in \mathcal{D}_n$ (Corollary 3.4) to get formulas for extremal mappings in G .

Let $D \subset \mathbb{C}^n$ be a domain. The *Lempert function* $\ell_D : D \times D \rightarrow [0, \infty)$ for D is given by

$$\begin{aligned} \ell_D(z, w) &= \inf \{ \rho(\sigma_1, \sigma_2) : \sigma_1, \sigma_2 \in \mathbb{D}, \exists f \in \mathcal{O}(\mathbb{D}, D) : f(\sigma_1) = z, f(\sigma_2) = w \} \\ &= \inf \{ \rho(0, \sigma) : \sigma \in \mathbb{D}, \exists f \in \mathcal{O}(\mathbb{D}, D) : f(0) = z, f(\sigma) = w \} \end{aligned}$$

and the Kobayashi-Royden pseudometric $\kappa_D : D \times \mathbb{C}^n \rightarrow [0, \infty)$ for D is

$$\begin{aligned} \kappa_D(z, X) &= \inf \left\{ \frac{|\alpha|}{1-|\sigma|^2} : \alpha \in \mathbb{C}, \sigma \in \mathbb{D}, \exists f \in \mathcal{O}(\mathbb{D}, D) : f(\sigma) = z, \alpha f'(\sigma) = X \right\} \\ &= \inf \{ \alpha > 0 : \exists f \in \mathcal{O}(\mathbb{D}, D) : f(0) = z, \alpha f'(0) = X \}. \end{aligned}$$

We say that a holomorphic map $f : \mathbb{D} \rightarrow D$ is a ℓ_D -*extremal map* if $\rho(\sigma_1, \sigma_2) = \ell_D(f(\sigma_1), f(\sigma_2))$ for some $\sigma_1, \sigma_2 \in \mathbb{D}$ such that $\sigma_1 \neq \sigma_2$. We call f a κ_D -*extremal map* if $\kappa_D(f(\sigma), f'(\sigma)) = \frac{1}{1-|\sigma|^2}$ for some $\sigma \in \mathbb{D}$. We often use the following basic fact: given $\sigma_1, \sigma_2 \in \mathbb{D}$, $\sigma_1 \neq \sigma_2$ and $f \in \mathcal{O}(\mathbb{D}, D)$, the equality $\rho(\sigma_1, \sigma_2) = \ell_D(f(\sigma_1), f(\sigma_2))$ holds iff there is no map $g \in \mathcal{O}(\mathbb{D}, D)$ such that $g(\sigma_1) = f(\sigma_1)$, $g(\sigma_2) = f(\sigma_2)$ and $g(\mathbb{D}) \subset\subset D$. And analogously, given $\sigma \in \mathbb{D}$ and $f \in \mathcal{O}(\mathbb{D}, D)$, the equality $\kappa_D(f(\sigma), f'(\sigma)) = \frac{1}{1-|\sigma|^2}$ holds iff there is no map $g \in \mathcal{O}(\mathbb{D}, D)$ such that $g(\sigma) = f(\sigma)$, $g'(\sigma) = f'(\sigma)$ and $g(\mathbb{D}) \subset\subset D$.

For a convex tube domain $D \subset \mathbb{C}^n$ containing no complex affine lines let $\mathcal{G}(D)$ denote the family of all Borel-measurable maps $g = (g_1, \dots, g_n) : \mathbb{T} \rightarrow \mathbb{R}^n$ such that $g_1, \dots, g_n \in L^1(\mathbb{T}, \mathcal{L}^{\mathbb{T}})$ and with some $h \in \mathcal{H}^n$ there holds

$$g(\lambda) \in P_D(\bar{\lambda}h(\lambda)) \text{ for } \mathcal{L}^{\mathbb{T}}\text{-a.e. } \lambda \in \mathbb{T}.$$

It follows from Theorem 3.1 that if $g \in \mathcal{G}(D)$ and $\varphi \in \mathcal{M}^n$ is a map with boundary measure $g d\mathcal{L}^{\mathbb{T}}$, then either φ is a complex geodesic for D (if $\varphi(0) \in D$)

or its image lies in ∂D (in the opposite situation). Note that if $\varphi(\mathbb{D}) \subset \partial D$ and in addition the domain $\text{Re } D$ is strictly convex in the geometric sense, then φ is just a constant map.

In what follows, for a non-empty set $A = \{j_1, \dots, j_k\} \subset \{1, \dots, n\}$ with $j_1 < \dots < j_k$, by π_A we denote the projection $\mathbb{C}^n \rightarrow \mathbb{C}^k$ on the coordinates j_1, \dots, j_k .

In the following two propositions we present formulas for ℓ_G -extremal and κ_G -extremal maps in some classes of bounded, pseudoconvex, complete Reinhardt domains.

Proposition 5.1. *Let $G \subset \mathbb{C}^n$, $n \geq 2$, be a bounded, pseudoconvex, complete Reinhardt domain such that the domain $\log G$ is strictly convex in the geometric sense and let $f = (f_1, \dots, f_n) \in \mathcal{O}(\mathbb{D}, G)$ be a ℓ_G -extremal or a κ_G -extremal map. Set*

$$A := \{j \in \{1, \dots, n\} : f_j \not\equiv 0\}$$

and let k denote the number of elements of A .

Then $k > 0$ and there exist some functions $B_1, \dots, B_k \in \text{Aut}(\mathbb{D}) \cup \{1\}$ and a map $g \in \mathcal{G}(D_{\pi_A(G)})$ such that

$$\pi_A \circ f = (B_1 e^{\varphi_1}, \dots, B_k e^{\varphi_k}),$$

where $\varphi = (\varphi_1, \dots, \varphi_k) \in \mathcal{M}^k$ is a map with boundary measure $g d\mathcal{L}^{\mathbb{T}}$.

Proposition 5.2. *Let $G \subset \mathbb{C}^2$ be a bounded, pseudoconvex, complete Reinhardt domain and let $R_1, R_2 > 0$ be such that $\pi_{\{1\}}(G) = R_1 \cdot \mathbb{D}$ and $\pi_{\{2\}}(G) = R_2 \cdot \mathbb{D}$. If $f \in \mathcal{O}(\mathbb{D}, G)$ is a ℓ_G -extremal or a κ_G -extremal map, then there holds at least one of the following conditions:*

- (i) *there exists $j \in \{1, 2\}$ such that $\frac{1}{R_j} f_j \in \text{Aut}(\mathbb{D})$, or*
- (ii) *there exist some $B_1, B_2 \in \text{Aut}(\mathbb{D}) \cup \{1\}$ and $g \in \mathcal{G}(D_G)$ such that f is of the form*

$$f = (B_1 e^{\varphi_1}, B_2 e^{\varphi_2}),$$

where $\varphi = (\varphi_1, \varphi_2) \in \mathcal{M}^2$ is a map with boundary measure $g d\mathcal{L}^{\mathbb{T}}$.

Proof of Proposition 5.1. We present the proof only for the case when f is a ℓ_G -extremal map, because the proof for κ_G -extremal map is analogous. Let $\sigma_1, \sigma_2 \in \mathbb{D}$ be such that $\rho(\sigma_1, \sigma_2) = \ell_G(f(\sigma_1), f(\sigma_2))$ and $\sigma_1 \neq \sigma_2$. It is clear that $k > 0$. The domain $\pi_A(G)$ satisfies the same assumptions as G , namely it is a bounded, pseudoconvex, complete Reinhardt domain with $\text{Re } D_{\pi_A(G)}$ being strictly convex. Moreover, if $z = (z_1, \dots, z_n), w = (w_1, \dots, w_n) \in G$ are such that $z_j = w_j = 0$ for every $j \notin A$, then $\ell_G(z, w) = \ell_{\pi_A(G)}(\pi_A(z), \pi_A(w))$. In particular,

$$\ell_{\pi_A(G)}(\pi_A(f(\sigma_1)), \pi_A(f(\sigma_2))) = \ell_G(f(\sigma_1), f(\sigma_2)) = \rho(\sigma_1, \sigma_2),$$

what means that $\pi_A \circ f$ is a $\ell_{\pi_A(G)}$ -extremal map. Therefore, we need only to prove the conclusion for the domain $\pi_A(G)$ and the mapping $\pi_A \circ f$. The latter map has no components equal identically to zero, so in fact it is enough to prove the proposition under the additional assumption that $f_1, \dots, f_n \not\equiv 0$ and $A = \{1, \dots, n\}$.

Since f_j is bounded and $f_j \neq 0$ for every j , we may write (see [Koo, p. 76])

$$(23) \quad f = (B_1 e^{u_1 + \psi_1}, \dots, B_n e^{u_n + \psi_n})$$

for a (possibly infinite or identically equal to 1) Blaschke product B_j , a function $u_j \in \mathcal{M}^1$ with boundary measure of the form $\log |f_j^*| d\mathcal{L}^\mathbb{T}$ (note that the function $\log |f_j^*|$ belongs to $L^1(\mathbb{T}, \mathcal{L}^\mathbb{T})$) and a function $\psi_j \in \mathcal{M}^1$ with $\text{Im } \psi_j(0) = 0$ and with boundary measure being finite, negative and singular to $\mathcal{L}^\mathbb{T}$. Set $\varphi := (u_1 + \psi_1, \dots, u_n + \psi_n)$. For every j we have $\text{Re } \psi_j \leq 0$ on \mathbb{D} and, for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$,

$$\text{Re } \psi_j^*(\lambda) = 0, \text{Re } \varphi_j^*(\lambda) = \text{Re } u_j^*(\lambda), |B_j^*(\lambda)| = 1, |f_j^*(\lambda)| = e^{\text{Re } u_j^*(\lambda)}.$$

In particular, $\varphi(\mathbb{D}) \subset \overline{D_G}$.

We claim that either the map φ is a complex geodesic for D_G or the image of φ lies in ∂D_G . The idea of this claim comes from [Edi-Zwo]. Assume that $\varphi(\mathbb{D}) \not\subset \partial D_G$, then clearly $\varphi(\mathbb{D}) \subset D_G$. If φ is not a complex geodesic for D_G , then there exists a map $\tilde{\varphi} = (\tilde{\varphi}_1, \dots, \tilde{\varphi}_n) \in \mathcal{O}(\mathbb{D}, D_G)$ such that $\tilde{\varphi}(\sigma_1) = \varphi(\sigma_1)$, $\tilde{\varphi}(\sigma_2) = \varphi(\sigma_2)$ and $\tilde{\varphi}(\mathbb{D}) \subset \subset D_G$. Now the map $(B_1 e^{\tilde{\varphi}_1}, \dots, B_n e^{\tilde{\varphi}_n})$ maps σ_1, σ_2 to $f(\sigma_1), f(\sigma_2)$ and its image is relatively compact in G . It is a contradiction with the equality $\ell_G(f(\sigma_1), f(\sigma_2)) = \rho(\sigma_1, \sigma_2)$.

From the above claim we conclude that for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ there holds $\varphi^*(\lambda) \in \partial D_G$ and hence $f^*(\lambda) \in \partial G$. Note that from the above considerations it follows that the latter condition holds for every ℓ_G -extremal map - we will use this fact several times.

We are going to show that

$$(24) \quad B_1 e^{\psi_1}, \dots, B_n e^{\psi_n} \in \text{Aut}(\mathbb{D}) \cup \{1\}.$$

For this, suppose to the contrary that $B_j e^{\psi_j} \notin \text{Aut}(\mathbb{D}) \cup \{1\}$ for some j . We may assume that $j = 1$. Then $B_1 e^{\psi_1} \in \mathcal{O}(\mathbb{D}, \mathbb{D}) \setminus \text{Aut}(\mathbb{D})$, so there exists a function $\xi \in \mathcal{O}(\mathbb{D}, \mathbb{D})$ such that $\xi(\sigma_1) = B_1(\sigma_1) e^{\psi_1(\sigma_1)}$, $\xi(\sigma_2) = B_1(\sigma_2) e^{\psi_1(\sigma_2)}$ and $\xi(\mathbb{D}) \subset \subset \mathbb{D}$. The map

$$F := (\xi e^{u_1}, f_2, \dots, f_n) \in \mathcal{O}(\mathbb{D}, G)$$

maps σ_1, σ_2 to $f(\sigma_1), f(\sigma_2)$, so it is also a ℓ_G -extremal map. In particular, $F^*(\lambda) \in \partial G$ for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$. Observe that $0 < |\xi^*(\lambda) e^{u_1^*(\lambda)}| < |f_1^*(\lambda)|$. Hence, from the fact that $D_G \in \mathcal{D}_n$ and $\mathcal{L}^\mathbb{T}$ -almost every $f^*(\lambda)$ belongs to ∂G we conclude that $\partial \text{Re } D_G$ contains a non-trivial segment parallel to the vector e_1 . This contradicts strict convexity of $\text{Re } D_G$.

From (24) it follows that $B_j \in \text{Aut}(\mathbb{D}) \cup \{1\}$ and $\text{Re } \psi_j \equiv 0$ for every j . As $\text{Im } \psi_j(0) = 0$, we get $\psi_j \equiv 0$. Set $g_j := \log |f_j^*|$ and $g := (g_1, \dots, g_n)$. The boundary measure of φ is equal to $g d\mathcal{L}^\mathbb{T}$, so to complete the proof we need only to show that $g \in \mathcal{G}(D_G)$. If $\varphi(\mathbb{D}) \subset D_G$, then φ is a complex geodesic for $D_G \in \mathcal{D}_n$ and the conclusion follows directly from Corollary 3.4. In the opposite case, when $\varphi(\mathbb{D}) \subset \partial D_G$, the map φ is constant because of strict convexity of $\text{Re } D_G$. Thus, the map $g = \text{Re } \varphi^*$ is also constant (up to a set of $\mathcal{L}^\mathbb{T}$ measure zero) and its image lies in $\partial \text{Re } D_G$, so it belongs to $\mathcal{G}(D_G)$. \square

Proof of Proposition 5.2. We again consider only the case when $f = (f_1, f_2)$ is a ℓ_G -extremal map. Take $\sigma_1, \sigma_2 \in \mathbb{D}$ such that $\rho(\sigma_1, \sigma_2) = \ell_G(f(\sigma_1), f(\sigma_2))$ and

$\sigma_1 \neq \sigma_2$. If $f_1 \equiv 0$ or $f_2 \equiv 0$, then similarly as in the previous proof we can show that $f_2 = \pi_{\{2\}} \circ f$ is a $\ell_{R_2 \cdot \mathbb{D}}$ -extremal map or $f_1 = \pi_{\{1\}} \circ f$ is a $\ell_{R_1 \cdot \mathbb{D}}$ -extremal map. Then the condition (i) is satisfied. Thus, it remains to consider the situation when $f_1, f_2 \neq 0$. In that case there hold (23) with $B_1, B_2, u_1, u_2, \psi_1, \psi_2$ and φ as in the previous proof. Like there, either $\varphi(\mathbb{D}) \subset \partial D_G$ or φ is a complex geodesic for D_G , what allows us to conclude that $\varphi^*(\lambda) \in \partial D_G$ and $f^*(\lambda) \in \partial G$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$.

We claim that there hold any of the condition (i) from Proposition 5.2 or the condition (24) with $n = 2$, i.e.

$$(25) \quad B_1 e^{\psi_1}, B_2 e^{\psi_2} \in \text{Aut}(\mathbb{D}) \cup \{1\}$$

(cf. [Kli, Lemat 4.3.3]). Suppose that $B_1 e^{\psi_1} \notin \text{Aut}(\mathbb{D}) \cup \{1\}$. There exists a function $\xi \in \mathcal{O}(\mathbb{D}, \mathbb{D})$ such that $\xi(\sigma_1) = B_1(\sigma_1) e^{\psi_1(\sigma_1)}$, $\xi(\sigma_2) = B_1(\sigma_2) e^{\psi_1(\sigma_2)}$ and $\xi(\mathbb{D}) \subset \subset \mathbb{D}$. Consider the map

$$F := (F_1, f_2) := (\xi e^{u_1}, f_2).$$

Like previously, F is a ℓ_G -extremal map and $F^*(\lambda) \in \partial G$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$. Since for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$ we have $f^*(\lambda) \in \partial G$ and $|F_1^*(\lambda)| < |f_1^*(\lambda)|$, the fact that $D_G \in \mathcal{D}_2$ imply that

$$f_2^*(\lambda) \in \partial \pi_{\{2\}}(G) = R_2 \cdot \mathbb{T}.$$

Put $R := \sup_{\lambda \in \mathbb{D}} |f_1(\lambda)|$. As G is a complete Reinhardt domain, for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$ the bidisc $(|f_1^*(\lambda)| \cdot \mathbb{D}) \times (|f_2^*(\lambda)| \cdot \mathbb{D})$ lies in G . Therefore, the bidisc $(R \cdot \mathbb{D}) \times (R_2 \cdot \mathbb{D})$ also lies in G (here we rely on the assumption that $n = 2$). We have $F(\mathbb{D}) \subset (R \cdot \mathbb{D}) \times (R_2 \cdot \mathbb{D})$, so F is a $\ell_{(R \cdot \mathbb{D}) \times (R_2 \cdot \mathbb{D})}$ -extremal map. Hence either $\frac{1}{R_2} f_2 \in \text{Aut}(\mathbb{D})$ or $\frac{1}{R} F_1 \in \text{Aut}(\mathbb{D})$. But the image of the latter function is relatively compact in \mathbb{D} , so there must hold $\frac{1}{R_2} f_2 \in \text{Aut}(\mathbb{D})$. Applying the same reasoning we can show that if $B_2 e^{\psi_2} \notin \text{Aut}(\mathbb{D}) \cup \{1\}$, then $\frac{1}{R_1} f_1 \in \text{Aut}(\mathbb{D})$. It means that at least of the conditions (25) or (i) holds.

To complete the proof it suffices to prove that the condition (ii) follows from (25). For this, assume that (25) holds. As before, we need only to show that the map $g := (\log |f_1^*|, \log |f_2^*|) = \text{Re } \varphi^*$ belongs to $\mathcal{G}(D_G)$. If $\varphi(\mathbb{D}) \subset G_D$, then the conclusion follows from Corollary 3.4. In the opposite case, when $\varphi(\mathbb{D}) \subset \partial D_G$, take a vector $v \in \mathbb{R}^n$ such that $\langle x - \text{Re } \varphi(0), v \rangle < 0$ for every $x \in \text{Re } D_G$. From the maximum principle for harmonic functions it follows that $\langle \text{Re } \varphi - \text{Re } \varphi(0), v \rangle \equiv 0$. Defining $h(\lambda) := \lambda \cdot v$ we get $g(\lambda) = \text{Re } \varphi^*(\lambda) \in P_D(\bar{\lambda} h(\lambda))$ for $\mathcal{L}^{\mathbb{T}}$ -a.e. $\lambda \in \mathbb{T}$. The proof is complete. \square

Example 5.3. (cf. [Kli, Theorem 4.1.4]) Given numbers $p, q \in (0, \infty)$ and $a \in (0, 1)$, consider the domain

$$G := G_{a,p,q} := \{(z_1, z_2) \in \mathbb{C}^2 : |z_1|, |z_2| < 1, |z_1|^p |z_2|^q < a\}.$$

It is a bounded, pseudoconvex, complete Reinhardt domain in \mathbb{C}^2 with

$$D_G = \{(x_1, x_2) \in \mathbb{R}^2 : x_1, x_2 < 0, px_1 + qx_2 < \log a\} + i\mathbb{R}^2.$$

Take $g = (g_1, g_2) \in \mathcal{G}(D_G)$. Observe that up to a set of $\mathcal{L}^{\mathbb{T}}$ measure zero there hold any of $g_1 \equiv 0$, $g_2 \equiv 0$ or $pg_1 + qg_2 \equiv \log a$. Indeed, take $h = (h_1, h_2)$

as in the definition of the family $\mathcal{G}(D_G)$. If $h_1 \equiv 0$ or $h_2 \equiv 0$, then $g_2 \equiv 0$ or $g_1 \equiv 0$, respectively. In the opposite case, for $\mathcal{L}^\mathbb{T}$ -a.e. $\lambda \in \mathbb{T}$ we have

$$g(\lambda) \in P_{D_G}(\bar{\lambda}h(\lambda)) = \{(tp^{-1}\log a, (1-t)q^{-1}\log a) : t \in [0, 1]\},$$

so $\mathcal{L}^\mathbb{T}$ -almost everywhere on \mathbb{T} there holds $pg_1 + qg_2 = \log a$. From this observation it follows that if $\varphi = (\varphi_1, \varphi_2)$ is a holomorphic map with boundary measure $g d\mathcal{L}^\mathbb{T}$, then there holds any of $\operatorname{Re} \varphi_1 \equiv 0$, $\operatorname{Re} \varphi_2 \equiv 0$ or $p \operatorname{Re} \varphi_1 + q \operatorname{Re} \varphi_2 \equiv \log a$. In the case when $\operatorname{Re} \varphi_1 \not\equiv 0$ and $\operatorname{Re} \varphi_2 \not\equiv 0$ we therefore get

$$\varphi(\lambda) = (\psi(\lambda)p^{-1}\log a, (1-\psi(\lambda))q^{-1}\log a + i\beta), \quad \lambda \in \mathbb{D}$$

for a number $\beta \in \mathbb{R}$ and a holomorphic map $\psi : \mathbb{D} \rightarrow \cdot$, where

$$\cdot := \{\zeta \in \mathbb{C} : 0 < \operatorname{Re} \zeta < 1\}.$$

From these considerations and from Proposition 5.2 we conclude that if $f = (f_1, f_2) \in \mathcal{O}(\mathbb{D}, G_{a,p,q})$ is a $\ell_{G_{a,p,q}}$ -extremal or a $\kappa_{G_{a,p,q}}$ -extremal map, then one of following conditions hold:

- (i) $f_1 \in \operatorname{Aut}(\mathbb{D})$, or
- (ii) $f_2 \in \operatorname{Aut}(\mathbb{D})$, or
- (iii) f is of the form

$$f = (B_1 \exp(\psi p^{-1} \log a), B_2 \exp((1-\psi)q^{-1} \log a + i\beta))$$

for some $\psi \in \mathcal{O}(\mathbb{D}, \cdot)$, $\beta \in \mathbb{R}$ and $B_1, B_2 \in \operatorname{Aut}(\mathbb{D}) \cup \{1\}$ with $B_1 B_2 \neq 1$.

Acknowledgements. I would like to thank Łukasz Kosiński for bringing my attention to some important papers and for many comments that improved the final shape of the paper.

References

- [Bra-Sor] F. Bracci, A. Saracco, *Hyperbolicity in unbounded convex domains*, Forum Math. 21 (2009), no. 5, 815-825.
- [Edi-Zwo] A. Edigarian, W. Zwonek, *Schwarz lemma for the tetrablock*, Bull. Lond. Math. Soc. 41 (2009), no. 3, 506-514.
- [Jar-Pfl] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis*, Walter de Gruyter & Co., Berlin, 1993.
- [Kli] P. Kliś, *Odwzorowania ekstremalne* (PhD dissertation), 2012.
- [Koo] P. Koosis, *Introduction to H^p spaces* (2-nd edition), Cambridge University Press, Cambridge, 1998.
- [Lem] L. Lempert, *La métrique de Kobayashi et la représentation des domaines sur la boule*, Bull. Soc. Math. France 109 (1981), 427-474.
- [Zaj] S. Zając, *Complex geodesics in convex tube domains*, to appear in Ann. Scuola Norm. Sup. Pisa Cl. Sci.