

# A SHORT NOTE ON THE MULTIPLIER IDEALS OF MONOMIAL SPACE CURVES

HOWARD M THOMPSON

**ABSTRACT.** Thompson (2014) exhibits a formula for the multiplier ideal with multiplier  $\lambda$  of a monomial curve  $C$  with ideal  $I$  as an intersection of a term coming from the  $I$ -adic valuation, the multiplier ideal of the term ideal of  $I$ , and terms coming from certain specified auxiliary valuations. This short note shows it suffices to consider at most two auxiliary valuations. This improvement is achieved through a more intrinsic approach, reduction to the toric case.

## 1. INTRODUCTION

Let  $(Y, \Delta)$  be a pair, consisting of a normal variety  $Y$  over an algebraically closed field of characteristic zero and a  $\mathbb{Q}$ -divisor  $\Delta$  such that  $K_Y + \Delta$  is  $\mathbb{Q}$ -Cartier. Let  $\pi : X \rightarrow Y$  be a log resolution of the ideal sheaf  $\mathcal{I} \subseteq \mathcal{O}_Y$  that is also a log resolution of the pair  $(Y, \Delta)$ . That is,  $\pi$  is a proper birational morphism such that  $X$  is smooth, the union of the exceptional set of  $\pi$  and  $\pi^{-1}(\Delta)$  is a divisor with simple normal crossing support, and  $\mathcal{I} \cdot \mathcal{O}_X = \mathcal{O}_X(-F)$  is also a divisor with simple normal crossing support. In this setting, we define the multiplier ideal of  $\mathcal{I}^\lambda$  on the pair  $(Y, \Delta)$  to be

$$\mathfrak{J}((Y, \Delta), \mathcal{I}^\lambda) = \pi_* \mathcal{O}_X(K_X - \lfloor \pi^*(K_Y + \Delta) + \lambda F \rfloor).$$

This ideal sheaf on  $Y$  does not depend upon the choice of log resolution.

In recent years, researchers have begun to study which divisors on a log resolution contribute jumping numbers. See Alberich-Carramiñana, Álvarez Montaner and Dachs-Cadefau [1], Galindo and Monserrat [6], Hyry and Järvilehto [10], Naie [11], Naie [12], Smith and Thompson [14], and Tucker [18]. This paper refines the result of Thompson [17]

---

*Date:* March 1, 2022.

*2010 Mathematics Subject Classification.* Primary: 14F18; Secondary: 14H50, 14M25.

*Key words and phrases.* Multiplier ideals, toroidal varieties.

The author would like to thank Karen E. Smith and Rocío Blanco as well as too many others to name, for putting up with his near endless and largely fruitless discussion of this topic.

by finding a smaller set of divisors that contains all the divisors that contribute jumping numbers for a monomial space curve.

Section 2 of this paper recalls a strengthening of the notion of an embedded resolution of singularities known as a factorizing resolution and uses it to provide a proposition (Proposition 3 on the next page) about the structure of multiplier ideals.

Section 3 on the facing page of this paper recalls the Howald-Blickle Theorem (Proposition 4 on page 4) that provides a formula for the multiplier ideals of a monomial ideal on a normal affine toric variety, provides a reinterpretation (Proposition 6 on page 4) of that theorem, and provides a formula (Proposition 7 on page 4) for the multiplier ideals of a principal binomial ideal.

Section 4 on page 5 applies the ideas of the previous sections to refine the result of Thompson [17].

## 2. USING FACTORIZING RESOLUTIONS TO COMPUTE MULTIPLIER IDEALS

**Definition 1.** Let  $Z$  be a generically smooth subscheme of any variety  $Y$ . A *factorizing resolution* of  $Z$  is an embedded resolution  $\pi : X \rightarrow Y$  of  $Z$  such that

$$\mathcal{I}_Z \cdot \mathcal{O}_X = \mathcal{I}_{\tilde{Z}} \cdot \mathcal{L}$$

where  $\tilde{Z}$  is the strict transform of  $Z$ ,  $\mathcal{L}$  is an invertible sheaf, and the support of  $\mathcal{I}_Z \cdot \mathcal{O}_X$  is a simple normal crossings variety.

Recall that  $\pi$  is an embedded resolution of  $Z$  if it is proper birational morphism  $\pi : X \rightarrow Y$  such that:  $X$  is smooth and  $\pi$  is an isomorphism over the generic points of the components of  $Z$ , the exceptional locus  $\text{exc}(\pi)$  of  $\pi$  is a divisor with simple normal crossing support, and the strict transform  $\tilde{Z}$  is smooth and transverse to  $\text{exc}(\pi)$ . For an embedded resolution, we always have  $\mathcal{I}_Z \cdot \mathcal{O}_X = \mathcal{I}_{\tilde{Z}} \cap \mathcal{L}$  for some invertible sheaf  $\mathcal{L}$ . Here we require the intersection to be a product. Typically, this is achieved by blowing up embedded components of  $\mathcal{I}_Z \cdot \mathcal{O}_X$ . Here is a theorem on the existence of factorizing resolutions.

**Proposition 2.** (*Theorem 1.2 of Bravo [3], Section 3 of Eisenstein [5]*) *Let  $Z$  be a generically smooth subscheme of any variety  $Y$  over an algebraically closed field of characteristic zero such that there exists a birational morphism  $\mu_1 : Y' \rightarrow Y$  from a smooth variety  $Y'$  that is an isomorphism over the generic points of the components of  $Z$ . If  $D$  is a divisor on  $Y'$  with simple normal crossing support such that no component of the strict transform of  $Z$  is contained in  $D$ , then there exists a factorizing resolution  $\pi : X \rightarrow Y$  of  $Z$  that factors through  $\mu_1$ ,*

$\pi = \mu_2 \circ \mu_1$ , such that  $\tilde{Z} \cup \text{exc}(\pi) \cup \mu_2^{-1}(D)$  has simple normal crossing support.

Notice that if  $\pi : X \rightarrow Y$  is such a factoring resolution of  $Z$ , then the blowup of  $\tilde{Z}$  is a log resolution of  $Z$  and that the exceptional locus of this blowup consists of a collection of prime divisors in one-to-one correspondence with the components of  $Z$  with codimension at least two.

**Proposition 3.** *Let  $Z_1, \dots, Z_r$  be the components of  $Z$  and suppose  $e_i$  is the codimension of  $Z_i$  for all  $i$ . Fix a factorizing resolution  $\pi : X \rightarrow Y$  of  $Z$  that is also a log resolution of the pair  $(Y, \Delta)$  and let  $\mathfrak{b} = \pi_*(\mathcal{L})$  where  $\mathcal{I}_Z \cdot \mathcal{O}_X = \mathcal{I}_{\tilde{Z}} \cdot \mathcal{L}$  as above. Then,*

$$\mathfrak{J}((Y, \Delta), \mathcal{I}_Z^\lambda) = \mathfrak{J}((Y, \Delta), \mathfrak{b}^\lambda) \cap \bigcap_{i=1}^r \mathcal{I}_{Z_i}^{(\lfloor \lambda+1-e_i \rfloor)}$$

*Proof.* Since  $\mathcal{I}_Z \subseteq \mathfrak{b}$ , it is clear that  $\mathfrak{J}((Y, \Delta), \mathcal{I}_Z^\lambda) \subseteq \mathfrak{J}((Y, \Delta), \mathfrak{b}^\lambda)$ . Let us now show  $\mathfrak{J}((Y, \Delta), \mathcal{I}_Z^\lambda) \subseteq \mathcal{I}_{Z_i}^{(\lfloor \lambda+1-e_i \rfloor)}$  for each  $i$ . Since  $Z$  is generically smooth, the  $\mathcal{I}_{Z_i}$  are prime and the  $\mathcal{I}_{Z_i}^{(\lfloor \lambda+1-e_i \rfloor)}$  are primary. Since  $\mathcal{I}_{Z_i}^{(\lfloor \lambda+1-e_i \rfloor)}$  is primary, it suffices to check the inclusion generically along the corresponding component (that is, after localizing at  $\mathcal{I}_{Z_i}$ ). Because  $Z$  is generically reduced,  $\mathcal{I}_Z \cdot \mathcal{O}_{Z_i} = \mathcal{I}_{Z_i} \cdot \mathcal{O}_{Z_i}$ . It is simple calculation based on the fact that one can resolve  $\mathcal{O}_{Z_i}$  generically by blowing up  $\mathcal{O}_{Z_i}$ , and the relative canonical divisor for this blowup is  $(e_i - 1)E_i$ , where  $E_i$  is the resulting exceptional divisor.

On the other hand, the extension of the contraction of an ideal is contained in the ideal. So,  $\mathfrak{b} \cdot \mathcal{O}_X \subseteq \mathcal{L}$ . Thus,

$$\mathfrak{b} \cdot \mathcal{O}_X \cap \bigcap_{i=1}^r \mathcal{I}_{\tilde{Z}_i} \subseteq \mathcal{L} \cap \bigcap_{i=1}^r \mathcal{I}_{\tilde{Z}_i} = \mathcal{I}_Z \cdot \mathcal{O}_X.$$

Therefore, we see

$$\mathfrak{J}((Y, \Delta), \mathfrak{b}^\lambda) \cap \bigcap_{i=1}^r \mathcal{I}_{Z_i}^{(\lfloor \lambda+1-e_i \rfloor)} \subseteq \mathfrak{J}((Y, \Delta), \mathcal{I}_Z^\lambda)$$

using any log resolution of  $\mathfrak{b}$ ,  $Z$  and  $(Y, \Delta)$  that factors through  $\pi$ .  $\square$

### 3. EXPLOITING THE TORIC CASE

This paragraph is, essentially, a direct quote of Blickle [4]. Let  $(Y, \Delta)$  be a pair such that  $Y$  is a normal (affine) toric variety (say  $Y = \text{Spec } R$  for some normal semigroup ring  $R \subseteq \mathbb{k}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ ) and  $\Delta$  is a torus invariant  $\mathbb{Q}$ -divisor. Since  $K_Y + \Delta$  is  $\mathbb{Q}$ -Cartier and torus invariant,

there is a monomial  $\mathbf{x}^{\mathbf{u}}$  such that  $\operatorname{div} \mathbf{x}^{\mathbf{u}} = r(K_Y + \Delta)$  for some integer  $r$ . Set  $\mathbf{w} = \mathbf{u}/r$ . Blickle's version of Howald's [8] formula is the following.

**Proposition 4.** (*Theorem 1 of Blickle [4]*) *Let  $\mathfrak{a}$  be a monomial ideal on  $Y$ . Then, if  $\operatorname{Newt}(\mathfrak{a})$  is the Newton Polyhedron of  $\mathfrak{a}$ ,*

$$\mathfrak{J}((Y, \Delta), \mathfrak{a}^\lambda) = \langle \mathbf{x}^{\mathbf{v}} \in R \mid \mathbf{v} + \mathbf{w} \in \text{interior of } \lambda \operatorname{Newt}(\mathfrak{a}) \rangle$$

for all  $\lambda > 0$ .

This means that the multiplier ideals of a monomial ideal on a toric variety are contributed by (divisors supported on unions of) the Rees divisors of the ideal. (See Thompson [16] for a quick overview of the relationship between toric blowups and Newton polyhedra.) Other divisors that may appear on a log resolution do not contribute.

**Definition 5.** We will say a sheaf  $\mathcal{F}$  (respectively a Weil divisor  $D$ ) on  $X$  is *locally monomial* if  $X$  can be covered by open subschemes  $U$  such that each  $U$  is isomorphic to an open subscheme of a normal toric variety in such a way that  $\mathcal{F}(U)$  is identified with a torus invariant sheaf (resp. a torus invariant divisor).

**Proposition 6.** *Let  $(Y, \Delta)$  be a pair, consisting of a normal variety  $Y$  over an algebraically closed field of characteristic zero and a  $\mathbb{Q}$ -divisor  $\Delta$  such that  $K_Y + \Delta$  is  $\mathbb{Q}$ -Cartier. If  $\pi : X \rightarrow Y$  is a proper birational morphism such that  $\pi^{-1}(\Delta)$  and  $\mathcal{I} \cdot \mathcal{O}_X$  are locally monomial, then the multiplier ideals of  $\mathcal{I}$  are contributed by the Rees divisors of  $\mathcal{I} \cdot \mathcal{O}_X$ .*

*Proof.* Since the question is local on  $X$ , it suffices to consider the case where  $X$  is a normal affine toric variety and  $\mathfrak{a} = \mathcal{I} \cdot \mathcal{O}_X$  is a monomial ideal. Let  $\mu : X' \rightarrow X$  be a toric log resolution of the ideal  $\mathfrak{a}$  that is also a log resolution of the pair  $(X, \pi^{-1}(\Delta))$ . Evidently,  $\mu$  factors through the blowup of  $\mathfrak{a}$  and, as in the toric case, orders of vanishing on any exceptional divisor of  $\mu \circ \pi$  are determined by those on the blowup of  $\mathfrak{a}$ . This is just the fact that when one represents a polyhedron as an intersection of half-spaces it suffices to consider only the facet-defining half-spaces.  $\square$

Consider the case of any principal binomial ideal  $I = \langle \mathbf{x}^{\mathbf{v}_1} - \mathbf{x}^{\mathbf{v}_2} \rangle \subseteq \mathbb{k}[x_1, \dots, x_n]$ .

**Proposition 7.** *If  $I = \langle \mathbf{x}^{\mathbf{v}_1} - \mathbf{x}^{\mathbf{v}_2} \rangle$  and  $\mathfrak{t} = \langle \mathbf{x}^{\mathbf{v}_1}, \mathbf{x}^{\mathbf{v}_2} \rangle$  is the term ideal of  $I$ , then*

$$\mathfrak{J}(\mathbb{A}^n, I^\lambda) = \mathfrak{J}(\mathbb{A}^n, \mathfrak{t}^\lambda) \cap I^{(\lfloor \lambda \rfloor)}.$$

*Proof.* Let  $\mathbf{v}$  be a primitive lattice vector such that  $r\mathbf{v} = \mathbf{v}_1 - \mathbf{v}_2$  for some positive integer  $r$ . Cover the normalized blowup of  $\mathfrak{t}$  with affine

open toric varieties  $U_1, \dots, U_s$ . Now, consider the covering consisting of the open sets of the form  $\text{Spec } \mathbb{k}[x_1, \dots, x_n, \mathbf{x}^{\pm \mathbf{v}}] \setminus V(f)$  where  $f = \frac{\mathbf{x}^{r\mathbf{v}} - 1}{\mathbf{x}^{\mathbf{v}} - \zeta}$  for an  $r$ th root of unity  $\zeta$  and the open subsets obtained by removing the closure of  $V(\mathbf{x}^{r\mathbf{v}} - 1) \subseteq \text{Spec } \mathbb{k}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$  from each  $U_i$ .

Note that since  $\mathbf{v}$  is primitive,  $\mathbb{Z}\mathbf{v}$  splits from  $\mathbb{Z}^n$ . So,  $\mathbb{N}^n + \mathbb{Z}\mathbf{v} \cong S \times \mathbb{Z}$  where  $S$  is the image of  $\mathbb{N}^n$  in  $\mathbb{Z}^n / \mathbb{Z}\mathbf{v}$ . Each open set of the form  $\text{Spec } \mathbb{k}[x_1, \dots, x_n, \mathbf{x}^{\pm \mathbf{v}}] \setminus V(f)$  is of the form  $\text{Spec } \mathbb{k}[S][t]_{t+1}$  where  $t = \mathbf{x}^{\mathbf{v}} - 1$ .

And, on each open set of the form  $U_i \setminus V(\mathbf{x}^{r\mathbf{v}} - 1)$ ,  $I \cdot \mathcal{O}_X = \mathfrak{t} \cdot \mathcal{O}_X$  is monomial already. Thus, Proposition 6 on the preceding page applies. Moreover, it is clear that the components of the closure of  $V(\mathbf{x}^{r\mathbf{v}} - 1) \subseteq \text{Spec } \mathbb{k}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$  are smooth and meet the boundary transversely.

So, a toric desingularization of the blowup of  $\mathfrak{t} \cdot \mathcal{O}_X$  is a factorizing resolution of  $I$ . Now, apply Proposition 3 on page 3.  $\square$

This result is not new. Principal binomial ideals are nondegenerate. For an alternate proof, see Howald [9].

#### 4. APPLICATION TO THE MONOMIAL SPACE CURVE CASE

Using the previous ideas, one can refine the result of Thompson [17]. The case where the monomial space curve is contained in a smooth toric surface follows from the principal toric case by using adjunction and inversion of adjunction.

Let  $\mathbb{k}$  be a field of characteristic zero, let  $C = \{(t^{n_1}, t^{n_2}, t^{n_3})\} \subset \mathbb{A}_{\mathbb{k}}^3$  be a monomial space curve not contained in a smooth toric surface. Assume  $\mathbf{n} = [n_1 \ n_2 \ n_3] \in \mathbb{Z}_{>0}^3$  is a primitive vector, let  $\text{ord}_{\mathbf{n}}$  be the monomial valuation given by the standard pairing,  $\mathbf{x}^{\mathbf{m}} \mapsto \langle \mathbf{n}, \mathbf{m} \rangle$ , and let  $I \subset \mathbb{k}[x_1, x_2, x_3]$  be the ideal of  $C$ . We may assume there exist irreducible binomials  $f_1 = \mathbf{x}^{\mathbf{m}_1^+} - \mathbf{x}^{\mathbf{m}_1^-}$ ,  $f_2 = \mathbf{x}^{\mathbf{m}_2^+} - \mathbf{x}^{\mathbf{m}_2^-}$ , and  $f_3 = \mathbf{x}^{\mathbf{m}_3^+} - \mathbf{x}^{\mathbf{m}_3^-}$  such that  $\{f_1, f_2, f_3\}$  or  $\{f_1, f_2\}$  is a minimal generating set for  $I$ . Let  $\mathfrak{t} = (\mathbf{x}^{\mathbf{m}_1^+}, \mathbf{x}^{\mathbf{m}_1^-}, \mathbf{x}^{\mathbf{m}_2^+}, \mathbf{x}^{\mathbf{m}_2^-}, \mathbf{x}^{\mathbf{m}_3^+}, \mathbf{x}^{\mathbf{m}_3^-})$  be the term ideal of  $I$ . Let  $d_i = \text{ord}_{\mathbf{n}}(f_i)$  for  $i = 1, 2, 3$ . Order the generators so that  $d_1 < d_2 < d_3$  and order the  $n_i$  so that  $n_i | d_i$  for  $i = 1, 2$  (and  $n_3 | d_3$  when  $f_3$  is a minimal generator). See Section 3 of Shibuta and Takagi [13] for a more detailed setup. Let  $\mathbf{m}_1 = \mathbf{m}_1^+ - \mathbf{m}_1^-$  and let  $\mathbf{q} = [q_1 \ q_2 \ 0] \in \mathbb{N}^3$  be the primitive vector such that  $\langle \mathbf{q}, \mathbf{m}_1 \rangle = 0$ . And, let  $e_i = \text{ord}_{\mathbf{q}}(f_i)$  for  $i = 1, 2, 3$ .

**Proposition 8.** *Let  $\mathbf{a}_1 = (\mathbf{x}^{\mathbf{m}_1^+}, \mathbf{x}^{\mathbf{m}_1^-})$ , let  $\mathbf{a}_2 = (x_1^{n_2 n_3}, x_2^{n_1 n_3}, x_3^{n_1 n_2})$ , and let the toric variety  $X = X_{\Sigma}$  be the normalized blowup of  $\mathbf{a}_1 \mathbf{a}_2$ . Then the ideal sheaf  $I \cdot \mathcal{O}_X$  is locally monomial.*

*Proof.* The blowup of  $\mathbf{a}_1$  is the partial desingularization of the toric surface  $V(f_1)$  identified in González Pérez and Teissier [7], and the normalized blowup of  $\mathbf{a}_2$  is the partial desingularization of  $C$ . The fan  $\Sigma_1$  of the blowup of  $\mathbf{a}_1$  has two maximal cones  $\{\mathbf{v} \in \mathbb{R}_{\geq 0}^3 \mid \langle \mathbf{v}, \mathbf{m}_1 \rangle \leq 0\}$  and  $\{\mathbf{v} \in \mathbb{R}_{\geq 0}^3 \mid \langle \mathbf{v}, \mathbf{m}_1 \rangle \geq 0\}$ . The normalized blowup of  $\mathbf{a}_2$  is stellar subdivision along the ray  $\rho = \mathbb{R}_{\geq 0}\mathbf{n}$ . Note that  $\mathbf{n}$  is in the intersection of the two maximal cones of  $\Sigma_1$ . So, the two operations on fans, stellar subdivision along  $\mathbf{n}$  and cutting with the plane  $\{v \in \mathbb{R}_{\geq 0}^3 \mid \langle \mathbf{v}, \mathbf{m}_1 \rangle = 0\}$  commute. And,  $\Sigma$  is the stellar subdivision along  $\rho$  of  $\Sigma_1$ . (Any toric desingularization of  $X$  provides a common embedded desingularization of  $C$  and the surface  $V(f_1)$ .)

First, consider the affine open  $U_\rho$  of  $X$  and fix an element of the affine semigroup  $\mathbf{m}_\rho \in \mathbf{S}_\rho$  such that  $\langle \mathbf{m}_\rho, \mathbf{n} \rangle = 1$ . I claim,  $\{\mathbf{m}_1, \mathbf{m}_2\}$  is a basis of the kernel of the matrix  $\begin{bmatrix} n_1 & n_2 & n_3 \end{bmatrix}$ ,  $\mathbf{S}_\sigma = \mathbb{N}^3 + \mathbb{Z}\mathbf{m}_1 + \mathbb{Z}\mathbf{m}_2$ ,  $f_i = (\mathbf{x}^{\mathbf{m}_\rho})^{d_i}(\mathbf{x}^{\mathbf{m}_i} - 1)$  for each  $i = 1, 2, 3$ , and  $f_3 \in (f_1, f_2)\mathbb{k}[\mathbf{S}_\sigma]$ . So,

$$\begin{aligned} I \cdot \mathcal{O}_{U_\rho} &= ((\mathbf{x}^{\mathbf{m}_\rho})^{d_1}(\mathbf{x}^{\mathbf{m}_1} - 1), (\mathbf{x}^{\mathbf{m}_\rho})^{d_2}(\mathbf{x}^{\mathbf{m}_2} - 1)) \\ &= (\mathbf{x}^{d_1\mathbf{m}_\rho}) \cap (\mathbf{x}^{\mathbf{m}_1} - 1, \mathbf{x}^{d_2\mathbf{m}_\rho}) \cap (\mathbf{x}^{\mathbf{m}_1} - 1, \mathbf{x}^{\mathbf{m}_2} - 1) \end{aligned}$$

is monomial in  $\mathbf{x}^{\mathbf{m}_\rho}$ ,  $\mathbf{x}^{\mathbf{m}_1} - 1$ , and  $\mathbf{x}^{\mathbf{m}_2} - 1$ . Since  $d_1 < d_2$ , there is an embedded component supported on the intersection of the strict transform of the surface  $V(f_1)$  and the divisor  $D_\rho$ . Away from this embedded component,  $I \cdot \mathcal{O}_X = \mathfrak{t} \cdot \mathcal{O}_X$ . Thus, it suffices to check the closed points where the curve  $\overline{V(\mathbf{x}^{\mathbf{m}_1} - 1, \mathbf{x}^{\mathbf{m}_\rho})}$  meets  $X \setminus U_\rho$ .

Let  $p \in X$  be one of these two points, and let  $\sigma$  be the smallest cone of  $\Sigma$  such that  $p \in U_\sigma$ . Evidently,  $\rho \subsetneq \sigma$  since  $p \notin U_\rho$ . After possibly replacing  $\mathbf{m}_1$  with  $-\mathbf{m}_1$ , we may assume  $\mathbf{x}^{\mathbf{m}_1} - 1 \in \mathbf{m}_{X,p}$ . Since  $\mathbf{x}^{\mathbf{m}_1} - 1 \in \mathbf{m}_{X,p}$ ,  $p$  is not a torus-fixed point and  $\sigma$  is two-dimensional.

Let  $\sigma = \mathbb{R}_{\geq 0}^2 \begin{bmatrix} n_1 & n_2 & n_3 \\ r_1 & r_2 & r_3 \end{bmatrix}$ . Note that  $\mathbf{m}_1$  is a basis for the kernel of  $\begin{bmatrix} n_1 & n_2 & n_3 \\ r_1 & r_2 & r_3 \end{bmatrix}$ , and  $\langle \mathbf{n}, \mathbf{m}_i \rangle = 0$  for  $i = 1, 2, 3$ . After possibly replacing  $\mathbf{m}_2$  with  $-\mathbf{m}_2$  and  $\mathbf{m}_3$  with  $-\mathbf{m}_3$ , we may assume  $\mathbf{m}_2, \mathbf{m}_3 \in \mathbf{S}_\sigma = \mathbb{N}^3 + \mathbb{Z}\mathbf{m}_1$ . As in Proposition 7 on page 4, the affine semigroup is a product  $\mathbf{S}_\sigma \cong \mathbf{S} \times \mathbb{Z}\mathbf{m}_1$  where  $\mathbf{S}$  is the image of  $\mathbb{N}^3$  in the quotient  $\mathbb{Z}^3/\mathbb{Z}\mathbf{m}_1$ . Thus,  $\mathbb{k}[\mathbf{S}_\sigma] \cong \mathbb{k}[\mathbf{S}][t_3]_{t_3+1}$  where  $t_3 = \mathbf{x}^{\mathbf{m}_1} - 1$ .  $\square$

Recall  $\mathbf{q} = [q_1 \ q_2 \ 0] \in \mathbb{N}^3$  is the primitive vector such that  $\langle \mathbf{q}, \mathbf{m}_1 \rangle = 0$  and  $e_i = \text{ord}_{\mathbf{q}}(f_i)$  for  $i = 1, 2, 3$ . Here is the improvement to the main theorem of Thompson [17].

**Proposition 9.** (i) If  $I$  is a complete intersection or if  $e_2(d_3 - d_1) \leq e_1(d_3 - d_2)$ , then

$$\mathfrak{J}(I^\lambda) = I^{(\lfloor \lambda - 1 \rfloor)} \cap \mathfrak{J}(\mathfrak{t}^\lambda) \cap (f \mid \nu_1(f) \geq \lfloor a_1 \lambda - k_1 \rfloor)$$

where  $\nu_1$  is the valuation given by the generating sequence  $x_i \mapsto n_i$  for  $i = 1, 2, 3$ ,  $f_1 \mapsto d_2$ . Thus,  $a_1 = \nu_1(I) = d_2$  and  $k_1 = \nu_1(J_{R_{\nu_1}/\mathbb{k}[x]}) = n_1 + n_2 + n_3 + d_2 - d_1$  where  $J_{R_{\nu_1}/\mathbb{k}[x]}$  is the Jacobian of the discrete valuation ring  $R_{\nu_1}$  of  $\nu_1$  over  $\mathbb{k}[x]$ .

(ii) Otherwise,

$$\mathfrak{J}(I^\lambda) = I^{(\lfloor \lambda - 1 \rfloor)} \cap \mathfrak{J}(\mathfrak{t}^\lambda) \bigcap_{i=1,2} (f \mid \nu_i(f) \geq \lfloor a_i \lambda - k_i \rfloor)$$

where  $\nu_1$  is as before and  $\nu_2$  is given by the generating sequence  $x_1 \mapsto e_2 n_1 + (d_3 - d_2)q_1$ ,  $x_2 \mapsto e_2 n_2 + (d_3 - d_2)q_2$ ,  $x_3 \mapsto e_2 n_3$ ,  $f_1 \mapsto e_2 d_3$ . Thus,  $a_2 = \nu_2(I) = e_2 d_3$ , and  $k_2 = \nu_2(J_{R_{\nu_2}/\mathbb{k}[x]}) = e_2(n_1 + n_2 + n_3) + (d_3 - d_1)(q_1 + q_2) + e_2(d_3 - d_1) - e_1(d_3 - d_2)$  where  $J_{R_{\nu_2}/\mathbb{k}[x]}$  is the Jacobian of the discrete valuation ring  $R_{\nu_2}$  of  $\nu_2$  over  $\mathbb{k}[x]$ .

*Proof.* Apply Proposition 8 on page 5 and the convex geometry computation in the appendix.  $\square$

**Corollary 10.** (i) If  $I$  is a complete intersection or if  $e_2(d_3 - d_1) \leq e_1(d_3 - d_2)$ , then the log canonical threshold of  $I$  (at the origin) is

$$\text{lct}_0(I) = \min \left( \text{lct}_0(\mathfrak{t}), \frac{k_1 + 1}{a_1} \right).$$

(ii) Otherwise,

$$\text{lct}_0(I) = \min \left( \text{lct}_0(\mathfrak{t}), \frac{k_1 + 1}{a_1}, \frac{k_2 + 1}{a_2} \right).$$

Note that when  $e_2(d_3 - d_1) = e_1(d_3 - d_2)$ ,  $\nu_2$  is monomial in the  $\mathbf{x}$ -variables and both formulas apply. In Example 5.3 of Blanco and Encinas [2],  $e_2(d_3 - d_1) = e_1(d_3 - d_2)$ ,  $\nu_2$ . I do not know an example where  $e_2(d_3 - d_1) > e_1(d_3 - d_2)$ . A Macaulay2 package that implements this calculation, as presented in Thompson [17], is described in Teitler [15].

## APPENDIX

Recall that  $d_i = \text{ord}_{\mathbf{n}}(f_i)$  and let  $\mathbf{u}_i = \begin{bmatrix} d_i \\ e_i \end{bmatrix} = \begin{bmatrix} n_1 & n_2 & n_3 \\ r_1 & r_2 & r_3 \end{bmatrix} \mathbf{m}_i^-$  for  $i = 1, 2, 3$ . In the local monomial coordinates, we find the Rees valuations from the facets of the Newton polyhedron. It suffices to

consider the ideal  $(\mathbf{t}^{u_1}t_3, \mathbf{t}^{u_2}, \mathbf{t}^{u_3}) \subset \mathbb{k}[\mathbf{S}]ty_3]$  (see the conclusion of the proof of Proposition 9 on page 6). We know  $e_1 > e_2$  by examining Section 3 of Shibuta and Takagi [13]. And,  $r_2 = 0$  or  $r_3 = 0$ .

If  $r_2 = 0$ , then  $e_2 = 0$ ,  $\mathbf{t}^{u_3} \in (\mathbf{t}^{u_2})$ , and the facets of the Newton polyhedron  $\text{Newt}(\mathbf{t}^{u_1}t_3, \mathbf{t}^{u_2})$  are orthogonal to the rows of the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ e_1 & d_2 - d_1 & 0 \\ 1 & 0 & d_2 - d_1 \end{bmatrix}$$

This includes the complete intersection case. Note that the only two rows of our matrix that have a nonzero last entry are  $[0 \ 0 \ 1]$  and  $[1 \ 0 \ d_2 - d_1]$ . The other vectors correspond to valuations that are monomial in the original  $\mathbf{x}$ -variables. Our ideal  $(\mathbf{t}^{u_1}t_3, \mathbf{t}^{u_2}, \mathbf{t}^{u_3})$  has order zero on the valuation corresponding to  $[0 \ 0 \ 1]$ . And, the row  $[1 \ 0 \ d_2 - d_1]$  corresponds to  $\nu_1$ .

If  $\mathbf{t}^{u_2} \notin \overline{(\mathbf{t}^{u_1}, \mathbf{t}^{u_3})}$  and  $e_2 \neq 0$ , then the facets of the Newton polyhedron  $\text{Newt}(\mathbf{t}^{u_1}t_3, \mathbf{t}^{u_2}, \mathbf{t}^{u_3})$  are orthogonal to the rows of the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ e_1 - e_2 & d_2 - d_1 & 0 \\ 1 & 0 & d_2 - d_1 \\ e_2 & d_3 - d_2 & 0 \end{bmatrix}$$

and these rows all have nonnegative integer entries. In terms of the parameters introduced in Section 3 of Shibuta and Takagi [13],  $\alpha \leq \gamma$  in this case.

If  $\mathbf{t}^{u_2} \in \overline{(\mathbf{t}^{u_1}, \mathbf{t}^{u_3})}$  and  $e_2 \neq 0$ , then  $r_3 = 0$  and the facets of the Newton polyhedron  $\text{Newt}(\mathbf{t}^{u_1}t_3, \mathbf{t}^{u_2}, \mathbf{t}^{u_3})$  are orthogonal to the rows of the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ e_1 - e_3 & d_3 - d_1 & 0 \\ 1 & 0 & d_2 - d_1 \\ e_2 & d_3 - d_2 & e_2(d_3 - d_1) - e_1(d_3 - d_2) \end{bmatrix}$$

and these rows all have nonnegative integer entries. Note that the only three rows that have a nonzero last entry are  $[0 \ 0 \ 1]$ ,  $[1 \ 0 \ d_2 - d_1]$ ,



and

$$\begin{bmatrix} e_2 & d_3 - d_2 & e_2(d_3 - d_1) - e_1(d_3 - d_2) \end{bmatrix}$$

corresponding to the only bounded facet of  $\text{Newt}(\mathbf{t}^{\mathbf{u}_1}t_3, \mathbf{t}^{\mathbf{u}_2}, \mathbf{t}^{\mathbf{u}_3})$ . The other vectors correspond to valuations that are monomial in the original  $\mathbf{x}$ -variables. And, the bounded facet corresponds to  $\nu_2$ . For  $\nu_2$ , the orders of vanishing of the  $x$ -variables are given by the entries of

$$\begin{aligned} & \begin{bmatrix} e_2 & d_3 - d_2 & e_2(d_3 - d_1) - e_1(d_3 - d_2) \end{bmatrix} \begin{bmatrix} n_1 & n_2 & n_3 \\ q_1 & q_2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} e_2n_1 + (d_3 - d_2)q_1 & e_2n_2 + (d_3 - d_2)q_2 & e_2n_3 \end{bmatrix} \end{aligned}$$

and  $\nu_2(f_i) = e_2d_3$  for all  $i = 1, 2, 3$

$$\begin{aligned} & \begin{bmatrix} e_2 & d_3 - d_2 & e_2(d_3 - d_1) - e_1(d_3 - d_2) \end{bmatrix} \begin{bmatrix} d_1 & d_2 & d_3 \\ e_1 & e_2 & 0 \\ 1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} e_2d_3 & e_2d_3 & e_2d_3 \end{bmatrix} \end{aligned}$$

## REFERENCES

- [1] M. Alberich-Carramiñana, J. Álvarez Montaner, and F. Dachs-Cadefau, *Multiplier ideals in two-dimensional local rings with rational singularities* (2014), 32 pp., arXiv:1412.3605v1 [math.AG].  $\uparrow 1$
- [2] R. Blanco and S. Encinas, *A procedure for computing the log canonical threshold of a binomial ideal* (2014), 28 pp., arXiv:1405.3942v2 [math.AG].  $\uparrow 7$
- [3] A. Bravo, *A remark on strong factorizing resolutions*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **107** (2013), no. 1, 53–60, DOI 10.1007/s13398-012-0080-8. MR3031261  $\uparrow 2$
- [4] M. Blickle, *Multiplier ideals and modules on toric varieties*, Math. Z. **248** (2004), no. 1, 113–121, DOI 10.1007/s00209-004-0655-y. MR2092724 (2006a:14082)  $\uparrow 3, 4$
- [5] E. Eisenstein, *Generalizations of the restriction theorem for multiplier ideals* (2010), 17 pp., arXiv:1001.2841v1 [math.AG].  $\uparrow 2$
- [6] C. Galindo and F. Monserrat, *The Poincaré series of multiplier ideals of a simple complete ideal in a local ring of a smooth surface*, Adv. Math. **225** (2010), no. 2, 1046–1068, DOI 10.1016/j.aim.2010.03.008. MR2671187 (2012a:14039)  $\uparrow 1$
- [7] P. D. González Pérez and B. Teissier, *Embedded resolutions of non necessarily normal affine toric varieties*, C. R. Math. Acad. Sci. Paris **334** (2002), no. 5, 379–382, DOI 10.1016/S1631-073X(02)02273-2 (English, with English and French summaries). MR1892938 (2003b:14019)  $\uparrow 6$
- [8] J. A. Howald, *Multiplier ideals of monomial ideals*, Trans. Amer. Math. Soc. **353** (2001), no. 7, 2665–2671 (electronic), DOI 10.1090/S0002-9947-01-02720-9. MR1828466 (2002b:14061)  $\uparrow 4$

- [9] ———, *Multiplier Ideals of Sufficiently General Polynomials* (2003), 9 pp., arXiv:math/0303203v1 [math.AG]. ↑5
- [10] E. Hyry and T. Järvilehto, *Jumping numbers and ordered tree structures on the dual graph*, Manuscripta Math. **136** (2011), no. 3-4, 411–437, DOI 10.1007/s00229-011-0449-6. MR2844818 ↑1
- [11] D. Naie, *Jumping numbers of a unibranch curve on a smooth surface*, Manuscripta Math. **128** (2009), no. 1, 33–49, DOI 10.1007/s00229-008-0223-6. MR2470185 (2009j:14034) ↑1
- [12] ———, *Mixed multiplier ideals and the irregularity of abelian coverings of smooth projective surfaces*, Expo. Math. **31** (2013), no. 1, 40–72, DOI 10.1016/j.exmath.2012.08.005. MR3035120 ↑1
- [13] T. Shibuta and S. Takagi, *Log canonical thresholds of binomial ideals*, Manuscripta Math. **130** (2009), no. 1, 45–61, DOI 10.1007/s00229-009-0270-7. MR2533766 ↑5, 8
- [14] K. E. Smith and H. M. Thompson, *Irrelevant exceptional divisors for curves on a smooth surface*, Algebra, geometry and their interactions, Contemp. Math., vol. 448, Amer. Math. Soc., Providence, RI, 2007, pp. 245–254, DOI 10.1090/conm/448/08669, (to appear in print). MR2389246 (2009c:14004) ↑1
- [15] Z. Teitler, *Software for multiplier ideals* (2013), 7 pp., arXiv:1305.4435v1 [math.AG]. ↑7
- [16] H. M. Thompson, *Comments on toric varieties* (2003), 6 pp., arXiv:0310336 [math.AG]. ↑4
- [17] ———, *Multiplier ideals of monomial space curves*, Proc. Amer. Math. Soc. Ser. B **1** (2014), 33–41, DOI 10.1090/S2330-1511-2014-00001-8. MR3168880 ↑1, 2, 5, 6, 7
- [18] K. Tucker, *Jumping numbers on algebraic surfaces with rational singularities*, Trans. Amer. Math. Soc. **362** (2010), no. 6, 3223–3241, DOI 10.1090/S0002-9947-09-04956-3. MR2592954 (2011c:14106) ↑1

MATHEMATICS DEPARTMENT, 402 MURCHIE SCIENCE BUILDING, 303 EAST KEARSLEY STREET, FLINT, MI 48502-1950

*E-mail address:* hmthomps@umflint.edu

*URL:* <http://homepages.umflint.edu/~hmthomps/>