

Exactly conserved quasilocal operators for the XXZ spin chain

R. G. Pereira

Instituto de Física de São Carlos, Universidade de São Paulo, C.P. 369, São Carlos, SP, 13560-970, Brazil

V. Pasquier

Institut de Physique Théorique, DSM, CEA, URA2306 CNRS, Saclay, F-91191 Gif-sur-Yvette, France

J. Sirker

Department of Physics and Research Center OPTIMAS, Technical University Kaiserslautern, D-67663 Kaiserslautern, Germany
Department of Physics and Astronomy, University of Manitoba, Winnipeg, Manitoba, Canada R3T 2N2

I. Affleck

Department of Physics and Astronomy, University of British Columbia, Vancouver, B.C., Canada, V6T 1Z1

Abstract. We extend T. Prosen's construction of quasilocal conserved quantities for the XXZ model [Phys. Rev. Lett. **106**, 217206 (2011)] to the case of periodic boundary conditions. These quasilocal operators stem from a two-parameter transfer matrix which employs a highest-weight representation of the quantum group algebra inherent in the Yang-Baxter algebra. In contrast with the open chain, where the conservation law is weakly violated by boundary terms, the quasilocal operators in the periodic chain exactly commute with the Hamiltonian and other local conserved quantities.

1. Introduction

Although a precise definition of quantum integrability is yet to be formulated, the common view is that quantum integrable models are characterized by a macroscopic number of local conserved quantities [1]. The most familiar examples are Bethe ansatz solvable models in which a family of commuting operators can be derived by taking logarithmic derivatives of the transfer matrix with respect to a spectral parameter [2, 3]. Recently the physical consequences of nontrivial conservation laws have become relevant for nonequilibrium dynamics of cold atomic gases confined to one-dimensional geometries [4, 5]. It is generally believed that the long time average of local observables in integrable systems after a quantum quench is described by a generalized Gibbs ensemble (GGE) which incorporates conserved quantities besides the Hamiltonian [6, 7]. However, it is not clear what are all the conserved quantities that need to be included in the density matrix of the GGE. While the set of all projection operators onto eigenstates of the Hamiltonian — whose number increases exponentially with system size — is definitely more than necessary to describe the equilibration of local observables, the family of local conserved quantities derived from the transfer matrix — whose number scales only linearly with system size — may not be sufficient for this purpose [8, 9]. In fact, recent results indicate that the GGE that includes only local conserved quantities fails to describe the steady state after a quench in the XXZ model [10, 11, 12]. GGE expectation values for the XXZ model had been computed before in [13], where the small deviations from numerics were attributed to large relaxation times. New integrals of motion have also been constructed to explain why many-body localized systems (which are not integrable in the usual sense) do not thermalize [14, 15].

The question of additional conserved quantities beyond the local ones usually associated with integrability has become even more pertinent since the discovery of a conserved *quasilocal* operator for the XXZ model [16]. Using a matrix product ansatz, Prosen constructed a non-Hermitian operator that commutes with the Hamiltonian of an open XXZ chain up to boundary terms. The operator Z in Ref. [16] is not local in the usual sense because it cannot be written in the form $Q_n = \sum_{j=1}^N q_j^n$, where q_j^n is a local density acting on sites $j+1, \dots, j+n$ with n finite. Nevertheless, it is quasilocal in the sense that the operator norm defined at infinite temperature as $\langle Z^\dagger Z \rangle = 2^{-N} \text{Tr}\{Z^\dagger Z\}$ grows linearly with system size, as it does for any local operator. This quasilocal operator cannot be written as a linear combination of the local conserved quantities obtained from the transfer matrix because it has different symmetry properties. In particular, its imaginary part changes sign under spin inversion $\sigma_j^z \rightarrow -\sigma_j^z, \sigma_j^\pm \rightarrow \sigma_j^\mp$, whereas the local conserved quantities are all invariant under the same transformation. This symmetry is important because it implies that, unlike the local conserved quantities, the quasilocal operator has an overlap with the spin current operator and provides a nonzero Mazur bound [17, 18] for the spin Drude weight at high temperatures [19, 20, 21, 22, 23, 24, 25, 26]. This result establishes ballistic spin transport in the critical phase of the XXZ model at zero magnetic field, except at the

Heisenberg point, where the Mazur bound vanishes [16, 27, 28].

In Prosen's original construction [16], the conservation of the quasilocal operator followed from a set of cubic algebraic relations that the matrices in the ansatz had to satisfy. The conservation law in the open chain is broken by boundary terms, but it was argued that, due to Lieb-Robinson bounds, the boundary terms do not affect bulk correlators in the thermodynamic limit [29]. Initially the new conserved operator seemed unrelated to the integrability of the XXZ model. However, it was soon realized that the cubic algebraic relations can be reduced to the quadratic quantum group algebra $U_q[\text{SU}(2)]$ [30]. The latter arises naturally in the quantum inverse scattering method, where it is convenient to view the XXZ model as the integrable q -deformation of the Heisenberg model [31, 32, 33, 34]. More recently, Prosen and Ilievski [35] made an explicit connection with integrability using a highest-weight Yang-Baxter transfer operator to derive a continuous family of quasilocal operators $Z(\varphi)$ labeled by a complex parameter φ . This family contain the previously found operator as the particular choice $\varphi = \pi/2$.

In this work we provide an alternative derivation of quasilocal operators that works for the periodic chain. In this case the conservation law is not spoiled by boundary terms.‡ The paper is organized as follows. In section 2, we review the derivation of the local conserved quantities of the XXZ model within the standard approach of taking logarithmic derivatives of the transfer matrix. In section 3, we construct a family of two-parameter conserved quantities using an auxiliary transfer matrix with a highest-weight representation in the auxiliary space. The expansion of the conserved quantities about special values of the representation parameter, along with a discussion of the conditions that lead to quasilocality, is presented in section 4. Section 5 makes the point that quasilocal operators are obtained only at first order in the expansion about the special representation, as higher order operators are strictly nonlocal. Section 6 contains the calculation of the Mazur bound for the spin Drude weight using a single quasilocal conserved quantity. In section 7, we discuss the family of quasilocal operators obtained by varying the spectral parameter continuously. Finally, section 8 presents the conclusions.

2. Local conserved quantities

Our goal is to derive generating functions of operators that commute with the XXZ Hamiltonian [2]

$$H = \sum_{j=1}^N (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z), \quad (2.1)$$

where $\sigma^{x,y,z}$ denote the standard Pauli matrices. The Hamiltonian acts on the tensor product of vector spaces $V^{\otimes N} \equiv V_1 \otimes V_2 \otimes \dots \otimes V_N$, where $V_j = \mathbb{C}^2$ is the quantum space

‡ After this work had been submitted, a new paper by Prosen [36] appeared on the arXiv which also discusses the exact conservation of quasilocal operators for the periodic chain.

of the spin on site j .

In the general scheme of the quantum inverse scattering method, one starts by introducing an R matrix that depends on a complex spectral parameter z and satisfies the Yang-Baxter equation [2, 3]

$$R_{12}(zw^{-1})R_{1\mathcal{Q}}(z)R_{2\mathcal{Q}}(w) = R_{2\mathcal{Q}}(w)R_{1\mathcal{Q}}(z)R_{12}(zw^{-1}). \quad (2.2)$$

Here $R_{12}(z)$ acts nontrivially on $V_1 \otimes V_2$ and as the identity on a third, auxiliary space \mathcal{Q} with dimension $d_{\mathcal{Q}}$, to be specified below. For the XXZ model (or six-vertex model), we can write $R_{12}(z) = R(z) \otimes \mathbb{1}$ with the R matrix

$$R(z) = \begin{pmatrix} a & & & \\ & cz^{-1} & b & \\ & b & cz & \\ & & & a \end{pmatrix}, \quad (2.3)$$

with $a = zq - z^{-1}q^{-1}$, $b = z - z^{-1}$, $c = q - q^{-1}$. The parameter q is related to the anisotropy Δ in Eq. (2.1) by $\Delta = (q + q^{-1})/2$. We use the following notation for matrices that act on the tensor product of two spaces [3] :

$$R = R_{kl}^{ij} e_{ij}^{V_1} \otimes e_{kl}^{V_2}. \quad (2.4)$$

Here $e_{ij}^{V_1}$ are matrices acting on V_1 defined by $e_{ij}^{V_1} = \hat{e}_i \otimes \hat{e}_j$, where the set of vectors $\{\hat{e}_i\}$ forms an orthonormal basis of V_1 . In the example of the tensor product of two-dimensional spaces ($i, j = 1, 2$), the R matrix in Eq. (2.3) is written in the form

$$R(z) = \begin{pmatrix} R_{11}^{11} & R_{12}^{11} & R_{11}^{12} & R_{12}^{12} \\ R_{21}^{11} & R_{22}^{11} & R_{21}^{12} & R_{22}^{12} \\ R_{11}^{21} & R_{12}^{21} & R_{11}^{22} & R_{12}^{22} \\ R_{21}^{21} & R_{22}^{21} & R_{21}^{22} & R_{22}^{22} \end{pmatrix}. \quad (2.5)$$

The product of $e_{ij}^{V_1}$ matrices has the property $e_{ij}^{V_1} e_{kl}^{V_1} = \delta_{jk} e_{il}^{V_1}$. Using this property we can write down each element of the Yang-Baxter equation (2.2) corresponding to $e_{ij}^{V_1} \otimes e_{kl}^{V_2} \otimes e_{mn}^{\mathcal{Q}}$, with $i, j, k, l = 1, 2$ and $m, n = 1, \dots, d_{\mathcal{Q}}$:

$$\begin{aligned} & \sum_{a,b=1}^2 \sum_{c=1}^{d_{\mathcal{Q}}} [R_{12}(zw^{-1})]_{kb}^{ia} [R_{1\mathcal{Q}}(z)]_{mc}^{aj} [R_{2\mathcal{Q}}(w)]_{cn}^{bl} \\ &= \sum_{a,b=1}^2 \sum_{c=1}^{d_{\mathcal{Q}}} [R_{2\mathcal{Q}}(w)]_{mc}^{kb} [R_{1\mathcal{Q}}(z)]_{cn}^{ia} [R_{12}(zw^{-1})]_{bl}^{aj}. \end{aligned} \quad (2.6)$$

The Lax operator associated with a given site j can be introduced as an R matrix that acts on $V_j \otimes \mathcal{Q}$

$$L_j(z) = R_{j\mathcal{Q}}(z). \quad (2.7)$$

Then Eq. (2.2) implies the quadratic relation for Lax operators involving the R matrix in Eq. (2.3)

$$R_{12}(zw^{-1})L_1(z)L_2(w) = L_2(w)L_1(z)R_{12}(zw^{-1}). \quad (2.8)$$

It is convenient to express the solutions of Eq. (2.8) for an arbitrary auxiliary space in terms of operators K, S^+, S^- :

$$L_j(z) = \frac{1}{2} \left[\frac{(z - z^{-1})}{2} \mathbb{1}_j \otimes (K + K^{-1}) + \frac{(z + z^{-1})}{2} \sigma_j^z \otimes (K - K^{-1}) + (q - q^{-1})(z\sigma_j^+ \otimes S^- + z^{-1}\sigma_j^- \otimes S^+) \right]. \quad (2.9)$$

Written as a matrix in V_j (with entries that act on \mathcal{Q}), the Lax operator is

$$L_j(z) = \frac{1}{2} \begin{pmatrix} zK - z^{-1}K^{-1} & z(q - q^{-1})S^- \\ z^{-1}(q - q^{-1})S^+ & zK^{-1} - z^{-1}K \end{pmatrix}. \quad (2.10)$$

The Yang-Baxter equation (2.8) is then satisfied provided that the operators S^\pm, K acting on \mathcal{Q} obey the quantum group algebra $U_q[\text{SU}(2)]$ [34]

$$KS^+ = qS^+K, \quad (2.11)$$

$$KS^- = q^{-1}S^-K, \quad (2.12)$$

$$[S^+, S^-] = \frac{K^2 - K^{-2}}{q - q^{-1}}. \quad (2.13)$$

Choosing $\mathcal{Q} = \mathbb{C}^2$, we can use the spin-1/2 representation

$$K = q^{\tau^z/2}, \quad S^\pm = \tau^\pm, \quad (2.14)$$

where $\tau^{x,y,z}$ are Pauli matrices in the auxiliary space. The monodromy matrix for N spins (acting on $V_1 \otimes V_2 \otimes \dots \otimes V_N \otimes \mathcal{Q}$) is defined as

$$T_{\mathcal{Q}}(z) = L_N(z)L_{N-1}(z) \dots L_1(z). \quad (2.15)$$

The transfer matrix is

$$t_{\mathcal{Q}}(z) = \text{tr}_{\mathcal{Q}}\{T_{\mathcal{Q}}(z)\}, \quad (2.16)$$

where $\text{tr}_{\mathcal{Q}}$ denotes the trace over the auxiliary space \mathcal{Q} . It can be shown [2, 3] that the transfer matrix forms a one-parameter family of commuting operators in $V_1 \otimes V_2 \otimes \dots \otimes V_N$:

$$[t_{\mathcal{Q}}(z), t_{\mathcal{Q}}(w)] = 0, \quad \forall z, w \in \mathbb{C}. \quad (2.17)$$

The local conserved quantities Q_n are given by [2]

$$Q_{n+1} = \frac{d^n}{dz^n} \ln t_{\mathcal{Q}}(z) \Big|_{z=1}, \quad n \geq 1. \quad (2.18)$$

The operator Q_2 is proportional to the XXZ Hamiltonian in Eq. (2.1). The first nontrivial conserved quantity, Q_3 , coincides with the energy current operator [37]. In general, each Q_n can be written as a sum of operators that act on n neighbouring spins and is therefore local.

Consider the spin inversion transformation \mathcal{C} defined in the quantum space as $\mathcal{C}^{-1}\sigma_j^z\mathcal{C} = -\sigma_j^z$, $\mathcal{C}^{-1}\sigma_j^\pm\mathcal{C} = \sigma_j^\mp$, $\forall j$. The Lax operator in Eq. (2.9) transforms as

$$\begin{aligned} \tilde{L}_j(z) &= \mathcal{C}^{-1}L_j(z)\mathcal{C} \\ &= \frac{1}{2} \left[\frac{(z - z^{-1})}{2} \mathbb{1}_j \otimes (K + K^{-1}) - \frac{(z + z^{-1})}{2} \sigma_j^z \otimes (K - K^{-1}) + (q - q^{-1})(z\sigma_j^- \otimes S^- + z^{-1}\sigma_j^+ \otimes S^+) \right]. \end{aligned} \quad (2.19)$$

In the case of the spin-1/2 representation in Eq. (2.14), we can show that the transfer matrix is invariant under spin inversion using the following similarity transformation that acts on \mathcal{Q} :

$$W(z) = [W(z)]^{-1} = \begin{pmatrix} 0 & z \\ z^{-1} & 0 \end{pmatrix}. \quad (2.20)$$

This transformation is such that

$$[W(z)]^{-1} q^{\tau^z/2} W(z) = q^{-\tau^z/2}, \quad (2.21)$$

$$[W(z)]^{-1} \tau^+ W(z) = z^2 \tau^-, \quad (2.22)$$

$$[W(z)]^{-1} \tau^- W(z) = z^{-2} \tau^+. \quad (2.23)$$

It follows that

$$[W(z)]^{-1} \tilde{L}_j(z) W(z) = L_j(z). \quad (2.24)$$

As a result,

$$\begin{aligned} \tilde{t}_{\mathcal{Q}}(z) &= \mathcal{C}^{-1} t_{\mathcal{Q}}(z) \mathcal{C} \\ &= \text{tr}_{\mathcal{Q}} \{ \tilde{L}_N(z) \tilde{L}_{N-1}(z) \dots \tilde{L}_1(z) \} \\ &= \text{tr}_{\mathcal{Q}} \{ W^{-1} \tilde{L}_N W W^{-1} \tilde{L}_{N-1} W \dots W^{-1} \tilde{L}_1 W \} \\ &= \text{tr}_{\mathcal{Q}} \{ L_N(z) L_{N-1}(z) \dots L_1(z) \} \\ &= t_{\mathcal{Q}}(z). \end{aligned} \quad (2.25)$$

Since Eq. (2.25) is verified for all $z \neq 0, \infty$, we conclude that all the Q_n 's derived by expanding $t_{\mathcal{Q}}(z)$ about $z = 1$ are invariant under spin inversion. For instance, this is clearly the case for the XXZ Hamiltonian at zero magnetic field in Eq. (2.1).

3. Conserved quantities from two-parameter transfer matrix

The idea to obtain a generating function of conserved quantities which are *not* invariant under spin inversion is to introduce an auxiliary transfer matrix that commutes with $t_{\mathcal{Q}}(z)$ but employs a different representation of the quantum group algebra. Let us consider an auxiliary space \mathcal{A} with dimension $d_{\mathcal{A}}$. We denote the Lax operator defined in $V_j \otimes \mathcal{A}$ by

$$\mathcal{L}_j(z) = R_{j\mathcal{A}}(z). \quad (3.1)$$

By analogy with Eq. (2.15), we can define the corresponding monodromy matrix

$$T_{\mathcal{A}}(z) = \mathcal{L}_N(z) \mathcal{L}_{N-1}(z) \dots \mathcal{L}_1(z), \quad (3.2)$$

as well as the auxiliary transfer matrix

$$t_{\mathcal{A}}(z) = \text{tr}_{\mathcal{A}} \{ T_{\mathcal{A}}(z) \}. \quad (3.3)$$

We then apply the “train argument” [34] for the Yang-Baxter equation with an R matrix in $\mathcal{Q} \otimes \mathcal{A}$ as follows:

$$\begin{aligned}
T_{\mathcal{Q}}(z)T_{\mathcal{A}}(w)R_{\mathcal{Q}\mathcal{A}}(w/z) &= R_{N\mathcal{Q}}(z) \dots R_{1\mathcal{Q}}(z)R_{N\mathcal{A}}(w) \dots R_{1\mathcal{A}}(w)R_{\mathcal{Q}\mathcal{A}}(w/z) \\
&= R_{N\mathcal{Q}} \dots R_{2\mathcal{Q}}R_{N\mathcal{A}} \dots R_{2\mathcal{A}}R_{1\mathcal{Q}}R_{1\mathcal{A}}R_{\mathcal{Q}\mathcal{A}} \\
&= R_{N\mathcal{Q}} \dots R_{2\mathcal{Q}}R_{N\mathcal{A}} \dots R_{2\mathcal{A}}R_{\mathcal{Q}\mathcal{A}}R_{1\mathcal{A}}R_{1\mathcal{Q}} \\
&= R_{\mathcal{Q}\mathcal{A}}R_{N\mathcal{A}} \dots R_{1\mathcal{A}}R_{N\mathcal{Q}} \dots R_{1\mathcal{Q}} \\
&= R_{\mathcal{Q}\mathcal{A}}(w/z)T_{\mathcal{A}}(w)T_{\mathcal{Q}}(z).
\end{aligned} \tag{3.4}$$

Taking the trace of Eq. (3.4) over \mathcal{Q} and \mathcal{A} , we obtain

$$[\text{tr}_{\mathcal{Q}}\{T_{\mathcal{Q}}(z)\}, \text{tr}_{\mathcal{A}}\{T_{\mathcal{A}}(w)\}] = 0, \tag{3.5}$$

thus

$$[t_{\mathcal{Q}}(z), t_{\mathcal{A}}(w)] = 0 \quad \forall z, w \in \mathbb{C}. \tag{3.6}$$

Therefore, since the XXZ Hamiltonian is among the operators generated by $t_{\mathcal{Q}}(z)$, we can use $t_{\mathcal{A}}(z)$ as a generating function of conserved quantities.

We shall work with the highest weight representation of $U_q[\text{SU}(2)]$:

$$K|r\rangle = uq^r|r\rangle, \tag{3.7}$$

$$S^+|r\rangle = -a_r|r+1\rangle, \tag{3.8}$$

$$S^-|r\rangle = b_r|r-1\rangle, \tag{3.9}$$

where $u \in \mathbb{C}$ is arbitrary. The index r can be interpreted as positions in a lattice in the auxiliary space, and the operators S^+ and S^- perform hopping between nearest-neighbour sites. Eq. (2.13) imposes the relation

$$a_rb_{r+1} - a_{r-1}b_r = \frac{u^2q^{2r} - u^{-2}q^{-2r}}{q - q^{-1}}, \tag{3.10}$$

which is satisfied by the choice

$$a_r = v \frac{u^2q^r - u^{-2}q^{-r}}{q - q^{-1}}, \tag{3.11}$$

$$b_r = v^{-1} \frac{q^r - q^{-r}}{q - q^{-1}}, \tag{3.12}$$

where v is another arbitrary parameter which we set to 1 hereafter. In this representation the Casimir operator is a function of the parameter u :

$$C = (q - q^{-1})^2 S^+ S^- + q^{-1} K^2 + q K^{-2} = u^2 q^{-1} + u^{-2} q. \tag{3.13}$$

The dimension of the auxiliary space depends on the value of $\Delta = (q + q^{-1})/2$. When q is a root of unity, *i.e.* $q = e^{i\lambda}$ with $\lambda = l\pi/m$ and $l, m \in \mathbb{Z}$ coprimes, we have $b_0 = b_m = 0$. In these cases we can restrict the auxiliary space index r to $0 \leq r \leq m-1$ and the representation has finite dimension $d_{\mathcal{A}} = m$. Notice that for $q = e^{i\pi l/m}$ we have $\Delta = \cos(\pi l/m)$, hence $|\Delta| \leq 1$, which corresponds to the gapless phase of the XXZ model.

The matrices in Eqs. (3.7), (3.8) and (3.9) are functions of the complex parameter u . The Lax operator defined in Eq. (3.1) is a function of both u and the spectral parameter z . Similarly to Eq. (2.9), we can write

$$\begin{aligned} \mathcal{L}_j(z, u) = & i[\mathbb{1}_j \otimes A_0(z, u) + \sigma_j^z \otimes A_z(z, u) \\ & + \sigma_j^+ \otimes A_+(z, u) + \sigma_j^- \otimes A_-(z, u)], \end{aligned} \quad (3.14)$$

where

$$A_0(z, u) = \frac{(z - z^{-1})}{4i} [K(u) + K^{-1}(u)], \quad (3.15)$$

$$A_z(z, u) = \frac{(z + z^{-1})}{4i} [K(u) - K^{-1}(u)], \quad (3.16)$$

$$A_+(z, u) = \frac{z}{2i} (q - q^{-1}) S^-(u), \quad (3.17)$$

$$A_-(z, u) = \frac{z^{-1}}{2i} (q - q^{-1}) S^+(u). \quad (3.18)$$

In this notation, the conserved quantity defined in Eq. (3.3) reads (hereafter we omit the index \mathcal{A} in $\text{tr}_{\mathcal{A}}$)

$$t_{\mathcal{A}}(z, u) = i^N \sum_{\{\alpha_j\}} \text{tr}\{A_{\alpha_N} \dots A_{\alpha_2} A_{\alpha_1}\} \prod_{j=1}^N \sigma_j^{\alpha_j}, \quad (3.19)$$

where the sum is over all $\alpha_j \in \{0, z, +, -\}$ and we use the notation $\sigma_j^0 \equiv \mathbb{1}_j$.

The operator in Eq. (3.19) is translationally invariant due to the cyclic property of the trace. On the other hand, it is not necessarily invariant under spin reversal for general u . (The similarity between matrices used in Eq. (2.24) is not verified for arbitrary values of u .) Moreover, $t_{\mathcal{A}}(z, u)$ is not invariant under parity transformation \mathcal{P} , which we can define as the reflection about the link between sites $j = 1$ and $j = N$: $\mathcal{P}^{-1} \sigma_j^{\alpha_j} \mathcal{P} = \sigma_{N+1-j}^{\alpha_j}$. We have

$$\begin{aligned} \mathcal{P}^{-1} t_{\mathcal{A}}(z, u) \mathcal{P} &= i^N \sum_{\{\alpha_j\}} \text{tr}\{A_{\alpha_N} \dots A_{\alpha_2} A_{\alpha_1}\} \prod_{j=1}^N \sigma_{N+1-j}^{\alpha_j} \\ &= i^N \sum_{\{\alpha_j\}} \text{tr}\{A_{\alpha_1} \dots A_{\alpha_{N-1}} A_{\alpha_N}\} \prod_{j=1}^N \sigma_j^{\alpha_j}. \end{aligned} \quad (3.20)$$

We note that $t_{\mathcal{A}}$ in Eq. (3.19) can also be written as

$$t_{\mathcal{A}}(z, u) = i^N \sum_{\{\alpha_j\}} \text{tr}\{A_{\alpha_1}^t A_{\alpha_2}^t \dots A_{\alpha_N}^t\} \prod_{j=1}^N \sigma_j^{\alpha_j}, \quad (3.21)$$

where A_{α}^t denotes the transpose of A_{α} . We define the two-parameter conserved quantity which is odd under parity as

$$\mathcal{I}(z, u) = (-i)^N [\mathcal{P}^{-1} t_{\mathcal{A}}(z, u) \mathcal{P} - t_{\mathcal{A}}(z, u)]. \quad (3.22)$$

Therefore,

$$\mathcal{I}(z, u) = \sum_{\{\alpha_j\}} \text{tr}\{A_{\alpha_1} \dots A_{\alpha_N} - A_{\alpha_1}^t \dots A_{\alpha_N}^t\} \prod_{j=1}^N \sigma_j^{\alpha_j}. \quad (3.23)$$

For reference, let us comment on the particular cases $\Delta = 0$ and $\Delta = \pm 1$. For $\Delta = 0$ the XXZ model is equivalent to free fermions via a Jordan-Wigner transformation. This point corresponds to $m = 2$, $q = i$; in this case the generators of the quantum group algebra become

$$K = u \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}, \quad S^+ = \frac{u^{-2} - u^2}{2i} \sigma^-, \quad S^- = \sigma^+. \quad (3.24)$$

Note that, although the auxiliary space is two-dimensional $\mathcal{A} = \mathbb{C}^2$, the representation differs from Eq. (2.14) for general u . Only for $u = e^{-i\pi/4}$ do we recover a parity-invariant representation. On the other hand, at the ferromagnetic SU(2) point $\Delta = -1$ ($q = -1$, $m = 1$), the A_α matrices reduce to numbers and the conserved quantity in Eq. (3.23) vanishes identically. At the antiferromagnetic SU(2) point $\Delta = 1$ ($q = 1$, $m \rightarrow \infty$) the operator is not identically zero but the representation becomes infinite dimensional.

4. Quasilocal conserved quantities

Now we turn to the task of extracting quasilocal operators from $\mathcal{I}(z, u)$ in Eq. (3.23). In order to calculate the Mazur bound for the Drude weight at high temperatures [18], it is convenient to define the inner product between two operators A and B acting on $V^{\otimes N}$ based on the thermal average at infinite temperature:

$$\langle A^\dagger B \rangle = 2^{-N} \text{Tr}\{A^\dagger B\}, \quad (4.1)$$

where Tr denotes the trace over the quantum space $V^{\otimes N}$. From Eq. (4.1) it can be shown that the norm of $\mathcal{I}(z, u)$ reduces to

$$\langle \mathcal{I}^\dagger(z, u) \mathcal{I}(z, u) \rangle = 2 \text{tr}_{\mathcal{A} \otimes \mathcal{A}} \{[T_1(z, u, u)]^N - [T_2(z, u, u)]^N\}. \quad (4.2)$$

Here $T_1(z, u, \bar{u})$ and $T_2(z, u, \bar{u})$ are transfer matrices in $\mathcal{A} \otimes \mathcal{A}$

$$T_1(z, u, \bar{u}) = \sum_{\alpha=0,z,\pm} C_\alpha A_\alpha^*(z, u) \otimes A_\alpha(z, \bar{u}), \quad (4.3)$$

$$T_2(z, u, \bar{u}) = \sum_{\alpha=0,z,\pm} C_\alpha A_\alpha^*(z, u) \otimes A_\alpha^t(z, \bar{u}), \quad (4.4)$$

where

$$C_\alpha = \frac{1}{2} \text{Tr} \{ \sigma^\alpha (\sigma^\alpha)^\dagger \}. \quad (4.5)$$

In contrast with Prosen's construction for the open chain [16], where the norm is computed from the matrix element between boundary states, Eq. (4.2) involves the trace over the auxiliary space. The analogy with the open chain can be explored further if we notice that, by setting the spectral parameter to be $z = i$, the matrix A_z in Eq. (3.16) vanishes and the conserved quantity does not contain any σ_j^z operators, as

assumed in the original matrix product ansatz [16]. In the following we shall focus on the particular choice $z = i$. We return to the question of general values of z in section 7. For $z = i$ the nonvanishing operators in auxiliary space simplify to

$$A_0(z = i, u) = \sum_{r=0}^{m-2} \frac{uq^r + u^{-1}q^{-r}}{2} |r\rangle\langle r|, \quad (4.6)$$

$$A_+(z = i, u) = \sum_{r=0}^{m-2} \frac{q^{r+1} - q^{-r-1}}{2} |r\rangle\langle r+1|, \quad (4.7)$$

$$A_-(z = i, u) = \sum_{r=0}^{m-2} \frac{u^2q^r - u^{-2}q^{-r}}{2} |r+1\rangle\langle r|. \quad (4.8)$$

After fixing the value of the spectral parameter, we are still free to choose the value of u in the representation of the quantum group algebra. We notice that the condition $u^4 = 1$ is special because in this case $a_0 = 0$, then the state $|r = 0\rangle$ is annihilated by A_{\pm} and decouples from the other states. Hereafter we choose $u = 1$, but the result for the other roots is equivalent. For $u = 1$ the Casimir operator becomes $C = q + q^{-1}$. Interestingly, a similar kind of special representation appears in open spin chains where the quantum group is an actual symmetry commuting with the Hamiltonian [33]. In that case, the Casimir for a spin-1/2 representation takes the value $q^{N-1} + q^{-N+1}$ and becomes special if $q^N = -1$, *i.e.*, for values of q that obey a “root of unity condition” depending on the chain length [38].

Let us then analyze the operator

$$\mathcal{I}_0 \equiv \mathcal{I}(z = i, u = 1). \quad (4.9)$$

Setting $u = 1$ in Eqs. (4.6) through (4.8), we obtain (recall $q = e^{i\lambda}$)

$$A_0(1) \equiv A_0(z = i, u = 1) = \sum_{r=0}^{m-1} \cos(\lambda r) |r\rangle\langle r|, \quad (4.10)$$

$$A_+(1) \equiv A_+(z = i, u = 1) = i \sum_{r=0}^{m-2} \sin[\lambda(r+1)] |r\rangle\langle r+1|, \quad (4.11)$$

$$A_-(1) \equiv A_-(z = i, u = 1) = -i \sum_{r=0}^{m-2} \sin(\lambda r) |r+1\rangle\langle r|. \quad (4.12)$$

The transfer matrices in Eqs. (4.3) and (4.4) become

$$\begin{aligned} T_1(1) &\equiv T_1(z = i, u = 1, \bar{u} = 1) \\ &= \sum_{r,s=0}^{m-1} \cos(\lambda r) \cos(\lambda s) |r, s\rangle\langle r, s| \\ &\quad + \frac{1}{2} \sum_{r,s=0}^{m-2} \sin[\lambda(r+1)] \sin[\lambda(s+1)] |r, s\rangle\langle r+1, s+1| \\ &\quad + \frac{1}{2} \sum_{r,s=0}^{m-2} \sin(\lambda r) \sin(\lambda s) |r+1, s+1\rangle\langle r, s|, \end{aligned} \quad (4.13)$$

$$\begin{aligned}
T_2(1) &\equiv T_2(z = i, u = 1, \bar{u} = 1) \\
&= \sum_{r,s=0}^{m-1} \cos(\lambda r) \cos(\lambda s) |r, s\rangle \langle r, s| \\
&\quad + \frac{1}{2} \sum_{r,s=0}^{m-2} \sin[\lambda(r+1)] \sin[\lambda(s+1)] |r, s+1\rangle \langle r+1, s| \\
&\quad + \frac{1}{2} \sum_{r,s=0}^{m-2} \sin(\lambda r) \sin(\lambda s) |r+1, s\rangle \langle r, s+1|.
\end{aligned} \tag{4.14}$$

The transfer matrices are block diagonal in subspaces of Kronecker states $\{|r, (r+k)(\text{mod } m)\rangle\}$ with fixed $k = 0, \dots, m-1$ in the case of $T_1(1)$, or Kronecker states $\{|r, (-r+k)(\text{mod } m)\rangle\}$ in the case of $T_2(1)$. Since we are interested in the scaling of the operator norm in Eq. (4.2) with system size N as $N \rightarrow \infty$, we may restrict ourselves to the subspace in which the transfer matrices have their largest eigenvalue. This happens when $k = 0$ for both $T_1(1)$ and $T_2(1)$. Within the $k = 0$ subspace we denote $|r, \pm r\rangle \rightarrow |r\rangle$ and obtain the reduced transfer matrices

$$\begin{aligned}
\mathcal{T}_1 &= \sum_{r=0}^{m-1} \cos^2(\lambda r) |r\rangle \langle r| + \frac{1}{2} \sum_{r=0}^{m-2} \sin^2[\lambda(r+1)] |r\rangle \langle r+1| \\
&\quad + \frac{1}{2} \sum_{r=0}^{m-2} \sin^2(\lambda r) |r+1\rangle \langle r|,
\end{aligned} \tag{4.15}$$

$$\begin{aligned}
\mathcal{T}_2 &= \sum_{r=0}^{m-1} \cos^2(\lambda r) |r\rangle \langle r| - \frac{1}{2} \sum_{r=0}^{m-2} \sin(\lambda r) \sin[\lambda(r+1)] \times \\
&\quad \times [|r\rangle \langle r+1| + |r+1\rangle \langle r|].
\end{aligned} \tag{4.16}$$

It is useful to note that

$$\mathcal{T}_1 = \mathbb{1} - B^2 + \frac{1}{2} \Delta B^2, \tag{4.17}$$

$$\mathcal{T}_2 = \mathbb{1} - B^2 - \frac{1}{2} B \Delta B, \tag{4.18}$$

where B is the diagonal matrix $B = \sum_{r=0}^{m-1} \sin(r\lambda) |r\rangle \langle r|$ and Δ is the uniform hopping matrix on an open chain with length m

$$\Delta = \sum_{r=0}^{m-2} (|r\rangle \langle r+1| + |r+1\rangle \langle r|). \tag{4.19}$$

The matrix \mathcal{T}_2 is symmetric, thus its eigenvalues are all real. Since $B|0\rangle = 0$, we find that $|r=0\rangle$ is an eigenvector of \mathcal{T}_2 with eigenvalue 1. It is easy to verify that all the other eigenvalues are smaller than 1. § On the other hand, \mathcal{T}_1 is not symmetric. However, in Appendix A we show that \mathcal{T}_1 and \mathcal{T}_2 are similar and have exactly the same spectrum (see also Fig. 1). It also follows from Eq. (4.17) that $|r=0\rangle$ is the *right* eigenvector of

§ For $\lambda \in \mathbb{R}$, i.e. $|\Delta| \leq 1$, we can show that the largest eigenvalue of \mathcal{T}_2 is 1 using the Gershgorin circle theorem. In the gapped Neel phase $\Delta > 1$ the method used here only gives rise to nonlocal operators.




Figure 1. Absolute value of the eigenvalues of the transfer matrices $T_1(z = i, u, u)$ (solid blue lines) and $T_2(z = i, u, u)$ (dashed red lines) as a function of $u \in \mathbb{R}$ for $q = e^{i\pi/3}$. The quasilocal conserved quantity is obtained by expanding about $u = 1$, where T_1 and T_2 have the same spectrum and their largest eigenvalue is normalized to 1.

\mathcal{T}_1 with eigenvalue 1. We will also need the *left* eigenvector of \mathcal{T}_1 with eigenvalue 1. In Appendix B we show that the solution to the eigenvalue equation $\langle 0_L | \mathcal{T}_1 = \langle 0_L |$ yields

$$\langle 0_L | = \sum_{r=0}^{m-1} (1 - r/m) \langle r |. \quad (4.20)$$

The left eigenvector $\langle 0_L |$ is not normalized to unity but is such that $\langle 0_L | 0 \rangle = 1$.

When calculating the norm of the conserved quantity using Eq. (4.2), we can use the macroscopic number of transfer matrices to project the auxiliary space into the eigenvectors of \mathcal{T}_1 or \mathcal{T}_2 with eigenvalue 1. In Appendix C we show that

$$\lim_{n \rightarrow \infty} \mathcal{T}_1^n = \lim_{n \rightarrow \infty} \mathcal{T}_1^{n/2} \sum_r |r\rangle \langle r| \mathcal{T}_1^{n/2} = |0\rangle \langle 0_L|, \quad (4.21)$$

$$\lim_{n \rightarrow \infty} \mathcal{T}_2^n = \lim_{n \rightarrow \infty} \mathcal{T}_2^{n/2} \sum_r |r\rangle \langle r| \mathcal{T}_2^{n/2} = |0\rangle \langle 0|. \quad (4.22)$$

In particular, the projection allows us to compute traces involving an arbitrary matrix M

$$\lim_{n \rightarrow \infty} \text{tr}\{\mathcal{T}_1^n M\} = \langle 0_L | M | 0 \rangle, \quad (4.23)$$

$$\lim_{n \rightarrow \infty} \text{tr}\{\mathcal{T}_2^n M\} = \langle 0 | M | 0 \rangle. \quad (4.24)$$

It turns out that we do not get a quasilocal operator by simply setting $z = i, u = 1$. The reason is that, since \mathcal{T}_1 and \mathcal{T}_2 are related by a similarity transformation, the operator \mathcal{I}_0 in Eq. (4.9) actually has zero norm:

$$\langle \mathcal{I}_0^\dagger \mathcal{I}_0 \rangle = \text{tr}\{\mathcal{T}_1^N - \mathcal{T}_2^N\} = 0. \quad (4.25)$$

Nevertheless, the properties of the transfer matrices suggest that quasilocal operators can be generated by expanding $\mathcal{I}(z = i, u)$ about $u = 1$:

$$\mathcal{I}(z = i, u = 1 + \varepsilon) = \varepsilon \mathcal{I}_1 + \mathcal{O}(\varepsilon^2). \quad (4.26)$$

Let us then consider the operator

$$\begin{aligned}\mathcal{I}_1 &= \left. \frac{\partial \mathcal{I}(z, u)}{\partial u} \right|_{z=i, u=1} \\ &= \sum_{\{\alpha_j\}} \text{tr} \left\{ \frac{\partial}{\partial u} [A_{\alpha_1}(u) \dots A_{\alpha_N}(u) - A_{\alpha_1}^t(u) \dots A_{\alpha_N}^t(u)] \right\} \prod_{j=1}^N \sigma_j^{\alpha_j},\end{aligned}\quad (4.27)$$

with matrices $A_\alpha(u)$ given in Eqs. (4.6), (4.7) and (4.8). Using the transfer matrices in Eqs. (4.3) and (4.4), we can express the norm of \mathcal{I}_1 as follows:

$$\begin{aligned}\langle \mathcal{I}_1^\dagger \mathcal{I}_1 \rangle &= 2 \text{tr} \left\{ \frac{\partial^2}{\partial u \partial \bar{u}} [T_1(z, u, \bar{u})]^N \right\} \Big|_{z=i, u=\bar{u}=1} \\ &\quad - 2 \text{tr} \left\{ \frac{\partial^2}{\partial u \partial \bar{u}} [T_2(z, u, \bar{u})]^N \right\} \Big|_{z=i, u=\bar{u}=1}.\end{aligned}\quad (4.28)$$

The operators inside the trace in Eq. (4.28) contain a macroscopic number of transfer matrices. Once again, this allows us to restrict to the Kronecker spaces which contain eigenvectors with eigenvalue 1. Let us introduce a shorthand notation for the derivatives of the reduced transfer matrices:

$$\mathcal{T}_1^{(n, n')} \equiv \frac{\partial^n}{\partial u^n} \frac{\partial^{n'}}{\partial \bar{u}^{n'}} T_1(z = i, u, \bar{u}) \Big|_{u=\bar{u}=1}, \quad (4.29)$$

and likewise for $\mathcal{T}_2^{(n, n')}$. The derivatives in Eq. (4.28) yield

$$\begin{aligned}\frac{\langle \mathcal{I}_1^\dagger \mathcal{I}_1 \rangle}{N} &= 2 \text{tr} \left\{ (\mathcal{T}_1)^{N-1} \mathcal{T}_1^{(1,1)} + (\mathcal{T}_2)^{N-1} \mathcal{T}_2^{(1,1)} \right\} \\ &\quad + 2 \sum_{n=0}^{N-2} \text{tr} \left\{ \mathcal{T}_1^{(1,0)} (\mathcal{T}_1)^n \mathcal{T}_1^{(0,1)} (\mathcal{T}_1)^{N-2-n} \right\} \\ &\quad + 2 \sum_{n=0}^{N-2} \text{tr} \left\{ \mathcal{T}_2^{(1,0)} (\mathcal{T}_2)^n \mathcal{T}_2^{(0,1)} (\mathcal{T}_2)^{N-2-n} \right\}.\end{aligned}\quad (4.30)$$

In order for \mathcal{I}_1 to be quasilocal, the righthand side of Eq. (4.30) must approach a finite value in the limit $N \rightarrow \infty$. First consider the last two terms in Eq. (4.30). The derivatives of the A_α matrices at $u = 1$ are

$$A'_0(1) \equiv \left. \frac{\partial A_0}{\partial u} \right|_{u=1} = i \sum_{r=0}^{m-1} \sin(\lambda r) |r\rangle \langle r|, \quad (4.31)$$

$$A'_+(1) \equiv \left. \frac{\partial A_+}{\partial u} \right|_{u=1} = 0, \quad (4.32)$$

$$A'_-(1) \equiv \left. \frac{\partial A_-}{\partial u} \right|_{u=1} = -2 \sum_{r=0}^{m-2} \cos(\lambda r) |r+1\rangle \langle r|. \quad (4.33)$$

Thus,

$$\mathcal{T}_1^{(0,1)} = \mathcal{T}_1^{(1,0)}$$

$$= \frac{i}{2} \sum_{r=0}^{m-1} \sin(2\lambda r) |r\rangle \langle r| + \frac{i}{2} \sum_{r=0}^{m-2} \sin(2\lambda r) |r+1\rangle \langle r|, \quad (4.34)$$

$$\begin{aligned} \mathcal{T}_2^{(0,1)} &= i \sum_{r=0}^{m-1} \sin(\lambda r) \cos(\lambda r) |r\rangle \langle r| \\ &\quad + i \sum_{r=0}^{m-2} \sin(\lambda r) \cos[\lambda(r+1)] |r+1\rangle \langle r|, \end{aligned} \quad (4.35)$$

$$\begin{aligned} \mathcal{T}_2^{(1,0)} &= i \sum_{r=0}^{m-1} \sin(\lambda r) \cos(\lambda r) |r\rangle \langle r| \\ &\quad - i \sum_{r=0}^{m-2} \cos(\lambda r) \sin[\lambda(r+1)] |r+1\rangle \langle r|. \end{aligned} \quad (4.36)$$

We then notice that

$$\mathcal{T}_1^{(0,1)} |0\rangle = \mathcal{T}_1^{(1,0)} |0\rangle = 0, \quad (4.37)$$

$$\mathcal{T}_2^{(0,1)} |0\rangle = \langle 0 | \mathcal{T}_2^{(1,0)} = 0. \quad (4.38)$$

These relations are a result of the decoupling of state $|0\rangle$ from the other states at $u = 1$ (see comment around Eq. (4.9)). Together with the projection in Eqs. (4.21) and (4.22), these relations imply that the last two terms in Eq. (4.30) vanish in the thermodynamic limit.

We are left with the contributions in the first line of Eq. (4.30), which give

$$\lim_{N \rightarrow \infty} \frac{\langle \mathcal{I}_1^\dagger \mathcal{I}_1 \rangle}{N} = 2 \langle 0_L | \mathcal{T}_1^{(1,1)} | 0 \rangle + 2 \langle 0 | \mathcal{T}_2^{(1,1)} | 0 \rangle. \quad (4.39)$$

The derivatives of the reduced transfer matrices in Eq. (4.39) are

$$\mathcal{T}_1^{(1,1)} = \sum_{r=0}^{m-1} \sin^2(\lambda r) |r\rangle \langle r| + 2 \sum_{r=0}^{m-2} \cos^2(\lambda r) |r+1\rangle \langle r|, \quad (4.40)$$

$$\mathcal{T}_2^{(1,1)} = \sum_{r=0}^{m-1} \sin^2(\lambda r) |r\rangle \langle r| + 2 \sum_{r=0}^{m-2} \cos(\lambda r) \cos[\lambda(r+1)] |r+1\rangle \langle r|. \quad (4.41)$$

The only nonzero matrix element that contributes to the norm of \mathcal{I}_1 is $\langle 0_L | \mathcal{T}_1^{(1,1)} | 0 \rangle$. Using Eq. (4.20), we obtain

$$\lim_{N \rightarrow \infty} \frac{\langle \mathcal{I}_1^\dagger \mathcal{I}_1 \rangle}{N} = 4 \left(1 - \frac{1}{m} \right). \quad (4.42)$$

This proves \mathcal{I}_1 is quasilocal for $m > 1$.

Let us make some remarks about the conditions that lead to the norm's growing linearly with system size. This is expected to happen whenever we have a representation that becomes reducible for a specific value of a continuous parameter ($u = 1$ in our case, see Eq. (4.9)) and the subrepresentation obtained in this case (the single state $|r = 0\rangle$) is parity invariant. It is then clear that the two terms in Eq. (4.2) are equal to Λ^N , where Λ is the largest eigenvalue in the parity-invariant subspace (assuming it dominates the

norm). Expanding the operators around this specific value with $\delta u = \varepsilon \ll 1$, we find that the transfer matrices in Eq. (4.2) behave as (here $\nu = 1, 2$)

$$T_\nu(1 + \varepsilon) = \begin{pmatrix} \Lambda & \varepsilon B_\nu \\ \varepsilon C_\nu & D + \varepsilon F_\nu \end{pmatrix}, \quad (4.43)$$

where D, F_ν are matrices in the subspace orthogonal to the parity invariant subspace. As a result, the eigenvalues behave as $\text{Tr}\{T_\nu^N\} = (\Lambda + \varepsilon^2 A_\nu)^N \approx \Lambda^N + N\varepsilon^2 A_\nu$. The norm will then be linear in N as long as $A_1 \neq A_2$.

5. Nonlocal operators generated in the expansion of $\mathcal{I}(z, u)$

The expansion in Eq. (4.26) to higher orders in $\varepsilon = u - 1$ gives rise to the family of operators

$$\mathcal{I}_\ell = \left. \frac{\partial^\ell}{\partial u^\ell} \mathcal{I}(z = i, u) \right|_{u=1}. \quad (5.1)$$

But \mathcal{I}_1 is the only quasilocal operator in this series because in general the norm of \mathcal{I}_ℓ scales like N^ℓ for $N \rightarrow \infty$. To see this, consider the case of \mathcal{I}_2 :

$$\begin{aligned} \langle \mathcal{I}_2^\dagger \mathcal{I}_2 \rangle &= 2 \text{tr} \left\{ \frac{\partial^4}{\partial u^2 \partial \bar{u}^2} [T_1(z, u, \bar{u})]^N \right\} \Big|_{z=i, u=\bar{u}=1} \\ &\quad - 2 \text{tr} \left\{ \frac{\partial^4}{\partial u^2 \partial \bar{u}^2} [T_2(z, u, \bar{u})]^N \right\} \Big|_{z=i, u=\bar{u}=1}. \end{aligned} \quad (5.2)$$

When applying derivatives in Eq. (5.2), we can discard terms which contain $\mathcal{T}_1^{(0,1)}, \mathcal{T}_1^{(1,0)}, \mathcal{T}_2^{(0,1)}, \mathcal{T}_2^{(1,0)}$, since their contribution vanishes in the thermodynamic limit. The result for large N is

$$\begin{aligned} \frac{\langle \mathcal{I}_2^\dagger \mathcal{I}_2 \rangle}{N} &= 2 \text{tr} \left[\mathcal{T}_1^{(2,2)} (\mathcal{T}_1)^{N-1} \right] + 4 \sum_{n=0}^{N-2} \text{tr} \left[\mathcal{T}_1^{(1,1)} (\mathcal{T}_1)^n \mathcal{T}_1^{(1,1)} (\mathcal{T}_1)^{N-2-n} \right] \\ &\quad + 2 \sum_{n=0}^{N-2} \text{tr} \left[\mathcal{T}_1^{(2,0)} (\mathcal{T}_1)^n \mathcal{T}_1^{(0,2)} (\mathcal{T}_1)^{N-2-n} \right] - (\mathcal{T}_1 \rightarrow \mathcal{T}_2). \end{aligned} \quad (5.3)$$

In contrast with Eq. (4.30), the terms on the righthand side of Eq. (5.3) that involve sums do not vanish identically because the matrices $\mathcal{T}_1^{(1,1)}, \mathcal{T}_1^{(2,0)}, \mathcal{T}_1^{(0,2)}$ do not annihilate the state $|0\rangle$ (and likewise for \mathcal{T}_2). In fact, the result of the sum increases linearly with N since the trace does not decay with the number n of \mathcal{T}_1 's between the derivatives. The coefficient of the $\mathcal{O}(N^2)$ term in the norm of \mathcal{I}_2 stems from terms in the sums with $n \sim N$. In order to extract this coefficient, we insert another projection onto the eigenvectors of \mathcal{T}_1 or \mathcal{T}_2 with eigenvalue 1 and obtain

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{\langle \mathcal{I}_2^\dagger \mathcal{I}_2 \rangle}{N^2} &= 4 \langle 0_L | \mathcal{T}_1^{(1,1)} | 0 \rangle^2 - 4 \langle 0 | \mathcal{T}_2^{(1,1)} | 0 \rangle^2 + 2 \langle 0_L | \mathcal{T}_1^{(2,0)} | 0 \rangle \langle 0_L | \mathcal{T}_1^{(0,2)} | 0 \rangle \\ &\quad - 2 \langle 0 | \mathcal{T}_2^{(2,0)} | 0 \rangle \langle 0 | \mathcal{T}_2^{(0,2)} | 0 \rangle. \end{aligned} \quad (5.4)$$

It is easy to verify that the last two terms in Eq. (5.4) cancel out. Using the matrices in Eq. (4.40) and (4.41), we find

$$\lim_{N \rightarrow \infty} \frac{\langle \mathcal{I}_2^\dagger \mathcal{I}_2 \rangle}{N^2} = 16 \left(1 - \frac{1}{m} \right)^2. \quad (5.5)$$

For general $\ell \geq 1$, the expression for the norm of \mathcal{I}_ℓ contains terms in which a number ℓ of matrices $\mathcal{T}_1^{(1,1)}$ are distributed over the N sites of the chain. Since the contribution in the trace does not decrease with the separation between the $\mathcal{T}_1^{(1,1)}$'s, the norm grows with the number of ways to choose the positions of these matrices when they are far apart, therefore $\langle \mathcal{I}_\ell^\dagger \mathcal{I}_\ell \rangle \propto N^\ell$.

6. Mazur bound

Within linear response theory, the optical conductivity for a given model is related to the dynamical current-current correlation function via the Kubo formula. The real part of the optical conductivity can be written as

$$\sigma'(\omega) = 2\pi D \delta(\omega) + \sigma_{reg}(\omega), \quad (6.1)$$

where D is the Drude weight and $\sigma_{reg}(\omega)$ is the regular part. A nonzero Drude weight implies infinite dc conductivity, *i.e.*, ballistic transport. The connection between integrability and transport is made particularly clear by means of the Mazur bound [18] for the Drude weight at finite temperature T

$$D \geq \frac{1}{2LT} \sum_k \frac{|\langle J Q_k \rangle|^2}{\langle Q_k^\dagger Q_k \rangle}. \quad (6.2)$$

Here L is the system size, $\langle \rangle$ denotes the thermal average, J is the current operator and $\{Q_k\}$ is a set of operators that commute with the Hamiltonian and are orthogonalized in the form $\langle Q_k^\dagger Q_l \rangle = \delta_{kl} \langle Q_k^\dagger Q_k \rangle$. Although integrable models possess an infinite number of conserved quantities in the thermodynamic limit, it suffices to find one single operator that gives a nonzero contribution to the right hand side of Eq. (6.2) in order to establish ballistic transport.

The current operator is obtained from the continuity equation for the density of the conserved charge. The spin current operator for the XXZ model (2.1) reads

$$J = i \sum_j (\sigma_j^+ \sigma_{j+1}^- - \sigma_j^- \sigma_{j+1}^+). \quad (6.3)$$

This operator is odd under spin inversion:

$$\mathcal{C}^{-1} J \mathcal{C} = -J. \quad (6.4)$$

As discussed in the section 2, all the local conserved quantities derived from the transfer matrix t_Q are invariant under spin inversion. This includes the XXZ Hamiltonian at zero magnetic field. As a result, $\langle J Q_n \rangle = 0$ for all the local Q_n 's.

Let us now show that the quasilocal operator \mathcal{I}_1 in Eq. (4.27) provides a nonzero Mazur bound at zero magnetic field. Notice that, since $[\mathcal{I}_1, H] = 0$ exactly, there is no




Figure 2. Mazur bound from $\mathcal{I}(z = i, u)$ as a function of chain length N for $q = e^{i\pi/3}$ and three different values of the parameter u : $u = 1.01$ (red circles), $u = 1.1$ (green squares) and $u = 1.4$ (blue triangles). The dashed line represents the analytical result for $u \rightarrow 1$ and $N \rightarrow \infty$ in Eq. (6.7).

issue with the violation of the conservation law by boundary terms as in the open chain [29]. We need to calculate the overlap between J and \mathcal{I}_1

$$\begin{aligned}
 \langle J\mathcal{I}_1 \rangle &= 2^{-N} \text{Tr}\{J\mathcal{I}_1\} \\
 &= Ni \sum_{\{\alpha_j\}} \frac{\partial}{\partial u} \text{tr} \{A_{\alpha_1}(u) \dots A_{\alpha_N}(u) - A_{\alpha_1}^t(u) \dots A_{\alpha_N}^t(u)\} \Big|_{u=1} \times \\
 &\quad \times 2^{-N} \text{Tr}\left\{ \prod_{j=1}^N \sigma_j^{\alpha_j} (\sigma_1^+ \sigma_2^- - \sigma_1^- \sigma_2^+) \right\} \\
 &= \frac{Ni}{2} \frac{\partial}{\partial u} \text{tr}\{[A_-(u), A_+(u)](A_0(u))^{N-2}\} \Big|_{u=1}. \tag{6.5}
 \end{aligned}$$

For $N \rightarrow \infty$, the factors of $(A_0)^N$ project the auxiliary space into $|0\rangle$, which is the eigenvector of A_0 with eigenvalue 1. The nonzero contribution stems from applying to derivative to $A_-(u)$:

$$\langle J\mathcal{I}_1 \rangle = -\frac{Ni}{2} \langle 0|A_+(1)A'_-(1)|0\rangle = -N \sin \lambda. \tag{6.6}$$

Using the result for the norm in Eq. (4.42), we find that the contribution from \mathcal{I}_1 to the Mazur bound in the high temperature limit is of the form $D \geq D_{\mathcal{I}_1}/4T$ with

$$D_{\mathcal{I}_1} = \lim_{N \rightarrow \infty} \frac{2\langle J\mathcal{I}_1 \rangle^2}{N\langle \mathcal{I}_1^\dagger \mathcal{I}_1 \rangle} = \frac{\sin^2 \lambda}{2} \frac{m}{m-1}. \tag{6.7}$$

This result agrees with the bound obtained from the quasi conserved operator for the open chain [16].

Since $\mathcal{I}_0 = \mathcal{I}(z = i, u = 1)$ vanishes, the result in Eq. (6.7) can also be written as

$$D_{\mathcal{I}_1}(z = i) = \lim_{N \rightarrow \infty} \lim_{u \rightarrow 1} \frac{2|\langle J\mathcal{I}(z = i, u) \rangle|^2}{N\langle \mathcal{I}^\dagger(z = i, u) \mathcal{I}(z = i, u) \rangle}, \tag{6.8}$$

where the order of the limits matters. The role of the limit $u \rightarrow 1$ before $N \rightarrow \infty$ is illustrated in Fig. 2, where we calculate the Mazur bound for finite chains numerically

without using the projection into subspaces of largest eigenvalues. For $u^4 \neq 1$, the conserved quantity $\mathcal{I}(z = i, u)$ is nonlocal and its contribution to the Mazur bound decreases exponentially with system size. For small $|u - 1|$ and large finite N , the Mazur bound approaches a plateau that agrees with the analytical result in the thermodynamic limit.

7. Continuous family of quasilocal operators

The choice of the spectral parameter $z = i$ in Eq. (4.27) is not required to derive a quasilocal conserved quantity. Generalizing the results of section 4 to arbitrary values of z , we find that the norm of

$$\mathcal{I}_1(z) = \left. \frac{\partial}{\partial u} \mathcal{I}(z, u) \right|_{u=1} \quad (7.1)$$

can be computed from the reduced transfer matrices (*cf.* Eqs. (4.15) and (4.16))

$$\begin{aligned} \mathcal{T}_1(z) = \sum_{r=0}^{m-1} \left\{ [(\operatorname{Im} z)^2 \cos^2(\lambda r) + (\operatorname{Re} z)^2 \sin^2(\lambda r)] |r\rangle\langle r| \right. \\ \left. + \frac{|z|^2}{2} \sin^2[\lambda(r+1)] |r\rangle\langle r+1| + \frac{|z|^{-2}}{2} \sin^2(\lambda r) |r+1\rangle\langle r| \right\}, \quad (7.2) \end{aligned}$$

$$\begin{aligned} \mathcal{T}_2(z) = \sum_{r=0}^{m-1} \left\{ [(\operatorname{Im} z)^2 \cos^2(\lambda r) + (\operatorname{Re} z)^2 \sin^2(\lambda r)] |r\rangle\langle r| \right. \\ \left. - \frac{1}{2} \sin(\lambda r) \sin[\lambda(r+1)] [|z|^2 |r\rangle\langle r+1| + |z|^{-2} |r+1\rangle\langle r|] \right\}. \quad (7.3) \end{aligned}$$

The state $|r = 0\rangle$ is an eigenvector of $\mathcal{T}_2(z)$ and a right eigenvector of $\mathcal{T}_1(z)$, with eigenvalue $\Lambda = (\operatorname{Im} z)^2$. As discussed at the end of section 4, the condition for quasilocality is that Λ be the largest eigenvalue of both transfer matrices. As shown in [35], this condition is satisfied by a continuous set of values of z in the complex plane.

As a measure of quasilocality, we use the Mazur bound in Eq. (6.8) generalized to arbitrary z . A nonzero value of $D_{\mathcal{I}_1}(z)$ implies that the norm of $\mathcal{I}_1(z)$ is extensive. Fig. 3 illustrates the magnitude of the Mazur bound $D_{\mathcal{I}_1}(z)$ in the complex z plane for $q = e^{i\pi/3}$ ($\Delta = 1/2$). We find that $D_{\mathcal{I}_1}(z)$ is maximum at $z = \pm i$, where it assumes the value predicted by Eq. (6.7). The domain where $D_{\mathcal{I}_1}(z) > 0$ was discussed in [35]: writing $z = |z|e^{i\theta}$, the conserved operator $\mathcal{I}_1(z)$ is quasilocal inside the cone $||\theta| - \frac{\pi}{2}| < \frac{\pi}{2m}$.

Therefore, the Mazur bound obtained from a *single* quasilocal operator is maximized by the choice $z = \pm i$. However, the entire continuous family $\{\mathcal{I}_1(z)\}$ can be used to raise the bound. The idea is to replace the sum on the rhs of Eq. (6.2) by an integral over the spectral parameter z , using the orthogonality between different elements in the family, as done in [35].




Figure 3. Magnitude of the Mazur bound $D_{\mathcal{I}_1}(z)$ calculated using a single quasilocal operator $\mathcal{I}_1(z)$, as a function of the spectral parameter z , for $q = e^{i\pi/3}$. Brighter regions correspond to larger values of $D_{\mathcal{I}_1}(z)$.

8. Conclusion

We have described a method to derive quasilocal operators which commute with the Hamiltonian of the XXZ chain with periodic boundary conditions. The key to this procedure is to introduce an auxiliary transfer matrix $t_A(z, u)$ that depends on two parameters, namely the usual spectral parameter z and the representation parameter u . The latter is a parameter of the highest-weight representation for the quantum group algebra that arises in the Yang-Baxter relation for the Lax operator. For values of anisotropy $\Delta = \cos(\pi l/m)$, with l, m integers, the highest-weight representation has finite dimension m . The two-parameter conserved operator $\mathcal{I}(z, u)$ that has a nonzero overlap with the spin current operator is defined from a linear combination of the auxiliary transfer matrix and its conjugate under parity. The norm of $\mathcal{I}(z, u)$ can be calculated using the transfer matrices $T_1(z, u)$ and $T_2(z, u)$, which are related by a similarity transformation. A quasilocal operator is obtained by expanding $\mathcal{I}(z, u)$ about the special value $u = 1$ where the highest-weight representation becomes reducible and the eigenvector of $T_1(z, u = 1)$ and $T_2(z, u = 1)$ with the largest eigenvalue decouples from the other states. The remaining spectral parameter z labels a continuous family of quasilocal conserved quantities $\{\mathcal{I}_1(z)\}$. This is in contrast with the usual discrete set of local conserved quantities which are obtained by taking logarithmic derivatives of the transfer matrix $t_Q(z)$ (defined with a spin-1/2 representation in auxiliary space).

It has been shown that the quasilocal operators are important to set a nonzero lower bound for the Drude weight in the spin-1/2 XXZ chain [16]. An important open question is whether there exist other families of quasilocal operators beyond the ones derived by

this method. Additional conserved quantities may be expected from the observation that the Mazur bound computed from the set $\{\mathcal{I}_1(z)\}$ has a fractal Δ dependence [16, 29] which is perhaps absent in the actual Drude weight at high temperatures [25].

We note that, while here we have focused on the periodic chain, it should be possible to apply the same techniques to integrable models with open boundaries, taking into account reflection operators at the boundaries. In fact, in [39] a two-parameter family of transfer matrices has been constructed for the open asymmetric simple exclusion process (ASEP) (see Eqs. (47) and (48) of [39], which are the generalization of the conserved quantities to the open case). The effect of the boundary parameters on transport properties is an interesting open question.

The role of quasilocal conserved quantities in the GGE also remains to be clarified [10, 11, 12]. Remarkably, there is evidence that expectation values of local observables in post-quench steady states deviate from the predictions of the GGE even for $\Delta > 1$ [10, 12], *i.e.* in the gapped Néel phase, where the method described here does not yield any quasilocal operators.

Acknowledgements

We thank Fabian Essler, Andreas Klümper, and Grégoire Misguich for discussions. This work is supported by CNPq (R.G.P.), the SFB/TR 49 (J.S.), NSERC (J.S., I.A.), and CIFAR (I.A.).

Appendix A. Similarity between reduced transfer matrices \mathcal{T}_1 and \mathcal{T}_2

The transfer matrices defined in section 4 are

$$\mathcal{T}_1 = \begin{pmatrix} 1 & \frac{1}{2} \sin^2 \lambda & 0 & 0 & \dots & 0 \\ 0 & \cos^2 \lambda & \frac{1}{2} \sin^2 2\lambda & 0 & \dots & 0 \\ 0 & \frac{1}{2} \sin^2 \lambda & \cos^2 2\lambda & \frac{1}{2} \sin^2 3\lambda & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \cos^2(m-1)\lambda \end{pmatrix}, \quad (\text{A.1})$$

$$\mathcal{T}_2 = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \cos^2 \lambda & -\frac{1}{2} \sin \lambda \sin 2\lambda & \dots & 0 \\ 0 & -\frac{1}{2} \sin \lambda \sin 2\lambda & \cos^2 2\lambda & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \cos^2(m-1)\lambda \end{pmatrix}. \quad (\text{A.2})$$

First we note that the sign of the off-diagonal terms of \mathcal{T}_2 can be changed by applying the “ Z_2 gauge transformation” $\mathcal{S}|r\rangle = (-1)^r|r\rangle$. Defining $\tilde{\mathcal{T}}_2 = \mathcal{S}^{-1}\mathcal{T}_2\mathcal{S}$, we obtain

$$\tilde{\mathcal{T}}_2 = \mathbb{1} - B^2 + \frac{1}{2}B\Delta B, \quad (\text{A.3})$$

which is to be compared to Eqs. (4.17) and (4.18).

Let us then show that \mathcal{T}_1 is similar to $\tilde{\mathcal{T}}_2$. It suffices to show that they have the same characteristic polynomial. The characteristic polynomial for \mathcal{T}_1 reads

$$\det(\mathcal{T}_1 - x\mathbb{1}) = (1 - x) \det \left[(1 - x)\mathbb{1} - \tilde{B}^2 + \frac{1}{2}\tilde{\Delta}\tilde{B}^2 \right], \quad (\text{A.4})$$

where \tilde{B} is the diagonal matrix $\tilde{B} = \sum_{r=1}^{m-1} \sin(r\lambda)|r\rangle\langle r|$ and $\tilde{\Delta}$ is the uniform hopping matrix $\tilde{\Delta} = \sum_{r=1}^{m-2} (|r\rangle\langle r+1| + |r+1\rangle\langle r|)$ in the $(m-1)$ -dimensional space. Likewise, the characteristic polynomial for $\tilde{\mathcal{T}}_2$ (and also for \mathcal{T}_2 , given the similarity between them) is

$$\det(\tilde{\mathcal{T}}_2 - x\mathbb{1}) = (1 - x) \det \left[(1 - x)\mathbb{1} - \tilde{B}^2 + \frac{1}{2}\tilde{B}\tilde{\Delta}\tilde{B} \right]. \quad (\text{A.5})$$

Since \tilde{B} is invertible (all of its eigenvalues are nonzero for $\lambda = l\pi/m$ with l, m coprimes), we can apply a similarity transformation inside the determinant sign as follows:

$$\begin{aligned} \det(\tilde{\mathcal{T}}_2 - x\mathbb{1}) &= (1 - x) \det \left[\tilde{B}^{-1} \left((1 - x)\mathbb{1} - \tilde{B}^2 + \frac{1}{2}\tilde{B}\tilde{\Delta}\tilde{B} \right) \tilde{B} \right] \\ &= (1 - x) \det \left[(1 - x)\mathbb{1} - \tilde{B}^2 + \frac{1}{2}\tilde{\Delta}\tilde{B}^2 \right] \\ &= \det(\mathcal{T}_1 - x\mathbb{1}). \end{aligned} \quad (\text{A.6})$$

This shows that \mathcal{T}_1 and \mathcal{T}_2 are similar.

Appendix B. Left eigenvector of \mathcal{T}_1 with eigenvalue 1

Let $|0_L\rangle$ denote the left eigenvectors of \mathcal{T}_1 with eigenvalue 1, which obeys

$$\langle 0_L | \mathcal{T}_1 = \langle 0_L |. \quad (\text{B.1})$$

We expand $|0_L\rangle$ in the orthonormal basis of $\{|r\rangle\}$ vectors

$$|0_L\rangle = \sum_{r=0}^{m-1} v_r |r\rangle. \quad (\text{B.2})$$

Our problem is then to find the coefficients v_r . For short, we denote the matrix elements of \mathcal{T}_1 as $\langle i | \mathcal{T}_1 | j \rangle = t_{i,j}$. A useful identity is

$$t_{r,r+1}t_{r+1,r} = \frac{1}{4}(1 - t_{r,r})(1 - t_{r+1,r+1}). \quad (\text{B.3})$$

The eigenvalue equation (B.1) is satisfied identically for the column $r = 0$. This corresponds to the freedom of choosing the value of v_0 (or the normalization of $|0_L\rangle$). Let us turn to the next simplest equation, the one stemming from the column $r = m-1$:

$$v_{m-2}t_{m-2,m-1} + v_{m-1}t_{m-1,m-1} = v_{m-1}, \quad (\text{B.4})$$

from which we get

$$v_{m-1} = \frac{t_{m-2,m-1}}{(1 - t_{m-1,m-1})C_0} v_{m-2}, \quad (\text{B.5})$$

with $C_0 = 1$. Next, the equation for column $r = m - 2$ reads

$$v_{m-3}t_{m-3,m-2} + v_{m-2}t_{m-2,m-2} + v_{m-1}t_{m-1,m-2} = v_{m-2}. \quad (\text{B.6})$$

Using Eqs. (B.3) and (B.6), we obtain

$$v_{m-2} = \frac{t_{m-3,m-2}}{(1 - t_{m-2,m-2})C_1} v_{m-3}, \quad (\text{B.7})$$

with $C_1 = 1 - 1/(4C_0)$. In general, we find for $r = 1, \dots, m - 1$

$$v_{m-r} = \frac{t_{m-r-1,m-r}}{(1 - t_{m-r,m-r})C_{r-1}} v_{m-r-1}, \quad (\text{B.8})$$

with $C_0 = 1$ and

$$C_r = 1 - \frac{1}{4C_{r-1}}, \quad 1 \leq r \leq m - 2. \quad (\text{B.9})$$

This relation express C_r as a continued fraction and the solution can be readily seen to be

$$C_r = \frac{r + 2}{2r + 2}. \quad (\text{B.10})$$

In addition, we can use the explicit expression for the matrix elements of \mathcal{T}_1 in Eq. (A.1), which gives

$$\frac{t_{r,r+1}}{1 - t_{r,r}} = \frac{\frac{1}{2} \sin^2 \lambda r}{1 - \cos^2 \lambda r} = \frac{1}{2}. \quad (\text{B.11})$$

Thus Eq. (B.8) simplifies to

$$v_{m-r} = \frac{r}{r + 1} v_{m-r-1}, \quad r = 2, \dots, m. \quad (\text{B.12})$$

Writing the coefficients v_r , $r = 1, \dots, m - 1$, in terms of v_0 , we find

$$v_r = \left(1 - \frac{r}{m}\right) v_0. \quad (\text{B.13})$$

Finally, setting $v_0 = 1$ we obtain the vector in Eq. (B.2)

$$|0_L\rangle = \sum_{r=0}^{m-1} \left(1 - \frac{r}{m}\right) |r\rangle. \quad (\text{B.14})$$

Appendix C. Calculating traces in the thermodynamic limit

Let M_1 be an arbitrary $m \times m$ matrix (not necessarily Hermitean), where m is the dimension of the auxiliary space \mathcal{A} . We want to compute $\text{tr}\{M_1\}$ using the eigenvectors of \mathcal{T}_1 . Since \mathcal{T}_1 is non-Hermitean, its right and left eigenvectors are different:

$$\mathcal{T}_1 |R_j\rangle = \lambda_j |R_j\rangle, \quad (\text{C.1})$$

$$\langle L_j | \mathcal{T}_1 = \lambda_j \langle L_j |. \quad (\text{C.2})$$

Nonetheless, the left and right eigenvalues are equivalent because \mathcal{T}_1 and $(\mathcal{T}_1)^t$ have the same characteristic polynomial [40].

Right eigenvectors with different eigenvalues are not necessarily orthogonal, *i.e.* $\langle R_j | R_l \rangle \neq \delta_{j,l}$. Let $\{|r\rangle\}$ denote the orthonormal basis of vectors representing sites in the auxiliary space. We can expand the vectors $|r\rangle$ in the non-orthogonal eigenvector basis in the form

$$|r\rangle = \sum_j V_{r,j} |R_j\rangle, \quad (\text{C.3})$$

$$|r\rangle = \sum_j W_{r,j} |L_j\rangle. \quad (\text{C.4})$$

The transpose of Eq. (C.4) yields

$$\langle r| = \sum_j \langle L_j | W_{r,j} = \sum_j \langle L_j | W_{j,r}^t. \quad (\text{C.5})$$

The inverse transformation reads

$$|R_i\rangle = \sum_r (V^{-1})_{i,r} |r\rangle, \quad (\text{C.6})$$

$$|L_i\rangle = \sum_j (W^{-1})_{i,r} |r\rangle. \quad (\text{C.7})$$

It can be proved that the left eigenvectors are orthogonal to right eigenvectors with different eigenvalues (and degenerate eigenvectors can be orthogonalized) [40], so that

$$(W^{-1})^t V^{-1} = D, \quad (\text{C.8})$$

with

$$D_{j,l} = \langle L_j | R_l \rangle = d_j \delta_{j,l}. \quad (\text{C.9})$$

Thus the inverse of D is also diagonal:

$$(W^t V)_{j,l} = \frac{1}{d_j} \delta_{j,l}. \quad (\text{C.10})$$

The trace of M_1 can be written as

$$\begin{aligned} \text{Tr}\{M_1\} &= \sum_r \langle r | M_1 | r \rangle \\ &= \sum_{r,j,l} W_{j,r}^t V_{r,l} \langle L_j | M_1 | R_l \rangle \\ &= \sum_{j,l} (W^t V)_{j,l} \langle L_j | M_1 | R_l \rangle \\ &= \sum_j \frac{\langle L_j | M_1 | R_j \rangle}{d_j}. \end{aligned} \quad (\text{C.11})$$

In our case M_1 is a product of a large number ($\sim \mathcal{O}(N)$) of transfer matrices \mathcal{T}_1 on the left and on the right. In the thermodynamic limit the trace is dominated by the contributions from the eigenvector of \mathcal{T}_1 with eigenvalue $\lambda_j = 1$:

$$|R_1\rangle = |0\rangle, \quad \langle L_1| = \langle 0_L|. \quad (\text{C.12})$$

Thus

$$\lim_{N \rightarrow \infty} \text{Tr}\{M_1\} = \frac{\langle L_1 | M_1 | R_1 \rangle}{d_1} = \frac{\langle 0_L | \tilde{M}_1 | 0 \rangle}{\langle 0_L | 0 \rangle}, \quad (\text{C.13})$$

where \tilde{M}_1 is obtained from M_1 by dropping the factors of \mathcal{T}_1^N . Using eigenvectors normalized as in Eqs. (4.20), we can write simply

$$\lim_{N \rightarrow \infty} \text{Tr}\{M_1\} = \langle 0_L | \tilde{M}_1 | 0 \rangle. \quad (\text{C.14})$$

The following relation is also useful:

$$\begin{aligned} \sum_r |r\rangle\langle r| &= \sum_{r,j,l} V_{r,j} W_{l,r}^t |R_j\rangle\langle L_l| \\ &= \sum_{j,l} (W^t V)_{l,j} |R_j\rangle\langle L_l| \\ &= \sum_j \frac{|R_j\rangle\langle L_j|}{d_j}. \end{aligned} \quad (\text{C.15})$$

If there is a large number of \mathcal{T}_1 's on both sides we can project onto the eigenvectors with eigenvalue $\lambda_j = 1$

$$\lim_{n \rightarrow \infty} \left(\mathcal{T}_1^n \sum_r |r\rangle\langle r| \mathcal{T}_1^n \right) = \frac{|0\rangle\langle 0_L|}{\langle 0_L | 0 \rangle} = |0\rangle\langle 0_L|. \quad (\text{C.16})$$

We also need to calculate traces involving \mathcal{T}_2 . These are easier because \mathcal{T}_2 is symmetric and the eigenvector with eigenvalue 1 is simply $|0\rangle$. Thus the trace of a matrix with $\mathcal{O}(N)$ factors of \mathcal{T}_2 can be reduced to

$$\lim_{N \rightarrow \infty} \text{Tr}\{M_2\} = \langle 0 | \tilde{M}_2 | 0 \rangle. \quad (\text{C.17})$$

The equivalent of Eq. (C.16) for \mathcal{T}_2 is

$$\lim_{n \rightarrow \infty} \left(\mathcal{T}_2^n \sum_r |r\rangle\langle r| \mathcal{T}_2^n \right) = |0\rangle\langle 0|. \quad (\text{C.18})$$

- [1] B. Sutherland, *Beautiful Models* (World Scientific, Singapore, 2004).
- [2] V. E. Korepin, A. G. Izergin, and N. M. Bogoliubov, *Quantum inverse scattering method and correlation functions* (Cambridge Univ. Press, Cambridge, 1993).
- [3] A. R. Chowdhury and A. G. Choudhury, *Quantum Integrable Systems* (Chapman & Hall/CRC, 2004).
- [4] T. Kinoshita, T. Wenger, and D. S. Weiss, *Nature* **440**, 900 (2006).
- [5] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, *Nat. Phys.* **8**, 325 (2012).
- [6] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008).
- [7] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Rev. Mod. Phys.* **83**, 863 (2011).
- [8] J. Sirker, N.P. Konstantinidis, N. Sedlmayr, *Phys. Rev. A* **89**, 042104 (2014).
- [9] M. Fagotti, *J. Stat. Mech.* (2014) P03016.
- [10] B. Wouters, M. Brockmann, J. De Nardis, D. Fioretto, and J.-S. Caux, *Phys. Rev. Lett.* **113**, 117202 (2014).
- [11] M. Mierzejewski, P. Prelovšek, and T. Prosen, *Phys. Rev. Lett.* **113**, 020602 (2014).
- [12] B. Pozsgay, M. Mestyán, M. A. Werner, M. Kormos, G. Zaránd, G. Takács, *Phys. Rev. Lett.* **113**, 117203 (2014).

- [13] M. Fagotti, M. Collura, F. H. L. Essler, and P. Calabrese, *Phys. Rev. B* **89**, 125101 (2014).
- [14] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. Lett.* **111**, 127201 (2013).
- [15] D. A. Huse and V. Oganesyan, arXiv:1305.4915.
- [16] T. Prosen, *Phys. Rev. Lett.* **106**, 217206 (2011).
- [17] P. Mazur, *Physica* **43**, 533 (1969).
- [18] X. Zotos, F. Naef, and P. Prelovšek, *Phys. Rev. B* **55**, 11029 (1997).
- [19] B. N. Narozhny, A. J. Millis, and N. Andrei, *Phys. Rev. B* **58**, R2921(R) (1998).
- [20] X. Zotos, *Phys. Rev. Lett.* **82**, 1764 (1999).
- [21] J. V. Alvarez and C. Gros, *Phys. Rev. Lett.* **88**, 077203 (2002).
- [22] F. Heidrich-Meisner, A. Honecker, D. C. Cabra, and W. Brenig, *Phys. Rev. B* **68**, 134436 (2003).
- [23] J. Benz, T. Fukui, A. Klümper, and C. Scheeren, *J. Phys. Soc. Jpn. Suppl.* **74**, 181 (2005).
- [24] J. Sirker, R. G. Pereira, and I. Affleck, *Phys. Rev. Lett.* **103**, 216602 (2009); *Phys. Rev. B* **83**, 035115 (2011).
- [25] C. Karrasch, J. H. Bardarson, and J. E. Moore, *Phys. Rev. Lett.* **108**, 227206 (2012).
- [26] C. Karrasch, J. Hauschild, S. Langer, and F. Heidrich-Meisner, *Phys. Rev. B* **87**, 245128 (2013).
- [27] T. Prosen, *Phys. Rev. Lett.* **107**, 137201 (2011).
- [28] M. Žnidarič, *Phys. Rev. Lett.* **106**, 220601 (2011).
- [29] E. Ilievski and T. Prosen, *Comm. Math. Phys.* **318**, 809 (2013).
- [30] D. Karevski, V. Popkov, and G. M. Schütz, *Phys. Rev. Lett.* **110**, 047201 (2013).
- [31] M. Jimbo and T. Miwa, *Algebraic Analysis of Solvable Lattice Models* (American Mathematical Society, Providence, 1994).
- [32] L. Mezincescu, R. I. Nepomechie, *Mod. Phys. Lett. A*, **6**, 2497 (1991).
- [33] V. Pasquier and H. Saleur, *Nucl. Phys. B* **330**, 523 (1990).
- [34] L. D. Faddeev, *Int. J. Mod. Phys. A* **10**, 1845 (1995).
- [35] T. Prosen and E. Ilievski, *Phys. Rev. Lett.* **111**, 057203 (2013).
- [36] T. Prosen, *Nucl. Phys. B* **886**, 1177 (2014).
- [37] A. Klümper and K. Sakai, *J. Phys. A* **35**, 2173 (2002).
- [38] K. Fabricius and B. M. McCoy, *Prog. in Math. Phys.* **23**, 119 (2001).
- [39] A. Lazarescu and V. Pasquier, *J. Phys. A: Math. Theor.* **47** 295202 (2014).
- [40] M. Marcus and H. Minc, *Introduction to Linear Algebra* (Dover, New York, 1988).