# A note on the rational cuspidal curves

Piotr Nayar, Barbara Pilat

#### Abstract

In this short note we give an elementary combinatorial argument, showing that the Conjecture of J. Fernández de Bobadilla, I. Luengo, A. Melle-Hernández, A. Némethi (see [BL], Conjecture 1.2) follows from Theorem 5.4 in [BL] in the case of rational cuspidal curves with two critical points.

**2010 Mathematics Subject Classification.** Primary 14H50; Secondary 14B05, 57M25, 57R58.

**Key words and phrases.** Rational cuspidal curve, Alexander polynomial, infimum convolution.

#### 1 Introduction

In this short note we deal with irreducible algebraic curves  $C \subset \mathbb{C}P^2$ . Such a curve has a finite set of singular points  $\{z_i\}_{i=1}^n$  such that a neighbourhood of each singular point intersects C in a cone on a link  $K_i \subset S^3$ . We would like to know what possible configurations of links  $\{K_i\}_{i=1}^n$  arise in this way. We consider only the case in which each  $K_i$  is connected (in this case  $K_i$  is a knot), and thus C is a rational curve, meaning that there is a rational surjective map  $\mathbb{C}P^1 \to C$ . Such a curve is called rational cuspidal. We refer to [M] for a survey on rational cuspidal curves.

Suppose that z is a cuspidal singular point of a curve C and B is a sufficiently small ball around z. Let  $\Psi(t)=(x(t),y(t))$  be a local parametrization of  $C\cap B$  near z. For any polynomial G(x,y) we look at the order at 0 of an analytic map  $t\mapsto G(x(t),y(t))\in C$ . Let S be the set of integers, which can be realized as the order for some G. Then S is a semigroup of  $\mathbb{Z}_{\geq 0}$ . We call it the semigroup of the singular point, see [W] for the details and proofs. The gap sequence,  $G=\mathbb{Z}_{\geq 0}\backslash S$ , has precisely  $\mu/2$  elements, where the largest one is  $\mu-1$ . Here  $\mu$  stands for the Milnor number. Assume that K is the

link of the singular point z. The Alexander polynomial of K is of the form

$$\Delta_K(t) = \sum_{i=0}^{2m} (-1)^i t^{n_i},$$

where  $(n_i)_{i=0}^{2m}$  form an increasing sequence with  $n_0 = 0$  and  $n_{2m} = 2g$ , for g being the genus of K. Writing  $t^{2n_i} - t^{2n_{i-1}} = (t-1)(t^{2n_i-1} + t^{2n_i-2} + \ldots + t^{2n_{i-1}})$  yields the representation

$$\Delta_K(t) = 1 + (t - 1) \sum_{j=1}^k t^{g_j},\tag{1}$$

for some finite sequence  $0 < g_1 < g_2 < \ldots < g_k$ . We have the following lemma (see [W], Exercise 5.7.7), which relates the Alexander polynomial to the gap sequence of a singular point.

**Lemma 1.** The sequence  $g_1, \ldots, g_k$  in (1) is the gap sequence of the semi-group of the singular point. In particular,  $k = |G| = \mu/2$ , where  $\mu$  is the Milnor number, so |G| is the genus.

If we write  $t^{g_j} = (t-1)(t^{g_j-1} + t^{g_j-2} + \ldots + 1) + 1$ , we obtain

$$\Delta_K(t) = 1 + (t-1)g(K) + (t-1)^2 \sum_{j=0}^{\mu-2} k_j t^j,$$

where  $k_j = |\{m > j : m \notin S\}|$ . This motivates the following definition.

**Definition.** For any finite increasing sequence of positive integers G we define

$$I_G(m) = |\{k \in G \cup \mathbb{Z}_{<0} : k \ge m\}|,$$

where  $\mathbb{Z}_{<0}$  is the set of negative integers. We shall call  $I_G$  the gap function, because in most applications G will be a gap sequence of some semigroup.

Clearly, for  $j = 0, 1, ..., \mu - 2$  we have  $I_G(j + 1) = k_j$ . In [FLMN] the following conjecture was proposed.

Conjecture 1. Suppose that the rational cuspidal curve C of degree d has critical points  $z_1, \ldots, z_n$ . Let  $K_1, \ldots, K_n$  be the corresponding links of singular points and let  $\Delta_1, \ldots, \Delta_n$  be their Alexander polynomials. Let g be the genus of K. Let  $\Delta = \Delta_1, \ldots, \Delta_n$ , expanded as

$$\Delta(t) = 1 + \frac{(d-1)(d-2)}{2}(t-1) + (t-1)^2 \sum_{j=0}^{2g-2} k_l$$

Then for any  $j = 0, \ldots, d-3$  we have  $k_{d(d-j-3)} \leq (j+1)(j+2)/2$ , with equality for n = 1.

This conjecture was verified in the case n=1 by Borodzik and Livingston, see [BL].

We define the infimum convolution of two functions.

**Definition.** Let  $I_1, I_2, \ldots, I_n : \mathbb{Z} \to \mathbb{Z}_{>0}$ . We define

$$(I_1 \diamond I_2 \dots \diamond I_n)(k) = \min_{\substack{k_1, k_2, \dots, k_n \in \mathbb{Z} \\ k_1 + k_2 + \dots + k_n = k}} (I_1(k_1) + I_2(k_2) + \dots + I_n(k_n)).$$

In [BL] the authors gave the proof of the following theorem.

**Theorem 1.** ([BL], Theorem 5.4) Let C be a rational cuspidal curve of degree d. Let  $I_1, \ldots, I_n$  be the gap functions associated to each singular point on C. Then for any  $j \in \{-1, 0, \ldots, d-2\}$  we have

$$I_1 \diamond I_2 \diamond \ldots \diamond I_n(jd+1) = \frac{1}{2}(j-d+1)(j-d+2).$$

Note that we have  $|G_1| + |G_2| + \ldots + |G_n| = \frac{(d-1)(d-2)}{2}$ . Therefore, one can give an equivalent reformulation of the Conjecture 1.

**Conjecture 2.** Suppose that the rational cuspidal curve C of degree d has critical points  $z_1, \ldots, z_n$ . Let  $K_1, \ldots, K_n$  be the corresponding links of singular points and let  $\Delta_1, \ldots, \Delta_n$  be their Alexander polynomials. Moreover, let  $G_1, G_2, \ldots, G_n$  be the gap sequences of these points. Let  $g = |G_1| + |G_2| + \ldots + |G_n|$  be the genus of K. Let  $\Delta = \Delta_1, \ldots, \Delta_n$ , expanded as

$$\Delta(t) = 1 + (t - 1)g + (t - 1)^2 \sum_{j=0}^{2g-2} k_l$$

and let  $I = I_1 \diamond I_2 \diamond ... \diamond I_n$ . Then for any j = 0, ..., d-3 we have  $k_{d(d-j-3)} \leq I(d(d-j-3)+1)$ , with equality for n = 1.

In this note we give an elementary argument, showing that FLMN conjecture follows from [BL] for n=2. The idea of our proof is to forget about the specific structure of the problem coming from theory of singularities and to prove Conjecture 2 for general sets  $G_1, G_2$ . Namely, we have the following theorem.

**Theorem 2.** Let G, H be two finite sets of positive integers and let  $I_G, I_H : \mathbb{Z} \to \mathbb{Z}_{\geq 0}$  be their gap functions. Let us define the polynomials

$$\begin{split} &\Delta_G(t) = 1 + (t-1) \sum_{\substack{j=1 \\ j=1}}^{|G|} t^{g_j} = 1 + (t-1)|G| + (t-1)^2 \sum_{\substack{j \geq 0 \\ j}} k_j^G t^j \\ &\Delta_H(t) = 1 + (t-1) \sum_{\substack{j=1 \\ j=1}}^{|H|} t^{h_j} = 1 + (t-1)|H| + (t-1)^2 \sum_{\substack{j \geq 0 \\ j}} k_j^H t^j \end{split},$$

where  $k_j^G = I_G(j+1), k_j^H = I_H(j+1), j \ge 0$ . Take  $\Delta = \Delta_G \cdot \Delta_H$  and  $I = I_G \diamond I_H$ . Then

$$\Delta(t) = 1 + (t-1)(|G| + |H|) + (t-1)^2 \sum_{j>0} k_j t^j,$$

where  $k_j \leq I(j+1)$  for  $j \geq 0$ .

This gives the proof of Conjecture 1 in the case n=2.

It is natural to ask whether the above theorem is valid for arbitrary  $n \geq 2$ . Recently, after we found our elementary combinatorial argument for n=2, J. Bodnár and A. Némethi showed that the Conjecture 1 is false for  $n\geq 3$ , see [BN]. They also found yet another proof of Conjecture 1 in the case of two singularities.

### 2 Proof of the main result

In this section we give a proof of our main result.

*Proof.* Our goal is to express the numbers  $k_j$  in terms of the numbers  $k_j^G$  and  $k_j^H$ . We have

$$\Delta(t) = \Delta_G(t)\Delta_H(t) = 1 + (t-1)(|G| + |H|)$$

$$+ (t-1)^2 \Big[ |G| \cdot |H| + \sum_{j \ge 0} (k_j^G + k_j^H)t^j + (t-1)(|G| \sum_{j \ge 0} k_j^H t^j + |H| \sum_{j \ge 0} k_j^G t^j)$$

$$+ (t-1)^2 \Big( \sum_{j \ge 0} k_j^G t^j \Big) \Big( \sum_{j \ge 0} k_j^H t^j \Big) \Big] = 1 + (t-1)(|G| + |H|) + (t-1)^2 \Theta(t),$$

with

$$\Theta(t) = |G| \cdot |H| + k_0^G (1 - |H|) + k_0^H (1 - |G|) + k_0^G k_0^H + \sum_{j \ge 1} t^j k_j,$$

where

$$k_j = k_j^G(1 - |H|) + |H|k_{j-1}^G + k_j^H(1 - |G|) + |G|k_{j-1}^H + l_j$$

and

$$l_j = \sum_{u+v=j, \ u,v \ge 0} k_u^G k_v^H - 2 \sum_{u+v=j-1, \ u,v \ge 0} k_u^G k_v^H + \sum_{u+v=j-2, \ u,v \ge 0} k_u^G k_v^H.$$

Note that  $k_0^G = |G|$  and  $k_0^H = |H|$ . Therefore,

$$k_0 = |G| \cdot |H| + k_0^G (1 - |H|) + k_0^H (1 - |G|) + k_0^G k_0^H$$
  
= |G| \cdot |H| + |G|(1 - |H|) + |H|(1 - |G|) + |G| \cdot |H| = |G| + |H|.

Moreover, for  $k \geq 1$  we have

$$I_H(k) \ge |H| - (k-1), \qquad I_G(1-k) = |G| + (k-1).$$

Thus, for  $k \geq 1$  we obtain

$$I_G(1-k) + I_H(k) \ge |G| + |H|.$$

For  $k \leq -1$ 

$$I_G(1-k) \ge |G| + k, \qquad I_H(k) = |H| - k.$$

In this case we obtain

$$I_G(1-k) + I_H(k) \ge |G| + |H|$$

and we arrive at

$$I(1) = \min_{k \in \mathbb{Z}} (I_G(1-k) + I_H(k))$$
  
=  $I_G(1) + I_H(0) = I_G(0) + I_H(1) = |G| + |H| = k_0.$ 

Note that

$$\begin{split} l_{j} &= \sum_{u+v=j,\ u,v\geq 0} k_{u}^{G} k_{v}^{H} - \sum_{u+v=j,\ u\geq 0,v\geq 1} k_{u}^{G} k_{v-1}^{H} - \sum_{u+v=j,\ u\geq 1,v\geq 0} k_{u-1}^{G} k_{v}^{H} \\ &+ \sum_{u+v=j,\ u,v\geq 1} k_{u-1}^{G} k_{v-1}^{H} = \sum_{u+v=j,u,v\geq 1} (k_{u}^{G} - k_{u-1}^{G})(k_{v}^{H} - k_{v-1}^{H}) \\ &+ k_{0}^{G} k_{j}^{H} + k_{j}^{G} k_{0}^{H} - k_{0}^{G} k_{j-1}^{H} - k_{j-1}^{G} k_{0}^{H}. \end{split}$$

Thus,

$$\begin{split} k_j &= \sum_{u+v=j, u, v \geq 1} (k_u^G - k_{u-1}^G) (k_v^H - k_{v-1}^H) \\ &+ \Big( (k_0^G k_j^H + k_j^G k_0^H - k_0^G k_{j-1}^H - k_{j-1}^G k_0^H) \\ &+ (k_j^G (1 - |H|) + |H| k_{j-1}^G + k_j^H (1 - |G|) + |G| k_{j-1}^H) \Big). \end{split}$$

Observe that we have a miracle.

$$\begin{split} k_0^G k_j^H + k_j^G k_0^H - k_0^G k_{j-1}^H - k_{j-1}^G k_0^H \\ + k_j^G (1 - |H|) + |H| k_{j-1}^G + k_j^H (1 - |G|) + |G| k_{j-1}^H \\ = |G| k_j^H + k_j^G |H| - |G| k_{j-1}^H - k_{j-1}^G |H| \\ + k_j^G (1 - |H|) + |H| k_{j-1}^G + k_j^H (1 - |G|) + |G| k_{j-1}^H \\ = k_j^G + k_j^H. \end{split}$$

We get

$$k_j = k_j^G + k_j^H + \sum_{u+v=j, u, v \ge 1} (k_{u-1}^G - k_u^G)(k_{v-1}^H - k_v^H).$$

We are to prove that  $k_j \leq (I_G \diamond I_H)(j+1)$ . It suffices to prove that  $k_j \leq I_G(j+1-l) + I_H(l)$  for every  $l \in \mathbb{Z}$ . Thus, we have to deal with the inequality

$$k_j^G + k_j^H + \sum_{u+v=j,u,v>1} (k_u^G - k_{u-1}^G)(k_v^H - k_{v-1}^H) \le I_G(j+1-l) + I_H(l), \quad j \ge 1, l \in \mathbb{Z}.$$

Note that if u + v = j then we have either  $u \ge j - l + 1$  or  $v \ge l$ . Thus,

$$1_{u \in G} 1_{v \in H} 1_{u+v=j} \le 1_{u \in G \cap [i-l+1,j]} + 1_{v \in H \cap [l,j]}$$
.

In the above expression we have used the convention  $[a, b] = \emptyset$  for a > b. We obtain

$$\begin{split} \sum_{u+v=j,u,v\geq 1} (k_{u-1}^G - k_u^G)(k_{v-1}^H - k_v^H) &= \sum_{u+v=j,u,v\geq 0} (k_{u-1}^G - k_u^G)(k_{v-1}^H - k_v^H) \\ &= \sum_{u+v=j,u,v\geq 0} \mathbf{1}_{u\in G} \mathbf{1}_{v\in H} \leq \sum_{u+v=j,u,v\geq 0} \left( \mathbf{1}_{u\in G\cap[j-l+1,j]} + \mathbf{1}_{v\in H\cap[l,j]} \right) \\ &= (k_{j-l}^G - k_j^G) + (k_{l-1}^H - k_j^H), \end{split}$$

what finishes our proof.

## Acknowledgements

We are grateful to Maciej Borodzik for his valuable comments about the meaning of our result.

#### References

- [FLMN] J. Fernández de Bobadilla, I. Luengo, A. Melle-Hernández, A. Némethi, Classification of rational unicuspidal projective curves whose singularities have one Puiseux pair, Proceedings of Sao Carlos Workshop 2004 Real and Complex Singularities, Series Trends in Mathematics, Birkhäuser 2007, 31–46.
- [BL] M. Borodzik, C. Livingston, Heegaard Floer homology and rational cuspidal curves, arXiv:1304.1062
- [BN] J. Bodnár and A. Némethi, arXiv:1405.0437
- [M] K. Moe, Rational cuspidal curves, Master Thesis, University of Oslo 2008, permanent link at University of Oslo: https://www.duo.uio.no/handle/123456789/10759
- [W] C. T. C. Wall, Singular Points of Plane Curves, London Mathematical Society Student Texts, 63. Cambridge University Press, Cambridge, 2004.

Piotr Nayar, nayar@mimuw.edu.pl Institute of Mathematics, University of Warsaw, Banacha 2, 02-097 Warszawa, Poland.

Barbara Pilat, B.Pilat@mini.pw.edu.pl Faculty of Mathematics and Information Science, Warsaw University of Technology, Koszykowa 75, 00-662 Warszawa, Poland.