INEQUALITIES FOR SUMS OF RANDOM VARIABLES IN NONCOMMUTATIVE PROBABILITY SPACES

GHADIR SADEGHI¹ AND MOHAMMAD SAL MOSLEHIAN²

ABSTRACT. We extend the classical Hoeffding inequality to the noncommutative setting. Under certain conditions we establish an improved version of the noncommutative Bennett inequality and use it to prove an extension of Rosenthal inequality as follows: Let (\mathfrak{M}, τ) be a noncommutative probability space, \mathfrak{N} be a von Neumann subalgebra of \mathfrak{M} with the corresponding conditional expectation $\mathcal{E}_{\mathfrak{N}}$ and let subalgebras $\mathfrak{N} \subseteq \mathfrak{A}_j \subseteq \mathfrak{M}$ $(j = 1, \dots, n)$ be successively independent over \mathfrak{N} . If $1 \leq r \leq 2 \leq p < \infty$ and $x_j \in \mathfrak{A}_j$ are self-adjoint elements such that $\mathcal{E}_{\mathfrak{N}}(x_j) = 0$ for all $1 \leq j \leq n$, then

$$\left\|\sum_{j=1}^{n} x_{j}\right\|_{p}^{p} \leq C(p,r) \left\{\sum_{j=1}^{n} \|x_{j}\|_{p}^{p} + \left(\sum_{j=1}^{n} \|x_{j}^{r}\|\right)^{\frac{p}{r}}\right\}.$$

1. INTRODUCTION

By a noncommutative probability space (\mathfrak{M}, τ) we mean a von Neumann algebra \mathfrak{M} on a Hilbert space \mathfrak{H} with unit element 1 equipped with a faithful normal finite trace τ such that $\tau(1) = 1$. The modules |x| of $x \in \mathfrak{M}$ is defined by continuous functional calculus as $|x| = (x^*x)^{1/2}$.

For each self-adjoint operator $x \in \mathfrak{M}$, there exists a unique spectral measure E as a σ additive mapping with respect to the strong operator topology from the Borel σ -algebra $\mathcal{B}(\mathbb{R})$ of \mathbb{R} into the set of all orthogonal projections such that for every Borel function $f : \sigma(x) \to \mathbb{C}$ the operator f(x) is defined by $f(x) = \int f(\lambda) dE(\lambda)$, in particular, $x = \int \lambda dE(\lambda)$ and $\chi_B(x) = \int_B dE(\lambda) = E(B)$. In addition,

$$\tau\left(\chi_{[t,\infty)}\left(|x|\right)\right) = \tau\left(\chi_{[t,\infty)}\left(x\right)\right) + \tau\left(\chi_{[t,\infty)}\left(-x\right)\right) \,. \tag{1.1}$$

Further, if $x \ge 0$ and t > 0, then $\chi_{[t,\infty)}(x) t \le x$. Hence we get the inequality

$$\tau(\chi_{[t,\infty)}(x)) \le t^{-1}\tau(x) \,,$$

²⁰¹⁰ Mathematics Subject Classification. Primary 46L53, 47A30; Secondary 60B20,60B11.

Key words and phrases. Noncommutative probability space; von Neumann algebra; trace; noncommutative Rosenthal inequality; noncommutative Bennett inequality; noncommutative Hoeffding inequality.

which is called the Markov inequality in the literature. For a self-adjoint element $x \in \mathfrak{M}$, it follows from the Markov inequality that

$$\tau(\chi_{[t,\infty)}(x)) = \tau(\chi_{[e^t,\infty)}(e^x)) \le e^{-t}\tau(e^x),$$

from where we reach the exponential Chebyshev inequality as follows:

$$\tau(\chi_{[t,\infty)}(x)) \le e^{-t}\tau(e^x).$$
(1.2)

As in the commutative context, we use the notation $\operatorname{Prob}(x \ge t) := \tau(\chi_{[t,\infty)}(x)).$

For any Borel set $A \subseteq \mathbb{R}$, we define $\nu(A) = \tau(E(A))$. Then ν is a scalar-valued spectral measure for x and $\nu(\mathbb{R}) = 1$. In addition,

$$\tau(f(x)) = \tau\left(\int f(\lambda)dE(\lambda)\right) = \int f(\lambda)d\tau\left(E(\lambda)\right) = \int f(\lambda)d\nu(\lambda).$$
(1.3)

By the measurable functional calculus [4] there exists a *-homomorphism $\pi : L^{\infty}(\nu) \to \mathfrak{M}$ depending on x such that $\pi(f) = f(x)$ for all $f \in L^{\infty}(\nu)$.

For $1 \leq p < \infty$, $L^p(\mathfrak{M}, \tau)$ is defined as the completion of \mathfrak{M} with respect to the norm $||x||_p = \tau(|x|^p)^{\frac{1}{p}}$. The commutative cases of discussed spaces are usual L^p -spaces and the Schatten *p*-classes \mathcal{C}_p . For further information we refer the reader to [18, 13, 11, 16]. Let $x \in \mathfrak{M}$ be positive. For $p \geq 1$ and positive $x \in \mathfrak{M}$, from (1.3), we have

$$\|x\|_{p}^{p} = \tau(x^{p}) = \int_{0}^{\infty} \lambda^{p} d\nu(\lambda) = \int_{0}^{\infty} pt^{p-1}\nu([t,\infty))dt = \int_{0}^{\infty} pt^{p-1}\tau(\chi_{[t,\infty)}(x))dt.$$

Let \mathcal{P} be the lattice of projections of \mathfrak{M} . Set $p^{\perp} = 1 - p$ for $p \in \mathcal{P}$. Given a family of projections $(p_{\lambda})_{\lambda \in \Lambda} \subseteq \mathcal{P}$, we denote by $\vee_{\lambda \in \Lambda} p_{\lambda}$ (resp., $\wedge_{\lambda \in \Lambda} p_{\lambda}$) the projection from \mathfrak{H} onto the closed subspace generated by $p_{\lambda}(\mathfrak{H})$ (resp. onto the subspace $\cap_{\lambda \in \Lambda} p(\mathfrak{H})$). Consequently, $(\vee_{\lambda \in \Lambda} p_{\lambda})^{\perp} = \wedge_{\lambda \in \Lambda} p_{\lambda}^{\perp}$. Two projections p and q are said to be equivalent if there exists a partial isometry $u \in \mathfrak{M}$ such that $u^*u = p$ and $uu^* = q$. In this case, we write $p \sim q$. If p is equivalent to a projection $q_1 \leq q$, we write $p \prec q$. We need the following elementary properties of projections (see [12]).

Lemma 1.1. Let p and q be two projection of \mathfrak{M} . Then

(i) $p \lor q - q \sim p - p \land q$. (ii) If $p \land q = 0$ then $p \prec q^{\perp}$ (iii) If p and q are equivalent projections in \mathfrak{M} then $\tau(p) = \tau(q)$. (iv) If $(p_{\lambda})_{\lambda \in \Lambda}$ is a family of projections in \mathfrak{M} then $\tau(\vee_{\lambda \in \Lambda} p_{\lambda}) \leq \sum_{\lambda \in \Lambda} \tau(p_{\lambda})$.

Let \mathfrak{N} be a von Neumann subalgebra of \mathfrak{M} . Then there exists a map $\mathcal{E}_{\mathfrak{N}} : \mathfrak{M} \to \mathfrak{N}$ satisfying the following properties:

- (i) $\mathcal{E}_{\mathfrak{N}}$ is a normal contraction positive map projecting \mathfrak{M} onto \mathfrak{N} ;
- (ii) $\mathcal{E}_{\mathfrak{N}}(axb) = a\mathcal{E}_{\mathfrak{N}}(x)b$ for any $x \in \mathfrak{M}$ and $a, b \in \mathfrak{N}$;
- (iii) $\tau \circ \mathcal{E}_{\mathfrak{N}} = \tau$.

Moreover, $\mathcal{E}_{\mathfrak{N}}$ is the unique map satisfying (i) and (ii). The map $\mathcal{E}_{\mathfrak{N}}$ is called the conditional expectation of \mathfrak{M} with respect to \mathfrak{N} . We say that two subalgebras $\mathfrak{N} \subseteq \mathfrak{A}, \mathfrak{B} \subseteq \mathfrak{M}$ are independent over \mathfrak{N} if $\mathcal{E}_{\mathfrak{N}}(xy) = \mathcal{E}_{\mathfrak{N}}(x)\mathcal{E}_{\mathfrak{N}}(y)$ for all $x \in \mathfrak{A}, y \in \mathfrak{B}$. In particular, two random variable X and Y of a commutative von Neumann algebra $L^{\infty}(\mu)$ in which μ is a probability measure are independent if the algebras they generate are independent over the complex field \mathbb{C} . A sequence of subalgebras $\mathfrak{N} \subseteq \mathfrak{A}_1, \ldots, \mathfrak{A}_n \subseteq \mathfrak{M}$ is called successively independent over \mathfrak{N} if \mathfrak{A}_{k+1} is independent of the algebra $\mathfrak{M}(k)$ generated by $\mathfrak{A}_1, \ldots, \mathfrak{A}_k$. For further information the reader is referred to [19, 20].

In 1970, Rosenthal [14] presented an inequality to describe isomorphic types of some subspaces in L_p -spaces. It indeed gives a bound for the *p*-norm of independent mean 0 random variables. More precisely, it says that for any $p \ge 2$ there exists a constant c(p) such that for any $n \in \mathbb{N}$ and any independent mean 0 random variables f_1, \dots, f_n it holds that

$$\mathbb{E}\left|\sum_{k=1}^{n} f_{k}\right|^{p} \leq c(p)\left(\left(\sum_{k=1}^{n} \mathbb{E}|f_{k}|^{2}\right)^{p/2} + \sum_{k=1}^{n} \mathbb{E}|f_{k}|^{p}\right).$$
(1.4)

Burkholder [3] generalized Rosenthal's inequality in the context of martingales. Since then this inequality has been generalized and applied by many mathematicians; see, e.g., [8, 10] and references therein. Recently, Junge and Zeng [9] extended the Bennett and Bernstein inequalities to the noncommutative setting and derive an improved version of the Rosenthal inequality from Bernstein's inequality by using the properties of Gamma function. In probability theory, the Bennett inequality (Bernstein inequality, resp.) gives an upper bound on the probability that the sum of independent random variables deviates from its expected value (deviates from its mean, resp.) by more than a fixed amount, see [1].

In this paper, we establish an extension of the noncommutative Bennett inequality due to Junge and Zeng [9] and use it together some noncommutative techniques to prove the Rosenthal inequality with a parameter $1 \le r \le 2$. We also prove a noncommutative Hoeffding inequality as an extension of the Bernstein inequality. The Hoeffding inequality [6] gives a probability bound for the deviation between the average of n independent bounded random variables and its mean (see Corollary 3.4). There have been several generalizations and applications of this significant inequality, see [2, 17].

GH. SADEGHI, M.S. MOSLEHIAN

2. Bennet inequality

We provide an improved version of the noncommutative Bennett inequality based on the arguments of [9, Theorem 01].

Theorem 2.1. (Noncommutative Bennett inequality) Let $\mathfrak{N} \subseteq \mathfrak{A}_j \subseteq \mathfrak{M}$ be successively independent over \mathfrak{N} and $x_j \in \mathfrak{A}_j$ be self-adjoint and $1 \leq r \leq 2$ such that

- $\mathcal{E}_{\mathfrak{N}}(x_j) \leq 0$,
- $\mathcal{E}_{\mathfrak{N}}(|x_j|^r) \leq b_j^r$,
- $||x_j|| \leq M$,

for some M > 0 and all $1 \le j \le n$. Then for each $t \ge 0$,

$$\operatorname{Prob}\left(\sum_{j=1}^{n} x_{j} \ge t\right) = \tau\left(\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)\right) \le \exp\left(\frac{-b}{M^{r}}\Phi\left(\frac{tM^{r-1}}{b}\right)\right), \quad (2.1)$$

where $\Phi(\alpha) = (1+\alpha)\log(1+\alpha) - \alpha$ and $b = \sum_{j=1}^{n} b_{j}^{r}.$

Proof. Let $\lambda \geq 0$. We have

$$\begin{split} \mathcal{E}_{\mathfrak{N}}\left(e^{\lambda x_{n}}\right) &= \mathcal{E}_{\mathfrak{N}}\left(\sum_{k=0}^{\infty} \frac{(\lambda x_{n})^{k}}{k!}\right) = \sum_{k=0}^{\infty} \frac{\lambda^{k}}{k!} \mathcal{E}_{\mathfrak{N}}(x_{n}^{k}) \\ &= 1 + \mathcal{E}_{\mathfrak{N}}(x_{n}) + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \mathcal{E}_{\mathfrak{N}}(x_{n}^{k}) \\ &\leq 1 + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \mathcal{E}_{\mathfrak{N}}(x_{n}^{k}) \qquad (\text{by } \mathcal{E}_{\mathfrak{N}}(x_{n}) \leq 0) \\ &\leq 1 + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \mathcal{E}_{\mathfrak{N}}(|x_{n}|^{k}) \qquad (\text{by } x_{n}^{k} \leq |x_{n}|^{k}) \\ &= 1 + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \mathcal{E}_{\mathfrak{N}}(|x_{n}|^{r}|x_{n}|^{k-r}) \\ &\leq 1 + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \|x_{n}\|^{k-r} \mathcal{E}_{\mathfrak{N}}(|x_{n}|^{r}) \qquad (\text{by } |x_{n}|^{r}|x_{n}|^{k-r} \leq \|x_{n}\|^{k-r}|x_{n}|^{r}) \\ &\leq 1 + \sum_{k=2}^{\infty} \frac{\lambda^{k}}{k!} \|x_{n}\|^{k-r} \mathcal{E}_{\mathfrak{N}}(|x_{n}|^{r}) \\ &= 1 + \frac{b_{n}^{2}}{|x_{n}||^{r}} (e^{\lambda \|x_{n}\|} - 1 - \lambda \|x_{n}\|) \\ &\leq \exp\left(\frac{b_{n}^{r}}{\|x_{n}\|^{r}} (e^{\lambda \|x_{n}\|} - 1 - \lambda \|x_{n}\|)\right). \end{split}$$

Note that the function $f(s) := \exp\left(\frac{e^{\lambda s} - 1 - \lambda s}{s^r}\right)$ is increasing for s > 0. It follows that

$$\mathcal{E}_{\mathfrak{N}}\left(e^{\lambda x_{n}}\right) \leq \exp\left(\frac{b_{n}^{r}}{M^{r}}\left(e^{\lambda M}-1-\lambda M\right)\right).$$
(2.2)

It follows from (1.2) that

$$\tau\left(\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)\right) = \tau\left(\chi_{[\lambda t,\infty)}\left(\sum_{j=1}^{n} \lambda x_{j}\right)\right) \leq \exp(-\lambda t)\tau\left(e^{\sum_{j=1}^{n} \lambda x_{j}}\right).$$
(2.3)

Recall that the Golden– Thompson inequality [15] stating that for all self-adjoint elements $z_1, z_2 \in \mathfrak{M}$,

$$\tau(e^{z_1+z_2}) \le \tau(e^{z_1}e^{z_2}). \tag{2.4}$$

Hence

$$\tau\left(e^{\sum_{j=1}^{n}\lambda x_{j}}\right) \leq \tau\left(e^{\sum_{j=1}^{n-1}\lambda x_{j}}e^{\lambda x_{n}}\right) \qquad (by (2.4))$$

$$= \tau\left(\mathcal{E}_{\mathfrak{N}}\left(e^{\sum_{j=1}^{n-1}\lambda x_{j}}e^{\lambda x_{n}}\right)\right)$$

$$= \tau\left(\mathcal{E}_{\mathfrak{N}}\left(e^{\sum_{j=1}^{n-1}\lambda x_{j}}\right)\mathcal{E}_{\mathfrak{N}}\left(e^{\lambda x_{n}}\right)\right)$$

$$\leq \exp\left(\frac{b_{n}^{r}}{M^{r}}\left(e^{\lambda M}-1-\lambda M\right)\right)\tau\left(\mathcal{E}_{\mathfrak{N}}\left(e^{\sum_{j=1}^{n-1}\lambda x_{j}}\right)\right)$$

$$(by (2.2) \text{ and traciality of } \tau)$$

$$\leq \exp\left(\frac{\sum_{j=1}^{n}b_{j}^{r}}{M^{r}}\left(e^{\lambda M}-1-\lambda M\right)\right). \qquad (by \text{ iterating } n-2 \text{ times})$$

We infer from the latter inequality together with (2.3) that

$$\tau\left(\chi_{[t,\infty)}\left(\sum_{j=1}^n x_j\right)\right) \le \exp\left(-\lambda t + \frac{\sum_{j=1}^n b_j^r}{M^r}(e^{\lambda M} - 1 - \lambda M)\right).$$

By basic calculus method we find the minimizing value

$$\lambda = \frac{1}{M} \log \left(1 + \frac{tM^{r-1}}{\sum_{j=1}^{n} b_j^r} \right),$$

which yields (3.1).

Since both inequalities $\Phi(t) \ge \frac{t^2}{2+2t/3}$ and $\Phi(t) \ge \frac{t}{2}\operatorname{arcsinh}(\frac{t}{2})$ are valid for all $t \ge 0$, so one can get the Bernstein and Prohorov inequalities from the Bennett's inequality as follows.

Corollary 2.2. Under the same hypothesis of Theorem 2.1,

$$\tau\left(\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)\right) \leq \exp\left(-\frac{t^{2}M^{r-2}}{2b + (2/3)tM^{r-1}}\right)$$
(2.5)

and

$$au\left(\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)\right) \leq \exp\left(-\frac{t}{2M}\operatorname{arcsinh}\left(\frac{tM}{2b}\right)\right)$$

We can immediately deduce the following commutative Bernstein's Inequality.

Corollary 2.3. Let X_1, \dots, X_n be independent Bernoulli random variables taking values 1 and -1 with probability 1/2. Then

$$\operatorname{Prob}\left(\left|\frac{1}{n}\sum_{i=1}^{n}X_{i}\right| \geq \varepsilon\right) \leq 2e^{\frac{-n\varepsilon^{2}}{2+(2\varepsilon/3)}}.$$

3. Hoeffding inequality

In this section we provide a noncommutative version of Hoeffding's inequality and present some consequences.

Theorem 3.1. (Noncommutative Hoeffding inequality) Let $\mathfrak{N} \subseteq \mathfrak{A}_j \subseteq \mathfrak{M}$ be successively independent over \mathfrak{N} and let $x_j \in \mathfrak{A}_j$ be self-adjoint such that $a_j \leq x_j \leq b_j$ for some real numbers $a_j < b_j$ and $\mathcal{E}_{\mathfrak{N}}(x_j) = \mu$ for some $\mu \geq 0$ and all $1 \leq j \leq n$. Then

$$\operatorname{Prob}\left(|S_n - n\mu| \ge t\right) \le 2 \exp\left\{\frac{-2t^2}{\sum_{j=1}^n (b_j - a_j)^2}\right\}.$$
(3.1)

for any t > 0, where $S_n = \sum_{j=1}^n x_j$.

Proof. First we show that if $x \in \mathfrak{A}_j$ such that $a \leq x \leq b$ and $\mathcal{E}_{\mathfrak{N}}(x) = 0$, then

$$\mathcal{E}_{\mathfrak{N}}(e^{sx}) \le \exp\left\{\frac{s^2(b-a)^2}{8}\right\}$$
(3.2)

for any s > 0.

Let s > 0. Note that $t \mapsto e^{ts}$ is convex, therefore for any $a \le \alpha \le b$,

$$e^{s\alpha} \le e^{sb} \frac{\alpha - a}{b - a} + e^{sa} \frac{b - \alpha}{b - a}.$$

By the functional calculus we have

$$e^{sx} \le e^{sb}\frac{x-a}{b-a} + e^{sa}\frac{b-x}{b-a}.$$

Since $\mathcal{E}_{\mathfrak{N}}$ is a positive map and $\mathcal{E}_{\mathfrak{N}}(x) = 0$, we reach

$$\mathcal{E}_{\mathfrak{N}}(e^{sx}) \leq \frac{-a}{b-a}e^{sb} + \frac{b}{b-a}e^{sa} = e^{h(\alpha)},$$

where $\alpha = s(b-a)$, $h(\alpha) = -\gamma\alpha + \log(1-\gamma+\gamma e^{\alpha})$ and $\gamma = -a/(b-a)$. In addition, h(0) = h'(0) = 0 and $h''(\alpha) \leq \frac{1}{4}$ for all $\alpha > 0$. By Taylor's Theorem there exists a real number $\xi \in (0, \alpha)$ such that

$$h(\alpha) = h(0) + \alpha h'(0) + \frac{\alpha^2}{2}h''(\xi) \le \frac{\alpha^2}{8} = \frac{s^2(b-a)^2}{8}$$

Hence

$$\mathcal{E}_{\mathfrak{N}}(e^{sx}) \le \exp\left\{\frac{s^2(b-a)^2}{8}\right\}.$$

Second, for arbitrary value of $\mathcal{E}_{\mathfrak{N}}(x)$, setting $y := x - \mu$ we get $a - \mu \leq y \leq b - \mu$ and $\mathcal{E}_{\mathfrak{N}}(y) = 0$. Employing (3.2) we reach

$$\mathcal{E}_{\mathfrak{N}}\left(e^{s(x-\mu)}\right) \le \exp\left\{\frac{s^2(b-a)^2}{8}\right\}.$$
(3.3)

Next, by the same reasoning as in the proof of Theorem 2.1, we get from (3.3) that

$$\tau\left(e^{\lambda\sum_{j=1}^{n}(x_{j}-\mu)}\right) \leq \tau\left(\mathcal{E}_{\mathfrak{N}}\left(e^{\lambda\sum_{j=1}^{n-1}(x_{j}-\mu)}\right)\mathcal{E}_{\mathfrak{N}}\left(e^{\lambda(x_{n}-\mu)}\right)\right)$$
$$\leq e^{\frac{\lambda^{2}(b_{n}-a_{n})^{2}}{8}}\tau\left(\mathcal{E}_{\mathfrak{N}}\left(e^{\lambda\sum_{j=1}^{n-1}(x_{j}-\mu)}\right)\right)\leq\cdots$$
$$\leq \exp\left\{\frac{\lambda^{2}\sum_{j=1}^{n}(b_{j}-a_{j})^{2}}{8}\right\}.$$

Therefore (1.1) and the exponential Chebyshev inequality (1.2) yield that

$$\operatorname{Prob}(|S_n - n\mu| \ge t) = 2\operatorname{Prob}(S_n - n\mu \ge t) \le 2e^{-\lambda t} \exp\left\{\frac{\lambda^2 \sum_{j=1}^n (b_j - a_j)^2}{8}\right\}$$

This is minimized when $\lambda = \frac{4t}{\sum_{j=1}^{n} (b_j - a_j)^2}$. Thus

Prob
$$(|S_n - n\mu| \le 2 \exp\left\{\frac{-2t^2}{\sum_{j=1}^n (b_j - a_j)^2}\right\}$$
,

which is the desired inequality.

Remark 3.2. Under the same hypothesis of Theorem 3.1 we can show that the bound in the Hoeffding inequality is sharper than that in the Bernstein inequality (2.5) for r = 2. Let us assume that We may assume that $\mathcal{E}_{\mathfrak{N}}(x_j) = 0$ and $-1 \leq x_j \leq 1$. By the functional calculus

 $|x_j| \leq 1$ so $\mathcal{E}_{\mathfrak{N}}(|x_j|^2) \leq 1$ and $||x_j|| \leq 1$. So b = n and M = 1 in the notation of Theorem 2.1. Then Heoffding inequality gives rise to

$$\operatorname{Prob}\left(\sum_{j=1}^{n} x_j \ge t\right) \le \exp\left\{\frac{-t^2}{2n}\right\} \,.$$

and from Bernstein inequality we have

$$\operatorname{Prob}\left(\sum_{j=1}^{n} x_j \ge t\right) \le \exp\left\{\frac{-t^2}{2n + (2t/3)}\right\}.$$

The next result is the classical (commutative) version of the Hoeffding inequality.

Corollary 3.3 (Hoeffding's Inequality [6]). Let $a \leq X_1, \dots, X_n \leq b$ be independent random variables with the expectation $\mathbb{E}(X_i) = \mu$ $(i = 1, \dots, n)$. If $\overline{X}_n = (\sum_{i=1}^n X_i) / n$, then

$$\operatorname{Prob}(\left|\overline{X}_n - \mu\right| \ge t) \le 2e^{-2nt^2/(b-a)^2}.$$

In the special case, we immediately get

Corollary 3.4. Let $0 \le X_1, \dots, X_n \le 1$ be independent random variables with common mean μ . Then with probability at least $1 - \varepsilon$,

$$\left|\frac{1}{n}\sum_{i=1}^{n} X_i - \mu\right| \le \sqrt{\frac{\log(2/\varepsilon)}{2n}}.$$

4. Rosenthal inequality

In this section we intend to extend the Rosenthal inequality by using the noncommutative Bennet inequality with a parameter $1 \le r \le 2$. Our approach is simpler than that of [7] for the case of usual random variables.

Theorem 4.1. Let $1 \leq r \leq 2 \leq p < \infty$, $\mathfrak{N} \subseteq \mathfrak{A}_j \subseteq \mathfrak{M}$ be successively independent over \mathfrak{N} and $x_j \in \mathfrak{A}_j$ be self-adjoint such that $\mathcal{E}_{\mathfrak{N}}(x_j) = 0$. Then there exists a constant C(p, r) such that

$$\left\|\sum_{j=1}^{n} x_{j}\right\|_{p}^{p} \leq C(p,r) \left\{\sum_{j=1}^{n} \|x_{j}\|_{p}^{p} + \left(\sum_{j=1}^{n} \|x_{j}^{r}\|\right)^{\frac{p}{r}}\right\}.$$

Proof. We use the noncommutative Bennett inequality, but for this end, we replace $\Phi(\alpha)$ by $\alpha \log(1 + \alpha) - \alpha$, which is clearly smaller than $\Phi(\alpha)$ for any $\alpha \ge 0$.

Let us fix an arbitrary number $s \geq 0$ and consider $y_j = x_j \chi_{(-\infty,s]}(x_j) \in \mathfrak{A}_j$. It follows from $y_j \leq x_j$ and the positivity of $\mathcal{E}_{\mathfrak{N}}$ that $\mathcal{E}_{\mathfrak{N}}(y_j) \leq \mathcal{E}_{\mathfrak{N}}(x_j) = 0$. In addition, $\mathcal{E}_{\mathfrak{N}}$ is norm decreasing, so

$$\sum_{j=1}^{n} \mathcal{E}_{\mathfrak{N}}(|x_{j}|^{r}) \leq \sum_{j=1}^{n} ||x_{j}^{r}|| := B \quad \text{and} \quad b := \sum_{j=1}^{n} ||y_{j}^{r}|| \geq \sum_{j=1}^{n} \mathcal{E}_{\mathfrak{N}}(|y_{j}|^{r}).$$

Further, $b \leq B$ since $||y_j|| \leq ||x_j|| \leq M$, where $M := \max_{1 \leq j \leq n} ||x_j||$. It follows from the noncommutative Bennett inequality (3.1) that

$$\tau\left(\chi_{[t,\infty)}\sum_{j=1}^{n}y_{j}\right) \leq \exp\left\{\frac{-b}{M^{r}}\Phi\left(\frac{tM^{r-1}}{b}\right)\right\}$$

$$\leq \exp\left\{\frac{-b}{M^{r}}\left[\frac{tM^{r-1}}{b}\log\left(1+\frac{tM^{r-1}}{b}\right)-\frac{tM^{r-1}}{b}\right]\right\}$$

$$\leq \exp\left\{-\frac{t}{M}\left(\log\left(1+\frac{tM^{r-1}}{B}\right)-1\right)\right\}$$

$$(4.1)$$

for all t > 0. We have

$$\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_j\right) \prec \chi_{[t,\infty)}\left(\sum_{j=1}^{n} y_j\right) \lor \left(\lor_{j=1}^{n} \chi_{[s,\infty)}(x_j)\right)$$
(4.2)

for all t > 0. To show this we have to prove that

$$\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right) \wedge \left(\chi_{[0,t)}\left(\sum_{j=1}^{n} y_{j}\right) \wedge \left(\wedge_{j=1}^{n} \chi_{(-\infty,s)}(x_{j})\right)\right) = 0.$$

Let $\xi \in \chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)(\mathfrak{H}) \cap \left(\chi_{[0,t)}\left(\sum_{j=1}^{n} y_{j}\right)(\mathfrak{H}) \cap \left(\cap_{j=1}^{n} \chi_{(-\infty,s)}(x_{j})(\mathfrak{H})\right)\right).$ Then
$$\left\langle \left(\sum_{j=1}^{n} x_{j}\right)\xi,\xi\right\rangle \geq t$$

and

$$t > \left\langle \left(\sum_{j=1}^{n} y_j\right)\xi, \xi \right\rangle = \left\langle \left(\sum_{j=1}^{n} x_j \chi_{(-\infty,s)}(x_j)\right)\xi, \xi \right\rangle = \left\langle \left(\sum_{j=1}^{n} x_j\right)\xi, \xi \right\rangle \ge t$$

since $\xi \in \bigcap_{j=1}^n \chi_{(-\infty,s)}(x_j)(\mathfrak{H})$. Therefore

$$\chi_{[t,\infty)}\left(\sum_{j=1}^n x_j\right) \wedge \left(\chi_{[0,t)}\left(\sum_{j=1}^n y_j\right) \wedge \left(\wedge_{j=1}^n \chi_{(-\infty,s)}(x_j)\right)\right) = 0.$$

We deduce from Lemma 1.1(ii) that

$$\chi_{[t,\infty)}\left(\sum_{j=1}^n x_j\right) \prec \left(\chi_{[0,t)}\left(\sum_{j=1}^n y_j\right) \land \left(\land_{j=1}^n \chi_{(-\infty,s)}(x_j)\right)\right)^{\perp}.$$

and this gives us inequality (4.2). Using (4.2), we get

$$\tau \left(\chi_{[t,\infty)} \left(\sum_{j=1}^{n} x_{j} \right) \right) \leq \tau \left(\chi_{[t,\infty)} \left(\sum_{j=1}^{n} y_{j} \right) \right) + \tau \left(\vee_{j=1}^{n} \chi_{[s,\infty)}(x_{j}) \right)$$

$$(by \text{ Lemma } \mathbf{1.1}(ii - iii))$$

$$\leq \exp \left\{ -\frac{t}{M} \left(\log \left(1 + \frac{tM^{r-1}}{B} \right) - 1 \right) \right\} + \sum_{j=1}^{n} \tau \left(\chi_{[s,\infty)}(x_{j}) \right)$$

$$(by \text{ Lemma } \mathbf{1.1}(iv), \ (\mathbf{4.1}))$$

for any t > 0. An easy investigation shows that the latter inequality holds if we replace M by any number L with $L \ge M$. In addition, it holds for all $0 < L \le M$ since the function

$$f(\alpha) = \exp\left\{-\frac{t}{\alpha}\left(\log\left(1 + \frac{t\alpha^{r-1}}{B}\right) - 1\right)\right\}$$

is decreasing for any $\alpha > 0$. Thus

$$\tau\left(\chi_{[t,\infty)}\left(\sum_{j=1}^{n} x_{j}\right)\right) \leq \exp\left\{-\frac{t}{L}\left(\log\left(1+\frac{tL^{r-1}}{B}\right)-1\right)\right\} + \sum_{j=1}^{n} \tau\left(\chi_{[s,\infty)}(x_{j})\right) \quad (4.3)$$

for all t > 0, s > 0, L > 0.

Next, we deal with the modulus of $\sum_{j=1}^{n} x_j$. Inequality (4.3) together with (1.1) imply that

$$\tau\left(\chi_{[t,\infty)}\left(\left|\sum_{j=1}^{n} x_{j}\right|\right)\right) \leq 2\exp\left\{-\frac{t}{L}\left(\log\left(1+\frac{tL^{r-1}}{B}\right)-1\right)\right\} + \sum_{j=1}^{n}\tau\left(\chi_{[s,\infty)}(|x_{j}|)\right).$$

Now, by putting first L = s and $s = \frac{t}{\gamma}$, where $\gamma > 0$ we obtain that

$$\tau\left(\chi_{[t,\infty)}\left(\left|\sum_{j=1}^{n} x_{j}\right|\right)\right) \leq 2\exp\left\{-\gamma\left(\log\left(1+\frac{t^{r}}{\gamma^{r-1}B}\right)-1\right)\right\} + \sum_{j=1}^{n}\tau\left(\chi_{[t,\infty)}\left(\gamma|x_{j}|\right)\right).$$

10

For $p \geq 2$, we have

$$\begin{aligned} \left\|\sum_{j=1}^{n} x_{j}\right\|_{p}^{p} \\ &= \int_{0}^{\infty} p t^{p-1} \tau \left(\chi_{[t,\infty)}\left(\left|\sum_{j=1}^{n} x_{j}\right|\right)\right) dt \\ &\leq \sum_{j=1}^{n} \int_{0}^{\infty} p t^{p-1} \tau \left(\chi_{[t,\infty)}(\gamma | x_{j}|)\right) dt + 2p \int_{0}^{\infty} t^{p-1} \exp\left\{-\gamma \left(\log\left(1 + \frac{t^{r}}{\gamma^{r-1}B}\right) - 1\right)\right\} dt \\ &\leq \gamma^{p} \sum_{j=1}^{n} \|x_{j}\|_{p}^{p} + \frac{2p}{r} (\gamma^{r-1}B)^{\frac{p}{r}} e^{\gamma} \int_{0}^{\infty} \beta^{\frac{p-r}{r}} (1+\beta)^{-\gamma} d\beta \,, \end{aligned}$$

$$(4.4)$$

where we use the change of variable $\beta = \frac{t^r}{\gamma^{r-1}B}$. Next let γ be such that the last integral of the above inequality is convergent, i.e. let us choose $\gamma > \frac{p}{r}$. With this choice of γ inequality (4.4) implies the Rosenthal inequality with

$$C(p,r) = \max\left\{\gamma^p, \frac{2p}{r}\gamma^{\frac{p(r-1)}{r}}e^{\gamma}\int_0^\infty \beta^{\frac{p-r}{r}}(1+\beta)^{-\gamma}d\beta\right\}.$$

In the commutative setting, we have

Corollary 4.2. Let $1 \leq r \leq 2 \leq p < \infty$ and X_1, \dots, X_n be independent real random variables with expected values $\mathbb{E}(X_j) = 0$ $(j = 1, \dots, n)$. Then

$$\mathbb{E}\left(\left|\sum_{j=1}^{n} X_{j}\right|^{p}\right) \leq C(p) \left\{\sum_{j=1}^{n} \mathbb{E}\left(|X_{j}|^{p}\right) + \left(\sum_{j=1}^{n} \mathbb{E}\left(|X_{j}|^{r}\right)\right)^{p/r}\right\},\$$

where $C(p) = \min_{\gamma > p/2} \max\left\{\gamma^{p}, \frac{2p}{r}\gamma^{\frac{p(r-1)}{r}}e^{\gamma}\int_{0}^{\infty}\beta^{\frac{p-r}{r}}(1+\beta)^{-\gamma}d\beta\right\}.$

References

- G. Bennett, Probability inequalities for the sum of independent random variables, J. Amer. Statist. Assoc. 57 (1962), 33-45.
- [2] V. Bentkus, On Hoeffding's inequalities, Ann. Probab. 32 (2004), no. 2, 1650-1673.
- [3] D. L. Burkholder, Distribution function inequalities for martingales, Ann. Probab. 1 (1973), 19–42.
- [4] J. B. Conway, A Course in Functional Analysis, Second, Graduate Texts in Mathematics, springer– Verlag, New York, 1990.
- [5] J. Dixmier, von Neumann Algebras, North-Holland Mathematical Library, Vol. 27, North-Holland, Amesterdam, 1981.

GH. SADEGHI, M.S. MOSLEHIAN

- [6] W. Hoeffding, Probability inequalities for sums of bounded random variables, J. Amer. Statist. Assoc. 58 (1963), 13-30.
- [7] X.-L. Hu, An extension of Rosenthal's inequality, Appl. Math. Comput. 218 (2011), no. 8, 4638–4640.
- [8] W. B. Johnson, G. Schechtman and J. Zinn, Best constants in moment inequalities for linear combinations of independent and exchangeable random variables, Ann. Probab. 13 (1985), no. 1, 234–253.
- M. Junge and Q. Zeng, Noncommutative Bennett and Rosenthal inequalities, Ann. Probab. 41 (2013), no. 6, 4287–4316.
- [10] M. Junge and Q. Xu, Noncommutative Burkholder/Rosenthal inequalities. II. Applications, Israel J. Math. 167 (2008), 227–282.
- [11] M. S. Moslehian, M. Tominaga and K.-S. Saito, Schatten p-norm inequalities related to an extended operator parallelogram law, Linear Algebra Appl. 435 (2011), no. 4, 823–829.
- [12] E. Nelson, Note on noncommutative integration, J. Funct. Anal. 15 (1974), 103–116.
- B. de Pagter, Noncommutative Banach function spaces, Positivity, 197-227, Trends Math., Birkhüser, Basel, (2007).
- [14] H. P. Rosenthal, On the subspaces of $L_p(p > 2)$ spanned by sequences of independent random variables, Israel J. Math. 8 (1970), 273-303.
- [15] M. B. Ruskai, Inequalities for traces on von Neumann algebras, Comm. Math. Phys. 26 (1972), 280-289.
- [16] Gh. Sadeghi, Non-commutative Orlicz spaces associated to a modular on τ-measurable operators, J. Math. Anal. Appl. 395 (2012), no. 2, 705–715.
- [17] J. A. Tropp, User-friendly tail bounds for sums of random matrices, Found. Comput. Math. 12 (2012), no. 4, 389-434.
- [18] F. J. Yeadon, Noncommutative L^p-spaces, Proc. Camb. Phil. Soc. 77 (1975), 91–102.
- [19] Q. Xu, Embedding of C_q and R_q into noncommutative L_p -spaces, $1 \le p < q \le 2$, Math. Ann. **335** (2006), no. 1, 109-131.
- [20] Q. Xu, Operator spaces and noncommutative L_p , Lectures in the Summer School on Banach spaces and Operator spaces, Nankai University China, 2007.

¹ Department of Mathematics and Computer Sciences, Hakim Sabzevari University, P.O. Box 397, Sabzevar, Iran

E-mail address: ghadir540gmail.com, g.sadeghi@hsu.ac.ir

² DEPARTMENT OF PURE MATHEMATICS, CENTER OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN

E-mail address: moslehian@um.ac.ir, moslehian@member.ams.org