

The First and Second Most Symmetric Nonsingular Cubic Surfaces

H. Kaneta, S. Marcugini and F. Pambianco

February 29, 2024

Abstract

The first and second most symmetric nonsingular cubic surfaces are $x^3 + y^3 + z^3 + t^3 = 0$ and $x^2y + y^2z + z^2t + t^2x = 0$, respectively.

1 Introduction

Throughout this paper k stands for an algebraically closed field of characteristic zero. Let C_d be a nonsingular curve of degree $d \geq 4$ in the two-dimensional projective space. As is well known, its automorphism group $\text{Aut}(C_d)$ is a finite subgroup of $PGL_3(k)$ such that $|\text{Aut}(C_d)| \leq 42d(d-3)$. In fact the maximum value β_d of $|\text{Aut}(C_d)|$ is equal to 168, 360 or $6d^2$, and attained by the Klein quartic curve, the Wiman sextic curve or the Fermat curve according as $d = 4$, $d = 6$ or $d \notin \{4, 6\}$ [5]. Moreover the Klein quartic and the Wiman sextic give rise to highly symmetric MDS codes [2].

Let S_d be a nonsingular surface of degree $d \geq 3$ in the three-dimensional projective space. Its automorphism group $\text{Aut}(S_d)$ is a finite subgroup of $PGL_4(k)$ if $d \neq 4$, can be an infinite group if $d = 4$ [4]. Highly symmetric surfaces may be both of theoretical and of practical interest. Hosoh has completed the classification of the automorphism groups of nonsingular cubic surfaces [1]. According to his classification the maximum and the second maximum of $|\text{Aut}(S_3)|$ is attained by the semidirect product $(\mathbf{Z}_3)^3 \rtimes_s \mathbf{S}_4$ and \mathbf{S}_5 , respectively. It may happen that $|\text{Paut}(S_d)|$, the order of the projective automorphism group of S_d , has the finite upper bound for every $d \geq 3$. In this paper we will specify S_3 such that $\text{Aut}(S_3)$ is isomorphic to $(\mathbf{Z}_3)^3 \rtimes_s \mathbf{S}_4$ or \mathbf{S}_5 .

$M_{m,n}(k)$ stands for the set of all $m \times n$ matrices of entries in k . By definition $M_n(k) = M_{n,n}(k)$, $GL_n(k) = \{A = [a_{ij}] \in M_n(k) : \det A \neq 0\}$, and $PGL_n(k) = GL_n(k)/(E_n)$, where (E_n) is the subgroup $\{\lambda E_n : \lambda \in k^*\}$ (E_n is the unit matrix in $GL_n(k)$). The coset $A(E_n)$ containing an $A \in GL_n(k)$ will be denoted (A) . We denote by $k[x]$ the k -algebra of polynomials in $x = [x_1, \dots, x_n]$ over k . For an $A \in GL_n(k)$ and $f \in k[x]$ we define a polynomial $f_A \in k[x]$ to be $f_A(x) = f(\sum \alpha_{1j}x_j, \dots, \sum \alpha_{nj}x_j)$, where $A^{-1} = [\alpha_{ij}]$. As is well known, the map $T_A : k[x] \rightarrow k[x]$ assigning f_A to f is a k -algebra isomorphism of $k[x]$ such that $T_A T_B = T_{AB}$, that is, $(f_B)_A = f_{AB}$. \mathbf{S}_n denotes the symmetric group, the group of all permutations of n elements.

A homogeneous polynomial f of degree $d \geq 1$ defines a projective algebraic set $V(f) = \{(a) \in P^{n-1} : f(a) = 0\}$ of an $(n-1)$ -dimensional projective space P^{n-1} over k . $V(f)$ is called a hypersurface of degree d . If $n = 4$, $d = 3$, and $V(f)$ is nonsingular, the automorphism group $\text{Aut}(V(f))$ of the hypersurface $V(f)$ is a finite subgroup of $PGL_4(k)$ consisting of $(A) \in PGL_4(k)$ such that $f_A \sim f$ [4]. Let $a = [a_1, \dots, a_n] \in k^n$, $(a) \in V(f)$, and $A \in GL_n(k)$. Then (a) is a singular point of $V(f)$ if $f_{x_i}(a) = 0$ for all i . If (a) is a nonsingular point of $V(f)$, $V(\sum_{i=1}^n \gamma_i x_i)$ is the tangent plane to $V(f)$ at (a) , where

$\gamma_i = f_{x_i}(a)$. Clearly $(A) : V(f) \rightarrow V(f_A)$ is a bijection, and if $b = Aa$ with $(a) \in V(f)$, then $(f_A)_{x_j}(b) = \sum_{i=1}^n f_{x_i}(a)\alpha_{ij}$, where $A^{-1} = [\alpha_{ij}]$. Consequently $(b) = (A)(a)$ is a nonsingular point of $V(f_A)$ if and only if (a) is a nonsingular point of $V(f)$, and the tangent plane of $V(f_A)$ at (b) coincides with $(A)V(\sum_{i=1}^n \gamma_i x_i)$. In particular if (a) is a nonsingular point of $V(f)$, $f_A \sim f$ and $(A)(a) = (a)$, then $[f_{x_1}(a), \dots, f_{x_n}(a)]A \sim [f_{x_1}(a), \dots, f_{x_n}(a)]$.

In §1 it will be shown that the Fermat surface $S = V(x^3 + y^3 + z^3 + t^3)$ is the unique cubic nonsingular surface, up to projective equivalence, such that $\text{Aut}(S)$ is isomorphic to $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$. In §2 it will be shown that a surface $S' = V(x^2t + y^2z + z^2t + t^2x)$ is the unique cubic nonsingular surface such that $\text{Aut}(S')$ is isomorphic to \mathbf{S}_5 .

2 $\mathbf{Z}_3^3 \times_s \mathbf{S}_4$ -invariant nonsingular cubic surfaces

Let $\omega \in k^*$ be of order three. Any subgroup of $PGL_4(k)$ isomorphic to \mathbf{Z}_3 is conjugate to $\langle (\text{diag}[\omega, 1, 1, 1]) \rangle$ or $\langle (\text{diag}[\omega, \omega^2, 1, 1]) \rangle$. It can be verified easily that $X = [x_{ij}] \in GL_4(k)$ satisfies $\text{diag}[\omega, 1, 1, 1]X \sim X\text{diag}[\omega, 1, 1, 1]$ (resp. $\text{diag}[\omega, \omega^2, 1, 1]X \sim X\text{diag}[\omega, \omega^2, 1, 1]$) if and only if $x_{1i} = x_{i1} = 0$ for all $i \in [2, 4]$ (resp. $x_{1i} = x_{i1} = x_{2j} = x_{j2} = 0$ for all $i \in [2, 4]$ and all $j \in [3, 4]$). Consequently a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^2$ is conjugate to $\langle (A), (B) \rangle$, where $\text{ord}(A) = \text{ord}(B) = 3$ and $(B^j) \in \langle (A) \rangle$ if and only if $j \in 3\mathbf{Z}$. We may assume that $A = \text{diag}[\omega, \omega^2, 1, 1]$ or $A = \text{diag}[\omega, 1, 1, 1]$ and that $B = \text{diag}[b_1, b_2, b_3, b_4]$. Assume first that $A = \text{diag}[\omega, \omega^2, 1, 1]$. If $|\{b_1, b_2, b_3, b_4\}| = 3$, then we may assume that B is equal to one of $S - \{A\}$, where

$$S = \{\text{diag}[1, 1, \omega, \omega^2], \text{diag}[1, \omega, 1, \omega^2], \text{diag}[1, \omega, \omega^2, 1], \text{diag}[\omega, 1, 1, \omega^2], \\ \text{diag}[\omega, 1, \omega^2, 1], \text{diag}[\omega, \omega^2, 1, 1]\},$$

Unless $B = \text{diag}[1, 1, \omega, \omega^2]$, there exist integers i and j such that $(A^i B^j)$ is equal to $(\text{diag}[1, 1, \omega, 1])$ or $(\text{diag}[1, 1, 1, \omega])$. So $\langle (A), (B) \rangle$ is conjugate to

$$\langle (\text{diag}[\omega, \omega^2, 1, 1]), (\text{diag}[1, 1, \omega, \omega^2]) \rangle \text{ or } \langle (\text{diag}[\omega, \omega^2, 1, 1]), (\text{diag}[1, 1, \omega, 1]) \rangle,$$

provided $|\{b_1, b_2, b_3, b_4\}| = 3$. Clearly $\langle (A), (B) \rangle$ is conjugate to

$$\langle (\text{diag}[\omega, \omega^2, 1, 1]), (\text{diag}[1, 1, \omega, 1]) \rangle \text{ or } \langle (\text{diag}[\omega, 1, 1, 1]), (\text{diag}[1, 1, \omega, 1]) \rangle,$$

provided $|\{b_1, b_2, b_3, b_4\}| = 2$. Assume secondly that $A = \text{diag}[\omega, 1, 1, 1]$. According as $|\{b_1, b_2, b_3, b_4\}|$ is two or three, we may assume B is equal to one of $T - \{A\}$ or S , where

$$T = \{\text{diag}[\omega, 1, 1, 1], \text{diag}[1, \omega, 1, 1], \text{diag}[1, 1, \omega, 1], \text{diag}[1, 1, 1, \omega]\}.$$

Thus a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^2$ is conjugate to one of

$$\begin{aligned} &\langle (\text{diag}[\omega, \omega^2, 1, 1]), (\text{diag}[1, 1, \omega, \omega^2]) \rangle, \\ &\langle (\text{diag}[\omega, \omega^2, 1, 1]), (\text{diag}[1, 1, \omega, 1]) \rangle, \\ &\langle (\text{diag}[\omega, 1, 1, 1]), (\text{diag}[1, \omega, 1, 1]) \rangle. \end{aligned}$$

Even though the following lemma may be known, we shall give a proof for the sake of completeness.

Lemma 2.1. *A subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^3$ is conjugate to*

$$G_{27} = \langle (\text{diag}[\omega, 1, 1, 1]), (\text{diag}[1, \omega, 1, 1]), (\text{diag}[1, 1, \omega, 1]) \rangle.$$

Proof. Let $G = \langle (A), (B), (C) \rangle$ be a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^3$. Since $\langle (A), (B) \rangle$ is isomorphic to $(\mathbf{Z}_3)^2$, we may assume one of the following three cases: 1) $A = \text{diag}[\omega, \omega^2, 1, 1]$, $B = \text{diag}[1, 1, \omega, \omega^2]$. 2) $A = \text{diag}[\omega, \omega^2, 1, 1]$, $B = \text{diag}[1, 1, \omega, 1]$. 3) $A = \text{diag}[\omega, 1, 1, 1]$, $B = \text{diag}[1, \omega, 1, 1]$. In any case we may assume $C = \text{diag}[c_1, c_2, c_3, c_4] \neq E_4$ with $c_i^3 = 1$ but $(C) \notin \langle (A), (B) \rangle$. Let $m = |\{c_1, c_2, c_3, c_4\}|$. Assume the case 1). If $m = 2$, then $G = G_{27}$. If $m = 3$, then we may assume C is equal to one of $S - \{A, B\}$ so that G contains one of T , hence $G = G_{27}$. Assume the case 2). If $m = 2$, then we may assume C is one of $T - \{B\}$, hence $G = G_{27}$. If $m = 3$, then we may assume C is one of $S - \{\text{diag}[1, \omega, 1, \omega^2], \text{diag}[\omega, 1, 1, \omega^2], A\}$, hence $G = G_{27}$. Note that $(\text{diag}[1, \omega, 1, \omega^2]), (\text{diag}[\omega, 1, 1, \omega^2]) \in \langle (A), (B) \rangle$. Assume the case 3). If $m = 2$, then we may assume C is one of $T - \{A, B\}$, hence $G = G_{27}$. If $m = 3$, then we may assume C is one of $S - \{\text{diag}[\omega, \omega^2, 1, 1]\}$, hence $G = G_{27}$.

The canonical group representation $\hat{\cdot} : \mathbf{S}_4 \rightarrow GL_4(k)$ of \mathbf{S}_4 is the one such that $\hat{\sigma}x = y$ with $y_i = x_{\sigma^{-1}(i)}$ for any column vector $x \in k^4$. Clearly this representation is an isomorphism. Let $r \geq 2$ be an integer, $\delta \in k^*$ be of order r ,

$$D(r) = \{\text{diag}[\lambda_1, \lambda_2, \lambda_3, \lambda_4] \in GL_4(k) : \lambda_1^r = \lambda_2^r = \lambda_3^r = \lambda_4^r\}.$$

The factor group $(D(r)) = D(r)/k^*E_4$ is isomorphic to $\{\text{diag}[\delta^i, \delta^j, \delta^\ell, 1] : i, j, \ell \in [0, r-1]\}$ which is isomorphic to $(\mathbf{Z}_r)^3$. A map $\varphi_\sigma : (D(r)) \rightarrow (D(r))$ defined by $\varphi_\sigma((A)) = (\hat{\sigma}A\hat{\sigma}^{-1})$ is a group automorphism such that $\varphi_{\sigma\tau} = \varphi_\sigma \circ \varphi_\tau$. The factor group $D(r)\hat{\mathbf{S}}_4/k^*E_4$ is isomorphic to the semidirect product $(D(r)) \times_s \mathbf{S}_4$ such that $((A), \sigma)((A'), \sigma') = ((A)\varphi_\sigma(A'), \sigma\sigma')$. Clearly $(D(r)) \times_s \mathbf{S}_4$ is isomorphic to $(\mathbf{Z}_r)^3 \times_s \mathbf{S}_4$. We note that $\hat{\sigma}\text{diag}[a_1, a_2, a_3, a_4]\hat{\sigma}^{-1} = \text{diag}[b_1, b_2, b_3, b_4]$, where $b_i = a_{\sigma^{-1}(i)}$. We may skip the proof of the following lemma [6].

Lemma 2.2. *If $r \geq 3$, then the projective automorphism group of the surface $V(x^r + y^r + z^r + t^r)$ is $D(r)\hat{\mathbf{S}}_4/k^*E_4$, which can be identified with $(D(r)) \times_s \mathbf{S}_4$.*

Lemma 2.3. *Any $(\mathbf{Z}_3)^3$ -invariant nonsingular cubic surface is projectively equivalent to $V(x^3 + y^3 + z^3 + t^3)$. In particular any $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$ -invariant nonsingular cubic surface is projectively equivalent to $V(x^3 + y^3 + z^3 + t^3)$, and its automorphism group is conjugate to $(D(3)) \times_s \mathbf{S}_4$.*

Proof. By Lemma 1.1 it suffices to show that any G_{27} -invariant nonsingular cubic surface is $V(ax^3 + by^3 + cz^3 + dt^3)$, where a, b, c and d are nonzero constants. Let $V(f)$ be a G_{27} -invariant nonsingular cubic surface, where the homogeneous polynomial $f(x, y, z, t)$ of degree three has the form

$$\begin{aligned} & a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 \\ & + x^2(b_{12}y + b_{13}z + b_{14}t) + y^2(b_{21}x + b_{23}z + b_{24}t) + z^2(b_{31}x + b_{32}y + b_{34}t) \\ & + t^2(b_{41}x + b_{42}y + b_{43}z) + c_1yzt + c_2xzt + c_3xyt + c_4xyz. \end{aligned}$$

Let

$$A_1 = \text{diag}[\omega, 1, 1, 1], A_2 = \text{diag}[1, \omega, 1, 1], A_3 = \text{diag}[1, 1, \omega, 1].$$

It is evident that $\text{ord}(A_i) = 3$. Since G_{27} contains (A_i) ($i \in [1, 4]$), $f_{A_i^{-1}}$ is equal to one of $\{f, \omega f, \omega^2 f\}$. If $f_{A_1^{-1}}$ is equal to ωf or $\omega^2 f$, then f is divisible by x , hence $V(f)$ is singular. Assume $f_{A_1^{-1}} = f$. Then

$$f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + y^2(b_{23}z + b_{24}t) + z^2(b_{32}y + b_{34}t) + t^2(b_{42}y + b_{43}z) + c_1yzt.$$

Unless $f_{A_2^{-1}} = f$, $V(f)$ is singular. Therefore $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_{34}z^2t + b_{43}t^2z$. Unless $f_{A_3^{-1}} = f$, $V(f)$ is singular. Thus $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3$. Now $V(f)$ is nonsingular if and only if $a_1a_2a_3a_4 \neq 0$.

3 \mathbf{S}_5 -invariant nonsingular cubic surfaces

A subgroup of $PGL_4(k)$ isomorphic to the symmetric group \mathbf{S}_5 is one of three groups C_{5I} , C_{5II} , C_{5III} up to conjugacy [3]. We denote these groups by $G(1)$, $G(2)$ and $G(3)$, respectively. There exist group isomorphisms $\varphi_i : \mathbf{S}_5 \rightarrow G(i)$ such that $(E_1) = \varphi_i((123))$, $(E_2) = \varphi_i((12)(34))$, $(E_3) = \varphi_i((12)(45))$ and $(F) = \varphi_i((12))$ generate $G(i)$ ($i \in [1, 3]$) (cf [7, (2.14) in chap.3]). Note that $(123) = (12)(23)$. Let $\omega = \frac{-1+i\sqrt{3}}{2}$. Then $G(1) = \langle (E_1), (E_2), (E_3), (F) \rangle$, where

$$E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \omega & 0 \\ 0 & 0 & 0 & \omega^2 \end{bmatrix}, \quad E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ 0 & \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ 0 & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}, \quad E_3 = \begin{bmatrix} \frac{1}{4} & -\frac{\sqrt{15}}{4} & 0 & 0 \\ \frac{\sqrt{15}}{4} & \frac{1}{4} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix},$$

$$F = \frac{1+i}{\sqrt{2}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Lemma 3.1. *The $G(1)$ -invariant nonsingular cubic surface is $V(f)$, where*

$$f(x, y, z, t) = 3\sqrt{15}x^3 + 10(y^3 + z^3 + t^3) - 3\sqrt{15}xy^2 - 6(\sqrt{15}x + 5y)zt.$$

Moreover, $\text{Aut}(V(f)) = G(1)$.

Proof. Let $F' = \frac{\sqrt{2}}{1+i}F$. Assume that $V(f)$ is a $G(1)$ -invariant nonsingular cubic surface, where $f(x, y, z, t)$ is a cubic homogeneous polynomial of the form as in the proof of Lemma 1.3. Since $E_1^3 = E_4$, $f_{E_1^{-1}} = \omega^i f$ for some $i \in [0, 2]$. If $i = 1$, then $f(x, y, z, t) = b_{13}x^2z + b_{23}y^2z + b_{34}z^2t + t^2(b_{41}x + b_{42}y) + c_4xyz$. Since $f_{F'^{-1}} \sim f$, we see $b_{13} = b_{23} = 0$, hence $V(f)$ is singular at $(1, 0, 0, 0)$. Similarly $i \neq 2$, namely $i = 0$, so that $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_1x^2y + b_2y^2x + c_1yzt + c_2xzt$. We have $a_3a_4 \neq 0$, for $V(f)$ is nonsingular. If $a_1 = a_2 = 0$, the condition $f_{E_3^{-1}} \sim f$ yields $b_1 = b_2 = 0$, hence $V(f)$ is singular. Thus $[a_1, a_2] \neq [0, 0]$. Therefore $f_{F'^{-1}} = f$, hence $a_3 = a_4$. Now $f_{E_3^{-1}} = f$, and $f_{E_3^{-1}} = -f$ does not hold. The condition $f_{E_3^{-1}} = f$ is equivalent to $c_2 = \sqrt{15}c_1/5$ and

$$\begin{bmatrix} -65 & 15\sqrt{15} & \sqrt{15} & -15 \\ 15\sqrt{15} & -63 & 15 & \sqrt{15} \\ 3\sqrt{15} & 45 & -93 & 13\sqrt{15} \\ -45 & 3\sqrt{15} & 13\sqrt{15} & -35 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ b_1 \\ b_2 \end{bmatrix} = 0.$$

The general $[a_1, a_2, b_1, b_2]$ satisfying the second condition above is $\lambda[3\sqrt{15}, 10, 0, -3\sqrt{15}]$, for the rank of the above matrix is equal to three. Consequently we may assume $f(x, y, z, t) = 3\sqrt{15}x^3 + 10y^3 + a_3(z^3 + t^3) - 3\sqrt{15}xy^2 + c_1(yzt + \frac{\sqrt{15}}{5}xzt)$. Now the condition $f_{E_3^{-1}} \sim f$ implies $f_{E_3^{-1}} = f$. Using the formula $(a + b + c)^3 = a^3 + b^3 + c^3 + 3a^2(b + c) + 3b^2(a + c) +$

$3c^2(a+b) + 6abc$, we can write $f_{E_3^{-1}}(x, y, z, t)$ as

$$\begin{aligned} & 3\sqrt{15}x^3 + \left(-\frac{10}{27} + \frac{16}{27}a_3 - \frac{4}{27}c_1\right)y^3 + \left(\frac{80}{27} + \frac{7}{27}a_3 - \frac{4}{27}c_1\right)z^3 + \left(\frac{80}{27} + \frac{7}{27}a_3 - \frac{4}{27}c_1\right)t^3 \\ & + y^2\left\{\left(-\frac{\sqrt{15}}{3} + \frac{4\sqrt{15}}{45}c_1\right)x + \left(\frac{60}{27} + \frac{12}{27}a_3 + \frac{6}{27}c_1\right)(z+t)\right\} \\ & + z^2\left\{\left(-\frac{4\sqrt{15}}{3} - \frac{2\sqrt{15}}{45}c_1\right)x + \left(-\frac{120}{27} + \frac{30}{27}a_3 + \frac{6}{27}c_1\right)y + \left(\frac{240}{27} - \frac{6}{27}a_3 + \frac{6}{27}c_1\right)t\right\} \\ & + t^2\left\{\left(-\frac{4\sqrt{15}}{3} - \frac{2\sqrt{15}}{45}c_1\right)x + \left(-\frac{120}{27} + \frac{30}{27}a_3 + \frac{6}{27}c_1\right)y + \left(\frac{240}{27} - \frac{6}{27}a_3 + \frac{6}{27}c_1\right)z\right\} \\ & + \left(-\frac{240}{27} - \frac{48}{27}a_3 + \frac{3}{27}c_1\right)yzt + \left(-\frac{8\sqrt{15}}{3} + \frac{\sqrt{15}}{9}c_1\right)xzt \\ & + \left(\frac{4\sqrt{15}}{3} + \frac{2\sqrt{15}}{45}c_1\right)xyt + \left(\frac{4\sqrt{15}}{3} + \frac{2\sqrt{15}}{45}c_1\right)xyz. \end{aligned}$$

Now $a_3 = 10$, $c = -30$ and $f_{E_3^{-1}} = f$ holds, as desired. We can show that $V(f)$ is nonsingular.

Let $\varepsilon = -\frac{\sqrt{5}-1}{4} + i\frac{\sqrt{10+2\sqrt{5}}}{4}$, hence $\text{ord}(\varepsilon) = 5$, $\alpha = \frac{-\sqrt{5}+1}{2}$, $\beta = \alpha^2$, $\gamma = -\alpha$, and let

$$H = \begin{bmatrix} \varepsilon^4 & 0 & 0 & 0 \\ 0 & \varepsilon^2 & 0 & 0 \\ 0 & 0 & \varepsilon & 0 \\ 0 & 0 & 0 & \varepsilon^3 \end{bmatrix}, \quad I = \begin{bmatrix} 1 & \alpha & \beta & \gamma \\ \alpha & \beta & \gamma & 1 \\ \beta & \gamma & 1 & \alpha \\ \gamma & 1 & \alpha & \beta \end{bmatrix}.$$

Lemma 3.2. *(H) and (I) generate in $PGL_4(k)$ a group $G(1)'$ conjugate to $G(1)$. The $G(1)'$ -invariant nonsingular cubic surface is $V(x^2y + y^2z + z^2t + t^2x)$. Moreover $\text{Aut}(V(x^2y + y^2z + z^2t + t^2x)) = G(1)'$.*

Proof. The transpositions $(i \ i+1)$ ($i \in [1, 4]$) generate \mathbf{S}_5 , while $(12345)(j \ j+1)(12345)^{-1} = (j+1 \ j+2)$ ($j \in [1, 3]$). Therefore (12345) and (12) generate \mathbf{S}_5 . Clearly $(123)(34)(45) = (12345)$. In view of the group isomorphism $\varphi_1 : S_5 \rightarrow G(1)$, $\varphi_1((12345)) = (E_1FE_2FE_3)$ and $\varphi_1((12)) = (F)$. Let

$$K = E_1FE_2FE_3 = \begin{bmatrix} -\frac{1}{4} & \frac{\sqrt{15}}{4} & 0 & 0 \\ -\frac{\sqrt{15}}{12} & -\frac{1}{12} & \frac{2}{3} & \frac{2}{3} \\ \frac{\sqrt{15}\omega}{6} & \frac{\omega}{6} & \frac{2\omega}{3} & -\frac{\omega}{3} \\ \frac{\sqrt{15}\omega^2}{6} & \frac{\omega^2}{6} & -\frac{\omega^2}{3} & \frac{2\omega^2}{3} \end{bmatrix}, \quad T = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 1 & 1 \end{bmatrix}.$$

Now

$$\begin{aligned} T^{-1}FT &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}, \quad T^{-1}KT = \begin{bmatrix} -\frac{1}{4} & \frac{\sqrt{15}}{4} & 0 & 0 \\ -\frac{\sqrt{15}}{12} & -\frac{1}{12} & \frac{4}{3} & 0 \\ -\frac{\sqrt{15}}{12} & -\frac{1}{12} & -\frac{1}{6} & -i\frac{\sqrt{3}}{2} \\ -i\frac{\sqrt{5}}{4} & -i\frac{\sqrt{3}}{12} & -i\frac{\sqrt{3}}{6} & -\frac{1}{2} \end{bmatrix}, \\ \det(T^{-1}KT - \lambda E_4) &= \lambda^4 + \lambda^3 + \lambda^2 + \lambda + 1 = \prod_{i=1}^4 (\lambda - \varepsilon^i). \end{aligned}$$

If λ is an eigenvalue of $T^{-1}KT$ and $x \in k^4$ satisfies $(T^{-1}KT - \lambda E_4)x = 0$, then

$$[x_1, x_2, x_3, x_4] = x_1[1, (1+4\lambda)\frac{\sqrt{15}}{15}, (1+\lambda+3\lambda^2)\frac{\sqrt{15}}{15}, i(-\lambda^4+\lambda^3)\frac{\sqrt{3}}{5}].$$

So $S^{-1}T^{-1}KTS = \text{diag}[\varepsilon^4, \varepsilon^2, \varepsilon, \varepsilon^3]$, where

$$S = \begin{bmatrix} 1 & 1 & 1 & 1 \\ (1+4\varepsilon^4)\frac{\sqrt{15}}{15} & (1+4\varepsilon^2)\frac{\sqrt{15}}{15} & (1+4\varepsilon)\frac{\sqrt{15}}{15} & (1+4\varepsilon^3)\frac{\sqrt{15}}{15} \\ (1+\varepsilon^4+3\varepsilon^3)\frac{\sqrt{15}}{15} & (1+\varepsilon^2+3\varepsilon^4)\frac{\sqrt{15}}{15} & (1+\varepsilon+3\varepsilon^2)\frac{\sqrt{15}}{15} & (1+\varepsilon^3+3\varepsilon)\frac{\sqrt{15}}{15} \\ i(-\varepsilon+\varepsilon^2)\frac{\sqrt{3}}{5} & i(-\varepsilon^3+\varepsilon)\frac{\sqrt{3}}{5} & i(-\varepsilon^4+\varepsilon^3)\frac{\sqrt{3}}{5} & i(-\varepsilon^2+\varepsilon^4)\frac{\sqrt{3}}{5} \end{bmatrix}$$

so that $\det S = -i\frac{8\sqrt{3}}{25}\varepsilon^3(\varepsilon-1)^3(3\varepsilon^3+6\varepsilon^2+4\varepsilon+2)$. We denote the (i, j) -cofactor of $S = [s_{ij}]$ by \tilde{s}_{ij} . Note that $S^{-1}\text{diag}[1, 1, 1, -1]S = E_4 + S^{-1}\text{diag}[0, 0, 0, -2]S$ and that the i -th row of $S^{-1}\text{diag}[0, 0, 0, -2]S$ is equal to $-2\tilde{s}_{4i}/\det S \times (4\text{-th row of } S)$. Denote $TS\text{diag}[1, \varepsilon^2, \varepsilon^3, \varepsilon]$ by S' . By computation we get the components of $(3\varepsilon^3+6\varepsilon^2+4\varepsilon+2)S'^{-1}FTS'$ as follows.

$$\begin{bmatrix} 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon \\ 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 \\ -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 \\ \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 \end{bmatrix}.$$

Moreover, since $(2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1} = (7\varepsilon^3+4\varepsilon^2+\varepsilon+8)/5$, we can verify

$$\begin{aligned} (2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(2\varepsilon^4+\varepsilon^3+2) &= \varepsilon^3+\varepsilon^2+1 = \alpha, \\ (2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(-\varepsilon^4+\varepsilon^2) &= \varepsilon^3+\varepsilon^2+2 = \alpha^2, \\ (2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(\varepsilon^3+2\varepsilon^2+2\varepsilon) &= -\varepsilon^3-\varepsilon^2-1 = -\alpha. \end{aligned}$$

We have shown that $S'^{-1}KS' = H$ and $(S'^{-1}FS') = (I)$.

Finally we shall show that a cubic homogeneous polynomial $f(x, y, z, t)$ of the form as in the proof of Lemma 1.3 such that $f_{H^{-1}} \sim f$, $f_{I^{-1}} \sim f$ and that $V(f)$ is nonsingular is proportional to $x^2y + y^2z + z^2t + t^2x$. Since $H^5 = E_4$, $f_H^{-1} = \varepsilon^i f$ for some $i \in [0, 4]$. If $i = 1$, then $f(x, y, z, t) = a_2y^3 + b_{14}x^2t + b_3t^3x + c_1yzt$, hence $V(f)$ is singular at $(0, 0, 0, 1)$. Similarly, unless $i = 0$, $V(f)$ has a singular point. If $i = 0$, then $f(x, y, z, t) = b_1x^2y + b_2y^2z + b_3z^2t + b_4t^2x$. Denoting the coefficients of x^3 , y^3 , z^3 and t^3 in $f_{I^{-1}}$ by

b'_1, b'_2, b'_3 and b'_4 , respectively, and noting $\alpha^2 = \alpha + 1$, we obtain

$$b' = \alpha \begin{bmatrix} 1 & \alpha^3 & -\alpha^4 & \alpha \\ \alpha^3 & -\alpha^4 & \alpha & 1 \\ -\alpha^4 & \alpha & 1 & \alpha^3 \\ \alpha & 1 & \alpha^3 & -\alpha^4 \end{bmatrix} b.$$

Since $b' = 0$ and the rank of the matrix involved is equal to three, it follows that $b_1 = b_2 = b_3 = b_4 \neq 0$. Now let $f = x^2y + y^2z + z^2t + t^2x$. In order to see $f_{I^{-1}} = 5(4\alpha+3)f$, we note that denoting $(x + \alpha y + \beta z + \gamma t)^2(x + \alpha x + \beta y + \gamma z + t)$ by $g(x, y, z, t)$, we have $f_{I^{-1}}(x, y, z, t) = g(x, y, z, t) + g(t, x, y, z) + g(z, t, x, y) + g(y, z, t, x)$.

The group $G(2)$ is generated by (E_1) , (E_2) , (E_3) and (F) , where

$$\begin{aligned} E_1 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \omega & 0 \\ 0 & 0 & 0 & \omega^2 \end{bmatrix}, \quad E_2 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 0 & 0 & \sqrt{2} \\ 0 & -1 & \sqrt{2} & 0 \\ 0 & \sqrt{2} & 1 & 0 \\ \sqrt{2} & 0 & 0 & -1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \\ F &= \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}. \end{aligned}$$

Lemma 3.3. *There exist no $G(2)$ -invariant nonsingular cubic surfaces.*

Proof. Let $f(x, y, z, t)$ be a $G(2)$ -invariant cubic homogeneous polynomial of the form as in the proof of Lemma 1.3. Since $F^4 = E_4$, $f_{F^{-1}} = i^j f$ for some $j \in [0, 3]$. If $j = 0$ or $j = 2$, then $f = 0$. If $j = 1$, then $f(x, y, z, t)$ has the form

$$\begin{aligned} & a_1(x^3 - iy^3) + a_3(z^3 - it^4) \\ & + x^2(b_{12}y + b_{13}z + b_{14}t) + y^2(ib_{12}x + ib_{14}z - ib_{13}t) \\ & + z^2(b_{31}x + b_{32}y + b_{34}t) + t^2(ib_{32}x - ib_{31}y + ib_{34}z) \\ & + c_1yzt - ic_1xzt + c_3xyt - ic_3xyz. \end{aligned}$$

Since $E_3^2 = E_4$, $f_{E_3^{-1}} = \pm f$. The coefficients of t^3 , t^2z , t^2x and t^2y of $f_{E_3^{-1}}$ (resp. $\pm f$) are

$$a_3, b_{34}, \frac{\sqrt{3}}{2}b_{31} + \frac{1}{2}b_{32}, \frac{1}{2}b_{31} - \frac{\sqrt{3}}{2}b_{32} \text{ (resp. } \mp ia_3, \pm ib_{34}, \pm ib_{32}, \mp ib_{31}),$$

hence $a_3 = b_{34} = b_{31} = b_{32} = 0$. Consequently $V(f)$ is singular at $(0, 0, 0, 1)$.

Let $\nu_1 = \sqrt{\frac{3}{8}} + i\sqrt{\frac{5}{8}}$ and $\nu_2 = \sqrt{\frac{3}{8}} - i\sqrt{\frac{5}{8}}$. The group $G(3)$ is generated by (E_1) , (E_2) , (E_3) and (F) , where

$$\begin{aligned} E_1 &= \begin{bmatrix} \omega & 0 & 0 & 0 \\ 0 & \omega & 0 & 0 \\ 0 & 0 & \omega^2 & 0 \\ 0 & 0 & 0 & \omega^2 \end{bmatrix}, \quad E_2 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 0 & 0 & \sqrt{2} \\ 0 & 1 & \sqrt{2} & 0 \\ 0 & \sqrt{2} & -1 & 0 \\ \sqrt{2} & 0 & 0 & -1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} 0 & 0 & 0 & \nu_1 \\ 0 & 0 & \nu_2 & 0 \\ 0 & \nu_1 & 0 & 0 \\ \nu_2 & 0 & 0 & 0 \end{bmatrix}, \\ F &= \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{bmatrix}. \end{aligned}$$

Lemma 3.4. *There exist no $G(3)$ -invariant nonsingular cubic surfaces. More precisely, there exist no $G(3)$ -invariant cubic forms except zero.*

Proof. Let $E'_1 = \omega^{-1}E_1$ and $f(x, y, z, t)$ a $G(3)$ -invariant cubic form. If $f_{E_1'^{-1}} = \omega f$, then f has the form

$$x^2(b_{13}z + b_{14}t) + y^2(b_{23}z + b_{24}t) + c_3xyt + c_4xyz,$$

hence the condition $f_{F^{-1}} \sim f$ implies $f = 0$. Similarly, if $f_{E_1'^{-1}} = \omega^2 f$, then $f = 0$. Now assume $f_{E_1'^{-1}} = f$, so that $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_1x^2y + b_2y^2x + b_3z^2t + b_4t^2z$. Since $f_{F^{-1}} = i^j f$ for some $j \in [0, 3]$, we obtain

$$\begin{bmatrix} i^j & 1 \\ -1 & i^j \end{bmatrix} \begin{bmatrix} a_1 \\ a_4 \end{bmatrix} = 0, \quad \begin{bmatrix} i^j & -1 \\ 1 & i^j \end{bmatrix} \begin{bmatrix} a_2 \\ a_3 \end{bmatrix} = 0, \quad \begin{bmatrix} i^j & -1 \\ 1 & i^j \end{bmatrix} \begin{bmatrix} b_1 \\ b_4 \end{bmatrix} = 0, \quad \begin{bmatrix} i^j & 1 \\ -1 & i^j \end{bmatrix} \begin{bmatrix} b_2 \\ b_3 \end{bmatrix} = 0.$$

Therefore, unless $j = 1$ or $j = 3$, $f(x, y, z, t) = 0$. Even if $j = 1$ or $j = 3$, the condition $f_{E_3^{-1}} \sim f$ implies $f = 0$, because $\{i, -i\} \cap \{\nu_1, \nu_2, \nu_1^3, \nu_2^3\} = \emptyset$.

Remark 3.5. *Let $G = \text{Aut}(V(x^2y + y^2z + z^2t + t^2x))$, Σ the field automorphism of $\mathbf{Q}(\varepsilon)$ such that $\varepsilon^\Sigma = \varepsilon^2$.*

(1) *Denoting by e_i be the i -th column of the unit matrix E_4 , let $J = [e_2, e_3, e_4, e_1]$. Then $(J) \in G = G(1)'$.*

(2) Define $I' \in GL_4(k)$ as I , using $\alpha' = \frac{\sqrt{5}+1}{2}$, $\beta' = \alpha'^2$ and $\gamma' = -\alpha'$ in place of α , β and γ , respectively. Then $H^\Sigma = H^2$ and $I^\Sigma = I'$, for $\sqrt{5}^\Sigma = -\sqrt{5}$. On the other hand $JHJ^{-1} = H^2$ and $JIJ^{-1} = \beta I'$. Consequently there exists uniquely a group isomorphism $\psi : \mathbf{S}_5 \rightarrow PGL_4(k)$ such that $\psi((12345)) = (H^2)$ and $\psi((12)) = (I')$. The projective representations of \mathbf{S}_5 ψ and φ_1 are equivalent.

References

- [1] Hosoh T.: Automorphism groups of cubic surfaces, J. Algebra **192**, 651–677 (1997).
- [2] Kaneta, H., Marcugini, S. and Pambianco, F.: On arcs and curves with many automorphisms, Mediterr. J. Math. **2**, 71–102 (2005).
- [3] Maschke H.: Bestimmung aller ternären und quaternären Collineationsgruppe, welche mit symmetrischen und alternierenden Buchstabenvertauschungsgruppen holoedrisch isomorph sind, Math. Ann. **51**, 253–298 (1899).
- [4] Matsumura H. and Monsky P.: On the automorphisms of hypersurfaces, J. Math. Kyoto Univ. **3-3**, 347–361 (1964).
- [5] Pambianco F.: Characterization of the Fermat curve as the most symmetric nonsingular plane curve, Mathematische Zeitschrift, 2014, DOI 10.1007/s00209-014-1288-4.
- [6] Shioda, T.: Arithmetic and geometry of the Fermat curve, Algebraic Seminar (Singapore 1987), World Sci. Publications, Singapore(1988).
- [7] Suzuki M.: Group Theory I, Springer 1982.