

K3 SURFACES, MODULAR FORMS, AND NON-GEOMETRIC HETEROTIC COMPACTIFICATIONS

ANDREAS MALMENDIER AND DAVID R. MORRISON

ABSTRACT. We construct non-geometric compactifications by using the F-theory dual of the heterotic string compactified on a two-torus, together with a close connection between Siegel modular forms of genus two and the equations of certain K3 surfaces. The modular group mixes together the Kähler, complex structure, and Wilson line moduli of the torus yielding weakly coupled heterotic string compactifications which have no large radius interpretation.

INTRODUCTION

The traditional approach to producing low-dimensional physical models out of high-dimensional theories such as the string theories and M-theory has been to use a specific geometric compactification of the “extra” dimensions and derive an effective description of the lower-dimensional theory from the choice of geometric compactification. However, it has long been recognized that there are other possibilities: for example, one can couple perturbative string theory to an arbitrary superconformal two-dimensional theory (geometric or not) to obtain an effective perturbative string compactification in lower dimensions.

One way of making an analogous construction in non-perturbative string theory is to exploit the nonperturbative duality transformations which relate various compactified string theories (and M-theory) to each other. This idea was the basis of the construction of F-theory [49], and more recently was used in constructions involving the type II theories [28] and the heterotic theories [38]. We pursue a further non-geometric construction of heterotic compactifications in this paper.

Our construction of such models relies on a very concrete relationship between modular forms on the moduli space of certain K3 surfaces and the equations of those K3 surfaces [35, 16, 17]. The K3 surfaces in question have a large collection of algebraic curve classes on them, generating a lattice known as $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$. The presence of these classes restricts the form of moduli space,¹ which turns out to be a space admitting

2010 *Mathematics Subject Classification.* Primary 81T30, Secondary 11F03, 14J28, 14J81.

Key words and phrases. K3 surface, heterotic string, F-theory, Siegel modular forms.

¹We use the term “moduli space” here as mathematicians do, denoting the parameter space of the geometric objects.

modular forms. The modular forms in question are the Siegel modular forms of genus two, which have previously made some appearances in the study of string compactification.²

The close connection between modular forms and equations allows us to mimic the basic F-theory construction, and build an interesting class of non-geometric heterotic compactifications which have duals described in terms of families of K3 surfaces. The starting point is the heterotic string compactified on a torus, and we exploit the non-perturbative duality symmetries which this theory possesses. We give the construction in considerable detail.

The paper begins with a review of F-theory in Section 1 and then proceeds to give a construction of non-geometric heterotic compactifications in Section 2. These compactifications require certain $5 + 1$ -dimensional soliton solutions (serving as sources for scalar fields) whose heterotic construction is discussed in Section 3.

After a brief digression in Section 4 to interpret our construction in the context of the $\mathfrak{so}(32)$ -heterotic string, we specialize in Section 5 to the case of compactifications to six dimensions. There we find a surprise: although non-geometric techniques were used for the construction, the models we obtain are not new, but were already known (at least in dual form). We conclude the paper with a discussion of this surprise and its implications.

1. REVIEW OF F-THEORY

One of the fundamental interpretations of F-theory is in terms of the type IIB string, where it depends on three ingredients: an $\mathrm{SL}_2(\mathbb{Z})$ symmetry of the theory, a complex scalar field τ (the axio-dilaton) with positive imaginary part (in an appropriate normalization) on which $\mathrm{SL}_2(\mathbb{Z})$ acts by fractional linear transformations, and D7-branes, which serve as a source for the multi-valuedness of τ if τ is allowed to vary.

In a standard compactification of the type IIB string, τ is a constant and D7-branes are absent. Vafa's idea in proposing F-theory [49] was to simultaneously allow a variable τ and the D7-brane sources, arriving at a new class of models in which the string coupling is never weak.

Since we cannot use the axio-dilaton τ directly in these models, it would be natural to identify the physically relevant quantity with $\mathbb{H}/\mathrm{SL}_2(\mathbb{Z})$ (where \mathbb{H} denotes the complex upper half-plane), but this turns out to be slightly too simplistic. To obtain the full range of F-theory models, we need instead to consider some functions of τ which are not invariant under $\mathrm{SL}_2(\mathbb{Z})$, but rather transform in a specific way. A function $f(\tau)$ which satisfies

$$(1.1) \quad f\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^m f(\tau)$$

for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ is a *modular form of weight m for $\mathrm{SL}_2(\mathbb{Z})$* and such forms turn out to provide the flexibility we need for F-theory.

²For an early appearance, see [18] and references therein. More recently [36, 13], U-duality of type IIB compactifications on K3 surfaces and the connection to modular forms of genus two was used to construct non-geometric compactifications analogous to the ones studied in this paper.

A simple way to write down some modular forms of even weight $m = 2k$ for $\mathrm{SL}_2(\mathbb{Z})$ is to use what are called *normalized Eisenstein series*, defined as

$$(1.2) \quad E_{2k}(\tau) = \frac{1}{2\zeta(2k)} \sum_{(0,0) \neq (m,n) \in \mathbb{Z}^2} \frac{1}{(m\tau + n)^{2k}},$$

where $\zeta(2k) = \sum_{n \geq 1} n^{-2k}$ is Riemann's zeta function. These normalized Eisenstein series have a Fourier expansion (in $q = e^{2\pi i \tau}$) of the form

$$(1.3) \quad E_{2k}(\tau) = 1 + O(q),$$

which is the reason for including the normalization factor. It is known that $E_4(\tau)$ and $E_6(\tau)$ generate the entire ring of modular forms for $\mathrm{SL}_2(\mathbb{Z})$. The combination

$$(1.4) \quad \begin{aligned} \Delta_{12}(\tau) &= 4 \left(-\frac{1}{3} E_4(\tau) \right)^3 + 27 \left(-\frac{2}{27} E_6(\tau) \right)^2 \\ &= -\frac{4}{27} E_4(\tau)^3 + \frac{4}{27} E_6(\tau)^2 \end{aligned}$$

plays a special role in the theory.³ In particular, if we compactify the parameter space $\mathrm{SL}_2(\mathbb{Z})$ to $\overline{\mathrm{SL}_2(\mathbb{Z})}$, then $\Delta_{12}(\tau)$ extends to the compactification and vanishes on the boundary (which corresponds to the $q \rightarrow 0$ limit).

An F-theory compactification (regarded as a compactification of the type IIB string with variable axio-dilaton) takes as its starting point a compact space W , a complex line bundle \mathcal{L} on W and sections $f(w)$ and $g(w)$ of the associated bundles $\mathcal{L}^{\otimes 4}$ and $\mathcal{L}^{\otimes 6}$. Then there is a (possibly non-supersymmetric) F-theory model with a variable τ function and $\mathrm{SL}_2(\mathbb{Z})$ symmetry, in which $f(w)$ is identified with $-\frac{1}{3}E_4(\tau)$ and $g(w)$ is identified with $-\frac{2}{27}E_6(\tau)$. One must also insert seven-branes of various kinds along the zeros of

$$(1.5) \quad \Delta(w) := 4f(w)^3 + 27g(w)^2.$$

The geometry behind this construction is a beautiful story from 19th-century mathematics: the Weierstrass \wp -function. In order to define a doubly-periodic meromorphic function in the complex plane (with periods 1 and τ), Weierstrass introduced a function of $z \in \mathbb{C}$ and τ :

$$(1.6) \quad \wp(z, \tau) = \frac{1}{z^2} + \sum_{(m,n) \neq (0,0)} \left(\frac{1}{(z - m\tau - n)^2} - \frac{1}{(m\tau + n)^2} \right).$$

This has a Laurent expansion (using the normalized Eisenstein series as well as the special values $\zeta(4) = \pi^4/90$ and $\zeta(6) = \pi^6/945$):

$$(1.7) \quad \begin{aligned} \wp(z, \tau) &= z^{-2} + 6\zeta(4)E_4(\tau)z^2 + 10\zeta(6)E_6(\tau)z^4 + O(z^6), \\ &= z^{-2} + \frac{\pi^4}{15}E_4(\tau)z^2 + \frac{2\pi^6}{189}E_6(\tau)z^4 + O(z^6), \end{aligned}$$

³The convention in F-theory slightly differs from the one used in number theory, where the discriminant is identified with the 24th power of the Ramanujan tau function, defined to be $\eta^{24}(\tau) = q \prod_{n \geq 1} (1 - q^n)^{24}$. With our conventions for Δ_{12} , we have $\Delta_{12}(\tau) = -2^8 \eta^{24}(\tau)$.

from which Laurent expansions for $(\wp(z, \tau)')^2$ and $\wp(z, \tau)^3$ can be calculated:

$$(1.8) \quad (\wp(z, \tau)')^2 = 4z^{-6} - \frac{8\pi^4}{15} E_4(\tau) z^{-2} - \frac{32\pi^6}{189} E_6(\tau) + O(z),$$

$$(1.9) \quad \wp(z, \tau)^3 = z^{-6} + \frac{\pi^4}{5} E_4(\tau) z^{-1} + \frac{6\pi^6}{189} E_6(\tau) + O(z).$$

It follows that

$$(1.10) \quad (\wp(z, \tau)')^2 - 4\wp(z, \tau)^3 + \frac{4\pi^4}{3} E_4(\tau) \wp(z, \tau) = -\frac{8\pi^6}{27} E_6(\tau) + O(z) = -\frac{8\pi^6}{27} E_6(\tau).$$

(This is an exact expression since the left hand side is an entire holomorphic function which is bounded since it is doubly-periodic, and hence constant.) If we set $x := \frac{1}{\pi^2} \wp(z, \tau)$ and $y := \frac{1}{2\pi^3} \wp(z, \tau)'$, we find an equation for the elliptic curve with modular parameter τ :

$$(1.11) \quad y^2 = x^3 - \frac{1}{3} E_4(\tau) x - \frac{2}{27} E_6(\tau).$$

Conversely, if we start from an elliptic curve with an equation of the form

$$(1.12) \quad y^2 = x^3 + fx + g$$

which is nonsingular, then we can recover τ up to $\text{SL}_2(\mathbb{Z})$ transformation as

$$(1.13) \quad \tau = \frac{\int_{\gamma_2} \frac{dx}{\sqrt{x^3 + fx + g}}}{\int_{\gamma_1} \frac{dx}{\sqrt{x^3 + fx + g}}},$$

where (γ_1, γ_2) is an oriented basis of the first homology of the elliptic curve, such that $f = -\frac{\lambda^4}{3} E_4(\tau)$, $g = -\frac{2\lambda^6}{27} E_6(\tau)$ for some nonvanishing scale factor λ . The condition for nonsingularity of (1.12) is that the quantity

$$(1.14) \quad \Delta := 4f^3 + 27g^3$$

does not vanish. It is this close connection between geometry and modular forms which allows the construction of families of elliptic curves with certain knowledge of the behavior of τ in such families.

To understand when the corresponding F-theory models are supersymmetric, we follow the duality between F-theory and M-theory. That duality is based on a key fact: Compactifying M-theory on a torus whose complex structure is labeled by τ and whose area is A gives a model dual to the type IIB string compactified on a circle of radius⁴ $A^{-3/4}$ whose axio-dilaton has value τ [48, 2]. This then gives a connection between the F-theory construction and a dual geometric compactification of M-theory: if the F-theory model is further compactified on S^1 (which can be done without breaking any supersymmetry that might be present), a model will be obtained which is dual to M-theory compactified on the total space of the family

$$(1.15) \quad y^2 = x^3 + f(w)x + g(w)$$

⁴The radius is measured in Einstein frame.

of elliptic curves over W . (The insertions of seven-branes in the F-theory model go over to singular elliptic fibers in the M-theory model which may require special treatment, but for generic sections f and g the total space of the family is nonsingular and the compactification makes sense as it stands.)

One can then ask whether the geometric M-theory model breaks or preserves supersymmetry, and the answer is known: supersymmetry is preserved exactly when the total space of the family is a Calabi–Yau manifold, which happens exactly when the line bundle \mathcal{L} is the anti-canonical bundle of the base, i.e., $\mathcal{L} = \mathcal{O}_W(-K_W)$. In this way, we recover the familiar conditions for a supersymmetric F-theory compactification.

To complete the story, we need to know what types of seven-branes need to be inserted. The answer here comes from algebraic geometry, through work of Kodaira [34] and Néron [45] which classifies the possible singular limits in one-parameter families of elliptic curves and thereby gives a catalog of the different types of seven-branes which must be inserted. This catalog is by now well-known, but we will reproduce it in Table 1, in which labels for the types of seven-branes are given using Kodaira’s notation. The type of brane depends on the orders of vanishing of f , g , and Δ at the singular point P , and determines both the type of singularity which appears in the M-theory dual, and the transformation in $\mathrm{SL}_2(\mathbb{Z})$ which describes how τ changes when the singular point is encircled. The I_n type corresponds to a stack of n D7-branes, while the I_n^* type corresponds to a stack of $n+4$ D7-branes on top of an orientifold plane. The last line of the table (labeled “non-minimal”) can be avoided by a suitable choice of line bundle for the Weierstrass model.

2. NON-GEOMETRIC HETEROTIC MODELS

By analogy, we now wish to study as our basic theory the heterotic string compactified on T^2 to produce an eight-dimensional effective theory. (This will be our analogue of the type IIB string in the previous section.) This effective theory has a complex scalar field which, after symmetries are taken into account, takes its values in the Narain space⁵ [43]

$$(2.1) \quad \mathcal{D}_{2,18}/O(\Lambda^{2,18})$$

which is the quotient of the symmetric space for $O(2, 18)$,

$$(2.2) \quad \mathcal{D}_{2,18} := (O(2) \times O(18)) \backslash O(2, 18),$$

by the automorphism group $O(\Lambda^{2,18})$ of the unique integral even unimodular lattice $\Lambda^{2,18}$ of signature $(2, 18)$. (This discrete group is sometimes called $O(2, 18, \mathbb{Z})$.) In an appropriate limit, this space decomposes as a product of spaces parameterizing the Kähler and complex structures on T^2 as well as Wilson line expectation values around the two generators of $\pi_1(T^2)$ (see [44]). However, that decomposition is only preserved by a parabolic subgroup $\Gamma \subset O(\Lambda^{2,18})$, which is much smaller. Letting the moduli of the entire space vary arbitrarily (i.e., employing the full $O(\Lambda^{2,18})$ symmetry) will produce a

⁵To better match both the algebraic geometry and modular forms literature, we take the Narain lattices for compactifications of the heterotic string on a d -torus to have signature $(d, 16 + d)$, which is the opposite of the usual convention in string theory.

brane type	$\text{ord}_P(f)$	$\text{ord}_P(g)$	$\text{ord}_P(\Delta)$	singularity	transformation
I_0	≥ 0	≥ 0	0	none	$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
$I_n, n \geq 1$	0	0	n	A_{n-1}	$\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$
II	≥ 1	1	2	none	$\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$
III	1	≥ 2	3	A_1	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$
IV	≥ 2	2	4	A_2	$\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$
I_0^*	≥ 2	≥ 3	6	D_4	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$
$I_n^*, n \geq 1$	2	3	$n+6$	D_{n+4}	$\begin{pmatrix} -1 & -n \\ 0 & -1 \end{pmatrix}$
IV^*	≥ 3	4	8	E_6	$\begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$
III^*	3	≥ 5	9	E_7	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$
II^*	≥ 4	5	10	E_8	$\begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$
non-minimal	≥ 4	≥ 6	≥ 12	non-canonical	–

TABLE 1. Kodaira–Néron classification of singular fibers and monodromy

compactification which has a right to be called non-geometric, because the Kähler and complex structures on T^2 , and the Wilson line values, are not distinguished under the $O(\Lambda^{2,18})$ -equivalences but instead are mingled together. With no Kähler class, we lose track of geometry.⁶

The construction we will give of non-geometric heterotic compactifications actually uses an index 2 subgroup $O^+(\Lambda^{2,18}) \subset O(\Lambda^{2,18})$ defined by

$$(2.3) \quad O^+(\Lambda^{2,18}) := O^+(2, 18) \cap O(\Lambda^{2,18}),$$

where $O^+(p, q)$ denotes the subgroup of $O(p, q)$ preserving the orientation on positive p -planes. The group $O^+(\Lambda^{2,18})$ is the maximum subgroup of $O(\Lambda^{2,18})$ whose action preserves the complex structure on the symmetric space, and thus is the maximal subgroup for which modular forms can be holomorphic. The corresponding quotient

$$(2.4) \quad \mathcal{D}_{2,18}/O^+(\Lambda^{2,18})$$

is a degree two cover of the Narain moduli space. The group $O^+(\Lambda^{2,18})$ is still large enough to thoroughly mix the Kähler, complex, and Wilson line moduli.

The quotient space (2.4) is the parameter space for elliptically fibered K3 surfaces with a section: this is a statement of the duality between F-theory and the heterotic string

⁶Note that the heterotic dilaton is *not* affected by this group action, so weakly coupled models of these non-geometric heterotic strings will exist.

in eight dimensions [49], and the identification of the discrete group for this moduli problem as $O^+(\Lambda^{2,18})$ is well-known in the mathematics literature (see, for example, [26]).⁷ To use these K3 surfaces in a similar way to the way that elliptic curves were used in constructing F-theory, we would need a close connection between $O^+(\Lambda^{2,18})$ -modular forms and the equations of the corresponding elliptically fibered K3 surfaces; unfortunately such a connection is not known.

However, by making a simple and natural restriction on our heterotic theories, we can find and exploit such a connection. Namely, let us consider heterotic models with only a single nonzero Wilson line expectation value. For definiteness, we restrict to the $\mathfrak{e}_8 \oplus \mathfrak{e}_8$ heterotic string, and note that asking for an unbroken gauge algebra of $\mathfrak{e}_8 \oplus \mathfrak{e}_7$ will ensure that only a single Wilson line expectation value is nonzero. (There is a similar story for the $\mathfrak{so}(32)$ string which we will describe in Section 4.)

Let $L^{2,3}$ be the lattice of signature $(2, 3)$ which is the orthogonal complement of $E_8(-1) \oplus E_7(-1)$ in $\Lambda^{2,18}$. By insisting that the Wilson lines values associated to the⁸ $E_8(-1) \oplus E_7(-1)$ sublattice be trivial (which leaves the algebra $\mathfrak{e}_8 \oplus \mathfrak{e}_7$ unbroken), we restrict to those heterotic vacua parameterized by the space

$$(2.5) \quad \mathcal{D}_{2,3}/O(L^{2,3}).$$

The corresponding degree two cover is

$$(2.6) \quad \mathcal{D}_{2,3}/O^+(L^{2,3}),$$

and this space parameterizes elliptically fibered K3 surfaces with section which have one fiber of Kodaira type III^* or worse and another fiber of Kodaira type precisely⁹ II^* . Such K3 surfaces contain the lattice $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ inside their Néron–Severi lattice, and are often referred to as “lattice-polarized K3 surfaces”. The $\Lambda^{1,1}$ summand is generated by the classes of the fiber and the section of the elliptic fibration.¹⁰

As we will describe below, the modular forms for $O^+(L^{2,3})$ have the desired property: there is a close geometric connection to the corresponding lattice-polarized K3 surfaces. (A similar picture was developed in earlier work in the case of no nontrivial Wilson line expectation values, using modular forms for $O^+(\Lambda^{2,2})$ [38].)

Let \mathbb{H}_g denote the Siegel upper half-space of genus g , on which the Siegel modular group $\mathrm{Sp}_{2g}(\mathbb{Z})$ acts. As explained in Appendix A, for $g = 2$ there is a homomorphism $\mathrm{Sp}_4(\mathbb{R}) \rightarrow O^+(2, 3)$ which induces an isomorphism

$$(2.7) \quad \mathbb{H}_2 \cong \mathcal{D}_{2,3}.$$

⁷The slight mismatch in duality groups between $O(\Lambda^{d,16+d})$ for the heterotic string compactified on T^d and $O^+(\Lambda^{d,16+d})$ for the dual theory occurs for the type I' dual (interpreted as a real K3 surface) when $d = 1$ [14], the F-theory dual when $d = 2$ as indicated here, the M-theory dual when $d = 3$ (described in [6] and based on [12, 37, 21]), and the type IIA dual when $d = 4$ [41]. It would be interesting to have a more complete understanding of this mismatch.

⁸The sign change on E_8 and E_7 is due to our sign conventions about the Narain lattice.

⁹As we will see in Section 5, assuming that this fiber is *precisely* type II^* avoids “pointlike instantons” on the heterotic dual after compactification to dimension six or below, at least for general moduli.

¹⁰For the lattice embedding of $\Lambda^{1,1}$ to correspond to an elliptic fibration with section we also need to require that its image in the Néron–Severi lattice contains a pseudo-ample class.

By a result of Vinberg [50] reviewed in Appendix A, the image of $\mathrm{Sp}_4(\mathbb{Z}) \rightarrow O^+(L^{2,3})$ is a subgroup of index 2, and the ring of $O^+(L^{2,3})$ -modular forms turns out to correspond to the ring of Siegel modular forms with $g = 2$ of even weight.¹¹ Igusa [31] showed that this latter ring is a polynomial ring in four free generators of degrees 4, 6, 10 and 12. We explicitly describe Igusa's generators ψ_4 , ψ_6 , χ_{10} and χ_{12} in Section B.2. (Igusa later showed [32] that for the full ring of $\mathrm{Sp}_4(\mathbb{Z})$ -modular forms, one needs an additional generator χ_{35} , also described in Section B.2, which is algebraically dependent on the others. In fact, χ_{35}^2 is an explicit polynomial in ψ_4 , ψ_6 , χ_{10} and χ_{12} which is given in (B.16).)

The key geometric fact, due in different forms to Kumar [35] and to Clingher–Doran [16, 17], is the equation for an elliptically fibered K3 surface whose periods give the point $\underline{\tau}$ in the Siegel upper halfspace \mathbb{H}_2 , with the coefficients in the equation being Siegel modular forms of even weight. (This is analogous to the Weierstrass equation for the elliptic curve $\mathbb{C}/\langle 1, \tau \rangle$ with coefficients being Eisenstein series in τ). That equation is:

$$(2.8) \quad y^2 = x^3 - t^3 \left(\frac{1}{48} \psi_4(\underline{\tau}) t + 4 \chi_{10}(\underline{\tau}) \right) x + t^5 \left(t^2 - \frac{1}{864} \psi_6(\underline{\tau}) t + \chi_{12}(\underline{\tau}) \right).$$

Just as in the elliptic curve case, the statement has two parts: starting from $\underline{\tau}$, we obtain the equation of a K3 surface (2.8). But conversely, if we start with a K3 surface S with an equation of the form¹²

$$(2.9) \quad y^2 = x^3 + a t^4 x + b t^6 + c t^3 x + d t^5 + t^7,$$

and we determine a point in $\mathcal{D}_{2,3}$ by calculating the periods of the holomorphic 2-form on S over a basis of the orthogonal complement of $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ in $H^2(S, \mathbb{Z})$ (which in turn determines $\underline{\tau} \in \mathbb{H}_2$ using the isomorphism (2.7)), then for some nonzero scale factor λ ,

$$(2.10) \quad a = -\frac{\lambda^4}{48} \psi_4(\underline{\tau}), \quad b = -\frac{\lambda^6}{864} \psi_6(\underline{\tau}), \quad c = -4 \lambda^{10} \chi_{10}(\underline{\tau}), \quad d = \lambda^{12} \chi_{12}(\underline{\tau}).$$

We verify in Appendix C that the K3 surface defined by (2.8) agrees with the ones found by Kumar and by Clingher–Doran.

The strategy for constructing a non-geometric heterotic compactification is now clear. Start with compact manifold Z and a line bundle Λ on Z . Pick sections $a(z)$, $b(z)$,

¹¹Note that because the groups are different, families of genus two curves (such as were used in [36, 13]) are not equivalent to the families of elliptic K3 surfaces needed for our construction.

¹²The reader may wonder why the coefficients of x^3 , y^2 , and t^7 have been set equal to 1 in (2.9). The choice of coefficient 1 for x^3 and y^2 is a familiar one, and derives from an analysis by Deligne [19] of families of elliptic curves: assuming that all fibers are generalized elliptic curves of an appropriate kind, it follows that the coefficients of x^3 and y^2 never vanish, and then by a change of coordinates these coefficients can be set to 1. The story for t^7 is similar: we are assuming that the Kodaira fiber at $t = \infty$ is exactly II^* , and this implies that the t^7 term in the equation must always be present. Thus, we could allow a coefficient α for t^7 but it would never be allowed to vanish; as a consequence, the change of coordinates $(x, y, t) \mapsto (x/\alpha^2, y/\alpha^3, t/\alpha)$ would be well-defined, and would map $x^3 - y^2 + \alpha t^7$ to $\alpha^{-6}(x^3 - y^2 + t^7)$. In other words, by making such a change of coordinates and then rescaling the entire equation by α^6 , we may assume that the coefficient of t^7 is 1.

$c(z)$, and $d(z)$ of $\Lambda^{\otimes 4}$, $\Lambda^{\otimes 6}$, $\Lambda^{\otimes 10}$, and $\Lambda^{\otimes 12}$, respectively. Then there is a non-geometric heterotic compactification on Z with variable τ and $O^+(L^{2,3})$ symmetry for which

$$(2.11) \quad \begin{aligned} a(z) &= -\frac{1}{48} \psi_4(\tau), \\ b(z) &= -\frac{1}{864} \psi_6(\tau), \\ c(z) &= -4 \chi_{10}(\tau), \\ d(z) &= \chi_{12}(\tau). \end{aligned}$$

(We can eliminate the scale factor λ , if any, by making a change of coordinates $(x, y, t) \mapsto (\lambda^{14}x, \lambda^{21}y, \lambda^6t)$.) Appropriate five-branes must be inserted on Z as dictated by the geometry of the corresponding family of K3 surfaces

$$(2.12) \quad y^2 = x^3 + a(z)t^4x + b(z)t^6 + c(z)t^3x + d(z)t^5 + t^7.$$

We will explore these five-branes in the next section.

To understand when the non-geometric heterotic compactifications we have constructed are supersymmetric, we follow the duality between the heterotic string and F-theory. The heterotic compactification on T^2 with parameter $\tau \in \mathbb{H}_2$ is dual to the F-theory compactification on the elliptically fibered K3 surface S_τ defined by (2.8), where t is a local coordinate on the base \mathbb{P}^1 of the elliptic fibration. Note that at $t = \infty$, S_τ has a Kodaira fiber of type precisely II^* : it can be no worse because the coefficient of t^7 in (2.8) is 1. At $t = 0$, there is a Kodaira fiber of type III^* or worse.

When $a(z)$, $b(z)$, $c(z)$, and $d(z)$ are sections of line bundles over Z , we wish to determine whether F-theory compactified on the elliptically fibered manifold (2.12) is supersymmetric, and this in turn depends on whether the total space defined by (2.12) is itself a Calabi–Yau manifold. The base of the elliptic fibration on the total space is a \mathbb{P}^1 -bundle $\pi : W \rightarrow Z$ which takes the form $W = \mathbb{P}(\mathcal{O} \oplus \mathcal{M})$ for some line bundle \mathcal{M} that coincides with the normal bundle of $\Sigma_0 := \{t = 0\}$ in W . Restricting the various terms in (2.12) to Σ_0 , we find relations $\Lambda^{\otimes 4} \otimes \mathcal{M}^{\otimes 4} = \Lambda^{\otimes 10} \otimes \mathcal{M}^{\otimes 3} = (\mathcal{L}|_{\Sigma_0})^{\otimes 4}$ and $\mathcal{M}^{\otimes 7} = \Lambda^{\otimes 6} \otimes \mathcal{M}^{\otimes 6} = \Lambda^{\otimes 12} \otimes \mathcal{M}^{\otimes 5} = (\mathcal{L}|_{\Sigma_0})^{\otimes 6}$. It follows that that $\mathcal{M} = \Lambda^{\otimes 6}$ and $\mathcal{L}|_{\Sigma_0} = \Lambda^{\otimes 7}$ (up to torsion). In other words, our \mathbb{P}^1 -bundle must take the form $W = \mathbb{P}(\mathcal{O} \oplus \Lambda^{\otimes 6})$. This property can be traced back to the fact that the coefficient of t^7 in (2.12) is 1.

Now to check the condition for supersymmetry, note that

$$(2.13) \quad -K_W = \Sigma_0 + \Sigma_\infty + \pi^{-1}(-K_Z),$$

where $\Sigma_\infty := \{t = \infty\}$. Since Σ_0 and Σ_∞ are disjoint, it follows that the condition for supersymmetry $\mathcal{L} = \mathcal{O}_W(-K_W)$ is equivalent to $\Lambda = \mathcal{O}_Z(-K_Z)$.

Let us briefly comment on the relationship of our construction with the appearance of Siegel modular forms in string compactifications involving the “STUV” model, as described in [18] and the references therein. If we take the mirror of our family of lattice-polarized K3 surfaces [6, 20], we will obtain a family of K3 surfaces depending

on 17 complex parameters whose quantum Kähler moduli space is $\mathcal{D}_{2,3}/O^+(L^{2,3})$. The K3 surfaces in the new family all contain a lattice $L^{1,2}$ within $H^{1,1}$ with the property that $L^{2,3} \cong \Lambda^{1,1} \oplus L^{1,2}$. If X is a Calabi–Yau threefold which has a one-parameter family of such K3 surfaces on it, then the $(1,1)$ classes on X also include the lattice $L^{1,2}$. Type IIA string theory compactified on X is the dual theory of the heterotic STUV model, as discussed in [18] and elsewhere. It is natural that quantum corrections of this gravitational theory would respect the symmetry group $O^+(L^{2,3})$ and so would turn out to be related to Siegel modular forms as well.

3. FIVE-BRANES

The base W of an elliptic fibration maps naturally to the compactification $\overline{\mathbb{H}/\mathrm{SL}_2(\mathbb{Z})}$ of the parameter space $\mathbb{H}/\mathrm{SL}_2(\mathbb{Z})$, and if this map is nonconstant, there must be singular fibers (at which the j -invariant approaches ∞). In fact, for a generic elliptic fibration, all seven-branes will have $j \rightarrow \infty$, and those correspond to familiar seven-brane constructions in type IIB string theory (D7-branes, possibly combined with orientifold planes).

The situation for fibrations of lattice-polarized K3 surfaces is very different. There is a Satake–Baily–Borel compactification [47, 8] $\overline{\mathcal{D}_{2,3}/O^+(L^{2,3})}$ of the parameter space whose boundary has codimension two, and this implies that a one-parameter family of lattice-polarized K3 surfaces *need not reach the boundary!* That would suggest that it might be possible to have a family which never degenerates (i.e., with no brane insertions needed), but this is not the case: there are codimension one loci where some elements of $O^+(L^{2,3})$ have fixed points, and there must always be branes associated with these fixed loci.

To find group elements with fixed points, note that a reflection in a lattice element of square -2 has a fixed locus of codimension one, belongs to $O^+(L^{2,3})$, and does not belong to $SO^+(L^{2,3}) \cong \mathrm{Sp}_4(\mathbb{Z})$. As a consequence, such reflections must act as -1 on the $\mathrm{Sp}_4(\mathbb{Z})$ -modular forms of odd weight, and the fixed locus of any such reflection must be contained in the vanishing locus of any $\mathrm{Sp}_4(\mathbb{Z})$ -modular form of odd weight. The modular forms of odd weight are generated by Igusa’s form χ_{35} , so that form must vanish along the fixed loci of our reflections.

From the point of view of K3 geometry, if the periods are preserved by the reflection in δ with $\delta^2 = -2$, then δ must belong to the Néron–Severi lattice of the K3 surface. That is, the lattice $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ must be enlarged by adjoining δ . It is not hard to show (using methods of [46], for example), that there are only two ways this enlargement can happen (if we have adjoined a single element only): either the lattice is extended to $\Lambda^{1,1} \oplus E_8(-1) \oplus E_8(-1)$ or it is extended to $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1) \oplus \langle -2 \rangle$. In the former case, the fibers in the elliptic fibration become II^* , II^* and $4 I_1$, whereas in the latter case, the fibers become II^* , III^* , I_2 and $4 I_1$.

If we start with an elliptically fibered K3 surface (2.9), then it is easy to see what the condition is for the first enhancement: we want the fiber at $t = 0$ to go from type III^* to type II^* , and this is achieved by setting $c = 0$.

To see the second enhancement requires a computation. Starting with (2.9), we compute the discriminant of the elliptic fibration to be

$$(3.1) \quad \Delta = t^9 (4(at + c)^3 + 27t(t^2 + bt + d)^2)$$

The zeros of Δ/t^9 represent the location of the I_1 fibers, so to find out when they coincide, we calculate the discriminant of that polynomial of degree 5 in t (which will vanish precisely when there are multiple roots). That discriminant turns out to take the form

$$(3.2) \quad 2^8 3^{12} \ell(a, b, c, d)^3 q(a, b, c, d),$$

where, if we assign weights 4, 6, 10, 12 to a, b, c, d , respectively, then ℓ is the polynomial of weighted degree 20

$$(3.3) \quad \ell(a, b, c, d) := a^2 d - a b c + c^2,$$

and q is the polynomial of weighted degree 60

$$(3.4) \quad \begin{aligned} q(a, b, c, d) := & 11664 d^5 + 864 a^3 d^4 - 5832 b^2 d^4 + 16 a^6 d^3 + 216 a^3 b^2 d^3 \\ & - 2592 a^2 b c d^3 + 16200 a c^2 d^3 + 729 b^4 d^3 + 888 a^4 c^2 d^2 - 5670 a b^2 c^2 d^2 \\ & - 13500 b c^3 d^2 + 16 a^7 c^2 d + 216 a^4 b^2 c^2 d - 3420 a^3 b c^3 d + 4125 a^2 c^4 d \\ & + 729 a b^4 c^2 d + 6075 b^3 c^3 d - 16 a^6 b c^3 + 16 a^5 c^4 - 216 a^3 b^3 c^3 \\ & + 2700 a^2 b^2 c^4 - 5625 a b c^5 - 729 b^5 c^3 + 3125 c^6 \end{aligned}$$

(which we computed directly using computer algebra).

The role of the polynomial ℓ is easy to see: it vanishes on precisely those K3 surfaces for which f and g have a common zero (at $t = -c/a$). Those are cases in which two I_1 's are replaced by a fiber of type II , but such a change does not affect the lattice or the gauge algebra and so these are not the K3 surfaces we are looking for.

Since the two lattice enhancements occur at $c = 0$ and $q(a, b, c, d) = 0$, we predict that $c \cdot q(a, b, c, d)$ should vanish along the locus where there is some degeneration. Indeed it turns out (as verified in Section B.2) that

$$(3.5) \quad q \left(-\frac{1}{48} \psi_4, -\frac{1}{864} \psi_6, -4 \chi_{10}, \chi_{12} \right) = 2^{-8} \chi_{35}^2 / \chi_{10},$$

confirming the prediction.

Thus, a generic non-geometric compactification constructed from these lattice-polarized K3 surfaces will have two types of five-branes, corresponding to¹³ $c = 0$ and $q(a, b, c, d) = 0$. From the heterotic side, these five-brane solitons are easy to see. When $q(a, b, c, d) = 0$, we have an additional gauge symmetry enhancement to include $\mathfrak{su}(2)$, and the parameters of the theory include a Coulomb branch for that gauge theory on which the Weyl group $W_{\mathfrak{su}(2)} = \mathbb{Z}_2$ acts. There is thus a five-brane solution in which the field has a \mathbb{Z}_2 ambiguity encircling the location in the moduli space of enhanced gauge symmetry.

¹³The loci $\{c = 0\}$ and $\{q(a, b, c, d) = 0\}$ correspond to the well-studied Humbert surfaces H_1 and H_4 described in Appendix B.

The other five-brane solution is similar: at $c = 0$, there is an enhancement from \mathfrak{e}_7 to \mathfrak{e}_8 gauge symmetry, and a similar \mathbb{Z}_2 acts on the moduli space, leading to a solution with a \mathbb{Z}_2 ambiguity. These two brane solutions are the analogue of the simplest brane (a single D7-brane) in F-theory.

Finding a complete catalog of five-brane solutions for this theory is quite challenging. As we explain in Appendix A.2, the parameter space $\mathcal{D}_{2,3}/O^+(L^{2,3})$ for our construction is closely related to some other moduli spaces: the moduli space of homogeneous sextics in two variables, the moduli space of Abelian surfaces, and the moduli space of curves of genus two. A version of Kodaira's classification was given for curves of genus two by Namikawa and Ueno [42] and this can in principle be used to give a classification of degenerations of these lattice-polarized K3 surfaces. We illustrate how this works in a number of interesting cases in Appendix D.

4. THE $\mathfrak{so}(32)$ HETEROTIC STRING

It turns out that the total space of a lattice-polarized K3 surface of the form (2.9) with lattice polarization by $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ always admits a second elliptic fibration with a different polarization [17] (see also [38]), which can be related to the $\mathfrak{so}(32)$ heterotic string. To see this, consider the birational transformation

$$(4.1) \quad x = X^2T, \quad y = X^2Y, \quad t = X$$

applied to (2.9). (In applying the transformation, we make the substitution and then divide by the common factor of X^4 .) The result is the equation

$$(4.2) \quad \begin{aligned} Y^2 &= X^2T^3 + aX^2T + bX^2 + cXT + dX + X^3 \\ &= X^3 + (T^3 + aT + b)X^2 + (cT + d)X. \end{aligned}$$

To more easily see the structure, we introduce homogeneous coordinates $[S, T]$ on the base \mathbb{P}^1 and write the equation as

$$(4.3) \quad Y^2 = X^3 + S(T^3 + aS^2T + bS^3)X^2 + S^7(cT + dS)X.$$

It is a straightforward exercise to complete the cube and calculate the discriminant, which is

$$(4.4) \quad \Delta = -S^{16}(cT + dS)^2(T^6 + 2aS^2T^4 + 2bS^3T^3 + a^2S^4T^2 + (2ab - 4c)S^5T + (b^2 - 4d)S^6).$$

Since S divides the coefficient of X^2 and S^2 divides the coefficient of X in (4.3), we conclude that the fiber at $S = 0$ is type I_{10}^* , so the gauge algebra is enhanced to $\mathfrak{so}(28)$. In addition, at the point $[S, T] = [-c, d]$, the coefficient of X^2 is *not* divisible by $(cT + dS)$, so the Kodaira type is I_2 and there is an additional enhancement of the gauge algebra to $\mathfrak{su}(2)$. (For generic coefficients, the other factor in the discriminant contributes six fibers of type I_1 .)

Since the constant term in (4.3) vanishes, the section $X = Y = 0$ defines an element of order 2 in the Mordell-Weil group. It follows as in [5, 7] that the actual gauge group of this model is $(\text{Spin}(28) \times SU(2))/\mathbb{Z}_2$.

The intrinsic property of elliptically fibered K3 surfaces which leads to an equation of the form (4.2) is the requirement that there be a 2-torsion element in the Mordell–Weil group, and that one fiber in the fibration be of type I_n^* for some $n \geq 10$. Under these assumptions, we can choose coordinates so that the specified fiber is at $T = \infty$. If we were to simply ask that the fiber at $T = \infty$ be of type I_{10}^* or worse (as well as having a 2-torsion element), then a slight modification of the argument in section 4 of [3] or appendix A or [38] would show that the equation takes the form

$$(4.5) \quad Y^2 = X^3 + (\alpha T^3 + \beta T^2 + aT + b)X^2 + (cT + d)X.$$

However, if $\alpha = 0$ then it turns out that the coefficient f of the Weierstrass form vanishes to order at least 3 and the coefficient g vanishes to order at least 4, which means that the fiber no longer has type I_n^* . Thus, our requirement of being type I_n^* for some $n \geq 10$ implies that $\alpha \neq 0$ (or in a family, that α has no zeros). Then the coordinate change

$$(4.6) \quad (X, Y, T) \mapsto (\alpha^{-2}X, \alpha^{-3}Y, \alpha^{-1}(T - \frac{1}{3}\beta))$$

(followed by multiplying the equation by α^6) yields an equation of the form (4.2), i.e., one in which the coefficient of T^3X^2 is 1 and the coefficient of T^2X^2 is 0.

The lattice enhancements which we have discussed can also be interpreted for these models. When $c = 0$, the gauge group enhances to $\text{Spin}(32)/\mathbb{Z}_2$, and when $q(a, b, c, d) = 0$, there is an additional enhancement of the gauge algebra by a factor of $\mathfrak{su}(2)$.

5. SIX-DIMENSIONAL COMPACTIFICATIONS

We now specialize to six-dimensional non-geometric heterotic compactifications. The base Z is a Riemann surface with an effective anti-canonical divisor, so it must either be an elliptic curve or the Riemann sphere. In the first case $Z = T^2$, the line bundle Λ is trivial, and the entire construction is just a T^2 compactification of the eight-dimensional theory, with no monodromy or brane insertions needed. In particular, the parameters of the eight-dimensional theory do not vary, and the compactification is geometric.

More interesting is the case $Z = \mathbb{P}^1$. In this case, as derived at the end of Section 2, we have $\Lambda = \mathcal{O}_{\mathbb{P}^1}(2)$, $\Lambda^{\otimes 6} = \mathcal{O}_{\mathbb{P}^1}(12)$, and the base $W = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(12))$ of the non-geometric model coincides with the Hirzebruch surface \mathbb{F}_{12} , and is similar to models first studied in [40]. In particular, a Calabi-Yau three-fold $\tilde{\mathbf{X}} \rightarrow \mathbb{F}_{12}$ can be defined by the Weierstrass equation

$$(5.1) \quad \begin{aligned} 0 = & -y^2 z + x^3 + s^4 t^3 \left(a(u, v) t + c(u, v) s \right) x z^2 \\ & + s^5 t^5 \left(t^2 + b(u, v) s t + d(u, v) s^2 \right) z^3, \end{aligned}$$

where $[u : v]$ denotes the homogeneous coordinates for the \mathbb{P}^1 that constitutes the base of the Hirzebruch surface \mathbb{F}_{12} , and $[s : t]$ denotes the homogeneous coordinates of the \mathbb{P}^1 that constitutes the fiber, and the coefficients $a(u, v)$, $b(u, v)$, $c(u, v)$, and $d(u, v)$ have

degrees 8, 12, 20, and 24, respectively, as homogeneous polynomials on \mathbb{P}^1 . The two \mathbb{C}^* -torus actions that define \mathbb{F}_{12} are given by

$$(5.2) \quad (s, t, u, v) \sim (\lambda^{-12}s, t, \lambda u, \lambda v), \quad (s, t, u, v) \sim (\lambda s, \lambda t, u, v),$$

for $\lambda \in \mathbb{C}^*$. The model has a fiber of type II^* over the section σ_∞ of self-intersection -12 given by $s = 0$, and a fiber of type III^* over a disjoint section σ_0 given by $t = 0$ with $\sigma_0 = \sigma_\infty + 12F$ where F is the fiber class. The divisor class of $\Delta = 0$ is $[\Delta] = -12K_{\mathbb{F}_{12}}$ where $K_{\mathbb{F}_{12}} = -2\sigma_\infty - 14F$. The two curves $s = 0$ and $t = 0$ will account for a large portion of the divisor class $[\Delta]$. The remaining part $[\Delta']$ of the divisor not contained in σ_0 and σ_∞ is $[\Delta'] = [\Delta] - 10\sigma_\infty - 9\sigma_0$. It follows that what is left of the discriminant divisor $[\Delta']$ will not collide with σ_∞ since $\sigma_\infty \cdot [\Delta'] = 0$. On the other hand, the divisor $[\Delta']$ will collide with σ_0 in a total number of $\sigma_0 \cdot [\Delta'] = 60$ points counted with multiplicity.

Because of the choice of \mathbb{F}_{12} , from the $\mathfrak{e}_8 \oplus \mathfrak{e}_7$ heterotic perspective there are no instantons on the \mathfrak{e}_8 summand but the \mathfrak{e}_7 summand has instanton number 24. Since instantons for \mathfrak{e}_8 must be pointlike instantons, this allows the heterotic model to make sense perturbatively,¹⁴ with instantons of finite size on \mathfrak{e}_7 .

The enhancement from \mathfrak{e}_7 to \mathfrak{e}_8 thus occurs at the 20 points $\{c(z) = 0\}$. These points are regarded as responsible for the matter representation of \mathfrak{e}_7 [10], giving 20 half-hypermultiplets in the 56-dimensional representation.

The enhancement from $\mathfrak{e}_8 \oplus \mathfrak{e}_7$ to $\mathfrak{e}_8 \oplus \mathfrak{e}_7 \oplus \mathfrak{su}(2)$ occurs along the locus $q(a, b, c, d) = 0$, which consists of 120 points on \mathbb{P}^1 . At these points, the \mathbb{P}^1 fiber of \mathbb{F}_{12} is tangent to the residual discriminant divisor Δ' .

Similarly, our requirement of a fiber of type I_n^* , $n \geq 10$, in the $\mathfrak{so}(32)$ heterotic string will not allow for the “hidden obstructor” of [3] to occur. (Such “hidden obstructor” points occur when the coefficient of T^3X^2 in (4.2) vanishes; as previously discussed, this vanishing is inconsistent with the fiber being of type I_n^* .) Avoiding these “hidden obstructors” allows for a perturbative description in the $\mathfrak{so}(32)$ case as well. The twenty zeros of $c(z)$ give rise to half-hypermultiplets in the tensor product of the vector representation of $\text{Spin}(28)$ with the fundamental representation of $\text{SU}(2)$ (which is again a 56-dimensional quaternionic representation).

6. DISCUSSION

Something rather surprising has just happened: although we started with a construction for non-geometric heterotic compactifications, the resulting compactifications in six dimensions are actually the familiar F-theory duals of geometric compactifications of the heterotic string on K3 surfaces! How did this happen?

Recall the original description of the heterotic/F-theory duality in six dimensions, as described in [40] and further amplified in [25]: the large radius limit of the heterotic string corresponds to a degeneration limit of the F-theory space, in which the F-theory base actually splits into two components. This limit involves tuning a holomorphic parameter,

¹⁴Note that this is a difference from the case of unbroken gauge algebra $\mathfrak{e}_8 \oplus \mathfrak{e}_8$ considered in [38], where all instantons are pointlike no matter how the instanton numbers are distributed between the two summands.

and tuning holomorphic parameters away from constant has an interesting property: it is not possible to keep the parameter controllably close to the limiting value. Instead, once the holomorphically varying quantity is non-constant, it samples *all* values.

The conclusion, then, is that taking a heterotic compactification even a “small distance” from the large radius limit destroys the traditional semiclassical interpretation and no longer allows us to discuss the compactification as being that of a manifold with a bundle. This is not unlike what happens in type II compactifications, where the analysis of Π -stability [22, 23, 4] shows that going any distance away from large radius limit, no matter how small, necessarily changes the stability conditions on some D-brane classes and so destroys the semiclassical interpretation of the theory.

It would be interesting to check whether this phenomenon persists in compactifications to four dimensions. There, the presence of fluxes may alter the structure of the moduli space, which is here discussed using purely geometric considerations. It may be that when fluxes are involved, some truly new non-geometric models can be constructed. We leave this question for future work.

As pointed out to us by the referee, there may be interesting lessons from this work for double field theory (see, for example, [1, 9, 29] and references therein) and its heterotic extensions. We leave this for future work as well.

Acknowledgements. We would like to thank Chuck Doran and Sav Sethi for helpful discussions. The first author acknowledges the generous support of the Kavli Institute for Theoretical Physics, and the second author is grateful to the Kavli Institute for the Physics and Mathematics of the Universe for hospitality during the early stages of this project. The work of the second author is supported by National Science Foundation grants DMS-1007414 and PHY-1307513 and by the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan.

APPENDIX A. DISCRETE GROUPS AND MODULAR FORMS

A.1. Modular forms for $O^+(\Lambda^{2,2})$. Explicit generators for $O(\Lambda^{2,2})$ are given in [30], together with their actions on $\mathbb{H} \times \mathbb{H}$. It implies that we can identify $O^+(\Lambda^{2,2})$ with $P(\mathrm{SL}_2(\mathbb{Z}) \times \mathrm{SL}_2(\mathbb{Z})) \rtimes \mathbb{Z}_2$, where the automorphism \mathbb{Z}_2 acts to exchange ρ and τ .

The modular forms of weight d for this group must be functions of ρ and τ of bidegree (d, d) invariant under the exchange. We claim that this ring of modular forms is a free polynomial algebra on $E_4(\rho)E_4(\tau)$, $E_6(\rho)E_6(\tau)$, and $\Delta_{12}(\rho)\Delta_{12}(\tau)$.

To see why this is true, let $t_4 = E_4(\rho)E_4(\tau)$, $t_6 = E_6(\rho)E_6(\tau)$, and $t_{12} = \Delta_{12}(\rho)\Delta_{12}(\tau)$ be elements of bidegree $(4, 4)$, $(6, 6)$, and $(12, 12)$, respectively. If they are not algebraically independent, then there exists a nonvanishing homogeneous polynomial $P(T)$ in the graded ring $\mathbb{C}[T_4, T_6, T_{12}]$ satisfying $P(t_4, t_6, t_{12}) = 0$. We take as $P(T)$ the polynomial of minimal degree and write it in the form $P_0(T_4, T_6, T_{12})T_{12} + P_1(T_4, T_6)$. In the equation

$$P_0(t_4, t_6, t_{12})t_{12} + P_1(t_4, t_6) = 0$$

we take the limit $\rho \rightarrow i\infty$. Notice that under $\rho \rightarrow i\infty$ we have

$$E_4(\rho) \rightarrow 1, \quad E_6(\rho) \rightarrow 1 \quad \eta(\rho) \rightarrow 0.$$

Therefore, we get $P_1(E_4(\tau), E_6(\tau)) = 0$. But $E_4(\tau)$ and $E_6(\tau)$ are algebraically independent. It follows that $P_1(T_4, T_6) = 0$ and $P_0(T)$ is different from zero with $P_0(t_4, t_6, t_{12}) = 0$. Since $P_0(T)$ is of smaller degree than $P(T)$ we get a contradiction to the assumed minimality of $P(T)$.

Fix $k \geq 0$ even and let $n(k)$ denote the dimension of the space of modular forms M_k of weight k for $\mathrm{SL}_2(\mathbb{Z})$. It is well-known that

$$(A.1) \quad n(k) := \dim M_k = \begin{cases} \lfloor k/12 \rfloor & \text{for } k \equiv 2(4), \\ \lfloor k/12 \rfloor + 1 & \text{otherwise.} \end{cases}$$

Equivalently, the dimension $n(k)$ equals the number of nonnegative integer solutions to the equation $k = 4p + 6q$ as E_4 and E_6 generate the ring of modular forms. Then, the dimension of $M_k \otimes M_k$ is $n(k)^2$, and the dimension of the linear subspace of bi-degree (k, k) defined by $f(\rho, \tau) = f(\tau, \rho)$ is $\frac{1}{2}n(k)(n(k)+1)$. Let R denote the graded subring generated by the algebraically independent t_4, t_6, t_{12} . The dimension of the subspace R_k of bi-degree (k, k) equals the number of nonnegative integer solutions to the equation $k = 4p + 6q + 12r$. If we fix r , then the number of such solutions equals the dimension of M_{k-12r} . From Equation (A.1) it follows that for the dimension we have $\dim M_k = \dim M_{k-12} + 1$. Let $r_0 = \lfloor k/12 \rfloor$ then summing over possible values for r we obtain the dimension of R_k :

$$\dim R_k = \sum_{r=0}^{r_0} \dim M_{k-12r_0+12r} = \begin{cases} \sum_{r=1}^{r_0} r & \text{for } k \equiv 2(4) \\ \sum_{r=0}^{r_0} r & \text{otherwise} \end{cases} = \frac{1}{2} n(k) (n(k) + 1),$$

which agrees with the dimension of $\mathrm{Sym}^2(M_k)$. It follows that the elements t_4, t_6, t_{12} generate $\mathrm{Sym}^2(M_*)$.

A.2. The moduli spaces. It is worthwhile to straighten out several moduli spaces of relevance here. The key observation, due to Vinberg [50], is that under the natural homomorphism $\mathrm{Sp}_4(\mathbb{R}) \rightarrow O^+(2, 3)$, the arithmetic group $\mathrm{Sp}_4(\mathbb{Z})$ (which is a maximal discrete subgroup of $\mathrm{Sp}_4(\mathbb{R})$) maps to an index two subgroup $SO^+(L^{2,3}) \subset O^+(L^{2,3})$ where $O^+(2, 3)$ denotes the subgroup of index 2 of $O(2, 3)$ consisting of the elements whose spinor norm is equal to the determinant. The isomorphism

$$(A.2) \quad \mathbb{H}_2 / \mathrm{Sp}_4(\mathbb{Z}) \cong \mathcal{D}_{2,3} / SO^+(L^{2,3})$$

gives rise to an isomorphism between the algebra of Siegel modular forms of genus 2 and the algebra of automorphic forms of $\mathcal{D}_{2,3}$ with respect to the group $SO^+(L^{2,3})$. But the algebra of automorphic forms of $\mathcal{D}_{2,3}$ with respect to the group $O^+(L^{2,3})$ is the *even* part of the algebra of automorphic forms with respect to $SO^+(L^{2,3})$ and, hence, is isomorphic to the algebra of *even* Siegel modular forms of genus 2.

This means that the moduli space of principally polarized abelian surfaces $\mathbb{H}_2 / \mathrm{Sp}_4(\mathbb{Z})$ has a degree two map to the moduli space of K3 surfaces with lattice polarization $\mathcal{D}_{2,3} / O^+(L^{2,3})$. This can be understood at the level of modular forms as follows. The Siegel modular forms of even weight, generated by $\psi_4, \psi_6, \chi_{10}$ and χ_{12} , are invariant under




FIGURE 1. Relations among moduli spaces

$O^+(L^{2,3})$ and give homogenous coordinates on the Baily-Borel compactification of that space. On the other hand, the full ring of modular forms is invariant under $\mathrm{Sp}_4(\mathbb{Z})$, and the equation $\chi_{35}^2 = \mathcal{F}(\psi_4, \psi_6, \chi_{10}, \chi_{12})$ (where \mathcal{F} is given in Equation (B.16)) expresses $\mathbb{H}_2/\mathrm{Sp}_4(\mathbb{Z})$ as a double cover of $\mathcal{D}_{2,3}/O^+(L^{2,3})$.

This same phenomenon carries over to the moduli of genus two curves. We can express a genus two curve in the form $y^2 = f(x)$ (cf. Appendix B.3) and so there is a map $\mathcal{M}_2 \rightarrow \mathcal{U}_6$ from the moduli space of genus two curves to the moduli space of degree six polynomials or sextics. This also turns out to be a map of degree two. The coordinates on the moduli space of degree six polynomials were worked out by Clebsch: in Igusa's notation, they are $I_2(f), I_4(f), I_6(f), I_{10}(f)$ and given in Equations (B.24). On the other hand, the moduli space of genus-two curves has one additional invariant R defined in Appendix B.3. The point is that under the operation $f(x) \mapsto \tilde{f}(x) = f(-x)$ the odd invariant $R(f)$ is mapped to $R(\tilde{f}) = -R(f)$ whereas the even invariants $I_2(f), I_4(f), I_6(f), I_{10}(f)$ remain the same, i.e., $I_{2k}(f) = I_{2k}(\tilde{f})$ for $k = 2, 4, 6, 10$. The subtle point is that mapping $f \mapsto \tilde{f}$ and, hence, the ramification points $\theta_i \mapsto -\theta_i$ defines equivalent sextics, but different genus-two curves. In fact, genus-two curves invariant under this action are the ones with bigger automorphism group with an extra involution of order two and $R(f) = 0$. Equation (B.28) expresses $R^2 = \mathcal{F}'(I_2, I_4, I_6, I_{10})$ as a polynomial in terms of the *even* Igusa-Clebsch invariants. Therefore, this expresses \mathcal{M}_2 as a double cover of \mathcal{U}_6 . The diagram in Figure 1 summarizes the discussion.

APPENDIX B. THE MODULI SPACE OF PRINCIPALLY POLARIZED ABELIAN SURFACES

B.1. The Siegel modular three-fold. The Siegel three-fold is a quasi-projective variety of dimension 3 obtained from the Siegel upper half-plane of degree two which by definition is the set of two-by-two symmetric matrices over \mathbb{C} whose imaginary part is

positive definite, i.e.,

$$(B.1) \quad \mathbb{H}_2 = \left\{ \underline{\tau} = \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} \middle| \tau_1, \tau_2, z \in \mathbb{C}, \operatorname{Im}(\tau_1) \operatorname{Im}(\tau_2) > \operatorname{Im}(z)^2, \operatorname{Im}(\tau_2) > 0 \right\},$$

quotiented out by the action of the modular transformations $\Gamma_2 := \operatorname{Sp}_4(\mathbb{Z})$, i.e.,

$$(B.2) \quad \mathcal{A}_2 = \mathbb{H}_2 / \Gamma_2.$$

Each $\underline{\tau} \in \mathbb{H}_2$ determines a principally polarized complex abelian surface $\mathbf{A}_{\underline{\tau}} = \mathbb{C}^2 / \langle \mathbb{Z}^2 \oplus \underline{\tau} \mathbb{Z}^2 \rangle$ with period matrix $(\underline{\tau}, \mathbb{I}_2) \in \operatorname{Mat}(2, 4; \mathbb{C})$. Two abelian surfaces $\mathbf{A}_{\underline{\tau}}$ and $\mathbf{A}_{\underline{\tau}'}$ are isomorphic if and only if there is a symplectic matrix

$$(B.3) \quad M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_2$$

such that $\underline{\tau}' = M(\underline{\tau}) := (A\underline{\tau} + B)(C\underline{\tau} + D)^{-1}$. It follows that the Siegel three-fold \mathcal{A}_2 is also the set of isomorphism classes of principally polarized abelian surfaces. The sets of abelian surfaces that have the same endomorphism ring form subvarieties of \mathcal{A}_2 . The endomorphism ring of principally polarized abelian surface tensored with \mathbb{Q} is either a quartic CM field, an indefinite quaternion algebra, a real quadratic field or in the generic case \mathbb{Q} . Irreducible components of the corresponding subsets in \mathcal{A}_2 have dimensions 0, 1, 2 and are known as CM points, Shimura curves and Humbert surfaces, respectively.

The Humbert surface H_{Δ} with invariant Δ is the space of principally polarized abelian surfaces admitting a symmetric endomorphism with discriminant Δ . It turns out that Δ is a positive integer $\equiv 0, 1 \pmod{4}$. In fact, H_{Δ} is the image inside \mathcal{A}_2 under the projection of the rational divisor associated to the equation

$$(B.4) \quad a\tau_1 + bz + c\tau_3 + d(z^2 - \tau_1\tau_2) + e = 0,$$

with integers a, b, c, d, e satisfying $\Delta = b^2 - 4ac - 4de$ and $\underline{\tau} = \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} \in \mathbb{H}_2$. For example, inside of \mathcal{A}_2 sit the Humbert surfaces H_1 and H_4 that are defined as the images under the projection of the rational divisor associated to $z = 0$ and $\tau_1 = \tau_2$, respectively. Equivalently, these points are invariant under the \mathbb{Z}_2 -action generated by $\begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix} \in \Gamma_2$ with $A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, respectively. In fact, the singular locus of \mathcal{A}_2 has H_1 and H_4 as its two connected components. As analytic spaces, the surfaces H_1 and H_4 are each isomorphic to the Hilbert modular surface

$$(B.5) \quad \left((\operatorname{SL}_2(\mathbb{Z}) \times \operatorname{SL}_2(\mathbb{Z})) \rtimes \mathbb{Z}_2 \right) \backslash (\mathbb{H} \times \mathbb{H}).$$

For a more detailed introduction to Siegel modular form, Humbert surfaces, and the Satake compactification of the Siegel modular threefold we refer to Freitag's book [24].

B.2. Siegel modular forms. In general, we can define the Eisenstein series ψ_{2k} of degree g and weight $2k$ (where we assume $2k > g + 1$ for convergence) by setting

$$(B.6) \quad \psi_{2k}(\underline{\tau}) = \sum_{(C, D)} \det(C \cdot \underline{\tau} + D)^{-2k},$$

where the sum runs over non-associated bottom rows (C, D) of elements in $\mathrm{Sp}_{2g}(\mathbb{Z})$ where non-associated means with respect to the multiplication on the left by $\mathrm{GL}(g, \mathbb{Z})$. For $g = 1$ and $k > 1$, we have $\mathrm{Sp}_2(\mathbb{Z}) = \mathrm{SL}_2(\mathbb{Z})$, $\mathrm{GL}(1, \mathbb{Z}) = \mathbb{Z}_2$, and we obtain $\psi_{2k}(\tau) = E_{2k}(\tau)$ where $E_{2k}(\tau) = 1 + O(q)$ with $q = \exp(2\pi i\tau)$ are the standard normalized Eisenstein series. The reason is that the series E_{2k} be written as

$$(B.7) \quad E_{2k}(\tau) = \sum_{\substack{(c,d)=1 \\ (c,d) \equiv (-c,-d)}} \frac{1}{(c\tau + d)^{2k}} ,$$

where the sum runs over all pairs of co-prime integers up to simultaneous $\mathbb{Z}_2 = \mathrm{GL}(1, \mathbb{Z})$ action. The connection to the Eisenstein series G_{2k} is given by

$$(B.8) \quad G_{2k}(\tau) = 2 \zeta(2k) E_{2k}(\tau) = \sum_{(m,n) \in \mathbb{Z}^2 \setminus (0,0)} \frac{1}{(m\tau + n)^{2k}} .$$

In the following, we will always assume $g = 2$ in the definition of ψ_{2k} . For $z \rightarrow 0$, we then have that

$$(B.9) \quad \psi_{2k} \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} = E_{2k}(\tau_1) E_{2k}(\tau_2) + O(z^2) .$$

Following Igusa [31] we define a cusp form of weight 12 by

$$(B.10) \quad \chi_{12}(\underline{\tau}) = \frac{691}{2^{13} 3^8 5^3 7^2} (3^2 7^2 \psi_4^3(\underline{\tau}) + 2 \cdot 5^3 \psi_6^2(\underline{\tau}) - 691 \psi_{12}(\underline{\tau})) .$$

We find that for $z \rightarrow 0$ its asymptotic behavior is given by

$$(B.11) \quad \chi_{12} \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} = \eta^{24}(\tau_1) \eta^{24}(\tau_2) + O(z^2)$$

where $\eta(\tau)$ is the Dedekind η -function and we have used that

$$(B.12) \quad \begin{aligned} 1728 \eta^{24}(\tau_j) &= E_4^3(\tau_j) - E_6^2(\tau_j) , \\ 691 E_{12}(\tau_j) &= 441 E_4(\tau_j)^3 + 250 E_6(\tau_j)^2 . \end{aligned}$$

Igusa's 'original' definition [31, Sec. 8, p. 195] for χ_{12} is

$$(B.13) \quad \begin{aligned} \tilde{\chi}_{12}(\underline{\tau}) &= \frac{131 \cdot 593}{2^{13} 3^7 5^3 7^2 337} (3^2 7^2 \psi_4^3(\underline{\tau}) + 2 \cdot 5^3 \psi_6^2(\underline{\tau}) - 691 \psi_{12}(\underline{\tau})) \\ &= \frac{3 \cdot 131 \cdot 593}{337 \cdot 691} \chi_{12}(\underline{\tau}) = 1.00078 \dots \chi_{12}(\underline{\tau}) . \end{aligned}$$

But all results in [31] that connect χ_{12} to the Igusa-Clebsch coefficients are based on the asymptotic expansion in Equation (B.11). Hence, the definition in Equation (B.10) must be used. Using Igusa's definition [31, Sec. 8, p. 195] we also define a second cusp form of weight 10 by

$$(B.14) \quad \chi_{10}(\underline{\tau}) = -\frac{43867}{2^{12} 3^5 5^2 7 \cdot 53} (\psi_4(\underline{\tau}) \psi_6(\underline{\tau}) - \psi_{10}(\underline{\tau})) .$$

We see that for $z \rightarrow 0$ its asymptotic behavior is given by

$$(B.15) \quad \chi_{10} \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix} = \eta^{24}(\tau_1) \eta^{24}(\tau_2) (\pi z)^2 + O(z^4).$$

Hence, the vanishing divisor of the cusp form χ_{10} is the Humbert surface H_1 because a period point $\underline{\tau}$ is equivalent to a point with $z = 0$ if and only if $\chi_{10}(\underline{\tau}) = 0$.

Igusa proved [32, 33] that the ring of Siegel modular forms is generated by ψ_4 , ψ_6 , χ_{10} , χ_{12} and by one more cusp form χ_{35} of odd weight 35 whose square is the following polynomial [32, p. 849] in the even generators

$$(B.16) \quad \begin{aligned} \chi_{35}^2 = & \frac{1}{2^{12} 3^9} \chi_{10} \left(2^{24} 3^{15} \chi_{12}^5 - 2^{13} 3^9 \psi_4^3 \chi_{12}^4 - 2^{13} 3^9 \psi_6^2 \chi_{12}^4 + 3^3 \psi_4^6 \chi_{12}^3 \right. \\ & - 2 \cdot 3^3 \psi_4^3 \psi_6^2 \chi_{12}^3 - 2^{14} 3^8 \psi_4^2 \psi_6 \chi_{10} \chi_{12}^3 - 2^{23} 3^{12} 5^2 \psi_4 \chi_{10}^2 \chi_{12}^3 + 3^3 \psi_4^4 \chi_{12}^3 \\ & + 2^{11} 3^6 37 \psi_4^4 \chi_{10}^2 \chi_{12}^2 + 2^{11} 3^6 5 \cdot 7 \psi_4 \psi_6^2 \chi_{10}^2 \chi_{12}^2 - 2^{23} 3^9 5^3 \psi_6 \chi_{10}^3 \chi_{12}^2 \\ & - 3^2 \psi_4^7 \chi_{10}^2 \chi_{12}^2 + 2 \cdot 3^2 \psi_4^4 \psi_6^2 \chi_{10}^2 \chi_{12}^2 + 2^{11} 3^5 5 \cdot 19 \psi_4^3 \psi_6 \chi_{10}^3 \chi_{12}^2 \\ & + 2^{20} 3^8 5^3 11 \psi_4^2 \chi_{10}^4 \chi_{12}^2 - 3^2 \psi_4 \psi_6^4 \chi_{10}^2 \chi_{12}^2 + 2^{11} 3^5 5^2 \psi_6^3 \chi_{10}^3 \chi_{12}^2 - 2 \psi_4^6 \psi_6 \chi_{10}^3 \chi_{12}^2 \\ & - 2^{12} 3^4 \psi_4^5 \chi_{10}^4 + 2^2 \psi_4^3 \psi_6^3 \chi_{10}^3 + 2^{12} 3^4 5^2 \psi_4^2 \psi_6^2 \chi_{10}^4 + 2^{21} 3^7 5^4 \psi_4 \psi_6 \chi_{10}^5 \\ & \left. - 2 \psi_6^5 \chi_{10}^3 + 2^{32} 3^9 5^5 \chi_{10}^6 \right). \end{aligned}$$

Hence, $Q := 2^{12} 3^9 \chi_{35}^2 / \chi_{10}$ is a polynomial of degree 60 in the even generators. One then checks that

$$(B.17) \quad q \left(-\frac{1}{48} \psi_4, -\frac{1}{864} \psi_6, -4 \chi_{10}, \chi_{12} \right) = \frac{1}{2^{20} 3^9} Q(\psi_4, \psi_6, \chi_{10}, \chi_{12}),$$

where q was defined in Equation (3.4). It is known that the vanishing divisor of Q is the Humbert surface H_4 [27] because a period point $\underline{\tau}$ is equivalent to a point with $\tau_1 = \tau_2$ if and only if $Q(\underline{\tau}) = 0$. Accordingly, the vanishing divisor of χ_{35} is the formal sum $H_1 + H_4$ of Humbert surfaces, that constitutes the singular locus of \mathcal{A}_2 .

In accordance with Igusa [31, Theorem 3] we also introduce the following ratios of Siegel modular forms

$$(B.18) \quad \mathbf{x}_1 = \frac{\psi_4 \chi_{10}^2}{\chi_{12}^2}, \quad \mathbf{x}_2 = \frac{\psi_6 \chi_{10}^3}{\chi_{12}^3}, \quad \mathbf{x}_3 = \frac{\chi_{10}^6}{\chi_{12}^5},$$

as well as

$$(B.19) \quad \mathbf{y}_1 = \frac{\mathbf{x}_1^3}{\mathbf{x}_3} = \frac{\psi_4^3}{\chi_{12}}, \quad \mathbf{y}_2 = \frac{\mathbf{x}_2^2}{\mathbf{x}_3} = \frac{\psi_6^2}{\chi_{12}}, \quad \mathbf{y}_3 = \frac{\mathbf{x}_1^2 \mathbf{x}_2}{\mathbf{x}_3} = \frac{\psi_4^2 \psi_6 \chi_{10}}{\chi_{12}},$$

where we have suppressed the dependence of each Siegel modular form on $\underline{\tau}$. These ratios have the following asymptotic expansion as $z \rightarrow 0$ [31, pp. 180–182]

$$(B.20) \quad \begin{aligned} \mathbf{x}_1 &= E_4(\tau_1) E_4(\tau_2) (\pi z)^4 + O(z^5), \\ \mathbf{x}_2 &= E_6(\tau_1) E_6(\tau_2) (\pi z)^6 + O(z^7), \\ \mathbf{x}_3 &= \eta^{24}(\tau_1) \eta^{24}(\tau_2) (\pi z)^{12} + O(z^{13}), \end{aligned}$$

and

$$\begin{aligned}
 \mathbf{y}_1 &= j(\tau_1) j(\tau_2) + O(z^2) , \\
 \mathbf{y}_2 &= \left(1728 - j(\tau_1)\right) \left(1728 - j(\tau_2)\right) + O(z^2) , \\
 \mathbf{y}_3 &= \frac{E_4^2(\tau_1) E_4^2(\tau_2) E_6(\tau_1) E_6(\tau_2)}{\eta^{24}(\tau_1) \eta^{24}(\tau_2)} (\pi z)^2 + O(z^3) ,
 \end{aligned}
 \tag{B.21}$$

where we have set

$$\begin{aligned}
 j(\tau_j) &= \frac{1728 E_4^3(\tau_j)}{E_4^3(\tau_j) - E_6^2(\tau_j)} = \frac{E_4^3(\tau_j)}{\eta^{24}(\tau_j)} , \\
 1728 - j(\tau_j) &= \frac{1728 E_6^2(\tau_j)}{E_4^3(\tau_j) - E_6^2(\tau_j)} = \frac{E_6^2(\tau_j)}{\eta^{24}(\tau_j)} .
 \end{aligned}
 \tag{B.22}$$

Notice that for the asymptotic behavior in Equations (B.20) and (B.21) the right normalization of χ_{12} was essential.

B.3. Sextics and Igusa invariants. We write the equation defining a genus-two curve C by a degree-six polynomial or sextic in the form

$$C : y^2 = f(x) = a_0 \prod_{i=1}^6 (x - \theta_i) = \sum_{i=0}^6 a_i x^{6-i} .
 \tag{B.23}$$

The roots $(\theta_i)_{i=1}^6$ of the sextic are the six ramification points of the map $C \rightarrow \mathbb{P}^1$. Their pre-images on C are the six Weierstrass points. The isomorphism class of f consists of all equivalent sextics where two sextics are considered equivalent if there is a linear transformation in $\mathrm{GL}(2, \mathbb{C})$ which takes the set of roots to the roots of the other. The action of the linear transformations on the Weierstrass points defines a 7-dimensional irreducible linear representation of $\mathrm{SL}_2(\mathbb{C})$. The corresponding invariants are called the invariants of the sextic.

Clebsch defined such invariants I_2, I_4, I_6, I_{10} of weights 2, 4, 6, 10, respectively, now called the *Igusa-Clebsch invariants* of the sextic curve in (B.23), as follows

$$\begin{aligned}
 I_2(f) &= a_0^2 \sum_{i < j, k < l, m < n} D_{ij}^2 D_{kl}^2 D_{mn}^2 , \\
 I_4(f) &= a_0^4 \sum_{i < j < k, l < m < n} D_{ij}^2 D_{jk}^2 D_{ki}^2 D_{lm}^2 D_{mn}^2 D_{nl}^2 , \\
 I_6(f) &= a_0^6 \sum_{\substack{i < j < k, l < m < n \\ i < l', j < m', k < n' \\ l', m', n' \in \{l, m, n\}}} D_{ij}^2 D_{jk}^2 D_{ki}^2 D_{lm}^2 D_{mn}^2 D_{nl}^2 D_{il'}^2 D_{jm'}^2 D_{kn'}^2 , \\
 I_{10}(f) &= a_0^{10} \prod_{i < j} D_{ij}^2 ,
 \end{aligned}
 \tag{B.24}$$

where $D_{ij} = \theta_i - \theta_j$ and all indices take values in $\{1, \dots, 6\}$. In the following, we will often suppress the argument f . The invariants (I_2, I_4, I_6, I_{10}) are the same invariants as

(A', B', C', D') in [39, p. 319] and also the same invariants as (A, B, C, D) in [31, p. 176]. It follows from the work of Mestre [39] that the Igusa-Clebsch invariants are arithmetic invariants, i.e., polynomials in the coefficients a_0, \dots, a_6 with integer coefficients, or $I_k \in \mathbb{Z}[a_0, \dots, a_6]$ for $k \in \{2, 4, 6, 10\}$. Furthermore, a theorem by Bolza and Clebsch states that two sextics given by f and f' are isomorphic if and only if there is a $\rho \in \mathbb{C}^*$ such that $I_{2k}(f') = \rho^{-2k} I_{2k}(f)$ for $k = 1, 2, 3, 5$. Thus, the invariants of a sextic define a point in a weighted projective space $[I_2 : I_4 : I_6 : I_{10}] \in \mathbb{WP}_{(2,4,6,10)}^3$. It was shown in [31] that points in the projective variety $\text{Proj } \mathbb{C}[I_2, I_4, I_6, I_{10}]$ which are not on $I_{10} = 0$ form the variety \mathcal{U}_6 of moduli of sextics. Equivalently, points in this weighted projective space $\{[I_2 : I_4 : I_6 : I_{10}] \in \mathbb{WP}_{(2,4,6,10)}^3 : I_{10} \neq 0\}$ are in one-to-one correspondence with isomorphism classes of sextics.

Often the *Clebsch invariants* of a sextic are used as well. The Clebsch invariants (A, B, C, D) are related to the Igusa-Clebsch invariants by the equations

$$\begin{aligned}
 (B.25) \quad I_2 &= -120 A, \\
 I_4 &= -720 A^2 + 6750 B, \\
 I_6 &= 8640 A^3 - 108000 A B + 202500 C, \\
 I_{10} &= -62208 A^5 + 972000 A^3 B + 1620000 A^2 C \\
 &\quad - 3037500 A B^2 - 6075000 B C - 4556250 D.
 \end{aligned}$$

Conversely, the invariants (A, B, C, D) are polynomial expressions in the Igusa invariants (I_2, I_4, I_6, I_{10}) with rational coefficients. Mestre [39] also defined the following polynomials in the Clebsch invariants

$$\begin{aligned}
 (B.26) \quad A_{11} &= 2 C + \frac{1}{3} A B, \\
 A_{22} &= A_{31} = D, \\
 A_{33} &= \frac{1}{2} B D + \frac{2}{9} C (B^2 + A C), \\
 A_{23} &= \frac{1}{3} B (B^2 + A C) + \frac{1}{3} C (2 C + \frac{1}{3} A B), \\
 A_{12} &= \frac{2}{3} (B^2 + A C).
 \end{aligned}$$

According to [39] one can obtain from a sextic f three binary quadrics of the form

$$(B.27) \quad y_i(x) := \alpha_i x^2 + \beta_i x + \gamma_i$$

with $i = 1, 2, 3$ by an operation called ‘Überschiebung’ [39, p. 317]. To fix the normalization and order of the quadrics we remark that in the notation of [39] we have $I_{10} = (y_3 y_1)_2$. The quadrics y_i for $i = 1, 2, 3$ have the property that their coefficients are polynomial expressions in the coefficients of f with rational coefficients. Moreover, under the operation $f(x) \mapsto \tilde{f}(x) = f(-x)$ the quadrics change according to $y_i(x) \mapsto \tilde{y}_i(x) = y_i(-x)$ for $i = 1, 2, 3$. Hence, they are not invariants of the sextic. In contrast, $I_2(f), I_4(f), I_6(f), I_{10}(f)$ remain unchanged under this operation, i.e.,

$I_{2k}(f) = I_{2k}(\tilde{f})$ for $k = 2, 4, 6, 10$. This latter statement is easily checked since Equations (B.24) are invariant under $f \mapsto \tilde{f}$ or, equivalently, $D_{ij} \mapsto -D_{ij}$.

We define R to be $1/4$ times the determinant of the three binary quadrics y_i for $i = 1, 2, 3$ with respect to the basis $x^2, x, 1$. It is obvious that under the operation $f(x) \mapsto \tilde{f}(x) = f(-x)$ the determinant R changes its sign, i.e., $R(f) \mapsto R(\tilde{f}) = -R(f)$. A calculation shows that

$$(B.28) \quad R^2 = \frac{1}{2} \begin{vmatrix} A_{11} & A_{12} & A_{31} \\ A_{12} & A_{22} & A_{23} \\ A_{31} & A_{23} & A_{33} \end{vmatrix},$$

where A_{ij} are the Clebsch invariants (B.26). Like $R(f)^2$, the coefficients $A_{ij}(f)$ are invariant under the operation $f(x) \mapsto \tilde{f}(x) = f(-x)$ as they are polynomials in (I_2, I_4, I_6, I_{10}) . Bolza [11] described the possible automorphism groups of genus-two curves defined by sextics. In particular, he provided effective criteria for the cases when the automorphism group of the sextic curve in (B.23) is nontrivial. The results are as follows:

- (1) The curve has an extra involution other than the exchange of sheets $(x, y) \rightarrow (x, -y)$ if and only if $R^2 = 0$. The sextic is then isomorphic to $f(x) = x^6 + c_1 x^4 + c_2 x^2 + 1$ for some $c_1, c_2 \in \mathbb{C}$ with the extra involution $(x, y) \rightarrow (-x, y)$.
- (2) The automorphism group contains an element of order 5 if and only if $I_2 = I_4 = I_6 = 0, I_{10} \neq 0$. The sextic is then isomorphic to $f(x) = x(x^5 + 1)$ with the element of order 5 being $(x, y) \rightarrow (\zeta_5^2 x, \zeta_5 y)$ where $\zeta_5 = \exp(2\pi i/5)$.

B.4. The moduli space of genus-two curves. Suppose that C is an irreducible projective nonsingular curve. If the self-intersection is $C \cdot C = 2$ then C is a curve of genus two. For every curve C of genus two there exists a unique pair $(\text{Jac}(C), j_C)$ where $\text{Jac}(C)$ is an abelian surface, called the Jacobian variety of the curve C , and $j_C : C \rightarrow \text{Jac}(C)$ is an embedding. One can always regain C from the pair $(\text{Jac}(C), \mathcal{P})$ where $\mathcal{P} = [C]$ is the class of C in the Néron-Severi group $\text{NS}(\text{Jac}(C))$. Thus, if C is a genus-two curve, then $\text{Jac}(C)$ is a principally polarized abelian surface with principal polarization $\mathcal{P} = [C]$, and the map sending a curve C to its Jacobian variety $\text{Jac}(C)$ is injective. In this way, the variety of moduli of curves of genus two is also the moduli space of their Jacobian varieties with canonical polarization. Since we have $\mathcal{P}^2 = 2$, the transcendental lattice is $T(\text{Jac}(C)) = \Lambda^{2,2} \oplus \langle -2 \rangle$. Furthermore, Torelli's theorem states that the map sending a curve C to its Jacobian variety $\text{Jac}(C)$ induces a birational map from the moduli space \mathcal{M}_2 of genus-two curves to the complement of the Humbert surface H_1 in \mathcal{A}_2 , i.e., $\mathcal{A}_2 - \text{supp}(\chi_{10})_0$.

One can then ask what the Igusa-Clebsch invariants of a genus-two curve C defined by a sextic curve f are in terms of $\underline{\tau}$ such that $(\underline{\tau}, \mathbb{I}_2) \in \text{Mat}(2, 4; \mathbb{C})$ is the period matrix of the principally polarized abelian surface $\mathbf{A}_{\underline{\tau}} = \text{Jac}(C)$. Based on the asymptotic behavior

in Equations (B.20) and (B.21), Igusa [31] proved that the relations are as follows:

$$\begin{aligned}
 I_2(f) &= \frac{\chi_{12}(\underline{\tau})}{\chi_{10}(\underline{\tau})}, \\
 I_4(f) &= \frac{1}{2^4 3^2} \psi_4(\underline{\tau}), \\
 I_6(f) &= \frac{1}{2^6 3^4} \psi_6(\underline{\tau}) + \frac{1}{2^4 3^3} \frac{\psi_4(\underline{\tau}) \chi_{12}(\underline{\tau})}{\chi_{10}(\underline{\tau})}, \\
 I_{10}(f) &= \frac{1}{2 \cdot 3^5} \chi_{10}(\underline{\tau}).
 \end{aligned}
 \tag{B.29}$$

Thus, we find that the point $[I_2 : I_4 : I_6 : I_{10}]$ in weighted projective space equals

$$\left[2^3 3 (3r\chi_{12}) : 2^2 3^2 \psi_4(r\chi_{10})^2 : 2^3 3^2 \left(4\psi_4(3r\chi_{12}) + \psi_6(r\chi_{10}) \right) (r\chi_{10})^2 : 2^2 (r\chi_{10})^6 \right]$$

with $r = 2^{12} 3^5$. Substituting (B.29) into Equations (B.25), (B.26) it also follows that

$$\begin{aligned}
 R(f)^2 &= 2^{-41} 3^{-42} 5^{-20} \frac{Q(\psi_4(\underline{\tau}), \psi_6(\underline{\tau}), \chi_{10}(\underline{\tau}), \chi_{12}(\underline{\tau}))}{\chi_{10}(\underline{\tau})^3} \\
 &= 2^{-29} 3^{-33} 5^{-20} \frac{\chi_{35}(\underline{\tau})^2}{\chi_{10}(\underline{\tau})^4},
 \end{aligned}
 \tag{B.31}$$

where Q and R^2 were defined in Equation (B.16) and (B.28), respectively.

If $\underline{\tau}$ is equivalent to a point with $\tau_1 = \tau_2$ or $[\underline{\tau}] \in H_4 \subset \mathcal{A}_2$ then the corresponding sextic curve has an extra automorphism with $R(f)^2 = 0$. The transcendental lattice degenerates to $T(\mathbf{A}_{\underline{\tau}}) = \Lambda^{1,1} \oplus \langle 2 \rangle \oplus \langle -2 \rangle$. If $\underline{\tau}$ is equivalent to a point with $z = 0$ or $[\underline{\tau}] \in H_1 \subset \mathcal{A}_2$, then the principally polarized abelian surface is a product of two elliptic curves $\mathbf{A}_{\underline{\tau}} = E_{\tau_1} \times E_{\tau_2}$ because of Equations (B.32) and (B.20). The transcendental lattice degenerates to $T(\mathbf{A}_{\underline{\tau}}) = \Lambda^{2,2}$.

For $I_2 \neq 0$ we use the variables $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ from Equations (B.18) to write

$$(B.32) \quad [I_2 : I_4 : I_6 : I_{10}] = \left[1 : \frac{1}{2^4 3^2} \mathbf{x}_1 : \frac{1}{2^6 3^4} \mathbf{x}_2 + \frac{1}{2^4 3^3} \mathbf{x}_1 : \frac{1}{2 \cdot 3^5} \mathbf{x}_3 \right] \in \mathbb{WP}_{(2,4,6,10)}^3.$$

Since the invariants I_4, I_6, I_{10} vanish simultaneously at sextics with triple roots all such abelian surfaces are mapped to $[1 : 0 : 0 : 0] \in \mathbb{WP}_{(2,4,6,10)}^3$ with uniformizing affine coordinates $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ around it. Blowing up this point gives a variety that parameterizes genus-two curves with $I_2 \neq 0$ and their degenerations. In the blow-up space we have to introduce additional coordinates that are obtained as ratios of $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3$ and have weight zero. Those are precisely the coordinates $\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3$ already introduced in Equation (B.19). It turns out that the coordinate ring of the blown-up space is $\mathbb{C}[\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3]$.

If a Jacobian variety corresponds to a product of elliptic curves then $\underline{\tau}$ is equivalent to a point with $z = 0$, i.e., $\underline{\tau}$ is located on the Humbert surface H_1 . We then have $\chi_{10}(\underline{\tau}) = 0, \chi_{12}(\underline{\tau}) \neq 0$ and $[I_2 : I_4 : I_6 : I_{10}] = [1 : 0 : 0 : 0]$. Equations (B.20) and (B.21) imply $\mathbf{x}_1 = \mathbf{x}_2 = \mathbf{x}_3 = \mathbf{y}_3 = 0$ and $\mathbf{y}_1 = j(\tau_1) j(\tau_2)$ and $\mathbf{y}_2 = (1728 - j(\tau_1)) (1728 - j(\tau_2))$.

APPENDIX C. K3 FIBRATIONS

C.1. The work of Clingher-Doran. Clingher and Doran introduced the following four-parameter quartic family in \mathbb{P}^3 [17, Eq. (3)] with canonical $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ lattice polarization that generalizes a special two-parameter family of K3 surfaces introduced by Inose

$$(C.1) \quad \mathbf{Y}^2 \mathbf{Z} \mathbf{W} - 4 \mathbf{X}^3 \mathbf{Z} + 3 \alpha \mathbf{X} \mathbf{Z} \mathbf{W}^2 + \beta \mathbf{Z} \mathbf{W}^3 + \gamma \mathbf{X} \mathbf{Z}^2 \mathbf{W} - \frac{1}{2} (\delta \mathbf{Z}^2 \mathbf{W}^2 + \mathbf{W}^4) = 0.$$

They also find the parameters $(\alpha, \beta, \gamma, \delta)$ in terms of Siegel modular forms

$$(C.2) \quad (\alpha, \beta, \gamma, \delta) = (\psi_4, \psi_6, 2^{12} 3^5 \mathcal{C}_{10}, 2^{12} 3^6 \mathcal{C}_{12}) .$$

(A similar picture was developed in earlier work for the case of a $H \oplus E_8 \oplus E_8$ lattice polarization [15].)

Clingher and Doran determine an alternate elliptic fibration on (C.1) that has two disjoint sections and a singular fiber of Kodaira-type I_{10}^* . Here, we use a normalization consistent with F-theory and set

$$(C.3) \quad \mathbf{X} = \frac{T X^3}{2^9 3^5}, \quad \mathbf{Y} = \frac{X^2 Y}{2^{15/2} 3^{9/2}}, \quad \mathbf{W} = \frac{X^3}{2^{10} 3^6}, \quad \mathbf{Z} = \frac{X^2}{2^{16} 3^9},$$

and obtain from Equation (C.1) the Jacobian elliptic fibration

$$(C.4) \quad Y^2 = X^3 + \left(T^3 - \frac{\psi_4}{48} T - \frac{\psi_6}{864} \right) X^2 - \left(4 \mathcal{C}_{10} T - \mathcal{C}_{12} \right) X$$

with special fibers of Kodaira-types I_{10}^* , I_2 , and $6 I_1$, and the second section $(Y, X) = (0, 0)$. However, we are interested in the Jacobian elliptic fibration with two distinct special fibers of Kodaira-types II^* and III^* , respectively. Therefore, we set

$$(C.5) \quad \mathbf{X} = \frac{t x}{2^9 3^5}, \quad \mathbf{Y} = \frac{y}{2^{15/2} 3^{9/2}}, \quad \mathbf{W} = \frac{t^3}{2^{10} 3^6}, \quad \mathbf{Z} = \frac{t^2}{2^{16} 3^9},$$

and obtain from Equation (C.1) the Jacobian elliptic fibration

$$(C.6) \quad y^2 = x^3 - t^3 \left(\frac{\psi_4}{48} t + 4 \mathcal{C}_{10} \right) x + t^5 \left(t^2 - \frac{\psi_6}{864} t + \mathcal{C}_{12} \right) .$$

Clingher and Doran also state [17, Thm. 1.7] that

$$(C.7) \quad [I_2 : I_4 : I_6 : I_{10}] = [2^3 3 \delta : 2^2 3^2 \alpha \gamma^2, 2^3 3^2 (4 \alpha \delta + \beta \gamma) \gamma^2 : 2^2 \gamma^6]$$

which equals

$$(C.8) \quad \left[2^3 3 (3 r \mathcal{C}_{12}) : 2^2 3^2 \psi_4 (r \mathcal{C}_{10})^2 : 2^3 3^2 \left(4 \psi_4 (3 r \mathcal{C}_{12}) + \psi_6 (r \mathcal{C}_{10}) \right) (r \mathcal{C}_{10})^2 : 2^2 (r \mathcal{C}_{10})^6 \right]$$

with $r = 2^{12}3^5$. Equation (C.8) implies $\mathcal{C}_{10} = \chi_{10}$ and $\mathcal{C}_{12} = \chi_{12}$ by comparison with Equation (B.30). This choice makes Equation (C.6) also be in agreement with Equation (2.8).¹⁵

C.2. The work of Kumar. To relate this to Kumar's work, we must consider Igusa–Clebsch invariants. Kumar worked with the moduli space of curves of genus 2, which correspond to $\chi_{10} \neq 0$. Kumar's basic theorem [35, Theorem 11] states that a Weierstrass model for a family of K3 surfaces with $\Lambda^{1,1} \oplus E_8(-1) \oplus E_7(-1)$ lattice polarization is given by the equation

$$(C.9) \quad y^2 = x^3 + t^3 (a t + c) x + t^5 (e t^2 + b t + d) ,$$

where t is an affine coordinate on the base \mathbb{P}^1 , x, y are the affine coordinates on the fiber, and the parameters (a, b, c, d, e) are expressed in terms of the Igusa-Clebsch invariants from Section B.3 as follows:

$$(C.10) \quad \begin{aligned} a &= -\frac{I_4}{12} = -\frac{\psi_4(\underline{\mathcal{I}})}{2^6 3^3} , \\ b &= \frac{I_2 I_4 - 3 I_6}{108} = -\frac{\psi_6(\underline{\mathcal{I}})}{2^8 3^6} , \\ c &= -1 , \\ d &= \frac{I_2}{24} = \frac{\chi_{12}(\underline{\mathcal{I}})}{2^3 3 \chi_{10}(\underline{\mathcal{I}})} , \\ e &= \frac{I_{10}}{4} = \frac{\chi_{10}(\underline{\mathcal{I}})}{2^3 3^5} . \end{aligned}$$

Here, we used Equations (B.29) to express the parameters in terms of Siegel modular forms. The discriminant of the elliptic fiber in Equation (C.9) is

$$(C.11) \quad \begin{aligned} \Delta &= t^9 \left(27 e^2 t^5 + 54 e b t^4 + 54 e d t^3 + 27 b^2 t^3 \right. \\ &\quad \left. + 4 a^3 t^3 + 12 a^2 c t^2 + 54 b d t^2 + 12 a c^2 t + 27 d^2 t + 4 c^3 \right) . \end{aligned}$$

Generically, the fibration has a singular fiber of Kodaira-type III^* at $t = 0$ and a singular fiber of Kodaira-type II^* at $t = \infty$. Moreover, there are five singular fibers of Kodaira-type I_1 at those t where the degree-five part of the discriminant vanishes. The Mordell-Weil group is $\text{MW}(\pi) = \{\text{id}\}$, the Néron-Severi lattice has signature $(1, 16)$ and discriminant 2, and the transcendental lattice is $\Lambda^{2,2} \oplus \langle -2 \rangle$.

We know from Section B.2 that $\chi_{10}(\underline{\mathcal{I}}) \rightarrow 0$ as $z \rightarrow 0$ in $\underline{\mathcal{I}}$. For the Weierstrass equation to remain well-defined for $z \rightarrow 0$, we rescale Equation (C.9) as follows

$$(C.12) \quad x \mapsto \frac{x}{\mu^6 \chi_{10}^2(\underline{\mathcal{I}})} , \quad y \mapsto \frac{y}{\mu^9 \chi_{10}^3(\underline{\mathcal{I}})} , \quad t \mapsto \frac{9 t}{\chi_{10}(\underline{\mathcal{I}})}$$

¹⁵Clingher and Doran claim $\mathcal{C}_{12} = \tilde{\chi}_{12}$ instead of $\mathcal{C}_{12} = \chi_{12}$, but we believe that this is the same slip as the one discussed in Equation (B.13)

with $\mu = 2^{1/6}/3^{1/2}$. We then obtain the following rescaled parameters in Equations (C.9)

$$(C.13) \quad a = -\frac{\psi_4(\underline{\tau})}{48}, \quad b = -\frac{\psi_6(\underline{\tau})}{864}, \quad c = -4\chi_{10}(\underline{\tau}), \quad d = \chi_{12}(\underline{\tau}), \quad e = 1.$$

With this choice for the coefficients Equation (C.9) remains well-defined in the limit $z \rightarrow 0$. In fact, setting $z = 0$ we obtain

$$(C.14) \quad a = -3 \frac{E_4(\tau_1) E_4(\tau_2)}{2^4 3^2}, \quad b = -2 \frac{E_6(\tau_1) E_6(\tau_2)}{2^6 3^3}, \quad c = 0, \quad d = \eta(\tau_1)^{24} \eta(\tau_2)^{24}, \quad e = 1,$$

and after rescaling by $y \mapsto \lambda^{18}y$, $x \mapsto \lambda^{12}x$, $t \mapsto \lambda^6 t$ with $\lambda = \eta(\tau_1)^2 \eta(\tau_2)^2$ the Weierstrass equation

$$(C.15) \quad y^2 = x^3 - 3A t^4 x + t^5 (t^2 - 2Bt + 1),$$

with

$$(C.16) \quad A = \frac{E_4(\tau_1) E_4(\tau_2)}{2^4 3^2 \eta(\tau_1)^8 \eta(\tau_2)^8}, \quad B = \frac{E_6(\tau_1) E_6(\tau_2)}{2^6 3^3 \eta(\tau_1)^{12} \eta(\tau_2)^{12}}.$$

Equation (C.15) matches precisely the family presented in [40]. Therefore, this computation provides yet another independent check of the normalization of χ_{12} in Equation (B.11).

APPENDIX D. DEGENERATIONS AND FIVE-BRANES

In this section we consider certain degenerations of the multi-parameter family of K3 surfaces in Equation (C.9). As we have seen, the parameters a, b, c, d can be interpreted as Siegel modular forms of even degree using Equation (C.13) or, equivalently, as the Igusa-Clebsch invariants of a binary sextic using Equation (B.29). On the other hand, Namikawa and Ueno gave a geometrical classification of all (degenerate) fibers in pencils of curves of genus two in [42]. Given a family of curves of genus two over the complex line with affine coordinate $u \in \mathbb{C}$ which is smooth over $\mathbb{C} \setminus \{0\}$, a multi-valued holomorphic map into the Siegel upper half plane of degree two, i.e., the period map, can be defined that determines the family uniquely. Moreover, there are three invariants called ‘monodromy’, ‘modulus point’, and ‘degree’ which determine the singular fiber at $u = 0$ uniquely. To each singular fiber, which is labeled in a fashion similar to Kodaira’s classification of singular fibers of elliptic surfaces, Namikawa and Ueno give a one-parameter family of genus-two curves with a singular fiber of each given type over $u = 0$.

We note that this work of Namikawa and Ueno provides an important class of examples of degenerations in our situation, but cannot be complete. This is because they studied degenerations of genus two curves (with modular group $\mathrm{Sp}_4(\mathbb{Z})$) rather than of binary sextics (with modular group $O^+(L^{2,3})$). Nevertheless, their work provides an interesting first start at studying degenerations and the associated five-branes.

From their list, we took all families of genus-two curves from [42] that develop degenerations of type *III*, in particular, parabolic points of type [3] with monodromy of infinite order over $u = 0$. These families realize all singular fibers with modulus point $\begin{pmatrix} \tau_1 & z \\ z & \infty \end{pmatrix}$

for $z \neq 0$ or $z = 0$ and with $\tau_1 \neq \infty$. The families are listed in Table 2 along with the Namikawa-Ueno type of the singular fiber over $u = 0$ and the modulus point. For the families in the table we computed the Igusa-Clebsch invariants as polynomials in u and determined their asymptotic expansion as $u \rightarrow 0$. By means of Equation (C.9), each of the families of genus-two curves then determines a degenerating family of K3 surfaces as u approaches zero. The degeneration consists of two elliptic surfaces meeting along a rational curve. In the last column of Table 2 we list the Kodaira-types of the singular fibers of these two rational elliptic surfaces. Among these singular fibers, the stable models for the period points $\begin{pmatrix} \tau_1 & z \\ z & \infty \end{pmatrix}$ or $\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$ are given by families with degenerations of Namikawa-Ueno type $[I_{n-0-0}]$ or $[I_n - I_0 - m]$ with $m, n > 0$.

To determine one of the rational components in the degeneration limit, let a, b, c, d be polynomials in u . If $c(u), d(u) \rightarrow 0$ as $u \rightarrow 0$, we obtain a degeneration

$$(D.1) \quad y^2 = x^3 + a(0)t^4x + (t + b(0))t^6,$$

i.e., a $(4, 6, 12)$ -point at $u = 0$. Blowing up by setting $y = t^3y_1$, $x = t^2x_1$, $u = tu_1$, we obtain as the proper transform at $u_1 = 0$ the rational elliptic surface

$$(D.2) \quad y_1^2 = x_1^3 + a(0)x_1 + (t + b(0))$$

with singular fibers of Kodaira-types $II^*, 2I_1$. This rational elliptic surface further degenerates to an isotrivial rational elliptic surface with singular fibers II^*, II and $j = 0$ if $a(0) = 0$.

To determine the second rational component in the degeneration limit, we will have to consider different vanishing orders for the coefficients a, b, c, d corresponding to different Namikawa-Ueno types for the singular fiber of the family of genus-two curves over $u = 0$. As an example, we first consider the family of genus-two curves

$$(D.3) \quad y^2 = u^\kappa (x^3 + \alpha x + 1) ((x - \beta)^2 + u^n)$$

that develops a singularity of Namikawa-Ueno type $[I_{n-0-0}]$ and $[I_{n-0-0}^*]$ with modulus point $\begin{pmatrix} \tau_1 & * \\ * & \infty \end{pmatrix}$ for $\kappa = 0$ and $\kappa = 1$, respectively. One checks that the asymptotic behavior of the Igusa-Clebsch invariants for this family of genus-two curves is given by

$$(D.4) \quad \begin{aligned} a(u) &= a_0 u^{4\kappa} + a_1 u^{4\kappa+n} + O(u^{4\kappa+2n}), \\ b(u) &= b_0 u^{6\kappa} + b_1 u^{6\kappa+n} + O(u^{6\kappa+2n}), \\ c(u) &= c_0 u^{10\kappa+n} + c_1 u^{10\kappa+2n} + O(u^{10\kappa+3n}), \\ d(u) &= d_0 u^{12\kappa+n} + d_1 u^{12\kappa+2n} + O(u^{12\kappa+3n}), \end{aligned}$$

with $n > 0$, $\kappa \in \{0, 1\}$, and $a_0, b_0, c_0, d_0 \neq 0$ and generic. From our discussion above, it follows that the proper transforms in the coordinate chart (u_1, t, x_1, y_1) at $u_1 = 0$ are rational elliptic surfaces with singular fibers $II^*, 2I_1$ and II^*, II for $\kappa = 0$ and $\kappa = 1$, respectively. On the other hand, setting $y = u^{21\kappa+3n}y_2$, $x = u^{14\kappa+2n}x_2$, $t = u^{6\kappa+n}t_2$ we obtain the rational elliptic surface

$$(D.5) \quad y_2^2 = x_2^3 + t_2^3 (a_0 t_2 + c_0) x_2 + t_2^5 (b_0 t_2 + d_0)$$

with singular fibers $III^*, 3I_1$ as the proper transform at $u = 0$.

type	modulus	family of genus-two curves	rat. components
$[I_{n-0-0}]$	$\begin{pmatrix} \tau_1 & * \\ * & \infty \end{pmatrix}$	$y^2 = (x^3 + \alpha x + 1) ((x - \beta)^2 + u^n)$	$II^*, 2I_1$ $III^*, 3I_1$
$[I_{n-0-0}^*]$	$\begin{pmatrix} \tau_1 & * \\ * & \infty \end{pmatrix}$	$y^2 = u (x^3 + \alpha x + 1) ((x - \beta)^2 + u^n)$	II^*, II $III^*, 3I_1$
$[II_{n-0}]$	$\begin{pmatrix} \tau_1 & * \\ * & \infty \end{pmatrix}$	$y^2 = (x^4 + \alpha u x^2 + u^2) ((x - 1)^2 + u^{n-1})$	II^*, II III^*, III
$[II_{n-0}^*]$	$\begin{pmatrix} \tau_1 & * \\ * & \infty \end{pmatrix}$	$y^2 = u (x^4 + \alpha u x^2 + u^2) ((x - 1)^2 + u^{n-1})$	II^*, II III^*, III
$[I_n - I_0 - m]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x^3 + \alpha u^{4m} x + u^{6m}) ((x - 1)^2 + u^n)$	II^*, II $II^*, 2I_1$
$[I_n - I_0^* - m]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x^3 + \alpha u^{4m+2} x + u^{6m+3}) ((x - 1)^2 + u^n)$	II^*, II II^*, II
$[I_n - I_0^* - 0]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x^3 + \alpha u^2 x + u^3) ((x - 1)^2 + u^n)$	II^*, II III^*, II, I_1
$[I_0 - I_n^* - m]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x + u) (x^2 + u^{n+2})$ $\times ((x - 1)^3 + \alpha u^{4m} (x - 1) + u^{6m})$	II^*, II II^*, II
$[I_0 - I_n^* - 0]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x + u) (x^2 + u^{n+2})$ $\times ((x - 1)^3 + \alpha (x - 1) + 1)$	II^*, II III^*, II, I_1
$[I_0^* - I_n^* - m]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x + u) (x^2 + u^{n+2})$ $\times ((x - 1)^3 + \alpha u^{4m+2} (x - 1) + u^{6m+3})$	II^*, II II^*, II
$[I_0^* - I_n^* - 0]$	$\begin{pmatrix} \tau_1 & 0 \\ 0 & \infty \end{pmatrix}$	$y^2 = (x + u) (x^2 + u^{n+2})$ $\times ((x - 1)^3 + \alpha u^2 (x - 1) + u^3)$	II^*, II III^*, II, I_1

TABLE 2. Families of genus-two curves with degeneration of type III

For each example in Table 2, we constructed the corresponding family of degenerating K3 surfaces and recovered the two rational elliptic surfaces in the degeneration limit whose singular fibers are listed in the last column of Table 2. We recorded the leading exponents $\mu(a), \mu(b), \mu(c), \mu(d)$ in the asymptotic expansions of a, b, c, d in Table 3 where $\kappa = 1$ if there is an additional star-fiber and $\kappa = 0$ otherwise. In Table 3, we also recorded the exponents $\mu(y), \mu(x), \mu(u)$ used in the coordinate change $y = u^{\mu(y)} y_2$, $x = u^{\mu(x)} x_2$, $t = u^{\mu(t)} t_2$ that recovers the second rational component in the degeneration limit.

REFERENCES

1. Aldazabal, G., Marques, D., Nunez, C.: Double field theory: A pedagogical review. *Class. Quant. Grav.* **30**, 163,001 (2013)
2. Aspinwall, P.S.: Some relationships between dualities in string theory. *Nucl. Phys. Proc. Suppl.* **46**, 30–38 (1996)
3. Aspinwall, P.S.: Point-like instantons and the $\text{Spin}(32)/\mathbb{Z}_2$ heterotic string. *Nucl. Phys. B* **496**, 149–176 (1997)
4. Aspinwall, P.S., Douglas, M.R.: D-brane stability and monodromy. *JHEP* **05**, 031 (2002)

type	$\mu(a)$	$\mu(b)$	$\mu(c)$	$\mu(d)$	$\mu(y)$	$\mu(x)$	$\mu(t)$
$[I_{n-0-0}]$ $[I_{n-0-0}^*]$	4κ	6κ	$10\kappa + n$	$12\kappa + n$	$21\kappa + 3n$	$14\kappa + 2n$	$6\kappa + n$
$[II_{n-0}]$ $[II_{n-0}^*]$	$2 + 4\kappa$	$3 + 6\kappa$	$5 + 10\kappa$ $+ n$	$6 + 12\kappa$ $+ n$	$15 + 21\kappa$ $+ 3n$	$10 + 14\kappa$ $+ 2n$	$5 + 6\kappa$ $+ n$
$[I_n - I_0 - m]$ $[I_n - I_0^* - m]$	$2\kappa + 4m$	$3\kappa + 6m$	$6\kappa + 12m$ $+ n$	$6\kappa + 12m$ $+ n$	$18\kappa + 21m$ $+ 3n$	$12\kappa + 14m$ $+ 2n$	$6\kappa + 6m$ $+ n$
$[I_0 - I_n^* - m]$ $[I_0^* - I_n^* - m]$	$2\kappa + 4m$ $+ 2$	$3\kappa + 6m$ $+ 3$	$6\kappa + 12m$ $+ 6 + n$	$6\kappa + 12m$ $+ 6 + n$	$18\kappa + 21m$ $+ 18 + 3n$	$12\kappa + 14m$ $+ 12 + 2n$	$6\kappa + 6m$ $+ 6 + n$

TABLE 3. Exponents of coefficients for families of K3 surfaces

5. Aspinwall, P.S., Gross, M.: The $SO(32)$ heterotic string on a $K3$ surface. *Phys. Lett. B* **387**(4), 735–742 (1996)
6. Aspinwall, P.S., Morrison, D.R.: String theory on $K3$ surfaces. In: B. Greene, S.T. Yau (eds.) *Mirror Symmetry II*, pp. 703–716. International Press, Cambridge (1997)
7. Aspinwall, P.S., Morrison, D.R.: Non-simply-connected gauge groups and rational points on elliptic curves. *J. High Energy Phys.* **07**, 012 (1998)
8. Baily Jr., W.L., Borel, A.: Compactification of arithmetic quotients of bounded symmetric domains. *Ann. of Math.* (2) **84**, 442–528 (1966)
9. Berman, D.S., Thompson, D.C.: Duality symmetric string and M-theory. *Phys. Rept.* **566**, 1–60 (2014)
10. Bershadsky, M., Intriligator, K., Kachru, S., Morrison, D.R., Sadov, V., Vafa, C.: Geometric singularities and enhanced gauge symmetries. *Nuclear Phys. B* **481**, 215–252 (1996)
11. Bolza, O.: On binary sextics with linear transformations into themselves. *Amer. J. Math.* **10**(1), 47–70 (1887)
12. Borcea, C.: Diffeomorphisms of a $K3$ surface. *Math. Ann.* **275**(1), 1–4 (1986)
13. Braun, A.P., Fucito, F., Morales, J.F.: U-folds as $K3$ fibrations. *JHEP* **1310**, 154 (2013)
14. Cachazo, F.A., Vafa, C.: Type I' and real algebraic geometry. *ArXiv:hep-th/0001029*
15. Clingher, A., Doran, C.F.: Modular invariants for lattice polarized $K3$ surfaces. *Michigan Math. J.* **55**(2), 355–393 (2007)
16. Clingher, A., Doran, C.F.: Note on a geometric isogeny of $K3$ surfaces. *Int. Math. Res. Not. IMRN* **2011**(16), 3657–3687 (2011)
17. Clingher, A., Doran, C.F.: Lattice polarized $K3$ surfaces and Siegel modular forms. *Adv. Math.* **231**(1), 172–212 (2012)
18. Curio, G.: $N = 2$ string-string duality and holomorphic couplings. *Fortsch. Phys.* **46**, 75–146 (1998)
19. Deligne, P.: Courbes elliptiques: formulaire (d'après J. Tate). In: *Modular functions of one variable, IV* (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), *Lecture Notes in Math.*, vol. 476, pp. 53–73. Springer, Berlin (1975)
20. Dolgachev, I.V.: Mirror symmetry for lattice polarized $K3$ surfaces. *J. Math. Sci.* **81**(3), 2599–2630 (1996)
21. Donaldson, S.K.: Polynomial invariants for smooth four-manifolds. *Topology* **29**(3), 257–315 (1990)
22. Douglas, M.R.: D-branes, categories and $N = 1$ supersymmetry. *J. Math. Phys.* **42**, 2818–2843 (2001)
23. Douglas, M.R.: D-branes and $\mathcal{N} = 1$ supersymmetry. In: *Strings, 2001 (Mumbai)*, *Clay Math. Proc.*, vol. 1, pp. 139–152. Amer. Math. Soc., Providence, RI (2002)

24. Freitag, E.: Siegelsche Modulfunktionen, *Grundlehren der Mathematischen Wissenschaften*, vol. 254. Springer-Verlag, Berlin (1983)
25. Friedman, R., Morgan, J., Witten, E.: Vector bundles and F theory. *Commun. Math. Phys.* **187**, 679–743 (1997)
26. Gritsenko, V.A., Hulek, K., Sankaran, G.K.: The Kodaira dimension of the moduli of $K3$ surfaces. *Invent. Math.* **169**(3), 519–567 (2007)
27. Gritsenko, V.A., Nikulin, V.V.: Igusa modular forms and “the simplest” Lorentzian Kac-Moody algebras. *Mat. Sb.* **187**(11), 27–66 (1996). English translation in *Sb. Math.* **187**(11), 1601–1641 (1996)
28. Hellerman, S., McGreevy, J., Williams, B.: Geometric constructions of nongeometric string theories. *JHEP* **01**, 024 (2004)
29. Hohm, O., Lüst, D., Zwiebach, B.: The spacetime of double field theory: Review, remarks, and outlook. *Fortsch. Phys.* **61**, 926–966 (2013)
30. Hosono, S., Lian, B.H., Oguiso, K., Yau, S.T.: $c = 2$ rational toroidal conformal field theories via the Gauss product. *Comm. Math. Phys.* **241**(2-3), 245–286 (2003)
31. Igusa, J.: On Siegel modular forms of genus two. *Amer. J. Math.* **84**, 175–200 (1962)
32. Igusa, J.: Modular forms and projective invariants. *Amer. J. Math.* **89**, 817–855 (1967)
33. Igusa, J.i.: On the ring of modular forms of degree two over \mathbf{Z} . *Amer. J. Math.* **101**(1), 149–183 (1979)
34. Kodaira, K.: On compact analytic surfaces. II, III. *Ann. of Math. (2)* **77**, 563–626 (1963). **78**, 1–40 (1963)
35. Kumar, A.: $K3$ surfaces associated with curves of genus two. *Int. Math. Res. Not.* **2008**(6), Art. ID rnm165 (2008)
36. Martucci, L., Morales, J.F., Pacifici, D.R.: Branes, U-folds and hyperelliptic fibrations. *JHEP* **1301**, 145 (2013)
37. Matumoto, T.: On diffeomorphisms of a $K3$ surface. In: *Algebraic and topological theories* (Kinoshita, 1984), pp. 616–621. Kinokuniya, Tokyo (1986)
38. McOrist, J., Morrison, D.R., Sethi, S.: Geometries, non-geometries, and fluxes. *Adv. Theor. Math. Phys.* **14**, 1515–1583 (2010)
39. Mestre, J.F.: Construction de courbes de genre 2 à partir de leurs modules. In: *Effective methods in algebraic geometry* (Castiglione, 1990), *Progr. Math.*, vol. 94, pp. 313–334. Birkhäuser Boston (1991)
40. Morrison, D.R., Vafa, C.: Compactifications of F-theory on Calabi–Yau threefolds, II. *Nuclear Phys. B* **476**, 437–469 (1996)
41. Nahm, W., Wendland, K.: A hiker’s guide to $K3$: Aspects of $N = (4,4)$ superconformal field theory with central charge $c = 6$. *Commun. Math. Phys.* **216**, 85–138 (2001)
42. Namikawa, Y., Ueno, K.: The complete classification of fibres in pencils of curves of genus two. *Manuscripta Math.* **9**, 143–186 (1973)
43. Narain, K.S.: New heterotic string theories in uncompactified dimensions < 10 . *Phys. Lett. B* **169**, 41–46 (1986)
44. Narain, K.S., Sarmadi, M.H., Witten, E.: A note on toroidal compactification of heterotic string theory. *Nucl. Phys. B* **279**, 369–379 (1987)
45. Néron, A.: Modèles minimaux des variétés abéliennes sur les corps locaux et globaux. *Inst. Hautes Études Sci. Publ. Math.* **21**, 5–128 (1964)
46. Nikulin, V.V.: Integral symmetric bilinear forms and some of their applications. *Izv. Akad. Nauk SSSR Ser. Mat.* **43**(1), 111–177, 238 (1979). English translation: *Math USSR Izvestija* **14**, 103–167 (1980)
47. Satake, I.: On representations and compactifications of symmetric Riemannian spaces. *Ann. of Math. (2)* **71**, 77–110 (1960)
48. Schwarz, J.H.: An $SL(2, \mathbf{Z})$ multiplet of type IIB superstrings. *Phys. Lett. B* **360**, 13–18 (1995)

- 49. Vafa, C.: Evidence for F-theory. Nucl. Phys. B **469**, 403–418 (1996)
- 50. Vinberg, E.B.: On the algebra of Siegel modular forms of genus 2. Trans. Moscow Math. Soc. **2013**, 1–13 (2013). Translation of: Trudy Moskovskogo Matematicheskogo Obshchestva, tom **74**, vypusk 1 (2013)

DEPARTMENT OF MATHEMATICS AND STATISTICS, COLBY COLLEGE, WATERVILLE, ME 04901

Current address: Department of Mathematics and Statistics, Utah State University, Logan, UT 84322

E-mail address: andreas.malmendier@usu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106

E-mail address: drm@math.ucsb.edu