A class of new bi-invariant metrics on the Hamiltonian diffeomorphism groups

Guangcun Lu * Tie Sun

School of Mathematical Sciences, Beijing Normal University Laboratory of Mathematics and Complex Systems, Ministry of Education Beijing 100875, The People's Republic of China (gclu@bnu.edu.cn suntie@mail.bnu.edu.cn)

March 1, 2022

Abstract

In this paper, we construct infinitely many bi-invariant metrics on the Hamiltonian diffeomorphism group and study their basic properties and corresponding generalizations of the Hofer inequality and Sikorav one.

Key words: Hamiltonian diffeomorphism, Hofer's metric, bi-invariant Finsler metrics

1 Introduction and Main Results

1.1 The Hofer metric

In 1989, H. Hofer [12] constructed a remarkable bi-invariant Finsler metric on the group of compactly supported Hamiltonian diffeomorphisms $\operatorname{Ham}(M, \omega)$ of a symplectic manifold (M, ω) , nowadays known as Hofer metric. Since then the intrinsic geometry of it has been being a very active and fruitful research field in symplectic topology and Hamiltonian dynamics (see the books [14, 18, 27], and the surveys [10, 19, 28, 24] and references therein for current progress situation).

Especially, a recent celebrated result made by Buhovsky and Ostrover [5] is a positive answer to the uniqueness question of the Hofer metric raised by Eliashberg

^{*2010} Mathematics Subject Classification. Primary 53D22, 53D40, 53D25.

Partially supported by the NNSF 10971014 and 11271044 of China, PCSIRT, RFDPHEC (No. 200800270003) and the Fundamental Research Funds for the Central Universities (No. 2012CXQT09).

and Polterovich [9]. They showed that up to equivalence of metrics the Hofer metric is the only bi-invariant Finsler metric on the group of Hamiltonian diffeomorphisms of a closed symplectic manifold under a natural assumption. For studies of non-Finslerian bi-invariant metrics on $\text{Ham}(M, \omega)$ the readers may refer to [31, 29, 22].

Let us briefly review the construction of the Hofer metric following the notations in [27] without special statements. The readers who are familiar with it may directly read the next section. Let (M^{2n}, ω) be a connected symplectic manifold of dimension 2n without boundary. Denote by $\mathcal{A}(M)$ the space of all smooth functions on M with compact support (resp. zero-mean with respect to the canonical volume form ω^n) if M is open (resp. closed). A (time-dependent) smooth Hamiltonian function F on $M \times I$, where $I \subset \mathbb{R}$ is an interval, is called **normalized** if $F_t = F(\cdot, t)$ belongs to $\mathcal{A}(M)$ for all t, and $\bigcup_{t \in I} \operatorname{supp}(F_t)$ is contained in a compact subset of M in the case when M is open. Such a normalized F determines a (time-dependent) Hamiltonian vector field X_{F_t} on M via $i_{X_{F_t}}\omega = -dF_t$, and when I = [0, 1] the corresponding flow $\{f_t\}$ starting from the identity is called a **Hamiltonian isotopy** generated by F and is also denoted by $\{\phi_{E}^{t}\}$ for convenience. A diffeomorphism of M is said to be Hamiltonian if it can be represented as a time-one map of some Hamiltonian isotopy. Denote by $\operatorname{Ham}(M,\omega)$ the set of all Hamiltonian diffeomorphisms on (M,ω) . It is a subgroup of $\text{Symp}(M, \omega)$. The Lie algebra of $\text{Ham}(M, \omega)$ can be naturally identified with $\mathcal{A}(M)$ because of the following important fact by Banyaga (cf. [27, Prop.1.4.B]).

Lemma 1.1 ([1]) For every smooth path $\{f_t\}$ in $\operatorname{Ham}(M, \omega), t \in [a, b]$, there exists a unique (time-dependent) normalized Hamiltonian function $F: M \times [a, b] \to \mathbb{R}$ such that

$$\frac{d}{dt}f_t(x) = X_{F_t}(f_t(x)) \quad \forall (x,t) \in M \times [a,b].$$
(1.1)

Hence the tangent vector of the path $\{f_t\}$ at t = s is the function F_s . The adjoint action of $\operatorname{Ham}(M, \omega)$ on $\mathcal{A}(M)$ is given by $\operatorname{Ad}_f G = G \circ f^{-1}$ for $f \in \operatorname{Ham}(M, \omega)$ and $G \in \mathcal{A}(M)$. Any adjoint invariant norm $\|\cdot\|$ on $\mathcal{A}(M)$, i.e., $\|\operatorname{Ad}_f G\| = \|G \circ f^{-1}\| = \|G\|$ for any $f \in \operatorname{Ham}(M, \omega)$ and $G \in \mathcal{A}(M)$, defines a Finsler structure on $\operatorname{Ham}(M, \omega)$, and thus the length of a Hamiltonian path $\{f_t\}, t \in [a, b]$ with (unique) normalized Hamiltonian F by

$$\text{Length}\{f_t\} = \int_a^b \|F_t\| dt, \qquad (1.2)$$

which does not depend on the parametrization. Without loss of generality we could fix a = 0, b = 1 in the definition. For arbitrary $\phi, \varphi \in \text{Ham}(M, \omega)$, their pseudo-distance is defined by

$$d(\phi, \varphi) = \inf\{\text{Length}(\alpha)\}$$
(1.3)

where the infimum is taken over all smooth Hamiltonian path $\alpha : [a, b] \to \operatorname{Ham}(M, \omega)$ with $\alpha(a) = \phi$ and $\alpha(b) = \varphi$. It is a bi-invariant pseudo-metric. (If the norm $\|\cdot\|$ on $\mathcal{A}(M)$ is not adjoint invariant, d is only right-invariant.) The non-degeneracy axiom, $d(\phi, \varphi) > 0$ for $\phi \neq \varphi$, is not satisfied in general. When the norm $\|\cdot\|$ is chosen as the L_{∞} -norm,

$$||H||_{\infty} := \max H - \min H, \quad \text{for} \quad H \in \mathcal{A}(M),$$

Hofer showed in [12] that the corresponding pseudo-metric d_H is a genuine metric in the case $M = \mathbb{R}^{2n}$. Later, this result was generalized to some larger class of symplectic manifolds by Polterovich [26], and finally to general manifolds by Lalonde and McDuff [17]. Nowadays this bi-invariant Finsler metric on the group Ham (M, ω) is called the **Hofer metric**, and the function $\|\cdot\|_H = d_H(\cdot, \mathrm{id}_M) : \mathrm{Ham}(M, \omega) \to \mathbb{R}$ is called the **Hofer norm**. Let

$$\mathcal{H} = \mathcal{H}(M) = \{ H \in C^{\infty}(M \times [0,1]) \mid H_t = H(\cdot,t) \in \mathcal{A}(M) \; \forall t \in [0,1] \},$$
$$\mathcal{F} = \mathcal{F}(M) = \{ H \in C^{\infty}(M \times \mathbb{R}/\mathbb{Z}) \mid H_t = H(\cdot,t) \in \mathcal{A}(M) \; \forall t \in \mathbb{R} \}.$$

Every $\phi \in \operatorname{Ham}(M, \omega)$ can be written as ϕ_F^1 with $F \in \mathcal{F}$. Moreover it holds that

$$d_{H}(\phi,\varphi) = \inf \left\{ \int_{0}^{1} \|H_{t} - K_{t}\|_{\infty} dt \mid H \in \mathcal{H} \text{ generates } \varphi \& K \in \mathcal{H} \text{ generates } \phi \right\},\$$

$$d_{H}(\mathrm{id}_{M},\varphi) = \inf \left\{ \max_{t} \|F_{t}\|_{\infty} \mid F \in \mathcal{F} \text{ generates } \varphi \right\},\tag{1.4}$$

see [14, (5.9)] for the first one, and [27, Lemma 5.1.C] for the second.

1.2 New bi-invariant metrics

Our new bi-invariant metrics on $\operatorname{Ham}(M, \omega)$ will be constructed in a similar way to Hofer's. For a smooth path $f : [0, 1] \to \operatorname{Ham}(M, \omega)$ generated by a (time-dependent) normalized Hamiltonian function F, and each integer $k = 0, 1, 2 \cdots$, we define the **k-length** of f by

$$\operatorname{Length}_k(f) := \sum_{i=0}^k \int_0^1 \left\| \frac{\partial^i F_t}{\partial t^i} \right\|_\infty dt = \sum_{i=0}^k \int_0^1 \left(\max_x \frac{\partial^i F_t}{\partial t^i} - \min_x \frac{\partial^i F_t}{\partial t^i} \right) dt.$$

Clearly, Length₀ is the same as that of (1.2). However, unlike Length₀ the k-length $(k \ge 1)$ strongly depends on the choice of parametrization.

Call a continuous path $f: [0,1] \to \operatorname{Ham}(M,\omega)$ piecewise smooth if there exists a division $0 = t_0 < t_1 < \cdots < t_n = 1, n \in \mathbb{N}$, such that $f_i = f|_{[t_{i-1},t_i]}$ is smooth for $i = 1, \cdots, n$. Let $F = \{F^l\}_{l=1}^n$ be the corresponding normalized Hamiltonian function, that is,

$$\frac{d}{dt}f_t(x) = X_{F_t^i}(f_t(x)) \quad \forall (x,t) \in M \times [t_{i-1}, t_i], i = 1, \cdots, n.$$

Define its k-length by

$$\operatorname{Length}_{k}(f) := \sum_{i=1}^{n} \operatorname{Length}_{k}(f_{i}) = \sum_{l=1}^{n} \sum_{i=0}^{k} \int_{t_{l-1}}^{t_{l}} \left\| \frac{\partial^{i} F_{t}^{l}}{\partial t^{i}} \right\|_{\infty} dt.$$
(1.5)

For $\phi, \varphi \in \operatorname{Ham}(M, \omega)$, let $\Omega(\phi, \varphi)$ denote the space of all continuous and piecewise smooth paths $f : [0, 1] \to \operatorname{Ham}(M, \omega)$ from $f(0) = \phi$ to $f(1) = \varphi$. Then we define pseudo-distances between ϕ and φ by

$$d_k(\phi,\varphi) = \inf \left\{ \operatorname{Length}_k(f) \middle| f \in \Omega(\phi,\varphi) \right\}, \quad k = 0, 1, \cdots.$$
 (1.6)

Lemma 1.2 $d_0 = d_H$ on Ham (M, ω) .

Clearly, $d_0 \leq d_H$ since (1.2) does not depend on the parametrization. The converse inequality easily follows from the definition of d_0 and the triangle inequality for d_H .

Let us make some comments on the definition of d_k .

Remark 1.3 (i) For a positive integer k, the function $\text{Length}_k(\cdot)$ depends on the choice of parametrization of the path, and the derivative terms will in fact vanish when we take the infimum with respect to the path space with variant parametrization intervals. In fact, suppose that a smooth Hamiltonian path $\alpha : [0, 1] \to \text{Ham}(M, \omega)$ is generated by a normalized Hamiltonian F_t . For any b > 0, the reparametrized path

$$\beta_b(t) := \alpha(t/b) : [0, b] \to \operatorname{Ham}(M, \omega)$$

is generated by the Hamiltonian function $G(x,t) = \frac{1}{b}F(x,\frac{t}{b})$, and hence

$$\int_0^b \left\| \frac{\partial G_t}{\partial t} \right\|_\infty dt = \int_0^b \frac{1}{b^2} \left\| \frac{\partial F}{\partial t}(x, \frac{s}{b}) \right\|_\infty ds = \frac{1}{b} \int_0^1 \left\| \frac{\partial F}{\partial t}(x, t) \right\|_\infty dt \to 0$$

as $b \to +\infty$. This fact still holds for the higher order derivatives. It follows that

 $b \to +\infty \Longrightarrow \operatorname{Length}_k(\beta_b) \to \operatorname{Length}_0(\alpha).$

Thus if we define d_k as in (1.3) no new thing can be obtained. This is why we fix the parametrization interval of paths, [a, b] = [0, 1].

(ii) Since the definition of k-length involves the derivatives of a Hamiltonian function until k order the quasi-triangle inequality cannot be obtained if we restrict to smooth paths from [0, 1] to $\operatorname{Ham}(M, \omega)$ in the definition of d_k . It is this reason that we extend the space of all smooth Hamiltonian paths to include the piecewise smooth ones.

As expected d_k has the following properties.

Theorem 1.4 $d_H = d_0 \leq d_1 \leq d_2 \leq \cdots$, and

- (i) (symmetry) $d_k(\phi, \varphi) = d_k(\varphi, \phi)$,
- (ii) (quasi-triangle inequality) $d_k(\phi, \varphi) \leq 2^k (d_k(\phi, \theta) + d_k(\theta, \varphi)),$
- (iii) (non-degeneracy) $d_k(\phi, \varphi) \ge 0$, and $d_k(\phi, \varphi) = 0 \iff \phi = \varphi$,
- (iv) (bi-invariance) $d_k(\phi, \varphi) = d_k(\phi\theta, \varphi\theta) = d_k(\theta\phi, \theta\varphi),$

for any $\phi, \varphi, \theta \in \operatorname{Ham}(M, \omega)$ and $k = 0, 1, \cdots$.

This theorem shows that (1.6) gives a sequence of bi-invariant quasi-metrics $\{d_k\}_{k=0}^{\infty}$ on $\operatorname{Ham}(M, \omega)$. Recall that a **quasidistance** on a nonempty set X is a function $\rho : X \times X \to [0, +\infty)$ such that (i) $\rho(x, y) = \rho(y, x)$ for all $x, y \in X$, (ii) $\rho(x, y) = 0$ if and only if x = y, (iii) and there exists a finite constant $c \ge 1$ (quasi-triangle constant) such that $\rho(x, y) \le c(\rho(x, z) + \rho(z, y))$ for every $x, y, z \in X$. Such a pair (X, ρ) is called a **quasimetric** space. (See [20]). A **group norm** (resp. **quasinorm**) on a group G is a symmetric, nondegenerate and nonnegative function ψ which is subadditive (resp. C-subadditive for some finite constant $C \ge 1$, that is, $\psi(xy) \le C(\psi(x) + \psi(y))$ for all $x, y \in G$). (See [20, page 113]).

For every d_k , let us define a function

$$\|\cdot\|_k = d_k(\cdot, \mathrm{id}_M). \tag{1.7}$$

From Theorem 1.4 one easily derives:

Theorem 1.5 $\|\cdot\|_0 = \|\cdot\|_H$, and for every $k \in \mathbb{N}$, $\|\cdot\|_k$ is a quasinorm, precisely speaking it satisfies:

- (i) (Symmetry) $\|\phi^{-1}\|_k = \|\phi\|_k$,
- (ii) (The quasi-triangle inequality) $\|\phi\varphi\|_k \leq 2^k (\|\phi\|_k + \|\varphi\|_k),$
- (iii) (Non-degeneracy) $\|\phi\|_k \ge 0$, and $\|\phi\|_k = 0 \iff \phi = \mathrm{id}_M$,
- (iv) (Conjugate invariance) $\|\theta\varphi\theta^{-1}\|_k = \|\varphi\|_k$,

where $\phi, \varphi, \theta \in \operatorname{Ham}(M, \omega)$ are arbitrary.

This shows that every $\|\cdot\|_k$ is a conjugate invariant quasinorm on the group $\operatorname{Ham}(M,\omega)$. Apply Theorem A.1 to $G = \operatorname{Ham}(M,\omega)$ and $\psi = \|\cdot\|_k$, $k \in \mathbb{N}$, we get

Theorem 1.6 Define the function $||| \cdot |||_k : \operatorname{Ham}(M, \omega) \to [0, \infty)$ by

$$\|\|\phi\|\|_{k} := \inf\left\{ \left(\sum_{i=1}^{N} \|\phi_{i}\|_{k}^{\frac{1}{k+1}} \right)^{1+k} \middle| \qquad N \in \mathbb{N}, \ (\phi_{1}, \cdots, \phi_{N}) \in \operatorname{Ham}(M, \omega)^{(N)}, \\ \phi = \phi_{1} \cdots \phi_{N} \right\},$$
(1.8)

where $\operatorname{Ham}(M,\omega)^{(N)} = \underbrace{\operatorname{Ham}(M,\omega) \times \cdots \times \operatorname{Ham}(M,\omega)}_{N}, N \in \mathbb{N}.$ Then

$$\||\phi|||_k = |||\phi^{-1}|||_k, \quad \forall \phi \in \operatorname{Ham}(M, \omega),$$
(1.9)

$$\||\theta\phi\theta^{-1}||_{k} = |||\phi|||_{k}, \quad \forall \theta, \phi \in \operatorname{Ham}(M, \omega),$$
(1.10)

$$4^{-(1+k)} \|\phi\|_{k} \le \|\phi\|_{k} \le \|\phi\|_{k}, \quad \forall \phi \in \operatorname{Ham}(M, \omega),$$
(1.11)

and

$$\| \phi \varphi \|_{k}^{\beta} \leq \| \phi \|_{k}^{\beta} + \| \varphi \|_{k}^{\beta}, \quad \forall \phi, \varphi \in \operatorname{Ham}(M, \omega)$$
(1.12)

for each $\beta \in (0, \frac{1}{1+k}]$. Also, for each $N \in \mathbb{N}$, $\beta \in (0, \frac{1}{1+k}]$, and $\phi_i \in \operatorname{Ham}(M, \omega)$, $i = 1, \dots, N$, it holds that

$$\|\phi_1 \cdots \phi_N\|_k \le 4^{k+1} \left\{ \sum_{i=1}^N \|\phi_i\|_k^\beta \right\}^{\frac{1}{\beta}}, \qquad (1.13)$$

and hence for each sequence $(\phi_i)_{i \in \mathbb{N}} \subset \operatorname{Ham}(M, \omega)$,

$$\sup_{N \in \mathbb{N}} \|\phi_1 \cdots \phi_N\|_k \le 4^{k+1} \left\{ \sum_{i=1}^{\infty} \|\phi_i\|_k^{\beta} \right\}^{\frac{1}{\beta}}.$$
 (1.14)

(1.9) can be derived from (1.8) and the symmetry of $\|\cdot\|_k$, and (1.10) can be obtained by Remark A.2 and the conjugate invariance of $\|\cdot\|_k$.

Corollary 1.7 The function $||| \cdot |||_k$ defined in (1.8) is a conjugate invariant quasinorm on $\operatorname{Ham}(M, \omega)$ which is equivalent to $|| \cdot ||_k$; and for every $\beta \in (0, \frac{1}{1+k}]$, $||| \cdot |||_k^\beta$ is a conjugate invariant norm on $\operatorname{Ham}(M, \omega)$. Thus

$$\tilde{d}_k(\phi,\varphi) := \| \phi \varphi^{-1} \|_k \tag{1.15}$$

is a bi-invariant quasimetric on $\operatorname{Ham}(M, \omega)$; and for each $\beta \in (0, \frac{1}{1+k}]$,

$$\tilde{d}_k^\beta(\phi,\varphi) := (\tilde{d}_k(\phi,\varphi))^\beta \tag{1.16}$$

is a bi-invariant metric on $\operatorname{Ham}(M, \omega)$. They all induce the same topology as d_k .

Consider the commutator of two elements φ and ψ in $\operatorname{Ham}(M,\omega)$, $[\varphi,\psi] := \varphi \psi \varphi^{-1} \psi^{-1}$. It follows from Theorem 1.5 that

$$\|[\varphi,\psi]\|_{k} \le 2^{k+1} \min\{\|\varphi\|_{k}, \|\psi\|_{k}\}.$$
(1.17)

Similarly (1.9)-(1.12) lead to

$$\| [\varphi, \psi] \|_{k}^{\beta} \le 2 \min\{ \| \varphi \|_{k}^{\beta}, \| \psi \|_{k}^{\beta} \}$$
(1.18)

for each $\beta \in (0, \frac{1}{1+k}]$. For a non-empty subset $A \subset M$ let $e_k(A)$ (resp. $\tilde{e}_k(A)$) denote the **displacement energy** of it with respect to $\|\cdot\|_k$ (resp. $\|\cdot\|_k$), that is,

$$e_k(A) = \inf\{\|\vartheta\|_k \mid \vartheta \in \operatorname{Ham}(M,\omega) \& A \cap \vartheta(A) = \emptyset\},$$
(1.19)

$$\tilde{e}_k(A) = \inf\{ \| \vartheta \|_k \mid \vartheta \in \operatorname{Ham}(M, \omega) \& A \cap \vartheta(A) = \emptyset \}.$$
(1.20)

As in the proof of [9, Lemma 2.3.B] we may obtain:

Theorem 1.8 Let $U \subset M$ be a non-empty open subset. Then for any $\varphi, \psi \in \text{Ham}(M, \omega)$ with $\text{supp}(\varphi) \subset U$ and $\text{supp}(\psi) \subset U$ it holds that

$$\|[\varphi,\psi]\|_{k} \le 4^{k+1}e_{k}(U) \quad and \quad \|\|[\varphi,\psi]\|_{k}^{\beta} \le 4(\tilde{e}_{k}(U))^{\beta}$$
(1.21)

for each $\beta \in (0, \frac{1}{1+k}]$.

Motivated by the so-called "coarse" Hofer norm, for $f = \{f_t\} \in \Omega(\phi, \psi)$, if $F = \{F^l\}_{l=1}^n$ is the corresponding normalized Hamiltonian function, we use

Length^{*}_k(f) := |||F|||_k :=
$$\sum_{i=0}^{k} \max_{1 \le l \le n} \max_{t_{l-1} \le t \le t_l} \left\| \frac{\partial^i F^l}{\partial t^i}(x, t) \right\|_{\infty}$$
 (1.22)

(which is independent of the choices of divisions) to replace (1.5), and obtain another sequence of bi-invariant quasimetrics

$$d_k^*(\phi,\varphi) = \inf \left\{ \operatorname{Length}_k^*(f) \, \middle| \, f \in \Omega(\phi,\varphi) \right\}, \quad k = 0, 1, \cdots$$
 (1.23)

as in (1.6). Clearly, $d_k \leq d_k^*$ for any $k \in \mathbb{N} \cup \{0\}$. (1.4) implies $d_0^* = d_0 = d_H$. Let $\|\cdot\|_k^* = d_k^*(\cdot, \mathrm{id}_M)$. Correspondingly, we have also $\|\|\cdot\|_k^*$ as in (1.8), and \tilde{d}_k^* , e_k^* and \tilde{e}_k^* .

Theorem 1.9 All conclusions from Theorem 1.4 to Theorem 1.8 still hold for $\|\cdot\|_k^*$, d_k^* , $\|\cdot\|\|_k^*$ and \tilde{d}_k^* , e_k^* and \tilde{e}_k^* , but we need to add a factor 2 for the coefficients of inequalities in (ii) of Theorems 1.4, 1.5, (1.17) and the factor 4 in the first inequality of (1.21).

The following result shows that the Hofer inequality in [13] also holds for each d_k^* .

Theorem 1.10 For every $\varphi, \psi \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0)$,

$$d_k^*(\varphi,\psi) := \|\varphi\psi^{-1}\|_k^* \le C \operatorname{diameter}\left(\operatorname{supp}(\varphi\psi^{-1})\right)|\varphi - \psi|_{C^0},$$

where C is a constant and $C \leq 2^{3k+8}(k+1)^2(1+2^{k+1}+2^{2k+2}+2^{3k+3})$. (Note: if k=0 the constant C can be chosen as 128 as in the Hofer inequality.)

Similarly, for any subset $S \subset \mathbb{R}^{2n}$, we define the **coarse proper displacement** *k*-energy $e_{p,k}^*(S)$ of it as

$$e_{p,k}^*(S) = \{a > 0 \mid \text{for every bounded subset } A \subset S \exists \psi \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0) \\ \text{such that } \|\psi\|_k^* \leq a \text{ and } A \text{ and } \psi(A) \text{ are properly separated} \},$$

and get the following generalization of the Sikorav inequality.

Theorem 1.11 If $H \in \mathcal{H}(\mathbb{R}^{2n})$ satisfies $\operatorname{supp}(H) \subset U \times [0,1]$, then

$$\|\varphi_H\|_k^* \le 2^{2k+4}(k+1)\left(1+2^{k+1}+2^{2k+2}+2^{3k+3}\right)e_{p,k}^*(U).$$

Finally, let us discuss the corresponding question investigated by Eliashberg and Polterovich [9]. When 0 the function

$$\mathcal{A}(M) \ni H \to ||H||_p = \left(\int_M |H|^p \omega^n\right)^{\frac{1}{p}}$$

is an adjoint invariant quasinorm because

$$\int_{M} |G(f^{-1}(x))|^{p} \omega^{n} = \int_{M} (f^{-1})^{*} (|G(x)|^{p} \omega^{n}) = \int_{M} |G(x)|^{p} \omega^{n}$$
(1.24)

for any $G \in \mathcal{A}(M)$ and $f \in \text{Symp}(M, \omega)$, and

$$||H + G||_p \le K_p(||H||_p + ||G||_p), \quad \forall H, G \in \mathcal{A}(M),$$

where K_p is equal to 1 for $p \ge 1$, and $2^{\frac{1-p}{p}}$ for $0 (so <math>\|\cdot\|_p$ is only a quasinorm on $\mathcal{A}(M)$ in this case).

For a smooth path $f : [0,1] \to \operatorname{Ham}(M,\omega)$ generated by a (time-dependent) normalized Hamiltonian function F_t , and each integer $k = 0, 1, 2 \cdots$, we define the **(k,p)-length** of f by

$$\operatorname{Length}_{(k,p)}(f) := \sum_{i=0}^{k} \int_{0}^{1} \left\| \frac{\partial^{i} F_{t}}{\partial t^{i}} \right\|_{p} dt, \qquad (1.25)$$

and the **(k,p)-length** of $f \in \Omega(\phi, \varphi)$ by the sum of (k, p)-lengths of all smooth pieces of it. By the same proofs as those of Theorem 1.4 it is readily verified that

$$d_{(k,p)}(\phi,\varphi) = \inf \left\{ \operatorname{Length}_{(k,p)}(f) \, \middle| \, f \in \Omega(\phi,\varphi) \right\}$$
(1.26)

defines a pseudo quasimetric $d_{(k,p)}$ on $\operatorname{Ham}(M,\omega)$ for any $p > 0, k = 0, 1, \cdots$. Note that

$$d_{(0,p)}(\phi,\varphi) = \inf \left\{ \operatorname{Length}_{(0,p)}(f) \middle| f \in \Omega(\phi,\varphi) \text{ is smooth} \right\}.$$
(1.27)

Eliashberg and Polterovich [9, 27] showed for each $p \in [1, \infty)$ that the pseudo-distance $d_{(0,p)}$ is degenerate, and vanishes if M is closed. We have the following extension.

Theorem 1.12 For each $k \in \mathbb{N} \cup \{0\}$ and each $0 , the pseudo quasimetric <math>d_{(k,p)}$ is degenerate, and vanishes if M is closed.

Similarly, for a closed embedded Lagrangian submanifold L of (M, ω) let $\mathcal{L}(M, \omega, L)$ denote the space of Lagrangian submanifolds of (M, ω) which is Hamiltonian isotopic to L. For each $k = 0, 1, 2, \cdots$, define $\delta_k : \mathcal{L}(M, \omega, L) \times \mathcal{L}(M, \omega, L) \to \mathbb{R} \cup \{+\infty\}$ by

$$\delta_k(L_1, L_2) = \inf\{ \|\phi\|_k \, | \, \phi \in \operatorname{Ham}(M, \omega) \& \phi(L_1) = L_2 \}.$$

Then $\delta_k(L_1, L_2) = \delta_k(L_2, L_1)$ and $\delta_k(L_1, L_2) \leq 2^k (\delta_k(L_1, L_3) + \delta_k(L_2, L_3))$ for any $L_i \in \mathcal{L}, i = 1, 2, 3$. If (M, ω) is a tame symplectic manifold, Chekanov showed in [8] that $\delta_0 = \delta_H$ is non-degenerate, and so each δ_k is a Ham (M, ω) -invariant quasimetric.

The paper is organized as follows. In Section 2 we give proofs of Theorem 1.4, 1.10, 1.11, 1.12. Extensions of our metrics onto the group of symplectic diffeomorphisms will be discussed in Section 3. Finally, Section 4 outlines our constructions on the group of strictly contact diffeomorphisms as a concluding remark.

Acknowledgements. We would like to thank Professors L. Polterovich, H.Hofer and J. C. Sikorav for kind helps in understanding their papers.

2 Proofs

2.1 Proof of Theorem 1.4

In the following we always assume k > 0.

(i) For $\Omega(\phi, \varphi) \ni f : [0,1] \to \operatorname{Ham}(M, \omega)$, let $\overline{f} \in \Omega(\varphi, \phi)$ be defined by $\overline{f}(t) =$

 $f(1-t), \forall t \in [0,1]$. Then $\Omega(\phi, \varphi) \ni f \to \overline{f} \in \Omega(\varphi, \phi)$ is a bijection. By the assumption there exists a division $0 = t_0 < t_1 < \cdots < t_n = 1, n \in \mathbb{N}$, such that $f_i = f|_{[t_{i-1},t_i]}$ is smooth for $i = 1, \cdots, n$. By Lemma 1.1, for each $j = 1, \cdots, n$ there exists a (time-dependent) normalized Hamiltonian function $F_j : M \times [t_{j-1}, t_j] \to \mathbb{R}$ such that

$$\frac{d}{dt}f_j(t)(x) = X_{F_j}(f_j(t)(x), t)$$
(2.1)

for all $x \in M$ and $t \in [t_{j-1}, t_j]$. Set $s_i = 1 - t_{n-i}$, $i = 0, \dots, n$. Then $0 = s_0 < s_1 < \dots < s_n = 1$ is a division of [0, 1], and for any $t \in [s_{j-1}, s_j]$ it holds that

$$\bar{f}_j(t) = \bar{f}|_{[s_{j-1},s_j]}(t) = f(1-t) = f|_{[t_{n-j},t_{n-j+1}]}(1-t) = f_{n-j+1}(1-t).$$

From this and (2.1) it follows that

$$\frac{d}{dt}\bar{f}_{j}(t)(x) = -\frac{d}{ds}f_{n-j+1}(s)(x)|_{s=1-t}
= -X_{F_{n-j+1}}(f_{n-j+1}(s)(x),s)|_{s=1-t} = X_{G_{j}}(\bar{f}_{j}(t)(x),t),$$

where $G_j: M \times [s_{j-1}, s_j] \to \mathbb{R}$ is given by $G_j(x, t) = -F_{n-j+1}(x, 1-t)$. Hence

$$\frac{\partial^{i}(G_{j}(x,s))}{\partial s^{i}} = (-1)^{i+1} \times (F_{n-j+1})_{2}^{(i)}(x,1-s) \quad \forall s \in [s_{j-1},s_{j}]$$

for $i = 0, \dots, k$, where $(F_{n-j+1})_2^{(i)}$ stands for the *i*th partial derivative of F_{n-j+1} with respect to the second variable. By the definition we have

$$\begin{aligned} \operatorname{Length}_{k}(\bar{f}) &= \sum_{j=1}^{n} \operatorname{Length}_{k}(\bar{f}_{j}) \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{s_{j-1}}^{s_{j}} \left\| \frac{\partial^{i}(G_{j}(x,s))}{\partial s^{i}} \right\|_{\infty} ds \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{s_{j-1}}^{s_{j}} \left\| (-1)^{i+1}(F_{n-j+1})_{2}^{(i)}(x,1-s) \right\|_{\infty} ds \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{s_{j-1}}^{s_{j}} \left\| (F_{n-j+1})_{2}^{(i)}(x,1-s) \right\|_{\infty} ds \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{t_{n-j}}^{t_{n-j+1}} \left\| (F_{n-j+1})_{2}^{(i)}(x,t) \right\|_{\infty} dt \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{t_{j-1}}^{t_{j}} \left\| (F_{j})_{2}^{(i)}(x,t) \right\|_{\infty} dt \\ &= \sum_{j=1}^{n} \operatorname{Length}_{k}(f_{j}) = \operatorname{Length}_{k}(f). \end{aligned}$$

Thus $d_k(\phi, \varphi) = d_k(\varphi, \phi)$.

(ii) Let $\Omega(\phi, \theta) \ni f : [0, 1] \to \operatorname{Ham}(M, \omega), \ \Omega(\theta, \varphi) \ni g : [0, 1] \to \operatorname{Ham}(M, \omega)$. We define the product path $g \sharp f : [0, 1] \to \operatorname{Ham}(M, \omega)$ of f and g by

$$g \sharp f(t) = \begin{cases} f(2t) & 0 \le t \le 1/2 \\ g(2t-1) & 1/2 \le t \le 1, \end{cases}$$

then $g \sharp f$ is a piecewise smooth Hamiltonian path connecting ϕ and φ , i.e. $g \sharp f \in \Omega(\phi, \varphi)$.

By assumption there exist divisions

$$0 < t_0 < t_1 < \dots < t_n = 1$$
 and $0 < t'_0 < t'_1 < \dots < t'_m = 1$,

such that for $i = 1, \dots, n$, $f_i = f|_{[t_{i-1}, t_i]}$ is smooth; for $j = 1, \dots, m$, $g_j = g|_{[t'_{j-1}, t'_j]}$ is also smooth. Denote the Hamiltonian functions generating $\{f_i\}_{i=1}^n$, $\{g_j\}_{j=1}^m$ by $\{F_i\}_{i=1}^n$ and $\{G_j\}_{j=1}^m$ respectively. For $0 \le i \le n$, set $s_i = \frac{t_i}{2}$, for $n+1 \le i \le n+m$, set $s_i = \frac{t'_{i-n}+1}{2}$, then

$$0 = s_0 < s_1 < \dots < s_n = \frac{1}{2} < s_{n+1} < \dots < s_{n+m} = 1$$

is a division of [0, 1], and at this time $(g \sharp f)_i = g \sharp f|_{[s_{i-1}, s_i]}$ is smooth for $i = 1, \dots, n + m$. Denote the Hamiltonian function generating $(g \sharp f)_i$ by $H_i : M \times [s_{i-1}, s_i] \to \mathbb{R}$, then

$$H_i(x,s) = 2F_i(x,2s), \quad 1 \le i \le n,$$

$$H_i(x,s) = 2G_{i-n}(x,2s-1), \quad n+1 \le i \le n+m.$$

By definition we have

$$d_k(\phi,\varphi) \leq \operatorname{Length}_k(g\sharp f) = \sum_{j=1}^{n+m} \operatorname{Length}_k\left((g\sharp f)_j\right)$$
$$= \sum_{j=1}^{n+m} \sum_{i=0}^k \int_{s_{j-1}}^{s_j} \left\| \frac{\partial^i(H_j(x,s))}{\partial s^i} \right\|_{\infty} ds$$
$$= \sum_{j=1}^n \sum_{i=0}^k \int_{s_{j-1}}^{s_j} \left\| \frac{\partial^i(2F_j(x,2s))}{\partial s^i} \right\|_{\infty} ds$$
$$+ \sum_{j=n+1}^{n+m} \sum_{i=0}^k \int_{s_{j-1}}^{s_j} \left\| \frac{\partial^i(2G_{j-n}(x,2s-1))}{\partial s^i} \right\|_{\infty} ds$$

$$\begin{split} &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{\frac{t_{j-1}}{2}}^{\frac{t_{j}}{2}} 2^{i+1} \left\| (F_{j})_{2}^{(i)}(x,2s) \right\|_{\infty} ds \\ &+ \sum_{j=n+1}^{n+m} \sum_{i=0}^{k} \int_{\frac{t'_{j-1}-n+1}{2}}^{\frac{t'_{j-n}+1}{2}} 2^{i+1} \left\| (G_{j-n})_{2}^{(i)}(x,2s-1) \right\|_{\infty} ds \\ &= \sum_{j=1}^{n} \sum_{i=0}^{k} \int_{t_{j-1}}^{t_{j}} 2^{i} \left\| (F_{j})_{2}^{(i)}(x,s) \right\|_{\infty} ds + \sum_{j=1}^{m} \sum_{i=0}^{k} \int_{t'_{j-1}}^{t'_{j}} 2^{i} \left\| (G_{j})_{2}^{(i)}(x,s) \right\|_{\infty} ds \\ &\leq 2^{k} \left(\sum_{j=1}^{n} \sum_{i=0}^{k} \int_{t_{j-1}}^{t_{j}} \left\| \frac{\partial^{i}(F_{j}(x,s))}{\partial s^{i}} \right\|_{\infty} ds + \sum_{j=1}^{m} \sum_{i=0}^{k} \int_{t'_{j-1}}^{t'_{j}} \left\| \frac{\partial^{i}(G_{j}(x,s))}{\partial s^{i}} \right\|_{\infty} ds \right) \\ &= 2^{k} \left(\sum_{j=1}^{n} \operatorname{Length}_{k}(f_{j}) + \sum_{j=1}^{m} \operatorname{Length}_{k}(g_{j}) \right) \end{split}$$

Take the infimum for all $f \in \Omega(\phi, \theta)$ and $g \in \Omega(\theta, \varphi)$ at the right hand of the above equation respectively, we obtain the desired triangle inequality $d_k(\phi, \varphi) \leq 2^k (d_k(\phi, \theta) + d_k(\theta, \varphi)).$

(iii) Because $d_k \ge d_0 = d_H$, the non-degeneracy of d_k is obvious.

(iv) Firstly, we prove the right-invariance of d_k . Let $f \in \Omega(\phi, \varphi)$ be generated by $\{F_i, 1 \leq i \leq n\}$ as above. Then $(f \circ \theta)(t) := f(t)\theta$ defines an element $f \circ \theta$ in $\Omega(\phi\theta, \varphi\theta)$, and $f \circ \theta|_{[t_{i-1}, t_i]}$ also correspond to (time-dependent) Hamiltonian functions $F_i, i = 1, \dots, n$. So $\text{Length}_k(f) = \text{Length}_k(f \circ \theta)$. Hence,

$$\begin{aligned} d_k(\phi,\varphi) &= \inf_f \{ \operatorname{Length}_k(f) \} = \inf_f \{ \operatorname{Length}_k(f \circ \theta) \} \\ &= \inf_{g \in \Omega(\phi\theta,\varphi\theta)} \{ \operatorname{Length}_k(g) \} = d_k(\phi\theta,\varphi\theta). \end{aligned}$$

Next we prove the left-invariance of d_k . Let f as above. Then $\theta \circ f(t) := \theta(f_t(x))$ defines an element $\theta \circ f$ in $\Omega(\theta\phi, \theta\varphi)$, and $\theta \circ f|_{[t_{i-1},t_i]}$ correspond to Hamiltonian functions $F_i(\theta^{-1}(\cdot), t), i = 1, \cdots, n$. So $\text{Length}_k(f) = \text{Length}_k(\theta \circ f)$. By the same argument as above, we get $d_k(\phi, \varphi) = d_k(\theta\phi, \theta\varphi)$.

2.2 Proof of Theorems 1.10, 1.11

Following [30, 13] we first prove

Lemma 2.1 Assume $\psi_1, \psi_2, \cdots, \psi_m \in \text{Ham}(\mathbb{R}^{2n}, \omega_0)$ have properly separated supports. Then

$$\|\psi_1\psi_2\cdots\psi_m\|_k^* \le 2(k+1)\max_j \|\psi_j\|_k^*.$$

Proof. Given $\varepsilon > 0$, by (1.23) we have $f_j \in \Omega(\mathrm{id}, \psi_j)$ such that $\mathrm{Length}_k^*(f_j) < \|\psi_j\|_k^* + \varepsilon$, $j = 1, \cdots, m$. Let H_j be the corresponding normalized Hamiltonian functions of $f_j, j = 1, \cdots, m$. Then $\||H_j\||_k = \mathrm{Length}_k^*(f_j) \leq \|\psi_j\|_k^* + \varepsilon$. Through a

refinement, we could suppose that there exists a common division of time $0 = t_0 < t_1 < \cdots < t_n = 1$ such that $H_j^i := H_j|_{[t_{i-1},t_i]}$ is smooth for every $1 \le j \le m, 1 \le i \le n$.

Using the same notations as those of [14], we set $S_j = \operatorname{supp}(\psi_j)$, choose R > 0such that $\operatorname{supp}(H_j) \subset B_R(x_j^*) \times [0, 1]$ with $x_j^* \in S_j$, and then vectors v_j such that the sets $B_R(S_j + v_j)$ are disjoint. Let $\tau \in \operatorname{Ham}(M, \omega)$ be the map associated to the S_j guaranteed by Lemma 8 on the page 175 of [14], and let $\hat{\psi}_j = \tau \psi_j \tau^{-1}$ and

$$\hat{f}_j(t) = \tau f_j(t) \tau^{-1}, \ j = 1, \cdots, m.$$

Then the corresponding normalized Hamiltonian function with \hat{f}_j is given by

$$\hat{H}_j(x,t) = H_j(\tau^{-1}(x),t) = H_j(x-v_j,t),$$

and thus the corresponding normalized Hamiltonian function with

$$\hat{f}_1 \hat{f}_2 \cdots \hat{f}_m \in \Omega(\mathrm{id}, \hat{\psi}_1 \hat{\psi}_2 \cdots \hat{\psi}_m)$$

is $\hat{H} = \hat{H}_1 + \dots + \hat{H}_m$. It follows that

$$\|\psi_1\psi_2\cdots\psi_m\|_k^* = \|\hat{\psi}_1\hat{\psi}_2\cdots\hat{\psi}_m\|_k^* \le |||\hat{H}|||_k.$$

By the definition in (1.22) we have $s_i \in [0, 1], i = 0, \dots, k$, such that

$$\begin{split} |||\hat{H}|||_{k} &= \sum_{i=0}^{k} \max_{1 \leq l \leq n} \max_{t_{l-1} \leq t \leq t_{l}} \left(\sup_{x} \frac{\partial^{i} \hat{H}^{l}}{\partial t^{i}} - \inf_{x} \frac{\partial^{i} \hat{H}^{l}}{\partial t^{i}} \right) \\ &= \sum_{i=0}^{k} \left(\sup_{x} \frac{\partial^{i} \hat{H}}{\partial t^{i}}(x, s_{i}) - \inf_{x} \frac{\partial^{i} \hat{H}}{\partial t^{i}}(x, s_{i}) \right) \\ &= \sum_{i=0}^{k} \left[\max_{1 \leq j \leq m} \left(\sup_{x} \frac{\partial^{i} \hat{H}_{j}}{\partial t^{i}}(x, s_{i}) \right) - \min_{1 \leq j \leq m} \left(\inf_{x} \frac{\partial^{i} \hat{H}_{j}}{\partial t^{i}}(x, s_{i}) \right) \right] \\ &\leq \sum_{i=0}^{k} 2 \max_{1 \leq j \leq m} \left(\left\| \frac{\partial^{i} \hat{H}_{j}}{\partial t^{i}}(\cdot, s_{i}) \right\|_{\infty} \right) \\ &\leq \sum_{i=0}^{k} 2 \max_{1 \leq j \leq m} \left(\max_{t \in [0,1]} \left\| \frac{\partial^{i} \hat{H}_{j}}{\partial t^{i}}(\cdot, t) \right\|_{\infty} \right) \\ &\leq 2(k+1) \max_{1 \leq j \leq m} \left(\sum_{i=0}^{k} \max_{t \in [0,1]} \left\| \frac{\partial^{i} \hat{H}_{j}}{\partial t^{i}}(\cdot, t) \right\|_{\infty} \right) \\ &= 2(k+1) \max_{1 \leq j \leq m} ||\hat{H}_{j}|||_{k} \\ &\leq 2(k+1) \max_{j} \|\psi_{j}\|_{k}^{*} + 2(k+1)\varepsilon. \end{split}$$

This holds for every $\varepsilon > 0$ and the lemma is proved.

Proof of Theorem 1.11. Given $\varepsilon > 0$, as in the proof of [30, 13] we can construct maps $\psi_j, 0 \le j \le N$ satisfying $d_k^*(\psi_j, \psi_{j+1}) < \varepsilon$, and maps $\varphi_j(0 \le j \le 2N)$, $\alpha_j(1 \le 2N)$

 $j \leq N$, $\beta_j (0 \leq j \leq N)$ with $\|\varphi_0\|_k^* \leq e_{p,k}^*(U) + \varepsilon$. Then we have

$$\begin{aligned} \|\varphi_H\|_k^* &= \|\beta_N\|_k^* = \left\| (\prod_{j=1}^N \alpha_j \beta_j) (\prod_{j=0}^{N-1} \alpha_{j+1} \beta_j)^{-1} \right\|_k^* \\ &\leq 2^{k+2} (k+1) \left(\max_{1 \le j \le N} \|\alpha_j \beta_j\|_k^* + \max_{0 \le j \le N-1} \|\alpha_{j+1} \beta_j\|_k^* \right) \end{aligned}$$

(because $\|\phi\psi\|_k^* \leq 2^{k+1}(\|\phi\|_k^* + \|\psi\|_k^*)$ by Theorem 1.9). We can estimate

$$\begin{aligned} \|\alpha_{j}\beta_{j}\|_{k}^{*} &\leq 2^{k+1} \left[\|\varphi_{2j-1}\|_{k}^{*} + 2^{k+1} (\|\varphi_{2j-1}^{-1}\varphi_{2j}\|_{k}^{*} + \|\varphi_{2j}\|_{k}^{*}) \right] \\ &\leq 2^{k+1} \|\varphi_{0}\|_{k}^{*} + 2^{3k+3} (\|\varphi_{2j-1}\|_{k}^{*} + \|\varphi_{2j}\|_{k}^{*}) + 2^{2k+2} \|\varphi_{0}\|_{k}^{*} \\ &= (2^{k+1} + 2^{2k+2} + 2^{3k+4}) \|\varphi_{0}\|_{k}^{*}. \end{aligned}$$

Similarly we have

$$\begin{aligned} \|\alpha_{j+1}\beta_{j}\|_{k}^{*} &\leq 2^{k+1} \left[\|\varphi_{2j+1}\|_{k}^{*} + 2^{k+1} (\|\psi_{j+1}\psi_{j}^{-1}\psi_{j}\varphi_{2j+1}^{-1}\varphi_{2j}\psi_{j}^{-1}\|_{k}^{*} + \|\varphi_{2j}\|_{k}^{*}) \right] \\ &\leq 2^{k+1} \|\varphi_{0}\|_{k}^{*} + 2^{3k+3} (\|\psi_{j+1}\psi_{j}^{-1}\|_{k}^{*} + \|\varphi_{2j+1}^{-1}\varphi_{2j}\|_{k}^{*}) + 2^{2k+2} \|\varphi_{0}\|_{k}^{*} \\ &\leq (2^{k+1} + 2^{2k+2} + 2^{4k+5}) \|\varphi_{0}\|_{k}^{*} + 2^{3k+3} d_{k}^{*}(\psi_{j+1}, \psi_{j}). \end{aligned}$$

Summing up, we have

$$\begin{aligned} \|\varphi_H\|_k^* &\leq 2^{k+2}(k+1)(2^{k+2}+2^{2k+3}+2^{3k+4}+2^{4k+5})\|\varphi_0\|_k^* + 2^{4k+5}(k+1)\varepsilon \\ &\leq 2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})e_{p,k}^*(U) + \\ &\quad 2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})\varepsilon + 2^{4k+5}(k+1)\varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, the proof is finished.

Carefully checking the proof of Proposition 6 in [13] (or [14, Lemma 10]) and replacing E and e_p therein we still have for our $\|\cdot\|_k^*$ and $e_{p,k}^*$:

Lemma 2.2 Let $\psi \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0)$ with $\psi \neq \operatorname{id}$ and let $\delta > |\psi - \operatorname{id}|_{C^0}$. For every $Q \subset \mathbb{R}^{2n}$ open and satisfying $Q \cap \operatorname{supp}(\psi) \neq \emptyset$, there exists a $\varphi \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0)$ satisfying

- (i) $\varphi | Q = \psi | Q$
- (ii) $\operatorname{supp}(\psi) \subset U$

(iii) $\|\varphi\|_k^* \le 2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})e_{p,k}^*(U),$

where U is the intersection of $B_{\delta}(Q)$ with the convex hull of $\operatorname{supp}(\psi)$, and $B_{\delta}(Q) = \{x | \operatorname{dist}(x, Q) < \delta\}.$

Similarly, corresponding to [13, Corollary 7] or [14, Lemma 11] we have

Lemma 2.3 Let $U = (a_1, a_2) \times (b_1, b_2) \oplus \mathbb{R}^{2n-2}$. Then for $\psi \in \text{Ham}(\mathbb{R}^{2n}, \omega_0)$ with $\text{supp}(\psi) \subset U$ it holds that

$$\|\varphi\|_k^* \le 2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})(a_2-a_1)(b_2-b_1).$$

Proof of Theorem 1.10. Let $\{\varphi_j\}_{j \in I}$ be as in the proof of [14, Theorem 9]. Corresponding to the inequality (iii) on the page 180 of [14], we have

$$\|\varphi_j\|_k^* \le 2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})e_{p,k}^*(U_j).$$

And similar to [14, (5.32)] we have estimate

$$\begin{split} \|\prod_{j\in I}\varphi_j\|_k^* &\leq 2(k+1)\max_{j\in I} \|\varphi_j\|_k^* \\ &\leq 2(k+1)2^{2k+4}(k+1)(1+2^{k+1}+2^{2k+2}+2^{3k+3})\cdot 2(\varepsilon+\delta_1)R \\ &\leq 2^{2k+6}(k+1)^2(1+2^{k+1}+2^{2k+2}+2^{3k+3})R\delta. \end{split}$$

Thus

$$\begin{aligned} \|\psi\|_{k}^{*} &\leq 2^{k+1}(\|\psi\theta\|_{k}^{*} + \|\theta\|_{k}^{*}) \\ &= 2^{k+1}(\|\prod \hat{\varphi}_{j}\|_{k}^{*} + \|\prod \varphi_{j}\|_{k}^{*}) \\ &\leq 2^{3k+8}(k+1)^{2}(1+2^{k+1}+2^{2k+2}+2^{3k+3})R\delta. \end{aligned}$$

This holds true for every $\delta > |\psi - \mathrm{id}|_{C^0}$, and we conclude that

$$\|\psi\|_k^* \le 2^{3k+8}(k+1)^2(1+2^{k+1}+2^{2k+2}+2^{3k+3})R|\psi-\mathrm{id}|_{C^0},$$

where $R = \text{diameter}(\text{supp}(\psi))$. Finally we have the inequality:

$$d_k^*(\varphi,\psi) \le 2^{3k+8}(k+1)^2(1+2^{k+1}+2^{2k+2}+2^{3k+3}) \operatorname{diameter}\left(\operatorname{supp}(\varphi\psi^{-1})\right)|\varphi-\psi|_{C^0},$$

2.3 Proof of Theorem 1.12

When k = 0 and $p \in [1, \infty)$ this is the result in [9]. We shall assume that either k = 0 and $0 or <math>k \in \mathbb{N}$ and 0 below.

The proof ideas are same as those of [9]. Given a bi-invariant pseudo quasimetric ρ on Ham (M, ω) , the $(\rho$ -)displacement energy of a subset $A \subset M$ is defined by

$$e_{\rho}(A) = \inf\{\rho(\mathrm{id}, g) \mid g \in \mathrm{Ham}(M, \omega) \text{ such that } g(A) \cap A = \emptyset\}.$$
 (2.2)

As in the proof of [9, Th.2.2.A] we may obtain:

Claim 2.4 Every nonempty open subset of M has positive displacement energy with respect to a bi-invariant quasimetric on $\operatorname{Ham}(M, \omega)$.

So it suffices to prove that the displacement energy associated with $d_{(k,p)}$ (0 vanishes for some embedded open ball in <math>M. In fact, using Darboux theorem, we can choose a chart $M \supset U \ni w \rightarrow (x_1(w), \cdots, x_n(w), y_1(w), \cdots, y_n(w)) \in \mathbb{R}^{2n}$

so that the symplectic form $\omega = \sum_{i=1}^{n} dx_i \wedge dy_i$ on it. Replacing ω by $N\omega$ for some large N > 0 we may assume

$$U \supset B(0,4) = \left\{ (x,y) \in \mathbb{R}^{2n} \mid \sum_{i=1}^{n} (x_i^2 + y_i^2) < 16 \right\}, \quad A = B(0,1).$$

Consider a Hamiltonian isotopy $\{h_t\}, t \in [0, 1]$ such that when restricted to U, h_t is simply a shift by 2t along the y_1 coordinate. Assume that $\{h_t\}$ is generated by H, then $H(x, y, t) = 2x_1$ on U (H is not normalized). Clearly $h_1(A) \cap A = \emptyset$.

Fixed a smooth cut-off function $\delta : \mathbb{R} \to [0, 1]$, such that $\delta \equiv 1$ for $|t| \leq 1/4$, $\delta \equiv 0$ for $|t| \geq 3/4$. Set $\delta_m(t) = \delta(mt)$. When M is open, we can define a sequence of functions $\{G_m\}_{m \in \mathbb{N}}$ as follows:

$$G_m(x,y,t) = \begin{cases} 2x_1 \delta_m(\sqrt{|x|^2 + |y - (2t, 0, \cdots, 0)|^2} - 1) & (x,y) \in U\\ 0 & (x,y) \notin U. \end{cases}$$
(2.3)

 G_m is smooth on M. The support of G_m is contained in a neighborhood of $h_t(\partial A)$ for each $t \in [0,1]$ and tends to $h_t(\partial A)$ as $m \to \infty$. Since for every t the function $G_m(\cdot, t)$ coincides with H near $h_t(\partial A)$, we conclude that the Hamiltonian vector field of G_m is equal to X_H near $h_t(\partial A)$ for every m. Hence the Hamilton isotopy $\{\phi_{G_m}^t\}$ of G_m satisfies $\phi_{G_m}^t(\partial A) = h_t(\partial A)$ and so $\phi_{G_m}(\partial A) \cap \partial A = \emptyset$. But this obviously implies that $\phi_{G_m}(A) \cap A = \emptyset$.

Clearly, for $0 we have <math>\int_0^1 ||G_m(\cdot, t)||_p dt \to 0$ as $m \to \infty$. Next we prove that for each $1 \le i \le k$,

$$\left\| \frac{\partial^i G_m}{\partial t^i}(\cdot, t) \right\|_p^p \to 0 \quad \text{uniformly in } t \in [0, 1] \text{ as } m \to \infty$$

For simplicity we write $r = \sqrt{(|x|^2 + |y - (2t, 0, \dots, 0)|^2)}$. In U we have

$$\frac{\partial^{i} G_{m}}{\partial t^{i}}(x, y, t) = m^{i} x_{1} \delta_{m}^{(i)}(r-1) \cdot 2^{i+1} \left(\frac{t-y_{1}}{r}\right)^{i} + m^{i-1} L(x, y, t).$$

Here L is a combination of $\delta_m^{(j)}$, $1 \leq j \leq i-1$. Its coefficients are rational functions of x, y, t which are bounded in C^0 -norm near $h_t(\partial A)$. Hence,

$$\begin{aligned} \left\| \frac{\partial^{i} G_{m}}{\partial t^{i}}(\cdot,t) \right\|_{p}^{p} &= \int_{U} m^{ip} \left| x_{1} \delta_{m}^{(i)}(r-1) \cdot 2^{i+1} \left(\frac{t-y_{1}}{r} \right)^{i} + \frac{1}{m} L(x,y,t) \right|^{p} \omega^{n} \\ &\leq C m^{ip} \int_{\Sigma_{1} \cup \Sigma_{2}} \omega^{n} = C m^{ip} \big(\operatorname{Vol}(\Sigma_{1}) + \operatorname{Vol}(\Sigma_{2}) \big), \end{aligned}$$

where C > 0 is a constant depending on δ , Σ_1 is a region bounded by two spheres whose radiuses are $1 + \frac{1}{4m}$, $1 + \frac{3}{4m}$ respectively, and Σ_2 is a similar region with radiuses $1 - \frac{1}{4m}, 1 - \frac{3}{4m}$ respectively. So we have

$$\begin{aligned} \left\| \frac{\partial^{i} G_{m}}{\partial t^{i}}(\cdot,t) \right\|_{p}^{p} \\ &\leq \frac{\pi^{n}}{n!} C m^{ip} \left[\left(1 + \frac{3}{4m} \right)^{2n} - \left(1 + \frac{1}{4m} \right)^{2n} + \left(1 - \frac{1}{4m} \right)^{2n} - \left(1 - \frac{3}{4m} \right)^{2n} \right] \\ &= \frac{\pi^{n}}{n!} C m^{ip} \left[\frac{2}{m} + O\left(\frac{1}{m^{2}} \right) \right]. \end{aligned}$$

Since $ip \le kp < 1$, so the above expression tends to zero when $m \to \infty$.

When M is closed, we could find an open set V disjoint with the above U, and shrinking U properly we can also assume that V is symplectomorphic to U. Then we can define a function \widetilde{G}_m on V which is the same form as G_m in (2.3) but with a minus in front additionally. Define K_m to be the sum of G_m and \widetilde{G}_m . Then K_m is a normalized Hamiltonian function such that $\phi_{K_m}(A) \cap A = \emptyset$. Using the same estimates as above for G_m and \widetilde{G}_m respectively, we derive

$$\left\|\frac{\partial^i K_m}{\partial t^i}(\cdot, t)\right\|_p \to 0$$
 uniformly in $t \in [0, 1]$ as $m \to \infty$.

All these lead to the first conclusion.

Note that $\{\phi \in \operatorname{Ham}(M, \omega) | \rho(\operatorname{id}_M, \phi) = 0\}$ is also a normal subgroup of $\operatorname{Ham}(M, \omega)$ for any bi-invariant pseudo quasimetric ρ on $\operatorname{Ham}(M, \omega)$. The second claim follows from Banyaga's theorem as in [27].

3 Extensions

It is a natural question to extend bi-invariant pseudo-metric on $\operatorname{Ham}(M, \omega)$ to the group of symplectic diffeomorphisms. There exist different ways realizing this. We only use the method by Lalonde and Polterovich [16], and one by Banyaga [2].

3.1 The method by Lalonde and Polterovich

Each $\phi \in \text{Symp}(M, \omega)$ induces an isometry with respect to the Hofer metric,

$$C_{\phi} : \operatorname{Ham}(M, \omega) \to \operatorname{Ham}(M, \omega), \ f \mapsto C_{\phi} f := \phi f \phi^{-1}.$$

For $\alpha \in (0, \infty]$ it was shown in [16, Proposition 1.2.A] that

$$r_{\alpha}(\phi) := \sup\{d_H(f, C_{\phi}f) \mid f \in \operatorname{Ham}(M, \omega) \& \|f\|_H \le \alpha\}$$

$$(3.1)$$

defines a bi-invariant function r_{α} and $r_{\alpha}(\phi) \leq 2\alpha \ \forall \phi \in \text{Symp}(M, \omega)$. In particular, r_{α} is an bi-invariant norm on $\text{Symp}(M, \omega)$ if $\alpha \in (0, \infty)$. We say $\phi \in \text{Symp}(M, \omega)$ to be **bounded** if the function $(0, \infty) \ni \alpha \mapsto r_{\alpha}(\phi)$ is bounded, or equivalently C_{ϕ} is C^{0} -bounded, i.e.,

$$r_{\infty}(\phi) = \sup\{d_H(f, C_{\phi}f) \mid f \in \operatorname{Ham}(M, \omega)\} < \infty.$$
(3.2)

By the bi-invariance of the Hofer metric d_H the set of all bounded symplectomorphism in $\operatorname{Symp}(M, \omega)$ form a normal subgroup of $\operatorname{Symp}(M, \omega)$, denoted by $\operatorname{BI}(M)$. Set $\operatorname{BI}_0(M) = \operatorname{BI}(M) \cap \operatorname{Symp}_0(M)$, where $\operatorname{Symp}_0(M)$ is the connected component of $\operatorname{Symp}(M, \omega)$ containing the identity map. Every $\phi \in \operatorname{Ham}(M, \omega)$ is bounded and $r_{\infty}(\phi) \leq 2 \|\phi\|_H$ because

$$d_H(f, C_{\phi}f) = d_H(\operatorname{id}, [\phi, f]) \le 2 \|\phi\|_H, \quad \forall f \in \operatorname{Ham}(M, \omega).$$
(3.3)

Hence $\operatorname{Ham}(M, \omega)$ is a subgroup of $\operatorname{BI}_0(M)$. These motivated Lalonde and Polterovich to propose the so-called **bounded** isometry conjecture.

Conjecture 1 ([16, Conjecture 1.3.A]) $BI_0(M) = Ham(M, \omega)$.

They proved it in [16] for some cases, for example, M is any closed surface with area form or M is a product of closed surfaces of genus greater than 0 with product symplectic form. For recent progresses the reader may refer to [15, 11, 7, 25].

Now let us consider corresponding questions with metric d_k . For each $\phi \in$ Symp (M, ω) it is easy to check that C_{ϕ} is still an isometry of Ham (M, ω) with respect to the quasi-norm $\|\cdot\|_k = d_k(\mathrm{id}, \cdot)$. Corresponding to (3.1) we define

$$r_{\alpha,k}(\phi) := \sup\{d_k(f, C_{\phi}f) \mid f \in \operatorname{Ham}(M, \omega) \& \|f\|_k \le \alpha\}$$
(3.4)

for each $\alpha \in (0, \infty]$. It is also bi-invariant and satisfies

$$r_{\alpha,k}(\phi) \le 2^{k+1}\alpha, \ \forall \phi \in \operatorname{Symp}(M,\omega).$$
 (3.5)

Actually, the function $r_{\alpha,k}$ is a quasi-norm on $\text{Symp}(M,\omega)$ by the following:

Lemma 3.1 For every $k \ge 0$, $\alpha \in (0, \infty]$, the function $r_{\alpha,k}$ on $\text{Symp}(M, \omega)$ defined above is conjugate invariant, assumes the value 0 only at the identity, and satisfies the quasi-triangle inequality

$$r_{\alpha,k}(\phi\psi) \le 2^k (r_{\alpha,k}(\phi) + r_{\alpha,k}(\psi)).$$

Proof. (i) The conjugate invariance of $r_{\alpha,k}$. For $\phi, \varphi \in \text{Symp}(M, \omega)$ we have

$$\begin{aligned} r_{\alpha,k}(\varphi\phi\varphi^{-1}) &= \sup_{\{f|\|f\|_k \le \alpha\}} d_k(f,\varphi\phi\varphi^{-1}f\varphi\phi^{-1}\varphi^{-1}) \\ &= \sup_{\{f|\|f\|_k \le \alpha\}} d_k(\varphi^{-1}f\varphi,\phi\varphi^{-1}f\varphi\phi^{-1}) \\ &= \sup_{\{g|\|g\|_k \le \alpha\}} d_k(g,\phi g\phi^{-1}) \quad (\text{setting } g = \varphi^{-1}f\varphi) \\ &= r_{\alpha,k}(\phi). \end{aligned}$$

(ii) The non-degeneracy of $r_{\alpha,k}$. If $r_{\alpha,k}(\phi) = 0$, then $d_k(f, \phi f \phi^{-1}) = 0$ for each $f \in \overline{\operatorname{Ham}(M,\omega)}, \|f\|_k \leq \alpha$. By the non-degeneracy of d_k , we have $f = \phi f \phi^{-1}$. Suppose that f is generated by Hamiltonian function F. Then $\phi f \phi^{-1}$ is generated by $F \circ \phi^{-1}$. Hence $F = F \circ \phi^{-1}$. By the arbitrariness of F, we have $\phi = \operatorname{id}$. (iii) The quasi-triangle inequality holds for $r_{\alpha,k}$.

$$\begin{aligned} r_{\alpha,k}(\phi\psi) &= \sup_{\{f|\|f\|_{k} \leq \alpha\}} d_{k}(f,\phi\psi f\psi^{-1}\phi^{-1}) \\ &\leq 2^{k} \left(\sup_{\{f|\|f\|_{k} \leq \alpha\}} d_{k}(f,\phi f\phi^{-1}) + \sup_{\{f|\|f\|_{k} \leq \alpha\}} d_{k}(\phi f\phi^{-1},\phi\psi f\psi^{-1}\phi^{-1}) \right) \\ &= 2^{k} \left(r_{\alpha,k}(\phi) + r_{\alpha,k}(\psi) \right). \end{aligned}$$

If $r_{\infty,k}(\phi) < \infty$ we call ϕ a **k-bounded isometry**. Clearly, every k-bounded symplectomorphism is bounded. As in (3.3) for $\phi \in \text{Ham}(M, \omega)$ we have

$$d_k(f, C_{\phi}f) = d_k(\operatorname{id}, [\phi, f]) \le 2^{k+1} \|\phi\|_k, \quad \forall f \in \operatorname{Ham}(M, \omega)$$
(3.6)

and thus $r_{\infty,k}(\phi) \leq 2^{k+1} \|\phi\|_k$.

Let

$$BI_k(M) = \{ \phi \in Symp(M, \omega) \, | \, r_{\infty, k}(\phi) < \infty \} \text{ and} \\ BI_{k, 0}(M) = BI_k(M) \cap Symp_0(M),$$

which correspond to the sets BI(M) and $BI_0(M)$ respectively. It is obvious that

$$\operatorname{BI}(M) \supseteq \operatorname{BI}_1(M) \supseteq \operatorname{BI}_2(M) \supseteq \cdots \supseteq \operatorname{BI}_k(M) \supseteq \cdots \supseteq \operatorname{Ham}(M,\omega)$$

because (3.6) and $\|\cdot\|_H = \|\cdot\|_0 \le \|\cdot\|_1 \le \cdots \le \|\cdot\|_k \le \cdots$. It is probably that $\operatorname{BI}_k(M) \ne \operatorname{BI}(M)$ for some $k \in \mathbb{N}$ and M. Corresponding to Conjecture 1 we have

Conjecture 2 For every symplectic manifold M and every integer $m \in \mathbb{N} \cup \{0\}$,

$$\bigcap_{k=m}^{\infty} \operatorname{BI}_{k,0}(M) = \operatorname{BI}_{m-1,0}(M).$$
(BIC)_m

In particular, for every symplectic manifold M,

$$\bigcap_{k=0}^{\infty} \operatorname{BI}_{k,0}(M) = \operatorname{Ham}(M,\omega).$$
(WBIC)

Here (WBIC) means weak bounded isometry conjecture. Clearly, the proof of (WBIC) is more hopeful than one of Conjecture 1. In particular, all manifolds mentioned above satisfy $(BIC)_m$ and (WBIC).

In the following we shall point out that many results in [16] can be generalized to the case of our quasi-metrics. Carefully checking the proof of Theorem 1.3.F in [16] we immediately obtain the following generalization of it. **Proposition 3.2** Suppose that a symplectic manifold (M, ω) satisfies $BI_{k,0}(M) = Ham(M, \omega)$ for some $k \in \mathbb{N} \cup \{0\}$. Then every symplectomorphism $\phi \in Symp(M, \omega)$ which acts nontrivially on $H^1_c(M, \mathbb{R})$ is k-unbounded.

Since a k-bounded symplectomorphism is also bounded, the following two propositions are, respectively, direct consequences of Theorems 1.4.A, 1.3.C in [16].

Proposition 3.3 If a closed Lagrangian submanifold $L \subset (M, \omega)$ admits a Riemannian metric with non-positive sectional curvature, and the inclusion $L \hookrightarrow M$ induces an injection on fundamental groups, then $\phi(L) \cap L \neq \emptyset$ for every k-bounded $\phi \in \text{Symp}(M, \omega)$.

Proposition 3.4 Let S be a closed surface of genus greater than 0 and let (W, ω_W) be closed and weakly exact (i.e., $\omega_W|_{\pi_2(W)} = 0$). Suppose that $\phi \times \psi$ be a k-bounded symplectomorphism of $(S \times W, \omega_S \oplus \omega_W)$. Then the symplectomorphism ϕ is Hamiltonian.

Finally, we give the corresponding result of Theorem 5.1.A in [16].

Theorem 3.5 For the standard symplectic space $(\mathbb{R}^{2n}, \omega_0)$, any compactly supported symplectic diffeomorphism ϕ of \mathbb{R}^{2n} is k-bounded. Precisely, we have

$$r_{\infty,k}(\phi) = \sup\{d_k(f, C_{\phi}f) \mid f \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0)\} \le 2^{3k+2}E(\operatorname{supp}(\phi)),$$

where $E(\operatorname{supp}(\phi))$ is the cylindrical capacity of $\operatorname{supp}(\phi)$. Recall that cylindrical capacity E(X) of a bounded subset $X \subset \mathbb{R}^{2n}$ is defined by

$$\inf\left\{c>0 \mid \exists \phi \in \operatorname{Ham}(\mathbb{R}^{2n},\omega_0) \text{ such that } \phi(X) \subset B^2(c) \times \mathbb{R}^{2n-2}\right\},\$$

where $B^2(c)$ stands for a disc of area c.

This result can be proved along the proof lines of [16, Theorem 5.1.A]. For the sake of completeness we give the proof of it. Firstly, we give a corresponding lemma with Lemma 5.1.B in [16].

Lemma 3.6 For all maps $\phi \in \text{Symp}(\mathbb{R}^{2n}, \omega_0)$ and $f, g \in \text{Ham}(\mathbb{R}^{2n}, \omega_0)$,

$$d_k(f, C_{\phi}f) \le 2^k d_k(f, C_{g\phi g^{-1}}f) + 2^{3k+2} \|g\|_k.$$

Proof. The quasi-triangle inequality implies that

$$d_k(f, C_{\phi}f) \le 2^k (d_k(f, C_{g\phi g^{-1}}f) + d_k(C_{\phi}f, C_{g\phi g^{-1}}f)).$$

Write $I = d_k(C_{\phi}f, C_{q\phi q^{-1}}f)$. Then

$$I = d_k(\phi f \phi^{-1}, g \phi g^{-1} f g \phi^{-1} g^{-1}) = d_k(f, \phi^{-1} g \phi g^{-1} f g \phi^{-1} g^{-1} \phi)$$

= $d_k(C_{[g,\phi^{-1}]}f, f),$

By Lemma 3.1 we have

$$I \leq r_{\infty,k}([g,\phi^{-1}]) = r_{\infty,k}(g\phi^{-1}g^{-1}\phi)$$

$$\leq 2^{k}(r_{\infty,k}(g) + r_{\infty,k}(\phi^{-1}g^{-1}\phi))$$

$$= 2^{k+1}r_{\infty,k}(g) \leq 4^{k+1}||g||_{k}$$

because of (3.6).

Lemma 3.7 Suppose that K is a compact subset in \mathbb{R}^{2n} , and that $L \subset \mathbb{R}^{2n}$ is a hyperplane so that K lies on the left of L. Let g be a compactly supported Hamiltonian diffeomorphism such that g(K) sits in the right side of L, and let L' be an arbitrary hyperplane parallel to L such that g(K) lies between L and L'. Then there exists another compactly supported Hamiltonian diffeomorphism g' such that $||g'||_k = ||g||_k$ and g'(K) lies on the right of L'.

Proof. Let $(x_1, y_1, \dots, x_n, y_n)$ denote the coordinates in \mathbb{R}^{2n} . Without loss of generality we may assume that for some v > 0 and $\varepsilon > 0$,

$$L = \{x_1 = 0\} \text{ and } L' = \{x_1 = v\},\$$

$$x_1 < -\varepsilon \quad \forall (x_1, y_1, \cdots, x_n, y_n) \in K.$$

Choose a cut off function $\eta(t) : \mathbb{R} \to [0, 1]$ satisfying

$$\eta|_{(-\infty,-\varepsilon]} = 0 \quad \text{and} \quad \eta|_{[0,+\infty)} = 1.$$

Let $S: \mathbb{R}^{2n} \to \mathbb{R}^{2n}$ be the Hamiltonian diffeomorphism generated by the function

$$\mathbb{R}^{2n} \ni (x_1, y_1, \cdots, x_n, y_n) \mapsto H(x_1, y_1, \cdots, x_n, y_n) = \eta(x_1) \cdot vy_1 \in \mathbb{R}.$$

(It is not compactly supported!) It is easily checked that

$$S((x_1, y_1, \dots, x_n, y_n) = (x_1, y_1, \dots, x_n, y_n) \quad \text{if } x_1 < -\varepsilon,$$

$$S((x_1, y_1, \dots, x_n, y_n) = (x_1 + v, y_1, \dots, x_n, y_n) \quad \text{if } x_1 > 0.$$

Set $g' := SgS^{-1}$. It is also a compactly supported Hamiltonian diffeomorphism and $||g'||_k = ||g||_k$ by the bi-invariance of the quasi-metric d_k . Clearly

$$g'(K) = Sg(K) = g(K) + (v, 0, \cdots, 0)$$

lies on the right of L'.

Proof of Theorem 3.5. By the definition of the cylindrical capacity, for a sufficiently small $\delta > 0$, there exists a Hamiltonian diffeomorphism ψ such that

$$\psi(\operatorname{supp}(\phi)) \subset B^2(c) \times \mathbb{R}^{2n-2}, \text{ where } c < E(\operatorname{supp}(\phi)) + \delta.$$

By composing with a suitable Hamiltonian diffeomorphism we may assume that $\psi(\operatorname{supp}(\phi))$ sits in $Q \times \mathbb{R}^{2n-2}$, where Q is an open square in the (x_1, y_1) -plane with

area c. Note that the displacement energy of Q is just c. By the example shown in [27, p.17], there exists a Hamiltonian isotopy $\{g_t\} \subset \operatorname{Ham}(\mathbb{R}^2, \omega_0)$ with the time-1 map g satisfying $\operatorname{Length}(g_t) = c$ and $Q \cap g(Q) = \emptyset$. Since the Hamiltonian of the flow $\{g_t\}$ may be chosen to be autonomous, we have $\|g\|_k \leq \operatorname{Length}_k(g_t) = \operatorname{Length}(g_t) = c$. Then for $\tilde{g} := g \times id : \mathbb{R}^{2n} \to \mathbb{R}^{2n}$, we have

$$\|\widetilde{g}\|_k \le \|g\|_k \le c < E(\operatorname{supp}(\phi)) + \delta$$

Obverse that $\psi(\operatorname{supp}(\phi))$ and $\tilde{g}\psi(\operatorname{supp}(\phi))$ have a positive distance. We can construct a hyperplane L lying between $\psi(\operatorname{supp}(\phi))$ and $\tilde{g}\psi(\operatorname{supp}(\phi))$ such that $\psi(\operatorname{supp}(\phi))$ strictly sits in the left side of L. For an arbitrary fixed $f \in \operatorname{Ham}(\mathbb{R}^{2n}, \omega_0)$, since $\operatorname{supp}(f)$ is a compact set of \mathbb{R}^{2n} by the assumption we may choose a hyperplane L'parallel to L such that $\tilde{g}\psi(\operatorname{supp}(\phi))$ strictly lies between L and L' and that $\psi(\operatorname{supp}(f))$ strictly sits in the left side of L'. Applying Lemma 3.7 to $K = \psi(\operatorname{supp}(\phi))$ we get a Hamiltonian diffeomorphism \tilde{g}' such that $\|\tilde{g}'\|_k = \|\tilde{g}\|_k < E(\operatorname{supp}(\phi)) + \delta$ and that $\tilde{g}'(K)$ lies in the right side of L' and hence

$$\widetilde{g}'(\psi(\operatorname{supp}(\phi))) \cap \psi(\operatorname{supp}(f)) = \emptyset.$$

It follows that $C_{(\psi^{-1}\tilde{g}'\psi)\phi(\psi^{-1}\tilde{g}'\psi)^{-1}}f = f$ since $f(\operatorname{supp}(f)) = \operatorname{supp}(f)$. So Lemma 3.6 leads to

$$d_k(f, C_{\phi}f) \le 2^{3k+2} \|\psi^{-1}\widetilde{g}'\psi\|_k = 2^{3k+2} \|\widetilde{g}'\|_k < 2^{3k+2} E(\operatorname{supp}(\phi)) + 2^{3k+2}\delta.$$

Since δ can be chosen arbitrarily small, the desired estimate is obtained.

Finally, as in [16, 5.2] using Lemma 3.1 and Proposition 3.5 we deduce that the functions

$$D_k(f,g) = r_{\infty,k}(g^{-1}f), \quad k = 0, 1, 2, \cdots$$
 (3.7)

give a sequence of non-degenerate bi-invariant quasi-metrics on the group $\operatorname{Symp}_{c}(\mathbb{R}^{2n}, \omega_{0})$ of all compactly supported symplectomorphisms of $(\mathbb{R}^{2n}, \omega_{0})$.

3.2 Banyaga's method

Let (M, ω) be a closed (i.e., compact and without boundary) symplectic manifold. For a smooth path $[a, b] \ni t \mapsto \phi_t \in \text{Symp}(M, \omega)$, (which means the mapping $(x, t) \mapsto \phi_t(x)$ to be smooth), it determines a unique smooth family of symplectic vector fields, $\dot{\phi}_t(x) = \frac{d\phi_t}{dt} \circ \phi_t^{-1}(x)$, whose dual 1-form $i_{\phi_t}\omega$ is closed. When [a, b] = [0, 1] and $\phi_0 = \text{id}, \Phi := \{\phi_t\}$ is called a **symplectic isotopy** on (M, ω) .

Fix a Riemannian metric g on M. A symplectic vector field X on M is said to be a **harmonic vector field** if $i_X \omega$ is a harmonic form; and a smooth path $[a,b] \ni t \mapsto \phi_t \in \text{Symp}(M,\omega)$ is called **harmonic** if each form $i_{\phi_t}\omega$ is harmonic. In particular, a symplectic isotopy $\Phi = \{\phi_t\}$ on (M,ω) is called a **harmonic isotopy** if it is a harmonic path in $\text{Symp}(M,\omega)$. **Lemma 3.8 ([2, Lemma 1])** Any smooth path $[a,b] \ni t \mapsto \phi_t \in \text{Symp}(M,\omega)$ can be decomposed in a unique way as $\phi_t = \rho_t \psi_t$, where $[a,b] \ni t \mapsto \rho_t \in \text{Symp}(M,\omega)$ is a harmonic path and $[a,b] \ni t \mapsto \psi_t \in \text{Ham}(M,\omega)$ is a (smooth) Hamiltonian path. In particular, if ϕ_t is a Hamiltonian path, then $\phi_t = \psi_t$ and $\rho_t = \text{id}$.

Fix a basis $\{h_1, \dots, h_r\}$ of harmonic 1-forms, where $r = \dim H^1(M, \mathbb{R})$. The space of harmonic 1-forms on M is equipped with the following Euclidean metric:

$$|h| = \sum_{i} |\lambda_{i}| \quad \text{if } h = \sum_{i} \lambda_{i} h_{i}.$$
(3.8)

Banyaga defined in [2] the length of a symplectic isotopy $\Phi = \{\phi_t\}$ by

$$l_{HL}(\Phi) = \int_0^1 \left(|\mathscr{H}_t| + \left(\max_x U_t - \min_x U_t \right) \right) dt, \tag{3.9}$$

where \mathscr{H}_t and U_t are smooth families of harmonic 1-forms and functions respectively and satisfy the Hodge decomposition

$$i_{\dot{\phi}_t}\omega = \mathscr{H}_t + dU_t. \tag{3.10}$$

Clearly, for a Hamiltonian isotopy Φ the formula (3.9) reduces to (1.2). As for (1.2) we can prove that $l_{HL}(\Phi)$ is independent of the choice of parametrization of the path Φ . However we don't have $l_{HL}(\Phi) = l_{HL}(\Phi^{-1})$ in general, where $\Phi^{-1} = \{\phi_t^{-1}\}$.

In [2] the energy $e(\phi)$ of any $\phi \in \text{Symp}_0(M, \omega)$ is defined by

$$e(\phi) = \inf_{\Phi} (l_{HL}(\Phi)),$$

where Φ runs over all symplectic isotopies connecting the identity and ϕ . The so called **Hofer-like metric** is the map

$$\|\cdot\|_{HL} : \operatorname{Symp}_0(M,\omega) \to \mathbb{R} \cup \{\infty\}, \ \phi \mapsto \frac{1}{2} \big(e(\phi) + e(\phi^{-1}) \big),$$

which is actually a norm on $\operatorname{Symp}_0(M, \omega)$. Notice that the norm $\|\cdot\|_{HL}$ depends on the choice of the Riemannian metric g on M and the choice of the Euclidean norm $|\cdot|$ on the space of harmonic 1-forms. However, different choices for g and $|\cdot|$ yield equivalent metrics. The **Hofer-like distance** d_{HL} on each connected component of $\operatorname{Symp}(M, \omega)$ is defined by $d_{HL}(\phi, \psi) := \|\phi\psi^{-1}\|_{HL}$. It is right invariant, but not left invariant. When Φ is a Hamiltonian isotopy, (3.9) reduces to (1.2), thus $\|\phi\|_{HL} \leq \|\phi\|_H \ \forall \phi \in \operatorname{Ham}(M, \omega)$. Moreover the subgroup $\operatorname{Ham}(M, \omega)$ is closed in $\operatorname{Symp}(M, \omega)$ endowed with the metric topology defined by $\|\cdot\|_{HL}$. As expected by Banyaga, Buss and Leclercq [6] showed that the restriction of the Hofer-like metric to $\operatorname{Ham}(M, \omega)$ is equivalent to the Hofer metric.

Given any smooth symplectic path $\alpha : [a, b] \to \text{Symp}(M, \omega)$, we have a decomposition as (3.10), $i_{\dot{\alpha}(t)}\omega = \mathscr{H}_t + dU_t$. For every integer $k = 0, 1, 2, \cdots$, we define the *k*-length of α as

$$l_{HL,k}(\alpha) := \sum_{i=0}^{k} \int_{a}^{b} \left[\left| \frac{\partial^{i} \mathscr{H}_{t}}{\partial t^{i}} \right| + \left(\max_{x} \frac{\partial^{i} U_{t}}{\partial t^{i}} - \min_{x} \frac{\partial^{i} U_{t}}{\partial t^{i}} \right) \right] dt.$$
(3.11)

where $|\cdot|$ is as in (3.8). Obviously, $l_{HL,0}(\alpha) = l_{HL}(\alpha)$. But when $k \ge 1$, $l_{HL,k}$ depends on the choice of parametrization of the path which is different from l_{HL} .

A continuous path Φ : $[0,1] \rightarrow \text{Symp}_0(M,\omega)$ is called a **piecewise smooth** symplectic isotopy if there exists a division $0 = t_0 < t_1 < \cdots < t_n = 1, n \in \mathbb{N}$, such that for each $i = 1, \cdots, n$, $\Phi_i = \Phi|_{[t_{i-1}, t_i]}$ is smooth, and $\Phi(0) = \text{id}$. We define the *k*-length of Φ as

$$l_{HL,k}(\Phi) := \sum_{i=1}^{n} l_{HL,k}(\Phi_i).$$

For any $\phi \in \text{Symp}_0(M, \omega)$, let $\Omega(\phi)$ be the set of all piecewise smooth isotopies $\Phi : [0, 1] \to \text{Symp}_0(M, \omega)$ with $\Phi(1) = \phi$. Define the energy of ϕ by

$$e_k(\phi) := \inf \left\{ l_{HL,k}(\Phi) \, \middle| \, \Phi \in \Omega(\phi) \right\},$$

and the corresponding **Hofer-like** k-metric by

$$\|\phi\|_{HL,k} = \frac{1}{2} (e_k(\phi) + e_k(\phi^{-1})), \quad k = 0, 1, \cdots$$

We define the **Hofer-like** k-distance by

$$d_{HL,k}(\phi,\psi) := \|\phi\psi^{-1}\|_{HL,k}$$

on every connected component of $\operatorname{Symp}(M, \omega)$.

Proposition 3.9 On a closed symplectic manifold (M, ω) with a fixed Riemannian metric g, we have

- (i) $\|\cdot\|_{HL} = \|\cdot\|_{HL,0} \le \|\cdot\|_{HL,1} \le \|\cdot\|_{HL,2} \le \cdots$,
- (ii) for each $k, \|\cdot\|_{HL,k}$ is a quasi-norm on Symp₀ (M, ω) .

Proof. (i) Notice that the corresponding Lemma 1.2 still hold in the current context. Hence for $\phi \in \text{Symp}_0(M, \omega)$, we have $e_0(\phi) = e(\phi)$, so $\|\cdot\|_{HL,0} = \|\cdot\|_{HL}$.

(ii) <u>Symmetry</u>: By definition, if $\phi, \psi \in \text{Symp}(M, \omega)$ is in the same connect component, then $d_{HL,k}(\phi, \psi) = d_{HL,k}(\psi, \phi)$.

Non-degeneracy: By (i) and the non-degeneracy of $\|\cdot\|_{HL}$, this is obvious.

<u>The quasi-triangle inequality</u>: Since the proof is similar to the one of property (ii) in Theorem 1.4, we skip some details and only outline the ideas of the proof. For any two symplectomorphisms $\phi, \psi \in \text{Symp}_0(M, \omega)$, we choose

$$\Omega(\phi) \ni \Phi : [0,1] \to \operatorname{Symp}_0(M,\omega) \quad \text{and} \\ \Omega(\psi) \ni \Psi : [0,1] \to \operatorname{Symp}_0(M,\omega).$$

Define the concatenation product $\Phi * \Psi$ of Φ and Ψ by

$$\Phi * \Psi(t) = \begin{cases} \Psi(2t) & 0 \le t \le 1/2 \\ \Phi(2t-1)\psi & 1/2 \le t \le 1. \end{cases}$$

then $\Phi * \Psi \in \Omega(\phi \psi)$. By definition we have

$$e_k(\phi\psi) \leq l_{HL,k}(\Phi * \Psi) \leq 2^k (l_{HL,k}(\Psi) + l_{HL,k}(\Phi(t)\psi))$$

= 2^k (l_{HL,k}(\Phi) + l_{HL,k}(\Psi))

for all $\Phi \in \Omega(\phi)$, $\Psi \in \Omega(\psi)$. Taking the infimum respectively we get

$$e_k(\phi\psi) \leq 2^k (\inf\{l_{HL,k}(\Phi) | \Phi \in \Omega(\phi)\} + \inf\{l_{HL,k}(\Psi) | \Psi \in \Omega(\psi)\})$$

= 2^k(e_k(\phi) + e_k(\psi)).

That is, e_k (and so $\|\cdot\|_{HL,k}$) satisfies the quasi-triangle inequality.

Proposition 3.10 Let (M, ω) be a closed symplectic manifold. Then for each k the subgroup Ham (M, ω) is closed in Symp (M, ω) with respect to the metric topology defined by $\|\cdot\|_{HL,k}$.

Proof. The ideas are similar to those of Theorem 14.2.A in [27]. Suppose there exists a sequence $\{f_n\} \subset \operatorname{Ham}(M, \omega)$ and $\phi \in \operatorname{Symp}(M, \omega)$, satisfying $d_{HL,k}(f_n, \phi) \to 0$ when $n \to \infty$. We intend to prove $\phi \in \operatorname{Ham}(M, \omega)$.

Since $\lim_{n\to\infty} \|f_n \phi^{-1}\|_{HL,k} = 0$, by the definition of $\|\cdot\|_{HL,k}$, for $\forall \varepsilon > 0, \exists N_0 > 0$, such that for each $N \ge N_0$ we have

$$\inf\{l_{HL,k}(\Phi)|\Phi\in\Omega(f_N\phi^{-1})\}<\varepsilon,$$

and so a $\Phi^N \in \Omega(f_N \phi^{-1})$ such that $l_{HL}(\Phi^N) \leq l_{HL,k}(\Phi^N) < \varepsilon$.

Assume the division of Φ^N is given by $0 = t_0 < t_1 < \cdots < t_n = 1$. For each $i = 1, \cdots, n, \Phi_i^N = \Phi^N|_{[t_{i-1}, t_i]}$ is a smooth symplectic path. Next we translate Φ^N into a smooth symplectic isotopy through the precedure of reparametrization. We choose a increasing, surjective smooth function $s_i : [t_{i-1}, t_i] \to [t_{i-1}, t_i]$ for each i, and require s_i is constant near the ends of its interval of definition. Define $\tilde{\Phi}^N : [0, 1] \to \text{Symp}_0(M, \omega)$ by

$$\widetilde{\Phi}^N(t) = \Phi_i^N(s_i(t)), \quad \forall t \in [t_{i-1}, t_i], \ i = 1, \cdots, n,$$

then $\widetilde{\Phi}^N$ is a smooth symplectic isotopy. In fact if the harmonic 1-forms and Hamiltonian functions generated by Φ^N are $\{i\mathscr{H}_t^N\}_{i=1}^n$ and $\{iU_t^N\}_{i=1}^n$ respectively, and the harmonic 1-forms and Hamiltonian functions generated by $\widetilde{\Phi}^N$ are $\widetilde{\mathscr{H}}_t^N$ and \widetilde{U}_t^N respectively, then when $t \in [t_{i-1}, t_i]$, we have

$$\widetilde{\mathscr{H}}_t^N = s_i'(t) \cdot^i \mathscr{H}_{s_i(t)}^N , \qquad \widetilde{U}_t^N = s_i'(t) \cdot^i U_{s_i(t)}^N.$$
(3.12)

By the change of variable formula, we get $l_{HL}(\widetilde{\Phi}^N) = l_{HL}(\Phi^N)$, In particular,

$$\int_0^1 \left| \widetilde{\mathscr{H}}_t^N \right| dt = \sum_{i=1}^n \int_{t_{i-1}}^{t_i} \left| {}^i \mathscr{H}_t^N \right| dt.$$

Since $l_{HL}(\Phi^N) < \varepsilon$, we get $\int_0^1 |\widetilde{\mathscr{H}}_t^N| dt < \varepsilon$.

Recall that the flux is a surjective homomorphism from the universal covering space $\widetilde{\text{Symp}}_0(M,\omega)$ of $\operatorname{Symp}_0(M,\omega)$ to $H^1(M,\mathbb{R})$ given by

$$\operatorname{Flux}(\{\phi_t\}) = \left[\int_0^1 (i_{\dot{\phi}_t}\omega)dt\right] \in H^1(M,\mathbb{R}).$$
(3.13)

 $\Gamma_{\omega} := \operatorname{Flux}(\pi_1(\operatorname{Symp}_0(M, \omega))) \subset H^1(M, \mathbb{R})$ is called the **flux group** of (M, ω) , and is discrete as proved by Ono in [23]. Flux descends to a surjective homomorphism

flux : Symp₀ $(M, \omega) \to H^1(M, \mathbb{R})/\Gamma_\omega$

with kernel Ham (M, ω) (cf. [18]).

For any symplectic isotopy Φ , let $\mathscr{H}(\Phi)$ denote the harmonic representation of the cohomology class $\operatorname{Flux}(\Phi)$. The decomposition (3.10) implies that

$$\mathscr{H}(\Phi) = \int_0^1 \mathscr{H}_t dt.$$

It follows from this that

$$\begin{aligned} \left| \mathscr{H}(\widetilde{\Phi}^{N}) \right| &= \left| \int_{0}^{1} \widetilde{\mathscr{H}}_{t}^{N} dt \right| = \left| \int_{0}^{1} \sum_{i=1}^{r} \widetilde{\lambda}_{i}^{N}(t) h_{i} dt \right| \\ &= \sum_{i=1}^{r} \left| \int_{0}^{1} \widetilde{\lambda}_{i}^{N}(t) dt \right| \\ &\leq \int_{0}^{1} \sum_{i=1}^{r} |\widetilde{\lambda}_{i}^{N}(t)| dt = \int_{0}^{1} |\widetilde{\mathscr{H}}_{t}^{N}| dt < \varepsilon. \end{aligned}$$

where $\widetilde{\mathscr{H}}_{t}^{N}$ is decomposed as $\sum_{i=1}^{r} \widetilde{\lambda}_{i}^{N}(t)h_{i}$.

Starting from $\widetilde{\Phi}^N$, we could construct a smooth symplectic path $\widetilde{\Phi}^N \circ \phi$ connecting ϕ and f_N . Obviously we have $\mathscr{H}(\widetilde{\Phi}^N) = \mathscr{H}(\widetilde{\Phi}^N \circ \phi)$. Choose any symplectic isotopy Ψ from id to ϕ , and any Hamiltonian isotopy α^N from id and f_N , we get a loop $(-\alpha^N)\sharp(\widetilde{\Phi}^N \circ \phi)\sharp\Psi$, whose flux has the harmonic representation

$$\mathscr{H}(\Psi) + \mathscr{H}(\widetilde{\Phi}^N \circ \phi) \in \Gamma_{\omega}$$

because a Hamiltonian path has zero flux. Note that $|\mathscr{H}(\tilde{\Phi}^N)| < \varepsilon$, and that ε is arbitrary small. We deduce that $\mathscr{H}(\Psi) \in \Gamma_{\omega}$ since Γ_{ω} is discrete. Hence $\phi \in \ker(\operatorname{flux}) = \operatorname{Ham}(M, \omega)$.

Remark 3.11 By the method in [2], we can't obtain that $f_N^{-1}\phi$ is a Hamiltonian diffeomorphism for every N large enough, but could only get the distance from $\operatorname{flux}(f_N^{-1}\phi)$ to Γ_{ω} trends to zero as $N \to \infty$.

Han [11] also introduced a method constructing bi-invariant (quasi) metrics on $\operatorname{Symp}(M, \omega)$ from the Hofer metric. For a fixed positive number K, he defined $\|\phi\|_K = \min(\|\phi\|_H, K)$ if $\phi \in \operatorname{Ham}(M, \omega)$, and K otherwise. However, when the above defined quasi-metrics (or metrics) $r_{\alpha,k}$, $\|\cdot\|_K$ are restricted back to $\operatorname{Ham}(M, \omega)$, the induced topologies are in general different from that of the Hofer metric.

4 Concluding remarks

Extensions of the Hofer metric to contact geometry were also studied, see Banyaga and Donato [3], Banyaga and Spaeth [4] and Müller and Spaeth [21]. Our proceeding constructions can be completed in contact manifolds. Let (N, α) be a compact contact manifold of dimension 2n + 1. There exists a one-to-one correspondence between contact isotopies on (N, α) and elements of the space $C^{\infty}(N \times [0, 1])$, $\{f_t\} \leftrightarrow H$, where $i_{X_t}\alpha = H_t$ with $H_t = H(\cdot, t)$ and $X_t = (\frac{d}{dt}f_t) \circ f_t^{-1}$; H is called the contact Hamiltonian function of $\{f_t\}$. Call $\phi \in \text{Diff}(N)$ a strictly contact diffeomorphism if $\phi^*\alpha = \alpha$. A contact isotopy is said to be strictly if each contact diffeomorphism in the isotopy is strictly. Denote by $G_{\alpha}(N)$ the group of strictly contact diffeomorphisms which are strictly contact isotopic to the identity.

Consider a surjective homomorphism from the universal cover $\widetilde{G_{\alpha}(N)}$ of $G_{\alpha}(N)$ to \mathbb{R} given by

$$\{\phi_t\} \mapsto c(\{\phi_t\}) = \frac{1}{\operatorname{Vol}(N)} \int_0^1 \left(\int_N H_t(x)\nu_\alpha\right) dt,$$

where H_t is the contact Hamiltonian of $\{\phi_t\}$ and the canonical volume form $\nu_{\alpha} := \alpha \wedge (d\alpha)^n$. For each $k = 0, 1, 2, \cdots$, we define k-contact length of ϕ_t by

$$\operatorname{Length}_{c,k}(\{\phi_t\}) := |c(\phi_t)| + \sum_{i=0}^k \int_0^1 \left(\max_{x \in N} \frac{\partial^i H}{\partial t^i}(x,t) - \min_{x \in N} \frac{\partial^i H}{\partial t^i}(x,t) \right) dt,$$

and k-contact energy of $\phi \in G_{\alpha}(N)$ by

$$E_{c,k}(\phi) = \inf_{\phi_t} \left(\text{Length}_{c,k}(\{\phi_t\}) \right),$$

where $\{\phi_t\}$ takes over all piecewise smooth strictly contact isotopy from id to ϕ . If k = 0 it becomes the contact length and contact energy in [3, (17) and (20)]. Using the results in [3, 21] we may directly prove

Theorem 4.1 For each $k = 0, 1, \dots$, the mapping

$$d_{c,k}: G_{\alpha}(N) \times G_{\alpha}(N) \to [0,\infty), \quad (\phi,\psi) \mapsto E_{c,k}(\phi\psi^{-1})$$

is a bi-invariant quasimetric on $G_{\alpha}(N)$.

When the contact manifold (N, α) is regular, that is, the Reeb field R_{α} of α generates a free S^1 -action on N, the quotient manifold $B = N/S^1$ is a base of a principal S^1 - bundle $\pi : N \to B$ and B has a canonical symplectic form ω satisfying $\pi^* \omega = d\alpha$. In this case there exists an exact sequence

$$\{1\} \to S^1 \to G_{\alpha}(M) \xrightarrow{p} \operatorname{Ham}(B, \omega) \to \{1\}.$$

As in the proof of [4, Lemma 4.2] it is not hard to prove that $E_{c,k}(\phi) \ge ||p(\phi)||_k$ for any $\phi \in G_{\alpha}(N)$.

As in Hofer geometry it is an important topic to study geodesics of our metrics.

A Appendix: Semigroupoid Metrization Theorem

Given a semigroupoid (G, *), let $G^{(1)} = G$, $G^{(2)} = \{(a, b) \in G \times G : a * b \text{ is well-defined}\}$ and for each $N \in \mathbb{N}, N \ge 2$ let

$$G^{(N)} := \{ (a_1, \cdots, a_N) \in G \times \cdots \times G \mid (a_j, a_{j+1}) \in G^{(2)} \\ \forall j \in \{1, \cdots, N-1\} \}.$$

In particular, if (G, *) is a semigroup, $G^{(N)}$ is just the Cartesian product of N copies of G.

Theorem A.1 ([20, Cor.3.33]) Let (G, *) be a semigroupoid, and assume that $\psi : G \to [0, \infty]$ is a function with the property that there exists a finite constant $C \ge 1$ such that

$$\psi(a * b) \le C(\psi(a) + \psi(b)), \text{ for all } (a, b) \in G^{(2)}.$$
 (A.1)

Introduce

$$\alpha := \frac{1}{1 + \log_2 C} \in (0, 1] \tag{A.2}$$

and define the function $\psi_{\sharp}: G \to [0, \infty]$ by

$$\psi_{\sharp}(a) := \inf\left\{ \left(\sum_{i=1}^{N} \psi(a_i)^{\alpha} \right)^{\frac{1}{\alpha}} : N \in \mathbb{N}, \ (a_1, \cdots, a_N) \in G^{(N)}, \ a = a_1 * \cdots * a_N \right\}.$$
(A.3)

Then $\psi \approx \psi_{\sharp}$. More specifically, with C the same constant as in (A.1), one has

$$(2C)^{-2}\psi \le \psi_{\sharp} \le \psi \quad on \ G. \tag{A.4}$$

In particular, $\psi^{-1}(\{0\}) = \psi_{\sharp}^{-1}(\{0\})$. Furthermore, for every $\beta \in (0, \alpha]$ one has

$$\psi_{\sharp}(a*b)^{\beta} \le \psi_{\sharp}(a)^{\beta} + \psi_{\sharp}(b)^{\beta}, \quad \forall (a,b) \in G^{(2)}, \tag{A.5}$$

and $\psi_{\sharp} = \psi$ on G if and only if $\psi_{\sharp}(a * b)^{\alpha} \leq \psi_{\sharp}(a)^{\alpha} + \psi_{\sharp}(b)^{\alpha}$ for all $(a, b) \in G^{(2)}$. Finally, for each $N \in \mathbb{N}$ the original function ψ satisfies

$$\psi(a_1 \ast \cdots a_N) \le 4C^2 \left\{ \sum_{i=1}^N \psi(a_i)^\beta \right\}^{\frac{1}{\beta}}$$
(A.6)

whenever $a_1, \cdots, a_N \in G$ are such that

$$(a_i, a_{i+1}) \in G^{(2)}$$
 for every $i \in \{1, \cdots, N-1\}$. (A.7)

In particular, if $(a_i)_{i \in \mathbb{N}} \subset G$ is a sequence with the property that (A.7) holds for every number $N \in \mathbb{N}$ with $N \geq 2$, then for each finite number $\beta \in (0, \alpha]$ one has

$$\sup_{N \in \mathbb{N}} \psi(a_1 \ast \cdots a_N) \le 4C^2 \left\{ \sum_{i=1}^{\infty} \psi(a_i)^{\beta} \right\}^{\frac{1}{\beta}}.$$
 (A.8)

Remark A.2 When (G, *) is a group, a function $\psi : G \to [0, \infty]$ is said to be conjugate invariant provided $\psi(b * a * b^{-1}) = \psi(a) \ \forall a, b \in G$. In this case ψ_{\sharp} is also conjugate invariant. In fact, for any $b \in G$, $(a_1, \dots, a_N) \in G^{(N)}$, $a = a_1 * \dots * a_N$, since $b * a * b^{-1} = (b * a_1 * b^{-1}) \dots * (b * a_N * b^{-1})$ and

$$\psi(b * a_i * b^{-1}) = \psi(a_i) \quad \forall i = 1, \cdots, N,$$

it follows from the definition of $\psi_{\sharp}(a)$ in (A.3) that $\psi_{\sharp}(b * a * b^{-1}) = \psi_{\sharp}(a)$.

References

- A. Banyaga, Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique, Comment. Math. Helv. 53(1978), no. 2, 174-227 (French).
- [2] A. Banyaga, A Hofer-like metric on the group of symplectic diffeomorphisms, Proceedings of the 2007 AMS-SIAM Summer Research Conference "Symplectic Topology and Measure presrving Dynamical Systems", Snowbird, UT, Contemporary Math. 2010, 1-23.
- [3] A. Banyaga and P. Donato, Lengths of contact isotopies and extensions of the Hofer metric, Annals of Global Analysis and Geometry, 30(2006),299-312.
- [4] A. Banyaga and P. Spaeth, The group of strictly contact homeomorphisms, arXiv:0812.2461, 2008.
- [5] L. Buhovsky and Y. Ostrover, On the uniqueness of Hofer's geometry, Geom. Funct. Anal. Vol. 21(2011) 1296-1330.
- [6] G. Buss and R. Leclercq, Pseudo-distances on symplectomorphism groups and applications to flux theory, Math. Z. 272(2012), 1001-1022.
- [7] C. Campos-Apanco and A. Pedroza, Bounded symplectic diffeomorphisms and split flux groups, Proc. of Amer. Math. Soc. Vol. 140(2012) 2883-2892.
- [8] Yu. V. Chekanov, Invariant Finsler metrics on the space of Lagrangian embeddings. Math. Z. 234(2000), no. 3, 605C619.
- [9] Y. Eliashberg and L. Polterovich, Bi-invariant metrics on the group of Hamiltonian diffeomorphisms, Internat. J. Math. 4(1993), no. 5, 727-738.
- [10] M. Entov, Quasi-morphisms and quasi-states in symplectic topology, preprint, arXiv:1404.6408, 2014.
- Z. Han, The bounded isometry conjecture for the Kodaira-Thurston manifold and 4-torus, Israel J. Math. 176 (2010) 285-306.
- [12] H. Hofer, On the topological properties of symplectic maps, Proc. Roy. Soc. Edinburgh Sect. A 115 (1990), no. 1-2, 25-38.
- [13] H. Hofer, Estimates for the energy of a symplectic map, Comm. Math. Helv., 68 (1993), 48-72.

- [14] H. Hofer and E. Zehnder, Symplectic invariants and Hamiltonian dynamics. Birkhäuser Verlag, Basel, 1994.
- [15] F. Lalonde and C. Pestieau, Stabilisation of symplectic inequalities and applications, American Mathematical Society Translations (2) 196 (1999), 63-72.
- [16] F. Lalonde and L. Polterovich, Symplectic diffeomorphisms as isometries of Hofer's norm, Topology 36 (1997), no. 3, 711-727.
- [17] F. Lalonde and D. McDuff, *The geometry of symplectic energy*, Ann. of Math.
 (2) 141 (1995), no. 2, 349-371.
- [18] D. McDuff and D. Salamon, Introduction to Symplectic Topology, Clarendon Press Oxford, 1995.
- [19] D. McDuff, Loops in the Hamiltonian group: a survey. Symplectic topology and measure preserving dynamical systems, 127-148, Contemp. Math., 512, Amer. Math. Soc., Providence, RI, 2010.
- [20] D. Mitrea, I. Mitrea, M. Mitrea and S. Monniaux, Groupoid Metrization Theory, Springer Science Business Media, New York, 2013.
- [21] S. Müller and P. Spaeth, On topological contact dynamics I: symplectization and applications of the energy-capacity inequality, arXiv:1110.6705, 2011.
- [22] Y.-G. Oh, Spectral invariants, analysis of the Floer moduli space, and geometry of the Hamiltonian diffeomorphism group, Duke Math. J. 130(2005), 199-295.
- [23] K. Ono, Floer-Novikov cohomology and the flux conjecture, Geom. Funct. Anal. Vol. 16(2006), 981-1020.
- [24] Y. Ostrover, When symplectic topology meets Banach space geometry, preprint, arXiv:1404.6954,2014.
- [25] A. Pedroza, On the bounded isometry conjecture, C. R. Acad. Sci. Paris, Ser. I 349 (2011) 1097-1100.
- [26] L. Polterovich, Symplectic displacement energy for Lagrangian submanifolds, Ergodic Theory Dynam. Systems 13(1993), no. 2, 357-367.
- [27] L. Polterovich, The geometry of the group of symplectic diffeomorphisms, Lectures in Mathematics ETH Zürich, Birkhüser Verlag, Basel, 2001.
- [28] L. Polterovich, Geometry on the group of Hamiltonian diffeomorphisms, Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, 401-410 (electronic).
- [29] M. Schwarz, On the action spectrum for closed symplectically aspherical manifolds, Pacific J. Math. 193(2000), 1046-1095.
- [30] J. C. Sikorav, Systémes hamiltoniens et topologie symplectique, prepnnt, Dipartimento di Matematica dell' Universitá di Pisa, 1990
- [31] C. Viterbo, Symplectic topology as the geometry of generating functions, Math. Ann. 292(1992), 685-710.