

CHARACTERIZATION OF RIESZ SPACES WITH TOPOLOGICALLY FULL CENTER

ŞAFAK ALPAY AND MEHMET ORHON

ABSTRACT. Let E be a Riesz space and let E^\sim denote its order dual. The orthomorphisms $Orth(E)$ on E , and the ideal center $Z(E)$ of E , are naturally embedded in $Orth(E^\sim)$ and $Z(E^\sim)$ respectively. We construct two unital algebra and order continuous Riesz homomorphisms

$$\gamma : ((Orth(E))^\sim)_n \rightarrow Orth(E^\sim)$$

and

$$m : Z(E)'' \rightarrow Z(E^\sim)$$

that extend the above mentioned natural inclusions respectively. Then, the range of γ is an order ideal in $Orth(E^\sim)$ if and only if m is surjective. Furthermore, m is surjective if and only if E has a topologically full center. (That is, the $\sigma(E, E^\sim)$ -closure of $Z(E)x$ contains the order ideal generated by x for each $x \in E_+$.) As a consequence, E has a topologically full center $Z(E)$ if and only if $Z(E^\sim) = \pi \cdot Z(E)''$ for some idempotent $\pi \in Z(E)''$.

1. INTRODUCTION

Let E be a Banach lattice and let $Z(E)$ be its (ideal) center. In general $Z(E)$ is a subalgebra and sublattice of $Z(E')$, the center of the Banach dual E' of E . It is possible to extend this embedding to a contractive algebra and lattice homomorphism of $Z(E)''$ into $Z(E')$. It is natural to ask when the homomorphism would be onto $Z(E')$. It is clear that $Z(E)$ has to be large, since $Z(E')$ is always large. It turns out that the concept of largeness best suiting the center $Z(E)$ in this problem is that $Z(E)$ should be topologically full in the sense of Wickstead [17]. Namely, for each $x \in E_+$, the closure of $Z(E)x$ is the closed ideal generated by x . Then it is shown that the homomorphism is onto $Z(E')$ if and only if $Z(E)$ is topologically full [11, Corollary 2].

Our purpose in this paper is to consider the corresponding problem for a Riesz space E with point separating order dual E^\sim . In the case of Riesz spaces however, there are, in general, two different algebras that act on E

Date: February 10, 2014.

2000 Mathematics Subject Classification. Primary 47B38, 46B42; Secondary 47B60, 46H25.

Key words and phrases. Riesz space, ideal center, orthomorphism, Arens extension.

·
·

which should be considered. Namely, $Orth(E)$, the algebra of the orthomorphisms on E , and its subalgebra and (order) ideal $Z(E)$, the center of E . In the case of Banach lattices $Orth(E) = Z(E)$, therefore this problem does not arise. Recall that an orthomorphism on E is an order bounded operator on E that preserves bands, and $Z(E)$ is the ideal generated by the identity operator on E in the Riesz space $Orth(E)$. In fact $Orth(E)$ is an f -algebra. The ideal center $Z(E)$, on the other hand, is a normed AM -lattice where the identity operator is the order unit of the AM -lattice. Similar to the Banach lattice case, $Orth(E)$ is embedded in $Orth(E^\sim)$ as an f -subalgebra and sublattice. Also as before, $Z(E)$ is embedded in $Z(E^\sim)$ as a subalgebra and sublattice. We construct two unital algebra and order continuous lattice homomorphisms

$$\gamma : (Orth(E)^\sim)_n^\sim \rightarrow Orth(E^\sim)$$

and

$$m : Z(E)'' \rightarrow Z(E^\sim)$$

that extend the two embeddings mentioned above respectively. Then the range of γ is an order ideal in $Orth(E^\sim)$ if and only if m is onto $Z(E^\sim)$ (Corollary 2). Also, m is onto $Z(E^\sim)$ if and only if E has a topologically full center (Proposition 5). That is for each $x \in E_+$, the $\sigma(E, E^\sim)$ -closure of $Z(E)x$ in E contains the ideal generated by x . It follows that E has a topologically full center $Z(E)$ if and only if the ideal center of its order dual E^\sim is given as $Z(E^\sim) = \pi \cdot Z(E)''$ for some idempotent $\pi \in Z(E)''$.

We point out that the proofs of the above mentioned results differ from those used in the Banach lattice case [11]. The method used in this paper owes a lot to the work and results of Huijsmans and de Pagter [8] on the bidual of an f -algebra.

In the Banach lattice case it is shown that if $Z(E)$ is topologically full then it is maximal abelian [11, Corollary 3]. Wickstead [19] showed that the converse is not true. He constructed an interesting AM -lattice with a center that is maximal abelian but is not topologically full. In the case of Riesz spaces, even when $Z(E)$ is topologically full, it need not be maximal abelian. In fact, when $Z(E)$ is topologically full, its commutant in the order bounded operators is $Orth(E)$ (Corollary 4).

If E is a Riesz space, by E^\sim we will denote the Riesz space of order bounded linear functionals on E . All Riesz spaces considered in this paper are assumed to have separating order duals. E_n^\sim will denote the order continuous linear functionals in E^\sim . We let $L_b(E, F)$ denote the space of order bounded linear operators from the Riesz space E into the Riesz space F . When $T : E \rightarrow F$ is an order bounded operator between two Riesz spaces, the adjoint of T carries F^\sim into E^\sim and we will denote it by T' . The dual of a normed space will be denoted by E' . In all terminology concerning Riesz spaces we will adhere to the definitions in [1] and [20].

Let us recall that for any associative algebra A , a multiplication (called the Arens multiplication) can be introduced in the second algebraic dual

A^{**} of A [2]. This is accomplished in three steps: given $a, b \in A$, $f \in A^*$ and $F, G \in A^{**}$, one defines $f \cdot a \in A^*$, $F \cdot f \in A^*$, and $F \cdot G \in A^{**}$ by the equations

$$\begin{aligned}(f \cdot a)(b) &= f(ab) \\ (F \cdot f)(a) &= F(f \cdot a) \\ (F \cdot G)(f) &= F(G \cdot f)\end{aligned}$$

For any Archimedean f -algebra A , the space $(A^\sim)_n^\sim$ is an Archimedean f -algebra with respect to the Arens multiplication [8].

We denote for any $F \in (A^\sim)_n^\sim$ the mapping $f \rightarrow F \cdot f$ by ν_F . The map ν_F is an orthomorphism on A^\sim [8]. The mapping $\nu : (A^\sim)_n^\sim \rightarrow \text{Orth}(A^\sim)$, defined by $\nu(F) = \nu_F$ is an algebra and Riesz homomorphism for any Archimedean f -algebra A . Moreover ν is onto $\text{Orth}(A^\sim)$ if and only if $(A^\sim)_n^\sim$ has a unit element. In that case, ν is injective by Theorem 5.2 in [8]. The main purpose of this paper is to extend the latter result to an arbitrary Riesz space.

2. THE ARENS HOMOMORPHISM

Let E be a Riesz space and consider the bilinear map

$$(1) \quad \text{Orth}(E) \times E \rightarrow E$$

defined by $(\pi, x) \rightarrow \pi(x)$ for each $\pi \in \text{Orth}(E)$ and $x \in E$. Related to (1), we define the following bilinear maps:

$$(2) \quad E \times E^\sim \rightarrow (\text{Orth}(E))^\sim :: (x, f) \rightarrow \psi_{x,f} : \psi_{x,f}(\pi) = f(\pi x)$$

$$(3) \quad E^\sim \times ((\text{Orth}(E))^\sim)^\sim \rightarrow E^\sim :: (f, F) \rightarrow F \bullet f : F \bullet f(x) = F(\psi_{x,f})$$

where $x \in E$, $f \in E^\sim$, $\pi \in \text{Orth}(E)$ and $F \in ((\text{Orth}(E))^\sim)^\sim$. We call the map defined in (3) the *Arens extension* of the map in (1).

For an Archimedean unital f -algebra A , we have $A^{\sim\sim} = (A^\sim)_n^\sim$ by Corollary 3.4 in [8]. Since $\text{Orth}(E)$ is a unital f -algebra with point separating order dual, it is Archimedean. This enables us to use the identification $((\text{Orth}(E))^\sim)^\sim = ((\text{Orth}(E))^\sim)_n^\sim$ throughout this paper. It is straightforward to check that the Arens product on the f -algebra $((\text{Orth}(E))^\sim)_n^\sim$ is compatible with the Arens extension defined in (3), that is, E^\sim is a unital module over $((\text{Orth}(E))^\sim)_n^\sim$.

We use (3) to define a linear operator

$$\gamma : ((\text{Orth}(E))^\sim)_n^\sim \rightarrow L_b(E^\sim) \text{ by } \gamma(F)(f) = F \bullet f$$

for each $F \in ((\text{Orth}(E))^\sim)_n^\sim$ and $f \in E^\sim$. We will call γ the Arens homomorphism of the order bidual of $\text{Orth}(E)$.

Proposition 1. *γ is a unital algebra and order continuous Riesz homomorphism such that $\gamma(((\text{Orth}(E))^\sim)_n^\sim) \subset \text{Orth}(E^\sim)$.*

Proof. It follows from the definition of γ that γ is a positive order continuous unital algebra homomorphism. Also it is easily checked that $\gamma(\pi) = \pi'$ for each $\pi \in \text{Orth}(E)$. Let $F \in ((\text{Orth}(E))^\sim)_n^\sim$ such that $|F| \leq \pi$ for some $\pi \in \text{Orth}(E)$. Then $|\gamma(F)| \leq \gamma(|F|) \leq \gamma(\pi) = \pi'$ by the positivity of γ . Since $\pi' \in \text{Orth}(E^\sim)$, we see that $\gamma(F) \in \text{Orth}(E^\sim)$. Then, since γ is order continuous and the ideal generated by $\text{Orth}(E)$ is strongly order dense in $((\text{Orth}(E))^\sim)_n^\sim$, the range of γ is contained in $\text{Orth}(E^\sim)$. That γ is a Riesz homomorphism follows from the fact that it is an algebra homomorphism by Corollary 5.5 in [9]. \square

If A is an f -algebra then the range of the homomorphism $\nu : (A^\sim)_n^\sim \rightarrow \text{Orth}(A^\sim)$ is contained in the range of the Arens homomorphism γ .

Proposition 2. *Let A be an f -algebra with point separating order dual. The range of the homomorphism $\nu : (A^\sim)_n^\sim \rightarrow \text{Orth}(A^\sim)$ is contained in the range of the Arens homomorphism γ .*

Proof. Let $P : A \rightarrow \text{Orth}(A)$ be the canonical embedding of A into $\text{Orth}(A)$ (i.e., $P(a)(b) = ab$ for all $a, b \in A$). It is well known that $P(A)$ is a sublattice and an algebra ideal in $\text{Orth}(A)$. For each $\mu \in (\text{Orth}(A))^\sim$, define $\hat{\mu} \in A^*$ by $\hat{\mu}(a) = \mu(P(a))$. Since positivity is preserved, it is clear that $\hat{\mu} \in A^\sim$ for each $\mu \in (\text{Orth}(A))^\sim$. Given $a \in A$, $f \in A^\sim$, we have $\psi_{a,f} \in (\text{Orth}(A))^\sim$. Then

$$\hat{\psi}_{a,f}(b) = \psi_{a,f}(P(b)) = f(P(b)(a)) = f(ab) = (f \cdot a)(b)$$

for each $b \in A$. That is $\hat{\psi}_{a,f} = f \cdot a$ for all $a \in A$, $f \in A^\sim$. Now let $F \in (A^\sim)_n^\sim$. Define $\hat{F} \in ((\text{Orth}(A))^\sim)^*$ by $\hat{F}(\mu) = F(\hat{\mu})$ for each $\mu \in (\text{Orth}(A))^\sim$. Since $0 \leq \mu$ implies $0 \leq \hat{\mu}$, we have $0 \leq \hat{F}$ whenever $0 \leq F$. That is $\hat{F} \in ((\text{Orth}(A))^\sim)^\sim = ((\text{Orth}(A))^\sim)_n^\sim$. For each $F \in (A^\sim)_n^\sim$, $f \in A^\sim$ and $a \in A$, we have

$$\begin{aligned} \nu_F(f)(a) &= F \cdot f(a) = F(f \cdot a) = F(\hat{\psi}_{a,f}) = \hat{F}(\psi_{a,f}) \\ &= \hat{F} \bullet f(a) = \gamma(\hat{F})(f)(a). \end{aligned}$$

Hence $\nu_F = \gamma(\hat{F})$. \square

Let us check the behavior of γ in some specific examples.

Example 1. *Let ω denote all sequences and let $A = l^1$ the f -subalgebra of ω consisting of absolutely summable sequences. Then $A^\sim = l^\infty$ and $\text{Orth}(A) = \text{Orth}(A^\sim) = l^\infty$. Since $(A^\sim)_n^\sim = l^1$, ν is the inclusion map $l^1 \rightarrow l^\infty$ so that ν is one-to-one and not onto. On the other hand $((\text{Orth}(A))^\sim)_n^\sim = (l^\infty)''$ and γ is the band projection of $(l^\infty)''$ onto l^∞ . Thus γ is onto and not one-to-one.*

Example 2. *Let $A = c_0$ be the f -subalgebra of ω consisting of the sequences convergent to zero. Then $A^\sim = l^1$ and $\text{Orth}(A) = \text{Orth}(A^\sim) = l^\infty$. Since $(A^\sim)_n^\sim = l^\infty$, ν is the identity map on l^∞ so that ν is one-to-one and onto. On the other hand $((\text{Orth}(A))^\sim)_n^\sim = (l^\infty)''$ and γ is the band projection of $(l^\infty)''$ onto l^∞ . Thus γ is onto and not one-to-one.*

Example 3. Consider $C[0, 1]$ with the product $*$ defined by $a * b = iab$ with $i(x) = x$ for all $x \in [0, 1]$. Then $A = (C[0, 1], *)$ is an Archimedean f -algebra. As shown in [8], $(A^\sim)_n^\sim$ is not semi-prime. Therefore v is not one-to-one and not onto [8]. On the other hand $\text{Orth}(A) = Z(C[0, 1]) = C[0, 1]$ and $\text{Orth}(A^\sim) = Z(C[0, 1]') = C[0, 1]''$. Since $(C[0, 1]^\sim)_n^\sim = C[0, 1]''$, γ is the identity map on $C[0, 1]''$. Therefore γ is one-to-one and onto.

3. RIESZ SPACES WITH TOPOLOGICALLY FULL CENTER

We start with the following definition that is due to Wickstead [17] in the case of Banach lattices.

Definition 1. Suppose E is a Riesz space. Then E is said to have a topologically full center if for each $x \in E_+$ the $\sigma(E, E^\sim)$ -closure of $Z(E)x$ contains the ideal generated by x .

Banach lattices with topologically full center were initiated in [17]. The class of Riesz spaces and the class Banach lattices with topologically full center are quite large. For example, in a σ -Dedekind complete Riesz space E each positive element generates a projection band. Therefore for each $x \in E_+$, $Z(E)x$ is an ideal and $Z(E)$ is topologically full. Also Banach lattices with a quasi-interior point or with a topological orthogonal system have topologically full center [19]. However not all Riesz spaces have topologically full center.

Example 4. (Zaanen [20, p.664]) Let E be the Riesz space of piecewise affine continuous functions on $[0, 1]$. Clearly the ideal generated by the constant 1 function equals E . But, as shown by Zaanen, $Z(E)$ is trivial, that is, it consists of the scalar multiples of the identity. Therefore E does not have a topologically full center.

The first example of an AM-space that has trivial center was given in [5]. A thorough study of Banach lattices with trivial center was undertaken in [18]. We refer the reader to [18] for further examples of Banach lattices with trivial center as well as a careful treatment of the following example of Goulet de Rugy mentioned above.

Example 5. ([5]) Let K be a compact Hausdorff space with a point $p \in K$ such that $\{p\}$ is not a G_δ -set in K (e.g., [12, Example 4, p.140]). Let $C_0(K)$ denote the elements of $C(K)$ that vanish at p . Let H denote the positive unit ball of $C_0(K)'$ with the relative $\sigma(C_0(K)', C_0(K))$ -topology. Let $E = \{f \in C(H) : f(r\mu) = rf(\mu) \text{ for all } r \in [0, 1] \text{ and for each } \mu \in H \text{ with } \|\mu\| = 1\}$. Then E is an AM-space (without order unit). As a sublattice of $l^\infty(H \setminus \{0\})$, one has $E^d = \{0\}$. Therefore $Z(E)$ is embedded in $l^\infty(H \setminus \{0\})$ [15]. Then one may compute that $Z(E)$ consists of continuous bounded functions on $H \setminus \{0\}$ that are constant on the rays (i.e., $g(r\mu) = g(\mu)$ for all $r \in (0, 1]$, for each $\mu \in H$ with $\|\mu\| = 1$). It follows by an argument in [5, p.371] that if there is a non-constant function in $Z(E)$ then $\{0\}$ is a G_δ -set in H . That

in turn implies that $\{p\}$ would be a G_δ -set in K . Therefore $Z(E)$ is trivial and E does not have a topologically full center.

$Z(E)$ is an Archimedean unital f -algebra with order unit. The order unit norm induced on $Z(E)$ is an algebra and lattice norm. $\widehat{Z(E)}$, the norm completion of $Z(E)$, is an AM-space and a partially ordered Banach algebra where the order unit and the algebra unit coincide. Therefore by the Stone algebra theorem $\widehat{Z(E)} \cong C(K)$ (isometric algebra and lattice homomorphism) for some compact Hausdorff space K . (Here $C(K)$ denotes the real valued continuous functions on K .) Then $Z(E)' = \widehat{Z(E)}' = \widehat{Z(E)}^\sim = Z(E)^\sim$ and $Z(E)^{\sim\sim} = Z(E)''$. $Z(E)''$ is an AM-space and with the Arens product, it is a partially ordered Banach algebra (an Archimedean f -algebra with unit) where the order unit and the algebra unit coincide. Therefore $Z(E)'' \cong C(S)$ for some hyperstonian space S . That is the Arens product on $Z(E)''$ coincides with the pointwise product on $C(S)$ [2].

Given the bilinear map

$$(4) \quad Z(E) \times E \rightarrow E$$

defined by $(T, x) \rightarrow Tx$ for each $T \in Z(E)$ and $x \in E$, we define the following bilinear maps:

$$(5) \quad E \times E^\sim \rightarrow Z(E)' :: (x, f) \rightarrow \mu_{x,f} = \psi_{x,f}|_{Z(E)}$$

$$(6) \quad E^\sim \times Z(E)'' \rightarrow E^\sim :: (f, F) \rightarrow F \circ f : F \circ f(x) = F(\mu_{x,f})$$

where $x \in E$, $f \in E^\sim$ and $F \in Z(E)''$. The Arens product on $Z(E)''$ is compatible with the bilinear map defined in (6). That is, E^\sim is a unital module over $Z(E)''$. (6) allows us to define a linear operator $m : Z(E)'' \rightarrow L_b(E^\sim)$ where $m(F)(f) = F \circ f$ for all $f \in E^\sim$ and $F \in Z(E)''$. It is easily checked that $m(T) = \gamma(T)$ whenever $T \in Z(E)$.

We have the following analogue of Proposition 1 for the map m .

Proposition 3. *m is a unital algebra and order continuous lattice homomorphism such that $m(Z(E)'') \subset Z(E^\sim)$.*

Proof. That m is a positive order continuous algebra homomorphism is immediate from the definition of m . That it is a lattice homomorphism follows as in Proposition 1.

For each $F \in Z(E)''$, there is a net $\{T_\alpha\}$ in $Z(E)$ such that $\|T_\alpha\| \leq \|F\|$ and $T_\alpha \rightarrow F$ in $\sigma(Z(E)'', Z(E)')$ -topology. Let $f \in E^\sim$ and $x \in E_+$. Then

$$-\|F\|f(x) \leq F \circ f(x) = \lim_{\alpha} f(T_\alpha x) \leq \|F\|f(x).$$

So $F \in Z(E^\sim)$. □

We will call the map $m : Z(E)'' \rightarrow Z(E^\sim)$ the Arens homomorphism of the bidual of $Z(E)$ (into $Z(E^\sim)$).

Proposition 4. *Let A be a unital f -algebra with point separating order dual. Then the Arens homomorphism of the bidual of $Z(A)$ is onto $Z(A^\sim)$.*

Proof. Let $F \in Z(A^\sim)$ with $0 \leq F \leq 1$. Since A is unital, Theorem 5.2 in [8] implies that the algebra and lattice homomorphism $\nu : (A^\sim)_n^\sim \rightarrow Orth(A^\sim)$ of [8] is one-to-one and onto. Therefore with a slight abuse of notation we will identify $(A^\sim)_n^\sim$ with $Orth(A^\sim)$. In the duality $\langle (A^\sim)_n^\sim, A^\sim \rangle$, A is dense in $(A^\sim)_n^\sim$ with respect to the weak topology. Since the locally solid convex topology $|\sigma|((A^\sim)_n^\sim, A^\sim)$ is compatible with the duality $\langle (A^\sim)_n^\sim, A^\sim \rangle$ [1, Theorem 11.13, p.170], there is a net $\{a_\alpha\}$ in A such that $\{a_\alpha\}$ converges to F in the $|\sigma|((A^\sim)_n^\sim, A^\sim)$ -topology. Lattice operations are continuous in the locally solid convex $|\sigma|((A^\sim)_n^\sim, A^\sim)$ -topology and A is a sublattice of $(A^\sim)_n^\sim$ with $1 \in A$. Therefore we may suppose that $0 \leq a_\alpha \leq 1$. Hence, by the Alaoglu Theorem, there is $T \in Z(A)''$ with $0 \leq T \leq 1$ such that a subnet $\{a_\beta\}$ converges to T in the $\sigma(Z(A)'', Z(A)')$ -topology. Then, when $f \in A^\sim$ and $b \in A$,

$$T \circ f(b) = T(\mu_{b,f}) = \lim_{\beta} \mu_{b,f}(a_\beta) = \lim_{\beta} \psi_{b,f}(a_\beta) = \lim_{\beta} f(a_\beta b).$$

On the other hand, since $\{a_\beta\}$ converges to F in the $|\sigma|((A^\sim)_n^\sim, A^\sim)$ -topology implies convergence also in the $\sigma((A^\sim)_n^\sim, A^\sim)$ -topology, we have

$$F \cdot f(b) = F(f \cdot b) = \lim_{\beta} f \cdot b(a_\beta) = \lim_{\beta} f(a_\beta b).$$

Therefore $m(T) = F$. □

Note that if E is a Riesz space and if we set $A = Orth(E)$, then $Z(A) = Z(E)$. Also, since then A is a unital f -algebra with point separating order dual, Proposition 4 will be true for A . In what follows we will use these facts repeatedly.

Corollary 1. *Let E be a Riesz space. Let $A = Orth(E)$ and m_A denote the Arens homomorphism of the bidual of $Z(A)$ onto $Z(A^\sim)$ (where $Z(A) = Z(E)$). Then*

(1) *the following diagram is commutative*

$$\begin{array}{ccccc} Z(E)'' & \xrightarrow{m} & Z(E^\sim) & \xrightarrow{i} & Orth(E^\sim) \\ \parallel & & & & \uparrow \gamma \\ Z(A)'' & \xrightarrow{m_A} & Z(A^\sim) & \xrightarrow{i} & (A^\sim)_n^\sim \end{array}$$

where i denotes the natural inclusion map;

(2) $\gamma(Z((A^\sim)_n^\sim)) = m(Z(E)'')$.

Proof. (1) Suppose $F \in Z(A)''$ with $0 \leq F \leq 1$. Then, as in the proof of Proposition 4, let $0 \leq a_\alpha \leq 1$ be a net in $Z(A) \subset A$ such that $\{a_\alpha\}$ converges to F in the $\sigma(Z(A)'', Z(A)')$ -topology and also to $m_A(F) \in (A^\sim)_n^\sim$ in the $\sigma((A^\sim)_n^\sim, A^\sim)$ -topology. Since $(A^\sim)_n^\sim = (Orth(E)^\sim)_n^\sim$, for each $x \in E$ and $f \in E^\sim$, we have

$$\begin{aligned} \gamma(m_A(F))(f)(x) &= m_A(F) \bullet f(x) = m_A(F)(\psi_{x,f}) = \lim_{\alpha} \psi_{x,f}(a_\alpha) \\ &= \lim_{\alpha} \mu_{x,f}(a_\alpha) = F(\mu_{x,f}) = m(F)(f)(x). \end{aligned}$$

(2) As in proof of Proposition 4, we may identify $(A^\sim)_n^\sim$ with $Orth(A^\sim)$ via the homomorphism ν . Then $Z(A^\sim) = Z((A^\sim)_n^\sim)$. Since, by Proposition 4, m_A is onto $Z(A^\sim)$, from part (1), we have $\gamma(Z((A^\sim)_n^\sim)) = m(Z(E)''')$. \square

Corollary 2. *Let E be a Riesz space. Then $\gamma((Orth(E)^\sim)_n^\sim)$ is an order ideal in $Orth(E^\sim)$ if and only if $m(Z(E)''') = Z(E^\sim)$.*

Proof. Let $A = Orth(E)$. Suppose $\gamma((A^\sim)_n^\sim)$ is an order ideal in $Orth(E^\sim)$. Since γ is an order continuous lattice homomorphism, the kernel $Ker(\gamma)$ is a band in $(A^\sim)_n^\sim$. Let $(1 - e)$ be the band projection of $(A^\sim)_n^\sim$ onto $Ker(\gamma)$. Then $e \cdot (A^\sim)_n^\sim$ is algebra and lattice isomorphic to the order ideal $\gamma((A^\sim)_n^\sim)$ with $\gamma(e) = 1$. Let $T \in Z(E^\sim)$ with $0 \leq T \leq 1 = \gamma(e)$. So there is $F \in (A^\sim)_n^\sim$ such that $\gamma(F) = T$. Moreover, since γ is an algebra and lattice homomorphism, we may choose F so that $0 \leq F \leq e \leq 1$. That is we may suppose that $F \in Z((A^\sim)_n^\sim)$. Then Corollary 1 part (2) implies there is $G \in Z(E)'''$ such that $m(G) = \gamma(F) = T$.

Conversely, suppose $m(Z(E)''') = Z(E^\sim)$. Suppose $0 \leq T \leq \gamma(F)$ for some $0 \leq F \in (A^\sim)_n^\sim$ and for some $T \in Orth(E^\sim)$. Since $Orth(E^\sim)$ is Dedekind complete, there is $\hat{T} \in Z(Orth(E^\sim)) = Z(E^\sim)$ such that $0 \leq \hat{T} \leq 1$ and $T = \hat{T}\gamma(F)$. By Corollary 1 part (2), there is $G \in Z((A^\sim)_n^\sim)$ such that $\gamma(G) = \hat{T}$. Then $T = \gamma(G)\gamma(F) = \gamma(G \cdot F)$ where $G \cdot F \in (A^\sim)_n^\sim$. \square

Corollary 2 gives the first part of the result stated in the abstract. We will complete the result by showing that the only Riesz spaces that satisfy Corollary 2 are necessarily those with a topologically full center. For Banach lattices a proof of this was given in [11]. Initially we will give the proof of the sufficiency.

Proposition 5. *Let E be a Riesz space. Then E has a topologically full center if and only if the Arens homomorphism $m : Z(E)''' \rightarrow Z(E^\sim)$ is surjective. Then there exists an idempotent $\pi \in Z(E)'''$ such that $Z(E^\sim) = \pi \cdot Z(E)'''$ and $Ker(m) = (1 - \pi) \cdot Z(E)'''$.*

Proof. (Sufficiency) Suppose m is surjective. To show that E has topologically full center it is sufficient to show that each $\sigma(E, E^\sim)$ -closed $Z(E)$ submodule of E is an ideal. This is equivalent to showing that each $\sigma(E^\sim, E)$ -closed $Z(E)$ submodule of E^\sim is an ideal. Let M be a $\sigma(E^\sim, E)$ -closed $Z(E)$ submodule of E^\sim and let $T \in Z(E)'''$. There is a net $\{T_\alpha\}$ in $Z(E)$ that converges to T in the $\sigma(Z(E)''', Z(E)')$ -topology. For each $x \in E$ and $f \in M$ we have

$$T_\alpha \circ f(x) = \mu_{x,f}(T_\alpha) \rightarrow T(\mu_{x,f}) = T \circ f(x).$$

Hence M is a $Z(E)'''$ -submodule of E^\sim . Since $Z(E^\sim) = m(Z(E)''')$, M is a $Z(E^\sim)$ -submodule. E^\sim is Dedekind complete. It is well known that a subspace of E^\sim is an ideal in E^\sim if and only if it is a $Z(E^\sim)$ -submodule of E^\sim (e.g., [19]). Hence M is an ideal in E^\sim . Since m is order continuous (Proposition 3), $Ker(m)$ is a band in $Z(E)'''$. Hence there exists a band

projection $\pi \in Z(E)''$ such that $Ker(m) = (1 - \pi) \cdot Z(E)''$ and $Z(E^\sim) = \pi \cdot Z(E)''$. \square

The proof of the converse requires some preparatory results.

Given a Riesz space E , let A be a unital subalgebra of $Z(E)$. Let A° denote the polar of A in $Z(E)'$ and let A^{oo} denote the polar of A° in $Z(E)''$. By standard duality theory, we have that $A' = Z(E)'/A^\circ$ and $A'' = A^{oo} \subset Z(E)''$. Since A is a normed algebra, A'' is a Banach algebra with the Arens product [2]. In fact A'' is a subalgebra of $Z(E)''$ when $Z(E)''$ has its Arens product.

Lemma 1. *Let A be a unital subalgebra of $Z(E)$. Then A^{oo} is a subalgebra of $Z(E)''$ and the algebra product on A^{oo} is identical with the Arens product on A'' under the canonical isomorphism of A^{oo} with A'' .*

Proof. Let $f \in A^0$ and $a \in A$. It is easily checked that $f \cdot a \in A^0$. Let $F, G \in A^{oo}$. There is a net $\{a_\alpha\}$ in A that converges to F in the $\sigma(Z(E)'', Z(E)')$ -topology. Let $f \in A^0$. Then

$$F \cdot G(f) = F(G \cdot f) = \lim_{\alpha} G \cdot f(a_\alpha) = \lim_{\alpha} G(f \cdot a_\alpha) = 0$$

and $F \cdot G \in A^{oo}$.

On the other hand, denote the isomorphism of A'' onto A^{oo} by $F \rightarrow \widehat{F}$. Given $f \in A'$, let $\widehat{f} \in Z(E)'$ denote any extension of f on $Z(E)$. When $F, G \in A''$, let $\{a_\alpha\}$ and $\{b_\beta\}$ be nets in A that converge to \widehat{F} and \widehat{G} respectively in the $\sigma(Z(E)'', Z(E)')$ -topology. Then, evidently, the respective nets also converge to F and G respectively in the $\sigma(A'', A')$ -topology. For any $f \in A'$, we have

$$F \cdot G(f) = \lim_{\alpha} \lim_{\beta} f(a_\alpha b_\beta) = \lim_{\alpha} \lim_{\beta} \widehat{f}(a_\alpha b_\beta) = \widehat{F} \cdot \widehat{G}(\widehat{f}).$$

\square

Lemma 2. *Let I be a closed algebra ideal in $Z(E)$ and consider $A = Z(E)/I$ with the quotient norm. Then A is a normed algebra and with the Arens product A'' may be identified with the subalgebra of $Z(E)''$ given by*

$$(I^{oo})^d = \{F \in Z(E)'' : |F| \wedge |G| = 0 \text{ for all } G \in I^{oo}\}.$$

Proof. Let $\widehat{Z(E)} = C(K')$ for some compact Hausdorff space K' and let \overline{I} denote the closure of I in $C(K')$. There is a closed subset K of K' such that $\overline{I} = \{a \in C(K') : a(K) = \{0\}\}$ and $C(K')/\overline{I} = C(K)$. Furthermore A is a subalgebra of $C(K)$. In fact $C(K)$ is the completion of A . Since \overline{I} is an order ideal in $\widehat{Z(E)}$, $A' \cong I^0 = (\overline{I})^0$ is a band in $Z(E)'$. Then $A'' \cong Z(E)''/I^{oo} = (I^{oo})^d$, since I^{oo} is a band in $Z(E)''$. It remains to check that the Arens product of A'' is identical with the product on the subalgebra $(I^{oo})^d$. Let $F, G \in Z(E)''$ and $\widehat{F}, \widehat{G} \in A''$ such that $\widehat{F} = F|_{I^0}$, $\widehat{G} = G|_{I^0}$. Let $\{a_\alpha\}$ and $\{b_\beta\}$ be nets in $Z(E)$ that converge to F and G respectively in the $\sigma(Z(E)'', Z(E)')$ -topology. Let $[a] = a + I$ for each $a \in Z(E)$. Then,

it follows that, $\{[a_\alpha]\}$ and $\{[b_\beta]\}$ converge to \widehat{F} and \widehat{G} respectively in the $\sigma(A'', A')$ -topology. Then, for $f \in I^0$, we have

$$\widehat{F} \cdot \widehat{G}(f) = \lim_{\alpha} \lim_{\beta} f([a_\alpha][b_\beta]) = \lim_{\alpha} \lim_{\beta} f(a_\alpha b_\beta) = F \cdot G(f).$$

□

Let J be an ideal in E^\sim . For any $F \in Z(E)''$, $m(F)|_J \in Z(J)$. Each operator in $Z(J)$ has a unique extension to an operator in the ideal center of the band generated by J . Since bands in E^\sim are projection bands, we have $Z(J) = Z(E^\sim)|_J$.

Suppose J is an ideal in E^\sim that separates the points of E . Let A be a unital subalgebra of $Z(E)$.

Definition 2. E is called cyclic with respect to A for the dual pair $\langle E, J \rangle$ if there is $u \in E_+$ such that $(Au)^0 = \{0\}$ in J .

Lemma 3. Let E be a Riesz space and J be an ideal in E^\sim that separates the points of E . Suppose E is cyclic with respect to a unital f -subalgebra A of $Z(E)$ for the dual pair $\langle E, J \rangle$. Then $m(A'')|_J = Z(J)$.

Proof. The norm completion \widehat{A} of A is a unital closed subalgebra of $\widehat{Z(E)}$. Therefore $\widehat{A} = C(K)$ for some compact Hausdorff space K . Furthermore $A^\sim = A' = C(K)'$ and $A'' = C(K)'' = C(S)$ for some hyperstonian space S . (We again mention that the usual multiplication on $C(S)$ is the Arens extension of the product on $C(K)$ as shown in [2]. Also the usual $C(S)$ -module structure of $C(K)'$ via its ideal center is the Arens homomorphism of the bidual of $C(K)$ onto the ideal center of $C(K)'$ (e.g., Proposition 4).) In the rest of the proof, by Lemma 1, we consider A'' as a subalgebra of $Z(E)''$.

Let $u \in E_+$ be a cyclic vector and $f \in J_+$. Then $\mu_{u,f} \in Z(E)'_+$ and $\mu_{u,f}|_A = \widehat{\mu}_{u,f} \in C(K)'_+$. Let P_f be the band projection of $C(K)'$ onto the band $B(\widehat{\mu}_{u,f})$ generated by $\widehat{\mu}_{u,f}$. By the Lebesgue Decomposition Theorem and the Radon-Nikodym Theorem, we have

$$B(\widehat{\mu}_{u,f}) = P_f \cdot C(K)' = L^1(\widehat{\mu}_{u,f}) = \{\mu \in C(K)' : |\mu| \ll \widehat{\mu}_{u,f}\}.$$

The first equality above follows by Proposition 4.

Suppose $e \circ f = 0$ for some idempotent $e \in A'' = C(S)$. Let $\{a_\alpha\}$ be a net in A that converges to e in the $\sigma(A'', A')$ -topology. Then for each $a \in A$,

$$\begin{aligned} e \cdot \widehat{\mu}_{u,f}(a) &= e(\widehat{\mu}_{u,f} \cdot a) = \lim_{\alpha} \widehat{\mu}_{u,f} \cdot a(a_\alpha) = \lim_{\alpha} \widehat{\mu}_{u,f}(aa_\alpha) = \lim_{\alpha} \mu_{u,f}(a_\alpha a) \\ &= \lim_{\alpha} f(a_\alpha a u) = \lim_{\alpha} \mu_{au,f}(a_\alpha) = e(\mu_{au,f}) = e \circ f(au) = \widehat{\mu}_{u,e \circ f}(a) = 0. \end{aligned}$$

Hence $0 \leq e \leq 1 - P_f$. Conversely, since u is a cyclic vector, $\widehat{\mu}_{u,(1-P_f) \circ f} = (1 - P_f) \cdot \widehat{\mu}_{u,f} = 0$ implies that $(1 - P_f) \circ f = 0$. Therefore $1 - P_f = \sup\{e \in C(S) : e \circ f = 0 \text{ and } e = e^2\}$. That is, $a \circ f = 0$ for some $a \in A''$ if and only if $P_f \cdot a = 0$.

Let $T \in Z(E^\sim)$ with $0 \leq T \leq 1$. Let $f \in J_+$. Then $0 \leq \widehat{\mu}_{u,Tf} \leq \widehat{\mu}_{u,f}$ in A' . Therefore, by the Radon-Nikodym Theorem, there is $a_f \in L^\infty(\widehat{\mu}_{u,f}) \subset C(S) = A''$ such that $\widehat{\mu}_{u,Tf} = a_f \cdot \widehat{\mu}_{u,f} = \widehat{\mu}_{u,a_f \circ f}$. Since u is a cyclic vector it follows that $Tf = a_f \circ f$. Suppose $g \in J$ such that $0 \leq g \leq f$. E^\sim is Dedekind complete, there is $G \in Z(E^\sim)$ with $0 \leq G \leq 1$ such that $g = Gf$. Therefore

$$m(a_g)(g) = Tg = T(Gf) = G(Tf) = Gm(a_f)(f) = m(a_f)(Gf) = m(a_f)(g).$$

That is $(a_g - a_f) \circ g = 0$. Therefore $(a_g - a_f) \cdot P_g = 0$. Now suppose $f, g \in J_+$ and $h = f \vee g$. Then $(a_f - a_h) \cdot P_f = (a_g - a_h) \cdot P_g = 0$. Hence $a_g \cdot P_f = a_f \cdot P_g$ for all $f, g \in J_+$. Since S is Stonian and $0 \leq a_f \leq P_f \leq 1$ for all $f \in J_+$, there is a unique $a \in A''$ such that $P_f \cdot a = a_f$ and $(1 - \sup\{P_f : f \in J_+\}) \cdot a = 0$. Then

$$Tf = a_f \circ f = (a \cdot P_f) \circ f = a \circ (P_f \circ f) = a \circ f$$

for all $f \in J_+$. □

Now we are ready to complete the proof of Proposition 5.

Proof. (Necessity) Suppose $x \in E_+$. Let $I(x)$ denote the ideal generated by x in E and let $Ann(x) = \{T \in Z(E) : Tx = 0\}$. $Ann(x)$ is a closed order and algebra ideal in $Z(E)$. Consider the map from $Z(E)$ into $Z(I(x))$ defined by $T \rightarrow T|_{I(x)}$. Clearly the map is a norm reducing positive algebra homomorphism. Since the kernel of this map is equal to $Ann(x)$, the map is also a lattice homomorphism. It induces a norm reducing lattice and algebra homomorphism of $Z(E)/Ann(x)$ into $Z(I(x))$ where $T + Ann(x) = [T] \rightarrow T|_{I(x)}$. The induced map is an isometry. For example, if $T \in Z(E)_+$ with norm $\|T|_{I(x)}\|$ in $Z(I(x))$, then $(T \wedge \|T|_{I(x)}\|1)|_{I(x)} = T|_{I(x)}$ and $\|[T]\| \leq \|T|_{I(x)}\|$. Hence let $A = Z(E)|_{I(x)}$ be the normed unital f -subalgebra of $Z(I(x))$. Then since $A \cong Z(E)/Ann(x)$ (isometric, lattice and algebra homomorphism), we may think of A'' as a subalgebra of $Z(E)''$ (c.f., Lemma 2). Equivalently, we may think of A'' as a subalgebra of $Z(I(x))''$ (c.f., Lemma 1).

Let $J = \{f|_{I(x)} : f \in E^\sim\}$. Clearly J is an ideal in $I(x)^\sim$ that separates the points of $I(x)$. Let $P_x \in Z(E^\sim)$ denote the band projection of E^\sim onto $(I(x)^0)^d$. The map $f|_{I(x)} \rightarrow P_x(f)$ is a lattice homomorphism of J onto the band $(I(x)^0)^d$. Let m_x be the Arens homomorphism of $Z(I(x))''$ into $Z(I(x)^\sim)$ and let m be the same for $Z(E)''$ into $Z(E^\sim)$. We claim that on elements of A'' , m_x restricted to J agrees with m restricted to $(I(x)^0)^d$. Namely, let $F \in A''$, $f \in E^\sim$ and $y \in I(x)$. Choose a net $\{a_\alpha\}$ in A that converges to F in the $\sigma(A'', A')$ -topology. Then

$$\begin{aligned} m_x(F)(f|_{I(x)})(y) &= F \circ f|_{I(x)}(y) = F(\mu_{y,f|_{I(x)}}) = F([\mu_{y,f|_{I(x)}}]) \\ &= \lim_{\alpha} [\mu_{y,f|_{I(x)}}](a_\alpha) = \lim_{\alpha} \mu_{y,f|_{I(x)}}(a_\alpha) = \lim_{\alpha} f(a_\alpha y) \end{aligned}$$

where $[\mu_{y,f}|_{I(x)}] = \mu_{y,f}|_{I(x)} + A^0 \in A' = Z(I(x))'/A^0$. On the other hand

$$\begin{aligned} m(F)(P_x(f))(y) &= P_x(m(F)(f))(y) = m(F)(f)(y) = F \circ f(y) \\ &= F(\mu_{y,f}) = \lim_{\alpha} \mu_{y,f}(a_{\alpha}) = \lim_{\alpha} f(a_{\alpha}y). \end{aligned}$$

In the last string of equalities, the first equality follows because $Z(E^{\sim})$ is commutative. The second equality follows because the range of $1 - P_x$ is $I(x)^0$ in E^{\sim} . Finally the fifth equality follows because $\mu_{y,f} \in \text{Ann}(x)^0 = A'$ in $Z(E)'$. Hence, the claim is verified. In what follows we will keep the notation that we established in this initial part of the proof.

Suppose that E has topologically full center. This means that $I(x)$ is cyclic with respect to the unital f -subalgebra A of $Z(I(x))$ for the duality $\langle I(x), J \rangle$. Given $T \in Z(E^{\sim})$ with $0 \leq T \leq 1$, let $\widehat{T} = T|_{(I(x)^0)^d} \in Z(J)$ where $\widehat{T}(f|_{I(x)})(y) = T(P_x(f))(y)$ for each $f \in E^{\sim}$ and $y \in I(x)$. Then by Lemma 3 there is $a_x \in A''$ with $0 \leq a_x \leq 1$ such that $m_x(a_x) = \widehat{T}$ on J . Then we have that

$$m(a_x)(P_x(f))(y) = m_x(a_x)(f|_{I(x)})(y) = \widehat{T}(f|_{I(x)})(y) = T(P_x(f))(y)$$

for all $f \in E^{\sim}$ and $y \in I(x)$. Since the range of $1 - P_x$ is $I(x)^0$ in E^{\sim} , we have that

$$m(a_x)(f)(y) = T(f)(y)$$

for all $f \in E^{\sim}$ and $y \in I(x)$.

Let $1 - e_x = \sup\{e \in Z(E)'' : e = e^2, m(e)(f)(x) = 0 \text{ for all } f \in E^{\sim}\}$. Since m is order continuous, we have $m(1 - e_x)(f)(x) = 0$ for all $f \in E^{\sim}$. Hence $F \circ f(x) = 0$ for all $f \in E^{\sim}$ for some $F \in Z(E)''$ if and only if $e_x \cdot F = 0$. (Note that $\{\mu_{x,f} : f \in E^{\sim}\}$ is an ideal in $Z(E)'$ and $1 - e_x$ is the band projection of $Z(E)''$ onto the band in $Z(E)''$ that annihilates this ideal.) Now repeating the argument in the proof of Lemma 3, we find a unique $a \in Z(E)''$ such that $0 \leq a \leq 1$, and $e_x \cdot a = a_x$ for each $x \in E_+$. Then, for each $f \in E^{\sim}$ and $x \in E_+$, we have

$$T(f)(x) = m(a_x)(f)(x) = m(a \cdot e_x)(f)(x) = m(e_x)m(a)(f)(x) = m(a)(f)(x).$$

Therefore $m(a) = T$. \square

We will conclude this paper by stating some immediate consequences of Proposition 5.

Corollary 3. *Let A be an f -algebra with point separating order dual such that $(A^{\sim})_n^{\sim}$ has a unit. Then A has topologically full center.*

Proof. By Theorem 5.2 [8], the homomorphism ν of $(A^{\sim})_n^{\sim}$ is onto $\text{Orth}(A^{\sim})$. Then Proposition 2 implies that the Arens homomorphism $\gamma : (\text{Orth}(A^{\sim})_n^{\sim}) \rightarrow \text{Orth}(A^{\sim})$ is also onto. Hence, by Corollary 2, m is onto $Z(A^{\sim})$. Therefore, by Proposition 5, A has topologically full center. \square

Remark 1. *Characterizations of f -algebras A such that $(A^\sim)_n^\sim$ has unit are given in [8],[4],[10]. Related to Corollary 3, we mention that we do not know any examples of semi-prime f -algebras that do not have topologically full center.*

Corollary 4. *Let E be a Riesz space with topologically full center. Suppose T is an order bounded operator on E . Then T commutes with $Z(E)$ if and only if T is in $Orth(E)$.*

Proof. T is order bounded implies $T' : E^\sim \rightarrow E^\sim$. If T commutes with $Z(E)$, then T' commutes with $m(Z(E)^\sim)$. Since $Z(E)$ is topologically full, $m(Z(E)^\sim) = Z(E^\sim)$. Therefore T' commutes with $Z(E^\sim)$. That is, T' commutes with the band projections on E^\sim . Since E^\sim is Dedekind complete, each band in E^\sim is a projection band. So T' is band preserving and therefore $T' \in Orth(E^\sim)$. By a result in [16, Theorem 3.3], $T \in Orth(E)$. \square

In view of Examples 4 and 5, the corollary may fail if $Z(E)$ is not topologically full. On the other hand, the result may be true even when $Z(E)$ is not topologically full. The example constructed by Wickstead in [19] shows this. We refer the reader to the introduction for more detailed information. Corollary 4 shows that $Orth(E)$ is maximal abelian when $Z(E)$ is topologically full. On the other hand, Wickstead's example in [19] shows that the converse is not true.

Corollary 5. *Let E be a Riesz space with topologically full center. Then the following are equivalent:*

- (1) $E^{\sim\sim} = (E^\sim)_n^\sim$.
- (2) m is continuous when its domain has the $\sigma(Z(E)^\sim, Z(E)')$ -topology and its range has the $\sigma(E^\sim, E^{\sim\sim})$ -operator topology.

We leave the straightforward proof of the corollary to the interested reader.

Before stating our final corollary, we want to discuss its content and fix some notation. Let K be a hyperstonian space. That is K is a Stonian compact Hausdorff space and $C(K)$ is a dual Banach space. Let $C(K)_*$ denote the predual of $C(K)$. Recall that $C(K)_* = C(K)'_n$, the order continuous linear functionals on $C(K)$. Hence $C(K)_*$ is a band in the Dedekind complete Banach lattice $C(K)'$. Since $Z(C(K)') = C(K)''$, there is an idempotent $p \in C(K)''$ such that p is the band projection on $C(K)'$ with range $C(K)_*$. That is

$$p \cdot C(K)' = C(K)'_n = C(K)_*.$$

Let E be a Riesz space. Its order dual E^\sim is a Dedekind complete Riesz space. Therefore E^\sim has a topologically full center $Z(E^\sim)$. Furthermore $Z(E^\sim)$ is itself Dedekind complete as a Banach lattice. In fact, it is familiar that $Z(E^\sim) = C(K)$ for some hyperstonian space K . (This will become clear in the proof of the corollary.) Let $m : Z(E^\sim)'' \rightarrow Z(E^{\sim\sim})$ be the Arens homomorphism of the bidual of $Z(E^\sim)$. Since $Z(E^\sim)$ is topologically

full, we have $m(Z(E^\sim)') = Z(E^{\sim\sim})$ and $\text{Ker}(m) = (1 - \pi) \cdot Z(E^\sim)'$ for some idempotent $\pi \in Z(E^\sim)' = C(K)'$ (Proposition 5). We will show that $p \circ E^{\sim\sim} = (E^\sim)_n^\sim$.

Corollary 6. *Let E be a Riesz space with point separating order dual E^\sim . Let m be the Arens homomorphism of the bidual of $Z(E^\sim)$ in $Z(E^{\sim\sim})$. Then*

- (1) $Z(E^\sim)$ is topologically full and $Z(E^\sim) = C(K)$ for some hyperstonian space K .
- (2) There is an idempotent $\pi \in C(K)''$ such that

$$Z(E^{\sim\sim}) = \pi \cdot C(K)'' \text{ and } \text{Ker}(m) = (1 - \pi) \cdot C(K)''.$$
- (3) There is an idempotent $p \in C(K)''$ with $p \leq \pi$ such that

$$p \cdot C(K)' = C(K)_* = C(K)'_n \text{ and } p \circ E^{\sim\sim} = (E^\sim)_n^\sim.$$
- (4) $E^{\sim\sim} = (E^\sim)_n^\sim$ if and only if $p = \pi$.

Proof. 1. Since E^\sim is Dedekind complete, $Z(E^\sim) = C(K)$ is topologically full and K is a Stonian compact Hausdorff space. (It is well known that K is hyperstonian. We include a proof for the sake of completeness.) To show that K is hyperstonian, it is sufficient to see that the order continuous linear functionals on $C(K)$ separate the points of $C(K)$ [12]. Consider $E \subset (E^\sim)_n^\sim \subset E^{\sim\sim}$. Take positive elements $x \in E$, $f \in E^\sim$ and $a_\tau, a \in C(K)$ such that $\{a_\tau\}$ is an increasing net with $\sup a_\tau = a$ in $C(K)$. Then, since E^\sim is Dedekind complete, $\sup a_\tau f = af$ in E^\sim . Therefore $a_\tau f(x) \uparrow af(x)$, since x is an order continuous linear functional on E^\sim . Consider $\mu_{f,x} \in C(K)'$ in the definition process of m , we have $\mu_{f,x}(b) = x(bf) = bf(x)$ for all $b \in C(K)$. Hence it follows that $\mu_{f,x} \in C(K)'_n$ for all $x \in E$ and $f \in E^\sim$. Also it is clear that these linear functionals separate the points of the center $Z(E^\sim) = C(K)$. Therefore K is hyperstonian and $C(K)_* = C(K)'_n$ is the predual of $C(K)$.

2. The existence of π is clear from Proposition 5. An equivalent means of defining $\pi \in C(K)''$ is by observing that π is the supremum of the band projections on $C(K)'$ obtained by considering the bands generated by each linear functional of the form $\mu_{f,x''} \in C(K)'$ when $f \in E^\sim$ and $x'' \in E^{\sim\sim}$.

3. Let $p \in C(K)'' = Z(C(K)')$ be the band projection onto the band $C(K)'_n = C(K)_*$. An equivalent means of defining p would be to observe that p is the supremum of the band projections on $C(K)'$ obtained by considering the bands generated by each linear functional of the form $\mu_{f,x} \in C(K)'$ when $f \in E^\sim$ and $x \in E \subset E^{\sim\sim}$. Hence we have $p \leq \pi$. It remains to show that $p \circ E^{\sim\sim} = (E^\sim)_n^\sim$. Note that for each $f \in E^\sim$ and each $x'' \in E^{\sim\sim}$, we have $\mu_{f,p \circ x''} = p \cdot \mu_{f,x''}$. (Namely, let $\{a_\alpha\}$ be a net in $C(K)$ that converges to p in $\sigma(C(K)'', C(K)')$ -topology. Then

$$\mu_{f,p \circ x''}(a) = p \circ x''(af) = p(\mu_{af,x''}) = \lim_\alpha \mu_{af,x''}(a_\alpha) = \lim_\alpha x''(a_\alpha af)$$

and

$$p \cdot \mu_{f,x''}(a) = p(\mu_{f,x''} \cdot a) = \lim_\alpha \mu_{f,x''} \cdot a(a_\alpha) = \lim_\alpha \mu_{f,x''}(aa_\alpha) = \lim_\alpha x''(a_\alpha af)$$

for each $a \in C(K)$. Here the second set of displayed equalities follow from the definition of the Arens product on the bidual of $C(K)$ when $C(K)$ is considered as a unital f -algebra [8].) But $p \cdot \mu_{f,x''} \in C(K)_* = C(K)'_n$ for each $f \in E^\sim$. By reversing the process we used in part (1), it follows that $p \circ x'' \in (E^\sim)_n^\sim$. Conversely if $x'' \in (E^\sim)_n^\sim$, the process we used in part (1) shows that $\mu_{f,x''} \in C(K)_*$. Therefore

$$\mu_{f,x''} = p \cdot \mu_{f,x''} = \mu_{f,p \circ x''}$$

for each $f \in E^\sim$. That is, $p \circ x'' = x''$ for all $x'' \in (E^\sim)_n^\sim$. So $p \circ E^{\sim\sim} = (E^\sim)_n^\sim$.

Now (4) is clear from parts (2) and (3). \square

Remark 2. *The article [8] has initiated considerable research on Arens product on the biduals of lattice ordered algebras, we include a partial list [3], [6], [7], [13], [14].*

REFERENCES

- [1] Aliprantis, C.D. and Burkinsaw, O., *Positive Operators*, Academic Press, London, 1985
- [2] Arens, R., "Operations induced in function classes", *Monatsh. Math.*, **55**, 1-19, 1951
- [3] Bernau, S.J and Huijsmans, C.B., "The order bidual of almost f -algebras and d -algebras", *Trans. Amer. Math. Soc.*, **347**, 4259-4275, 1995
- [4] Boulabiar, K. and Jaber, J., "The order bidual of f -algebras revisited", *Positivity*, **15**, 271-279, 2011
- [5] Goulet de Rugy, A., "La structure idéale des M-espaces", *J. Math. Pures Appl.*, **55**, 331-373, 1972
- [6] Grobler, J.J., "Commutativity of the Arens product in lattice ordered algebras", *Positivity*, **3**, 357-364, 1999
- [7] Huijsmans, C.B., "The order bidual of lattice ordered algebras.II", *J. Operator Theory*, **22**, 277-290, 1989
- [8] Huijsmans, C.B. and de Pagter, B., "The order bidual of lattice ordered algebras", *J. Func. Anal.*, **59**, 41-64, 1984
- [9] Huijsmans, C.B. and de Pagter, B., "Subalgebras and Riesz subspaces of an f -algebra", *Proc. London Math. Soc.*, **48**, 41-64, 1984
- [10] Jaber, J., " f -algebras with a σ -bounded approximate unit", *Positivity*, **18**, 161-170, 2014
- [11] Orhon, M., "The ideal center of the dual of a Banach lattice", *Positivity*, **14**, 841-847, 2010
- [12] Schaefer, H.H., *Banach lattices and positive operators*, Springer, Berlin, 1974
- [13] Scheffold, E., "Der Bidual von F-Banachverbandsalgebren", *Acta Sci. Math. (Szeged)*, **55**, 167-179, 1991
- [14] Scheffold, E., "Über symmetrische Operatoren auf Banachverbanden und Arens-Regularität", *Czechoslovak Math. J.*, **48(123)**, 747-753, 1998
- [15] Wickstead, A.W., "The structure space of a Banach lattice", *J. Math. Pures Appl.*, **56**, 39-54, 1977
- [16] Wickstead, A.W., "Representation and duality of multiplication operators on archimedean Riesz spaces", *Comp. Math.*, **35**, 225-238, 1977
- [17] Wickstead, A.W., "Extremal structures of cones of operators", *Quart. J. Math. Oxford (2)*, **32**, 239-253, 1981
- [18] Wickstead, A.W., "Banach lattices with trivial centre", *Proc. R. Ir. Acad.*, **88A**, 71-83, 1988

- [19] Wickstead, A.W., "Banach lattices with topologically full centre", *Vladikavkaz Mat. Zh.*, **11**, 50-60, 2009
- [20] Zaanen, A.C., *Riesz Spaces II*, North Holland, Amsterdam, 1983

MATHEMATICS DEPARTMENT, MIDDLE EAST TECHNICAL UNIVERSITY, 06531 ANKARA,
TURKIYE
E-mail address: `safak@metu.edu.tr`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW HAMPSHIRE,
DURHAM, NH 03824, USA
E-mail address: `mo@unh.edu`