

A Note on Weak Hypercyclicity and Linear Fractional Composition Operator

Arman Shokrollahi

Department of Mathematics, West Virginia University, Morgantown, WV 26506, USA.

Abstract: This paper discusses the existence of a sufficient condition for an operator to be weakly hypercyclic. We establish a weak hypercyclicity criterion, and thereupon we can answer questions 5.3 and 5.8 posed by Chan and Sanders in [4]. Lastly, we show that for specific type of composition operator, weak hypercyclicity and hypercyclicity are equivalent.

MSC: 47A16; 47S99; 46A45

Keywords: Banach space • chaos • composition operator • Hardy space • hypercyclicity • linear operator • norm hypercyclic • weak hypercyclicity.

1. Introduction

In this note, we try to briefly discuss a sufficient condition in terms of norm and weakly open sets for an operator on a reflexive Banach space to be weakly hypercyclic. This gives a useful criterion for weak hypercyclicity of operators. We apply this to specific classes of weakly hypercyclic operators including bilateral weighted shifts on $\ell^p(\mathbb{Z})$ with $1 \leq p < \infty$. This, in turn, provides a large class of weakly hypercyclic bilateral shift which are not norm hypercyclic and it answers question 5.3 in [4] in a different way than what Sanders presented in [14]. In the last section, we show hypercyclicity and weak hypercyclicity are equivalent for a composition operator on the space $H(U)$ of all complex-valued functions holomorphic on the open unit disk U or the Hardy space H^2 that is the collection of functions $f \in H(U)$ with $\sum_{n=1}^{\infty} |\hat{f}(n)|^2 < \infty$. So, there are other classes of operators for which norm hypercyclicity and weak hypercyclicity are equivalent and it answers question 5.8 in [4]. The authors in [1–4, 6, 8–11, 13] have already studied weak hypercyclicity and composition operators in various ways. Dodson [7] has also prepared a survey in this regard.

The following theorem summarizes all the necessary conditions which have been obtained for weakly hypercyclic operators. For a proof, see [4] and [6].

Theorem 1.1.

Let T be a bounded operator on Banach space X . If T is a weakly hypercyclic operator, then

- (i) for every norm open set G and weakly open set W , there is some integer $n \geq 0$ such that $T^n G \cap W \neq \emptyset$,
- (ii) the set of weakly hypercyclic vectors for T is norm dense in X ,
- (iii) T^* has no eigenvalue,
- (iv) every component of spectrum T intersect the unit circle.

The following theorem proposes a sufficient condition in terms of open and weakly set for an operator to be weakly hypercyclic. $\text{Ball}_N(X)$ and $\text{Ball}_N^0(X)$ represent the subsets $\{x \in X : \|x\| \leq N\}$ and $\{x \in X : \|x\| < N\}$ of X , respectively, for any integer N .

Theorem 1.2.

Let T be a bounded operator on a reflexive Banach space X such that

- (i) for every norm open set G and weakly open set W , there is some integer $n \geq 0$ such that $T^n G \cap W \neq \emptyset$,
- (ii) there exists some integer M such that if G is a norm open set with $G \cap \text{Ball}(X) \neq \emptyset$ and W is a weakly open set with $W \cap \text{Ball}(X) \neq \emptyset$, then there is some integer $n \geq 0$ such that $T^n(G \cap \text{Ball}(X)) \cap W \cap \text{Ball}_M(X) \neq \emptyset$.

Then, T is weakly hypercyclic.

Proof. The reflexivity of X and Alaoglu's Theorem imply that, for every integer N , $\text{Ball}_N(X)$ is weakly compact and metrizable. Let Γ_N be a collection of countable bases of $\text{Ball}_N(X)$, then $\Gamma = \bigcup \Gamma_N$ is also countable. Suppose $\Gamma = \{W_1, W_2, \dots, W_N, \dots\}$. For each W_j , there is a weakly open set V_j and an integer $N \geq 1$ such that $W_j = V_j \cap \text{Ball}_N(X)$. Let $A = \{(m, j) : T^{-m}V_j \cap \text{Ball}^0(X) \neq \emptyset\}$. A is nonempty. In fact, condition (i) ensures that there is some integer m such that $(m, j) \in A$ for every j . Note that by condition (ii), for every norm open set G and $(m, j) \in A$, there is some integer n such that the set $T^n(G \cap \text{Ball}(X)) \cap T^{-m}V_j \cap \text{Ball}_M(X)$ is nonempty, and consequently $G \cap \text{Ball}(X) \cap T^{-n}[T^{-m}V_j \cap \text{Ball}_M(X)] \neq \emptyset$. So, the set $\bigcup_{n \in \mathbb{N}} T^{-n}[T^{-m}V_j \cap \text{Ball}_M(X)]$ is norm dense in $\text{Ball}(X)$. The Baire's Category Theorem implies that the set

$$WHY := \bigcap_{(m,j) \in A} \bigcup_n T^{-n}[T^{-m}V_j \cap \text{Ball}_M(X)]$$

is also norm dense in $\text{Ball}(X)$. Need to be mentioned that to use the Baire's Category Theorem, it would not be difficult to show that $T^{-n}[T^{-m}W_j \cap \text{Ball}_M(X)] \cap \text{Ball}(X)$ is open in $\text{Ball}(X)$.

Now, we claim every element of the above set is weakly hypercyclic for T . Choosing x in the last set, we show that x is a weakly hypercyclic vector for T . For, suppose W is an arbitrary weakly open set in X , then by condition (i), $T^m \text{Ball}^0(X) \cap W \neq \emptyset$, for some integer $m \geq 0$. There is some integer $N > M\|T\|^m$ such that $T^m \text{Ball}^0(X) \cap W \cap \text{Ball}_N(X) \neq \emptyset$ and consequently $T^{-m}(W \cap \text{Ball}_N(X)) \cap \text{Ball}^0(X) \neq \emptyset$. Since Γ_N is a basis of $\text{Ball}_N(X)$ and $\Gamma_N \subseteq \Gamma$, there exists some $W_j \in \Gamma_N$ with $W_j \cap \text{Ball}_N(X) \subseteq W \cap \text{Ball}_N(X)$. Hence, $T^{-m}(V_j \cap \text{Ball}_N(X)) \cap \text{Ball}^0(X) \neq \emptyset$ when $T^{-m}V_j \cap \text{Ball}^0(X) \neq \emptyset$ for some j . It states that $(m, j) \in A$ and since $x \in WHY(T)$, there is some integer n such that $T^{n+m}x \in V_j$ and $T^n x \in \text{Ball}_M(X)$ when $\|T^{n+m}x\| \leq M\|T\|^m \leq N$. So, $T^{n+m}x \in V_j \cap \text{Ball}_N(X)$ which is a subset of $W \cap \text{Ball}_N(X)$. Thus, $T^{n+m}x \in W$, and so $\text{Orb}(T, x)$ is

weakly dense in X , and T is weakly hypercyclic. (the subject of orbits has been studied in many literature, for example see [12].) \square

2. Weak Hypercyclicity

Theorem 2.1 (Weak Hypercyclicity Criterion).

Let X be a reflexive, separable Banach space, $T \in B(X)$, and there exist two dense subsets Y and Z in X , a sequence $\{n_k\}$ of integers and positive integer M such that

1. $T^{n_k}y \xrightarrow{wk} 0$ for every $y \in Y$,
2. for every $y \in Y$ with $\|y\| \leq 1$, there is some integer N such that

$$\sup\{\|T^{n_k}y\| : k \geq N\} \leq M,$$

3. there exists a linear map $S_k : Z \rightarrow X$ such that for every $z \in Z$,

$$S_k z \rightarrow 0, \quad \text{and} \quad T^{n_k} S_{n_k} z \rightarrow z.$$

Then, T is weakly hypercyclic.

Proof. We apply the preceding theorem (Theorem 1.2). Suppose G is a norm open set and W is a weakly open set. Choose $z \in G \cap Z$, $y \in W \cap Y$, and let $y_k = y + S_k z$, then $y_k \rightarrow y$ in norm topology while $T^{n_k} y_k \xrightarrow{wk} z$. It gives condition (i) of Theorem 1.2.

Again suppose G is a norm open set and W is a weakly open set with $W \cap \text{Ball}(X) \neq \emptyset$ and $G \cap \text{Ball}(X) \neq \emptyset$. Note that $W \cap \text{Ball}^0(X)$ and $G \cap \text{Ball}^0(X)$ are also nonempty. Let $z \in G \cap Y \cap \text{Ball}^0(X)$ and $y \in W \cap Y \cap \text{Ball}^0(X)$, and y_k be the above sequence in the last paragraph. Then for sufficiently large k , $\|y_k\| \leq 1$, $\|T^{n_k} y_k\| \leq M + 1$ and $y_k \rightarrow y$, $T^{n_k} y_k \xrightarrow{wk} z$ as $k \rightarrow \infty$. It states $T^n G \cap W \cap \text{Ball}_{M+1}(X) \neq \emptyset$. Thus, the necessary conditions of part (ii) of Theorem 1.2 are fulfilled and the proof is completed. \square

3. The Unilateral and Bilateral Weighted Shift

Let $\{e_j : j \in \mathbb{N}\}$ be the canonical basis of $\ell^p(\mathbb{N})$. Then, the operator $T : \ell^p(\mathbb{N}) \rightarrow \ell^p(\mathbb{N})$ defined by $T(e_j) = w_j e_{j-1}$ for $j \geq 2$ and $T(e_1) = 0$, for some positive and bounded sequence $\{w_j : j \in \mathbb{N}\}$, is called a *unilateral backward shift*. Also, if $\{e_j : j \in \mathbb{Z}\}$ is the standard basis of $\ell^p(\mathbb{Z})$, then we define the bilateral backward shift T on $\ell^p(\mathbb{Z})$ by $T(e_j) = w_j e_{j-1}$ for all $j \in \mathbb{Z}$ and for some positive and bounded weights $\{w_j : j \in \mathbb{Z}\}$.

Salas [13] introduced the norm hypercyclic unilateral and bilateral weighted shift in terms of the sequence of their weights.

Theorem 3.1 (Salas Theorem [13]). 1. The unilateral weighted shift T with weight sequence $\{w_j : j \geq 1\}$ is hypercyclic if and only if $\sup\{w_1 w_2 \cdots w_n : n \geq 1\} = \infty$.

2. The bilateral weighted shift T with weight sequence $\{w_j : j \in \mathbb{Z}\}$ is hypercyclic if and only if for any given $\epsilon > 0$ and $q \in \mathbb{N}$, there exists an arbitrary large n such that for all $|j| < q$,

$$\prod_{s=1}^n w_{j+s} > \frac{1}{\epsilon}, \quad \text{and} \quad \prod_{s=0}^{n-1} w_{j-s} < \epsilon.$$

The authors in [4] (Theorem 4.1) showed that $\sup\{w_1 w_2 \cdots w_n : n \geq 1\} = \infty$ is equivalent to weak hypercyclicity, hence, weakly hypercyclic unilateral weighted shift is norm hypercyclic. They also introduced a sufficient condition for a bilateral weighted shift on $\ell^p(\mathbb{Z})$ with $2 \leq p < \infty$ to be weakly hypercyclic. This condition is weaker than Salas' condition and more difficult to state. Someone might claim that the following conjecture provides a sufficient condition for weak hypercyclicity of bilateral weighted shift, based on weak hypercyclicity criterion.

Conjecture 1. *The bilateral weighted shift $T(e_j) = w_j e_{j-1}$ on $\ell^p(\mathbb{Z})$ with $1 \leq p < \infty$ is weakly hypercyclic if and only if there exists some sequence $\{n_k\}$ of integers such that*

1. $\sup\{w_j w_{j-1} \cdots w_{j-n_k+1} : k \geq 1, j \in \mathbb{Z}\} < \infty$,
2. for all $j \in \mathbb{Z}$, $w_j w_{j+1} \cdots w_{j+n_k} \rightarrow \infty$ as $k \rightarrow \infty$.

This conjecture is false, because there does not exist a shift that satisfies both conditions (1) and (2) of the conjecture. For a proof, suppose there is an increasing sequence (n_k) satisfying both conditions. Let $\alpha := \sup\{w_j w_{j-1} \cdots w_{j-n_k+1} : k \geq 1, j \in \mathbb{Z}\} < \infty$. If $\alpha = 0$, then $w_j = 0$ for some j , and so condition (2) fails to hold. Assume $\alpha > 0$, then by condition (2), there is n_k such that

$$w_0 w_1 \cdots w_{n_k} > \alpha N. \quad (3.1)$$

For $j = n_k$, we have

$$w_{n_k} w_{n_k-1} \cdots w_1 = w_j w_{j-1} \cdots w_{j-n_k+1} \leq \alpha. \quad (3.2)$$

Combining (3.1) and (3.2) yields

$$\alpha N < w_0 w_1 \cdots w_{n_k} = w_0 (w_1 \cdots w_{n_k}) \leq w_0 \alpha,$$

and therefore, $w_0 > N$. Since this inequality holds for any integer $N \geq 1$, we get $w_0 = \infty$ which is a contradiction.

4. The Linear Fractional Composition Operator

Every holomorphic self-map φ of U induces a linear composition operator $C : H(U) \rightarrow H(U)$ by $C(f)(z) = f(\varphi(z))$ for every $f \in H(U)$ and $z \in U$. Shapiro introduced a complete characterization of hypercyclic operators on $H(U)$. In fact, he showed that C_φ is hypercyclic if and only if φ has no fixed point in U . This condition as we see in the theorem below is equivalent to weak hypercyclicity as well.

Theorem 4.1.

Let φ be a holomorphic self-map on U . Then C_φ is hypercyclic on $H(U)$ if and only if C_φ is weakly hypercyclic.

Proof. Clearly hypercyclicity implies weak hypercyclicity. To prove the other direction, it is enough to show that if C_φ is weakly hypercyclic, then φ has no fixed point in U . Suppose φ has a fixed point $p \in U$ and $\text{Orb}(C_\varphi, f)$ is weakly dense for some $f \in H(U)$. Let g be an arbitrary vector in $H(U)$ and $\epsilon > 0$. The linear functional $\Lambda_p : H(U) \rightarrow \mathbb{C}$ by $\Lambda_p(f) = f(p)$ is continuous on $H(U)$, so there is some positive integer n such that $|\Lambda_p(f \circ \varphi_n - g)| < \epsilon$ when $|f(p) - g(p)| < \epsilon$. It states that any function in $H(U)$ must have the value $f(p)$ at p . But the weakly closure of this orbit cannot be all of $H(U)$ and consequently no C_φ -orbit is weakly dense, i.e., C_φ is not weakly hypercyclic. \square

Now, suppose φ has a linear fractional self-map of U with no fixed point in U . Then, we say φ is *parabolic* if φ has only one fixed point which must lie on the unit circle. Parabolic maps are conjugate to translations of the right half-plane into itself. Also, we say φ is *hyperbolic* if it has two fixed points, one of them lies on the unit circle and the other one is out of \bar{U} which in the automorphism case, both the fixed points must lie on ∂U [3]. Bourdon and Shapiro [2, 3] characterized the hypercyclic composition operator on Hardy space, H^2 . The obtained results state that for a linear fractional self-map φ of U , C_φ is hypercyclic on H^2 unless φ is a parabolic non-automorphism [3].

Theorem 4.2.

Let φ be a linear fractional self-map of U . Then C_φ is hypercyclic on H^2 if and only if C_φ is weakly hypercyclic.

Proof. It is clear that if φ is not parabolic non-automorphism, then hypercyclicity and weak hypercyclicity for C_φ are equivalent. Let φ be parabolic non-automorphism, so it has only one fixed point which lies on ∂U . Without loss of generality, we may take this fixed point to be $+1$. Set

$$\sigma(z) = \frac{1+z}{1-z}, \quad \text{and} \quad \Phi = \sigma \circ \varphi \circ \sigma^{-1}.$$

Then, σ is a linear fractional mapping of U onto the open right half-plane \mathbf{P} , and one can easily check that $\Phi(w) = w + a$ where $\text{Re}(a) > 0$ and $w \in \mathbf{P}$. An easy computation shows that for each $z \in U$,

$$1 - |\varphi_n(z)|^2 = \frac{4\text{Re}(\sigma(z) + na)}{|1 + \sigma(z) + na|^2},$$

and

$$\varphi_n(z) - \varphi_n(0) = \frac{2(\sigma(z) - \sigma(0))}{(\sigma(z) + na + 1)(\sigma(0) + na + 1)}.$$

In addition, for each pair of points $z, w \in U$, and $f \in H^2$, the following estimate holds,

$$|f(z) - f(w)| \leq 2 \|f\| \frac{|z - w|}{(\min\{1 - |w|, 1 - |z|\})^{3/2}}.$$

By substituting z and w by $\varphi_n(z)$ and $\varphi_n(0)$, respectively, and using the last estimate, we get

$$|f(\varphi_n(z)) - f(\varphi_n(0))| \leq \frac{M}{\sqrt{n}},$$

where the constant M depends on f , z , and φ (for more details see [2, 3]).

Now, suppose $\text{Orb}(C_\varphi, f)$ is weakly dense for some vector $f \in H^2$. The linear functional $\Lambda_p : H^2 \rightarrow \mathbb{C}$ by $\Lambda_p(g) = g(p)$ is bounded for every $p \in U$. Let g be an arbitrary vector in H^2 , $z \in U$, and $\epsilon > 0$. There exists a large enough n such that

$$|f(\varphi_n(z)) - f(\varphi_n(0))| < \frac{\epsilon}{4},$$

and

$$|\Lambda_z(f \circ \varphi_n - g)| < \frac{\epsilon}{4}, \quad \text{and} \quad |\Lambda_0(f \circ \varphi_n - g)| < \frac{\epsilon}{4}.$$

Therefore,

$$|f(\varphi_n(z)) - g(z)| < \frac{\epsilon}{4}, \quad \text{and} \quad |f(\varphi_n(0)) - g(0)| < \frac{\epsilon}{4}.$$

Thus, by the triangle inequality, we can deduce $|g(z) - g(0)| < \epsilon$, and consequently $g \equiv g(0)$.

Hence, only constant functions can be weak cluster points of the C_φ -orbit of an H^2 function, and thereof C_φ is not weakly hypercyclic. This fulfills the proof. \square

References

- [1] J. Bes, and A. Peris, Hereditarily Hypercyclic Operators, *J. Functional Analysis* **167** (1999), 94–112.
- [2] P. S. Bourdon, and J. H. Shapiro, Cyclic composition operators on H^2 , *Proc. Symp. Pure Math.* **51** (1990), 43–53.
- [3] P. S. Bourdon, and J. H. Shapiro, “Cyclic Phenomena for Composition Operators”, *Memories Amer. Math. Soc.*, 1997.
- [4] K. C. Chan, and R. Sanders, A weakly hypercyclic operator that is not norm hypercyclic, *J. Operator Theory* **52** (2004), 39–59.
- [5] K. C. Chan, and J. H. Shapiro, The cyclic behavior of translation operators on Hilbert spaces of entire functions, *Indiana Univ. Math. J.* **40** (1991), 1421–1449.
- [6] S. J. Dilworth, and V. G. Troitsky, Spectrum of a weakly hypercyclic operator meets the unit circle, 2002. [arXiv: [math/0208193v1](https://arxiv.org/abs/math/0208193v1)]
- [7] C. T. J. Dodson, A review of some recent work on hypercyclicity, 2012. [arXiv: [1211.4390](https://arxiv.org/abs/1211.4390)]
- [8] N. S. Feldman, Perturbations of hypercyclic vectors, *J. Math. Anal. Appl.* **273** (2002), 67–74.
- [9] G. Godefroy, and J. H. Shapiro, Operators with dense invariant cyclic vector manifolds, *J. Func. Anal.* **98** (1991), 229–269.

-
- [10] C. Kitai, “Invariant Closed Sets for Linear Operators”, Ph.D. Thesis, Univ. of Toronto, 1982.
 - [11] G. R. MacLane, Sequences of derivatives and normal families, *J. d’Analyse Mathématique* **2** (1952), 72–87.
 - [12] S. Rolewicz, On orbits of elements, *Studia Math* **32** (1969), 17–22.
 - [13] H. N. Salas, Hypercyclic weighted shifts, *Trans. Amer. Math. Soc.* **347** (1995) 993–1004.
 - [14] R. Sanders, Weakly supercyclic operators, *J. Math. Anal. Appl.* **292** (2004), 148–159.
 - [15] J. H. Shapiro, [Notes on the dynamics of linear operators](http://www.joelshapiro.org/), 2001. <http://www.joelshapiro.org/>