

Gravitational tests of the Generalized Uncertainty Principle

Fabio Scardigli^{a,b*} and Roberto Casadio^{c†}

^a*Dipartimento di Matematica, Politecnico di Milano,
Piazza Leonardo da Vinci 32, 20133 Milano, Italy*

^b*Institute of Physics, Academia Sinica, Taipei 115, Taiwan*

^c*Dipartimento di Fisica, Università di Bologna and I.N.F.N., Sezione di Bologna,
via Irnerio 46, 40126 Bologna, Italy*

Abstract

We compute the corrections to the Schwarzschild metric necessary to reproduce the Hawking temperature derived from a Generalized Uncertainty Principle (GUP), so that the GUP deformation parameter is directly linked to the deformation of the metric. Using this modified Schwarzschild metric, we compute corrections to the standard General Relativistic predictions for the light deflection and perihelion precession, both for planets in the solar system and for binary pulsars. This analysis allows us to set bounds for the GUP deformation parameter from well-known astronomical measurements.

PACS 04.60 - Quantum theory of gravitation.

1 Introduction

Research on generalizations of the uncertainty principle of quantum mechanics has nowadays a long history [1]. One of the main lines of investigation focuses on understanding how the Heisenberg Uncertainty Principle (HUP) should be modified once gravity is taken into account. Given the pivotal rôle played by gravitation in these arguments, it is not surprising that the most relevant modifications to the HUP have been proposed in string theory, loop quantum gravity, deformed special relativity, and studies of black hole physics [2, 3, 4, 5, 6, 7], just to mention some of the most notable frameworks.

An interesting novelty, emerged during the last decade or so, is a lively debate on the *measurable* features of various kinds of Generalized Uncertainty Principles (GUPs). From more theoretical shores, the discussion has therefore landed on the ground of experimental predictions about the *size* of these modifications, and several experiments have been proposed to test GUPs in the laboratory. Among the more elaborated proposals are those, for example, of the groups of Brukner and Cerdonio [8].

*Corresponding author. E-mail: fabio@phys.ntu.edu.tw

†E-mail: casadio@bo.infn.it

Studies that aim at putting bounds on the deforming parameter of the GUP, heretofore denoted by β , date back at least to Brau [9], and can be roughly divided into three different categories (actually, only two, as we will see). In the first group one finds papers such as those of Brau [9], Vagenas [10], Nozari [11], which use a specific representation of the operators in the deformed fundamental commutator¹

$$[\hat{X}, \hat{P}] = i\hbar \left(1 + \beta \frac{\hat{P}^2}{m_p^2} \right), \quad (1.1)$$

in order to compute corrections to quantum mechanical predictions, such as energy shifts in the spectrum of the hydrogen atom, or to the Lamb shift, the Landau levels, Scanning Tunneling Microscope, charmonium levels, etc. The bounds so obtained on β are very stringent, but the drawback of this approach is a potentially strong dependence of the expected shifts on the specific representation chosen for the variables X and P in the fundamental commutator (1.1).

In the second group, we can find the works of, e.g., Chang [12], Nozari and Pedram [13], where a deformation of classical Newtonian mechanics is introduced by modifying the standard Poisson brackets in a way that resembles the quantum commutator,

$$[\hat{x}, \hat{p}] = i\hbar (1 + \beta_0 \hat{p}^2) \quad \Rightarrow \quad \{X, P\} = (1 + \beta_0 P^2), \quad (1.2)$$

where $\beta_0 = \beta/m_p^2$. In particular, Chang in Ref. [12] computes the precession of the perihelion of Mercury directly from this GUP-deformed Newtonian mechanics, and interprets it as an extra contribution to the well known precession of 43"/century due to General Relativity (GR). He then compares this global result with the observational data, and the very accurate agreement between the GR prediction and observations leaves Chang not much room for possible extra contributions to the precession. In fact, he obtains the tremendously small bound $\beta \lesssim 10^{-66}$. A problem with this approach is that a GUP-deformed Newtonian mechanics is simply superposed to the usual GR theory linearly. One may argue that a modification of GR at order β should likewise be considered, but this is however omitted in Ref. [12]. In other words, it is not clear why the two structures, GR and GUP-modified Newtonian mechanics, should coexist independently, and why the two different precession errors add into a final single precession angle. As a matter of fact, in the limit $\beta \rightarrow 0$, Ref. [12] recovers the Newtonian mechanics, to which GR corrections must be added as an extra structure. Clearly, the physical relevance of this approach and the bound that follows for β , remain therefore questionable.

Finally, a third group of works on the evaluation of β contains, for example, papers by Ghosh [14] and Pramanik [15]. They use a covariant formalism, first defined in Minkowski space, with the metric $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$, which can be easily generalized to curved space-times via the standard procedure $\eta_{\mu\nu} \rightarrow g_{\mu\nu}$. These papers should however be considered as belonging to the second group. In fact, a closer look reveals that they also start from a deformation of classical Poisson brackets, although posited in covariant form. From the deformed covariant Poisson brackets, they obtain interesting consequences, like a β -deformed geodesic equation, which leads to a violation of the Equivalence Principle. They do not deform the field equations or the metric. In Appendix A, however, we show that the violation of the Equivalence Principle is completely due to the deformation of the Poisson brackets, and has nothing to do with the covariant formalism, or

¹We shall work with $c = k_B = 1$, but explicitly show the Newton constant G_N and Planck constant \hbar . We also recall that the Planck length is defined as $\ell_p^2 = G_N \hbar / c^3$, the Planck energy as $\mathcal{E}_p \ell_p = \hbar c / 2$, and the Planck mass as $m_p = \mathcal{E}_p / c^2$, so that $G_N = \ell_p / 2 m_p$ and $\hbar = 2 \ell_p m_p$.

with a deformation of the GR field equations or solutions, or of the geodesic equation. Nonetheless, the Ghosh-Pramanik formalism remains covariant when $\beta \rightarrow 0$ and reproduces standard GR results in the limit $\beta \rightarrow 0$ (this differs, in general, from the results obtained by papers in the second group).

The novelties of our approach, when compared with the previous ones, are many and various. The main point is to start directly from a quantum mechanical effect, the Hawking evaporation, for which the GUP is necessarily relevant, rather than postulating specific representations of canonical operators or modifications of the classical equations of motion. We connect the deformation of the Schwarzschild metric directly to the uncertainty relation, without relying on a specific representation of commutators. We leave the Poisson brackets and classical Newtonian mechanics untouched, and recover GR, and standard quantum mechanics, in the limit $\beta \rightarrow 0$. In particular, we preserve the Equivalence Principle, and the equation of motion of a test particle is still given by the standard geodesic equation. In the present work, this is obtained by deforming a specific solution of the standard GR field equations, namely the Schwarzschild metric. In Appendix B, we display a non-relativistic analog of this procedure. A further, more profound, step in this direction would be to formulate from scratch the deformed field equations of GR, not just assume a deformed solution (as we did), or a deformed kinematics (as Chang, Ghosh, etc. did). This task is left for future developments.

2 Deforming the Schwarzschild metric

In this section, we start from a known way of deriving the Hawking temperature directly from the metric of a black hole, and then show how the GUP modifies the Hawking temperature. These two steps will pave the road to a deformation of the Schwarzschild metric, constructed so as to reproduce the GUP-modified Hawking temperature.

2.1 Standard mass-temperature relation

We consider here a space-time with a metric that locally has the form

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = F(r) dt^2 - F(r)^{-1} dr^2 - r^2 d\Omega^2, \quad (2.1)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$. For the typical cases we shall consider later on, one has

$$F(r) = 1 - \frac{2G_N M}{r} + \frac{G_N Q^2}{r^2} + \Lambda r^2, \quad (2.2)$$

however, we do not require any specific form for $F(r)$ for the moment. Note that the time-like coordinate is chosen as $x^0 = t$, the parameters M (mass), Q (electric charge), Λ (cosmological constant, up to a factor) are real and continuous, with $\Lambda < 0$ corresponding to a de Sitter space-time, and $\Lambda > 0$ to an anti de Sitter space-time. The horizons (if any), are located at the positive zeros of the function $F(r)$ (see, for example, Ref. [16]).

We shall loosely follow the derivation in Ref. [17]. Suppose $r = r_H$ is an horizon, so that $F(r_H) = 0$, and consider $r \geq r_H$. After the Wick rotation $t \rightarrow i\tau$ the metric reads

$$ds^2 = -[F(r) d\tau^2 + F(r)^{-1} dr^2 + r^2 d\Omega^2]. \quad (2.3)$$

In the region just outside the horizon, $r \gtrsim r_H$, we perform the coordinate transformation $(\tau, r) \rightarrow (\alpha, R)$ defined by

$$\begin{cases} R d\alpha = F(r)^{1/2} d\tau \\ dR = F(r)^{-1/2} dr, \end{cases} \quad (2.4)$$

so that the Euclidean metric becomes

$$ds^2 = -[R^2 d\alpha^2 + dR^2 + r^2(R) d\Omega^2] . \quad (2.5)$$

The first two terms in ds^2 represent the length element squared of flat 2-dimensional Euclidean space in polar coordinates. The Euclidean time τ is therefore proportional to the polar angle α . Now, denote the period of τ by Θ (the period of α is of course 2π). From the second equation we see that R is a function of r only. Therefore, integrating the first of the Eq. (2.4) over a full period, we get

$$R(r) \int_0^{2\pi} d\alpha = \sqrt{F(r)} \int_0^\Theta d\tau \implies 2\pi R(r) = \Theta \sqrt{F(r)} . \quad (2.6)$$

We are interested in what happens just outside the horizon, therefore we can expand $F(r)$ around r_H . Namely, for $(r - r_H)$ small we obtain

$$\sqrt{F(r)} = [F(r_H) + F'(r_H)(r - r_H) + \dots]^{1/2} \simeq \sqrt{F'(r_H)} \sqrt{r - r_H} . \quad (2.7)$$

Eq. (2.6) thus becomes

$$2\pi R(r) \simeq \Theta \sqrt{F'(r_H)} \sqrt{r - r_H} . \quad (2.8)$$

The second of the equations (2.4) likewise becomes

$$dR(r) \simeq \frac{dr}{\sqrt{F'(r_H)} \sqrt{r - r_H}} , \quad (2.9)$$

which yields

$$R(r) \simeq \frac{2}{\sqrt{F'(r_H)}} \sqrt{r - r_H} . \quad (2.10)$$

This, together with Eq. (2.8), implies

$$\Theta = \frac{4\pi}{F'(r_H)} . \quad (2.11)$$

Now, Θ is the period of the Euclidean time, which means, by general principles of QFT, that a quantized scalar field outside the horizon lives in a heat bath with temperature $T = \hbar \Theta^{-1}$. To conclude, the temperature of the black hole horizon as seen by a distant observer is in general given by

$$T = \hbar \frac{F'(r_H)}{4\pi} . \quad (2.12)$$

In particular, for a Schwarzschild black hole the function $F(r)$ is given by $Q = \Lambda = 0$ in Eq. (2.1) above, the horizon is at $r_H = 2 G_N M$, and we get

$$T_H = \frac{\hbar}{8\pi G_N M} , \quad (2.13)$$

which is the well-known Hawking temperature.

2.2 GUP modified mass-temperature relation

The most common form of deformation of the Heisenberg uncertainty relation (and the form of GUP that we are going to study in this paper) is without doubt the following

$$\Delta x \Delta p \geq \frac{\hbar}{2} \left(1 + \beta \frac{4 \ell_p^2}{\hbar^2} \Delta p^2 \right) = \frac{\hbar}{2} \left[1 + \beta \left(\frac{\Delta p}{m_p} \right)^2 \right], \quad (2.14)$$

which, for mirror-symmetric states (with $\langle \hat{p} \rangle^2 = 0$), can be equivalently written in terms of commutators as

$$[\hat{x}, \hat{p}] = i\hbar \left[1 + \beta \left(\frac{\hat{p}}{m_p} \right)^2 \right], \quad (2.15)$$

since $\Delta x \Delta p \geq (1/2)|\langle [\hat{x}, \hat{p}] \rangle|$. The dimensionless parameter β is usually assumed to be of order one, in the most common quantum gravity formulations.

As is well known from the argument of the Heisenberg microscope [19], the size δx of the smallest detail of an object, theoretically detectable with a beam of photons of energy E , is roughly given by

$$\delta x \simeq \frac{\hbar}{2E}, \quad (2.16)$$

since larger and larger energies are required to explore smaller and smaller details. From the uncertainty relation (2.14), we see that the GUP version of the standard Heisenberg formula (2.16) is

$$\delta x \simeq \frac{\hbar}{2E} + 2\beta \ell_p^2 \frac{E}{\hbar}. \quad (2.17)$$

which relates the (average) wavelength of a photon to its energy E . (The standard dispersion relation $E = pc$ is assumed.) Conversely, with the relation (2.17) one can compute the energy E of a photon with a given (average) wavelength $\lambda \simeq \delta x$. Following the arguments of Refs. [20, 21, 22, 23, 24, 25, 26], we can consider an ensemble of unpolarized photons of Hawking radiation just outside the event horizon of a Schwarzschild black hole. From a geometrical point of view, it is easy to see that the position uncertainty of such photons is of the order of the unmodified Schwarzschild radius

$$R_H = 2 G_N M. \quad (2.18)$$

An equivalent argument comes from considering the average wavelength of the Hawking radiation, which is of the order of the geometrical size of the hole. We can estimate the uncertainty in photon position as

$$\delta x \simeq 2\mu R_H, \quad (2.19)$$

where the proportionality constant μ is of order unity and will be fixed soon. According to the equipartition principle, the average energy E of unpolarized photons of the Hawking radiation is simply related with their temperature by

$$E = T. \quad (2.20)$$

Inserting Eqs. (2.19) and (2.20) into the formula (2.17), we have

$$4\mu G_{\text{N}} M \simeq \frac{\hbar}{2T} + 2\beta G_{\text{N}} T . \quad (2.21)$$

In order to fix μ , we consider the semiclassical limit $\beta \rightarrow 0$, and require that formula (2.21) predicts the standard semiclassical Hawking temperature (2.13), that is $T(\beta \rightarrow 0) = T_{\text{H}}$. This fixes $\mu = \pi$, so that we have

$$M = \frac{\hbar}{8\pi G_{\text{N}} T} + \beta \frac{T}{2\pi} . \quad (2.22)$$

This is the mass-temperature relation predicted by the GUP for a Schwarzschild black hole. Of course this relation can be easily inverted, to get

$$T = \frac{\pi}{\beta} \left(M - \sqrt{M^2 - \frac{\beta}{\pi^2} m_{\text{p}}^2} \right) , \quad (2.23)$$

where we used $\hbar/G_{\text{N}} = 4m_{\text{p}}^2$. Since, however, the term proportional to β is small, especially for solar mass black holes with $M \gg m_{\text{p}}$, we can expand in powers of β , namely

$$T = T_{\text{H}} \left(1 + \frac{\beta m_{\text{p}}^2}{4\pi^2 M^2} + \dots \right) . \quad (2.24)$$

To zero order in β , we recover the usual Hawking formula (2.13). Already from here we can extract an interesting estimate of β , since the previous expansion is valid only if

$$\frac{\beta m_{\text{p}}^2}{4\pi^2 M^2} \ll 1 , \quad (2.25)$$

which means $\beta \ll 1.3 \cdot 10^{78}$ for a solar mass black hole with $M \simeq 10^{38} m_{\text{p}}$.

2.3 GUP modified Schwarzschild metric

We can legitimately wonder what kind of (deformed) metric would predict a Hawking temperature like the one inferred from the GUP relation (2.22), for a given β . Since we are interested only in small corrections to the Hawking formula, we can consider a deformation of the Schwarzschild metric of the kind

$$F(r) = 1 - \frac{2G_{\text{N}} M}{r} + \varepsilon \frac{G_{\text{N}}^2 M^2}{r^2} , \quad (2.26)$$

and we shall look for the lowest order correction in ε . Note however that, since $R_{\text{H}}/r \sim 10^{-5}$ on the surface of the Sun, the term proportional to ε can still be considered small even if ε is relatively large. The horizons are now given by solutions of the equation

$$r^2 - 2G_{\text{N}} M r + \varepsilon G_{\text{N}}^2 M^2 = 0 . \quad (2.27)$$

We choose the root closest to the unmodified Schwarzschild radius (2.18), namely

$$r_{\text{H}} = R_{\text{H}} \frac{1 + \sqrt{1 - \varepsilon}}{2} , \quad (2.28)$$

which is valid for $\varepsilon \leq 1$ (and possibly negative). Then

$$F'(r) = \frac{2 G_N M}{r^2} - 2 \varepsilon \frac{G_N^2 M^2}{r^3} \quad (2.29)$$

and

$$F'(r_H) = \frac{2}{G_N M} \frac{\sqrt{1-\varepsilon}}{(1 + \sqrt{1-\varepsilon})^2} \simeq \frac{1}{R_H} \left(1 - \frac{\varepsilon^2}{16} + \dots \right), \quad (2.30)$$

where the last expansion is valid for $|\varepsilon| \ll 1$. Hence, the temperature predicted by this deformed Schwarzschild metric is

$$T(\varepsilon) = \hbar \frac{F'(r_H)}{4\pi} = \frac{\hbar}{2\pi G_N M} \frac{\sqrt{1-\varepsilon}}{(1 + \sqrt{1-\varepsilon})^2}, \quad (2.31)$$

which must coincide with the temperature $T(\beta)$ predicted by Eq. (2.23), for any given β . That is, $T(\varepsilon)$ should solve Eq. (2.22),

$$M = \frac{\hbar}{8\pi G_N T(\varepsilon)} + \frac{\beta}{2\pi} T(\varepsilon), \quad (2.32)$$

which yields a relation between β and ε ,

$$\beta(\varepsilon) = -\pi^2 \frac{G_N M^2}{\hbar} \frac{\varepsilon^2}{1-\varepsilon}. \quad (2.33)$$

For $|\varepsilon| \ll 1$, to the lowest order in ε , we thus get

$$\beta = -\pi^2 \varepsilon^2 \frac{M^2}{4 m_p^2}, \quad (2.34)$$

where we recall that both β and ε are dimensionless. It is now of great interest to note that Eq. (2.33) forces us to admit that $\beta < 0$, since $\varepsilon \leq 1$. Although quite unexpected, this is a suggestion of fundamental importance. It seems that a metric is able to reproduce the GUP-deformed Hawking temperature *only if* the deforming parameter β is *negative*. We already encountered a situation like this when we studied the uncertainty relation formulated on a crystal lattice [27]. This could be a further hint that the physical space-time has actually a lattice or granular structure at the level of the Planck scale.

3 Light deflection by deformed Schwarzschild metric

Having established a connection between the GUP parameter β and the deformation ε of the Schwarzschild metric, we are now in a position to compute the physical (possible observable) consequences of such a deformed metric. To begin with, we examine the unbound orbits around a massive body, namely the light deflection by the Sun. Our treatment roughly follows that of Ref. [28].

With reference to Fig. 1, we consider a polar coordinate system centred in the Sun, with (ϕ, r) labelling the position of the incoming photon, $\phi = \phi(r)$ describes the photon orbit, R_\odot is the




Figure 1: Deflection of light by the Sun (refer to the text).

radius of the Sun, r_0 the minimum distance between the photon and Sun. The photon orbit is thus described by

$$\phi(r) - \phi(\infty) = \int_{\infty}^r \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 F(r_0) - F(r) \right]^{-1/2} dr . \quad (3.1)$$

The global deflection angle of the orbit from a straight line is just twice its change from ∞ to r_0 , of course minus π ,

$$\Delta\phi \equiv 2 |\phi(r_0) - \phi(\infty)| - \pi . \quad (3.2)$$

This integral can be evaluated exactly by using elliptic integrals, which, however, can only be numerically computed by expanding in a suitable small parameter. It is both easier and more useful to expand before integrating. Care must be taken in choosing the right small parameter, which will also help in identifying the finite part of the integral. Here physical intuition comes in help. In fact, if the central body had a negligible mass, then $F(r) \rightarrow 1$ (the Minkowskian limit), and the trajectory of the photon would be a (almost) straight line. Departure from a straight line increases as the central mass $M \sim R_H$ increases, as well as if the minimum distance from the source r_0 decreases. The right parameter by which expanding the integral (3.1) is thus the ratio R_H/r_0 .

Now, the argument of the integral can be written as

$$\frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 F(r_0) - F(r) \right]^{-1/2} = \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 - 1 \right]^{-1/2} \times \left[F(r_0) + \frac{F(r_0) - F(r)}{(r/r_0)^2 - 1} \right]^{-1/2} , \quad (3.3)$$

and we find, to first order in ε and to the second order in R_H/r_0 ,

$$\begin{aligned} \left[F(r_0) + \frac{F(r_0) - F(r)}{(r/r_0)^2 - 1} \right]^{-1/2} &= \left[1 - \frac{2 G_N M}{r_0} + \varepsilon \frac{G_N^2 M^2}{r_0^2} - \frac{2 G_N M r_0}{r(r+r_0)} + \varepsilon \frac{G_N^2 M^2}{r^2} \right]^{-1/2} \\ &\simeq 1 + \frac{G_N M}{r_0} + (3 - \varepsilon) \frac{G_N^2 M^2}{2 r_0^2} + \frac{G_N M r_0}{r(r+r_0)} \left(1 + \frac{3 G_N M}{r_0} \right) \\ &\quad - \varepsilon \frac{G_N^2 M^2}{2 r^2} + \frac{3 G_N^2 M^2 r_0^2}{2 r^2 (r+r_0)^2} , \end{aligned} \quad (3.4)$$

where we employed the expansion $(1 + \delta)^{-1/2} \simeq 1 - \frac{\delta}{2} + \frac{3}{8} \delta^2$. The integral (3.1) now becomes

$$\phi(r) - \phi(\infty) \simeq A + B + C + D , \quad (3.5)$$

where

$$A = \int_{\infty}^r \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 - 1 \right]^{-1/2} \left[1 + \frac{G_N M}{r_0} + (3 - \varepsilon) \frac{G_N^2 M^2}{2 r_0^2} \right] dr , \quad (3.6)$$

$$B = \int_{\infty}^r \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 - 1 \right]^{-1/2} \frac{G_N M r_0}{r (r + r_0)} \left(1 + \frac{3 G_N M}{r_0} \right) dr , \quad (3.7)$$

$$C = - \int_{\infty}^r \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 - 1 \right]^{-1/2} \frac{\varepsilon G_N^2 M^2}{2 r^2} dr \quad (3.8)$$

and

$$D = \int_{\infty}^r \frac{1}{r} \left[\left(\frac{r}{r_0} \right)^2 - 1 \right]^{-1/2} \frac{3 G_N^2 M^2 r_0^2}{2 r^2 (r + r_0)^2} dr . \quad (3.9)$$

The integrals are all elementary, and we obtain

$$\begin{aligned} \phi(r) - \phi(\infty) \simeq & - \left[1 + \frac{G_N M}{r_0} + (3 - \varepsilon) \frac{G_N^2 M^2}{2 r_0^2} \right] \arcsin \left(\frac{r_0}{r} \right) \\ & + \frac{G_N M}{r_0} \left(1 + \frac{3 G_N M}{r_0} \right) \left[\sqrt{1 - \left(\frac{r_0}{r} \right)^2} \left(1 + \frac{r}{r + r_0} \right) + \arcsin \left(\frac{r_0}{r} \right) - 2 \right] \\ & - \frac{\varepsilon G_N^2 M^2}{2 r_0^2} \left[\frac{r_0}{2r} \sqrt{1 - \left(\frac{r_0}{r} \right)^2} - \frac{1}{2} \arcsin \left(\frac{r_0}{r} \right) \right] \\ & - \frac{3 G_N^2 M^2}{2 r_0^2} \left[\sqrt{1 - \left(\frac{r_0}{r} \right)^2} \left(2 - \frac{r_0}{2r} - \frac{r^2}{3 (r + r_0)^2} + \frac{11 r}{3 (r + r_0)} \right) + \frac{7}{2} \arcsin \left(\frac{r_0}{r} \right) - \frac{16}{3} \right] \end{aligned}$$

and finally

$$\Delta \phi \simeq \frac{2 R_H}{r_0} + \frac{R_H^2}{16 r_0^2} (15\pi - 16 - 3\pi \varepsilon) , \quad (3.10)$$

which shows that ours is indeed an expansion in R_H/r_0 . Notice also that the term of second order in R_H/r_0 does not vanish when $\varepsilon \rightarrow 0$, since we properly expanded the integrand to this order.

Our result can now be compared with the deflection angle of a light ray (or a photon) just grazing the Sun surface, which is usually given in the form

$$\Delta \phi = \frac{1}{2} (1 + \gamma) \frac{2 R_H}{r_0} , \quad (3.11)$$

where $r_0 = R_\odot$, and $R_H = 2 G_N M_\odot$. The best measurements presently available for the parameter γ from the light bending close to the surface of the Sun, are given by the the development of the very-long-baseline radio interferometry (VLBI, see Ref. [29]). A 2004 analysis of almost 2 million VLBI observations of 541 radio sources, made by 87 VLBI sites, yielded $\gamma - 1 \simeq (-1.7 \pm 4.5) \cdot 10^{-4}$ [30]. A 2009 analysis updated to through 2008 data yielded $\gamma - 1 \simeq (-1.6 \pm 1.5) \cdot 10^{-4}$ [31]. Good measurements were also performed by the optical astrometry satellite Hipparcos (at the level of 0.1 percent), and significative improvements are expected from the just launched satellite Gaia (see Ref. [32]). On comparing Eq. (3.10) with (3.11), we immediately get

$$\gamma - 1 = \frac{G_N M}{8 r_0} (15\pi - 16 - 3\pi\varepsilon) , \quad (3.12)$$

or

$$|\gamma - 1| = \frac{G_N M}{8 r_0} |15\pi - 16 - 3\pi\varepsilon| \lesssim 1.6 \cdot 10^{-4} . \quad (3.13)$$

For $M = M_\odot$ and $r_0 = R_\odot$, this means $-60.7 < \varepsilon < 67.3$, which, together with the mathematical constrain $\varepsilon \leq 1$, gives a range for the ε values of about

$$-65 \lesssim \varepsilon \leq 1 . \quad (3.14)$$

Assuming the "worst" situation, namely $\varepsilon \simeq -65$, then Eq. (2.33) gives the upper bound for the GUP parameter

$$|\beta| = \frac{M^2}{4 m_p^2} \frac{\pi^2 \varepsilon^2}{1 - \varepsilon} \lesssim 5.3 \cdot 10^{78} , \quad (3.15)$$

which is comparable with the limit obtained from general considerations on the validity of the low β expansion (2.25). This not-so-tight bound has to do with the well-known fact that light deflection is not the most precise test of GR (since light deflection is still a "Newtonian" phenomenon, see Ref. [33]). A much better estimation will be obtained in the next Section, where the perihelion precession is considered.

4 Perihelion precession by deformed Schwarzschild metric

Here, we consider a particle bound in a orbit around a massive body, typically a planet around the Sun. Again, we roughly follow the treatment of Ref. [28]. In Fig. 2 we can see the relevant geometrical parameters for an elliptic orbit in a polar coordinates system, with the radial coordinate r which at aphelia and perihelia takes, respectively, the maximum value r_+ and minimum value r_- ; e is the eccentricity, a the semi-major axis, and L the *semilatus rectum*. These geometrical parameters are related by

$$r_\pm = (1 \pm e) a \quad (4.1)$$

$$L = (1 - e^2) a \quad (4.2)$$

$$\frac{2}{L} = \frac{1}{r_+} + \frac{1}{r_-} . \quad (4.3)$$

The two relevant constants of motion of the system, E and J , can be interpreted respectively as the energy per unit mass, $(1 - E)/2$, and an angular momentum per unit mass (see Ref. [28]). The




Figure 2: Elements of the ellipse used in the text for the calculations of precession of planetary orbits.

constants E and J can be further expressed as functions of $F(r_-)$ and $F(r_+)$. The angle swept out by the position vector when it increases from r_- to r is then given by the integral

$$\phi(r) - \phi(r_-) = \int_{r_-}^r \left[\frac{r_-^2 \left(\frac{1}{F(r)} - \frac{1}{F(r_-)} \right) - r_+^2 \left(\frac{1}{F(r)} - \frac{1}{F(r_+)} \right)}{r_-^2 r_+^2 \left(\frac{1}{F(r_+)} - \frac{1}{F(r_-)} \right)} - \frac{1}{r^2} \right]^{-1/2} \frac{dr}{r^2 \sqrt{F(r)}} . \quad (4.4)$$

The total change in ϕ at every lap is just twice the change as r increases from r_- to r_+ . This would equal 2π if the orbit were a closed ellipse, so the total orbital precession in each revolution is given by

$$\Delta\phi = 2|\phi(r_+) - \phi(r_-)| - 2\pi . \quad (4.5)$$

As before, the exact value of the integral can be expressed via elliptic functions, but then one has to expand the elliptic functions to obtain useful results. It is much faster to expand before integrating, where, in analogy with the case of light deflection, the small parameter is now given by R_H/r_- . Since $R_H/r \lesssim R_H/r_-$ all along the orbit, this also implies that one can expand in R_H/r . From the definition of $F(r)$, to second order in R_H/r , we have

$$\frac{1}{F(r)} \simeq 1 + \frac{2G_N M}{r} + (4 - \varepsilon) \frac{G_N^2 M^2}{r^2} , \quad (4.6)$$

so the expression in the square brackets inside the integral in Eq. (4.4) is actually a quadratic function of $1/r$. Moreover, it vanishes for $r = r_{\pm}$, and we can therefore write

$$\frac{r_-^2 \left(\frac{1}{F(r)} - \frac{1}{F(r_-)} \right) - r_+^2 \left(\frac{1}{F(r)} - \frac{1}{F(r_+)} \right)}{r_-^2 r_+^2 \left(\frac{1}{F(r_+)} - \frac{1}{F(r_-)} \right)} - \frac{1}{r^2} = C \left(\frac{1}{r_-} - \frac{1}{r} \right) \left(\frac{1}{r} - \frac{1}{r_+} \right) , \quad (4.7)$$

so that the equation (4.4) for the trajectory becomes

$$\phi(r) - \phi(r_-) = \frac{1}{\sqrt{C}} \int_{r_-}^r \left[\left(\frac{1}{r_-} - \frac{1}{r} \right) \left(\frac{1}{r} - \frac{1}{r_+} \right) \right]^{-1/2} \frac{dr}{r^2 \sqrt{F(r)}} . \quad (4.8)$$

We can find the constant C by considering the limit $r \rightarrow \infty$, and hence $F(r)^{-1} \rightarrow 1$,

$$C = \frac{r_+^2 F(r_-) [F(r_+) - 1] - r_-^2 F(r_+) [F(r_-) - 1]}{r_- r_+ [F(r_-) - F(r_+)]} . \quad (4.9)$$

By inserting the expressions for $F(r_{\pm})$ from (2.26), and recalling the formula (4.3) for L , with a bit of algebra, we get the exact expression

$$C = \left(1 - \frac{2R_H}{L} + \varepsilon \frac{R_H^2}{L^2} - \varepsilon^2 \frac{R_H^3}{8aL^2}\right) \left(1 - \varepsilon \frac{R_H}{2L}\right)^{-1} \quad (4.10)$$

We can now expand to second order in R_H/L ,

$$C^{-1/2} \simeq 1 + \left(\frac{4-\varepsilon}{2}\right) \frac{G_N M}{L} + \left(6 - 3\varepsilon - \frac{1}{8}\varepsilon^2\right) \frac{G_N^2 M^2}{L^2} , \quad (4.11)$$

and, to second order in R_H/r ,

$$[F(r)]^{-1/2} \simeq 1 + \frac{G_N M}{r} + (3 - \varepsilon) \frac{G_N^2 M^2}{2r^2} . \quad (4.12)$$

The integral in Eq. (4.8) is largely simplified if we choose a clever change of variable. Let us consider the closed orbit predicted by Newtonian mechanics. Taking the pole in the direction of the *semilatus rectum*, the polar equation of the trajectory reads

$$\frac{1}{r} = \frac{1}{L} - \frac{e}{L} \sin \psi , \quad (4.13)$$

and this will be our change of variable in the integral (4.8), with

$$\frac{dr}{r^2} = \frac{e}{L} \cos \psi \, d\psi . \quad (4.14)$$

On recalling Eq. (4.3) and

$$\frac{e}{L} = \frac{1}{2} \left(\frac{1}{r_-} - \frac{1}{r_+} \right) , \quad (4.15)$$

we have

$$\frac{1}{r_-} - \frac{1}{r} = \frac{e}{L} (1 + \sin \psi) \quad (4.16)$$

$$\frac{1}{r} - \frac{1}{r_+} = \frac{e}{L} (1 - \sin \psi) , \quad (4.17)$$

and

$$\left(\frac{1}{r_-} - \frac{1}{r} \right) \left(\frac{1}{r} - \frac{1}{r_+} \right) = \frac{e^2}{L^2} \cos^2 \psi . \quad (4.18)$$

When $r = r_-$, then $\psi = -\pi/2$. Inserting everything back into Eq. (4.8), to second order in R_H/r , we get

$$\begin{aligned} \phi(r) - \phi(r_-) &\simeq \left[1 + \left(\frac{4-\varepsilon}{2} \right) \frac{G_N M}{L} + \left(6 - 3\varepsilon - \frac{1}{8}\varepsilon^2 \right) \frac{G_N^2 M^2}{L^2} \right] \\ &\times \int_{-\pi/2}^{\psi} \left[1 + \frac{G_N M}{L} (1 - e \sin \psi) + (3 - \varepsilon) \frac{G_N^2 M^2}{2L^2} (1 - e \sin \psi)^2 \right] d\psi . \end{aligned} \quad (4.19)$$

At the aphelion $r = r_+$ and $\psi = \pi/2$. The computation of the integral to the same order in R_H/r is elementary, and yields

$$\begin{aligned}\phi(r_+) - \phi(r_-) &\simeq \left[1 + \left(\frac{4-\varepsilon}{2} \right) \frac{G_N M}{L} + \left(6 - 3\varepsilon - \frac{1}{8} \varepsilon^2 \right) \frac{G_N^2 M^2}{L^2} \right] \\ &\times \left[1 + \frac{G_N M}{L} + (3-\varepsilon) \frac{G_N^2 M^2}{2 L^2} \left(1 + \frac{e^2}{2} \right) \right] \pi \\ &\simeq \pi \left[1 + \left(\frac{6-\varepsilon}{2} \right) \frac{G_N M}{L} + \frac{G_N^2 M^2}{L^2} N(\varepsilon, e) \right],\end{aligned}\quad (4.20)$$

where

$$N(\varepsilon, e) = \frac{1}{2} \left[19 - 8\varepsilon + (3-\varepsilon) \frac{e^2}{2} - \frac{\varepsilon^2}{4} \right]. \quad (4.21)$$

Finally the total precession after a single lap is given by

$$\Delta\phi \simeq 2\pi \left(\frac{6-\varepsilon}{2} \right) \frac{G_N M}{L} + 2\pi \frac{G_N^2 M^2}{L^2} N(\varepsilon, e). \quad (4.22)$$

In particular, we note that, to first order in R_H/L , we can write

$$\Delta\phi \simeq \frac{6\pi G_N M}{L} \left(1 - \frac{\varepsilon}{6} \right), \quad (4.23)$$

which, of course, reproduces the usual GR prediction in the limit $\varepsilon \rightarrow 0$. This relation should now be compared with known observational data.

4.1 Solar system data

The perihelion precession for Mercury is by far the best known and measured GR precession in the Solar system. Referring to Ref. [29] for the latest most accurate and comprehensive data, we can report the relation

$$\langle \dot{\omega} \rangle = \frac{6\pi G_N M}{L} \left[\frac{1}{3} (2 + 2\gamma - \bar{\beta}) + 3 \cdot 10^{-4} \frac{J_2}{10^{-7}} \right], \quad (4.24)$$

where $\langle \dot{\omega} \rangle$ is the measured perihelion shift, J_2 a dimensionless measure of the quadrupole moment of the Sun, and γ and $\bar{\beta}$ are the usual Eddington-Robertson expansion parameters. The latest data from helioseismology give $J_2 = (2.2 \pm 0.1) \cdot 10^{-7}$. The measured perihelion shift of Mercury is known very accurately: after the perturbing effects of other planets have been accounted for, the excess shift is known to about 0.1% from radar observations of Mercury between 1966 and 1990 [34]. The solar oblateness effect due to the quadrupole moment is then smaller than the observational error, so it can be neglected. Substituting standard orbital elements and physical constants for Mercury and the Sun, we obtain

$$\langle \dot{\omega} \rangle = \left(1 + \frac{2\gamma - \bar{\beta} - 1}{3} \right) 42.98''/\text{century}, \quad (4.25)$$

where we can place a bound of $|2\gamma - \bar{\beta} - 1| \lesssim 3 \cdot 10^{-3}$.

Parameter	Value
e (Eccentricity)	0.6171334(5)
P_b (days) (Orbital period)	0.322997448911(4)
$\langle\dot{\omega}\rangle$ (deg/year) (Periastron shift)	4.226598(5)
γ (s) (Time dilation-gravitational redshift)	$4.2992(8) \times 10^{-3}$
\dot{P}_b (s/s) (Orbital period decay)	$-2.423(1) \times 10^{-12}$

Table 1: Orbital parameters of PRS B 1913+16. Figures in parentheses represent estimated uncertainties in the last quoted digit. Data from Ref.[36].

Comparing with $\Delta\phi$ from Eq. (4.23), we get

$$|\varepsilon| \lesssim 6 \cdot 10^{-3} , \quad (4.26)$$

which, replaced in Eq. (2.33), yields the lower bound

$$|\beta| = \frac{M^2}{4m_p^2} \frac{\pi^2 \varepsilon^2}{1 - \varepsilon} \lesssim 3 \cdot 10^{72} . \quad (4.27)$$

We can also consider the most recent data from the Messenger spacecraft [35], which orbited Mercury in 2011-2013, and improved very much the knowledge of its orbit. We can actually push this bound even lower, to $|2\gamma - \bar{\beta} - 1| \lesssim 7.8 \cdot 10^{-5}$, although the knowledge of J_2 would have to improve simultaneously. If just the error in $|2\gamma - \bar{\beta} - 1|$ were taken into account, this would imply

$$|\varepsilon| = 2 |2\gamma - \bar{\beta} - 1| \lesssim 1.56 \cdot 10^{-4} \quad (4.28)$$

and therefore

$$|\beta| \lesssim 2 \cdot 10^{69} . \quad (4.29)$$

But of course this limit should not be considered completely reliable in this contest, since the less accurate bound on J_2 cannot be brutally neglected, at least in principle. As expected, we gain here at least six orders of magnitude, showing once again that the perihelion shift is one of the most precise tests of GR, a true GR effect not present at all in Newtonian gravity (as it is well known). We can however try to put this limit on a firmer ground by looking for even larger effects of this kind. Where? Of course, in Binary Pulsars!

4.2 Pulsar PRS B 1913+16 data

Clearly, binary pulsars are very good candidates for measurements of periastron shifts. Among the known pulsar systems, the best tested pair is the Pulsar PRS B 1913+16. Discovered in 1974 by Hulse and Taylor, this system has become, after 40 years of observations, one of the most reliable celestial laboratories for precise GR measurements. For example, prediction of GR for the period decay rate due to emission of gravitational waves coincides with the measured value up to an error on the 14th decimal figure.

The state of the art on this system is described in Ref. [36]. In Table 1, we report the orbital parameters of interest for us. The parameters e and P_b are called Keplerian parameters, since

they are well defined quantities also in the Newtonian theory. On the contrary, $\langle\dot{\omega}\rangle$, γ , \dot{P}_b are known as Damour-Deruelle post-Keplerian parameters [37], quantities typically well defined in GR only. In Ref. [38], Taylor and Weisberg have shown that each GR post-Keplerian parameter can be expressed in terms of the Keplerian parameters and of the unknown masses of the pulsar and its companion, m_1 , m_2 . In fact,

$$\begin{aligned}\langle\dot{\omega}\rangle &= 3 G_N^{2/3} c^{-2} (P_b/2\pi)^{-5/3} (1-e^2)^{-1} (m_1+m_2)^{2/3} \\ &= 2.113323(2) \left[\frac{(m_1+m_2)}{M_\odot} \right]^{2/3} \text{ deg yr}^{-1}\end{aligned}\quad (4.30)$$

$$\begin{aligned}\gamma &= G_N^{2/3} c^{-2} e (P_b/2\pi)^{1/3} m_2 (m_1+2m_2)(m_1+m_2)^{-4/3} \\ &= 0.002936679(2) \left[\frac{m_2 (m_1+2m_2)(m_1+m_2)^{-4/3}}{M_\odot^{2/3}} \right] \text{ s}\end{aligned}\quad (4.31)$$

$$\begin{aligned}\dot{P}_b &= -\frac{192 \pi G_N^{5/3}}{5c^5} \left(\frac{P_b}{2\pi} \right)^{-5/3} \left(1 + \frac{73}{24} e^2 + \frac{37}{96} e^4 \right) (1-e^2)^{-7/2} m_1 m_2 (m_1+m_2)^{-1/3} \\ &= -1.699451(8) \cdot 10^{-12} \left[\frac{m_1 m_2 (m_1+m_2)^{-1/3}}{M_\odot^{5/3}} \right].\end{aligned}\quad (4.32)$$

To compute the numerical coefficients, in the second line of each equation we have substituted values for P_b and e from Table 1, and used the constants $G_N M_\odot/c^3 = 4.925490947 \cdot 10^{-6} \text{ s}$ and $1 \text{ Julian year} = 86400 \cdot 365.25 \text{ s}$. The figures in parentheses represent uncertainties in the last quoted digit, determined by propagating the uncertainties listed in Table 1. In each case, the errors are dominated by the experimental uncertainty in orbital eccentricity, e . Furthermore, we note that the analytical expression for the periastron angular velocity $\langle\dot{\omega}\rangle$ given above by Taylor and Weisberg, can be obtained from the standard GR prediction for the shift-per-lap $6\pi G_N M/L$, simply by dividing this for the period P_b , and of course with $M = m_1 + m_2$ (other quantities being expressed as function of P_b and e).

To get $\langle\dot{\omega}\rangle^{\text{GR}}$, the GR theoretical prediction of the periastron shift, the strategy is the following: first, insert the observational values for γ and \dot{P}_b from Table 1 into Eqs. (4.31) and (4.32), and solve for m_1 , m_2 . Then, substitute the values of m_1 , m_2 thus found into Eq. (4.30) to compute $\langle\dot{\omega}\rangle^{\text{GR}}$, and compare this prediction with the observed value of $\langle\dot{\omega}\rangle^{\text{Obs}}$ again given in Table 1. The relative error can then be defined as

$$\tilde{\varepsilon} = \frac{\langle\dot{\omega}\rangle^{\text{Obs}} - \langle\dot{\omega}\rangle^{\text{GR}}}{\langle\dot{\omega}\rangle^{\text{GR}}}, \quad (4.33)$$

that is $\langle\dot{\omega}\rangle^{\text{GR}}(1 + \tilde{\varepsilon}) = \langle\dot{\omega}\rangle^{\text{Obs}}$, and, on comparing with $\Delta\phi$ in Eq. (4.23), we get $|\varepsilon| = 6|\tilde{\varepsilon}|$.

There is one further issue we should care of: measurements are now so precise that the observed value of \dot{P}_b in Table 1 should be corrected for the relative acceleration between the pulsar reference frame and the solar system centre-of-mass frame (see Ref. [39]). Such relative acceleration is mainly due to the fact that the pulsar and our solar system are located in different arms of our Galaxy, at different distances from the galactic centre. The small additional kinematic contribution to the observed \dot{P}_b is

$$\Delta\dot{P}_{b,gal} = (-0.027 \pm 0.005) \cdot 10^{-12} \quad (4.34)$$

We can solve the system of the two equations (4.31) and (4.32) for the two unknowns m_1 and m_2 . The best result is obtained by combining, from Table 1, the lower bound for $\dot{P}_b = (-2.423 - 0.001) \cdot 10^{-12}$, and the upper bound of the correction term $\Delta\dot{P}_{b,gal} = (-0.027 + 0.005) \cdot 10^{-12}$, in order to compute the value $(\dot{P}_b - \Delta\dot{P}_{b,gal})$ to be inserted in the LHS of Eq. (4.32). We then get

$$\tilde{\varepsilon} = 8.9 \cdot 10^{-5} , \quad (4.35)$$

which means $|\varepsilon| \simeq 5.4 \cdot 10^{-4}$. Recalling now that $M = m_1 + m_2 = 2.828 \cdot M_\odot$, this translates into the bound

$$|\beta| \lesssim 2 \cdot 10^{71} , \quad (4.36)$$

which is tighter than the bound (4.27) coming from "standard" Mercury observations, but weaker than the "Messenger bound" of the previous section. However, note that we do not have the caveat of the error bounds on J_2 here.

Finally, once again, it can be easily checked that the expansion (4.20) is fully convergent also when the data of the pulsar PSR B 1913+16 are inserted, as well as it is convergent in the solar system field.

5 Conclusions

We have shown that a suitable deformation of the Schwarzschild metric can reproduce the Hawking temperature for a black hole, when this is computed from a Generalized Uncertainty Principle. We found in this way an analytic relation between the deformation parameter of the metric, ε , and the usual GUP deformation parameter β . In particular, when $\beta \rightarrow 0$, we correctly recover GR, and standard quantum mechanics. Neither the geodesic equation, nor the equivalence principle are violated, for any value of β or ε .

Well-known astronomical measurements, in the Solar system as well as in binary pulsar systems, allowed us to put constraints on the parameter β . This direction seems to point towards promising research: at present we just deformed the Schwarzschild solution, but a future possibility is to deform the full field equations of GR, in order to get, among other things, a more stringent bound on the GUP parameter β . We would like to conclude by emphasizing that, although in the existing literature one can find bounds on β much tighter than those obtained in this paper, they seem to depend, at least partially, either on a specific representation of the deformed commutator, or on a deformation of Poisson brackets implying a violation of the equivalence principle. The line of reasoning followed in the present paper avoids these possible difficulties.

A Modified Poisson brackets and equivalence principle

We shall show here that a deformation of the classical Newtonian (i.e. non covariant) Poisson brackets implies a violation of the equivalence principle. In Refs. [14, 15] as well as [12, 13], Poisson brackets are deformed in the same fashion as the quantum commutators derived from a GUP.

Now, considering just a one dimensional system, to keep the calculation simple, we can write the Poisson brackets for one pair of canonical variables as

$$\{q, p\} = 1 + \beta p^2 , \quad \{q, q\} = \{p, p\} = 0 . \quad (\text{A.1})$$

It is then easy to show that for any regular function $H(q, p)$, the following hold

$$\{q, H\} = (1 + \beta p^2) \frac{\partial H}{\partial p} , \quad \{p, H\} = -(1 + \beta p^2) \frac{\partial H}{\partial q} . \quad (\text{A.2})$$

A point-like particle of mass m moving in a Newtonian potential is described by the Hamiltonian (we assume $M \gg m$)

$$H = \frac{p^2}{2m} - \frac{G_N M m}{q} , \quad (\text{A.3})$$

and therefore evolves according to the equations of motion

$$\dot{q} = \{q, H\} = (1 + \beta p^2) \frac{p}{m} \quad (\text{A.4})$$

$$\dot{p} = \{p, H\} = -(1 + \beta p^2) \frac{G_N M m}{q} .$$

From the first equation, we get

$$m \dot{q} = p + \beta p^3 , \quad (\text{A.5})$$

which implies

$$m \ddot{q} = (1 + 3\beta p^2) \dot{p} , \quad (\text{A.6})$$

and, using now the second of Eqs. (A.4), we have, to first order in β ,

$$\ddot{q} \simeq -(1 + 4\beta p^2) \frac{G_N M}{q^2} . \quad (\text{A.7})$$

Eq. (A.5) can be solved for p to the first order in β , yielding

$$p \simeq m \dot{q} - \frac{\beta (m \dot{q})^3}{1 + 3\beta (m \dot{q})^2} . \quad (\text{A.8})$$

Finally, to first order in β , we have the equation of motion

$$\ddot{q} \simeq -(1 + 4\beta (m \dot{q})^2) \frac{G_N M}{q^2} . \quad (\text{A.9})$$

Clearly, the trajectory of a test particle of mass m will depend on m , which signals a violation of the Equivalence Principle. This violation has nothing to do with GR, or the geodesic equation, or the covariant formalism, but strictly followed from the modified Poisson brackets (A.1).

B Non-relativistic analogue

We display here a non-relativistic analogue of the deformation procedure followed in the main text. For a given metric like the one in Eq. (2.1) we know that, in a situation of weak, stationary field, and for slowly moving particles, we can define the effective (Newtonian) potential (see, e.g., [28])

$$V(r) = \frac{1}{2} [F(r) - 1] , \quad (\text{B.1})$$

and the Hamiltonian of a particle of mass m , moving in this potential, can be written as

$$H = \frac{p^2}{2m} + m V(q) . \quad (\text{B.2})$$

We assume undeformed, standard Poisson brackets

$$\{q, p\} = 1 , \quad \{q, q\} = \{p, p\} = 0 , \quad (\text{B.3})$$

therefore the equations of motion are the usual ones,

$$\begin{aligned} \dot{q} &= \{q, H\} = \frac{\partial H}{\partial p} = \frac{p}{m} \\ \dot{p} &= \{p, H\} = -\frac{\partial H}{\partial q} = -m \frac{\partial V}{\partial q} , \end{aligned} \quad (\text{B.4})$$

which yield the equation of motion for q

$$\ddot{q} = -\frac{\partial V}{\partial q} . \quad (\text{B.5})$$

The above clearly preserves the equivalence principle for any kind of potential $V(q)$. In particular, by choosing a deformed metric as in Eq. (2.26), we have

$$\ddot{q} = -\frac{G_N M}{q^2} + \varepsilon \frac{G_N^2 M^2}{q^3} , \quad (\text{B.6})$$

which does not depend on m and the equivalence principle is not violated.

References

- [1] H.S. Snyder, Phys.Rev. **71**, 38 (1947).
C.N. Yang, Phys. Rev. **72**, 874 (1947).
F. Karolyhazy, Nuovo Cim. A **42**, 390 (1966).
- [2] D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. B **197**, 81 (1987).
D.J. Gross, P.F. Mende, Phys. Lett. B **197**, 129 (1987).
D. Amati, M. Ciafaloni, G. Veneziano, Phys. Lett. B **216**, 41 (1989).
K. Konishi, G. Paffuti, P. Provero, Phys. Lett. B **234**, 276 (1990).
G. Veneziano, Proceedings of PASCOS 90, *Quantum Gravity near the Planck scale*, Boston 1990, p.486 (unpublished).
- [3] M. Maggiore, Phys. Lett. B **304**, 65 (1993).
- [4] A. Kempf, G. Mangano, R.B. Mann, Phys. Rev. D **52**, 1108 (1995).
M. Bojowald, A. Kempf, Phys. Rev. D **86**, 085017 (2012) [arXiv:1112.0994].
- [5] F. Scardigli, Phys. Lett. B **452**, 39 (1999).
- [6] R.J. Adler, D.I. Santiago, Mod. Phys. Lett. **A14**, 1371 (1999).

- [7] F. Scardigli and R. Casadio, *Class. Quantum Grav.* **20**, 3915 (2003).
- [8] C. Brukner et al., *Nature Physics* **8** 393 (2012).
F. Marin, M. Cerdonio, et al., *Nature Physics* **9**, 71 (2013).
- [9] F. Brau, *J. Phys. A* **32**, 7691 (1999).
- [10] S. Das, E. Vagenas, *Phys. Rev. Lett.* **101**, 221301 (2008).
A.F. Ali, S. Das, E. Vagenas, *Phys. Rev. D* **84**, 044013 (2011).
- [11] P. Pedram, K. Nozari, and S.H. Taheri, *JHEP* 1103, 093 (2011).
- [12] L.N. Chang, D. Minic, N. Okamura, and T. Takeuchi, *Phys. Rev. D* **66**, 026003 (2002).
- [13] K. Nozari, S. Akhshabi, *Chaos Solitons Fractals* **37** 324 (2008).
P. Pedram, *Phys. Lett. B* **718** 638 (2012).
- [14] S. Pramanik, S. Ghosh, *Int. J. Mod. Phys. A* **28** 27, 1350131 (2013).
S. Ghosh, *Class. Quant. Grav.* **31** 025025 (2014).
- [15] S. Pramanik, *Implication of Geodesic Equation in Generalized Uncertainty Principle framework*, arXiv:1404.2567 [hep-th].
- [16] M. Walker, *J. Math. Phys. (N.Y.)* **11**, 2280 (1970).
- [17] A. Zee, *Quantum Field Theory in a Nutshell*, Princeton University Press (2003).
- [18] L. J. Garay, *Int. J. Mod. Phys. A* **10**, 145 (1995).
- [19] W. Heisenberg, *Zeitschrift für Physik*, **43**, 172 (1927).
- [20] F. Scardigli, *Nuovo Cim. B* **110**, 1029 (1995).
- [21] R.J. Adler, P. Chen and D.I. Santiago, *Gen. Rel. Grav.* **33** 2101 (2001).
- [22] M. Cavaglia and S. Das, *Class. Quant. Grav.* **21**, 4511 (2004).
- [23] M. Cavaglia, S. Das and R. Maartens, *Class. Quant. Grav.* **20**, L205 (2003).
- [24] L. Susskind, J. Lindesay, *An Introduction to Black Holes, Information, and the String Theory Revolution* (World Scientific, Singapore, 2005). See chapter 10.
- [25] K. Nouicer, *Class. Quant. Grav.* **24**, 5917 (2007).
- [26] F. Scardigli, *Glimpses on the micro black hole planck phase*, [arXiv:0809.1832].
- [27] P. Jizba, H. Kleinert, and F. Scardigli, *Phys. Rev. D* **81**, 084030 (2010).
- [28] S. Weinberg, *Gravitation and Cosmology*, John Wiley & Sons, New York (1972).
- [29] C.M. Will, *The Confrontation between General Relativity and Experiment*, *Living Rev. Relativity*, **9**, (2006), 3. Update in arXiv:1403.7377v1 [Online Article: <http://www.livingreviews.org/lrr-2006-3>]

- [30] S.S. Shapiro, J.L. Davis, D.E. Lebach, and J.S. Gregory, *Measurement of the solar gravitational deflection of radio waves using geodetic very-long-baseline interferometry data, 1979 1999*, Phys. Rev. Lett., **92** 121101, (2004).
- [31] S.B. Lambert, C. Le Poncin-Lafitte, *Determining the relativistic parameter γ using very long baseline interferometry*, Astron. Astrophys., 499, 331 (2009). [arXiv:0903.1615].
- [32] <http://en.wikipedia.org/wiki/Hipparcos>
<http://sci.esa.int/gaia/>
- [33] See the paper of Johann Georg von Soldner (1801) translated and reproduced in S. Jaki, Found. Phys. **8**, Issue 11-12, 927 (1978).
- [34] I.I. Shapiro, *Solar system tests of general relativity: Recent results and present plans*, in N.Ashby, D.F.Bartlett, and W.Wyss, eds., General Relativity and Gravitation, Proceedings of the 12th International Conference on General Relativity and Gravitation, University of Colorado at Boulder, July 28, 1989, 313330, (Cambridge University Press, Cambridge, U.K., New York, U.S.A., 1990).
- [35] A.K. Verma, A. Fienga, J. Laskar, H. Manche, and M. Gastineau, *Use of MESSENGER radio-science data to improve planetary ephemeris and to test general relativity*, Astron. Astrophys., **561**, A115 (2014).
- [36] J.M. Weisberg, D.J. Nice, J.H. Taylor, Astrophysical Journal, **722**, 1030 (2010). [arXiv:1011.0718]
- [37] T. Damour, N. Deruelle, Ann. Inst. H. Poincare (Phys. Theorique), **44**, 263 (1986).
- [38] J.H. Taylor and J.M. Weisberg, Astrophysical Journal, **345**, 434 (1989).
- [39] T. Damour and J.H. Taylor, Astrophysical Journal, **366**, 501 (1991).