Yet another proof of Hawking and Ellis's Lemma 8.5.5

S. Krasnikov*

Abstract

The fact that the null generators of a future Cauchy horizon are past complete was first proved by Hawking and Ellis [1]. Then Budzyński, Kondracki, and Królak outlined a proof free from the error found in the original one [2]. Finally, Minguzzi has recently published his version of the proof [3] patching a previously unnoticed hole in the preceding two. I am not aware of any flaws in that last proof, but it is quite difficult. In this note I present a simpler one.

1 Introduction

Let H^+ denote a future Cauchy horizon. A lemma by Hawking and Ellis says

Lemma 8.5.5 of [1]. If $H^+(\mathcal{Q})$ is compact for a partial Cauchy surface \mathcal{Q} , then the null geodesic generating segments of $H^+(\mathcal{Q})$ are geodesically complete in the past direction.

The lemma itself has never been doubted (to my knowledge), but the proof offered in [1] was found to be flawed, see [3] and references therein. To improve this situation — which is important, because the lemma is a popular tool in mathematical relativity — Minguzzi has recently published a new, more accurate, proof of the lemma (or, to be precise, of some strengthening of it).

In this comment I present yet another proof of the same fact. The reason for doing this is that my version is much simpler (partly because its major part is replaced by a reference to a lemma proved elsewhere).

^{*}Email: S.V.Krasnikov@mail.ru

2 The proposition and its proof

In a spacetime M consider a past inextendible null curve γ totally imprisoned in a compact set K. Pick a smooth unit timelike future directed vector field τ on M and define (uniquely up to an additive constant) the "arc length parameter" l on γ by the requirement

$$g(\partial_l, \boldsymbol{\tau}) = -1. \tag{1}$$

In addition to l define on γ an affine parameter s so that ∂_s is future directed and s=0 at l=0. Now γ is characterized by the (evidently negative) function

$$h \equiv g(\partial_s, \boldsymbol{\tau}),$$

which relates l to s:

$$h = -\frac{\mathrm{d}l}{\mathrm{d}s}, \qquad s(l) = \int_{l}^{0} \frac{\mathrm{d}\check{l}}{h(\check{l})}.$$
 (2)

As is proven in [4]

$$h'/h$$
 is bounded on γ . (3)

Further, a few minor changes — a past inextendible $\gamma(l)$ with $l \in (-\infty, 0]$ instead of the future inextendible $\gamma(l)$ with $l \in [0, \infty)$ and an arbitrary compact \mathcal{K} instead of some specific \mathcal{L} — leave [4, Lemma 8] valid while bringing it to the following form.

Lemma 8 of [4]. Assume h(l) is such that there exists a smooth function f(l) defined at non-positive l and obeying for some non-negative constants $c_1, \underline{f}, \overline{f}$ the inequalities

$$\underline{f} \le f \le \overline{f}, \qquad |f'/f| < \infty$$

and

$$h'/h < -f'/f - c_1 f, \qquad \forall l \le 0. \tag{4}$$

Then there is a timelike past inextendible curve γ_{\varkappa_0} which is obtained by moving each point of γ to the past along the integral curves of τ and which is totally imprisoned in a compact set \mathcal{O} .

Proposition. If K is a subset of the boundary of a globally hyperbolic past set M^{in} , then γ is past complete.

Proof. Suppose the lemma is false and γ is past incomplete. This would mean that the affine parameter s is bounded from below and, correspondingly, the integral (2) converges at $l \to -\infty$. Which allows one to define the following smooth positive function on $(-\infty, 0]$

$$f(l) \equiv \frac{1}{h} \left[-\int_{l}^{0} \frac{d\tilde{l}}{h(\tilde{l})} + 2 \int_{-\infty}^{0} \frac{d\tilde{l}}{h(\tilde{l})} \right]^{-1}.$$
 (5)

f so defined satisfies the equation

$$f'/f + h'/h = -f \tag{6}$$

and consequently, condition (4) holds.

As h is negative, the boundedness of the integral (2) provides a simple estimate

$$\infty > -\int_{-\infty}^{0} \frac{\mathrm{d}\check{l}}{h(\check{l})} > -\frac{1/h(l)}{(|h'/h|)^{\max}}, \qquad \forall l \in (-\infty, 0],$$

which implies, due to (3), that 1/h is bounded. It follows then from (5) that f is bounded too. Finally, the just proven boundedness of f combined with (6) and (3) implies the boundedness of f'/f. Thus all the conditions of Lemma 2 are fulfilled and the corresponding variation transforms γ into a past inextendible timelike curve γ_{\varkappa_0} . The latter being timelike lies entirely in the closed (due to the globally hyperbolicity of M^{in} , to which $\gamma_{\varkappa_0}(0)$ belongs) set $J^-(\gamma_{\varkappa_0}(0)) \subset M^{\text{in}}$ (the inclusion follows from the fact that M^{in} is a past set). Thus, γ_{\varkappa_0} is totally imprisoned in the compact subset $\mathcal{O} \cap J^-(\gamma_{\varkappa_0}(0))$ of the globally hyperbolic spacetime M^{in} , which is forbidden by [1, proposition 6.4.7].

References

- [1] S. W. Hawking and G. F. R. Ellis *The Large scale structure of spacetime* (Cambridge, Cambridge University Press, 1973).
- [2] R. Budzyński, W. Kondracki, A. Krolak New properties of Cauchy and event horizons, arXiv:gr-qc/0011033.

- [3] E. Minguzzi, J. Math. Phys. **55**, 082503 (2014).
- [4] S. Krasnikov, Physical Review D **90**, 024067 (2014).