Waveguides with asymptotically diverging twisting

David Krejčiřík

Department of Theoretical Physics, Nuclear Physics Institute ASCR, 25068 Řež, Czech Republic; krejcirik@ujf.cas.cz

15 July 2014

Abstract

We provide a class of unbounded three-dimensional domains of infinite volume for which the spectrum of the associated Dirichlet Laplacian is purely discrete. The construction is based on considering tubes with asymptotically diverging twisting angle.

1 Introduction

Given an open set Ω in \mathbb{R}^d , with $d \geq 1$, we denote by $-\Delta_D^{\Omega}$ the Dirichlet Laplacian in $L^2(\Omega)$. The significant features of Ω as regards the spectrum of $-\Delta_D^{\Omega}$ are conveniently described by means of the classification introduced by Glazman [8, Sec. 49] (see also [3, Sec. X.6.1]): Ω is *quasi-conical* if it contains arbitrarily large balls; Ω is *quasi-cylindrical* if it is not quasi-conical but contains an infinite set of identical disjoint balls; Ω is *quasi-bounded* if it is neither quasiconical nor quasi-cylindrical.

By choosing a singular sequence of functions supported in the enlarging balls, it is easy to see (cf. [3, Thm. X.6.5]) that the spectrum of $-\Delta_D^{\Omega}$ is purely essential whenever Ω is quasi-conical, namely, $\sigma(-\Delta_D^{\Omega}) = \sigma_{ess}(-\Delta_D^{\Omega}) = [0, \infty)$. The other extreme case is given by quasi-bounded Ω , for which the spectrum is "typically" purely discrete, *i.e.* $\sigma_{ess}(-\Delta_D^{\Omega}) = \emptyset$, at least under some regularity assumptions about Ω (precise conditions can be stated in terms of capacity, cf. [3, Thm. VIII.3.1]). These regularity assumptions are automatically satisfied if Ω is bounded or, more generally, of finite volume. In general, it is well known that the discreteness of the spectrum of $-\Delta_D^{\Omega}$ is equivalent to the compactness of the resolvent, which in turn is equivalent to the compactness of the embedding

$$W_0^{1,2}(\Omega) \hookrightarrow L^2(\Omega)$$
. (1)

The spectral picture is most complicated for quasi-cylindrical domains, where both essential and discrete spectra might be present. It is not difficult to see that $\sigma_{\rm ess}(-\Delta_D^{\Omega}) \neq \emptyset$ whenever Ω is quasi-cylindrical, while the existence of discrete eigenvalues is considered as a non-trivial property here.

A distinguished class of quasi-cylindrical domains is represented by unbounded *tubes*, *i.e.* tubular neighbourhoods of unbounded submanifolds in \mathbb{R}^d . The motivation for this restriction is at least two-fold. First, the tubular geometry is rich enough to demonstrate the mathematical complexity of the class of quasi-cylindrical domains. Second, the Dirichlet Laplacian in tubes is a reasonable model for the Hamiltonian in quantum-waveguide nanostructures, where the unboundedness is required to have non-trivial transport properties. The latter has lead to an enormous number of research papers in recent years, in which the interplay between the geometry of a tube and the spectrum of the associated Dirichlet Laplacian is investigated in various settings; we especially refer to the survey articles [2, 11, 10, 14, 9] with many references. As the most recent result, let us point out the repulsive effect of *twisting* in three-dimensional tubes about a spatial curve, which can be mathematically formalised in terms of Hardy-type inequalities [4, 10, 13, 1].

The objective of the present note is to show that, apart from expectable quasi-cylindrical settings, the tubular geometry gives rise to an interesting class of quasi-bounded domains with purely discrete spectrum, too. To the best of our knowledge, this class of domains has not been considered yet in connection with the compactness property of the Sobolev embedding (1). Our construction is based on considering tubes with a twisting angle diverging at the infinity of \mathbb{R}^d . For simplicity, we restrict to d = 3 in this note.

We would like to emphasise that the waveguides considered in this note have a *uniform cross-section*. This makes the model very different from the recent study [5], where the cross-section is allowed to vary along the reference line and shrinks to a point at the infinity of the waveguide. The latter leads to cusp-type domains for which compactness of (1) follows easily by an equivalent characterisation of quasi-boundedness

$$\lim_{\substack{x \in \Omega \\ |x| \to \infty}} \operatorname{dist}(x, \partial \Omega) = 0.$$
(2)

In the present model the validity of (2) is not that evident.

2 Twisted waveguides

A (non-bent) twisted tube Ω is obtained by translating and rotating a bounded open connected set $\omega \subset \mathbb{R}^2$ about a straight line in \mathbb{R}^3 . More precisely, following [10], Ω is defined as the image of a straight tube $\Omega_0 := \mathbb{R} \times \omega$ via the mapping

$$\mathcal{L}(x) := \left(x_1, x_2 \cos \theta(x_1) + x_3 \sin \theta(x_1), -x_2 \sin \theta(x_1) + x_3 \cos \theta(x_1)\right), \quad (3)$$

i.e. $\Omega := \mathcal{L}(\Omega_0)$. Here $\theta : \mathbb{R} \to \mathbb{R}$ is the rotation angle, for which we merely assume

$$\theta \in W^{1,\infty}_{\text{loc}}(\mathbb{R})\,. \tag{4}$$

In this note, we are interested in spectral properties of the Dirichlet Laplacian $-\Delta_D^{\Omega}$ in $L^2(\Omega)$ in the situation when the rotation angle explodes at infinity, *i.e.*,

$$\lim_{|x_1| \to \infty} |\dot{\theta}(x_1)| = +\infty.$$
(5)

This situation is briefly discussed (without any proofs) in [12, Sec. 6.2].

Let us now give some intuition for the results established below.

1. If the tube Ω is untwisted (*i.e.* either θ is constant or ω is rotationally invariant with respect to the origin in \mathbb{R}^2), then Ω can be identified with the straight tube Ω_0 and the spectrum is easily found by a separation of variables,

$$\sigma(-\Delta_D^{\Omega_0}) = [E_1, \infty), \qquad (6)$$

where E_1 denotes the lowest eigenvalue of $-\Delta_D^{\omega}$ in $L^2(\omega)$.

2. If Ω is twisted but $\dot{\theta}$ vanishes at infinity, then the spectrum (6) is not changed [10, Thm. 4.1 & Sec. 6.1]. However, there is a fine difference reflected in the fact that $-\Delta_D^{\Omega}$ satisfies a Hardy-type inequality [4, 10, 13]

$$-\Delta_D^{\Omega_0} - E_1 \ge \rho \,, \tag{7}$$

where $\rho: \Omega \to [0, \infty)$ is a non-trivial function.

- 3. This Hardy inequality turns into a Poincaré-type inequality (*i.e.* ρ bounded from below by a positive constant) if $|\dot{\theta}|$ is bounded from below by a positive constant [10, Sec. 6.1]. Consequently, the spectrum of $-\Delta_D^{\Omega}$ starts strictly above E_1 in this case.
- 4. For instance, if Ω is constantly twisted, *i.e.* $\dot{\theta}(x_1) = \beta$, then the spectral problem can be solved by a Floquet-type decomposition and it is shown in [6] that

$$\sigma(-\Delta_D^{\Omega}) = [\lambda_1, \infty), \qquad (8)$$

where λ_1 is the lowest eigenvalue of $-\Delta_D^{\omega} - \beta^2 \partial_{\tau}^2$ in $L^2(\omega)$, with $\partial_{\tau} := x_3 \partial_2 - x_2 \partial_3$ being the transverse angular derivative. Note that $\lambda_1 > E_1$ if, and only if, $\beta \neq 0$ and ω is not rotationally invariant with respect to the origin in \mathbb{R}^2 .

Our objective is to show that the repulsive effect in the situation (5) is so strong that the threshold of the essential spectrum may even diverge to infinity, so that the spectrum of $-\Delta_D^{\Omega}$ becomes purely discrete.

3 Quasi-cylindrical realisation

First, however, let us show that the condition (5) by itself is not sufficient to guarantee the absence of essential spectrum.

Theorem 1. Let $0 \in \omega$. Then

$$\sigma(-\Delta_D^\Omega) \supset [\mu_1, \infty) \,,$$

where μ_1 denotes the first eigenvalue of the Dirichlet Laplacian on the disc D_r of radius $r := \text{dist}(0, \partial \omega)$.

Proof. By the definition of Ω , the cylinder $\mathbb{R} \times D_r$ clearly lies inside Ω . The spectrum of the Dirichlet Laplacian in the cylinder coincides with the interval $[\mu_1, \infty)$. Applying the Weyl criterion for $-\Delta_D^{\Omega}$ with a singular sequence for the cylinder, we end up with the claim.

Since $\mathbb{R} \times D_r \subset \Omega \subset \mathbb{R} \times D_R$ whenever $0 \in \omega$, where $R := \sup_{t \in \omega} |t|$, it follows that Ω is quasi-cylindrical in this case. The geometrical setting of Theorem 1 is illustrated by Figure 1.



Figure 1: Quasi-cylindrical realisation: Waveguide of elliptical cross-section with a centre of rotation in the interior of the ellipse. The embedded cylindrical channel is responsible for the existence of essential spectrum.

4 Quasi-bounded realisation

Now we give a sufficient condition which guarantees the absence of essential spectrum.

Theorem 2. Let $\omega \subset \{(t_1, t_2) \in \mathbb{R}^2 \mid t_1 > 0\}$ and assume (4) and (5). Then

$$\sigma(-\Delta_D^{\Omega}) = \sigma_{\rm disc}(-\Delta_D^{\Omega})\,.$$

Proof. It follows from (5) that for any $n \in \mathbb{N}$ there exists $s_n \in (0, \infty)$ such that $|\dot{\theta}(x_1)| > n$ for every $|x_1| > s_n$. Fix $n \in \mathbb{N}$ and define $I_{\text{int}} := (-s_n, s_n)$, $I_{\text{ext}} := (-\infty, -s_n) \cup (s_n, +\infty)$. We also define $\Omega_{\text{int}} := \mathcal{L}(I_{\text{int}} \times \omega)$ and $\Omega_{\text{ext}} := \mathcal{L}(I_{\text{ext}} \times \omega)$. Imposing a supplementary Neumann condition at the cuts $\{\pm s_n\} \times \omega$, we obtain a lower bound

$$-\Delta_D^{\Omega} \ge -\Delta_{DN}^{\Omega_{\mathrm{int}}} \oplus -\Delta_{DN}^{\Omega_{\mathrm{ext}}}$$
 .

Here $-\Delta_{DN}^{\Omega_{\text{int}}}$ denotes the operator in $L^2(\Omega_{\text{int}})$ associated with the quadratic form $\psi \mapsto \|\nabla \psi\|_{L^2(\Omega_{\text{int}})}^2$ with the domain being the restriction of $H_0^1(\Omega)$ to Ω_{int} , and similarly for $-\Delta_{DN}^{\Omega_{\text{ext}}}$. The spectrum of $-\Delta_{DN}^{\Omega_{\text{int}}}$ is purely discrete. Hence, by the minimax principle,

$$\inf \sigma_{\mathrm{ess}}(-\Delta_D^{\Omega}) \ge \inf \sigma_{\mathrm{ess}}(-\Delta_{DN}^{\Omega_{\mathrm{ext}}}) \ge \inf \sigma(-\Delta_{DN}^{\Omega_{\mathrm{ext}}}) \,.$$

On Ω_{ext} the magnitude of the velocity of rotation of the cross-section ω about $\mathbb{R} \times \{0\}$ is greater than n. Hence, the longitudinal distance on which the cross-section turns by 180 degrees is estimated from above by π/n . That is, by our hypothesis, the rays on Ω_{ext} parallel to the axis of rotation and emanating from the cuts $\{\pm s_n\} \times \omega$ will hit a Dirichlet boundary of Ω_{ext} at most at the distance π/n . The same argument shows that the rays on Ω_{ext} parallel to the axis of rotation and emanating from any cut $\{x_1\} \times \omega$, with $|x_1| > s_n + \pi/n$, will hit a Dirichlet boundary of Ω_{ext} in both directions at most at the distance π/n . Consequently, for any ψ from the form domain of $-\Delta_{DN}^{\Omega_{\text{ext}}}$, we have

$$\|\nabla\psi\|_{L^2(\Omega_{\text{ext}})}^2 \ge \|\partial_1\psi\|_{L^2(\Omega_{\text{ext}})}^2 \ge \left(\frac{\pi}{2\pi/n}\right)^2 \|\psi\|_{L^2(\Omega_{\text{ext}})}^2.$$

By the minimax principle, it thus follows

$$\inf \sigma(-\Delta_{DN}^{\Omega_{\text{ext}}}) \ge \frac{n^2}{4} \,.$$

Since n can be chosen arbitrarily large, we conclude with $\sigma_{\text{ess}}(-\Delta_D^{\Omega}) = \varnothing$. \Box

It follows from Theorem 2 that the embedding (1) is compact. Since the quasi-boundedness is a necessary condition for the compactness of (1), we see that Ω is quasi-bounded under the hypothesis of Theorem 2. The geometrical setting of Theorem 2 is illustrated by Figure 2.



Figure 2: Quasi-bounded realisation: Waveguide of elliptical cross-section with a centre of rotation in the exterior of the ellipse.

5 Conclusion

Condition (5) leads to a new class of waveguides, whose detailed spectral analysis constitutes an interesting open problem.

For instance, in the quasi-cylindrical setting (Theorem 1), one could study the existence of discrete spectra or Hardy-type inequalities and the rate of accumulation of discrete eigenvalues to the threshold of the essential spectrum.

In the quasi-bounded setting (Theorem 2), an interesting question is to establish some sort of Weyl-type asymptotics for the discrete eigenvalues. Note that Ω has an infinite volume regardless of the choice of θ . In fact, the volume is "locally preserved" for twisted tubes, *i.e.* $|\mathcal{L}((a_1, a_2) \times \omega)| = |(a_1, a_2) \times \omega|$ for any $a_1 < a_2$. This makes the present model very different from [5], where the cross-section shrinks to a point at the infinity of the waveguide. Sharp semiclassical estimates of eigenvalues in distinct types of domains of infinite volume has been established recently in [7].

Acknowledgment

I would like to thank Rupert L. Frank who raised the question of waveguides with asymptotically diverging twisting after my talk in the *Isaac Newton Institute for Mathematical Sciences* in Cambridge, April 2007. The research was partially supported by the project RVO61389005 and the GACR grant No. 14-06818S. The author also acknowledges the award from the *Neuron fund for support of science*, Czech Republic, May 2014.

References

- Ph. Briet, H. Hammedi, and D. Krejčiřík, Hardy inequalities in globally twisted waveguides, arXiv:1406.2841 [math-SP] (2014).
- [2] P. Duclos and P. Exner, Curvature-induced bound states in quantum waveguides in two and three dimensions, Rev. Math. Phys. 7 (1995), 73–102.
- [3] D. E. Edmunds and W. D. Evans, Spectral theory and differential operators, Oxford University Press, 1987.
- [4] T. Ekholm, H. Kovařík, and D. Krejčiřík, A Hardy inequality in twisted waveguides, Arch. Ration. Mech. Anal. 188 (2008), 245–264.
- [5] P. Exner and D. Barseghyan, Spectral estimates for Dirichlet Laplacians and Schrödinger operators on geometrically nontrivial cusps, J. Spectr. Theory 3 (2013), 465–484.
- [6] P. Exner and H. Kovařík, Spectrum of the Schrödinger operator in a perturbed periodically twisted tube, Lett. Math. Phys. 73 (2005), 183–192.
- [7] L. Geisinger and T. Weidl, Sharp spectral estimates in domains of infinite volume, Rev. Math. Phys. 23 (2011), 615–641.
- [8] I. M. Glazman, Direct methods of qualitative spectral analysis of singular differential operators, Israel Program for Scientific Translations, 1965.
- [9] S. Haag, J. Lampart, and S. Teufel, *Generalised quantum waveguides*, arXiv:1402.1067 [math-ph] (2014).
- [10] D. Krejčiřík, Twisting versus bending in quantum waveguides, Analysis on Graphs and its Applications, Cambridge, 2007 (P. Exner et al., ed.), Proc. Sympos. Pure Math., vol. 77, Amer. Math. Soc., Providence, RI, 2008, pp. 617–636. See arXiv:0712.3371v2 [math-ph] (2009) for a corrected version.
- [11] D. Krejčiřík and J. Kříž, On the spectrum of curved quantum waveguides, Publ. RIMS, Kyoto University 41 (2005), no. 3, 757–791.
- [12] D. Krejčiřík and H. Šediváková, The effective Hamiltonian in curved quantum waveguides under mild regularity assumptions, Rev. Math. Phys. 24 (2012), 1250018.
- [13] D. Krejčiřík and E. Zuazua, The Hardy inequality and the heat equation in twisted tubes, J. Math. Pures Appl. 94 (2010), 277–303.
- [14] J. Wachsmuth and S. Teufel, Effective Hamiltonians for constrained quantum systems, Mem. Amer. Math. Soc. 1083 (2013).