

A LEHTO–VIRTANEN-TYPE THEOREM AND A RESCALING PRINCIPLE FOR AN ISOLATED ESSENTIAL SINGULARITY OF A HOLOMORPHIC CURVE IN A COMPLEX SPACE

YŪSUKE OKUYAMA

To the memory of Professor Shoshichi Kobayashi

ABSTRACT. We establish a Lehto–Virtanen-type theorem and a rescaling principle for an isolated essential singularity of a holomorphic curve in a complex space, which are useful for establishing a big Picard-type theorem and a big Brody-type one for holomorphic curves.

1. INTRODUCTION

Let V be a complex space. For a holomorphic mapping $f : \mathbb{D} \setminus \{0\} \rightarrow V$, we say that f has an *isolated essential singularity* at the origin if f does not extend to a holomorphic mapping from $\mathbb{D}$ to a complex space. Our aim is to contribute to the study of the Kobayashi hyperbolicity and the Brody one of a complex space by establishing a Lehto–Virtanen-type theorem and a rescaling principle for an isolated essential singularity of a holomorphic curve in a complex space.

Notation 1.1. Set $\mathbb{D}(a, r) := \{z \in \mathbb{C} : |z - a| < r\}$ for every $a \in \mathbb{C}$ and every $r > 0$, and set $\mathbb{D}(r) := \mathbb{D}(0, r)$ for every $r > 0$. Then $\mathbb{D}(1) = \mathbb{D}$. For every metric δ on a complex space V , set $\text{diam}_\delta(S) := \sup\{\delta(a, a') : a, a' \in S\}$ for a non-empty subset S in V . Finally, for a complex space V , let d_V be the Kobayashi pseudometric on V .

For the foundation of hyperbolic complex spaces, see the books [5, 6].

1.1. A Lehto–Virtanen-type theorem and a big Picard-type theorem. The following is a generalization of Lehto–Virtanen [8].

Theorem 1 (a Lehto–Virtanen-type theorem). *Let V be a complex space equipped with a metric δ inducing the topology of V , and $f : \mathbb{D} \setminus \{0\} \rightarrow V$ be a holomorphic mapping having an isolated essential singularity at the origin. If $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$, then there exists a sequence (z_n) in $\mathbb{D} \setminus \{0\}$ tending to 0 as $n \rightarrow \infty$ such that $\lim_{n \rightarrow \infty} f(z_n)$ exists in V and that $\liminf_{n \rightarrow \infty} \text{diam}_\delta(f(\partial\mathbb{D}(|z_n|))) > 0$.*

Date: December 11, 2021.

2010 Mathematics Subject Classification. Primary 32Q45; Secondary 32H02.

Key words and phrases. holomorphic curve, isolated essential singularity, complex space, rescaling principle, Kobayashi hyperbolicity, Brody hyperbolicity.

Partially supported by JSPS Grant-in-Aid for Young Scientists (B), 24740087.

Remark 1.2. There always exists a metric δ on V inducing the topology of V . When V is Kobayashi hyperbolic, we can set $\delta = d_V$ in Theorem 1.

In this article, we do *not* require the relative compactness of Y in Z in the following definition.

Definition 1.3 (a hyperbolically imbedded complex subspace). A complex subspace Y in a complex space Z is called a hyperbolically imbedded complex subspace in Z if for every distinct points $p, q \in \overline{Y}$, there exist open neighborhoods U_p, U_q of p, q in Z , respectively, such that $d_Y(U_p \cap Y, U_q \cap Y) > 0$.

The following, which will be established as an application of Theorem 1, slightly generalizes Kobayashi [6, Theorem VII.3.6].

Theorem 2 (a big Picard-type theorem). *Let Y be a hyperbolically imbedded complex subspace in a complex space Z , and $f : \mathbb{D} \setminus \{0\} \rightarrow Y$ be a holomorphic mapping. If $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$ (as a subset in Z), then f extends to a holomorphic mapping from $\mathbb{D}$ to Z .*

A Kobayashi hyperbolic complex space is a hyperbolically imbedded complex subspace in itself. Hence Theorem 2 is a generalization of Kwack [7], which can be restated as follows.

Theorem 1.4 (Kwack [7]). *If there is a holomorphic mapping from $\mathbb{D} \setminus \{0\}$ to a complex space V having an isolated essential singularity at the origin and satisfying $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$, then V is not Kobayashi hyperbolic.*

Remark 1.5. The corresponding little Picard-type theorem, which does not require the assumption $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$, follows from the definition of the Kobayashi hyperbolicity: see Fact 2.8 below.

1.2. A rescaling principle and a big Brody-type theorem. When a complex space is compact, it admits a nice metric (see Theorem 2.9).

Theorem 1 together with a Zalcman-type theorem (see Theorem 2.13 or 2.14) also establishes the following rescaling principle for an isolated essential singularity of a holomorphic curve.

Theorem 3 (a rescaling principle). *Let V be a compact complex space equipped with a metric satisfying the conditions in Theorem 2.9, and $f : \mathbb{D} \setminus \{0\} \rightarrow V$ be a holomorphic mapping. Then, f has an isolated essential singularity at the origin if and only if there are sequences (z_k) and (ρ_k) in $\mathbb{C}$ and $(0, \infty)$, respectively, and a non-constant holomorphic mapping $g : X \rightarrow V$, where X is either $\mathbb{C}$ or $\mathbb{C} \setminus \{0\}$, such that $\lim_{k \rightarrow \infty} z_k = 0$ on $\mathbb{C}$, that $\lim_{k \rightarrow \infty} \rho_k = 0$ in $\mathbb{R}$, and that $\lim_{k \rightarrow \infty} f(z_k + \rho_k v) = g(v)$ locally uniformly on X .*

Remark 1.6. Originally, similar results to Theorems 1 and 3 have been established in [11, Lemma 3.1, Theorem 1] for quasiregular mappings from a punctured ball to a (compact) Riemannian manifold having an isolated essential singularity at the puncture, and are applied not only to deduce “big” results from their corresponding “little” ones (e.g., Holopainen–Pankka’s big Picard-type theorem [3] from Holopainen–Rickman’s little one [4] for quasiregular mappings when the target is compact) but also to establish

the density of repelling periodic points in the Julia set for *local uniformly quasiregular* dynamics including complex dynamics [10].

Remark 1.7 (a holomorphic mapping exceptional in Julia's sense). In Theorem 3, when V is a compact Hermitian complex manifold equipped with an Hermitian metric δ_V , the sequence (z_k) can identically equal 0 (and then $X = \mathbb{C} \setminus \{0\}$) if the mapping f is *exceptional in Julia's sense* in that

$$\limsup_{z \rightarrow 0} |z| f^\#(z) < \infty,$$

where we set $f^\#(z) := \lim_{w \rightarrow z} \delta_V(f(z), f(w))/|z - w|$ on $\mathbb{D} \setminus \{0\}$. Conversely, if $\limsup_{z \rightarrow 0} |z| f^\#(z) = \infty$, then the case $X = \mathbb{C}$ can occur. Some examples of both non-exceptional f and exceptional f in Julia's sense have been known and studied in the Nevanlinna theory: see Lehto–Virtanen [8].

When V is compact, the following improvement of Theorem 1.4 immediately follows from Theorem 3 (see also Remark 2.7).

Corollary 1.8 (a big Brody-type theorem). *If there is a holomorphic mapping from $\mathbb{D} \setminus \{0\}$ to a compact complex space V having an isolated essential singularity at the origin, then V is not Brody hyperbolic.*

Remark 1.9. Corollary 1.8 is also a consequence of Theorem 1.4 and the equivalence between the Kobayashi hyperbolicity and the Brody one for compact complex spaces, the latter of which is known as Brody's theorem [2]. See Fact 2.8 and Remark 2.16 below.

1.3. Organization of this article. In Section 2, we gather some background materials from the Kobayashi hyperbolic geometry as well as a proof of a Zalcman-type theorem (Theorem 2.13) using the original Zalcman's lemma (Theorem 2.14) in the setting that V is not necessarily an Hermitian compact complex manifold, which we hope might be of independent interest. We show Theorems 1, 2, and 3 in Sections 3, 4, and 5, respectively. Section 6 is devoted to some details on Remark 1.7.

2. BACKGROUND

Let V be a complex space. Let us recall the definition of the Kobayashi pseudometric d_V on V (cf. [6, §VII] or [12, §2]).

Definition 2.1 (the Kobayashi pseudometric d_V on V). For every $p, q \in V$, a chain of holomorphic disks from p to q is a pair of a finite sequence $(f_j)_{j=1}^m$ of holomorphic mappings from $\mathbb{D}$ to V and a finite sequence $((a_j, b_j))_{j=1}^m$ in $\mathbb{D} \times \mathbb{D}$ such that $f_1(a_1) = p$, $f_m(b_m) = q$ and that for every $j \in \{1, 2, \dots, m-1\}$, $f_j(b_j) = f_{j+1}(a_{j+1})$ (the $m \in \mathbb{N}$ depends on each chain of holomorphic disks from p to q). Let $\rho_{\mathbb{D}}$ be the Poincaré (, Bergman or hyperbolic) metric on $\mathbb{D}$. Then $d_V(p, q)$ is defined by the infimum of the sum $\sum_{j=1}^m \rho_{\mathbb{D}}(a_j, b_j)$, where the infimum is taken over all chains $((f_j)_{j=1}^m, ((a_j, b_j))_{j=1}^m)$ of holomorphic disks from p to q .

Example 2.2. For every $a \in \mathbb{C}$ and every $R > 0$, the Kobayashi pseudometric $d_{\mathbb{D}(a, R)}$ on $\mathbb{D}(a, R)$ coincides with the Poincaré metric $\rho_{\mathbb{D}(a, R)}$ on $\mathbb{D}(a, R)$, which is a Kähler metric given by $ds_{\mathbb{D}(a, R)} = R|dz|/(R^2 - |z - a|^2)$ on

$\mathbb{D}(a, R)$. The following comparison is useful: for every $a \in \mathbb{C}$, every $R > 0$, every $r \in (0, R)$, and every $z, w \in \overline{\mathbb{D}(a, r)}$,

$$(2.1) \quad \frac{1}{R}|z - w| \leq d_{\mathbb{D}(a, R)}(z, w) \leq \frac{R}{R^2 - r^2}|z - w|.$$

Example 2.3 (cf. [6, Propositions IV.1.1 and VI.2.1]). The Kobayashi pseudometric $d_{\mathbb{D} \setminus \{0\}}$ on $\mathbb{D} \setminus \{0\}$ coincides with the hyperbolic metric on $\mathbb{D} \setminus \{0\}$, which is a Kähler metric given by $ds_{\mathbb{D} \setminus \{0\}} = |dz|/(-|z| \log |z|)$ on $\mathbb{D} \setminus \{0\}$. In particular, the arc length of the circle $\partial\mathbb{D}(r)$ with respect to $d_{\mathbb{D} \setminus \{0\}}$ is $O(1/(-\log r))$ as $r \rightarrow 0$.

Fact 2.4. The Kobayashi pseudometrics on complex spaces enjoy the *non-increasing* property under holomorphisms in that for complex spaces X, Y , a holomorphic mapping $f : X \rightarrow Y$, and points $x, x' \in X$,

$$d_Y(f(x), f(x')) \leq d_X(x, x').$$

In particular, the Kobayashi pseudometrics are invariant under biholomorphisms between complex spaces.

Definition 2.5 (the Kobayashi hyperbolicity and the Brody one). A complex space V is said to be Kobayashi hyperbolic (resp. Brody hyperbolic) if the Kobayashi pseudometric d_V is a metric on V (resp. if there is no non-constant holomorphic mapping from $\mathbb{C}$ to V).

Fact 2.6. If a complex space V is Kobayashi hyperbolic, then d_V induces the topology of V .

Remark 2.7. For a non-constant holomorphic mapping $g : \mathbb{C} \setminus \{0\} \rightarrow V$, $g \circ \exp : \mathbb{C} \rightarrow V$ is non-constant and holomorphic. Conversely, for a non-constant holomorphic mapping $g : \mathbb{C} \rightarrow V$, $g|(\mathbb{C} \setminus \{0\}) : \mathbb{C} \setminus \{0\} \rightarrow V$ is non-constant and holomorphic. Hence, a complex space V is Brody hyperbolic if and only if there is no non-constant holomorphic mapping from X to V , where X is either $\mathbb{C}$ or $\mathbb{C} \setminus \{0\}$.

Fact 2.8 (a little Picard-type theorem and Brody's theorem). If a complex space V is Kobayashi hyperbolic, then it is also Brody hyperbolic: this “little Picard-type theorem” is almost by the definition of the Kobayashi hyperbolicity and that of the Brody one.

Brody's theorem asserts that the converse is also true if in addition V is compact, that is, a Brody hyperbolic compact complex space is Kobayashi hyperbolic.

When a complex space is compact, it admits a nice metric.

Theorem 2.9 (cf. [12, Subsection 4.1]). For every compact complex space V , there is a metric δ on V satisfying that

- (1) the distance δ induces the topology of V , and that
- (2) there is an open covering $\{U_x : x \in V\}$ of V such that for every $x \in V$, U_x is a Kobayashi hyperbolic subdomain in V containing x and satisfies $\delta \leq d_{U_x}$ on $U_x(\times U_x)$.

The following *Lipschitz continuity on disks* of holomorphic curves from a domain in $\mathbb{C}$ into compact complex spaces is useful: the boundedness (2.2) is

a special case of the “only if” part of a Marty-type theorem (Theorem 2.12) below, and the equality (2.3) is by the invariance under biholomorphisms of the Kobayashi (pseudo)metrics (cf. Fact 2.4).

Theorem 2.10 (cf. [12, Subsection 2.3]). *Let V be a compact complex space equipped with a metric δ satisfying the conditions in Theorem 2.9. Then for every open disk $\mathbb{D}(a, r)$ and every holomorphic mapping f from an open neighborhood of $\mathbb{D}(a, r)$ in $\mathbb{C}$ to V , we have*

$$(2.2) \quad L_{f, \mathbb{D}(a, r)} := \sup_{w, w' \in \mathbb{D}(a, r), w \neq w'} \frac{\delta(f(w), f(w'))}{d_{\mathbb{D}(a, r)}(w, w')} < \infty,$$

which satisfies the invariance

$$(2.3) \quad L_{f \circ \phi, \mathbb{D}(b, s)} = L_{f, \mathbb{D}(a, r)}$$

for every biholomorphism $\phi : \mathbb{D}(b, s) \rightarrow \mathbb{D}(a, r) = \phi(\mathbb{D}(b, s))$.

Definition 2.11. For complex spaces X and Y , let $\mathcal{O}(X, Y)$ be the set of all holomorphic mappings from X to Y .

We conclude this section with a generalization of Marty’s theorem [9, Théorème 5] and that of Zalcman’s lemma [13], both of which characterize the (non-)normality on D of a family in $\mathcal{O}(D, V)$ for a domain D in $\mathbb{C}$ and a compact complex space V .

Theorem 2.12 (a Marty-type theorem). *Let D be a domain in $\mathbb{C}$, V a compact complex space equipped with a metric δ satisfying the conditions in Theorem 2.9, and $\mathcal{F}$ a family in $\mathcal{O}(D, V)$. Then, $\mathcal{F}$ is normal on D if and only if $\sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(a, r)} < \infty$ for every $a \in D$ and every $r > 0$ satisfying $\mathbb{D}(a, r) \subseteq D$.*

Proof. By the Arzelà–Ascoli theorem, $\mathcal{F}$ is normal on D if and only if $\mathcal{F}$ is locally equicontinuous on D . Let us show that $\mathcal{F}$ is locally equicontinuous on D if and only if $\sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(a, r)} < \infty$ for every $a \in D$ and every $r > 0$ satisfying $\mathbb{D}(a, r) \subseteq D$. The “if” part is straightforward by (2.1).

Suppose that $\mathbb{D}(a, r) \subseteq D$ but $\sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(a, r)} = \infty$ for some $a \in D$ and some $r > 0$. Then there are a sequence (f_k) in $\mathcal{F}$ and sequences $(z_k), (w_k)$ in $\mathbb{D}(a, r)$ such that $z_k \neq w_k$ for every $k \in \mathbb{N}$ and that

$$(2.4) \quad \lim_{k \rightarrow \infty} \frac{\delta(f_k(z_k), f_k(w_k))}{d_{\mathbb{D}(a, r)}(z_k, w_k)} = \infty.$$

Increasing $r > 0$ slightly and taking a subsequence if necessary, the limit $b := \lim_{k \rightarrow \infty} z_k$ exists in $\mathbb{D}(a, r)$, and then by $\text{diam}_\delta(V) < \infty$, we also have $\lim_{k \rightarrow \infty} w_k = b$.

Suppose contrary that $\mathcal{F}$ is locally equicontinuous on D . Then by the Arzelà–Ascoli theorem, taking a subsequence if necessary, there exists a locally uniform limit $f := \lim_{k \rightarrow \infty} f_k$ on D . Then there exists $\epsilon > 0$ so small that for every $k \in \mathbb{N}$, $f_k(\mathbb{D}(b, \epsilon)) \subset U_{f(b)}$. Hence, for every $k \in \mathbb{N}$ so large that $z_k, w_k \in \mathbb{D}(b, \epsilon)$, we have

$$(2.5) \quad \delta(f_k(z_k), f_k(w_k)) \leq d_{U_{f(b)}}(f_k(z_k), f_k(w_k)) \leq d_{\mathbb{D}(b, \epsilon)}(z_k, w_k)$$

by the property (2) of U_x for $x = f(b)$ in Theorem 2.9 and the non-increasing property under holomorphisms of the Kobayashi (pseudo)metrics (cf. Fact 2.4). By (2.5) and (2.1), we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{\delta(f_k(z_k), f_k(w_k))}{d_{\mathbb{D}(a,r)}(z_k, w_k)} &\leq \lim_{k \rightarrow \infty} \frac{d_{\mathbb{D}(b,\epsilon)}(z_k, w_k)}{d_{\mathbb{D}(a,r)}(z_k, w_k)} \\ &\leq \lim_{k \rightarrow \infty} \left(\frac{r}{|z_k - w_k|} \cdot \frac{\epsilon \cdot |z_k - w_k|}{\epsilon^2 - (\epsilon/2)^2} \right) = \frac{r}{3\epsilon/4} < \infty, \end{aligned}$$

which contradicts (2.4). Now the proof of the “only if” part is complete. $\square$

Theorem 2.13 (a Zalcman-type theorem). *Let D be a domain in $\mathbb{C}$, V a compact complex space equipped with a metric δ satisfying the conditions in Theorem 2.9, and $\mathcal{F}$ a family in $\mathcal{O}(D, V)$. Then, $\mathcal{F}$ is not normal at a point $a \in D$, that is, not normal on any open neighborhood of a in D , if and only if there are sequences (f_k) , (z_k) , and (ρ_k) in $\mathcal{F}$, D , and $(0, \infty)$, respectively, and a non-constant $g \in \mathcal{O}(\mathbb{C}, V)$ such that $\lim_{k \rightarrow \infty} z_k = a$, that $\lim_{k \rightarrow \infty} \rho_k = 0$, and that $\lim_{k \rightarrow \infty} f_k(z_k + \rho_k v) = g(v)$ locally uniformly on $\mathbb{C}$.*

We will divide the proof of Theorem 2.13 into two parts: a deduction of Theorem 2.13 from the *original* Zalcman’s lemma (Theorem 2.14 below), and a proof of this original one. The former part, say, a *double rescaling argument*, also motivates a part of the proof of Theorem 3.

The following is due to Zalcman in the case where $V = \mathbb{P}^1$ equipped with the Fubini-Study metric.

Theorem 2.14. *Let V be a compact complex space equipped with a metric δ satisfying the conditions in Theorem 2.9, and $\mathcal{F}$ a family in $\mathcal{O}(\mathbb{D}, V)$. Then, $\mathcal{F}$ is not normal on $\mathbb{D}$ if and only if there are sequences (f_k) , (z_k) , and (ρ_k) in $\mathcal{F}$, $\mathbb{D}$, and $(0, \infty)$, respectively, and a non-constant $g \in \mathcal{O}(\mathbb{C}, V)$ such that $\lim_{k \rightarrow \infty} z_k$ exists in $\mathbb{D}$, that $\lim_{k \rightarrow \infty} \rho_k = 0$, and that $\lim_{k \rightarrow \infty} f_k(z_k + \rho_k v) = g(v)$ locally uniformly on $\mathbb{C}$.*

Proof of Theorem 2.13 using Theorem 2.14. The “if” part of Theorem 2.13 is straightforward from that of the original Zalcman’s lemma (Theorem 2.14).

Suppose that $\mathcal{F}$ is not normal at a point $a \in D$. Then by (2.1) and the Arzelà–Ascoli theorem, $\limsup_{r \rightarrow 0} \sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(a,r)} = \infty$, that is, there are sequences (r_k) in $(0, \infty)$ and (f_k) in $\mathcal{F}$ such that $\lim_{k \rightarrow \infty} r_k = 0$ and that

$$(2.6) \quad \lim_{k \rightarrow \infty} L_{f_k, \mathbb{D}(a, r_k)} = \infty.$$

Fix $\epsilon > 0$ small enough. Then for every $k \in \mathbb{N}$, a holomorphic mapping $g_k \in \mathcal{O}(\mathbb{D}(1 + \epsilon), V)$ is defined by

$$g_k(w) := f_k(a + r_k w),$$

and then $L_{g_k, \mathbb{D}} = L_{f_k, \mathbb{D}(a, r_k)}$ by (2.3). Hence by (2.6) and a Marty-type theorem (Theorem 2.12), the family $\{g_k : k \in \mathbb{N}\}$ is not normal on $\mathbb{D}(1 + \epsilon/2)$, so that by Theorem 2.14, there are sequences (w_j) , (η_j) , and (k_j) in $\mathbb{C}$, $(0, \infty)$, and $\mathbb{N}$, respectively, and a non-constant $g \in \mathcal{O}(\mathbb{C}, V)$ such that $\lim_{j \rightarrow \infty} w_j$

exists in $\mathbb{D}(1 + \epsilon/2)$, that $\lim_{j \rightarrow \infty} \eta_j = 0$, that $\lim_{j \rightarrow \infty} k_j = \infty$, and that

$$g(v) = \lim_{j \rightarrow \infty} g_{k_j}(w_j + \eta_j v)$$

locally uniformly on $\mathbb{C}$. For every $j \in \mathbb{N}$, we have $g_{k_j}(w_j + \eta_j v) = f_{k_j}((a + r_{k_j} w_j) + (r_{k_j} \eta_j) v)$ on $\mathbb{D}(\eta_j^{-1}(\epsilon/2))$, and set $z_j := a + r_{k_j} w_j$ and $\rho_j := r_{k_j} \eta_j$.

Now the proof of the “only if” part of Theorem 2.13 is complete since $\lim_{j \rightarrow \infty} z_j = a$ and $\lim_{j \rightarrow \infty} \rho_j = 0$. $\square$

Proof of Theorem 2.14. If $\mathcal{F}$ is not normal on $\mathbb{D}$, then by (2.1) and the Arzelà–Ascoli theorem, we have $\sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(r)} = \infty$ for some $r \in (0, 1)$. Increasing $r \in (0, 1)$ slightly if necessary, we can choose sequences $(z'_k), (w'_k)$ in $\mathbb{D}(r)$ and a sequence (f_k) in $\mathcal{F}$ such that both $\{z'_k : k \in \mathbb{N}\}$ and $\{w'_k : k \in \mathbb{N}\}$ are relatively compact in $\mathbb{D}(r)$, that $z'_k \neq w'_k$ for every $k \in \mathbb{N}$, and that $\lim_{k \rightarrow \infty} \delta(f_k(z'_k), f_k(w'_k))/d_{\mathbb{D}(r)}(z'_k, w'_k) = \infty$. Then, by (2.1), we also have $\lim_{k \rightarrow \infty} \delta(f_k(z'_k), f_k(w'_k))/|z'_k - w'_k| = \infty$.

For every $k \in \mathbb{N}$, set

$$M_k := \sup_{z, w \in \mathbb{D}(r): z \neq w} \left(\frac{r^2 - |z|^2}{r^2} \cdot \frac{\delta(f_k(z), f_k(w))}{|z - w|} \right).$$

Then $\lim_{k \rightarrow \infty} M_k = \infty$. For every $k \in \mathbb{N}$, choose distinct $z_k, w_k \in \mathbb{D}(r)$ satisfying that

$$(2.7) \quad \frac{r^2 - |z_k|^2}{r^2} \cdot \frac{\delta(f_k(z_k), f_k(w_k))}{|z_k - w_k|} \geq \frac{M_k}{2},$$

and set

$$\rho_k := \frac{|z_k - w_k|}{\delta(f_k(z_k), f_k(w_k))} \left(\leq \frac{2}{M_k} \right) \quad \text{and} \quad R_k := \frac{r - |z_k|}{\rho_k} \left(\geq \frac{M_k}{2} \cdot \frac{r}{2} \right).$$

Then $\rho_k \rightarrow 0$ and $R_k \rightarrow \infty$ as $k \rightarrow \infty$. Taking a subsequence if necessary, we can also assume that $\lim_{k \rightarrow \infty} z_k$ exists in $\overline{\mathbb{D}(r)}$.

For every $k \in \mathbb{N}$ and every $v \in \mathbb{D}(R_k)$, $z_k + \rho_k v \in \mathbb{D}(r)$. Hence for each $k \in \mathbb{N}$, a holomorphic mapping $g_k \in \mathcal{O}(\mathbb{D}(R_k), V)$ is defined by

$$g_k(v) := f_k(z_k + \rho_k v)$$

and satisfies that for every distinct $x, y \in \mathbb{D}(R_k)$,

$$\begin{aligned} \frac{\delta(g_k(x), g_k(y))}{|x - y|} &= \frac{\delta(f_k(z_k + \rho_k x), f_k(z_k + \rho_k y))}{|(z_k + \rho_k x) - (z_k + \rho_k y)|} \cdot \rho_k \\ &\leq \frac{r^2 M_k}{r^2 - |z_k + \rho_k x|^2} \cdot \frac{2}{M_k} \frac{r^2 - |z_k|^2}{r^2} = 2 \frac{r^2 - |z_k|^2}{r^2 - |z_k + \rho_k x|^2} \\ &\leq 2 \frac{(r + |z_k|)(r - |z_k|)}{(r + |z_k + \rho_k x|)(r - |z_k| - \rho_k |x|)} \leq 2 \cdot \frac{2r}{r} \cdot \frac{1}{1 - |x|/R_k} = \frac{4}{1 - |x|/R_k}. \end{aligned}$$

Hence $\{g_k : k \in \mathbb{N}\}$ is locally equicontinuous on $\mathbb{C}$.

By the Arzelà–Ascoli theorem, taking a subsequence if necessary, there exists the locally uniform limit $g := \lim_{k \rightarrow \infty} g_k$ on $\mathbb{C}$, which is in $\mathcal{O}(\mathbb{C}, V)$. It remains to show that g is non-constant: For every $k \in \mathbb{N}$, set

$$v_k := \frac{z_k - w_k}{\rho_k} \in \mathbb{C} \setminus \{0\}.$$

By the definition of ρ_k , we have not only $\sup_{k \in \mathbb{N}} |v_k| \leq \text{diam}_\delta V < \infty$ but also, for every $k \in \mathbb{N}$,

$$(2.8) \quad \frac{\delta(g_k(0), g_k(v_k))}{|v_k|} = \frac{\delta(f_k(z_k), f_k(w_k))}{|z_k - w_k|} \cdot \rho_k = 1.$$

Taking a subsequence if necessary, we can assume that the limit $v^* := \lim_{k \rightarrow \infty} v_k$ exists in $\mathbb{C}$.

Suppose contrary that g is constant on $\mathbb{C}$. Then by (2.8), we must have $v^* = 0$. Fix $R > 1$. For every $k \in \mathbb{N}$ large enough, we have $v_k \in \mathbb{D}(R)$ and $g_k(\mathbb{D}(R)) \subset U_{g(0)}$, so that

$$(2.9) \quad \delta(g_k(0), g_k(v_k)) \leq d_{U_{g(0)}}(g_k(0), g_k(v_k)) \leq d_{\mathbb{D}(R)}(0, v_k)$$

by the property (2) of U_x for $x = g(0)$ in Theorem 2.9 and the non-increasing property under holomorphisms of the Kobayashi (pseudo)metrics (cf. Fact 2.4). By (2.9) and (2.1), we have

$$\begin{aligned} \limsup_{k \rightarrow \infty} \frac{\delta(g_k(0), g_k(v_k))}{|v_k|} &\leq \limsup_{k \rightarrow \infty} \frac{d_{\mathbb{D}(R)}(0, v_k)}{|v_k|} \\ &\leq \limsup_{k \rightarrow \infty} \frac{R}{R^2 - |v_k|^2} = \frac{1}{R} < 1, \end{aligned}$$

which contradicts (2.8). Hence g is non-constant on $\mathbb{C}$, and the proof of the “only if” part is complete.

Suppose now that there are sequences (f_k) , (z_k) , and (ρ_k) in $\mathcal{F}$, $\mathbb{D}$, and $(0, \infty)$, respectively, and a non-constant $g \in \mathcal{O}(\mathbb{C}, V)$ such that the limit $a := \lim_{k \rightarrow \infty} z_k$ exists in $\mathbb{D}$, that $\lim_{k \rightarrow \infty} \rho_k = 0$, and that $\lim_{k \rightarrow \infty} f_k(z_k + \rho_k v) = g(v)$ locally uniformly on $\mathbb{C}$. Fix $r > 0$ so small that $\mathbb{D}(a, r) \Subset \mathbb{D}$.

Suppose contrary that $\mathcal{F}$ is normal on $\mathbb{D}$. Then by a Marty-type theorem (Theorem 2.12), we have $\sup_{f \in \mathcal{F}} L_{f, \mathbb{D}(a, r)} < \infty$. In particular, by (2.1), we have

$$C := \sup_{f \in \mathcal{F}} \left(\sup_{z, w \in \mathbb{D}(a, r/2), z \neq w} \frac{\delta(f(z), f(w))}{|z - w|} \right) < \infty.$$

For every distinct $x, y \in \mathbb{C}$, since $\lim_{k \rightarrow \infty} (z_k + \rho_k x) = \lim_{k \rightarrow \infty} (z_k + \rho_k y) = a$, we have

$$\frac{\delta(g(x), g(y))}{|x - y|} = \lim_{k \rightarrow \infty} \frac{\delta(f_k(z_k + \rho_k x), f_k(z_k + \rho_k y))}{|(z_k + \rho_k x) - (z_k + \rho_k y)|} \cdot \rho_k \leq C \cdot \lim_{k \rightarrow \infty} \rho_k = 0,$$

so that $g(x) = g(y)$. This contradicts that g is non-constant on $\mathbb{C}$, and the proof of the “if” part is also complete. $\square$

Remark 2.15. For a much shorter proof of (the “only if” part of) Theorem 2.13 in the case where V is an Hermitian compact complex manifold, see Berteloot [1, §2.1].

Remark 2.16. Brody’s theorem (cf. Fact 2.8) in the setting that U is not necessarily an Hermitian compact complex manifold follows from the original Zalcman’s lemma (Theorem 2.14) by showing that *if a compact complex space V is not Kobayashi hyperbolic, then the family $\mathcal{O}(\mathbb{D}, V)$ is not normal on $\mathbb{D}$* : Fix $p, q \in V$. We can fix $r \in (0, 1)$ so close to 1 that $d_V(p, q) = \inf \sum_{j=1}^m d_{\mathbb{D}}(a_j, b_j)$, where the infimum is taken over all such

chains $((f_j)_{j=1}^m, ((a_j, b_j))_{j=1}^m)$ of holomorphic disks from p to q that satisfy $a_j \equiv 0$ and $b_j \in \mathbb{D}(r)$ for every $j \in \{1, 2, \dots, m\}$. Let us equip V with a metric δ satisfying the conditions in Theorem 2.9.

Suppose now that $\mathcal{O}(\mathbb{D}, V)$ is normal on $\mathbb{D}$. Then by a Marty-type theorem (Theorem 2.12), we have $C(r) := \sup_{f \in \mathcal{O}(\mathbb{D}, V)} L_{f, \mathbb{D}(r)} < \infty$, and then using also the triangle inequality for δ , we have $d_V(p, q) \geq \delta(p, q)/C(r)$. This implies that $d_V(p, q) > 0$ if $p \neq q$, i.e., V is Kobayashi hyperbolic.

Remark 2.17. For another proof of Brody's theorem (cf. Fact 2.8) similar to the original Brody's argument and in the setting that U is not necessarily an Hermitian compact complex manifold, see Yamanoi [12, §2.3].

3. PROOF OF THEOREM 1

Let V be a complex space equipped with a metric δ inducing the topology of V , and $f : \mathbb{D} \setminus \{0\} \rightarrow V$ a holomorphic mapping having an isolated essential singularity at the origin. Suppose that $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$. Then we can fix a sequence (z_n) in $\mathbb{D} \setminus \{0\}$ tending to 0 as $n \rightarrow \infty$ such that the limit $a := \lim_{n \rightarrow \infty} f(z_n)$ exists in V . Fix an open neighborhood W of a in V equivalent to an analytic subset in an open subset Ω in $\mathbb{C}^d$ for some $d \in \mathbb{N}$, and fix a subdomain $W' \Subset W$ containing a .

If $\liminf_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z_n|)) > 0$, then we are done. So, suppose that $\liminf_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z_n|)) = 0$. Taking a subsequence if necessary, we can even assume that

$$(3.1) \quad \lim_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z_n|)) = 0.$$

Then for every $n \in \mathbb{N}$ large enough, $f(\partial \mathbb{D}(|z_n|)) \subset W'$. For every $n \in \mathbb{N}$ large enough, since the origin is an isolated essential singularity of f , by Riemann's extension theorem, the following maximum

$$r'_n := \max\{r \in (0, |z_n|) : f(\partial \mathbb{D}(r)) \not\subset W'\} > 0$$

exists, and then $f(\overline{\mathbb{D}(|z_n|)} \setminus \mathbb{D}(r'_n)) \subset \overline{W'}$ by the continuity of f . Fix a sequence (z'_n) in $\mathbb{D} \setminus \{0\}$ tending to 0 as $n \rightarrow \infty$ such that for every $n \in \mathbb{N}$ large enough, $z'_n \in \partial \mathbb{D}(r'_n)$ and $f(z'_n) \in \overline{W'} \setminus W'$. By the compactness of $\overline{W'} \setminus W'$, taking a subsequence if necessary, the limit $b := \lim_{n \rightarrow \infty} f(z'_n)$ exists in $\overline{W'} \setminus W'$. It remains to show that $\liminf_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z'_n|)) > 0$.

Suppose contrary that $\liminf_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z'_n|)) = 0$. Taking a subsequence if necessary, we can even assume that

$$(3.1') \quad \lim_{n \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|z'_n|)) = 0.$$

Since a and b are distinct points in W , which we identify with an analytic subset in an open subset Ω in $\mathbb{C}^d$, there exists an affine coordinate system $w = (w_1, \dots, w_d)$ on Ω such that $w(a) = 0$ and that $w_1(b) \neq 0$. Set $w \circ f = (f_1, \dots, f_d) : f^{-1}(W) \rightarrow w(W)$. Then, for every $n \in \mathbb{N}$ large enough, under the assumptions (3.1) and (3.1'), we have both

$$(3.2) \quad f_1(\partial \mathbb{D}(|z_n|)) \subset \mathbb{D}(|w_1(b)|/3) \quad \text{and} \quad f_1(\partial \mathbb{D}(|z'_n|)) \subset \mathbb{D}(w_1(b), |w_1(b)|/3).$$

Fix such $n \in \mathbb{N}$ as satisfies (3.2). Let ℓ be a line segment in the ring domain $\mathbb{D}(|z_n|) \setminus \overline{\mathbb{D}(|z'_n|)} (\Subset f^{-1}(W))$ having one end point in $\partial \mathbb{D}(|z_n|)$ and

the other in $\partial\mathbb{D}(|z'_n|)$. Then the path $f_1(\ell)$ in $w_1(W)$ joins the closed curves $f_1(\partial\mathbb{D}(|z_n|))$ and $f_1(\partial\mathbb{D}(|z'_n|))$, so by (3.2), we may fix $y_0 \in \ell$ such that

$$(3.3) \quad f_1(y_0) \notin \overline{\mathbb{D}(|w_1(b)|/3)} \cup \overline{\mathbb{D}(w_1(b), |w_1(b)|/3)}.$$

Since f_1 is a holomorphic function on $f^{-1}(W)$ and takes the value $f_1(y_0)$ at least at $y_0 \in \mathbb{D}(|z_n|) \setminus \overline{\mathbb{D}(|z'_n|)}$, by the residue theorem,

$$(3.4) \quad 1 \leq \frac{1}{2i\pi} \int_{\partial(\mathbb{D}(|z_n|) \setminus \overline{\mathbb{D}(|z'_n|)})} \frac{f'_1(z)dz}{f_1(z) - f(y_0)} \\ = \frac{1}{2i\pi} \int_{(f_1)_*(\partial\mathbb{D}(|z_n|))} \frac{dw_1}{w_1 - f(y_0)} - \frac{1}{2i\pi} \int_{(f_1)_*(\partial\mathbb{D}(|z'_n|))} \frac{dw_1}{w_1 - f(y_0)},$$

where the boundary $\partial(\mathbb{D}(|z_n|) \setminus \overline{\mathbb{D}(|z'_n|)})$ is canonically oriented. On the other hand, by (3.2) and (3.3), the argument principle yields

$$\frac{1}{2i\pi} \int_{(f_1)_*(\partial\mathbb{D}(|z_n|))} \frac{dw_1}{w_1 - f(y_0)} = \frac{1}{2i\pi} \int_{(f_1)_*(\partial\mathbb{D}(|z'_n|))} \frac{dw_1}{w_1 - f(y_0)} = 0,$$

which contradicts (3.4).

Hence $\liminf_{n \rightarrow \infty} \text{diam}_\delta f(\partial\mathbb{D}(|z'_n|)) > 0$, and the proof is complete. $\square$

Remark 3.1. The final residue theoretic argument applied to f_1 can be replaced by a more topological argument (for f_1) as in [11, Proof of Lemma 3.1]. In [11, Lemma 3.1], the target Riemannian n -manifold M of a quasiregular mapping $f : \mathbb{B}^n \setminus \{0\} \rightarrow M$ was assumed to be compact, but this assumption can be relaxed as $\bigcap_{r>0} \overline{f(\mathbb{B}^n(r) \setminus \{0\})} \neq \emptyset$ as in Theorem 1. Moreover, in [11, Lemma 3.1], we only claimed that $\limsup_{r \rightarrow 0} \text{diam}(f(\partial\mathbb{B}^n(r))) > 0$, but this assertion can be so strengthen that there exists a sequence (x_j) in $\mathbb{B}^n \setminus \{0\}$ tending to 0 as $j \rightarrow \infty$ such that $\lim_{j \rightarrow \infty} f(x_j)$ exists in M and that $\liminf_{j \rightarrow \infty} \text{diam}(f(\partial\mathbb{B}^n(|x_j|))) > 0$, as in Theorem 1.

4. PROOF OF THEOREM 2

Recall that d_V denotes the Kobayashi pseudometric on a complex space V and that, for every metric δ on V , we set $\text{diam}_\delta(S) = \sup\{\delta(a, a') : a, a' \in S\}$ for a non-empty subset S in V .

Let Y be a complex subspace in a complex space Z , fix a metric δ on Z inducing the topology of Z , and let $f : \mathbb{D} \setminus \{0\} \rightarrow Y$ be a holomorphic mapping.

Suppose that $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$ (as a subset in Z) and that f does not extend to a holomorphic mapping from $\mathbb{D}$ to Z . We claim that Y is not a hyperbolically imbedded complex subspace in Z .

Under the above assumption, by Theorem 1, there exists a sequence (z_n) in $\mathbb{D} \setminus \{0\}$ tending to 0 as $n \rightarrow \infty$ such that the limit

$$a := \lim_{n \rightarrow \infty} f(z_n)$$

exists in $\overline{Y}$ and that $r_0 := \liminf_{n \rightarrow \infty} \text{diam}_\delta(f(\partial\mathbb{D}(|z_n|))) > 0$. Fix a relatively compact open neighborhood W of a in $\{p \in Z : \delta(p, a) < r_0\}$. Then,

taking a subsequence of (z_n) if necessary, there is a sequence (w_n) in $\mathbb{D} \setminus \{0\}$ such that for every $n \in \mathbb{N}$,

$$w_n \in \partial\mathbb{D}(|z_n|)$$

and $f(w_n) \in \partial W$. By the compactness of ∂W , taking a subsequence of (z_n) if necessary, the limit

$$b := \lim_{n \rightarrow \infty} f(w_n)$$

also exists in $\overline{Y} \setminus \{a\}$.

Now, for every open neighborhoods U_a, U_b of a, b in Z , respectively, we have

$$\begin{aligned} d_Y(Y \cap U_a, Y \cap U_b) &\leq \limsup_{n \rightarrow \infty} d_Y(f(z_n), f(w_n)) \\ &\leq \limsup_{n \rightarrow \infty} d_{\mathbb{D} \setminus \{0\}}(z_n, w_n) \leq \limsup_{n \rightarrow \infty} \text{diam}_{d_{\mathbb{D} \setminus \{0\}}}(\partial\mathbb{D}(|z_n|)) = 0, \end{aligned}$$

where the second inequality is by the non-increasing property of Kobayashi pseudometrics under holomorphisms (see Fact 2.4) and the final equality is by a direct computation (cf. Example 2.3).

Hence Y is not a hyperbolically imbedded complex subspace in Z , and the proof is complete. $\square$

5. PROOF OF THEOREM 3

Let V be a compact complex space and $f : \mathbb{D} \setminus \{0\} \rightarrow V$ be a holomorphic mapping having an isolated essential singularity at the origin, and fix a metric δ on V satisfying the conditions in Theorem 2.9.

Let us first show the “only if” part of Theorem 3. We study the cases that **(i)** $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} = \infty$ and that **(ii)** $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} < \infty$, separately.

Suppose first that $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} = \infty$. In this case, the following “double rescaling” argument is similar to that in the proof of Theorem 2.13 using Theorem 2.14 in Section 2. Choose a sequence (y_k) in $\mathbb{D} \setminus \{0\}$ such that $\lim_{k \rightarrow \infty} y_k = 0$ and that

$$(5.1) \quad \lim_{k \rightarrow \infty} L_{f, \mathbb{D}(y_k, |y_k|/2)} = \infty.$$

Fix $\epsilon > 0$ small enough. Then for every $k \in \mathbb{N}$, a holomorphic mapping $g_k : \mathbb{D}(1 + \epsilon) \rightarrow V$ is defined by

$$g_k(w) := f\left(y_k + \frac{|y_k|}{2}w\right),$$

so that $L_{g_k, \mathbb{D}} = L_{f, \mathbb{D}(y_k, |y_k|/2)}$ by (2.3). Hence by (5.1) and a Marty-type theorem (Theorem 2.12), the family $\{g_k : k \in \mathbb{N}\}$ is not normal on $\mathbb{D}(1 + \epsilon/2)$, so that by Theorem 2.14, there are sequences (w_j) , (η_j) , and (k_j) in $\mathbb{C}$, $(0, \infty)$, and $\mathbb{N}$, respectively, and a non-constant $g \in \mathcal{O}(\mathbb{C}, V)$ such that $\lim_{j \rightarrow \infty} w_j$ exists in $\mathbb{D}(1 + \epsilon/2)$, that $\lim_{j \rightarrow \infty} \eta_j = 0$, that $\lim_{j \rightarrow \infty} k_j = \infty$, and that

$$g(v) = \lim_{j \rightarrow \infty} g_{k_j}(w_j + \eta_j v)$$

locally uniformly on $\mathbb{C}$. For every $j \in \mathbb{N}$, we have $g_{k_j}(w_j + \eta_j v) = f((y_{k_j} + (|y_{k_j}|/2)w_j) + ((|y_{k_j}|/2)\eta_j)v)$ on $\mathbb{D}(\eta_j^{-1}(\epsilon/2))$, and set $z_j := y_{k_j} + (|y_{k_j}|/2)w_j$ and $\rho_j := (|y_{k_j}|/2)\eta_j$. We are done in this case (i) since $\lim_{j \rightarrow \infty} z_j = 0$ and $\lim_{j \rightarrow \infty} \rho_j = 0$.

Suppose next that $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} < \infty$. By the compactness of V , we have $\bigcap_{r>0} \overline{f(\mathbb{D}(r) \setminus \{0\})} \neq \emptyset$, so that by Theorem 1, there is a sequence (w_k) in $\mathbb{D} \setminus \{0\}$ tending to 0 as $k \rightarrow \infty$ such that $a := \lim_{k \rightarrow \infty} f(w_k)$ exists in V and that $\liminf_{k \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|w_k|)) > 0$.

For every $k \in \mathbb{N}$, a holomorphic mapping $g_k : \mathbb{D}(|w_k|^{-1}) \setminus \{0\} \rightarrow V$ is defined by

$$g_k(v) := f(|w_k|v).$$

Then for every $v \in \mathbb{C} \setminus \{0\}$,

$$\limsup_{k \rightarrow \infty} L_{g_k, \mathbb{D}(v, |v|/2)} = \limsup_{k \rightarrow \infty} L_{f, \mathbb{D}(|w_k|v, |w_k|v/2)} \leq \limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} < \infty,$$

where the equality is by (2.3) and the next inequality is by $\lim_{k \rightarrow \infty} |w_k| = 0$. Hence by (2.1) and the Arzelà–Ascoli theorem, the family $\{g_k : k \in \mathbb{N}\}$ is normal on $\mathbb{C} \setminus \{0\}$ so, taking a subsequence if necessary, the locally uniform limit $g := \lim_{k \rightarrow \infty} g_k$ exists on $\mathbb{C} \setminus \{0\}$, which is in $\mathcal{O}(\mathbb{C} \setminus \{0\}, V)$. It remains to show that g is non-constant on $\mathbb{C} \setminus \{0\}$. If g is constant, then since for every $k \in \mathbb{N}$,

$$g_k(\partial \mathbb{D}) = f(\partial \mathbb{D}(|w_k|)) \ni f(w_k),$$

we have $g \equiv a = \lim_{k \rightarrow \infty} f(w_k)$ on $\mathbb{C} \setminus \{0\}$, and in turn

$$0 = \text{diam}_\delta(\{a\}) = \limsup_{k \rightarrow \infty} \text{diam}_\delta(g_k(\partial \mathbb{D})) = \limsup_{k \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|w_k|)),$$

which contradicts that $\liminf_{k \rightarrow \infty} \text{diam}_\delta f(\partial \mathbb{D}(|w_k|)) > 0$. Hence g is non-constant on $\mathbb{C} \setminus \{0\}$, and we are also done in this case (ii) by setting $z_k \equiv 0$ and $\rho_k := |w_k|$ for each $k \in \mathbb{N}$. Now the proof of the “only if” part of Theorem 3 is complete.

Suppose now that there are sequences (z_k) and (ρ_k) in $\mathbb{C}$ and $(0, \infty)$, respectively, and a non-constant holomorphic mapping $g : X \rightarrow V$, where X is either $\mathbb{C}$ or $\mathbb{C} \setminus \{0\}$, such that $\lim_{k \rightarrow \infty} z_k = 0$ in $\mathbb{C}$, that $\lim_{k \rightarrow \infty} \rho_k = 0$ in $\mathbb{R}$, and that $\lim_{k \rightarrow \infty} f(z_k + \rho_k v) = g(v)$ locally uniformly on X . If f extends to a holomorphic mapping from $\mathbb{D}$ to V , then for every $v \in X$, $g(v) = \lim_{k \rightarrow \infty} f(z_k + \rho_k v) = f(0)$, which contradicts that g is non-constant on X . Now the proof of the “if” part of Theorem 3 is also complete. $\square$

Remark 5.1. The proof of Theorem 3 is similar to [11, Proof of Theorem 1] for quasiregular mappings. In the holomorphic curve case, however, the *Lipschitz continuity on disks* (2.2) of holomorphic curves and the invariance of the Kobayashi pseudometrics under biholomorphisms between complex spaces make the argument much simpler than that in the quasiregular case.

6. ON REMARK 1.7

Let V be a compact Hermitian manifold equipped with an Hermitian metric δ_V . Then δ_V satisfies the properties in Theorem 2.9 (cf. [12, §2.3]). Let

$f : \mathbb{D} \setminus \{0\} \rightarrow V$ be a holomorphic curve having an isolated essential singularity at the origin, and recall that $f^\#(z) := \lim_{w \rightarrow z} \delta_V(f(z), f(w))/|z - w|$ on $\mathbb{D} \setminus \{0\}$. Then $f^*ds_V = f^\#|dz|$ on $\mathbb{D}$, where ds_V is the arc-length element of δ_V on V .

Let us see that, *in the proof of the “only if” part of Theorem 3, the case $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} < \infty$ occurs if and only if f is exceptional in Julia’s sense*: by (2.1), for every $z \in \mathbb{D}(2/3) \setminus \{0\}$, we have

$$\begin{aligned} L_{f, \mathbb{D}(z, |z|/2)} &\geq \lim_{w \rightarrow z} \frac{\delta_V(f(z), f(w))}{d_{\mathbb{D}(z, |z|/2)}(z, w)} \\ &\geq \lim_{w \rightarrow z} \frac{f^\#(z)}{(|z|/2)/(|z|/2)^2 - |w - z|^2} = \frac{|z|}{2} \cdot f^\#(z). \end{aligned}$$

Hence f is exceptional in Julia’s sense if $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} < \infty$.

If $\limsup_{z \rightarrow 0} L_{f, \mathbb{D}(z, |z|/2)} = \infty$, then there are sequences (z_n) , (z'_n) , and (z''_n) in $\mathbb{C} \setminus \{0\}$ such that $\lim_{n \rightarrow \infty} z_n = 0$, that $\lim_{n \rightarrow \infty} L_{f, \mathbb{D}(z_n, |z_n|/2)} = \infty$, and that for every $n \in \mathbb{N}$, the points z'_n, z''_n are in $\mathbb{D}(z_n, |z_n|/2)$ and distinct and satisfies

$$(6.1) \quad \frac{1}{2} L_{f, \mathbb{D}(z_n, |z_n|/2)} \leq \frac{\delta_V(f(z'_n), f(z''_n))}{d_{\mathbb{D}(z_n, |z_n|/2)}(z'_n, z''_n)}.$$

For every $n \in \mathbb{N}$, there is also $w_n \in \mathbb{D}(z_n, |z_n|/2)$ on the Euclidean line segment joining z'_n, z''_n such that

$$(6.2) \quad \delta_V(f(z'_n), f(z''_n)) \leq f^\#(w_n)|z'_n - z''_n|.$$

For every $n \in \mathbb{N}$, we have $|z_n|/2 \leq |w_n| \leq 3|z_n|/2$, and then by (6.1), (6.2), and (2.1),

$$\frac{1}{2} L_{f, \mathbb{D}(z_n, |z_n|/2)} \leq \frac{f^\#(w_n)|z'_n - z''_n|}{d_{\mathbb{D}(z_n, |z_n|/2)}(z'_n, z''_n)} \leq \frac{f^\#(w_n)}{1/(|z_n|/2)} \leq |w_n| f^\#(w_n)$$

so

$$\limsup_{z \rightarrow \infty} |z| f^\#(z) \geq \limsup_{n \rightarrow \infty} |w_n| f^\#(w_n) \geq \frac{1}{2} \limsup_{n \rightarrow \infty} L_{f, \mathbb{D}(z_n, |z_n|/2)} = \infty.$$

Hence f is not exceptional in Julia’s sense.

Acknowledgement. The author thanks the referee for a careful scrutiny, which helped us improve this article. The author also thanks Professor Katsutoshi Yamanoi for discussions on the Kobayashi hyperbolic geometry of complex spaces, and Professor Pekka Pankka for invaluable comments.

REFERENCES

- [1] BERTELOOT, F. Méthodes de changement d’échelles en analyse complexe, *Ann. Fac. Sci. Toulouse Math. (6)*, **15**, 3 (2006), 427–483.
- [2] BRODY, R. Compact manifolds in hyperbolicity, *Transactions of the American Mathematical Society*, **235** (1978), 213–219.
- [3] HOLOPAINEN, I. and PANKKA, P. A big Picard theorem for quasiregular mappings into manifolds with many ends, *Proc. Amer. Math. Soc.*, **133**, 4 (2005), 1143–1150 (electronic).
- [4] HOLOPAINEN, I. and RICKMAN, S. A Picard type theorem for quasiregular mappings of $\mathbf{R}^n$ into n -manifolds with many ends, *Rev. Mat. Iberoamericana*, **8**, 2 (1992), 131–148.

- [5] KOBAYASHI, S. *Hyperbolic complex spaces*, Vol. 318 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, Springer-Verlag, Berlin (1998).
- [6] KOBAYASHI, S. *Hyperbolic manifolds and holomorphic mappings*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition (2005), An introduction.
- [7] KWACK, M. H. Generalization of the big Picard theorem, *Annals of Mathematics* (1969), 9–22.
- [8] LEHTO, O. and VIRTANEN, K. I. On the behaviour of meromorphic functions in the neighbourhood of an isolated singularity, *Ann. Acad. Sci. Fenn. Ser. A. I.*, **1957**, 240 (1957), 9.
- [9] MARTY, F. Recherches sur la répartition des valeurs d’une fonction méromorphe, *Ann. Fac. Sci. Toulouse Sci. Math. Sci. Phys. (3)*, **23** (1931), 183–261.
- [10] OKUYAMA, Y. and PANKKA, P. Accumulation of periodic points for local uniformly quasiregular mappings, *RIMS Kôkyûroku Bessatsu*, **B43** (2013), 121–139.
- [11] OKUYAMA, Y. and PANKKA, P. Rescaling principle for isolated essential singularities of quasiregular mappings, *Proc. Amer. Math. Soc.* (to appear).
- [12] YAMANOI, K. Kobayashi hyperbolicity and higher dimensional Nevanlinna theory, *preprint* (2014).
- [13] ZALCMAN, L. A heuristic principle in complex function theory, *Amer. Math. Monthly*, **82**, 8 (1975), 813–817.

DIVISION OF MATHEMATICS, KYOTO INSTITUTE OF TECHNOLOGY, SAKYO-KU, KYOTO 606-8585, JAPAN

E-mail address: okuyama@kit.ac.jp