Absence of a quantum limit to charge diffusion in bad metals

Nandan Pakhira^{1, *} and Ross H. McKenzie^{1, †}

¹School of Mathematics and Physics, The University of Queensland, Brisbane, QLD 4072, Australia.

Good metals are characterised by diffusive transport of coherent quasi-particle states and the resistivity is much less than the Mott-Ioffe-Regel (MIR) limit, $\frac{ha}{a^2}$, where a is the lattice constant. In bad metals, such as many strongly correlated electron materials, the resistivity exceeds the Mott-Ioffe-Regel limit and the transport is incoherent in nature. Hartnoll, loosely motivated by holographic duality (AdS/CFT correspondence) in string theory, recently proposed a lower bound to the charge diffusion constant, $D \geq \hbar v_F^2/(k_B T)$, in the incoherent regime of transport, where v_F is the Fermi velocity and T the temperature. Using dynamical mean field theory (DMFT) we calculate the charge diffusion constant in a single band Hubbard model at half filling. We show that in the strongly correlated regime the Hartnoll's bound is violated in the crossover region between the coherent Fermi liquid region and the incoherent (bad metal) local moment region. The violation occurs even when the bare Fermi velocity v_F is replaced by its low temperature renormalised value, v_F^* . The bound is satisfied at all temperatures in the weakly and moderately correlated systems as well as in strongly correlated systems in the high temperature region where the resistivity is close to linear in temperature. Our calculated charge diffusion constant, in the incoherent regime of transport, also strongly violates a proposed quantum limit of spin diffusion, $D_s \sim 1.3\hbar/m$, where m is the fermion mass, experimentally observed and theoretically calculated in a cold degenerate Fermi gas in the unitary limit of scattering.

PACS numbers: 71.27.+a, 05.60.Gg, 67.10.Jn

I. INTRODUCTION

Good metals like copper and gold are characterised by high optical reflectivity, electrical and thermal conductivity. The transport in these systems can be characterised by diffusive transport of coherent quasi-particle states, where the mean-free path is much larger than the lattice constant. The low temperature resistivity in good metals is well within the Mott-Ioffe-Regel (MIR) limit, $\frac{ha}{c^2} \sim 250 \ \mu\Omega - \text{cm}$, where a is the lattice constant. However, in a large class of strongly correlated systems like 3d-transition metal oxide compounds and most notably in the strange metal regime of doped cuprates (high T_c superconductors) at optimal doping the resistivity far exceeds the MIR limit¹ and hence cannot be characterised by diffusive transport of coherent quasi-particle states in the limit of weak scattering. Other signatures of a bad metal include a thermopower of order k_B/e , the absence of a Drude peak in the optical conductivity, and a nonmonotonic temperature dependence of the Hall constant and thermopower.²⁻⁴

There have been a range of theoretical attempts to understand the incoherent regime of transport, especially for the strange metal phase of doped cuprates (high T_c superconductors) at optimal doping. Recently, there is a string theory based approach to understand transport in the incoherent regime^{5,6}. String theory, originally proposed as a possible theory for quantum gravity, is mathematically consistent but yet has no experimental verification. In the following paragraph we briefly describe how a string theory based approach has been proposed to describe transport in condensed matter systems.

Maldacena conjectured⁷ that the large N limit of certain supersymmetric conformal field theory (CFT) has correspondence to super gravity in anti-de Sitter spaces in higher dimension. This is known as the AdS/CFTcorrespondence or gauge/gravity duality. The most famous example of AdS/CFT correspondence states that IIB string theory in the product space $AdS_5 \times S^5$ is dual to large N_c limit of $\mathcal{N} = 4$ supersymmetric $SU(N_c)$ Yang-Mills theory on the four dimensional boundary. Further AdS/CFT correspondences relate fluid dynamics to event horizon dynamics of a black hole in anti-de Sitter space. In the hydrodynamic regime (long wavelength limit) of the correspondence, Einstein's equations of general relativity reduce to the Navier-Stokes equation for fluid mechanics. Classical fluids are characterised by transport coefficients such as shear viscosity and diffusion constant. Using the AdS/CFT correspondence Kovtun et al.⁸ calculated the ratio, η/s , of the shear viscosity (η) and the entropy density (s) and proposed a lower bound $\frac{\eta}{s} \geq \frac{\hbar}{4\pi k_{\rm P}}$. Such a bound is respected in classical fluids like water, the quark-gluon plasma (QGP) created in the relativistic heavy ion collider $(RHIC)^9$, and in experiments on cold degenerate Fermi gases in the unitary limit¹⁰. However, some violations of this bound have been reported¹¹. Inspired by the bound on the viscosity and also using the AdS/CFT correspondence Hartnoll recently $proposed^{12}$ a lower bound for the diffusion constant.

$$D \gtrsim \mathcal{D}_H \equiv \frac{\hbar v_F^2}{k_B T},\tag{1}$$

in the incoherent regime of transport in strongly correlated electron systems. But, except near quantum critical point, condensed matter systems are probably neither relativistic nor conformal¹³. So, this proposal needs to be tested against model based calculations. The temperature dependent diffusion constant, D(T), is related to the temperature dependent conductivity, $\sigma(T)$, through the Nernst-Einstein relation

$$\sigma(T) = e^2 \frac{\partial n}{\partial \mu} D(T).$$
 (2)

where $\kappa_e(T) = \frac{\partial n}{\partial \mu}$ is the *charge compressibility*. For completeness we give a derivation of this relation in the Appendix. Because of the above relation knowledge of the charge diffusion constant, D(T), may help us to better understand the electrical conductivity, $\sigma(T)$.

Experiments on cold degenerate Fermi gases in the unitary scattering limit show a quantum limit to the spin diffusion constant^{14,15}, $D_s \simeq 1.3\hbar/m$. This bound is also supported by theoretical calculations¹⁶. However, experiments on a two-dimensional Fermi gas found a value D_s that was more than two orders of magnitude smaller than the proposed bound¹⁷. But spin diffusion in charge neutral systems such as cold atomic gases has no obvious relation to charge diffusion in charged quantum fluids such as strongly correlated electron systems. For example, in a Mott insulator the charge diffusion constant is zero but the spin diffusion constant is non-zero. Hence, it is conceivable that in a bad metallic phase close to the Mott insulator that the charge diffusion constant is much smaller than the spin diffusion constant. In the present article we do a model based calculation of the charge diffusion constant in a Hubbard model, and explore the possible existence of a lower bound to the charge diffusion constant and its possible relation to spin diffusion in atomic gases.

The organization of the paper is as follows. In Sec. II we introduce the single band Hubbard model and its solution under single site dynamical mean field theory (DMFT). We also briefly describe DMFT self-consistency using iterated perturbation theory (IPT) as a solver for the impurity problem arising under single site DMFT. In Sec. III we briefly introduce calculation of transport and thermodynamic quantities under the single site DMFT approximation. Then in Sec. IV we show our results for a single band Hubbard model on the Bethe lattice at half-filling. We find clear violation of Hartnoll's proposed bound, even when the bare Fermi velocity v_F is replaced by the low-temperature renormalised velocity, v_F^* . Finally, in Sec. V we conclude and briefly consider how relaxing some of our assumptions may modify the results.

II. MODEL BASED CALCULATIONS

We consider the single band Hubbard model with nearest neighbor hopping, described by the Hamiltonian

$$H = -t \sum_{\langle i,j \rangle,\sigma} (c_{i\sigma}^{\dagger} c_{j\sigma} + H.c.) - \mu \sum_{i,\sigma} n_{i\sigma} + U \sum_{i} n_{i\uparrow} n_{i\downarrow} (3)$$

where $n_{i\sigma} = c^{\dagger}_{i\sigma}c_{i\sigma}$, t is the hopping amplitude μ is the chemical potential and U is Coulomb repulsion for a doubly occupied site. This is probably the simplest model which incorporates nontrivial strong correlation effects. But this model has an exact solution only in one dimension and study of this model in higher dimension involves various approximations. Static mean field descriptions like the Hartree-Fock decomposition of the quartic term $Un_{i\uparrow}n_{i\downarrow} \simeq U\langle n_{i\uparrow}\rangle n_{i\downarrow} + Un_{i\uparrow}\langle n_{i\downarrow}\rangle$ only shifts the local chemical potential. Because of the complete neglect of the quantum fluctuations this approximation does not generate any new energy scale (e.g. the Fermi liquid coherence scale) which can become relevant at low temperature regions. However, as in the case of classical mean field theory for the Ising model, in the limit of of large dimension, $d \to \infty$, (or large connectivity z) the model reduces to an effective single impurity model provided the scaling $t \to t^*/\sqrt{2d}$ is made on a d-dimensional hyper-cubic lattice¹⁸. Under this approximation we neglect all spatial fluctuations yet fully retain quantum dynamics for the single site. The self-energy $\Sigma(\omega)$ only depends on frequency and not wave vector. This is known as the dynamical mean field theory 19 (DMFT). It has been found DMFT gives a good description of the Mott metal-insulator transition with increasing correlation strength, U, and the crossover from a Fermi liquid to bad metal with increasing temperature². Furthermore, DMFT has been found to give a quantitative description of the temperature dependence of the resistivity²⁰ and the frequency dependent optical conductivity²¹ for organic charge transfer salts that are described by a two dimensional Hubbard model at half-filling²². Combining DMFT with electronic structure calculations based on density functional theory has given an excellent description of properties of a diverse range of transition metal and rare earth compounds²³.

A. Dynamical mean field theory

As a consequence of the scaling, $t \to t^*/\sqrt{2d}$, all the self energy diagrams, arising under skeletal graph expansion of the irreducible self energy and involving non local Green's functions vanishes in the limit $d \to \infty$. Then the self energy becomes local and involves only the local Green's function. The lattice problem for the Hubbard model then can be mapped onto an effective single impurity Anderson model¹⁹:

$$H_{imp} = \sum_{l,\sigma} (\tilde{\epsilon}_l - \mu) c_{l\sigma}^{\dagger} c_{l\sigma} + \sum_{l,\sigma} (V_l c_{l\sigma}^{\dagger} d_{0\sigma} + H.c.) -\mu \sum_{\sigma} n_{d0\sigma} + U n_{d0\uparrow} n_{d0\downarrow}, \qquad (4)$$

where $n_{d0\sigma} = d^{\dagger}_{0\sigma} d_{0\sigma}$. The operators $d^{\dagger}_{0\sigma}$ and $d_{0\sigma}$ characterise a given site i = 0 and $\{c^{\dagger}_{l\sigma}, c_{l\sigma}\}$ characterise the effective bath arising from electrons at all other sites. $\tilde{\epsilon}_l$

and V_l are effective parameters characterising the dispersion of the bath and its coupling to the local site. $\tilde{\epsilon}_l$ and V_l or equivalently the bath Green's function, given by $G_0(\omega)$,

$$G_0^{-1}(\omega) = \omega + \mu - \int_{-\infty}^{+\infty} \frac{\Delta(\epsilon) \ d\epsilon}{\omega + \mu - \epsilon}$$
$$\Delta(\epsilon) = \sum_{l\sigma} V_l^2 \delta(\epsilon - \tilde{\epsilon}_l) \tag{5}$$

can be calculated self consistently by solving the impurity problem iteratively. The solution of the impurity problem is the toughest part and usually involves use of numerical methods such as quantum Monte Carlo (QMC), exact diagonalization (ED), or the numerical renormalization group (NRG).

We use iterated perturbation theory $(IPT)^{24,25}$ as it is easy to implement, computationally cheap, and captures the essential physics in the parameter regime we are interested in, $U < 0.8U_c$ where U_c is the critical value of ${\cal U}$ at which the Mott metal-insulator transition occurs. For example, Bulla²⁶ showed that for the Bethe lattice at half-filling the results of IPT and NRG are similar except extremely close to the Mott transition. Indeed in the proximity of the Mott transition, Terletska *et al.*²⁷ found that the temperature dependent resistivity calculated from IPT was in agreement with that found by continuous time QMC (CT-QMC). Also, recently Arsenault $et \ al.^{28}$ showed that for lattices with a van Hove singularity in the density of states (DOS), even in the proximity of the Mott transition IPT with a modified self-consistency condition matches with results from CT-QMC. So, for the single band Hubbard model results from the IPT are generic in nature. In the next section we review DMFT self consistency using IPT.

B. Iterated Perturbation Theory

The iterated perturbation theory (IPT) is a semianalytical method. The irreducible self-energy in IPT is approximated using second order polarization bubble involving bath Green's function, $G_0(\omega)$. The self-energy under this approximation can be shown (using moment expansion) to smoothly interpolates between the atomic limit t = 0 and the weak coupling limit $U \rightarrow 0$. In the following paragraph we elaborate DMFT self-consistency method using IPT as impurity solver. We work with real, not imaginary, frequencies and so no analytic continuation is necessary.

(i) For a given lattice density of states $N_0(\epsilon)$ and self energy $\Sigma(\omega)$ the *local* Green's function is given by

$$G(\omega) = \int_{-\infty}^{+\infty} \frac{N_0(\epsilon)d\epsilon}{\omega^+ + \mu - \Sigma(\omega^+) - \epsilon},$$
 (6)

where μ is the *local* chemical potential.

(ii) From knowledge of the *local* Green's function $G_{\rm loc}(\omega)$

we can calculate the *bath hybridization* function, $\Delta(\omega)$ by using-

$$\Delta(\omega) = \omega^{+} + \mu - \Sigma(\omega) - G^{-1}(\omega).$$
(7)

(iii) Subsequently using *bath hybridization* we can calculate the *bath Green's function* as

$$G_0(\omega) = \frac{1}{\omega + \tilde{\mu}_0 - \Delta(\omega)}.$$
(8)

The parameter $\tilde{\mu}_0 = \mu - Un$ is the bath chemical potential and it vanishes at half filling for the particle-hole symmetric case, which we consider in the present study. (iv) The fully interacting Green's function can be calculated using the Dyson's equation

$$G(\omega) = \frac{1}{G_0^{-1}(\omega) - \tilde{\mu}_0 + \mu - \Sigma(\omega)}.$$
 (9)

(v) The new self-energy can be calculated following the IPT ansatz 25 as

$$\Sigma(\omega) = Un + \frac{A\Sigma^{(2)}(\omega)}{1 - B\Sigma^{(2)}(\omega)}$$
(10)

where,

$$A = \frac{n(1-n)}{n_0(1-n_0)} ; \qquad B = \frac{U(1-n) - \mu + \mu_0}{n_0(1-n_0)U^2}$$
(11)

and

$$n = -\frac{1}{\pi} \int_{-\infty}^{+\infty} d\omega \ n_F(\omega) \ \text{Im}[G(\omega^+)], \qquad (12)$$

$$n_0 = -\frac{1}{\pi} \int_{-\infty}^{+\infty} d\omega \ n_F(\omega) \ \mathrm{Im}[G_0(\omega^+)] \qquad (13)$$

are the *local* and *bath* particle numbers, respectively. $\Sigma^{(2)}(\omega)$ is the self energy from second order perturbation theory and is given by

$$\Sigma^{(2)}(\omega) = U^2 \int_{-\infty}^{+\infty} \prod_{i=1}^{3} \left(d\epsilon_i \rho_0(\epsilon_i) \right) \left[\frac{n_F(-\epsilon_1) n_F(\epsilon_2) n_F(-\epsilon_3)}{\omega + i\eta - \epsilon_1 + \epsilon_2 - \epsilon_3} + \frac{n_F(\epsilon_1) n_F(-\epsilon_2) n_F(\epsilon_3)}{\omega + i\eta - \epsilon_1 + \epsilon_2 - \epsilon_3} \right]$$
(14)

where $\rho_0(\omega) = -\frac{1}{\pi} \text{Im}[G_0(\omega^+)]$ and $\eta \to 0^+$. We iterate (i) - (v) until the desired self-consistency in self-energy and other physical quantities are achieved. Hence, we focus solely on the case of half-filling (n = 1). Due to particle-hole symmetry $\mu = \frac{U}{2}$ for all U and T. This speeds up computation significantly, as it is not necessary to self-consistently determine μ from Eq. (12)

C. Bethe lattice

We choose a Bethe lattice (Cayley tree) because it makes computation even faster because the local Green's function, $G(\omega)$, has an exact analytical form. Particlehole symmetry also simplifies the calculations. The Bethe lattice produces qualitatively similar results to the hyper-cubic lattice¹⁹ and lower dimensional Hubbard models^{4,21}. In the limit of infinite coordination number $(z \to \infty)$, the density of states has *semi-circular* form²⁹

$$N_0(\epsilon) = \frac{2}{\pi W^2} \sqrt{W^2 - \epsilon^2} \Theta(W - |\epsilon|)$$
(15)

where $\Theta(x)$ is the familiar unit step function, $W = 2t^*$ is the half-band width and the hopping amplitude in this case is scaled as $t \to t^*/\sqrt{z}$. Most importantly the *local* Green's function has the exact analytical form

:

$$G(\omega) = \frac{2}{W^2} \left[\zeta - \sqrt{\zeta^2 - W^2} \right], \qquad (16)$$

$$\zeta(\omega) \equiv \omega + i\eta + \mu - \Sigma(\omega). \tag{17}$$

It can be easily verified that in this case the bath hybridization function, $\Delta(\omega) = \frac{W^2}{4}G(\omega) \equiv t^{*2}G(\omega)$, is proportional to the *local* Green's function.

III. TRANSPORT PROPERTIES

Using the self-consistent self energy we can calculate various quantities like dc conductivity, charge compressibility, and diffusivity.

A. dc Conductivity

In the limit of $d \to \infty$ all vertex corrections to twobody correlation functions drop out³⁰ and the temperature dependent dc conductivity, $\sigma(T)$, can be calculated using the simple polarization bubble as^{19,31}

$$\sigma(T) = \frac{\pi e^2}{\hbar} \frac{1}{\nu} \int_{-\infty}^{+\infty} d\epsilon \, \Phi_{xx}(\epsilon) \int_{-\infty}^{+\infty} d\omega \left(-\frac{\partial n_F(\omega)}{\partial \omega} \right) A^2(\omega, \epsilon) \mathbb{1} 8$$

where $\nu = a^d$ is the volume of the unit cell of a *d*dimensional hyper-cubic lattice with lattice constant *a*,

$$A(\omega, \epsilon) = -\frac{1}{\pi} \operatorname{Im} \left[\frac{1}{\omega + \mu - \Sigma(\omega) - \epsilon} \right], \quad (19)$$

$$n_F(\omega) = \frac{1}{e^{\beta\omega} + 1} \tag{20}$$

are the spectral density and Fermi function, respectively and

$$\Phi_{xx}(\epsilon) = \frac{1}{N} \sum_{\mathbf{k}} \left(\frac{\partial \epsilon_{\mathbf{k}}}{\partial k_x} \right)^2 \delta(\epsilon - \epsilon_{\mathbf{k}})$$
(21)

is the transport density of states. N is the number of lattice sites.

Because of its tree like structure the Bethe lattice has no loop and no energy dispersion relation in **k**. But, by invoking the f-sum rule it can be shown that³²⁻³⁴

$$\Phi_{xx}(\epsilon) = \frac{1}{3d} (W^2 - \epsilon^2) N_0(\epsilon), \qquad (22)$$

is the correct transport density of states in the limit of $d \to \infty$. It is interesting to mention that for a Bethe lattice with coordination number z the connectivity K = z - 1 while that for the hyper-cubic lattice is 2d. So, in the limit of large coordination number we can take the connectivity to be equal to 2d and we can always do the mapping $z \leftrightarrow 2d$.

B. Charge Compressibility

The inverse of the charge compressibility can be interpreted as the energy cost to add or remove a particle from a system. For the non-interacting system (U = 0) at zero temperature (T = 0), $\kappa_e = N_0(E_F)$, where $N_0(E_F)$ is the density of states at the Fermi level.

In a general many-body system the local particle number is given by

$$n = \frac{1}{\nu} \int_{-\infty}^{+\infty} d\omega \ n_F(\omega) \sum_{\mathbf{k}} A(\mathbf{k}, \omega), \tag{23}$$

where, the spectral function is

$$A(\mathbf{k},\omega) = -\frac{1}{\pi} \operatorname{Im} \left[\frac{1}{\omega + \mu - \epsilon_{\mathbf{k}} - \Sigma_{\mathbf{k}}(\omega)} \right].$$
(24)

The self-energy, $\Sigma_{\mathbf{k}}(\omega)$, in the limit of $d \to \infty$ is independent of wave vector \mathbf{k} and is given by $\Sigma(\omega)$. Hence, differentiating with respect to μ , the charge compressibility, $\kappa_e(T) = \frac{\partial n}{\partial \mu}$, under the DMFT approximation is given by

$$\kappa_e(T) = \frac{1}{\pi} \operatorname{Im} \int_{-\infty}^{+\infty} d\omega \ n_F(\omega) \left(1 - \frac{\partial \Sigma(\omega)}{\partial \mu} \right) \\ \times \int_{-\infty}^{+\infty} \frac{N_0(\epsilon) \ d\epsilon}{(\omega + \mu - \Sigma(\omega) - \epsilon)^2}.$$
(25)

The effect of the derivative $\frac{\partial \Sigma(\omega)}{\partial \mu}$ on the charge compressibility of a Fermi liquid was discussed previously by Luttinger³⁶ and by Hotta and Fujimoto³⁷. Here it does not have a closed analytical form and we evaluate it beginning with the IPT expression (10). The charge compressibility $\kappa_e(T)$ is then given by

$$\kappa_e(T) = \frac{\tilde{J} + \tilde{K}}{1 + U(\tilde{J} + \tilde{K})},\tag{26}$$

where the \tilde{J} term in the denominator is associated with the Hartree term in the self energy (10). For the Bethe lattice

$$\tilde{J} = -\frac{1}{\pi} \operatorname{Im} \int_{-\infty}^{+\infty} d\omega \, n_F(\omega) \left[2 - \frac{2\zeta}{\sqrt{\zeta^2 - 1}} \right]$$
(27)
$$\tilde{K} = \frac{1}{\pi} \operatorname{Im} \int_{-\infty}^{+\infty} d\omega \, n_F(\omega) \left[2 - \frac{2\zeta}{\sqrt{\zeta^2 - 1}} \right] \frac{\partial \tilde{\Sigma}_2(\omega)}{\partial \mu}$$
(28)

and $\zeta(\omega)$ is given by Eq. (17). If we define $\frac{\partial \rho_{\Sigma}(\omega)}{\partial \mu} = -\frac{1}{\pi} \text{Im} \frac{\partial \tilde{\Sigma}_2(\omega)}{\partial \mu}$ then

$$\frac{\partial \rho_{\Sigma}(\omega)}{\partial \mu} = \int d\epsilon_1 d\epsilon_2 \left[2 \frac{\partial \rho_{\mathcal{G}}(\epsilon_1)}{\partial \mu} \rho_{\mathcal{G}}(\omega - \epsilon_1 + \epsilon_2) \rho_{\mathcal{G}}(\epsilon_2) + \rho_{\mathcal{G}}(\epsilon_1) \rho_{\mathcal{G}}(\omega - \epsilon_1 + \epsilon_2) \frac{\partial \rho_{\mathcal{G}}(\epsilon_2)}{\partial \mu} \right] \times \left[n_F(-\epsilon_1) n_F(-\omega + \epsilon_1 - \epsilon_2) n_F(\epsilon_2) + n_F(\epsilon_1) n_F(\omega - \epsilon_1 + \epsilon_2) n_F(-\epsilon_2) \right], \quad (29)$$

where $\rho_{\mathcal{G}}(\omega) = -\frac{1}{\pi} \text{Im} G_0(\omega)$ and for Bethe lattice

$$\frac{\partial \rho_{\mathcal{G}}(\omega)}{\partial \mu} = -\frac{1}{\pi} \operatorname{Im} \frac{2 - \frac{2\zeta}{\sqrt{\zeta^2 - 1}}}{4(\omega + \mu_0 - \Delta(\omega))^2}.$$
 (30)

The expression in Eq. (29) is calculated by using standard FFT routine and the real part of $\frac{\partial \tilde{\Sigma}_2(\omega)}{\partial \mu}$ can be calculated using Hilbert transform. We note in passing that we find for most parameter regimes that the expression (26) is dominated by the \tilde{J} terms and the \tilde{K} terms involve only a small correction.

C. Diffusivity

As mentioned earlier the diffusivity, D(T) can be calculated using the Nernst-Einstein relation in Eq. (2). To compare to the limit of diffusion constant, proposed by Hartnoll, we need to find the Fermi velocity v_F . First, one has to decide whether this should be the bare Fermi velocity, i.e. the band structure value, or a renormalised value v_F^* associated with a low temperature Fermi liquid state. In that case, $v_F^* = Zv_F$ where Z is the quasiparticle renormalization factor which can be calculated from the self energy, $\Sigma(\omega) = \Sigma_R(\omega) + i\Sigma_I(\omega)$:

$$Z = \left(1 - \frac{\partial \Sigma_R(\omega)}{\partial \omega}\Big|_{\omega \to 0}\right)^{-1}.$$
 (31)

The Hartnoll bound, \mathcal{D}_H then gets renormalized to $\mathcal{D}_H^* = Z^2 \mathcal{D}_H$. Note that as the Mott transition is approached this decreases the lower bound by several orders of magnitude, making it harder to violate. It is not completely clear to us from the arguments of Hartnoll whether one should use v_F or v_F^* , particularly as he is concerned with incoherent transport, i.e., outside the Fermi liquid

regime. Here, we use the latter but note that this choice makes the bound much less stringent.

Second, there is the issue of how to evaluate the Fermi velocity in the DMFT approximation, in the limit of infinite dimensionality, $d \to \infty$. Since, $v_{\mathbf{k}}^2 = \left(\frac{\partial \epsilon_{\mathbf{k}}}{\partial \mathbf{k}}\right)^2$ appears in the expression for *transport density of states* in Eq. (21) we define

$$\hbar v_F^2 = \frac{1}{\hbar} \frac{\Phi_{xx}(\epsilon = 0)}{N_0(\epsilon = 0)}$$
(32)

in the limit of $d \to \infty$. This definition of $\hbar v_F^2$ gives the correct Fermi velocity³¹ for the hyper-cubic lattice in the limit of $d \to \infty$. Also, by pure dimensional analysis for any lattice structure $\hbar v_F = \lambda W a$ (λ being a numerical constant of order one for a given lattice). Hartnoll's proposed quantum bound for diffusion constant on the Bethe lattice is then given by

$$\mathcal{D}_H = \frac{W^2 a^2}{3d\hbar k_B T}.$$
(33)

Including the renormalisation of the Fermi velocity the dimensionless scaled diffusivity is then given by

$$\frac{D(T)}{\mathcal{D}_{H}^{*}} = \pi \left(\frac{k_{B}T}{W}\right) \frac{1}{Z^{2}\tilde{\kappa}_{e}(T)} \int_{-W}^{+W} d\epsilon \int_{-\infty}^{+\infty} d\omega (W^{2} - \epsilon^{2}) N(\epsilon) \times A^{2}(\epsilon, \omega) \left(-\frac{\partial n_{F}(\omega)}{\partial \omega}\right), \quad (34)$$

where $\tilde{\kappa}_e(T) = \frac{\partial \tilde{n}}{\partial \tilde{\mu}}$ is the dimensionless charge compressibility and $\mu = \tilde{\mu}W$, $\tilde{n} = n\nu$. The advantage of calculating scaled diffusivity is that it does not depend on universal constants such as \hbar or material dependent constants such as the lattice constant, a, and the unit cell volume, ν , and the temperature appears only as a dimensionless scaled quantity.

We now turn to comparison with the proposed bound for the spin diffusion constant. In a similar spirit we use $\frac{1}{m} = \frac{1}{\hbar^2} \frac{\partial^2 \epsilon_k}{\partial k_x^2}$ as a generalized definition for inverse mass. Then for the hyper-cubic lattice we get $\frac{Wa^2}{d\hbar^2}$ as an effective inverse mass averaged over the Fermi surface at half-filling. If we take this to be same in the Bethe lattice as well then we will have

$$\mathcal{D}_A = \frac{\alpha W a^2}{d\hbar} \tag{35}$$

with $\alpha = 1.3$ a dimensionless constant.

In an interacting system the bare mass m gets renormalized to an effective mass $m^* = m/Z$ where Z is the quasi-particle weight. Thus the bound \mathcal{D}_A will get renormalized to

$$\mathcal{D}_A^* = \alpha \frac{\hbar}{m^*} = Z \mathcal{D}_A \tag{36}$$

The scaled diffusivity in this case is then given by

$$\frac{D(T)}{\mathcal{D}_{A}^{*}} = \frac{\pi}{3\alpha Z} \frac{1}{\tilde{\kappa}_{e}(T)} \int_{-W}^{+W} d\epsilon \int_{-\infty}^{+\infty} d\omega (W^{2} - \epsilon^{2}) N(\epsilon) \times A^{2}(\epsilon, \omega) \left(-\frac{\partial n_{F}(\omega)}{\partial \omega}\right).$$
(37)

IV. RESULTS

We consider the case of half filling, n = 1, *i.e.* each site on the average is occupied by one electron. We study spectral and transport properties as a function of correlation strength U and temperature, T (enters as $k_B T$ with dimension of energy). Henceforth, unless stated otherwise, all the energy scales will be measured in units of half-bandwidth, W.

A. Spectral function

In Fig. 1 we show the evolution of the spectral function,

$$A_d(\omega) = -\frac{1}{\pi} \operatorname{Im} \left[G(\omega^+) \right], \qquad (38)$$

as a function of U and T. Similar results has been obtained previously by other authors 24 . For completeness we show these results here because they illustrate the essential physics (the destruction of quasi-particles) behind violation of the MIR limit and Hartnoll's bound. In panel (a) of Fig. 1 we show the spectral function for a weakly correlated system U = 0.5. The spectral function is dominated by a broad central peak and a very small smearing of the non-interacting (U = 0) band edges at $\omega = \pm 1$. The integrated spectral weight is dominated by the contribution from the central peak and $A_d(0) \simeq 2/\pi$, as in the non-interacting case. At finite temperature, due to particle-hole excitations across the Fermi surface, the spectral weight at the Fermi energy $\omega = 0$ gets transferred to finite frequency but the central peak still remains intact.

As we increase the correlation strength (U = 1) side bands develop on either side of the central peak as shown in the panel (b) of Fig. 1. The side bands eventually develop into high energy Hubbard bands at $\omega = \pm \frac{U}{2}$ as shown in panel (c) of Fig. 1 for U = 2. The Hubbard bands are well separated from the central peak which arises due to Kondo resonance effects in the effective single impurity Anderson model¹⁹. The width and height of the Kondo resonance is controlled by the effective Kondo temperature T_K . Since, $T_K \sim W \exp(-\Gamma_{\text{eff}}/U)$, where $\Gamma_{\rm eff}$ is an effective hybridization strength, the width of the Kondo resonance decreases while its height increases with increasing U as shown in panel (d) for U = 2.5. For very large $U > U_c$ the Kondo resonance gets completely killed and we enter into the Mott insulating state characterised by fully gapped spectral function at the Fermi

energy. The numerical value of U_c depends on the numerical technique that one uses and $U_c \simeq 3.3 - 3.4$ for the IPT based impurity solver^{24,26}.

At finite temperature in the moderately correlated regime like U = 1.0 the central peak, despite getting broadened, remains intact even for temperatures as high as $T \sim W$. The side bands thermally broaden out. For the strongly correlated regime of U = 2 and U = 2.5the central peak (quasi-particle peak) as well as the integrated spectral weight under it decreases with increasing temperature and eventually the central peak gets completely destroyed for temperatures $T \gg T_K$. This corresponds to the finite temperature crossover from the strong coupling regime into the local moment regime of the effective Anderson impurity model. The crossover region becomes increasingly sharp as evident in panel (d), which corresponds to the fragile nature of the quasiparticle state.



FIG. 1: (Color online) Energy dependent spectral function, $A_d(\omega)$, for various values of the interaction strength U and temperature T. Panel (a) : weakly correlated regime. (Green) dashed line is the density of states for the non-interacting case. Panel (b) : moderately correlated regime. Panel (c) and (d) : strongly correlated regime. With increasing U, the integrated spectral weight under central peak (Kondo resonance) gets transferred to high energy Hubbard bands which corresponds to destruction of quasi-particle states. In these cases, there is a temperature dependent crossover between strong coupling regime (Fermi liquid) and local moment regime (bad metals). All energies and temperatures are measured in units of W, the half-bandwidth.

B. Quasi-particle weight

In Fig. 2 we show the continuous decrease of the quasiparticle renormalization factor, Z, with increasing U. This also tracks the continuous destruction of coherent quasi-particle states.Comparison of Z against results from numerical renormalization group (NRG) based calculations by Bulla²⁶ validates the qualitative correctness of IPT based approach though Z begins to differs by 50% in the strong correlation regime (U = 2.5). We also note that for U = 2.5, Z > 0.2 and so in some sense for that regime the system is not extremely correlated. Yet we will see that even in this regime Hartnoll's bound is violated. For comparison, in the doped Hubbard model on the square lattice (with U = 16t = 3.5W at 15% doping n = 0.85) DMFT gives $Z \simeq 0.2^4$.

C. Charge compressibility

In Fig. 2 we also show the evolution of the zero temperature charge compressibility, κ_e , as a function of correlation strength, U. The charge compressibility continuously goes to zero with increasing, U. As mentioned earlier, $1/\kappa_e$ can be thought of as the energy required to add/remove an electron to/from the systems. Hence it gets increasingly harder to add or remove an electron into the system as we increase U, *i.e.*, the system increasingly becomes incompressible and finally at $U = U_c$ the system becomes completely incompressible. Note that at U = 0, $\kappa_e = N_0(E_F) = 2/\pi$, as it should be. A similar decrease in charge compressibility with increasing Uwas observed in exact diagonalization calculations for the Hubbard model on the triangular lattice at half-filling³⁵.



FIG. 2: (Color online) The zero-temperature quasi-particle renormalization factor Z and charge compressibility κ_e as a function of U. The system becomes increasingly incompressible with increasing U and the quasi-particle weight Z smoothly decreases, as the transition to the Mott insulator is approached at $U_c \simeq 3.4^{26}$. U is measured in units of W.

In Fig. 3 we show the temperature dependence of the charge compressibility for a range of values of U. In the

non-interacting case (U = 0) we can show that

$$\kappa_e(T) = \int_{-\infty}^{+\infty} \left(-\frac{\partial n_F(\omega)}{\partial \omega} \right) N_0(\omega) \tag{39}$$

This expression is similar to that for the Pauli spin susceptibility, $\chi(T)$, and the associated temperature dependence is shown in Figure 3 as a dashed line. The steady decrease in the charge compressibility withincreasing temperature is largely due to the broadening of the Fermi-Dirac distribution function.

Using standard integral expressions involving the Fermi function 38

$$\int_{-\infty}^{+\infty} H(\epsilon) n_F(\epsilon) d\epsilon = \int_{-\infty}^{\mu} H(\epsilon) d\epsilon + \frac{\pi^2}{6} \left(k_B T \right)^2 H'(\mu) + \cdot \left(40 \right)$$

the expression for the charge compressibility in Eq. (39) reduces to

$$\kappa_e(T) = N_0(\mu) + \frac{\pi^2}{6} (k_B T)^2 \left. \frac{d^2 N_0(\omega)}{d\omega^2} \right|_{\omega=\mu} + \cdots . (41)$$

For the Bethe lattice this reduces to

$$\kappa_e(T) \simeq \frac{2}{\pi W} \left[1 - \frac{\pi^2}{6} \left(\frac{k_B T}{W} \right)^2 + \cdots \right].$$
(42)

In the interacting case we expect in the Fermi liquid state

$$\kappa_e(T) \simeq \kappa_e(0) \left[1 - \frac{\delta}{Z^2} \left(\frac{k_B T}{W} \right)^2 + \cdots \right],$$
(43)

with $\delta \sim 1$. So, because of the thermal broadening effects, just as in the case of the spin susceptibility, the charge compressibility in the Fermi liquid state will decrease quadratically in temperature at low temperatures. This quadratic dependence is shown in Fig. 3 as dotted lines that have been fitted to the low temperature behaviour. This explains the rapid decrease of the charge compressibility at low temperatures in the coherent Fermi liquid state, because the temperature scale for the decrease is that of the coherence temperature which close to the Mott transition becomes very small^{2,19}.

D. Resistivity

In Fig. 4 we show the temperature dependence of the resistivity, $\rho(T)$, scaled by the Mott-Ioffe-Regel limit, $\rho_{MIR} = ha/e^2$, for various correlation strengths, U. In the weakly correlated regime (U = 0.5) the resistivity is well within the Mott-Ioffe-Regel limit in the entire temperature range up to W. Hence, the transport can be characterised by weak scattering of coherent quasiparticle states. At very low temperatures ($T < T_K \ll$



FIG. 3: (Color online) Temperature dependence of the charge compressibility, κ_e , for various correlation strengths U. The dashed line is the non-interacting (U = 0) case calculated using Eq. (39) and dotted lines are quadratic fits to the Fermi liquid form Eq. (43). The apparent kink-like behaviour for U = 2.5 is due to the sharp crossover between the Fermi liquid and the bad metal (local moments fixed point). Both T and U are measured in units of W.

W) the resistivity is proportional to T^2 as expected in the coherent Fermi-liquid regime. The T^2 behaviour is due to the fact that in the Fermi liquid regime the imaginary part of the self-energy, or equivalently the inverse of quasi-particle life time (τ_{qp}^{-1}) is proportional to T^2 (or ω^2 at T = 0). At high temperatures $(T \gg T_K)$, the resistivity is roughly linear in T and this corresponds to the incoherent (bad metal) regime of transport³.

As we increase U, the resistivity smoothly crosses the Mott-Ioffe-Regel limit and in the strongly correlated regime (U = 2.0 and above) the resistivity far exceeds the MIR limit. This is due to the sharp crossover from the strong coupling (Fermi liquid) regime to the local moment (bad metal) regime in the strong correlation regime and is consistent with the picture of fragile quasi-particles states in the strong correlation regime.

In elemental crystals one can distinguish metals and insulators by the temperature dependence of the resistivity. It is monotonically increasing (decreasing) with increasing temperature for metals (insulators). However, this criteria is unreliable for strongly correlated electron materials. For example, a non-monotonic temperature dependence of the resistivity in a bad metal is seen experimentally in a number of organic charge salts. (See for example the inset of Figure 2(a) of Ref. 39). Thus it is important to note that even though the derivative $d\rho(T)/dT$ changes sign for some curves in Fig. 4 there is no metal-insulator transition, i.e. all the curves are for the metallic phase. This is evident from the finite temperature spectral function, $A_d(\omega)$ at $\omega = 0$ and the non-zero charge compressibility κ_e .

It should also be stressed that for the given choice of U we are still far away from Mott transition at $U_c \simeq$

3.4. This is also evident from the relatively large quasiparticle weight $Z \sim 0.2$ even for U = 2.5 where the MIR limit is violated by a factor of 100. So, the transport in this bad metal phase is incoherent in nature.



FIG. 4: (Color online) Resistivity shows violation of the Mott-Ioffe-Regel limit ($\rho_{MIR} = ha/e^2$) with increasing U. This is consistent with the picture that transport becomes increasingly incoherent with increasing correlation effects. Both T and U are measured in units of W.

E. Diffusivity

Finally, using the Nernst-Einstein relation we calculate the charge diffusivity. In Fig. 5 we show the scaled diffusivity, $D(T)/D_H^*$, as a function of temperature (T) for various correlation strengths (U). In the weakly correlated regime (U = 0.5) and moderately correlated regime (U = 1.0 and U = 1.5) the scaled diffusivity satisfies Hartnoll's bound. However in the strongly correlated regime, U = 2.0 and above, the scaled diffusivity shows violation of Hartnoll's bound in the low temperature region. But at high temperatures $T \gg T_K$ the scaled diffusivity satisfies Hartnoll's bound. It is important to mention that the kink like behaviour at around $T \sim 0.05$ for U = 2.5 is closely related to the sharp crossover between the Fermi liquid fixed point and the local moment fixed point in the effective single impurity Anderson model. As we can see from Fig. 5 the violation of Hartnoll's bound in strongly correlated systems is in the crossover region between coherent (Fermi liquid) regime and incoherent (local moment) regime. The magnitude of violation increases with increasing U due to increased sharpness in crossover region. In the high temperature region, where the resistivity is roughly linear in T the scaled diffusivity is well above the Hartnoll's bound, provided one uses the renormalised Fermi velocity.



FIG. 5: (Color online) Scaled diffusivity shows violation of Hartnoll's bound in the strongly correlated incoherent regime of transport in the low temperature region. However, in strongly correlated systems at large temperatures $(T \gg T_K)$ and for weakly correlated systems at all temperatures the bound is respected. Kink like behaviour for U = 2.5 is due to sharp crossover between the Fermi liquid and the local moment regime. Both T and U are measured in units of W.

Finally, we compare the charge diffusivity to the quantum limit of the spin diffusion constant, $D_s \simeq 1.3\hbar/m$, experimentally observed¹⁴ and theoretically calculated¹⁶ in the degenerate Fermi gas in the unitary limit. In Fig. 6 we show the scaled diffusivity, $D(T)/D_A^*$, as a function of temperature for various U. The scaled diffusivity also violates the quantum limit of spin diffusion constant, $D_s \simeq 1.3\hbar/m$. The violation is severe in the strongly correlated regime. All the temperature dependence is ultimately due to inherent temperature dependence of self energy, $\Sigma(\omega)$.

It is important to mention that spin diffusion in charge neutral systems like the degenerate Fermi gas in the unitary limit has no clear relation to charge diffusion in electron liquids. In charged systems there are dynamical screening effects while such screening effects are not present in neutral atomic gases such as the strongly interacting degenerate Fermi gas at Feshbach resonance. Most interestingly in a Mott insulator the charge diffusion constant is zero while the spin diffusion constant is finite. The differences illustrate different mechanisms for charge and spin transport in strongly correlated systems and one should not necessarily expect any simple relationship between the spin and charge diffusion constants.

For a degenerate non-interacting Fermi gas in three dimensions the charge diffusion constant is given by $D = \frac{1}{3} \frac{\hbar}{m} k_F \ell$, where k_F is the Fermi wave vector and ℓ is the mean free path. In the weak scattering limit $k_F \ell \gg 1$ and $D \gg \frac{1}{3} \frac{\hbar}{m}$. So, just like the upper limit (MIR) of resistivity we can define lower limit for the charge diffusion constant $D_{lim} = \frac{1}{3} \frac{\hbar}{m}$. In the weakly interacting quasiparticle regime of transport the limit will be renormalized

to $D_{lim}^* = \frac{1}{3} \frac{\hbar}{m^*}$. The quantum spin diffusion limit in degenerate Fermi gas will roughly correspond to $k_F \ell \sim 4$ for charge diffusion in a condensed matter system and the diffusion will correspond to transport through coherent quasi-particle states.



FIG. 6: (Color online) Scaled diffusion constant for charge transport also violates quantum limit for spin diffusion constant in the incoherent regime of transport. Inset : Detailed plot near the origin shows small yet non-vanishing scaled diffusivity for U = 2.5. D_H/D_A traces out the temperature dependence, $1/3\alpha k_B T$, of the Hartnoll bound showing how for T < 0.3 it is larger than the proposed bound on the spin diffusion constant for cold atoms. Both T and U are measured in units of W.

V. CONCLUSIONS

We have studied the conductivity, charge compressibility, and charge diffusivity in a single band Hubbard model using single site dynamical mean field theory. The calculated resistivity far exceeds the MIR limit in the strong correlation regime. The transport in the weakly correlated region can be characterized by diffusive scattering of coherent quasi-particle states but in the strongly correlated bad metal state the transport is incoherent. The charge compressibility decreases with increasing U which corresponds to the fact that in the correlated regime, the energy cost to create a charge fluctuation increases with increasing U. Then using the Nernst-Einstein relation we calculated the charge diffusivity in the system. In the weakly and moderately correlated systems the scaled diffusivity respects Hartnoll's bound at all temperatures. However, in the strongly correlated systems the bound is violated in the crossover region between the coherent Fermi liquid regime and incoherent local moment regime. In the high temperature region $(T \gg T_K)$ particularly in the region where resistivity is roughly linear in T the bound is found to be respected for all interaction strengths.

We also compared the calculated charge diffusivity against the quantum limit of spin diffusion observed in the degenerate Fermi gas in the unitary limit. The calculated diffusivity strongly violates the quantum limit of spin diffusion in the incoherent regime. So, within the single site DMFT approximation we do not observe any quantum limit to charge diffusion in the strongly correlated incoherent regime.

Hartholl's proposed bound is based on the AdS/CFT correspondence and various conservation laws in fluids. But within single site DMFT approximation there is energy conservation but no momentum conservation at a given site. On the other hand, the spatial fluctuations neglected in single site DMFT can be systematically incorporated through other approximations such the dynamical cluster approximation $(DCA)^{41,42}$ in which momentum is conserved within the cluster, bath as well as at the boundary of the cluster should be able to address this issue. One might also consider how vertex corrections could modify the results. For a doped Hubbard model it was found in a 4 site DCA calculation that the vertex corrections to the optical conductivity were not significant, except very close to the Mott insulator⁴³. A study of the doped two-dimensional Hubbard model using a two-particle self-consistent approach found that vertex corrections altered the calculated resistivity by less than a factor of two^{44} .

As has been pointed out⁴⁰ the interacting many electron system in strongly correlated materials so far has no gravity description and hence the strongly coupled gauge theory has no dual description. To be more precise, holographic quantum systems associated with known gravity descriptions have no direct relation with strongly correlated electron systems. Furthermore, the AdS/CFT correspondence only describes conformally invariant field theories and except for one dimensional systems at a quantum critical point, it is not clear that strongly correlated electron systems are conformally invariant¹³.

Acknowledgments

We acknowledge helpful discussions with X. Deng, J. K. Freericks, S. Hartnoll, H. R. Krishnamurthy, B. J.

- * Electronic address: npakhira@gmail.com
- [†] Electronic address: r.mckenzie@uq.edu.au; URL: condensedconcepts.blogspot.com
- ¹ N. E. Hussey, K. Takenaka, and H. Takagi, Phil. Mag. 84, 2847 (2004).
- ² J. Merino and R. H. McKenzie, Phys. Rev. B **61**, 7996 (2000).
- ³ X. Deng *et al.*, Phys. Rev. Lett. **110**, 086401 (2013).
- ⁴ W. Xu, K. Haule, and G. Kotliar, Phys. Rev. Lett. **111**, 036401 (2013).
- ⁵ S. Sachdev and M. Müller, J. Phys : Condens. Matter 21, 164216 (2009).

Powell, T. V. Ramakrishnan, D. Tanaskovic, A. Taraphder, N. Vidhyadhiraja, and J. Vucicevic. This work was supported by an Australian Research Council Discovery Project grant.

APPENDIX: Derivation of the Nernst-Einstein relation

Fick's law for diffusion is given by

$$\mathbf{j}_m(\mathbf{r}) = -D(T)\boldsymbol{\nabla}n(\mathbf{r}) \tag{44}$$

where D(T) is the diffusion constant, $\mathbf{j}_m(\mathbf{r})$ is the mass current and $n(\mathbf{r})$ is the local particle number. On the other hand Ohm's law for electrical conductivity is given by

$$\mathbf{j}_e(\mathbf{r}) = \sigma(T)\mathbf{E}(\mathbf{r}) \tag{45}$$

where $\sigma(T)$ is the electrical conductivity, $\mathbf{j}_e(\mathbf{r})$ is the *electric current* and $\mathbf{E}(\mathbf{r})$ is the external electric field. We have

$$\mathbf{j}_e(\mathbf{r}) = e\mathbf{j}_m(\mathbf{r}) = -eD(T)\frac{\partial n}{\partial \mu}\boldsymbol{\nabla}\mu(\mathbf{r})$$
(46)

where $\mu(\mathbf{r}) = \mu_0 + e\phi(\mathbf{r})$ is the chemical potential in the presence of external field and μ_0 is that in the absence of external field and $\phi(\mathbf{r})$ is the electric potential. Then

$$\nabla \mu(\mathbf{r}) = e \nabla \phi(\mathbf{r}) = -e \mathbf{E}(\mathbf{r}). \tag{47}$$

Combining Eq. (46) and (47) gives

$$\mathbf{j}_e(\mathbf{r}) = e^2 \frac{\partial n}{\partial \mu} D(T) \mathbf{E}(\mathbf{r}).$$
(48)

Comparing this with Ohm's law in Eq. (45) we finally get

$$\sigma(T) = e^2 \frac{\partial n}{\partial \mu} D(T).$$
(49)

- ⁶ T. Faulkner, N. Iqbal, H. Liu, J. McGreevy, and D. Vegh, Science **329**, 1043 (2010).
- J. M. Maldacena, Adv. Theor. Math. Phys. 2, 231 (1998).
- ⁸ P. K. Kovtun, D. T. Son, and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005).
- ⁹ E. Shuryak, Prog. Part. Nucl. Phys. **53**, 273 (2004).
- ¹⁰ C. Cao *et al.*, Science **331**, 58 (2011).
- ¹¹ T. Schäfer, Ann. Rev. Nuc. Part. Sci. **64**, 125 (2014).
- ¹² S. A. Hartnoll, Nature Phys. **11**, 54 (2014).
- ¹³ P. W. Anderson, Phys. Today. **66**, 9 (2013).
- ¹⁴ A. Sommer, M. Ku, G. Raoti, and M. W. Zwierlein, Nature 472, 201 (2011).

- ¹⁵ A. B. Bardon *et al.*, Science **344**, 722 (2014).
- ¹⁶ T. Enss and R. Haussmann, Phys. Rev. Lett. **109**, 195303 (2012).
- ¹⁷ M. Koschorreck, D. Pertot, E. Vogt, and M. Köhl, Nature. Phys. 9, 405 (2013).
- ¹⁸ W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).
- ¹⁹ A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. 68, 13 (1996).
- ²⁰ P. Limelette *et al.*, Phys. Rev. Lett. **91**, 016401 (2003).
- ²¹ J. Merino, M. Dumm, N. Drichko, M. Dressel, and R. H. McKenzie, Phys. Rev. Lett. **100**, 086404 (2008).
- ²² B. J. Powell and R. H. McKenzie, Rep. Prog. Phys. 74, 056501 (2011).
- ²³ G. Kotliar *et al.*, Rev. Mod. Phys. **78**, 865 (2006).
- ²⁴ X. Y. Zhang, M. J. Rozenberg, and G. Kotliar, Phys. Rev. Lett. **70**, 1666 (1993).
- ²⁵ H. Kajueter and G. Kotliar, Phys. Rev. Lett. **77**, 131 (1996).
- ²⁶ R. Bulla, Phys. Rev. Lett. **83**, 136 (1999).
- ²⁷ H. Terletska, J. Vučičević, D. Tanasković, and V. Dobrosavljević, Phys. Rev. Lett. **107**, 026401 (2011).
- ²⁸ L.-F. Arsenault, P. Sémon, and A.-M. S. Tremblay, Phys. Rev. B 86, 085133 (2012).
- ²⁹ E. Economou, *Green's Functions in Quantum Physics* (Springer Berlin Heidelberg, 2010).
- ³⁰ A. Khurana, Phys. Rev. Lett. **64**, 1990 (1990).

- ³¹ T. Pruschke, D. L. Cox, and M. Jarrell, Phys. Rev. B 47, 3553 (1993).
- ³² W. Chung and J. K. Freericks, Phys. Rev. B 57, 11955 (1998).
- ³³ A. Chattopadhyay, A. J. Millis, and S. Das Sarma, Phys. Rev. B **61**, 10738 (2000).
- ³⁴ L.-F. Arsenault and A.-M. S. Tremblay, Phys. Rev. B 88, 205109 (2013).
- ³⁵ J. Kokalj and R. H. McKenzie, Phys. Rev. Lett. **110**, 206402 (2013).
- ³⁶ J. M. Luttinger, Phys. Rev. **119**, 1153 (1960).
- ³⁷ T. Hotta and S. Fujimoto, Phys. Rev. B **54**, 5381 (1996).
- ³⁸ N. Ashcroft and N. Mermin, *Solid State Physics* (Saunders College, 1976).
- ³⁹ Y. Kurosaki, Y. Shimizu, K. Miyagawa, K. Kanoda, and G. Saito, Phys. Rev. Lett. **95**, 177001 (2005).
- ⁴⁰ Hong Liu, Phys. Today **65**, 68 (2012).
- ⁴¹ T. Maier, M. Jarrell, T. Pruschke, and M. H. Hettler, Rev. Mod. Phys. **77**, 1027 (2005).
- ⁴² N. Lin, E. Gull, and A. J. Millis, Phys. Rev. B 82, 045104 (2010).
- ⁴³ N. Lin, E. Gull, and A. J. Millis, Phys. Rev. B 80, 161105 (2009).
- ⁴⁴ D. Bergeron, V. Hankevych, B. Kyung, and A.-M. S. Tremblay, Phys. Rev. B 84, 085128 (2011).