

MOTIVIC BIRATIONAL COVERS AND FINITE FILTRATIONS ON CHOW GROUPS

PABLO PELAEZ

ABSTRACT. We introduce a functorial tower of localizing subcategories in the Morel-Voevodsky motivic stable homotopy category. We study the tower for the spectrum HG representing motivic cohomology with coefficients in an abelian group G . When $G = \mathbb{Z}/\ell$ is a field, we show that the tower induces an interesting finite filtration on the Chow groups of smooth schemes over a perfect field. For $G = \mathbb{Z}$ or $G = \mathbb{Q}$, we show that the tower induces an interesting finite filtration on the motivic cohomology groups of smooth schemes over an algebraically closed field. With rational coefficients, this finite filtration satisfies several of the properties of the still conjectural Bloch-Beilinson-Murre filtration.

1. INTRODUCTION

The main goal of this paper is to construct an interesting finite filtration on the Chow groups (with coefficients in a finite field \mathbb{Z}/ℓ) of algebraic cycles modulo rational equivalence, and as well an interesting finite filtration on the motivic cohomology groups (with integral or rational coefficients). Considering rational coefficients, we obtain a finite filtration which satisfies several of the properties of the still conjectural Bloch-Beilinson-Murre filtration [1], [2], [14].

Our approach can be sketched quickly as follows. For a smooth scheme Y of finite type over a perfect field k , the Chow groups can be computed in Voevodsky's triangulated category of motives [31]. Thus, a standard adjointness argument implies that the Chow groups can be computed as well in the Morel-Voevodsky motivic stable homotopy category \mathcal{SH} :

$$CH^q(Y)_R \cong \mathrm{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), HR);$$

where HR represents in \mathcal{SH} motivic cohomology with R -coefficients. Since \mathcal{SH} is a triangulated category, it is possible to construct the filtration by considering a tower in \mathcal{SH} (see §5):

$$\cdots \rightarrow wb_{-3}^c(HR) \rightarrow wb_{-2}^c(HR) \rightarrow wb_{-1}^c(HR) \rightarrow HR$$

and defining the p -component of the filtration $F^p CH^q(Y)$ to be the image of the induced map:

$$\mathrm{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), wb_{-p}^c HR) \rightarrow \mathrm{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), HR) \cong CH^q(Y)_R$$

2010 *Mathematics Subject Classification.* Primary 14C25, 14C35, 14F42, 19E15; Secondary 18G55, 55P42.

Key words and phrases. Bloch-Beilinson-Murre filtration, Chow Groups, Filtration on the Chow Groups, Filtration on Motivic Cohomology, Localizing Subcategories, Motivic Cohomology, Motivic Homotopy Theory, Orthogonal Subcategories.

Research partially supported by DGAPA-UNAM grant IA100814.

With this definition, it is not at all obvious that the filtration $F^\bullet CH^q(Y)$ is finite. This is proved in 5.2.3. It is also interesting to observe that the filtration is defined (and is finite) for any coefficient ring R (not just the rationals) and for any smooth k -scheme of finite type (not necessarily projective).

Now, we describe the contents of the paper. In §1.1, the main results are stated and proved using the constructions of the remaining sections. In §1.2-1.3, we fix the notation and introduce the basic definitions that will be used in the rest of the paper. In §2, we prove some general results for localizing and orthogonal subcategories of a triangulated category which is the homotopy category of a Quillen model category, see 2.1.10, 2.2.11, 2.2.13, 2.2.16. In §3, we introduce the weakly birational covers and the weakly birational tower which will be used to construct the filtrations we are interested in, see 3.2.1, 3.2.4, 3.2.11, 3.2.15. In §4, we study the weakly birational tower with rational coefficients and show that in this case, all the components of the tower are motives and the corresponding maps in the tower are as well maps of motives, see 4.2.1. We also prove some auxiliary results of independent interest for the slice filtration with rational coefficients, see 4.1.13, 4.1.15, 4.1.16. Finally, in §5, we study the weakly birational tower for motivic cohomology and describe the main properties of the induced filtration, see 5.1.7, 5.1.8, 5.2.1, 5.2.3.

1.1. Main Results. We will consider a base scheme of the form $X = \text{Spec}(k)$, where k is a perfect field. We will write $H\mathbb{F}$ (resp. $H\mathbb{F}^{p,q}$) for the spectrum representing in \mathcal{SH} motivic cohomology with coefficients in a finite field $\mathbb{F} = \mathbb{Z}/\ell$, where $\ell \neq \text{char}(k)$; (resp. $\Sigma^{p-q}(\mathbb{G}_m^q \wedge H\mathbb{F})$). Let $\theta_{-n*}^{H\mathbb{F}} : wb_{-n}^c H\mathbb{F} \rightarrow H\mathbb{F}$ be the universal map defined in (5.1.3).

Theorem 1.1.1. *Let $Y \in \text{Sm}_X$, and $q \geq 0$. Then, there exists a non-trivial finite decreasing filtration F^\bullet on $CH^q(Y)_{\mathbb{F}}$ where the n -component is given by the image of $\theta_{-n*}^{H\mathbb{F}}$:*

$$\text{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), wb_{-n}^c H\mathbb{F}) \xrightarrow{\theta_{-n*}^{H\mathbb{F}}} \text{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), H\mathbb{F}) = CH^q(Y)_{\mathbb{F}}.$$

In addition, the filtration F^\bullet is functorial in Y with respect to morphisms in Sm_X and satisfies:

- (1) $F^0 CH^q(Y)_{\mathbb{F}} = CH^q(Y)_{\mathbb{F}}$,
- (2) $F^{q+1} CH^q(Y)_{\mathbb{F}} = 0$.

Proof. Since $CH^q(Y)_{\mathbb{F}} \cong \text{Hom}_{\mathcal{SH}}(\Sigma^{-q}(\mathbb{G}_m^{-q} \wedge Y_+), H\mathbb{F})$, we deduce all the properties (except for the non-triviality) by combining 5.2.1 and 5.2.3.

To show that the filtration is in general non-trivial, we will assume that $q = \ell m$ for some $m \geq 1$, and that there exists $a \in CH^m(Y)_{\mathbb{F}}$ such that $0 \neq a^\ell \in CH^q(Y)_{\mathbb{F}}$. Hence by [33, Lemma 9.8], we deduce that $0 \neq a^\ell = P^m(a) \in CH^q(Y)_{\mathbb{F}}$, where $P^m : H\mathbb{F} \rightarrow H\mathbb{F}^{2m(\ell-1), m(\ell-1)}$ is the motivic operation P^m constructed by Voevodsky in

[33, p. 33]. By adjointness, $P^m(a)$ is the following composition in \mathcal{SH} :

$$\begin{array}{c}
 \Sigma^{-m(\ell-1)}(\mathbb{G}_m^{-m(\ell-1)} \wedge \Sigma^{-m}(\mathbb{G}_m^{-m} \wedge Y_+)) \\
 \downarrow \Sigma^{-m(\ell-1)}(\mathbb{G}_m^{-m(\ell-1)} \wedge a) \\
 \Sigma^{-m(\ell-1)}(\mathbb{G}_m^{-m(\ell-1)} \wedge H\mathbb{F}) = H\mathbb{F}^{-2m(\ell-1), -m(\ell-1)} \\
 \downarrow \Sigma^{-m(\ell-1)}(\mathbb{G}_m^{-m(\ell-1)} \wedge P^m) \\
 H\mathbb{F}
 \end{array}$$

Finally, by 5.3.2 we deduce that $0 \neq P^m(a) \in F^{m(\ell-1)}CH^q(Y)_{\mathbb{F}}$. \square

Let $Y \in Sm_X$. We will write $H^{*,*}(Y, R)$ for $\oplus_{p,q \in \mathbb{Z}} H^{p,q}(Y, R)$, where:

$$H^{p,q}(Y, R) = \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), HR)$$

is the motivic cohomology of Y of degree p and weight q , with R -coefficients. In the next result we will consider motivic cohomology with integer and rational coefficients. We will write R for \mathbb{Z} or \mathbb{Q} ; and $\mathbf{1}$ for $\Sigma_T^\infty X_+$, the sphere spectrum in \mathcal{SH} .

Theorem 1.1.2. *Let the base scheme $X = \text{Spec}(k)$, with k an algebraically closed field; and $Y \in Sm_X$. Then, there exists a non-trivial decreasing filtration F^\bullet on $H^{*,*}(Y, R)$, where the n -component in degree p and weight q is given by the image of θ_{-n*}^{HR} :*

$$\begin{array}{ccc}
 \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), wb_{-n}^c HR) & \xrightarrow{\theta_{-n*}^{HR}} & \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), HR) \\
 & & \parallel \\
 & & H^{p,q}(Y, R).
 \end{array}$$

In addition, the filtration F^\bullet is functorial in Y with respect to morphisms in Sm_X and satisfies:

- (1) $F^0 H^{*,*}(Y, R) = H^{*,*}(Y, R)$,
- (2) $F^{q+1} H^{p,q}(Y, R) = 0$.

Proof. Combining 5.2.1 and 5.2.3, we deduce all the properties, except for the non-triviality of the filtration.

To show the non-triviality of the filtration, by 5.1.8 and 5.1.5 it suffices to construct a non-trivial map $HR \rightarrow \mathbb{G}_m^q \wedge HR$, for $q > 0$. To obtain such a map, it suffices to show the existence of a non-trivial map $a : \mathbf{1} \rightarrow \mathbb{G}_m^q \wedge HR$; since $a \wedge HR$ will produce the non-trivial map we were looking for.

By definition of the sphere spectrum, $\text{Hom}_{\mathcal{SH}}(\mathbf{1}, \mathbb{G}_m^q \wedge HR) \cong H^{q,q}(X, R)$, where $X = \text{Spec}(k)$. Now, a theorem of Nesterenko-Suslin and Suslin-Voevodsky [18], [27, Thm. 3.4] implies that $H^{q,q}(X, R)$ is isomorphic to the degree q Milnor K -theory of k with R coefficients, $K_q^M(k)_R$. However, it is well-known that $K_q^M(k)_{\mathbb{Z}}$ is uniquely divisible [36, Ch. III, §7, Ex. 7.3(a)]. Thus, $H^{q,q}(X, \mathbb{Z}) \neq 0$ and $H^{q,q}(X, \mathbb{Q}) \neq 0$, as we wanted. \square

With rational coefficients, the filtration in 1.1.2 is not only functorial with respect to maps in Sm_X , but in fact it is functorial with respect to Suslin-Voevodsky

transfers. Let $DM_{\mathbb{Q}}$ denote Voevodsky's big (admitting infinite direct sums) triangulated category of motives with rational coefficients.

Theorem 1.1.3. *Let $R = \mathbb{Q}$, and $X = \text{Spec}(k)$ with k a perfect field. Then the filtration F^{\bullet} in 1.1.2 satisfies:*

- (1) F^{\bullet} is functorial with respect to maps in $DM_{\mathbb{Q}}$.
- (2) F^{\bullet} is functorial with respect to Suslin-Voevodsky transfers.

Proof. It suffices to prove (1), since there is a functor $Cor(X) \rightarrow DM_{\mathbb{Q}}$ [30, Thm 3.2.6], where $Cor(X)$ is the Suslin-Voevodsky category of finite correspondences over X ; having same objects as Sm_X , morphisms $c(Y, Z)$ given by the group of finite relative cycles on $Y \times_X Z$ over Y [28], and composition as in [35, (2.1) p. 673].

Finally, (1) follows directly from 4.2.1. \square

When k is an algebraically closed field (e.g. $k = \mathbb{C}$), combining 1.1.2 and 1.1.3 we deduce that the filtration F^{\bullet} with rational coefficients satisfies several of the properties of the still conjectural Bloch-Beilinson-Murre filtration [1], [2], [14], [9, §2], which was originally defined for the Chow groups with rational coefficients, but admits a natural extension to motivic cohomology of arbitrary degree and weight. Namely, our filtration F^{\bullet} is non-trivial, finite and compatible with the action of correspondences (since it admits Suslin-Voevodsky transfers). Unfortunately, the methods of this paper are not sufficient to show that the filtration is non-trivial when restricted to the Chow groups with rational coefficients.

1.2. Definitions and Notation. In this paper X will denote a Noetherian separated base scheme of finite Krull dimension, Sch_X the category of schemes of finite type over X and Sm_X the full subcategory of Sch_X consisting of smooth schemes over X regarded as a site with the Nisnevich topology. All the maps between schemes will be considered over the base X . Given $Y \in Sch_X$, all the closed subsets Z of Y will be considered as closed subschemes with the reduced structure.

Let \mathcal{M} be the category of pointed simplicial presheaves on Sm_X equipped with the motivic Quillen model structure [25] constructed by Jardine [10, Thm. 1.1], which is Quillen equivalent to the one defined originally by Morel-Voevodsky [10, Thm. 1.2], taking the affine line \mathbb{A}_X^1 as interval [13, p. 86 Thm. 3.2]. Given a map $f : W \rightarrow Y$ in Sm_X , we will abuse notation and denote by f the induced map $f : W_+ \rightarrow Y_+$ in \mathcal{M} between the corresponding pointed simplicial presheaves represented by W and Y , respectively.

We define T in \mathcal{M} as the pointed simplicial presheaf represented by $S^1 \wedge \mathbb{G}_m$, where \mathbb{G}_m is the multiplicative group $\mathbb{A}_X^1 - \{0\}$ pointed by 1, and S^1 denotes the simplicial circle. Given an arbitrary integer $r \geq 1$, let S^r (resp. \mathbb{G}_m^r) denote the iterated smash product of S^1 (resp. \mathbb{G}_m) with r -factors: $S^1 \wedge \cdots \wedge S^1$ (resp. $\mathbb{G}_m \wedge \cdots \wedge \mathbb{G}_m$); $S^0 = \mathbb{G}_m^0$ will be by definition equal to the pointed simplicial presheaf represented by the base scheme X . Let $Spt(\mathcal{M})$ denote Jardine's category of symmetric T -spectra on \mathcal{M} equipped with the motivic model structure defined in [10, Thm. 4.15]. The smash product of symmetric T -spectra endows $Spt(\mathcal{M})$ with the structure of a symmetric monoidal model category [10, Prop. 4.14]. Hence, it is possible to consider (commutative) ring spectra with unit in $Spt(\mathcal{M})$ and modules in $Spt(\mathcal{M})$ over a given (commutative) ring spectrum.

We will use the language of triangulated categories. Our main reference will be [16]. Given a triangulated category, we will write Σ^1 (resp. Σ^{-1}) to denote its

suspension (resp. desuspension) functor; and for $n > 0$, Σ^n (resp. Σ^{-n}) will be the composition of Σ^1 (resp. Σ^{-1}) iterated n -times (resp. $-n$ -times). To simplify the notation, we will write Σ^0 for the identity functor.

We will use the following notation in all the categories under consideration: $*$ will denote the terminal object, and \cong will denote that a map or a functor is an isomorphism. If a category is simplicial, we will write $\text{Map}(-, -)$ for its simplicial set of morphisms.

1.3. Quillen model categories [24]. We will follow the conventions of [5], in particular all the Quillen model categories that we will consider will be closed under arbitrary limits and colimits. Given a Quillen model category \mathcal{A} , unless explicitly stated otherwise we will write $\mathbf{Ho}(\mathcal{A})$ for its homotopy category.

1.3.1. Bousfield localization. For details and definitions about Bousfield localization we refer the reader to Hirschhorn's book [5]. Let us just mention the following theorem of Hirschhorn, which guarantees the existence of left and right Bousfield localizations.

Theorem 1.3.2 (see [5, Thms. 4.1.1 and 5.1.1]). *Let \mathcal{A} be a Quillen model category which is cellular and proper. Let L be a set of maps in \mathcal{A} and let K be a set of objects in \mathcal{A} . Then:*

- (1) *The left Bousfield localization of \mathcal{A} with respect to L exists.*
- (2) *The right Bousfield localization of \mathcal{A} with respect to the class of K -colocal equivalences exists.*

The following result guarantees the existence of left and right Bousfield localizations for the motivic stable homotopy category $\text{Spt}(\mathcal{M})$.

Theorem 1.3.3. *The Quillen model category $\text{Spt}(\mathcal{M})$ is:*

- (1) *cellular (see [8], [6, Cor. 1.6] or [19, Thm. 2.7.4]).*
- (2) *proper (see [10, Thm. 4.15]).*

2. LOCALIZING AND ORTHOGONAL SUBCATEGORIES

Most of the results in this section could be stated in the language of triangulated categories [16, Ch. 8-9]. However, we will work within the framework of Quillen model categories which has the advantage that in the applications to motivic homotopy theory, the constructions admit a natural extension to the motivic unstable homotopy category [23].

In this section, \mathcal{A} will be a simplicial [25, II.1, II.2], stable [7, Ch. 7], proper and cellular Quillen model category. In particular, by 1.3.2 all the left and right Bousfield localizations that we will consider in this section exist. We will write R (resp. Q) for a fibrant (resp. cofibrant) replacement functor in \mathcal{A} .

Let \mathcal{T} denote the homotopy category of \mathcal{A} . We will consider \mathcal{T} as a triangulated category where the distinguished triangles are given by the cofibre sequences [25, I.3] in \mathcal{A} and the suspension functor is induced by smashing with respect to the simplicial circle S^1 [7, Def. 6.1.1].

2.1. Localizing subcategories. Let \mathcal{T}' be a subcategory of \mathcal{T} . Recall that \mathcal{T}' is a localizing subcategory of \mathcal{T} ; if \mathcal{T}' is a full triangulated subcategory of \mathcal{T} , and \mathcal{T}' is closed under arbitrary coproducts.

2.1.1. *Generators.* Let \mathcal{G} be a class of objects in $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$. We will follow the notation of Neeman [17] and write

$$(2.1.2) \quad \text{Loc}(\mathcal{G})$$

for the intersection of all the localizing subcategories of \mathcal{T} which contain \mathcal{G} . We will say that $\text{Loc}(\mathcal{G})$ is the localizing subcategory of \mathcal{T} generated by \mathcal{G} .

Lemma 2.1.3. *Let K be an object in $\text{Loc}(\mathcal{G})$ such that for every $G \in \mathcal{G}$ and every $n \in \mathbb{Z}$: $\text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, K) = 0$. Then $K = *$.*

Proof. Let ${}^\perp K$ denote the full subcategory of \mathcal{T} which consists of objects $L \in \mathcal{T}$ satisfying $\text{Hom}_{\mathcal{T}}(\Sigma^n L, K) = 0$, for every $n \in \mathbb{Z}$. We observe that ${}^\perp K$ is triangulated and closed under arbitrary coproducts. By hypothesis $\mathcal{G} \subseteq {}^\perp K$, hence $\text{Loc}(\mathcal{G}) \subseteq {}^\perp K$. Thus, we conclude that $K = *$. \square

Corollary 2.1.4. *Let $f : M \rightarrow N$ be a map in $\text{Loc}(\mathcal{G})$ such that for every $G \in \mathcal{G}$ and every $n \in \mathbb{Z}$, the induced map of abelian groups:*

$$f_* : \text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, M) \rightarrow \text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, N)$$

is an isomorphism. Then f is an isomorphism.

Proof. Complete f to a distinguished triangle in $\text{Loc}(\mathcal{G})$: $M \rightarrow N \rightarrow K$. Since the maps f_* above are all isomorphisms, we conclude that for every $G \in \mathcal{G}$ and every $n \in \mathbb{Z}$: $\text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, K) = 0$. By 2.1.3, we deduce that $K \cong *$ in $\text{Loc}(\mathcal{G})$. Thus $f : M \rightarrow N$ is an isomorphism. \square

2.1.5. *Presentation of localizing subcategories.* From now on we will assume that \mathcal{G} in (2.1.2) is a set. Let $\mathbf{R}_{\mathcal{G}}\mathcal{A}$ denote the right Bousfield localization of \mathcal{A} with respect to the set:

$$(2.1.6) \quad \Sigma^{\pm\infty}\mathcal{G} = \{\Sigma^n G : G \in \mathcal{G}; n \in \mathbb{Z}\}$$

Remark 2.1.7. By [5, 5.1.1.(1)(a) and 3.1.8] we can assume that the objects in $\Sigma^{\pm\infty}\mathcal{G}$ are cofibrant in \mathcal{A} .

Let C be a functorial cofibrant replacement in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$. Recall that \mathcal{A} is a simplicial Quillen model category with fibrant replacement functor R .

Proposition 2.1.8. *The homotopy category of $\mathbf{R}_{\mathcal{G}}\mathcal{A}$, $\mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$ is a triangulated category and the Quillen adjunction $(id, id, \varphi) : \mathbf{R}_{\mathcal{G}}\mathcal{A} \rightarrow \mathcal{A}$ induces an adjunction between triangulated functors:*

$$(C, R, \varphi) : \mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A}) \rightarrow \mathcal{T} = \mathbf{Ho}(\mathcal{A})$$

Proof. By construction, the suspension functor Σ^1 of $\mathbf{Ho}(\mathcal{A})$ is induced by the Quillen adjunction $(S^1, \Omega_{S^1}, \varphi) : \mathcal{A} \rightarrow \mathcal{A}$. Since $\Sigma^{\pm\infty}\mathcal{G} = \Sigma^1(\Sigma^{\pm\infty}\mathcal{G})$, it follows from [5, 3.3.20(2)(b)] that $(S^1, \Omega_{S^1}, \varphi) : \mathbf{R}_{\mathcal{G}}\mathcal{A} \rightarrow \mathbf{R}_{\mathcal{G}}\mathcal{A}$ is a Quillen equivalence. Thus, $\mathbf{R}_{\mathcal{G}}\mathcal{A}$ is a stable model category and its homotopy category $\mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$ is a triangulated category.

Since the identity functor $id : \mathcal{A} \rightarrow \mathbf{R}_{\mathcal{G}}\mathcal{A}$ is a right Quillen functor, we conclude that (C, R, φ) is an adjunction. By [7, Props. 6.4.1 and 7.1.12] we deduce that C and R are triangulated functors. \square

Proposition 2.1.9. *Let $f : M \rightarrow N$ be a map in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$. Then f is a weak equivalence in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$ if and only if for every $G \in \mathcal{G}$ and every $n \in \mathbb{Z}$, the induced map of abelian groups:*

$$f_* : \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, M) \rightarrow \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, N)$$

is an isomorphism.

Proof. (\Rightarrow): Assume that f is a weak equivalence in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$, equivalently a $\Sigma^{\pm\infty}\mathcal{G}$ -colocal equivalence in \mathcal{A} [5, 5.1.1(1)(a)]. By 2.1.7, we can assume that all the objects in $\Sigma^{\pm\infty}\mathcal{G}$ are cofibrant in \mathcal{A} . Hence, for every H in $\Sigma^{\pm\infty}\mathcal{G}$ the following maps are weak equivalences of simplicial sets [5, 3.1.8]:

$$\mathrm{Map}(H, RX) \xrightarrow{(Rf)_*} \mathrm{Map}(H, RY)$$

Since \mathcal{A} is a simplicial model category and H is cofibrant, we deduce that $\mathrm{Map}(H, RX)$ and $\mathrm{Map}(H, RY)$ are both Kan complexes. Thus, for every $r \geq 0$ and every $H \in \Sigma^{\pm\infty}\mathcal{G}$ the following diagram commutes, where the top row and the vertical maps are all isomorphisms of abelian groups:

$$\begin{array}{ccc} \pi_r \mathrm{Map}(H, RX) & \xrightarrow[\cong]{(Rf)_*} & \pi_r \mathrm{Map}(H, RY) \\ \cong \downarrow & & \downarrow \cong \\ \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^r H, X) & \xrightarrow{f_*} & \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^r H, Y) \end{array}$$

Therefore, for every $G \in \mathcal{G}$ (see 2.1.6) and every $n \in \mathbb{Z}$ the map:

$$\mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, X) \xrightarrow{f_*} \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, Y)$$

is an isomorphism of abelian groups.

(\Leftarrow): Fix $H \in \Sigma^{\pm\infty}\mathcal{G}$. Let ω_0, η_0 denote the base points of $\mathrm{Map}(H, R)$ and $\mathrm{Map}(H, RY)$ respectively. It suffices to show that the map:

$$\mathrm{Map}(H, RX) \xrightarrow{(Rf)_*} \mathrm{Map}(H, RY)$$

is a weak equivalence of simplicial sets [5, 5.1.1(1)(a) and 3.1.8].

Since \mathcal{A} is a pointed simplicial model category, lemma 6.1.2 in [7] implies that for $k \geq 0$ the following diagram commutes:

$$\begin{array}{ccc} \pi_{k, \omega_0} \mathrm{Map}(H, RX) & \xrightarrow{(Rf)_*} & \pi_{k, \eta_0} \mathrm{Map}(H, RY) \\ \cong \downarrow & & \downarrow \cong \\ \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^k H, X) & \xrightarrow{f_*} & \mathrm{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^k H, Y) \end{array}$$

By hypothesis the bottom row is an isomorphism of abelian groups, thus we deduce that all the maps in the top row are also isomorphisms. Therefore, for every $H \in \Sigma^{\pm\infty}\mathcal{G}$, the induced map of simplicial sets:

$$\mathrm{Map}(H, RX) \xrightarrow{(Rf)_*} \mathrm{Map}(H, RY)$$

is a weak equivalence when it is restricted to the path component of $\text{Map}(H, RX)$ containing ω_0 . We observe that $\Sigma^{-1}H$ is also in $\Sigma^{\pm\infty}\mathcal{G}$, hence the following map:

$$\text{Map}(S^1, \text{Map}(\Sigma^{-1}H, RX)) \longrightarrow \text{Map}(S^1, \text{Map}(\Sigma^{-1}H, RY))$$

is a weak equivalence of simplicial sets, since taking S^1 -loops kills the path components that do not contain the base point. Then, we observe that the rows in the following commutative diagram are isomorphisms, since \mathcal{A} is a simplicial stable model category:

$$\begin{array}{ccc} \text{Map}(S^1, \text{Map}(\Sigma^{-1}H, RX)) & \xrightarrow{\cong} & \text{Map}(H, RX) \\ \downarrow & & \downarrow (Rf)_* \\ \text{Map}(S^1, \text{Map}(\Sigma^{-1}H, RY)) & \xrightarrow[\cong]{} & \text{Map}(H, RY) \end{array}$$

Hence, the two out of three property for weak equivalences implies that the $(Rf)_*$ is a weak equivalence of simplicial sets. \square

Theorem 2.1.10. *Assume that all the objects in \mathcal{G} are compact in $\text{Loc}(\mathcal{G})$ in the sense of Neeman [15, Def. 1.6]. Then, $\text{Loc}(\mathcal{G})$ is naturally equivalent as a triangulated category to the homotopy category $\mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$.*

Proof. Combining 2.1.3 with the compactness of the objects in \mathcal{G} , we conclude that $\text{Loc}(\mathcal{G})$ is compactly generated in the sense of Neeman [15, Def. 1.7]. Thus, by Neeman's version of Brown representability [15, Thm. 4.1], we deduce that the inclusion $i : \text{Loc}(\mathcal{G}) \rightarrow \mathcal{T}$ admits a right adjoint r .

Let η denote the counit of the adjunction $(i, r, \varphi) : \text{Loc}(\mathcal{G}) \rightarrow \mathcal{T}$, and let $K \in \mathbf{R}_{\mathcal{G}}\mathcal{A}$ be an arbitrary cofibrant and fibrant object. We observe that for every $G \in \mathcal{G}$ and every $n \in \mathbb{Z}$, $\Sigma^n G$ is in $\text{Loc}(\mathcal{G})$. Since i is a full embedding, we deduce:

$$\text{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, i \circ r(K)) = \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(i(\Sigma^n G), i \circ r(K)) = \text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, r(K))$$

Thus, by adjointness:

$$\begin{aligned} \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, i \circ r(K)) &= \text{Hom}_{\text{Loc}(\mathcal{G})}(\Sigma^n G, r(K)) \\ &\cong \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(i(\Sigma^n G), K) = \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, K) \end{aligned}$$

Hence, we deduce that for every $G \in \mathcal{G}$, and every $n \in \mathbb{Z}$, the induced map of abelian groups:

$$\eta_*^K : \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, i \circ r(K)) \rightarrow \text{Hom}_{\mathbf{Ho}(\mathcal{A})}(\Sigma^n G, K)$$

is an isomorphism. Thus, by 2.1.9 we conclude that the map $R(\eta^K) : R(i \circ r(K)) \rightarrow K$ in $\mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$ is an isomorphism. This implies that the triangulated functor $R \circ i : \text{Loc}(\mathcal{G}) \rightarrow \mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$ is essentially surjective on objects. Thus, it only remains to check that $R \circ i$ is fully faithful.

By 2.1.7, we can assume that the objects in $\Sigma^{\pm\infty}\mathcal{G}$ are cofibrant in \mathcal{A} . Thus, it follows from [5, 5.1.6] that the objects in $\text{Loc}(\mathcal{G})$ are all cofibrant in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$. Then, by [5, 3.5.2(2)] we deduce that $R \circ i$ is fully faithful, since all the objects in $\text{Loc}(\mathcal{G})$ are cofibrant in $\mathbf{R}_{\mathcal{G}}\mathcal{A}$ and hence colocal in \mathcal{A} [5, 5.1.5 and 5.1.6]. \square

Corollary 2.1.11. *Assume that all the objects in \mathcal{G} are compact in $\text{Loc}(\mathcal{G})$ in the sense of Neeman [15, Def. 1.6]. Then, $\mathbf{Ho}(\mathbf{R}_{\mathcal{G}}\mathcal{A})$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6], with $\Sigma^{\pm\infty}\mathcal{G}$ (see 2.1.6) as set of compact generators [15, Def. 1.8].*

Proof. By 2.1.10 it suffices to show that $\text{Loc}(\mathcal{G})$ is compactly generated with set of generators given by $\Sigma^{\pm\infty}\mathcal{G}$. But this follows directly from 2.1.3 since we are assuming that \mathcal{G} is a set, and that all its objects are compact. \square

2.2. Orthogonal subcategories. Given a triangulated subcategory \mathcal{T}' of \mathcal{T} , let \mathcal{T}'^\perp (resp. ${}^\perp\mathcal{T}'$) denote the full subcategory of \mathcal{T} with objects K such that for every $A \in \mathcal{T}'$, $\text{Hom}_{\mathcal{T}}(A, K) = 0$ (resp. $\text{Hom}_{\mathcal{T}}(K, A) = 0$).

Lemma 2.2.1. *Let \mathcal{T}' be triangulated subcategory of \mathcal{T} , then \mathcal{T}'^\perp and ${}^\perp\mathcal{T}'$ are full triangulated subcategories of \mathcal{T} .*

Proof. By definition \mathcal{T}'^\perp and ${}^\perp\mathcal{T}'$ are full subcategories of \mathcal{T} . Thus, it suffices to check that they are triangulated. But this follows immediately from the fact that the functor $\text{Hom}_{\mathcal{T}}(A, -)$ (resp. $\text{Hom}_{\mathcal{T}}(-, A)$) is homological (resp. cohomological) for every $A \in \mathcal{T}$ (see [16, Def. 1.1.7]). \square

2.2.2. Presentation of orthogonal subcategories. From now on we will assume that \mathcal{G} in (2.1.2) is a set. Our goal is to find a presentation for $\text{Loc}(\mathcal{G})^\perp$. Let $\mathbf{L}_{\mathcal{G}}\mathcal{A}$ denote the left Bousfield localization of \mathcal{A} with respect to the set of maps:

$$(2.2.3) \quad M_{\mathcal{G}} = \{\Sigma^n G \rightarrow * : G \in \mathcal{G}, n \in \mathbb{Z}\}$$

Remark 2.2.4. Taking $F = U = \text{id}$ in [5, 3.3.20.(1)(b)] we can assume that the objects $\Sigma^n G$ appearing as domains in $M_{\mathcal{G}}$ are cofibrant in \mathcal{A} .

Let L be a functorial fibrant replacement in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$. Recall that \mathcal{A} is a simplicial Quillen model category with cofibrant replacement functor Q .

Proposition 2.2.5. *The homotopy category of $\mathbf{L}_{\mathcal{G}}\mathcal{A}$, $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$ is a triangulated category and the Quillen adjunction $(\text{id}, \text{id}, \varphi) : \mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}}\mathcal{A}$ induces an adjunction between triangulated functors:*

$$(Q, L, \varphi) : \mathcal{T} = \mathbf{Ho}(\mathcal{A}) \rightarrow \mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$$

Proof. By construction, the suspension functor Σ^1 of $\mathbf{Ho}(\mathcal{A})$ is induced by the Quillen adjunction $(S^1, \Omega_{S^1}, \varphi) : \mathcal{A} \rightarrow \mathcal{A}$. Since $M_{\mathcal{G}} = \Sigma^1(M_{\mathcal{G}})$, it follows from [5, 3.3.20(1)(b)] that $(S^1, \Omega_{S^1}, \varphi) : \mathbf{L}_{\mathcal{G}}\mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}}\mathcal{A}$ is a Quillen equivalence. Thus, $\mathbf{L}_{\mathcal{G}}\mathcal{A}$ is a stable model category and its homotopy category $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$ is a triangulated category.

Since the identity functor $\text{id} : \mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}}\mathcal{A}$ is a left Quillen functor, we conclude that (Q, L, φ) is an adjunction. By [7, Props. 6.4.1 and 7.1.12] we deduce that Q and L are triangulated functors. \square

Example 2.2.6. Let W be a set consisting of maps in \mathcal{A} . Let $\mathbf{L}_W\mathcal{A}$ denote the left Bousfield localization of \mathcal{A} with respect to the set of maps:

$$sW = \{\Sigma^n f : f \in W, n \in \mathbb{Z}\}$$

$\mathbf{L}_W\mathcal{A}$ admits a description as in 2.2.2, 2.2.3. For this, consider the set of objects \mathcal{G}_W in \mathcal{A} which complete the maps in W to cofibre sequences in \mathcal{A} (distinguished triangles in $\mathbf{Ho}(\mathcal{A})$). Namely, there is a bijection of sets $W \rightarrow \mathcal{G}_W$, $f \mapsto G_f$ such that for every $f : E \rightarrow F$ in W , G_f fits in a cofibre sequence in \mathcal{A} : $E \rightarrow F \rightarrow G_f$. Let $\mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ be the left Bousfield localization of \mathcal{A} with respect to the set of maps $M_{\mathcal{G}_W}$ (see 2.2.3).

Let L be a functorial fibrant replacement in $\mathbf{L}_W\mathcal{A}$. Recall that \mathcal{A} is a simplicial Quillen model category with cofibrant replacement functor Q .

Lemma 2.2.7. (1) *The homotopy category of $\mathbf{L}_W\mathcal{A}$, $\mathbf{Ho}(\mathbf{L}_W\mathcal{A})$ is a triangulated category and the Quillen adjunction $(id, id, \varphi) : \mathcal{A} \rightarrow \mathbf{L}_W\mathcal{A}$ induces an adjunction between triangulated functors:*

$$(Q, L, \varphi) : \mathcal{T} = \mathbf{Ho}(\mathcal{A}) \rightarrow \mathbf{Ho}(\mathbf{L}_W\mathcal{A})$$

- (2) *Let $G \in \mathcal{G}_W$ and $\Sigma^n G \rightarrow *$ a map in $M_{\mathcal{G}_W}$ (see 2.2.3). Then $\Sigma^n G \rightarrow *$ becomes an isomorphism in $\mathbf{Ho}(\mathbf{L}_W\mathcal{A})$.*
- (3) *Let $G \in \mathcal{G}_W$. Then the map $G \rightarrow *$ is an isomorphism in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}_W}\mathcal{A})$.*
- (4) *Let f be a map in sW (see 2.2.6). Then f becomes an isomorphism in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}_W}\mathcal{A})$.*

Proof. (1): It follows using the same argument as in 2.2.5.

(2) By 2.2.7(1) it suffices to show that $G \rightarrow *$ becomes an isomorphism in $\mathbf{Ho}(\mathbf{L}_W\mathcal{A})$. By construction of \mathcal{G}_W , there exists a map $f : E \rightarrow F$ in W , such that $E \xrightarrow{f} F \longrightarrow G$ is a distinguished triangle in $\mathbf{Ho}(\mathcal{A})$. Thus, by 2.2.7(1) it suffices to check that f becomes an isomorphism in $\mathbf{Ho}(\mathbf{L}_W\mathcal{A})$. This follows directly from the definition of $\mathbf{L}_W\mathcal{A}$ and the universal property of left Bousfield localizations [5, 3.1.1.(1)(a) and 3.3.19.(1)].

(3): This follows directly from the construction of $\mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ and [5, 3.1.1.(1)(a) and 3.3.19.(1)].

(4): By 2.2.5 we can assume that f is in W (see 2.2.6). By construction there is a distinguished triangle in $\mathbf{Ho}(\mathcal{A})$ with $G \in \mathcal{G}_W$: $E \xrightarrow{f} F \longrightarrow G$. Combining 2.2.7(3) and 2.2.5 we deduce that f becomes an isomorphism in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}_W}\mathcal{A})$. \square

Proposition 2.2.8. *The identity functor:*

$$(id, id, \varphi) : \mathbf{L}_W\mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}_W}\mathcal{A}$$

is a Quillen equivalence.

Proof. Since $\mathbf{L}_W\mathcal{A}$, $\mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ are left Bousfield localizations of \mathcal{A} , we deduce that they are simplicial model categories with the same cofibrant replacement functor Q . Thus, it suffices to show that they have the same class of weak equivalences. By [5, 9.7.4], it is enough to check that the fibrant objects in $\mathbf{L}_W\mathcal{A}$ and $\mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ are the same.

For this, it suffices to show [5, 3.1.6.(c) and 4.1.1.(2)] that the identity functor $id : \mathbf{L}_W\mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ (resp. $id : \mathbf{L}_{\mathcal{G}_W}\mathcal{A} \rightarrow \mathbf{L}_W\mathcal{A}$) is a left Quillen functor. We consider the following diagram, where the arrows are left Quillen functors:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{id} & \mathbf{L}_{\mathcal{G}_W}\mathcal{A} \\ id \downarrow & & \\ \mathbf{L}_W\mathcal{A} & & \end{array}$$

By 2.2.7(4) (resp. 2.2.7(2)), the maps in sW (resp. $M_{\mathcal{G}_W}$) become isomorphisms in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}_W}\mathcal{A})$ (resp. $\mathbf{Ho}(\mathbf{L}_W\mathcal{A})$). Thus, the universal property of left Bousfield localizations [5, 3.3.19.(1) and 3.1.1.(1)] implies that $id : \mathbf{L}_W\mathcal{A} \rightarrow \mathbf{L}_{\mathcal{G}_W}\mathcal{A}$ (resp. $id : \mathbf{L}_{\mathcal{G}_W}\mathcal{A} \rightarrow \mathbf{L}_W\mathcal{A}$) is a left Quillen functor. \square

The next two lemmas characterize the fibrant objects in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$. Recall that L is a fibrant replacement functor in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$.

Lemma 2.2.9. *Let K be an arbitrary object in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$. Then LK is in $\text{Loc}(\mathcal{G})^{\perp}$.*

Proof. We will assume that the objects $\Sigma^n G$ appearing as domains in $M_{\mathcal{G}}$ (see (2.2.3)) are cofibrant in \mathcal{A} (see 2.2.4). The universal property of left Bousfield localizations [5, 3.3.19.(1) and 3.1.1.(1)] implies that for every $n \in \mathbb{Z}$ and every $G \in \mathcal{G}$, the map $\Sigma^n G \rightarrow *$ is an isomorphism in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$. By adjointness:

$$\text{Hom}_{\mathcal{T}}(\Sigma^n G, LK) \cong \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(\Sigma^n G, K) = 0$$

for every $n \in \mathbb{Z}$ and every $G \in \mathcal{G}$. Thus, we conclude that \mathcal{G} is contained in ${}^{\perp}LK$, where ${}^{\perp}LK$ is the full subcategory of $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$ with objects A such that for every $n \in \mathbb{Z}$, $\text{Hom}_{\mathcal{T}}(\Sigma^n A, LK) = 0$.

We observe that ${}^{\perp}LK$ is a full triangulated subcategory of \mathcal{T} (see 2.2.1) and is closed under arbitrary coproducts. Thus, $\text{Loc}(\mathcal{G}) \subseteq {}^{\perp}LK$, i.e. $LK \in \text{Loc}(\mathcal{G})^{\perp}$. \square

Lemma 2.2.10. *Let K be a fibrant object in \mathcal{A} . Assume that K is in $\text{Loc}(\mathcal{G})^{\perp}$. Then K is also fibrant in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$.*

Proof. We will assume that the objects $\Sigma^n G$ appearing as domains in $M_{\mathcal{G}}$ (see (2.2.3)) are cofibrant in \mathcal{A} (see 2.2.4). By [5, 3.1.4.(1)(a) and 4.1.1] it suffices to show that the map: $*$ $= \text{Map}(*, K) \rightarrow \text{Map}(\Sigma^n G, K)$, is a weak equivalence of simplicial sets for every $n \in \mathbb{Z}$ and every $G \in \mathcal{G}$.

Let ω be a base point for $\text{Map}(\Sigma^n G, K)$. Since \mathcal{A} is a simplicial model category, we deduce $\pi_0 \text{Map}(\Sigma^n G, K) \cong \text{Hom}_{\mathcal{T}}(\Sigma^n G, K)$ which is zero by hypothesis when $G \in \mathcal{G}$. Thus, we conclude that $\text{Map}(\Sigma^n G, K)$ has only one connected component, and as a consequence it suffices to show that for $r \geq 1$ the homotopy groups $\pi_{r,\omega} \text{Map}(\Sigma^n G, K) = *$.

By [8, Lem. 6.1.2], we conclude $\pi_{r,\omega} \text{Map}(\Sigma^n G, K) \cong \text{Hom}_{\mathcal{T}}(\Sigma^{n+r} G, K)$ which vanishes by hypothesis. Hence the result follows. \square

The following theorem is the main result of this section.

Theorem 2.2.11. *Assume that \mathcal{G} is a set. Then, $\text{Loc}(\mathcal{G})^{\perp}$ is naturally equivalent as a triangulated category to the homotopy category $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$.*

Proof. Combining 2.2.5 and 2.2.9, we deduce that the triangulated functor $L : \mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A}) \rightarrow \mathcal{T} = \mathbf{Ho}(\mathcal{A})$, factors through the inclusion $j : \text{Loc}(\mathcal{G})^{\perp} \rightarrow \mathcal{T}$, which is also a triangulated functor.

We will abuse notation and write $L : \mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A}) \rightarrow \text{Loc}(\mathcal{G})^{\perp}$ for the corresponding factorization. By [5, 3.5.2(1)], we conclude that L is fully faithful. Finally, lemma 2.2.10 implies that L is essentially surjective on objects, and so an equivalence of triangulated categories. \square

Corollary 2.2.12. *Assume that \mathcal{G} is a set. Then, $\text{Loc}(\mathcal{G})^{\perp}$ is a localizing subcategory of $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$.*

Proof. Recall (see 1.3) that \mathcal{A} is in particular closed under arbitrary coproducts. Thus, by 2.2.11 we deduce that $\text{Loc}(\mathcal{G})^{\perp}$ is closed under arbitrary coproducts. Therefore, the result follows from 2.2.1. \square

We are particularly interested in conditions that guarantee that $\text{Loc}(\mathcal{G})^\perp$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6].

Recall that L is a functorial fibrant replacement in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$.

Theorem 2.2.13. *Assume that \mathcal{G} is a set and that $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6], with set of compact generators \mathcal{K} [15, Def. 1.8]. Then, $\text{Loc}(\mathcal{G})^\perp$ is a compactly generated triangulated category in the sense of Neeman, with set of compact generators:*

$$LK = \{LK : K \in \mathcal{K}\}.$$

Proof. By 2.2.11, it suffices to show that $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$ is compactly generated with generators $\{LK : K \in \mathcal{K}\}$.

Let $B \in \mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$ such that for every $K \in \mathcal{K}$: $\text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, B) = 0$. Then $B \cong *$ in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$. In effect, we may assume that B is fibrant in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$, then by adjointness:

$$\text{Hom}_{\mathcal{T}}(K, B) \cong \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(K, B) = \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, B) = 0$$

Since \mathcal{K} is a set of compact generators for \mathcal{T} , we deduce that $B \cong *$ in \mathcal{T} . By [5, 3.3.3.(1)(a)] we conclude that $B \cong *$ in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$.

It only remains to show that the objects in $LK = \{LK : K \in \mathcal{K}\}$ are compact in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$. Let B_λ be a family of objects in $\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})$ indexed by a set Λ . We need to show:

$$\text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, \oplus_{\lambda \in \Lambda} B_\lambda) \cong \oplus_{\lambda \in \Lambda} \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, B_\lambda)$$

We may assume that $\oplus_{\lambda \in \Lambda} B_\lambda$ is fibrant in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$. By 2.2.10, we deduce that for every $\lambda \in \Lambda$, B_λ is also fibrant in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$. By adjointness and the compactness of K in \mathcal{T} :

$$\begin{aligned} \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, \oplus_{\lambda \in \Lambda} B_\lambda) &= \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(K, \oplus_{\lambda \in \Lambda} B_\lambda) \cong \text{Hom}_{\mathcal{T}}(K, \oplus_{\lambda \in \Lambda} B_\lambda) \\ &\cong \oplus_{\lambda \in \Lambda} \text{Hom}_{\mathcal{T}}(K, B_\lambda) \cong \oplus_{\lambda \in \Lambda} \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(K, B_\lambda) \\ &= \oplus_{\lambda \in \Lambda} \text{Hom}_{\mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A})}(LK, B_\lambda) \end{aligned}$$

Hence the result follows. \square

We obtain the following interesting corollaries, where L is a functorial fibrant replacement in $\mathbf{L}_{\mathcal{G}}\mathcal{A}$.

Corollary 2.2.14. *Assume that \mathcal{G} is a set and that $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6], with set of compact generators \mathcal{K} [15, Def. 1.8]. Then, $\text{Loc}(\mathcal{G})^\perp = \text{Loc}(LK)$ (see 2.1.2).*

Proof. It follows from 2.2.13 and [15, Thm. 2.1(2.1.2)]. \square

With the notation of 2.1.5, 2.1.6 let $\mathbf{R}_{L\mathcal{K}}\mathcal{A}$ be the right Bousfield localization of \mathcal{A} with respect to the set: $\{\Sigma^n LK : K \in \mathcal{K}, n \in \mathbb{Z}\}$.

Corollary 2.2.15. *Assume that \mathcal{G} is a set and that $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6], with set of compact generators \mathcal{K} [15, Def. 1.8]. Then, $\text{Loc}(\mathcal{G})^\perp$ is naturally equivalent as a triangulated category to the homotopy category $\mathbf{Ho}(\mathbf{R}_{L\mathcal{K}}\mathcal{A})$.*

Proof. By 2.2.14, $\text{Loc}(\mathcal{G})^\perp = \text{Loc}(LK)$. Hence, the result follows from 2.2.13 and 2.1.10. \square

Combining 2.2.11 and 2.2.15 we obtain two different presentations for $Loc(\mathcal{G})^\perp$, one as a left Bousfield localization and the other as a right Bousfield localization:

Corollary 2.2.16. *Assume that \mathcal{G} is a set and that $\mathcal{T} = \mathbf{Ho}(\mathcal{A})$ is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.6], with set of compact generators \mathcal{K} [15, Def. 1.8]. Then,*

$$Loc(\mathcal{G})^\perp \cong \mathbf{Ho}(\mathbf{L}_{\mathcal{G}}\mathcal{A}) \cong \mathbf{Ho}(\mathbf{R}_{L\mathcal{K}}\mathcal{A})$$

are naturally equivalent as triangulated categories.

3. WEAKLY BIRATIONAL COVERINGS AND THE WEAKLY BIRATIONAL TOWER

In [21] we introduced the birational (resp. weakly birational) motivic stable homotopy categories. In this section, after recalling their definition we show that we can apply to them the formalism of §2 and then use this formalism to construct the (weakly) birational coverings and the birational tower in the Morel-Voevodsky motivic stable homotopy category.

Recall that $Spt(\mathcal{M})$ is Jardine's category of symmetric T -spectra on \mathcal{M} equipped with the motivic model structure [10, theorem 4.15]. We will write \mathcal{SH} for its homotopy category, which is triangulated.

3.1. Birational motivic stable homotopy categories. We will use Jardine's notation [10, p. 506-507]. Namely, let F_n denote the left adjoint to the n -evaluation functor:

$$\begin{array}{ccc} Spt(\mathcal{M}) & \xrightarrow{ev_n} & \mathcal{M} \\ (E^m)_{m \geq 0} & \longmapsto & E^n \end{array}$$

Notice that $F_0(A)$ is just the usual infinite suspension spectrum $\Sigma_T^\infty A$. The sphere spectrum $\mathbf{1}$ is $F_0(S^0)$.

3.1.1. Generators for \mathcal{SH} . Recall that \mathcal{SH} is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.7] with set of compact generators [19, Prop. 3.1.5]:

$$(3.1.2) \quad \mathcal{G}_{\mathcal{SH}} = \{F_n(S^p \wedge \mathbb{G}_m^q \wedge U_+) : U \in Sm_X; n, p, q \geq 0\}.$$

where U_+ denotes the simplicial presheaf represented by U with a disjoint base point.

Remark 3.1.3. The desuspension functor $\Sigma^{-1} : \mathcal{SH} \rightarrow \mathcal{SH}$ can be written in this notation as $E \mapsto F_1(\mathbb{G}_m \wedge E)$. If $q > 0$, we will write $\mathbb{G}_m^{-q} \wedge E$ for $F_q(S^q \wedge E)$.

3.1.4. Codimension. Let $Y \in Sch_X$, and Z a closed subscheme of Y . The *codimension* of Z in Y (see [24, section 7.5]), $codim_Y Z$ is the infimum (over the generic points z_i of Z) of the dimensions of the local rings \mathcal{O}_{Y, z_i} .

Since X is Noetherian of finite Krull dimension and Y is of finite type over X , $codim_Y Z$ is always finite.

3.1.5. Open immersions. Let $n \geq 0$ be an integer, and consider the following set of open immersions which have a closed complement of codimension at least $n + 1$

$$\begin{aligned} B_n = \{ & \iota_{U,Y} : U \rightarrow Y \text{ open immersion} \mid \\ & Y \in Sm_X; Y \text{ irreducible}; (codim_Y Y \setminus U) \geq n + 1 \} \end{aligned}$$

3.1.6. *Open immersions with smooth complement.* Let $n \geq 0$, and consider the following set of open immersions with smooth closed complement of codimension at least $n + 1$

$$WB_n = \{\iota_{U,Y} : U \rightarrow Y \text{ open immersion} \mid \\ Y, Z = Y \setminus U \in Sm_X; Y \text{ irreducible}; (\text{codim}_Y Z) \geq n + 1\}$$

Now we will study the left Bousfield localizations of $Spt(\mathcal{M})$ with respect to a suitable set of maps induced by the families of open immersions B_n, WB_n described above.

3.1.7. *First presentation.* Let $n \in \mathbb{Z}$ be an arbitrary integer.

- (1) We will write $B_n Spt(\mathcal{M})$ (resp. $WB_n Spt(\mathcal{M})$) for the left Bousfield localization of $Spt(\mathcal{M})$ with respect to the set of maps:

$$sB_n = \{F_p(\mathbb{G}_m^b \wedge \iota_{U,Y}) : b, p, r \geq 0, b - p \geq n - r; \iota_{U,Y} \in B_r\} \\ (\text{resp. } sWB_n = \{F_p(\mathbb{G}_m^b \wedge \iota_{U,Y}) : b, p, r \geq 0, b - p \geq n - r; \iota_{U,Y} \in WB_r\}).$$

- (2) We will write $b^{(n)}$ (resp. $wb^{(n)}$) for a fibrant replacement functor in $B_n Spt(\mathcal{M})$ (resp. $WB_n Spt(\mathcal{M})$) and $\mathcal{SH}(B_n)$ (resp. $\mathcal{SH}(WB_n)$) for its associated homotopy category.

For $n \neq 0$ we will call $\mathcal{SH}(B_n)$ (resp. $\mathcal{SH}(WB_n)$) the codimension $(n+1)$ -birational motivic stable homotopy category (resp. codimension $(n+1)$ -weakly birational motivic stable homotopy category), and for $n = 0$ we will call it the birational motivic stable homotopy category (resp. weakly birational motivic stable homotopy category).

3.1.8. *Second presentation.*

- (1) Let \mathcal{G}_{B_n} (resp. \mathcal{G}_{WB_n}) be the set of objects in $Spt(\mathcal{M})$ which complete the maps in sB_n (resp. sWB_n) to cofibre sequences in $Spt(\mathcal{M})$, i.e. distinguished triangles in \mathcal{SH} . Namely, there is a bijection of sets $sB_n \rightarrow \mathcal{G}_{B_n}$ (resp. $sWB_n \rightarrow \mathcal{G}_{WB_n}$), $f \mapsto G_f$ such that for every $f : E \rightarrow F$ in sB_n (resp. sWB_n), G_f fits in a cofibre sequence in $Spt(\mathcal{M})$: $E \rightarrow F \rightarrow G_f$.
- (2) Let $\mathbf{L}_{\mathcal{G}_{B_n}} Spt(\mathcal{M})$ (resp. $\mathbf{L}_{\mathcal{G}_{WB_n}} Spt(\mathcal{M})$) be the left Bousfield localization of $Spt(\mathcal{M})$ with respect to the set of maps $M_{\mathcal{G}_{B_n}}$ (resp. $M_{\mathcal{G}_{WB_n}}$), see 2.2.3.
- (3) We will write $Loc(\mathcal{G}_{B_n})$ (resp. $Loc(\mathcal{G}_{WB_n})$) for the localizing subcategory of \mathcal{SH} generated by \mathcal{G}_{B_n} (resp. \mathcal{G}_{WB_n}).
- (4) We will write B_n^\perp (resp. WB_n^\perp) for $Loc(\mathcal{G}_{B_n})^\perp$ (resp. $Loc(\mathcal{G}_{WB_n})^\perp$) in \mathcal{SH} .

Theorem 3.1.9. *The identity functor:*

$$id : B_n Spt(\mathcal{M}) \rightarrow \mathbf{L}_{\mathcal{G}_{B_n}} Spt(\mathcal{M}) \\ (\text{resp. } id : WB_n Spt(\mathcal{M}) \rightarrow \mathbf{L}_{\mathcal{G}_{WB_n}} Spt(\mathcal{M})).$$

is a Quillen equivalence.

Proof. We just need to consider the first claim, the proof being exactly the same in the second case. By 2.2.6 and 2.2.8, it suffices to check that $B_n Spt(\mathcal{M})$ (see 3.1.7) is also the left Bousfield localization of $Spt(\mathcal{M})$ with respect to the set of maps $\{\Sigma^r f : f \in sB_n; r \geq 0\}$. This follows directly from Lemma 2.5 in [21] and the universal property of left Bousfield localizations [5, 3.1.1.(1)(a) and 3.3.19.(1)]. \square

3.1.10. *Third presentation.* Recall that $b^{(n)}$ (resp. $wb^{(n)}$) is a fibrant replacement functor in $B_n Spt(\mathcal{M})$ (resp. $WB_n Spt(\mathcal{M})$). By 3.1.9, $b^{(n)}$ (resp. $wb^{(n)}$) is also a fibrant replacement functor in $\mathbf{L}_{\mathcal{G}_{B_n}} Spt(\mathcal{M})$ (resp. $\mathbf{L}_{\mathcal{G}_{WB_n}} Spt(\mathcal{M})$). We will write $\mathbf{R}_{b^{(n)}\mathcal{G}_{\mathcal{SH}}} Spt(\mathcal{M})$ (resp. $\mathbf{R}_{wb^{(n)}\mathcal{G}_{\mathcal{SH}}} Spt(\mathcal{M})$) for the right Bousfield localization of $Spt(\mathcal{M})$ with respect to the set (see 3.1.2 and 2.1.5, 2.1.6):

$$\begin{aligned} & \{\Sigma^m(b^{(n)}G) : G \in \mathcal{G}_{\mathcal{SH}}, m \in \mathbb{Z}\} \\ & (\text{resp. } \{\Sigma^m(wb^{(n)}G) : G \in \mathcal{G}_{\mathcal{SH}}, m \in \mathbb{Z}\}). \end{aligned}$$

Recall that B_n^\perp, WB_n^\perp is respectively $Loc(\mathcal{G}_{B_n})^\perp$ and $Loc(\mathcal{G}_{WB_n})^\perp$.

Theorem 3.1.11. *The following:*

$$\begin{aligned} \mathcal{SH}(B_n) &\cong B_n^\perp \cong \mathbf{Ho}(\mathbf{R}_{b^{(n)}\mathcal{G}_{\mathcal{SH}}} Spt(\mathcal{M})) \\ (\text{resp. } \mathcal{SH}(WB_n) &\cong WB_n^\perp \cong \mathbf{Ho}(\mathbf{R}_{wb^{(n)}\mathcal{G}_{\mathcal{SH}}} Spt(\mathcal{M}))). \end{aligned}$$

are naturally equivalent as triangulated categories.

Proof. We just need to consider the first claim, the proof being exactly the same in the second case. By 3.1.1, \mathcal{SH} is compactly generated with set of compact generators $\mathcal{G}_{\mathcal{SH}}$. Hence the result follows by combining 3.1.9 and 2.2.16. \square

Recall that $\mathcal{G}_{\mathcal{SH}}$ is a set of compact generators for \mathcal{SH} (see 3.1.2).

Corollary 3.1.12. *The inclusion (see 3.1.8(4)):*

$$\begin{aligned} j_n : B_n^\perp &\rightarrow \mathcal{SH} \\ (\text{resp. } j_n : WB_n^\perp &\rightarrow \mathcal{SH}) \end{aligned}$$

is a triangulated functor which commutes with arbitrary coproducts. In addition, B_n^\perp (resp. WB_n^\perp) is a compactly generated triangulated category in the sense of Neeman [15, Def. 1.7] with set of compact generators:

$$\begin{aligned} b^{(n)}\mathcal{G}_{\mathcal{SH}} &= \{\Sigma^m(b^{(n)}G) : G \in \mathcal{G}_{\mathcal{SH}}, m \in \mathbb{Z}\} \\ (\text{resp. } wb^{(n)}\mathcal{G}_{\mathcal{SH}} &= \{\Sigma^m(wb^{(n)}G) : G \in \mathcal{G}_{\mathcal{SH}}, m \in \mathbb{Z}\}). \end{aligned}$$

Proof. We just need to consider the first claim, the proof being exactly the same in the second case. It is clear that the inclusion $B_n^\perp \rightarrow \mathcal{SH}$ is a triangulated functor and that it respects arbitrary coproducts. By 3.1.1, \mathcal{SH} is compactly generated with set of compact generators $\mathcal{G}_{\mathcal{SH}}$. The compactness of B_n^\perp follows from 3.1.11 and 2.2.13. \square

3.1.13. *Voevodsky's effective categories.* For every integer $q \in \mathbb{Z}$, consider the following family of symmetric T -spectra:

$$(3.1.14) \quad C_{\text{eff}}^q = \{F_n(S^r \wedge \mathbb{G}_m^s \wedge U_+) \mid n, r, s \geq 0; s - n \geq q; U \in Sm_X\}$$

Voevodsky [32] defines the q -effective motivic stable homotopy category $\Sigma_T^q \mathcal{SH}^{\text{eff}}$ to be $Loc(C_{\text{eff}}^q)$ (see 2.1.2), i.e. $\Sigma_T^q \mathcal{SH}^{\text{eff}}$ is the smallest full triangulated subcategory of \mathcal{SH} which contains C_{eff}^q and is closed under arbitrary coproducts. When $q = 0$ we will simply write $\mathcal{SH}^{\text{eff}}$ for $\Sigma_T^0 \mathcal{SH}^{\text{eff}}$.

Definition 3.1.15. Let $E \in Spt(\mathcal{M})$ be a symmetric T -spectrum. We will say that E is n -orthogonal, if for all $K \in \Sigma_T^n \mathcal{SH}^{\text{eff}}$: $\text{Hom}_{\mathcal{SH}}(K, E) = 0$. Let $\mathcal{SH}^\perp(n)$ denote the full subcategory of \mathcal{SH} consisting of the n -orthogonal objects.

Theorem 3.1.16. *Let n be an arbitrary integer. Then the full triangulated subcategories of \mathcal{SH} : WB_n^\perp and $\mathcal{SH}^\perp(n+1)$ are identical (see 3.1.8(4)). In addition, if the base scheme $X = \text{Spec}(k)$, with k a perfect field, then the full triangulated subcategories of \mathcal{SH} : B_n^\perp and $\mathcal{SH}^\perp(n+1)$ are identical.*

Proof. By 3.1.9 and 2.2.11, WB_n^\perp is naturally isomorphic as triangulated category to $\mathbf{Ho}(WB_n \text{Spt}(\mathcal{M})) = \mathcal{SH}(WB_n)$. On the other hand, by [21, Thms. 1.4(2) and 3.6], $\mathcal{SH}^\perp(n+1)$ is also naturally isomorphic as triangulated category to $\mathbf{Ho}(WB_n \text{Spt}(\mathcal{M})) = \mathcal{SH}(WB_n)$.

Then, the result follows combining 2.2.9, 2.2.10 and [21, Prop. 3.5]. The case of a perfect field follows from Prop. 2.12 in [21]. \square

Recall that for $q > 0$, $\mathbb{G}_m^{-q} \wedge E$ is $F_q(S^q \wedge E)$ (see 3.1.3).

Corollary 3.1.17. *Let E be an arbitrary symmetric T -spectrum in \mathcal{SH} , and n, q be arbitrary integers. Then E is in WB_n^\perp if and only if $\mathbb{G}_m^q \wedge E$ is in WB_{n+q}^\perp .*

Proof. By 3.1.16, it follows that WB_n^\perp is equal to $\mathcal{SH}^\perp(n+1)$, which by definition is $(\Sigma_T^{n+1} \mathcal{SH}^{\text{eff}})^\perp$ (see 3.1.15). To conclude, we observe that the triangulated functor $\mathbb{G}_m^q : \mathcal{SH} \rightarrow \mathcal{SH}$, $E \mapsto \mathbb{G}_m^q \wedge E$; admits an inverse ($E \mapsto \mathbb{G}_m^{-q} \wedge E$) and maps $\Sigma_T^{n+1} \mathcal{SH}^{\text{eff}}$ surjectively onto $\Sigma_T^{n+q+1} \mathcal{SH}^{\text{eff}}$. \square

3.2. The weakly birational tower. In this section q will be an arbitrary integer. By construction there is an inclusion $sWB_{q+1} \subseteq sWB_q$ of sets of maps in $\text{Spt}(\mathcal{M})$ (see 3.1.5, 3.1.6, 3.1.7(1)), and hence an inclusion $\mathcal{G}_{WB_{q+1}} \subseteq \mathcal{G}_{WB_q}$ of sets of objects in $\text{Spt}(\mathcal{M})$ (see 3.1.8(1)).

Thus, we deduce that $\text{Loc}(\mathcal{G}_{WB_{q+1}}) \subseteq \text{Loc}(\mathcal{G}_{WB_q})$; and as a consequence (see 3.1.8(4)):

$$WB_q^\perp \subseteq WB_{q+1}^\perp.$$

which are localizing subcategories of \mathcal{SH} (see 2.2.12 and 3.1.8(4)). Therefore, we obtain the following tower of localizing subcategories in \mathcal{SH} :

$$(3.2.1) \quad \cdots \subseteq WB_{q-1}^\perp \subseteq WB_q^\perp \subseteq WB_{q+1}^\perp \subseteq \cdots$$

Proposition 3.2.2. *The inclusion, $j_q : WB_q^\perp \rightarrow \mathcal{SH}$ admits a right adjoint:*

$$p_q : \mathcal{SH} \rightarrow WB_q^\perp,$$

which is also a triangulated functor.

Proof. The result follows by combining 3.1.12 and theorem 4.1 in [15]. \square

Remark 3.2.3. Since the inclusion $j_q : WB_q^\perp \rightarrow \mathcal{SH}$ is a full embedding, we deduce that the unit of the adjunction $\text{id} \xrightarrow{\tau} p_q j_q$ is a natural isomorphism.

3.2.4. Weakly birational cover. We define $wb_q^c = j_q p_q$. Then clearly $wb_q^c \circ wb_{q+1}^c = wb_q^c$ and there exists a canonical natural transformation $wb_q^c \rightarrow wb_{q+1}^c$.

The following proposition is well-known.

Proposition 3.2.5. *The counit $wb_q^c = j_q p_q \xrightarrow{\theta_q} \text{id}$ of the adjunction constructed in 3.2.2 satisfies the following universal property:*

For any symmetric T -spectrum E in \mathcal{SH} and for any $F \in WB_q^\perp$, the map $\theta_q^E : wb_q^c E \rightarrow E$ in \mathcal{SH} induces an isomorphism of abelian groups:

$$\mathrm{Hom}_{\mathcal{SH}}(F, wb_q^c E) \xrightarrow[\cong]{\theta_q^E} \mathrm{Hom}_{\mathcal{SH}}(F, E)$$

Proof. If $F \in WB_q^\perp$, then $\mathrm{Hom}_{\mathcal{SH}}(F, E) = \mathrm{Hom}_{\mathcal{SH}}(j_q F, E)$. By adjointness:

$$\mathrm{Hom}_{\mathcal{SH}}(j_q F, E) \cong \mathrm{Hom}_{WB_q^\perp}(F, p_q E).$$

Since WB_q^\perp is a full subcategory of \mathcal{SH} , we deduce that:

$$\mathrm{Hom}_{WB_q^\perp}(F, p_q E) = \mathrm{Hom}_{\mathcal{SH}}(j_q F, j_q p_q E) = \mathrm{Hom}_{\mathcal{SH}}(F, wb_q^c E).$$

This finishes the proof. \square

Corollary 3.2.6. *Let E be an arbitrary symmetric T -spectrum in \mathcal{SH} . Then, the natural map $\theta_q^E : wb_q^c E \rightarrow E$ is an isomorphism in \mathcal{SH} if and only if E belongs to WB_q^\perp .*

Proof. By 2.2.12, we conclude that WB_q^\perp is a localizing subcategory of \mathcal{SH} .

First, we assume that θ_q^E is an isomorphism in \mathcal{SH} . Then, by construction $wb_q^c E$ is in WB_q^\perp (see 3.2.2 and 3.2.4). Hence, we deduce that E is also in WB_q^\perp since it is a localizing subcategory of \mathcal{SH} .

Finally, we assume that E is in WB_q^\perp which is a localizing subcategory of \mathcal{SH} . Thus, θ_q^E is a map in WB_q^\perp since $wb_q^c E$ is in WB_q^\perp by construction. Therefore, by 3.2.5 we deduce that θ_q^E is an isomorphism in WB_q^\perp , and hence an isomorphism in \mathcal{SH} . \square

Theorem 3.2.7. *There exist triangulated functors:*

$$wb_{q+1/q} : \mathcal{SH} \longrightarrow \mathcal{SH}$$

together with natural transformations:

$$\begin{aligned} \pi_{q+1} : wb_{q+1}^c &\longrightarrow wb_{q+1/q} \\ \sigma_{q+1} : wb_{q+1/q} &\longrightarrow \Sigma^1 wb_q^c \end{aligned}$$

such that for any symmetric T -spectrum E in \mathcal{SH} the following conditions hold:

(1) *There is a distinguished triangle in \mathcal{SH} :*

$$(3.2.8) \quad wb_q^c E \longrightarrow wb_{q+1}^c E \xrightarrow{\pi_{q+1}} wb_{q+1/q} E \xrightarrow{\sigma_{q+1}} \Sigma^1 wb_q^c E$$

(2) *$wb_{q+1/q} E$ is in WB_{q+1}^\perp .*

(3) *$wb_{q+1/q} E$ is in $(WB_q^\perp)^\perp$. Namely, for any F in WB_q^\perp :*

$$\mathrm{Hom}_{\mathcal{SH}}(F, wb_{q+1/q} E) = 0.$$

Proof. By 3.1.12, the triangulated categories WB_q^\perp and WB_{q+1}^\perp are compactly generated. Thus, the result follows from propositions 9.1.19 and 9.1.8 in [16]. \square

Theorem 3.2.9. *There exist triangulated functors:*

$$wb_{>q} : \mathcal{SH} \longrightarrow \mathcal{SH}$$

together with natural transformations:

$$\begin{aligned}\pi_{>q} : id &\longrightarrow wb_{>q} \\ \sigma_{>q} : wb_{>q} &\longrightarrow \Sigma^1 wb_q^c\end{aligned}$$

such that for any symmetric T -spectrum E in \mathcal{SH} the following conditions hold:

(1) There is a distinguished triangle in \mathcal{SH} :

$$(3.2.10) \quad wb_q^c E \longrightarrow E \xrightarrow{\pi_{>q}} wb_{>q} E \xrightarrow{\sigma_{>q}} \Sigma^1 wb_q^c E$$

(2) $wb_{>q} E$ is in $(WB_q^\perp)^\perp$. Namely, for any F in WB_q^\perp :

$$\mathrm{Hom}_{\mathcal{SH}}(F, wb_{>q} E) = 0.$$

Proof. By 3.1.1 and 3.1.12, the triangulated categories \mathcal{SH} and WB_q^\perp are compactly generated. Thus, the result follows from propositions 9.1.19 and 9.1.8 in [16]. \square

Theorem 3.2.11. *For any symmetric T -spectrum E in \mathcal{SH} , there exists the following commutative diagram in \mathcal{SH} :*

$$\begin{array}{ccccccc} wb_q^c E & \longrightarrow & wb_{q+1}^c E & \xrightarrow{\pi_{q+1}} & wb_{q+1/q} E & \xrightarrow{\sigma_{q+1}} & \Sigma^1 wb_q^c E \\ \parallel & & \downarrow & & \downarrow & & \parallel \\ wb_q^c E & \longrightarrow & E & \xrightarrow{\pi_{>q}} & wb_{>q} E & \xrightarrow{\sigma_{>q}} & \Sigma^1 wb_q^c E \\ \downarrow & & \downarrow \pi_{>q+1} & & \downarrow & & \downarrow \\ * & \longrightarrow & wb_{>q+1} E & \xlongequal{\quad} & wb_{>q+1} E & \longrightarrow & * \\ \downarrow & & \downarrow \sigma_{>q+1} & & \downarrow & & \downarrow \\ \Sigma^1 wb_q^c E & \longrightarrow & \Sigma^1 wb_{q+1}^c E & \xrightarrow{\Sigma^1 \pi_{q+1}} & \Sigma^1 wb_{q+1/q} E & \xrightarrow{\Sigma^1 \sigma_{q+1}} & \Sigma^2 wb_q^c E \end{array}$$

where all the rows and columns are distinguished triangles in \mathcal{SH} .

Proof. The result follows from 3.2.7, 3.2.9 and the octahedral axiom applied to the following commutative diagram:

$$\begin{array}{ccc} wb_q^c E & \longrightarrow & wb_{q+1}^c E \\ & \searrow & \swarrow \\ & E & \end{array}$$

\square

3.2.12. *The spectral sequence.* By 3.2.11, for every symmetric T -spectrum E in \mathcal{SH} there is a tower in \mathcal{SH} :

$$(3.2.13) \quad \begin{array}{ccccccc} \cdots & \longrightarrow & wb_{-1}^c(E) & \longrightarrow & wb_0^c(E) & \longrightarrow & wb_1^c(E) \longrightarrow \cdots \\ & & & \searrow \theta_{-1}^E & \downarrow \theta_0^E & \swarrow \theta_1^E & \\ & & & & E & & \end{array}$$

We will call (3.2.13) the weakly birational tower of E .

Remark 3.2.14. By 3.2.2 and 3.2.4, the weakly birational tower (3.2.13) is functorial with respect to morphisms in \mathcal{SH} .

Theorem 3.2.15. *Let G, K be arbitrary symmetric T -spectra in \mathcal{SH} . Then there is a spectral sequence of homological type with term $E_{p,q}^1 = \text{Hom}_{\mathcal{SH}}(G, \Sigma^{q-p} \text{wb}_{p/p-1} K)$ and where the abutment is given by the associated graded group for the filtration F_\bullet of $\text{Hom}_{\mathcal{SH}}(G, K)$ defined by the image of $\theta_{p*}^K : \text{Hom}_{\mathcal{SH}}(G, \text{wb}_p^c K) \rightarrow \text{Hom}_{\mathcal{SH}}(G, K)$, or equivalently the kernel of $\pi_{>q} : \text{Hom}_{\mathcal{SH}}(G, K) \rightarrow \text{Hom}_{\mathcal{SH}}(G, \text{wb}_{>q} K)$.*

Proof. Since \mathcal{SH} is a triangulated category, the result follows from 3.2.11 and 3.2.13. \square

Remark 3.2.16. The constructions of this section can be carried out as well using the tower (see 3.1.8):

$$(3.2.17) \quad \cdots \subseteq B_{q-1}^\perp \subseteq B_q^\perp \subseteq B_{q+1}^\perp \subseteq \cdots$$

and it follows from 3.1.16 that both constructions are canonically isomorphic when the base scheme X is of the form $\text{Spec}(k)$ with k a perfect field.

4. RATIONAL COEFFICIENTS

The main result of this section is 4.2.1, which describes the weakly birational tower with rational coefficients.

4.1. \mathbb{Q} -local spectra. Recall that a symmetric T -spectrum E in $\text{Spt}(\mathcal{M})$ is \mathbb{Q} -local if the abelian group $\text{Hom}_{\mathcal{SH}}(\Sigma^p(\mathbb{G}_m^q \wedge Y_+), E)$ is a \mathbb{Q} -vector space for every Y in Sm_X and every p, q in \mathbb{Z} [12, Remark 4.3.3], and that a map $f : E \rightarrow F$ in $\text{Spt}(\mathcal{M})$ is a *rational weak equivalence* if for every Y in Sm_X and every p, q in \mathbb{Z} the induced map $f_* \otimes \mathbb{Q} : \text{Hom}_{\mathcal{SH}}(\Sigma^p(\mathbb{G}_m^q \wedge Y_+), E) \otimes \mathbb{Q} \rightarrow \text{Hom}_{\mathcal{SH}}(\Sigma^p(\mathbb{G}_m^q \wedge Y_+), F) \otimes \mathbb{Q}$ is an isomorphism of \mathbb{Q} -vector spaces [12, Remark 4.3.3].

The work of Morel [12, §5.2 p.246 and Remark 4.3.3] shows that the left Bousfield localization of $\text{Spt}(\mathcal{M})$ with respect to the rational weak equivalences exists. We will write $\text{Spt}(\mathcal{M})_{\mathbb{Q}}$ for this left Bousfield localization, and $\mathcal{SH}_{\mathbb{Q}}$ for its homotopy category.

Recall that $\mathbf{1}$ is the sphere spectrum in \mathcal{SH} . The rational Moore spectrum $\mathbf{1}_{\mathbb{Q}}$ in $\text{Spt}(\mathcal{M})$ is defined to be the homotopy colimit of the filtering diagram:

$$\mathbf{1} \xrightarrow{2} \mathbf{1} \xrightarrow{3} \cdots \xrightarrow{n} \mathbf{1} \xrightarrow{n+1} \cdots$$

where $\mathbf{1} \xrightarrow{n} \mathbf{1}$ is the composition of the sum map with the diagonal $\mathbf{1} \xrightarrow{\Delta} \oplus_{i=1}^n \mathbf{1} \xrightarrow{\Sigma} \mathbf{1}$. Let $u : \mathbf{1} \rightarrow \mathbf{1}_{\mathbb{Q}}$ be the canonical map. A concrete model for a \mathbb{Q} -localization functor is given by $E \mapsto E \wedge \mathbf{1}_{\mathbb{Q}}$, i.e. $E \wedge \mathbf{1}_{\mathbb{Q}}$ is \mathbb{Q} -local in \mathcal{SH} and the map $E \wedge (\mathbf{1} \xrightarrow{u} \mathbf{1}_{\mathbb{Q}})$ is a rational weak equivalence.

Lemma 4.1.1. *Let E be a \mathbb{Q} -local symmetric T -spectrum in \mathcal{SH} . Then the map $E \wedge u : E \rightarrow E \wedge \mathbf{1}_{\mathbb{Q}}$ is an isomorphism in \mathcal{SH} .*

Proof. Since \mathcal{SH} is a compactly generated category in the sense of Neeman (see 3.1.1); it suffices to show that for every compact generator $G \in \mathcal{G}_{\mathcal{SH}}$ (see 3.1.2) the map induced by $E \wedge u$, $\text{Hom}_{\mathcal{SH}}(G, E) \rightarrow \text{Hom}_{\mathcal{SH}}(G, E \wedge \mathbf{1}_{\mathbb{Q}})$ is an isomorphism of abelian groups. We observe that $E \wedge \mathbf{1}_{\mathbb{Q}}$ is the homotopy colimit of the diagram:

$$E \xrightarrow{2} E \xrightarrow{3} \cdots \xrightarrow{n} E \xrightarrow{n+1} \cdots$$

Thus, by [15, Lem. 2.8] we just need to show that for every integer $n \geq 1$, the maps $\mathrm{Hom}_{\mathcal{SH}}(G, E) \xrightarrow{n} \mathrm{Hom}_{\mathcal{SH}}(G, E)$ are isomorphisms of abelian groups. This follows directly from the fact that E is \mathbb{Q} -local (see 4.1). \square

By construction, $\mathbf{1}_{\mathbb{Q}}$ is \mathbb{Q} -local in \mathcal{SH} . Thus, by 4.1.1 we conclude that $\mathbf{1}_{\mathbb{Q}} \wedge \mathbf{1}_{\mathbb{Q}} \cong \mathbf{1}_{\mathbb{Q}}$ in \mathcal{SH} . Hence, we deduce that $\mathbf{1}_{\mathbb{Q}}$ is a commutative ring spectrum in \mathcal{SH} ; with unit $u : \mathbf{1} \rightarrow \mathbf{1}_{\mathbb{Q}}$.

Remark 4.1.2. Combining [4, Prop. 1.4.3] and [4, p.99 (2.3.22.1)-(2.3.22.2)]; we can assume that $\mathbf{1}_{\mathbb{Q}}$ is a commutative ring spectrum with unit in $Spt(\mathcal{M})$.

Proposition 4.1.3. *The following conditions on a symmetric T -spectrum E are equivalent:*

- (1) E is \mathbb{Q} -local (see 4.1),
- (2) The map $E \wedge u : E \rightarrow E \wedge \mathbf{1}_{\mathbb{Q}}$ is an isomorphism in \mathcal{SH} .
- (3) E is a module in \mathcal{SH} over $\mathbf{1}_{\mathbb{Q}}$.
- (4) There exists a $\mathbf{1}_{\mathbb{Q}}$ -module in $Spt(\mathcal{M})$, E' which is isomorphic to E in \mathcal{SH} .

Proof. (1) \Rightarrow (2): This follows from 4.1.1.

(2) \Rightarrow (4): It suffices to show that $E' = E \wedge \mathbf{1}_{\mathbb{Q}}$ is a $\mathbf{1}_{\mathbb{Q}}$ -module in $Spt(\mathcal{M})$, and this holds since $\mathbf{1}_{\mathbb{Q}}$ is a commutative ring spectrum with unit in $Spt(\mathcal{M})$ (see 4.1.2).

(4) \Rightarrow (3): Obvious.

(3) \Rightarrow (1): Let $\mu_E : E \wedge \mathbf{1}_{\mathbb{Q}} \rightarrow E$ be the map inducing the $\mathbf{1}_{\mathbb{Q}}$ -module structure of E in \mathcal{SH} . We observe that the composition $\mu_E \circ (E \wedge u) : E \rightarrow E$ is the identity on E . Hence E is a direct summand of $E \wedge \mathbf{1}_{\mathbb{Q}}$, which is \mathbb{Q} -local by construction. Thus we conclude that E is also \mathbb{Q} -local, since this property is clearly closed under direct summands (see 4.1). \square

4.1.4. Orthogonality and weakly birational covers.

Lemma 4.1.5. *Let E be an arbitrary symmetric T -spectrum in \mathcal{SH} . Assume that E belongs to $\mathcal{SH}^{\perp}(q)$ for some integer $q \in \mathbb{Z}$ (see 3.1.15). Then $E \wedge \mathbf{1}_{\mathbb{Q}}$ also belongs to $\mathcal{SH}^{\perp}(q)$.*

Proof. Combining 3.1.15, 2.2.12 and 3.1.13 we deduce that $\mathcal{SH}^{\perp}(q)$ is a localizing subcategory of \mathcal{SH} . By construction, $E \wedge \mathbf{1}_{\mathbb{Q}}$ is the homotopy colimit of the diagram:

$$E \xrightarrow{2} E \xrightarrow{3} \dots \xrightarrow{n} E \xrightarrow{n+1} \dots$$

To conclude, we observe that by definition of the homotopy colimit (see [16, Def. 1.6.4]) any localizing subcategory (see 2.1) is closed under homotopy colimits. \square

Given an arbitrary integer $n \in \mathbb{Z}$, consider the weakly n -birational cover of E (see 3.2.4, 3.2.5), $\theta_n^E : wb_n^c(E) \rightarrow E$. Assume that E is \mathbb{Q} -local in \mathcal{SH} . By 4.1.3, we conclude that $E \wedge \mathbf{1}_{\mathbb{Q}} \cong E$ in \mathcal{SH} . Thus, taking the \mathbb{Q} -localization of θ_n^E we obtain a commutative triangle in \mathcal{SH} :

$$(4.1.6) \quad \begin{array}{ccc} wb_n^c(E) & \xrightarrow{\theta_n^E} & E \\ \downarrow wb_n^c(E) \wedge u & \nearrow \theta_n^E \wedge \mathbf{1}_{\mathbb{Q}} & \\ wb_n^c(E) \wedge \mathbf{1}_{\mathbb{Q}} & & \end{array}$$

Combining 4.1.5, 3.1.16 and the universal property of θ_n^E (see 3.2.5), we conclude that there is a map $r : wb_n^c(E) \wedge \mathbf{1}_{\mathbb{Q}} \rightarrow wb_n^c(E)$ such that the following triangle in \mathcal{SH} commutes:

$$(4.1.7) \quad \begin{array}{ccc} wb_n^c(E) & \xrightarrow{\theta_n^E} & E \\ r \uparrow & \nearrow \theta_n^E \wedge \mathbf{1}_{\mathbb{Q}} & \\ wb_n^c(E) \wedge \mathbf{1}_{\mathbb{Q}} & & \end{array}$$

Lemma 4.1.8. *Let E be \mathbb{Q} -local spectrum in \mathcal{SH} . Then the composition map $r \circ (wb_n^c(E) \wedge u) : wb_n^c(E) \rightarrow wb_n^c(E)$ is the identity on $wb_n^c(E)$, i.e. $wb_n^c(E)$ is a direct summand of $wb_n^c(E) \wedge \mathbf{1}_{\mathbb{Q}}$. Hence, $wb_n^c(E)$ is \mathbb{Q} -local and the map $wb_n^c(E) \wedge u : wb_n^c(E) \rightarrow wb_n^c(E) \wedge \mathbf{1}_{\mathbb{Q}}$ is an isomorphism in \mathcal{SH} .*

Proof. Since the direct summand of a \mathbb{Q} -local object in \mathcal{SH} is \mathbb{Q} -local (see 4.1), by 4.1.3 it suffices to show that $r \circ (wb_n^c(E) \wedge u) = id$. This follows from the universal property (3.2.5) of θ_n^E and the commutativity of (4.1.7), (4.1.6). \square

4.1.9. *Voevodsky's slice tower for orthogonal spectra.* Recall that the q -effective motivic stable homotopy category $\Sigma_T^q \mathcal{SH}^{eff}$ (see 3.1.13) is the localizing subcategory of \mathcal{SH} , of the form $Loc(C_{eff}^q)$ (see 2.1.2 and 3.1.14). Voevodsky [32] defines the slice filtration in \mathcal{SH} to be the family of localizing subcategories:

$$\dots \subseteq \Sigma_T^{q+1} \mathcal{SH}^{eff} \subseteq \Sigma_T^q \mathcal{SH}^{eff} \subseteq \Sigma_T^{q-1} \mathcal{SH}^{eff} \subseteq \dots$$

The work of Neeman [15], [16] shows that the inclusion $\Sigma_T^q \mathcal{SH}^{eff} \xrightarrow{i_q} \mathcal{SH}$ admits a right adjoint $\mathcal{SH} \xrightarrow{r_q} \Sigma_T^q \mathcal{SH}^{eff}$ [19, Prop. 3.1.12]; and that the functors, $\mathcal{SH} \xrightarrow{f_q} \mathcal{SH}$, $\mathcal{SH} \xrightarrow{s_q} \mathcal{SH}$ are triangulated, where f_q is defined as the composition $i_q \circ r_q$; and s_q is characterized by the fact that for every $E \in \mathcal{SH}$, there exists a distinguished triangle in \mathcal{SH} [19, Thms. 3.1.16, 3.1.18]:

$$(4.1.10) \quad f_{q+1}E \rightarrow f_qE \rightarrow s_qE.$$

We will refer to f_qE as the $(q-1)$ -connective cover of E , and to s_qE as the q -slice of E .

Remark 4.1.11. It follows directly from the definition that s_qE is in $\mathcal{SH}^{\perp}(q+1)$.

Remark 4.1.12. Since \mathcal{SH} is compactly generated in the sense of Neeman (see 3.1.1) with set of compact generators $\mathcal{G}_{\mathcal{SH}}$ (see 3.1.2), and by construction $\mathcal{G}_{\mathcal{SH}}$ is the union of C_{eff}^q for $q \in \mathbb{Z}$ (see 3.1.14), we conclude that E is the homotopy colimit of its slice tower: $\dots \rightarrow f_nE \rightarrow f_{n-1}E \rightarrow f_{n-2}E \rightarrow \dots$.

The sphere spectrum $\mathbf{1}$ is contained in the localizing subcategory \mathcal{SH}^{eff} (see 3.1.13). Thus, $\mathbf{1}_{\mathbb{Q}}$ is also contained in \mathcal{SH}^{eff} since localizing subcategories are closed under homotopy colimits. Consider the triangulated functor, $-\wedge \mathbf{1}_{\mathbb{Q}} : \mathcal{SH} \rightarrow \mathcal{SH}$.

Theorem 4.1.13. *Let E be an arbitrary symmetric T -spectrum in \mathcal{SH} , and $p \in \mathbb{Z}$ an arbitrary integer. Then $s_p(E \wedge \mathbf{1}_{\mathbb{Q}}) \cong s_p(E) \wedge \mathbf{1}_{\mathbb{Q}}$.*

Proof. Since $\mathbf{1}_{\mathbb{Q}}$ belongs to \mathcal{SH}^{eff} , we deduce that $(\Sigma_T^q \mathcal{SH}^{eff}) \wedge \mathbf{1}_{\mathbb{Q}} \subseteq (\Sigma_T^q \mathcal{SH}^{eff})$ for every integer $q \in \mathbb{Z}$. Furthermore, it is clear that the functor $-\wedge \mathbf{1}_{\mathbb{Q}}$ commutes with filtered homotopy colimits.

Therefore, by [22, Rmk. 2.13 and Thm. 2.12] it suffices to show that $s_q(E) \wedge \mathbf{1}_{\mathbb{Q}}$ is in $\mathcal{SH}^\perp(q+1)$ for every integer $q \in \mathbb{Z}$. This follows from 4.1.5 since by construction $s_q(E)$ belongs to $\mathcal{SH}^\perp(q+1)$ (see 4.1.11). \square

Proposition 4.1.14. *Let E be a \mathbb{Q} -local symmetric T -spectrum in \mathcal{SH} . Then for every integer $p \in \mathbb{Z}$, $s_p E$ is a module in \mathcal{SH} over $s_0(\mathbf{1}) \wedge \mathbf{1}_{\mathbb{Q}}$.*

Proof. We observe that $E \cong E \wedge \mathbf{1}_{\mathbb{Q}}$ in \mathcal{SH} , since E is \mathbb{Q} -local in \mathcal{SH} (see 4.1.3). Thus, by 4.1.13 we deduce that $s_p E \cong s_p(E \wedge \mathbf{1}_{\mathbb{Q}}) \cong s_p(E) \wedge \mathbf{1}_{\mathbb{Q}}$ in \mathcal{SH} .

To finish the proof, it suffices to show that $s_p E$ is a module in \mathcal{SH} over $s_0 \mathbf{1}$. This follows from [19, Thm. 3.6.14(6)]. \square

Let $\mathbf{H}_{\mathfrak{B}} = KGL^{(0)} \in Spt(\mathcal{M})$ denote the Beilinson motivic cohomology spectrum constructed by Riou in [26]. By the work of Cisinski-Dégliise [3, Cor. 14.2.6], we conclude that $\mathbf{H}_{\mathfrak{B}}$ is a commutative cofibrant ring spectrum in $Spt(\mathcal{M})$. In addition, in [20, Thm. 4.1] we show that $s_0(\mathbf{1}) \wedge \mathbf{1}_{\mathbb{Q}}$ is equipped with a unique structure of $\mathbf{H}_{\mathfrak{B}}$ -algebra in $Spt(\mathcal{M}_X)$.

Corollary 4.1.15. *Let E be a \mathbb{Q} -local symmetric T -spectrum in \mathcal{SH} . Then for every integer $p \in \mathbb{Z}$, $s_p E$ is a module in $Spt(\mathcal{M})$ over $\mathbf{H}_{\mathfrak{B}}$.*

Proof. Combining 4.1.14 and [20, Thm. 4.1], we deduce that $s_p E$ is a module in \mathcal{SH} over $\mathbf{H}_{\mathfrak{B}}$. Then the result follows from [3, Cor 14.2.16(i)-(v)], since E is \mathbb{Q} -local in \mathcal{SH} and hence $s_p E \cong s_p(E \wedge \mathbf{1}_{\mathbb{Q}}) \cong s_p(E) \wedge \mathbf{1}_{\mathbb{Q}}$ (see 4.1.13) is also \mathbb{Q} -local. \square

Let $e : \mathbf{1} \rightarrow \mathbf{H}_{\mathfrak{B}}$ be the unit map of the ring spectrum $\mathbf{H}_{\mathfrak{B}}$ in $Spt(\mathcal{M})$.

Theorem 4.1.16. *Let E be a \mathbb{Q} -local symmetric T -spectrum in \mathcal{SH} . Assume that E belongs to the localizing subcategory $\mathcal{SH}^\perp(n)$ for some integer $n \in \mathbb{Z}$. Then E is a module in $Spt(\mathcal{M})$ over $\mathbf{H}_{\mathfrak{B}}$.*

Proof. By [3, Cor 14.2.16(i)-(v)], it suffices to show that $E \wedge e : E \rightarrow E \wedge \mathbf{H}_{\mathfrak{B}}$ is an isomorphism in \mathcal{SH} . Since $E \cong \text{hocolim}_{p \leq n} f_p E$ in \mathcal{SH} (see 4.1.12), it suffices to show that $f_p E \wedge e$ is an isomorphism in \mathcal{SH} for every integer $p \leq n$.

We observe that $f_p E \cong * \cong s_p E$ for $p \geq n$, since by hypothesis E is in $\mathcal{SH}^\perp(n)$. Hence, the slice tower of E is of the form:

$$* = f_n E \rightarrow s_{n-1} E \cong f_{n-1} E \rightarrow f_{n-2} E \rightarrow f_{n-3} E \rightarrow \cdots$$

Thus, by 4.1.10 it suffices to show that $s_p E \wedge e$ is an isomorphism in \mathcal{SH} for every integer $p \leq n$. This follows from 4.1.15 and [3, Cor 14.2.16(i)-(v)]. \square

4.2. The weakly birational tower for \mathbb{Q} -local spectra. Given symmetric T -spectrum E in \mathcal{SH} , we consider its weakly birational tower (3.2.13):

$$\begin{array}{ccccccc} \cdots & \longrightarrow & wb_{-1}^c(E) & \longrightarrow & wb_0^c(E) & \longrightarrow & wb_1^c(E) \longrightarrow \cdots \\ & & \searrow \theta_{-1}^E & & \downarrow \theta_0^E & & \swarrow \theta_1^E \\ & & & & E & & \end{array}$$

Recall that $wb_n^c E$ is the weakly n -birational cover of E (see 3.2.4, 3.2.5).

Theorem 4.2.1. *Let E be a \mathbb{Q} -local symmetric T -spectrum in \mathcal{SH} (see 4.1), and $n \in \mathbb{Z}$ an arbitrary integer. Then $wb_n^c E$ satisfies the following properties:*

- (1) $wb_n^c E$ is \mathbb{Q} -local in \mathcal{SH} .

- (2) $wb_n^c E$ is a module in $Spt(\mathcal{M})$ over $\mathbf{H}_{\mathfrak{B}}$.
 (3) The natural map $wb_n^c E \rightarrow wb_{n+1}^c E$ is a map of $\mathbf{H}_{\mathfrak{B}}$ -modules in $Spt(\mathcal{M})$.

Proof. (1): This follows from 4.1.8.

(2): By construction $wb_n^c E$ is in WB_n^\perp (see 3.2.2 and 3.2.4). Hence, the result follows by combining 4.1.16 and 3.1.16.

(3): By [3, Cor. 14.2.6] $\mathbf{H}_{\mathfrak{B}}$ is a commutative ring in $Spt(\mathcal{M})$, with unit $u : \mathbf{1} \rightarrow \mathbf{H}_{\mathfrak{B}}$. Thus the bottom row in the following commutative diagram in \mathcal{SH} is a map of $\mathbf{H}_{\mathfrak{B}}$ -modules in $Spt(\mathcal{M})$:

$$\begin{array}{ccc} wb_n^c E & \longrightarrow & wb_{n+1}^c E \\ (wb_n^c E) \wedge e \downarrow & & \downarrow (wb_{n+1}^c E) \wedge e \\ wb_n^c E \wedge \mathbf{H}_{\mathfrak{B}} & \longrightarrow & wb_{n+1}^c E \wedge \mathbf{H}_{\mathfrak{B}} \end{array}$$

Hence it suffices to show that the vertical maps are isomorphisms in \mathcal{SH} . This follows from [3, Cor 14.2.16(i)-(v)], since by (2) above $wb_n^c E$ and $wb_{n+1}^c E$ are modules in $Spt(\mathcal{M})$ over $\mathbf{H}_{\mathfrak{B}}$. \square

5. THE WEAKLY BIRATIONAL TOWER FOR MOTIVIC COHOMOLOGY

Throughout this section, the base scheme X will be of the form $\text{Spec}(k)$ with k a perfect field. We will study the weakly birational tower (3.2.13) for Voevodsky's T -spectrum HR , which represents in \mathcal{SH} motivic cohomology [29, 6.1] with coefficients in an abelian group or a commutative ring R . Our goal is to show that in some cases the tower is not trivial, and that it induces an interesting finite filtration on the Chow groups.

5.1. Main properties.

Lemma 5.1.1. *HR belongs to WB_0^\perp and to \mathcal{SH}^{eff} .*

Proof. By 3.1.11 and [21, Def. 3.1 and Thm. 3.7], we are reduced to show that HR is isomorphic in \mathcal{SH} to its zero slice $s_0(HR)$ [32]. This follows from [34] in characteristic zero, and [11, Lem. 10.4.1] in positive characteristic. \square

Proposition 5.1.2. *Let $n \geq 0$ be an arbitrary integer. Then the natural map (see 3.2.13), $\theta_n^{HR} : wb_n^c(HR) \rightarrow HR$ is an isomorphism in \mathcal{SH} .*

Proof. Combining 3.2.6 and 3.2.1, it suffices to show that HR is in WB_0^\perp . This follows from 5.1.1. \square

Hence, we conclude that the weakly birational tower (3.2.13) for motivic cohomology with R coefficients is as follows:

$$(5.1.3) \quad \begin{array}{ccccccc} \cdots & \longrightarrow & wb_{-3}^c(HR) & \longrightarrow & wb_{-2}^c(HR) & \longrightarrow & wb_{-1}^c(HR) \\ & & & \searrow \theta_{-3}^{HR} & \searrow \theta_{-2}^{HR} & & \downarrow \theta_{-1}^{HR} \\ & & & & & & HR \end{array}$$

5.1.4. *Non-triviality of the tower.* Our goal is to show that for $n < 0$, the weakly birational covers $wb_n^c(HR)$ and the layers $wb_{n/(n-1)}(HR)$ are interesting, i.e. they are not isomorphic in \mathcal{SH} to either HR or $*$.

First we introduce some notation; let $HR^{p,q}$, A_{HR} denote $\Sigma^{p-q} \wedge \mathbb{G}_m^q \wedge HR$ and $\bigoplus_{p,q \in \mathbb{Z}} HR^{p,q}$, respectively. By adjointness, there are isomorphisms:

$$(5.1.5) \quad \mathrm{Hom}_{\mathcal{SH}}(HR^{p,q}, HR) \cong \mathrm{Hom}_{\mathcal{SH}}(HR, HR^{-p,-q}) = \mathcal{A}^{-p,-q}$$

$$(5.1.6) \quad \mathrm{Hom}_{\mathcal{SH}}(A_{HR}, HR) \cong \prod_{p,q \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{SH}}(HR, HR^{-p,-q}) = \prod_{p,q \in \mathbb{Z}} \mathcal{A}^{-p,-q}$$

where $\mathcal{A}^{-p,-q}$ is the group of bistable operations in motivic cohomology with R coefficients of degree $-p$ and weight $-q$ [33].

The non-triviality of $wb_n^c(HR)$ and $wb_{n/(n-1)}(HR)$ will follow from the existence of non-trivial elements in $\mathcal{A}^{p,q}$ for appropriate p and q . This information is codified in the spectral sequence of the tower (5.1.3) evaluated in A_{HR} . Namely:

Theorem 5.1.7. *Let $p \leq 0$. Consider the spectral sequence of the tower (5.1.3) with term (see 3.2.15):*

$$E_{p,q}^1 = \mathrm{Hom}_{\mathcal{SH}}(A_{HR}, \Sigma^{q-p} wb_{p/p-1} HR) = \prod_{j \in \mathbb{Z}} \mathcal{A}^{j,-p}$$

given by all bistable operations in motivic cohomology with R coefficients of weight $-p$. Then the spectral sequence degenerates. Therefore, the term $E_{p,q}^1$ is the associated graded $gr_p F_\bullet$ for the filtration F_\bullet of $\mathrm{Hom}_{\mathcal{SH}}(A_{HR}, HR) = \prod_{j,k \in \mathbb{Z}} \mathcal{A}^{-j,-k}$ given by the image of

$$\theta_{n*}^{HR} : \mathrm{Hom}_{\mathcal{SH}}(A_{HR}, wb_n^c HR) \rightarrow \mathrm{Hom}_{\mathcal{SH}}(A_{HR}, HR),$$

which can be written explicitly as $\prod_{j \in \mathbb{Z}, k \leq n} \mathcal{A}^{-j,-k}$, i.e. coincides with all bistable operations in motivic cohomology with R coefficients of weight $\geq -n$.

Proof. The existence of the spectral sequence with term

$$E_{p,q}^1 = \mathrm{Hom}_{\mathcal{SH}}(A_{HR}, \Sigma^{q-p} wb_{p/p-1} HR),$$

follows from 3.2.15. Combining 5.3.2, 5.3.3 and 5.3.5; we deduce that the spectral sequence degenerates and that (see 5.1.5 and 5.1.6):

$$E_{p,q}^1 = \prod_{j \in \mathbb{Z}} \mathrm{Hom}_{\mathcal{SH}}(HR^{j,p}, HR) = \prod_{j \in \mathbb{Z}} \mathcal{A}^{j,-p}$$

Since the spectral sequence degenerates, we conclude that $E_{p,q}^1$ coincides with the associated graded $gr_p F_\bullet$ for the filtration given by the image of θ_{n*}^{HR} . Finally, combining 5.3.2, 5.3.3 and (5.1.5)-(5.1.6) we deduce that the image of θ_{n*}^{HR} is $\prod_{j \in \mathbb{Z}, k \leq n} \mathcal{A}^{-j,-k}$. \square

Theorem 5.1.8. *Let $q > 0$ be an arbitrary integer. Assume that there is an integer p such that $\mathcal{A}^{p,q} \neq 0$, i.e. there exist non-trivial bistable motivic operations of degree p and weight q . Then:*

- (1) *For every integer $0 < n \leq q$, $wb_{-n}^c HR$ is not isomorphic to $*$. In addition, the natural map $\theta_{-n}^{HR} : wb_{-n}^c HR \rightarrow HR$ is not an isomorphism in \mathcal{SH} .*
- (2) *The layer $wb_{-q/(-q-1)} HR$ is not isomorphic to $*$ in \mathcal{SH} .*

Proof. (1): On the one hand, combining 5.3.2 with 5.1.5, we conclude that $wb_{-n}^c HR$ and $*$ are not isomorphic in \mathcal{SH} . On the other hand, by 5.3.4 we deduce that θ_{-n}^{HR} is not an isomorphism in \mathcal{SH} .

(2): This follows by combining 5.3.2 and 5.3.5. \square

5.2. The filtration in the motivic cohomology of a scheme. Recall that X is of the form $\text{Spec}(k)$ with k a perfect field and that $H^{p,q} = \Sigma^{p-q} \wedge \mathbb{G}_m^q \wedge HR$.

Let $Y \in Sm_X$. We will write $H^{p,q}(Y, R)$ for $\text{Hom}_{\mathcal{SH}}(\Sigma_T^\infty Y_+, HR^{p,q})$, i.e. for the motivic cohomology of Y with coefficients in R of degree p and weight q . By adjointness: $H^{p,q}(Y, R) \cong \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), HR)$. Therefore, the abutment of the spectral sequence of the tower (5.1.3) evaluated in $\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+)$ induces an interesting filtration in $H^{p,q}(Y, R)$. Namely:

Theorem 5.2.1. *Let p, q be arbitrary integers, and let Y be in Sm_X . Then there exists a decreasing filtration F^\bullet on $H^{p,q}(Y, R)$ where the n -component is given by the image of θ_{-n*}^{HR} :*

$$\begin{array}{c} \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), wb_{-n}^c HR) \\ \downarrow \theta_{-n*}^{HR} \\ \text{Hom}_{\mathcal{SH}}(\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+), HR) = H^{p,q}(Y, R). \end{array}$$

In addition, the filtration F^\bullet is functorial in Y with respect to morphisms in Sm_X .

Proof. The existence of the filtration follows directly from 3.2.15. The functoriality is clear from the definition. \square

5.2.2. Finiteness of the filtration. Our goal is to show that for Y in Sm_X , the filtration F^\bullet on $H^{p,q}(Y, R)$ defined in 5.2.1 is concentrated in the range $0 \leq n \leq q$.

Theorem 5.2.3. *Let p, q be arbitrary integers, and let Y be in Sm_X . Then the decreasing filtration F^\bullet on $H^{p,q}(Y, R)$ constructed in 5.2.1 satisfies the following properties:*

- (1) $F^0 H^{p,q}(Y, R) = H^{p,q}(Y, R)$,
- (2) $F^{q+1} H^{p,q}(Y, R) = 0$.

Proof. By 5.1.2 we deduce that θ_0^{HR} is an isomorphism in \mathcal{SH} . This proves the first claim. For the second claim, we observe that $\Sigma^{-p+q}(\mathbb{G}_m^{-q} \wedge Y_+)$ is in $\Sigma_T^{-q} \mathcal{SH}^{eff}$ (see 3.1.14). Thus, it suffices to show that $wb_{-q-1}^c HR$ belongs to $\mathcal{SH}^\perp(-q)$ (see 3.1.15), or equivalently to WB_{-q-1}^\perp (see 3.1.16). This follows from 3.2.2 and 3.2.4. \square

5.3. Auxiliary results.

Lemma 5.3.1. *Let $q > 0$ be an arbitrary integer. Then, for every integer p : $\text{Hom}_{\mathcal{SH}}(HR^{p,q}, HR) = 0$.*

Proof. By 5.1.1, we conclude that $HR^{p,q}$ is in $\Sigma_T^q \mathcal{SH}^{eff}$. Combining 3.1.16 and 3.1.15, we observe that it suffices to show that HR is in WB_{q-1}^\perp . But this follows from 3.2.6 and 5.1.2. \square

Lemma 5.3.2. *Let $q \leq n$ be arbitrary integers. Then for every integer p , the natural map; $\theta_n^{HR} : wb_n^c HR \rightarrow HR$ induces an isomorphism:*

$$\text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_n^c HR) \rightarrow \text{Hom}_{\mathcal{SH}}(HR^{p,q}, HR).$$

Proof. By 3.2.5 and 3.2.1 it suffices to show that $HR^{p,q}$ is in WB_q^\perp . Then the result follows by combining 5.1.1 and 3.1.17. \square

Lemma 5.3.3. *Let $q \geq n + 1$ be arbitrary integers. Then, for every integer p : $\text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_n^c HR) = 0$.*

Proof. By construction $wb_n^c HR$ is in WB_n^\perp (see 3.2.2 and 3.2.4), and therefore also in WB_{q-1}^\perp (see 3.2.1). Combining 3.1.16 and 3.1.15, we conclude that it suffices to show that $HR^{p,q}$ is in $\Sigma_T^q \mathcal{SH}^{eff}$. This follows from 5.1.1. \square

Corollary 5.3.4. *Let $n < 0$ be an arbitrary integer. Then the natural map $\theta_n^{HR} : wb_n^c HR \rightarrow HR$ is not an isomorphism in \mathcal{SH} .*

Proof. Taking $p = q = 0$ in 5.3.3 we deduce that the identity map on HR does not factor through θ_n^{HR} . Hence we conclude that θ_n^{HR} is not an isomorphism in \mathcal{SH} . \square

Consider the distinguished triangle in \mathcal{SH} (see 3.2.7):

$$wb_n^c HR \longrightarrow wb_{n+1}^c HR \xrightarrow{\pi_{n+1}} wb_{n+1/n} HR \xrightarrow{\sigma_{n+1}} \Sigma^1 wb_n^c HR$$

Lemma 5.3.5. *Let q be an arbitrary integer. Then, the natural map π_q induces an isomorphism:*

$$\text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_q^c HR) \rightarrow \text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_{q/(q-1)} HR).$$

Proof. Since \mathcal{SH} is a triangulated category, it suffices to show that

$$\text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_{q-1}^c HR) = \text{Hom}_{\mathcal{SH}}(HR^{p,q}, \Sigma^1 wb_{q-1}^c HR) = 0.$$

On the one hand, by 5.3.3 we conclude that $\text{Hom}_{\mathcal{SH}}(HR^{p,q}, wb_{q-1}^c HR) = 0$. On the other hand, by adjointness and 5.3.3 we conclude that:

$$\text{Hom}_{\mathcal{SH}}(HR^{p,q}, \Sigma^1 wb_{q-1}^c HR) \cong \text{Hom}_{\mathcal{SH}}(HR^{p-1,q}, wb_{q-1}^c HR) = 0.$$

This finishes the proof. \square

ACKNOWLEDGEMENTS

The author would like to thank warmly Chuck Weibel for his interest in this work, as well as for all the advice and support during all these years.

REFERENCES

- [1] A. A. Beilinson. Height pairing between algebraic cycles. In *K-theory, arithmetic and geometry (Moscow, 1984-1986)*, volume 1289 of *Lecture Notes in Math.*, pages 1–25. Springer, Berlin, 1987.
- [2] S. Bloch. *Lectures on algebraic cycles*. Duke University Mathematics Series, IV. Duke University Mathematics Department, Durham, N.C., 1980.
- [3] D.-C. Cisinski and F. Déglise. Triangulated categories of mixed motives. 2009.
- [4] D.-C. Cisinski and F. Déglise. Mixed Weil cohomologies. *Adv. Math.*, 230(1):55–130, 2012.
- [5] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [6] J. Hornbostel. Localizations in motivic homotopy theory. *Math. Proc. Cambridge Philos. Soc.*, 140(1):95–114, 2006.
- [7] M. Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [8] M. Hovey. Spectra and symmetric spectra in general model categories. *J. Pure Appl. Algebra*, 165(1):63–127, 2001.

- [9] U. Jannsen. Motivic sheaves and filtrations on Chow groups. In *Motives (Seattle, WA, 1991)*, volume 55 of *Proc. Sympos. Pure Math.*, pages 245–302. Amer. Math. Soc., Providence, RI, 1994.
- [10] J. F. Jardine. Motivic symmetric spectra. *Doc. Math.*, 5:445–553 (electronic), 2000.
- [11] M. Levine. The homotopy coniveau tower. *J. Topol.*, 1(1):217–267, 2008.
- [12] F. Morel. On the motivic π_0 of the sphere spectrum. In *Axiomatic, enriched and motivic homotopy theory*, volume 131 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 219–260. Kluwer Acad. Publ., Dordrecht, 2004.
- [13] F. Morel and V. Voevodsky. \mathbf{A}^1 -homotopy theory of schemes. *Inst. Hautes Études Sci. Publ. Math.*, (90):45–143 (2001), 1999.
- [14] J. P. Murre. On a conjectural filtration on the Chow groups of an algebraic variety. I. The general conjectures and some examples. *Indag. Math. (N.S.)*, 4(2):177–188, 1993.
- [15] A. Neeman. The Grothendieck duality theorem via Bousfield’s techniques and Brown representability. *J. Amer. Math. Soc.*, 9(1):205–236, 1996.
- [16] A. Neeman. *Triangulated categories*, volume 148 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2001.
- [17] A. Neeman. Colocalizing subcategories of $\mathbf{D}(R)$. *J. Reine Angew. Math.*, 653:221–243, 2011.
- [18] Y. P. Nesterenko and A. A. Suslin. Homology of the general linear group over a local ring, and Milnor’s K -theory. *Izv. Akad. Nauk SSSR Ser. Mat.*, 53(1):121–146, 1989.
- [19] P. Pelaez. Multiplicative properties of the slice filtration. *Astérisque*, (335):xvi+289, 2011.
- [20] P. Pelaez. On the orientability of the slice filtration. *Homology Homotopy Appl.*, 13(2):293–300, 2011.
- [21] P. Pelaez. Birational motivic homotopy theories and the slice filtration. *Doc. Math.*, 18:51–70, 2013.
- [22] P. Pelaez. On the functoriality of the slice filtration. *J. K-Theory*, 11(1):55–71, 2013.
- [23] P. Pelaez. The unstable slice filtration. *Trans. Amer. Math. Soc.*, 366(11):5991–6025, 2014.
- [24] D. Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [25] D. G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
- [26] J. Riou. Algebraic K -theory, \mathbf{A}^1 -homotopy and Riemann-Roch theorems. *J. Topol.*, 3(2):229–264, 2010.
- [27] A. Suslin and V. Voevodsky. Bloch-Kato conjecture and motivic cohomology with finite coefficients. In *The arithmetic and geometry of algebraic cycles (Banff, AB, 1998)*, volume 548 of *NATO Sci. Ser. C Math. Phys. Sci.*, pages 117–189. Kluwer Acad. Publ., Dordrecht, 2000.
- [28] A. Suslin and V. Voevodsky. Relative cycles and Chow sheaves. In *Cycles, transfers, and motivic homology theories*, volume 143 of *Ann. of Math. Stud.*, pages 10–86. Princeton Univ. Press, Princeton, NJ, 2000.
- [29] V. Voevodsky. \mathbf{A}^1 -homotopy theory. In *Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998)*, number Extra Vol. I, pages 579–604 (electronic), 1998.
- [30] V. Voevodsky. Triangulated categories of motives over a field. In *Cycles, transfers, and motivic homology theories*, volume 143 of *Ann. of Math. Stud.*, pages 188–238. Princeton Univ. Press, Princeton, NJ, 2000.
- [31] V. Voevodsky. Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic. *Int. Math. Res. Not.*, (7):351–355, 2002.
- [32] V. Voevodsky. Open problems in the motivic stable homotopy theory. I. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 3–34. Int. Press, Somerville, MA, 2002.
- [33] V. Voevodsky. Reduced power operations in motivic cohomology. *Publ. Math. Inst. Hautes Études Sci.*, (98):1–57, 2003.
- [34] V. Voevodsky. On the zero slice of the sphere spectrum. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):106–115, 2004.
- [35] V. Voevodsky. Cancellation theorem. *Doc. Math.*, (Extra volume: Andrei A. Suslin sixtieth birthday):671–685, 2010.
- [36] C. A. Weibel. *The K-book*, volume 145 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K -theory.

INSTITUTO DE MATEMÁTICAS, CIUDAD UNIVERSITARIA, UNAM, DF 04510, MÉXICO
E-mail address: `pablo.pelaez@im.unam.mx`