

# Multiply Warped Products with a Quarter-symmetric Connection

Quan Qu, Yong Wang \*

## Abstract

In this paper, we study the Einstein warped products and multiply warped products with a quarter-symmetric connection. We also study warped products and multiply warped products with a quarter-symmetric connection with constant scalar curvature. Then apply our results to generalized Robertson-Walker space-times with a quarter-symmetric connection and generalized Kasner space-times with a quarter-symmetric connection.

**Keywords:** Warped products; multiply warped products; quarter-symmetric connection; Ricci tensor; scalar curvature; Einstein manifolds.

## 1 Introduction

The (singly) warped product  $B \times_b F$  of two pseudo-Riemannian manifolds  $(B, g_B)$  and  $(F, g_F)$  with a smooth function  $b : B \rightarrow (0, \infty)$  is a product manifold of form  $B \times F$  with the metric tensor  $g = g_B \oplus b^2 g_F$ . Here,  $(B, g_B)$  is called the base manifold and  $(F, g_F)$  is called as the fiber manifold and  $b$  is called as the warping function. The concept of warped products was first introduced by Bishop and O'Neill [1] to construct examples of Riemannian manifolds with negative curvature. In [2], F. Dobarro and E. Dozo had studied the problem of showing when a Riemannian metric of constant scalar curvature can be produced on a product manifolds by a warped product construction from the viewpoint of partial differential equations and variational methods. In [3],

---

\*Corresponding author.

E-mail addresses: quq453@nenu.edu.cn (Q. Qu), wangy581@nenu.edu.cn (Y. Wang).

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin, 130024, China

Ehrlich, Jung and Kim got explicit solutions to warping function to have a constant scalar curvature for generalized Robertson-Walker space-times. In [4], explicit solutions were also obtained for the warping function to make the space-time as Einstein when the fiber is also Einstein.

One can generalize (singly) warped products to multiply warped products. A multiply warped product  $(M, g)$  is the product manifold  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  with the metric  $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ , where for each  $i \in \{1, \dots, m\}$ ,  $b_i : B \rightarrow (0, \infty)$  is smooth and  $(F_i, g_{F_i})$  is a pseudo-Riemannian manifold. In particular, when  $B = (c, d)$ , the metric  $g_B = -dt^2$  is negative and  $(F_i, g_{F_i})$  is a Riemannian manifold, we call  $M$  as the multiply generalized Robertson-Walker space-time.

Singly warped products have a natural generalization. A twisted product  $(M, g)$  is a product manifold of form  $M = B \times_b F$ , with a smooth function  $b : B \times F \rightarrow (0, \infty)$ , and the metric tensor  $g = g_B \oplus b^2 g_F$ . In [5], they showed that mixed Ricci-flat twisted products could be expressed as warped products. As a consequence, any Einstein twisted products are warped products. Similar to the definition of multiply warped product, a multiply twisted product  $(M, g)$  is a product manifold of form  $M = B \times b_1 F_1 \times b_2 F_2 \cdots \times b_m F_m$  with the metric  $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ , where for each  $i \in \{1, \dots, m\}$ ,  $b_i : B \times F_i \rightarrow (0, \infty)$  is smooth. So in this paper, we define the multiply twisted products as generalizations of twisted products and multiply warped products.

The definition of a semi-symmetric metric connection was given by H. Hayden in [6]. In 1970, K. Yano [7] considered a semi-symmetric metric connection and studied some of its properties. Then in 1975, Golab [8] introduced the idea of a quarter-symmetric linear connection in differentiable manifold which is a generalization of semi-symmetric connection. A linear connection  $\nabla$  on an  $n$ -dimensional Riemannian manifold  $(M, g)$  is called a quarter-symmetric connection if its torsion tensor  $T$  of the connection  $\nabla$  satisfies  $T(X, Y) = \pi(Y)\phi X - \pi(X)\phi Y$ , where  $\pi$  is a 1-form and  $\phi$  is a (1,1) tensor field. In particular, if  $\phi(X) = X$ , then the quarter-symmetric connection reduces to the semi-symmetric connection.

In [9], Dobarro and Ünal studied Ricci-flat and Einstein-Lorentzian multiply warped products and considered the case of having constant scalar curvature for multiply warped products and applied their results to generalized Kasner space-times. In [10], S. Sular and C. Özgür studied warped product manifolds with a semi-symmetric metric connection, they computed curvature of semi-symmetric metric connection and consid-

ered Einstein warped product manifolds with a semi-symmetric metric connection. In [11], they studied warped product manifolds with a semi-symmetric non-metric connection. In [12], we considered multiply warped products with a semi-symmetric metric connection, then apply our results to generalized Robertson-Walker spacetimes with a semi-symmetric metric connection and generalized Kasner spacetimes with a semi-symmetric metric connection. In [13], we studied curvature of multiply warped products with a semi-symmetric non-metric connection. In this paper, we will generalize our result to warped and multiply warped products with a quarter-symmetric connection.

This paper is arranged as follows. In Section 2, we give the definition of a quarter-symmetric connection and its curvature, then give the formula of the Levi-Civita connection and curvature of singly warped and multiply twisted product. In section 3, we first compute curvature of a singly warped product with a quarter-symmetric connection, then study the generalized Robertson-Walker space-times with a quarter-symmetric connection. In section 4, firstly we compute curvature of multiply twisted products with a quarter-symmetric connection, secondly we study the special multiply warped product with a quarter-symmetric connection, finally we consider the generalized Kasner space-times with a quarter-symmetric connection.

## 2 Preliminaries

Let  $M$  be a Riemannian manifold with Riemannian metric  $g$ . A linear connection  $\bar{\nabla}$  on a Riemannian manifold  $M$  is called a quarter-symmetric connection if the torsion tensor  $T$  of the connection  $\bar{\nabla}$

$$T(X, Y) = \bar{\nabla}_X Y - \bar{\nabla}_Y X - [X, Y] \quad (1)$$

satisfies

$$T(X, Y) = \pi(Y)\phi X - \pi(X)\phi Y \quad (2)$$

where  $\pi$  is a 1-form associated with the vector field  $P$  on  $M$  defined by  $\pi(X) = g(X, P)$  and  $\phi$  is a (1,1) tensor field.  $\bar{\nabla}$  is called a quarter-symmetric metric connection if it satisfies  $\bar{\nabla}g = 0$ .  $\bar{\nabla}$  is called a quarter-symmetric non-metric connection if it satisfies  $\bar{\nabla}g \neq 0$ .

If  $\nabla$  is the Levi-Civita connection of  $M$ , in the equation (2.4) in [14], let  $\varphi_1 = \lambda_1$ ,  $\varphi_2 = 0$ ,  $U = P$ ,  $f_1 = 0$ ,  $f_2 = \lambda_2 - \lambda_1$ ,  $U_2 = P$ ,  $\lambda_1 \neq 0$ ,  $\lambda_2 \neq 0$ , then we get a

linear connection  $\bar{\nabla}$  defined by

$$\bar{\nabla}_X Y = \nabla_X Y + \lambda_1 \pi(Y)X - \lambda_2 g(X, Y)P. \quad (3)$$

It is easy to see that:

when  $\lambda_1 = \lambda_2 = 1$ ,  $\bar{\nabla}$  is a semi-symmetric metric connection;  
when  $\lambda_1 = \lambda_2 \neq 1$ ,  $\bar{\nabla}$  is a quarter-symmetric metric connection;  
when  $\lambda_1 \neq \lambda_2$ ,  $\bar{\nabla}$  is a quarter-symmetric non-metric connection.

In [12], we considered the case . In this paper, we will consider cases and .

Let  $R$  and  $\bar{R}$  be the curvature tensors of  $\nabla$  and  $\bar{\nabla}$ , respectively. By the equation (3.13) in [14], we can get

$$\begin{aligned} \bar{R}(X, Y)Z &= R(X, Y)Z + \lambda_1 g(Z, \nabla_X P)Y - \lambda_1 g(Z, \nabla_Y P)X \\ &+ \lambda_2 g(X, Z)\nabla_Y P - \lambda_2 g(Y, Z)\nabla_X P \\ &+ \lambda_1 \lambda_2 \pi(P)[g(X, Z)Y - g(Y, Z)X] \\ &+ \lambda_2^2 [g(Y, Z)\pi(X) - g(X, Z)\pi(Y)]P \\ &+ \lambda_1^2 \pi(Z)[\pi(Y)X - \pi(X)Y] \end{aligned} \quad (4)$$

for any vector fields  $X, Y, Z$  on  $M$ .

**Remark 1.** When  $\lambda_1 = \lambda_2 = 1$ , we can get the equation (4) in [12].

## 2.1 Warped Product

Let  $(B, g_B)$  and  $(F, g_F)$  be two Riemannian manifolds and  $f : B \rightarrow (0, \infty)$  be a smooth function. The warped product is the product manifold  $B \times F$  with the metric tensor  $g = g_B \oplus f^2 g_F$ . The function  $f$  is called the warping function of the warped product, and the Hessian of  $f$  is defined by  $H^f(X, Y) = XYf - (\nabla_X Y)f$ .

We need the following two lemmas from [14], for later use:

**Lemma 2.1.** Let  $M = B \times_f F$  be a warped product,  $\nabla, \nabla^B$  and  $\nabla^F$  denote the Levi-Civita connection on  $M, B$  and  $F$ , respectively. If  $X, Y \in \Gamma(TB)$  and  $U, W \in \Gamma(TF)$ , then:

- (1)  $\nabla_X Y = \nabla_X^B Y$ ;
- (2)  $\nabla_X U = \nabla_U X = \frac{Xf}{f}U$ ;
- (3)  $\nabla_U W = -\frac{g(U, W)}{f} \text{grad}_B f + \nabla_U^F W$ .

**Lemma 2.2.** *Let  $M = B \times_f F$  be a warped product with curvature  $R$ , If  $X, Y, Z \in \Gamma(TB)$  and  $U, V, W \in \Gamma(TF)$ , then:*

- (1)  $R(X, Y)Z = R^B(X, Y)Z$ ;
- (2)  $R(V, X)Y = -\frac{H_B^f(X, Y)}{f}V$ ;
- (3)  $R(X, Y)V = R(V, W)X = 0$ ;
- (4)  $R(X, V)W = -\frac{g(V, W)}{f}\nabla_X^B \text{grad}_B f$ ;
- (5)  $R(V, W)U = R^F(V, W)U - \frac{|\text{grad}_B f|_B^2}{f^2}[g(W, U)V - g(V, U)W]$ .

## 2.2 Multiply Twisted Product

A multiply twisted product  $(M, g)$  is a product manifold of form  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  with the metric  $g = g_B \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ , where for each  $i \in \{1, \dots, m\}$ ,  $b_i : B \times F_i \rightarrow (0, \infty)$  is smooth. Similarly the Hessian of  $b_i$  is defined by  $H^{b_i}(X, Y) = XYb_i - (\nabla_X Y)b_i$ .

We need the following two lemmas from [12], for later use:

**Lemma 2.3.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product and let  $X, Y \in \Gamma(TB)$  and  $U \in \Gamma(TF_i), W \in \Gamma(TF_j)$ , then:*

- (1)  $\nabla_X Y = \nabla_X^B Y$ ;
- (2)  $\nabla_X U = \nabla_U X = \frac{Xb_i}{b_i}U$ ;
- (3)  $\nabla_U W = 0$  if  $i \neq j$ ;
- (4)  $\nabla_U W = U(\ln b_i)W + W(\ln b_i)U - \frac{g_{F_i}(U, W)}{b_i} \text{grad}_{F_i} b_i - b_i g_{F_i}(U, W) \text{grad}_B b_i + \nabla_U^{F_i} W$  if  $i = j$ .

**Lemma 2.4.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product and let  $X, Y, Z \in \Gamma(TB)$  and  $V \in \Gamma(TF_i), W \in \Gamma(TF_j), U \in \Gamma(TF_k)$ , then:*

- (1)  $R(X, Y)Z = R^B(X, Y)Z$ ;
- (2)  $R(V, X)Y = -\frac{H_B^{b_i}(X, Y)}{b_i}V$ ;
- (3)  $R(X, V)W = R(V, W)X = R(V, X)W = 0$  if  $i \neq j$ ;
- (4)  $R(X, Y)V = 0$ ;
- (5)  $R(V, W)X = VX(\ln b_i)W - WX(\ln b_i)V$  if  $i = j$ ;
- (6)  $R(V, W)U = 0$  if  $i = j \neq k$  or  $i \neq j \neq k$ ;
- (7)  $R(U, V)W = -g(V, W)\frac{g_B(\text{grad}_B b_i, \text{grad}_B b_k)}{b_i b_k}U$  if  $i \neq j \neq k$ ;
- (8)  $R(X, V)W = -\frac{g(V, W)}{b_i}\nabla_X^B(\text{grad}_B b_i) + [WX(\ln b_i)]V - g_{F_i}(W, V)\text{grad}_{F_i} X(\ln b_i)$  if  $i = j$ ;
- (9)  $R(V, W)U = g(V, U)\text{grad}_B(W(\ln b_i)) - g(W, U)\text{grad}_B(V(\ln b_i)) + R^{F_i}(V, W)U - \frac{|\text{grad}_B b_i|_B^2}{b_i^2}[g(W, U)V - g(V, U)W]$  if  $i = j = k$ .

**Remark 2.** *It is easy to see that Lemmas 2.1 and 2.2 are Corollaries of Lemma 2.3 and 2.4, respectively.*

Finally we define the curvature, Ricci curvature and scalar curvature as follows:

$$R(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]},$$

$$Ric(X, Y) = \sum_k \varepsilon_k \langle R(X, E_k)Y, E_k \rangle,$$

$$S = \sum_k \varepsilon_k Ric(E_k, E_k)$$

where  $E_k$  is an orthonormal base of  $M$  with  $\langle E_k, E_k \rangle = \varepsilon_k$ ,  $\varepsilon_k = \pm 1$ .

### 3 Warped Product with a Quarter-symmetric Connection

In this section, we firstly compute curvature, Ricci curvature and scalar curvature of singly warped product with a quarter-symmetric connection, then study the generalized Robertson-Walker space-times with a quarter-symmetric connection.

#### 3.1 Connection and Curvature

By Lemma 2.1 and the equation (3), we have the following two Propositions:

**Proposition 3.1.** *Let  $M = B \times_f F$  be a warped product. If  $X, Y \in \Gamma(TB)$ ,  $U, W \in \Gamma(TF)$  and  $P \in \Gamma(TB)$ , then:*

- (1)  $\bar{\nabla}_X Y = \bar{\nabla}_X^B Y$ ;
- (2)  $\bar{\nabla}_X U = \frac{Xf}{f} U$ ;
- (3)  $\bar{\nabla}_U X = \left[ \frac{Xf}{f} + \lambda_1 \pi(X) \right] U$ ;
- (4)  $\bar{\nabla}_U W = -f g_F(U, W) \text{grad}_B f + \nabla_U^F W - \lambda_2 g(U, W) P$ .

**Proposition 3.2.** *Let  $M = B \times_f F$  be a warped product. If  $X, Y \in \Gamma(TB)$ ,  $U, W \in \Gamma(TF)$  and  $P \in \Gamma(TF)$ , then:*

- (1)  $\bar{\nabla}_X Y = \nabla_X^B Y - \lambda_2 g(X, Y) P$ ;
- (2)  $\bar{\nabla}_X U = \frac{Xf}{f} U + \lambda_1 \pi(U) X$ ;
- (3)  $\bar{\nabla}_U X = \frac{Xf}{f} U$ ;
- (4)  $\bar{\nabla}_U W = -\frac{g(U, W)}{f} \text{grad}_B f + \bar{\nabla}_U^F W$ .

By Lemmas 2.12.2 and the equation (4), we have the following two Propositions:

**Proposition 3.3.** *Let  $M = B \times_f F$  be a warped product. If  $X, Y, Z \in \Gamma(TB)$ ,  $U, V, W \in \Gamma(TF)$  and  $P \in \Gamma(TB)$ , then:*

- (1)  $\overline{R}(X, Y)Z = \overline{R}^B(X, Y)Z$ ;
- (2)  $\overline{R}(V, X)Y = -\left[\frac{H_B^f(X, Y)}{f} + \lambda_2 \frac{Pf}{f}g(X, Y) + \lambda_1 \lambda_2 \pi(P)g(X, Y) + \lambda_1 g(Y, \nabla_X P) - \lambda_1^2 \pi(X)\pi(Y)\right]V$ ;
- (3)  $\overline{R}(X, Y)V = 0$ ;
- (4)  $\overline{R}(V, W)X = 0$ ;
- (5)  $\overline{R}(X, V)W = -g(V, W)\left[\frac{\nabla_X^B(\text{grad}_B f)}{f} + \lambda_1 \frac{Pf}{f}X + \lambda_2 \nabla_X P + \lambda_1 \lambda_2 \pi(P)X - \lambda_2^2 \pi(X)P\right]$ ;
- (6)  $\overline{R}(U, V)W = R^F(U, V)W - \left[\frac{|\text{grad}_B f|_B^2}{f^2} + (\lambda_1 + \lambda_2) \frac{Pf}{f} + \lambda_1 \lambda_2 \pi(P)\right]\left[g(V, W)U - g(U, W)V\right]$ .

**Proposition 3.4.** *Let  $M = B \times_f F$  be a warped product. If  $X, Y, Z \in \Gamma(TB)$ ,  $U, V, W \in \Gamma(TF)$  and  $P \in \Gamma(TF)$ , then:*

- (1)  $\overline{R}(X, Y)Z = R^B(X, Y)Z + \lambda_2 \left[g(X, Z) \frac{Yf}{f} - g(Y, Z) \frac{Xf}{f}\right]P + \lambda_1 \lambda_2 \pi(P)[g(X, Z)Y - g(Y, Z)X]$ ;
- (2)  $\overline{R}(V, X)Y = -\frac{H_B^f(X, Y)}{f}V - \lambda_1 \pi(V) \frac{Yf}{f}X - \lambda_2 g(X, Y) \nabla_V P - g(X, Y)[\lambda_1 \lambda_2 \pi(P)V - \lambda_2^2 \pi(V)P]$ ;
- (3)  $\overline{R}(X, Y)V = \lambda_1 \pi(V) \left[\frac{Xf}{f}Y - \frac{Yf}{f}X\right]$ ;
- (4)  $\overline{R}(V, W)X = \lambda_1 \frac{Xf}{f}[\pi(W)V - \pi(V)W]$ ;
- (5)  $\overline{R}(X, V)W = -g(V, W) \frac{\nabla_X^B(\text{grad}_B f)}{f} + \lambda_1 \frac{Xf}{f} \pi(W)V - \lambda_1 g(W, \nabla_V P)X - \lambda_2 g(V, W) \frac{Xf}{f}P - \lambda_1 \lambda_2 g(V, W) \pi(P)X + \lambda_1^2 \pi(W) \pi(V)X$ ;
- (6)  $\overline{R}(U, V)W = R^F(U, V)W - \frac{|\text{grad}_B f|_B^2}{f^2}[g(V, W)U - g(U, W)V] + \lambda_1 [g(W, \nabla_U P)V - g(W, \nabla_V P)U] + \lambda_2 [g(U, W) \nabla_V P - g(V, W) \nabla_U P] + \lambda_1 \lambda_2 \pi(P)[g(U, W)V - g(V, W)U] + \lambda_2^2 [g(V, W) \pi(U) - g(U, W) \pi(V)]P + \lambda_1^2 \pi(W)[\pi(V)U - \pi(U)V]$ .

By Propositions 3.3 and 3.4 and the definition of the Ricci curvature tensor, we have the following two Propositions:

**Proposition 3.5.** *Let  $M = B \times_f F$  be a warped product,  $\dim B = n_1$ ,  $\dim F = n_2$  and  $\dim M = \overline{n} = n_1 + n_2$ . If  $X, Y \in \Gamma(TB)$ ,  $V, W \in \Gamma(TF)$  and  $P \in \Gamma(TB)$ , then:*

- (1)  $\overline{Ric}(X, Y) = \overline{Ric}^B(X, Y) + n_2 \left[\frac{H_B^f(X, Y)}{f} + \lambda_2 \frac{Pf}{f}g(X, Y) + \lambda_1 \lambda_2 \pi(P)g(X, Y) + \lambda_1 g(Y, \nabla_X P) - \lambda_1^2 \pi(X)\pi(Y)\right]$ ;
- (2)  $\overline{Ric}(X, V) = \overline{Ric}(V, X) = 0$ ;

$$(3) \overline{Ric}(V, W) = \overline{Ric}^F(V, W) + \left\{ \frac{\Delta_B f}{f} + (n_2 - 1) \frac{|grad_B f|_B^2}{f^2} + [(\overline{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] \pi(P) \right. \\ \left. + \lambda_2 div_B P + [(\overline{n} - 1)\lambda_1 + (n_2 - 1)\lambda_2] \frac{Pf}{f} \right\} g(V, W).$$

where  $div_B P = \sum_{k=1}^{n_1} \varepsilon_k \langle \nabla_{E_k} P, E_k \rangle$ , and  $E_k, 1 \leq k \leq n_1$  is an orthonormal base of  $B$  with  $\varepsilon_k = g(E_k, E_k)$ .

**Proposition 3.6.** *Let  $M = B \times_f F$  be a warped product,  $\dim B = n_1, \dim F = n_2$  and  $\dim M = \overline{n} = n_1 + n_2$ . If  $X, Y \in \Gamma(TB)$ ,  $V, W \in \Gamma(TF)$  and  $P \in \Gamma(TF)$ , then:*

$$(1) \overline{Ric}(X, Y) = \overline{Ric}^B(X, Y) + n_2 \frac{H_B^f(X, Y)}{f} + [(\overline{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] \pi(P) g(X, Y) \\ + \lambda_2 g(X, Y) div_F P; \\ (2) \overline{Ric}(X, V) = [(\overline{n} - 1)\lambda_1 - \lambda_2] \pi(V) \frac{Xf}{f}; \\ (3) \overline{Ric}(V, X) = [\lambda_2 - (\overline{n} - 1)\lambda_1] \pi(V) \frac{Xf}{f}; \\ (4) \overline{Ric}(V, W) = Ric^F(V, W) + g(V, W) \left\{ \frac{\Delta_B f}{f} + (n_2 - 1) \frac{|grad_B f|_B^2}{f^2} + [(\overline{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] \pi(P) \right\} \\ + [(\overline{n} - 1)\lambda_1 - \lambda_2] g(W, \nabla_V P) + [\lambda_2^2 + (1 - \overline{n})\lambda_1^2] \pi(V) \pi(W) + \lambda_2 g(V, W) div_F P.$$

By Proposition 3.5 and the definition of the scalar curvature, we have the following:

**Proposition 3.7.** *Let  $M = B \times_f F$  be a warped product,  $\dim B = n_1, \dim F = n_2$  and  $\dim M = \overline{n} = n_1 + n_2$ . If  $P \in \Gamma(TB)$ , then the scalar curvature  $\overline{S}$  has the following expression:*

$$\overline{S} = \overline{S}^B + 2n_2 \frac{\Delta_B f}{f} + \frac{S^F}{f^2} + n_2(n_2 - 1) \frac{|grad_B f|_B^2}{f^2} + n_2(\overline{n} - 1)(\lambda_1 + \lambda_2) \frac{Pf}{f} \\ + [n_2(\overline{n} + n_1 - 1)\lambda_1 \lambda_2 - n_2(\lambda_1^2 + \lambda_2^2)] \pi(P) + n_2(\lambda_1 + \lambda_2) div_B P.$$

By Proposition 3.6 and the definition of the scalar curvature, we have the following:

**Proposition 3.8.** *Let  $M = B \times_f F$  be a warped product,  $\dim B = n_1, \dim F = n_2$  and  $\dim M = \overline{n} = n_1 + n_2$ . If  $P \in \Gamma(TF)$ , then the scalar curvature  $\overline{S}$  has the following expression:*

$$\overline{S} = \overline{S}^B + 2n_2 \frac{\Delta_B f}{f} + \frac{S^F}{f^2} + n_2(n_2 - 1) \frac{|grad_B f|_B^2}{f^2} + [\overline{n}(\overline{n} - 1)\lambda_1 \lambda_2 + (1 - \overline{n})(\lambda_1^2 + \lambda_2^2)] \pi(P) \\ + (\overline{n} - 1)(\lambda_1 + \lambda_2) div_F P.$$

### 3.2 Generalized Robertson-Walker Space-times with a Quarter-symmetric Connection

**Theorem 3.9.** *Let  $M = I \times_f F$  be a warped product, where  $I$  is an open interval in  $\mathbb{R}$ ,  $\dim I = 1$  and  $\dim F = \overline{n} - 1$  ( $\overline{n} \geq 3$ ). Then  $(M, \overline{\nabla})$  is Einstein if and only if  $(F, \nabla^F)$  is Einstein for  $P = \frac{\partial}{\partial t}$  or  $f$  is a constant on  $I$  for  $P \in \Gamma(TF)$ ,  $\lambda_2 \neq (\overline{n} - 1)\lambda_1$ .*



*Proof.* Assume that  $P = \frac{\partial}{\partial t}$ , Let  $f = e^{\frac{q}{2}}$ , by Proposition 3.5, we can write

$$\overline{Ric}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = (1 - \bar{n})\left[\frac{1}{4}(q')^2 + \frac{1}{2}q'' - \frac{1}{2}\lambda_2 q' + \lambda_1 \lambda_2 - \lambda_1^2\right]g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right); \quad (5)$$

$$\overline{Ric}\left(\frac{\partial}{\partial t}, V\right) = 0;$$

$$\begin{aligned} \overline{Ric}(V, W) = Ric^F(V, W) + e^q & \left\{ \frac{\bar{n}-1}{4}(q')^2 + \frac{1}{2}\left[(\bar{n}-1)\lambda_1 + (\bar{n}-2)\lambda_2\right]q' \right. \\ & \left. + \lambda_2^2 + (1-\bar{n})\lambda_1\lambda_2 + \frac{1}{2}q'' \right\} g_F(V, W) \end{aligned} \quad (6)$$

for any  $V, W \in \Gamma(TF)$ .

Since  $M$  is Einstein manifold, we have

$$\overline{Ric}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \alpha g_I\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right); \quad (7)$$

$$\overline{Ric}(V, W) = \alpha e^q g_F(V, W). \quad (8)$$

From equations (5) and (7), we get

$$\alpha = (1 - \bar{n})\left[\frac{1}{4}(q')^2 + \frac{1}{2}q'' - \frac{1}{2}\lambda_2 q' + \lambda_1 \lambda_2 - \lambda_1^2\right]. \quad (9)$$

Similarly, from equations (6) and (8), and by the use of (9), we obtain

$$Ric^F(V, W) = \left[\frac{1-\bar{n}}{2}(q')^2 - \frac{\bar{n}}{2}q'' + \left(\frac{1-\bar{n}}{2}\lambda_1 + \frac{1}{2}\lambda_2\right)q' + (\bar{n}-1)\lambda_1^2 - \lambda_2^2\right]e^q g_F(V, W)$$

which implies that  $(F, \nabla^F)$  is Einstein manifold.

Assume that  $P \in \Gamma(TF)$ , by Proposition 3.6, we have

$$\overline{Ric}\left(\frac{\partial}{\partial t}, V\right) = \frac{1}{2}q'[(\bar{n}-1)\lambda_1 - \lambda_2]\pi(V); \quad (10)$$

$$\overline{Ric}\left(V, \frac{\partial}{\partial t}\right) = \frac{1}{2}q'[\lambda_2 - (\bar{n}-1)\lambda_1]\pi(V) \quad (11)$$

for any  $V \in \Gamma(TF)$ .

Since  $M$  is an Einstein manifold, we can write

$$\overline{Ric}\left(\frac{\partial}{\partial t}, V\right) = \alpha g\left(\frac{\partial}{\partial t}, V\right) = 0 = \alpha g\left(V, \frac{\partial}{\partial t}\right) = \overline{Ric}\left(V, \frac{\partial}{\partial t}\right) \quad (12)$$

where  $\frac{\partial}{\partial t} \in \Gamma(TB)$  and  $V \in \Gamma(TF)$ .

Since  $\lambda_2 \neq (\bar{n}-1)\lambda_1, \pi(V) \neq 0$ , , using equations (10), (11), (12), then we can get  $q' = 0$ , which means  $q$  is a constant on  $I$ , then  $f$  is a constant on  $I$ .  $\square$

**Theorem 3.10.** *Let  $M = B \times_f I$  be a warped product, where  $I$  is an open interval in  $\mathbb{R}$ ,  $\dim I = 1$  and  $\dim B = \bar{n} - 1$  ( $\bar{n} \geq 3$ ). Then*

(1) *If  $(M, \bar{\nabla})$  is an Einstein manifold,  $P \in \Gamma(TB)$ ,  $\nabla^B P = 0$ , and  $f$  is a constant on  $B$ , then:*

$$\bar{S}^B = [(\bar{n} - 1)(\bar{n} - 2)\lambda_1\lambda_2 + \lambda_1^2 + (1 - \bar{n})\lambda_2^2]\pi(P).$$

*Furthermore, if  $(\bar{n} - 1)(\bar{n} - 2)\lambda_1\lambda_2 + \lambda_1^2 + (1 - \bar{n})\lambda_2^2 = 0$ , then  $\bar{S}^B = 0$ .*

(2) *If  $(M, \bar{\nabla})$  is Einstein manifold,  $P \in \Gamma(TI)$ , and  $\lambda_2 \neq (\bar{n} - 1)\lambda_1$  then  $f$  is a constant on  $B$ .*

(3) *If  $f$  is a constant on  $B$ ,  $(B, \nabla^B)$  is Einstein,  $P \in \Gamma(TI)$ , then  $(M, \bar{\nabla})$  is Einstein manifold; furthermore, if  $\lambda_2 = (\bar{n} - 1)\lambda_1$ , then  $\alpha = \alpha_B$ , where  $\alpha$  and  $\alpha_B$  denote the Einstein constant on  $M$  and  $B$ , respectively.*

*Proof.* Assume that  $(M, \bar{\nabla})$  is an Einstein manifold,  $P \in \Gamma(TB)$ , then

$$\overline{Ric}(X, Y) = \frac{\bar{s}}{\bar{n}}g(X, Y) \quad (13)$$

for any  $X, Y \in \Gamma(TB)$ . Consider that  $f$  is a constant on  $B$  and by Proposition 3.7, we can get

$$\overline{Ric}(X, Y) = \frac{1}{\bar{n}}g(X, Y)\{\bar{S}^B + [(2\bar{n} - 2)\lambda_1\lambda_2 - (\lambda_1^2 + \lambda_2^2)]\pi(P)\}.$$

Define  $\hat{S} = \sum_{k=1}^{\bar{n}-1} \frac{1}{\bar{n}} \overline{Ric}(E_k, E_k)$ , where  $E_k$  is an orthonormal base of  $B$ , then we have

$$\hat{S} = \frac{\bar{n} - 1}{\bar{n}} \left\{ \bar{S}^B + [(2\bar{n} - 2)\lambda_1\lambda_2 - (\lambda_1^2 + \lambda_2^2)]\pi(P) \right\}. \quad (14)$$

On the other hand, using Proposition 3.5, we can get

$$\overline{Ric}(X, Y) = \overline{Ric}^B(X, Y) + [\lambda_1\lambda_2\pi(P)g(X, Y) - \lambda_1^2\pi(X)\pi(Y)],$$

then we have

$$\hat{S} = \bar{S}^B + [(\bar{n} - 1)\lambda_1\lambda_2 - \lambda_1^2]\pi(P). \quad (15)$$

From equations (14) and (15), we can get

$$\bar{S}^B = [(\bar{n} - 1)(\bar{n} - 2)\lambda_1\lambda_2 + \lambda_1^2 + (1 - \bar{n})\lambda_2^2]\pi(P).$$

Assume that  $(M, \bar{\nabla})$  is Einstein manifold,  $P \in \Gamma(TI)$ , by Proposition 3.6, we obtain

$$\overline{Ric}(X, P) = [(\bar{n} - 1)\lambda_1 - \lambda_2]\pi(P)\frac{Xf}{f}; \quad (16)$$

$$\overline{Ric}(P, X) = [\lambda_2 - (\bar{n} - 1)\lambda_1]\pi(P)\frac{Xf}{f}. \quad (17)$$

Using the similar proof of Theorem 3.9., we can get  $Xf = 0$ , which means  $f$  is a constant on  $B$ .

If  $f$  is a constant on  $B$ , and  $(B, \nabla^B)$  is Einstein, and  $P \in \Gamma(TI)$ , then

$$Ric^B(X, Y) = \alpha_B g(X, Y). \quad (18)$$

By Theorem 3.6, we have

$$\overline{Ric}(X, Y) = \overline{Ric}^B(X, Y) + [(\bar{n} - 1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)g(X, Y). \quad (19)$$

So by equations (18) and (19), we can easily get

$$\overline{Ric}(X, Y) = \{\alpha_B + [(\bar{n} - 1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)\}g(X, Y)$$

which means  $(M, \overline{\nabla})$  is an Einstein manifold.

Furthermore, if  $\lambda_2 = (\bar{n} - 1)\lambda_1$ , then  $\overline{Ric}(X, Y) = \alpha_B g(X, Y)$ , and  $\alpha = \alpha_B$ .  $\square$

Now we specially study  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $I$  is an open interval in  $\mathbb{R}$ . Let begin with the following theorem:

**Theorem 3.11.** *Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = l$ . Then  $(M, \overline{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if the following conditions are satisfied*

- (1)  $(F, \nabla^F)$  is Einstein with the Einstein constant  $\alpha_F$ ;
- (2)  $l\left(\lambda_2 \frac{f'}{f} - \frac{f''}{f} + \lambda_1^2 - \lambda_1\lambda_2\right) = \alpha$ ;
- (3)  $\alpha_F - f f'' + (1 - l)(f')^2 + [\lambda_2^2 - l\lambda_1\lambda_2 - \alpha]f^2 + [l\lambda_1 + (l - 1)\lambda_2]f f' = 0$ .

*Proof.* By Proposition 3.5, we have

$$\begin{aligned} \overline{Ric}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) &= -l\left(\lambda_2 \frac{f'}{f} - \frac{f''}{f} + \lambda_1^2 - \lambda_1\lambda_2\right); \\ \overline{Ric}\left(\frac{\partial}{\partial t}, V\right) &= \overline{Ric}\left(V, \frac{\partial}{\partial t}\right) = 0; \\ \overline{Ric}(V, W) &= Ric^F(V, W) + g_F(V, W)\{-f f'' - (l - 1)f'^2 + (\lambda_2^2 - l\lambda_1\lambda_2)f^2 \\ &\quad + [l\lambda_1 + (l - 1)\lambda_2]f f'\}. \end{aligned}$$

Then by the Einstein condition, we get Theorem 3.11.  $\square$

**Remark 3.** When  $\lambda_1 = \lambda_2 = 1$ , we can get Corollary 21 in [12].

Considering the dimension of  $F$ , we get Corollaries 3.12 and 3.13 of Theorem 3.11:

**Corollary 3.12.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = 1$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if  
 $f'' = \lambda_2 f' + (\lambda_1^2 - \lambda_1 \lambda_2) f - \alpha f$

**Remark 4.** (1) From Theorem 3.11, we can also get: if  $\dim F = 1$ , then  $\alpha_F = 0$ ;

(2) When  $\lambda_1 = \lambda_2 = 1$ , we can get Corollary 23 in [12].

**Corollary 3.13.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = l > 1$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if the following conditions are satisfied:

- (1)  $(F, \nabla^F)$  is Einstein with the Einstein constant  $\alpha^F$ ;
- (2)  $f'' = \lambda_2 f' + (\lambda_1^2 - \lambda_1 \lambda_2 - \frac{\alpha}{l}) f$ ;
- (3)  $\frac{\alpha_F}{1-l} + (f')^2 + \left[ \frac{\alpha}{l} + \lambda_1 \lambda_2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} \right] f^2 + \left[ \frac{l}{1-l} \lambda_1 + \frac{(l-2)}{1-l} \lambda_2 \right] f f' = 0$ .

**Remark 5.** When  $\lambda_1 = \lambda_2 = 1$ , we can get Corollary 24 in [12].

By Corollary 3.12 and elementary methods for ordinary differential equations, we get:

**Theorem 3.14.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = 1$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if  
(1)  $\alpha < (\lambda_1 - \frac{1}{2} \lambda_2)^2$ ,  $f(t) = c_1 e^{((\lambda_2 + \sqrt{(2\lambda_1 - \lambda_2)^2 - 4\alpha})/2)t} + c_2 e^{((\lambda_2 - \sqrt{(2\lambda_1 - \lambda_2)^2 - 4\alpha})/2)t}$ ,  
(2)  $\alpha = (\lambda_1 - \frac{1}{2} \lambda_2)^2$ ,  $f(t) = c_1 e^{(\lambda_2/2)t} + c_2 t e^{(\lambda_2/2)t}$ ,  
(3)  $\alpha > (\lambda_1 - \frac{1}{2} \lambda_2)^2$ ,  $f(t) = c_1 e^{(\lambda_2/2)t} \cos((\sqrt{4\alpha - (2\lambda_1 - \lambda_2)^2}/2)t) + c_2 e^{(\lambda_2/2)t} \sin((\sqrt{4\alpha - (2\lambda_1 - \lambda_2)^2}/2)t)$ .

**Remark 6.** When  $\lambda_1 = \lambda_2 = 1$ , we can get Corollary 25 in [12].

As a Corollary of Theorem 3.14, we have

**Corollary 3.15.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = 1$ , and  $\lambda_2 = 2\lambda_1$ , then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if  
(1)  $\alpha < 0$ ,  $f(t) = c_1 e^{(\lambda_1 + \sqrt{-\alpha})t} + c_2 e^{(\lambda_1 - \sqrt{-\alpha})t}$ ,  
(2)  $\alpha = 0$ ,  $f(t) = c_1 e^{\lambda_1 t} + c_2 t e^{\lambda_1 t}$ ,  
(3)  $\alpha > 0$ ,  $f(t) = c_1 e^{\lambda_1 t} \cos(\sqrt{\alpha} t) + c_2 e^{\lambda_1 t} \sin(\sqrt{\alpha} t)$ .

**Theorem 3.16.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ ,  $\dim F = l > 1$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if one of the following conditions is satisfied:

- (1)  $\alpha = (\lambda_1^2 - \lambda_1 \lambda_2)l$ ,  $\alpha_F = c_2^2(l\lambda_1^2 - \lambda_2^2)$ ,  $f(t) = c_2$ ;
- (2)  $\lambda_1 = \lambda_2$ ,  $\alpha = 0$ ,  $\alpha_F = (l-1)c_2^2\lambda_1^2$ ,  $f(t) = c_1 e^{\lambda_1 t} + c_2$ ;
- (3)  $\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 \neq 0$ ,  $\lambda_2 \neq l\lambda_1$ ,  $\alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2 - (l^2+l)\lambda_1\lambda_1^3 - 2l^2\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}$ ,  $\alpha_F = 0$ ,  
 $f(t) = c_0 e^{(l\lambda_1^2 - \lambda_2^2)t/(l\lambda_1 - \lambda_2)}$ ;
- (4)  $\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 = 0$ ,  $\alpha = l(\lambda_1 - \frac{1}{2}\lambda_2)^2$ ,  $\alpha_F = 0$ ,  $f(t) = c_1 e^{\frac{\lambda_2}{2}t}$ .

*Proof.* Let  $\frac{\alpha}{l} = d_0$ ,  $\frac{\alpha_F}{1-l} = \bar{d}_0$ ,  $a_0 = \frac{\lambda_2 + \sqrt{(2\lambda_1 - \lambda_2)^2 - 4d_0}}{2}$ ,  $b_0 = \frac{\lambda_2 - \sqrt{(2\lambda_1 - \lambda_2)^2 - 4d_0}}{2}$ , then  $a_0 + b_0 = \lambda_2$ ,  $a_0 b_0 = d_0 + \lambda_1 \lambda_2 - \lambda_1^2$ .

$d_0 < (\lambda_1 - \frac{1}{2}\lambda_2)^2$ , then  $f(t) = c_1 e^{a_0 t} + c_2 e^{b_0 t}$ . By Corollary 3.13(3), then

$$\begin{aligned} & \bar{d}_0 + c_1^2 \left( a_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l} \lambda_1 a_0 + \frac{l-2}{1-l} \lambda_2 a_0 \right) e^{2a_0 t} \\ & + c_2^2 \left( b_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l} \lambda_1 b_0 + \frac{l-2}{1-l} \lambda_2 b_0 \right) e^{2b_0 t} \\ & + c_1 c_2 \left[ 4a_0 b_0 + \frac{2\lambda_2^2 - 2l\lambda_1^2}{1-l} + \left( \frac{l}{1-l} \lambda_1 + \frac{l-2}{1-l} \lambda_2 \right) (a_0 + b_0) \right] e^{(a_0 + b_0)t} = 0. \end{aligned} \quad (20)$$

1)  $b_0 = 0$ , we have  $d_0 = \lambda_1^2 - \lambda_1 \lambda_2 < (\lambda_1 - \frac{1}{2}\lambda_2)^2$ ,  $a_0 = \lambda_2$ ,  $\alpha = (\lambda_1^2 - \lambda_1 \lambda_2)l$ . By the equation (20), we get

$$\bar{d}_0 + c_1^2 \frac{l\lambda_1(\lambda_2 - \lambda_1)}{1-l} e^{2\lambda_2 t} + c_2^2 \frac{\lambda_2^2 - l\lambda_1^2}{1-l} + c_1 c_2 \frac{l(\lambda_2 + 2\lambda_1)(\lambda_2 - \lambda_1)}{1-l} e^{\lambda_2 t} = 0.$$

Since  $e^{2\lambda_2 t}$  and  $e^{\lambda_2 t}$  are linearly independent, we have

$$\begin{cases} c_1^2 \frac{l\lambda_1(\lambda_2 - \lambda_1)}{1-l} = 0 \\ \bar{d}_0 + c_2^2 \frac{\lambda_2^2 - l\lambda_1^2}{1-l} = 0 \\ c_1 c_2 \frac{l(\lambda_2 + 2\lambda_1)(\lambda_2 - \lambda_1)}{1-l} = 0 \end{cases}$$

1'.  $c_1 = 0$ ,  $c_2 \neq 0$ . We have  $\alpha = (\lambda_1^2 - \lambda_1 \lambda_2)l$ ,  $\alpha_F = c_2^2(l\lambda_1^2 - \lambda_2^2)$ ,  $f(t) = c_2$ .

2'.  $c_1 \neq 0$ ,  $c_2 = 0$ . We have  $\lambda_1 = \lambda_2$ ,  $\alpha = \alpha_F = 0$ ,  $f(t) = c_1 e^{\lambda_1 t}$ .

3'.  $c_1 \neq 0$ ,  $c_2 \neq 0$ . We have  $\lambda_1 = \lambda_2$ ,  $\alpha = 0$ ,  $\alpha_F = (l-1)c_2^2\lambda_1^2$ ,  $f(t) = c_1 e^{\lambda_1 t} + c_2$ .

It is easy to see that the conclusion of 2' is a special case of 3'.

2)  $b_0 \neq 0$ , 1'.  $c_1 = 0$ ,  $c_2 \neq 0$ . Since  $e^{2a_0 t}$ ,  $e^{2b_0 t}$  and  $e^{(a_0 + b_0)t}$  are linearly independent, we have  $\bar{d}_0 = 0$ ,  $b_0^2 + a_0 b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l} \lambda_1 b_0 + \frac{l-2}{1-l} \lambda_2 b_0 = 0$ , then  $\alpha_F = 0$ ,  $\frac{l\lambda_1 - \lambda_2}{1-l} b_0 =$

$\frac{l\lambda_1^2 - \lambda_2^2}{1-l}$ . If  $\lambda_2 = l\lambda_1$ , we have  $0 = l\lambda_1^2$ , this is a contradiction with  $\lambda_1 \neq 0$ . So  $\lambda_2 \neq l\lambda_1$  and  $b_0 = \frac{l\lambda_1^2 - \lambda_2^2}{l\lambda_1 - \lambda_2}$ . Considering that  $b_0 < \frac{\lambda_2}{2}$ , we get  $\frac{\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2}{l\lambda_1 - \lambda_2} > 0$ .  $a_0 = \frac{l\lambda_1(\lambda_2 - \lambda_1)}{l\lambda_1 - \lambda_2}$ ,  $d_0 = \frac{(3l+1)\lambda_1^2\lambda_2^2 - (l+1)\lambda_1\lambda_2^3 - 2l\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}$ ,  $\alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2 - (l^2+l)\lambda_1\lambda_2^3 - 2l^2\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}$  and  $d_0$  satisfies  $d_0 < (\lambda_1 - \frac{1}{2}\lambda_2)^2$ . So in this case, we obtain

$$\frac{\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2}{l\lambda_1 - \lambda_2} > 0, \alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2 - (l^2+l)\lambda_1\lambda_2^3 - 2l^2\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}, \alpha_F = 0, f(t) = c_2 e^{\frac{l\lambda_1^2 - \lambda_2^2}{l\lambda_1 - \lambda_2}t}.$$

2'.  $c_1 \neq 0, c_2 = 0$ . Using the same method we can get  $a_0 = \frac{l\lambda_1^2 - \lambda_2^2}{l\lambda_1 - \lambda_2}$ ,  $b_0 = \frac{l\lambda_1(\lambda_2 - \lambda_1)}{l\lambda_1 - \lambda_2}$  and  $\frac{\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2}{l\lambda_1 - \lambda_2} < 0$ ,  $\alpha = \frac{(3l^2+l)\lambda_1^2\lambda_2^2 - (l^2+l)\lambda_1\lambda_2^3 - 2l^2\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}$ ,  $\alpha_F = 0$ ,  $f(t) = c_1 e^{\frac{l\lambda_1^2 - \lambda_2^2}{l\lambda_1 - \lambda_2}t}$ .

So by 1' and 2', we get Theorem 3.16(3).

3'.  $c_1 \neq 0, c_2 \neq 0$ . Since  $e^{2a_0t}, e^{2b_0t}$  and  $e^{(a_0+b_0)t}$  are linearly independent, we have

$$a_0^2 + a_0b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1a_0 + \frac{l-2}{1-l}\lambda_2a_0 = 0; \quad (21a)$$

$$b_0^2 + a_0b_0 + \lambda_1^2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} + \frac{l}{1-l}\lambda_1b_0 + \frac{l-2}{1-l}\lambda_2b_0 = 0. \quad (21b)$$

(21a) - (21b) we get  $(a_0 - b_0)\lambda_2 + \frac{l\lambda_1}{1-l}(a_0 - b_0) + \frac{l-2}{1-l}\lambda_2(a_0 - b_0) = 0$ , since  $a_0 \neq b_0$  we have  $\lambda_2 + \frac{l\lambda_1}{1-l} + \frac{l-2}{1-l}\lambda_2 = 0$ , then  $\lambda_2 = l\lambda_1$ , using (21a) again we get  $0 = l\lambda_1^2$ , this is a contradiction with  $\lambda_1 \neq 0$ . So in case 3' we have no solution.

$d_0 = (\lambda_1 - \frac{1}{2}\lambda_2)^2$ , then  $f(t) = c_1 e^{\frac{\lambda_2}{2}t} + c_2 t e^{\frac{\lambda_2}{2}t}$ . By Corollary 3.13(3), we get

$$\begin{aligned} \overline{d_0} + \left(\frac{\lambda_2}{2}c_1 + c_2 + \frac{\lambda_2}{2}c_2t\right)^2 e^{\lambda_2t} + \frac{(5-l)\lambda_2^2 - 4l\lambda_1^2}{4(1-l)}(c_1 + c_2t)^2 e^{\lambda_2t} \\ + \left(\frac{l}{1-l}\lambda_1 + \frac{l-2}{1-l}\lambda_2\right)(c_1 + c_2t)\left(\frac{\lambda_2}{2}c_1 + c_2 + \frac{\lambda_2}{2}c_2t\right)e^{\lambda_2t} = 0 \end{aligned}$$

1'.  $c_1 = 0, c_2 \neq 0$ . The coefficient of  $e^{\lambda_2t}$  is  $c_2^2 = 0$ , this is a contradiction with  $c_2 \neq 0$ .

2'.  $c_1 \neq 0, c_2 = 0$ . Then  $\overline{d_0} = 0$  and from the coefficient of  $e^{\lambda_2t}$  we can get  $\frac{\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2}{l\lambda_1 - \lambda_2} = 0$ , so  $\alpha = d_0l = l(\lambda_1 - \frac{1}{2}\lambda_2)^2$ ,  $\alpha_F = 0$ ,  $f(t) = c_1 e^{\frac{\lambda_2}{2}t}$ .

3'.  $c_1 \neq 0, c_2 \neq 0$ . From the coefficient of  $t^2 e^{\lambda_2t}$  we get

$$\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_1^2 = 0, \quad (22)$$

by the equation (22), the coefficient of  $t e^{\lambda_2t}$  becomes

$$\frac{(l-3)\lambda_2 + 2l\lambda_1}{2(1-l)}c_2 - \frac{\lambda_2^2}{4}c_1 = 0, \quad (23)$$

the coefficient of  $e^{\lambda_2t}$  is  $\frac{l\lambda_1 - \lambda_2}{1-l}c_1c_2 + c_2^2 = 0$ , then  $c_2 = \frac{\lambda_2 - l\lambda_1}{1-l}c_1$ , so by equation (23) we get  $(2l-7-l^2)\lambda_2 = (4l^2-10l)\lambda_1$ , considering that  $2l-7-l^2 \neq 0, 4l^2-10l \neq 0$ , we obtain

$\lambda_2 = \frac{(4l^2-10l)}{(2l-7-l^2)}\lambda_1$ , using the equation (22) again, we have  $(l-1)^2(3l^2-15l+49) = 0$ , but  $l-1 \neq 0, 3l^2-15l+49 \neq 0$ , so we have no solution in this case.

$d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$ , let  $h_0 = \frac{1}{2}\sqrt{4d_0 - (2\lambda_1 - \lambda_2)^2}$ , then  $f(t) = e^{\frac{\lambda_2}{2}t}(c_1\cos(h_0t) + c_2\sin(h_0t))$ . By Corollary 3.13(3), we get

$$\begin{aligned} & \overline{d_0} + \left[ \left( \frac{\lambda_2}{2}c_1 + c_2h_0 \right) \cos(h_0t) + \left( \frac{\lambda_2}{2}c_2 - c_1h_0 \right) \sin(h_0t) \right]^2 e^{\lambda_2t} \\ & + \left( d_0 + \lambda_1\lambda_2 + \frac{\lambda_2^2 - \lambda_1^2}{1-l} \right) \left( c_1\cos(h_0t) + c_2\sin(h_0t) \right)^2 e^{\lambda_2t} + \left( \frac{l}{1-l}\lambda_1 + \frac{l-2}{1-l}\lambda_2 \right) \\ & \left( c_1\cos(h_0t) + c_2\sin(h_0t) \right) \left[ \left( \frac{\lambda_2}{2}c_1 + c_2h_0 \right) \cos(h_0t) + \left( \frac{\lambda_2}{2}c_2 - c_1h_0 \right) \sin(h_0t) \right] e^{\lambda_2t} = 0. \end{aligned}$$

The coefficient of  $\cos^2(h_0t)e^{\lambda_2t}$  is

$$\left[ d_0 + \frac{(l+1)\lambda_2^2 + (4-2l)\lambda_1\lambda_2 - 4\lambda_1^2}{4(1-l)} \right] c_1^2 + c_2^2 h_0^2 - \frac{l\lambda_1 - \lambda_2}{1-l} c_1 c_2 h_0 = 0. \quad (24)$$

Similarly the coefficient of  $\sin^2(h_0t)e^{\lambda_2t}$  is

$$\left[ d_0 + \frac{(l+1)\lambda_2^2 + (4-2l)\lambda_1\lambda_2 - 4\lambda_1^2}{4(1-l)} \right] c_2^2 + c_1^2 h_0^2 - \frac{l\lambda_1 - \lambda_2}{1-l} c_1 c_2 h_0 = 0. \quad (25)$$

If  $c_1 = 0, c_2 \neq 0$ , by the equation (24) we get  $c_2^2 h_0^2 = 0$ , then  $h_0 = 0$ , this is a contradiction with  $h_0 \neq 0$ ;

If  $c_1 \neq 0, c_2 = 0$ , by the equation (25) we get  $c_1^2 h_0^2 = 0$ , then  $h_0 = 0$ , this is a contradiction with  $h_0 \neq 0$ . So  $c_1 \neq 0, c_2 \neq 0$ , (24) + (25) we get

$$d_0 = \frac{[(2l-4)\lambda_1 - l\lambda_2](\lambda_2 - \lambda_1)}{4(1-l)}. \quad (26)$$

Considering  $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$ , we have

$$\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2 > 0. \quad (27)$$

The coefficient of  $\sin(h_0t)\cos(h_0t)e^{\lambda_2t}$  is

$$\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{1-l} c_1 c_2 + \frac{l\lambda_1 - \lambda_2}{1-l} h_0 (c_2^2 - c_1^2) = 0. \quad (28)$$

Using (24) - (25), we get

$$\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)} (c_1^2 - c_2^2) + \frac{2(l\lambda_1 - \lambda_2)}{1-l} h_0 c_1 c_2. \quad (29)$$

Let  $a = \frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)} < 0$ ,  $b = \frac{l\lambda_1 - \lambda_2}{1-l}h_0$ ,  $x = c_1^2 - c_2^2$ ,  $y = c_1c_2 \neq 0$ , from equations (28) and (29) we obtain

$$ax + 2by = 0, \quad (30a)$$

$$2ay - bx = 0. \quad (30b)$$

Through  $(30a) \times b + (30b) \times a$ , then  $a^2 + b^2 = 0$ , which means  $\left[\frac{\lambda_2^2 - 2l\lambda_1^2 + l\lambda_1\lambda_2}{2(1-l)}\right]^2 + \left[\frac{l\lambda_1 - \lambda_2}{1-l}h_0\right]^2 = 0$ , then we get

$$d_0 = \frac{(3l+1)\lambda_1^2\lambda_2^2 - (l+1)\lambda_1\lambda_2^3 - 2l\lambda_1^3\lambda_2}{(l\lambda_1 - \lambda_2)^2}. \quad (31)$$

Using equations (26) and (31), we have  $(\lambda_1 - \lambda_2)[(2-l)\lambda_1 - \lambda_2](\lambda_2^2 + l\lambda_1\lambda_2 - 2l\lambda_2^2) = 0$ . By inequality (27), we get  $\lambda_1 = \lambda_2$  or  $\lambda_2 = (2-l)\lambda_1$ .

1)  $\lambda_1 = \lambda_2$ , by the equation (26) we get  $d_0 = 0$ , this is a contradiction with  $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2 = \frac{1}{4}\lambda_1^2 > 0$ .

2)  $\lambda_2 = (2-l)\lambda_1$ , by the equation (26) we get  $d_0 = \frac{l^2-4}{4}\lambda_1^2 < \frac{l^2}{4}\lambda_1^2$ . On the other hand,  $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2 = \frac{l^2}{4}\lambda_1^2$ , this is a contradiction.

In a word, we have no solution in the case of  $d_0 > (\lambda_1 - \frac{1}{2}\lambda_2)^2$ .  $\square$

**Remark 7.** When  $\lambda_1 = \lambda_2 = 1$ , we get Theorem 26 in [12].

**Proposition 3.17.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2g_F$ ,  $P = \frac{\partial}{\partial t}$ . Then  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$  if and only if  $(F, \nabla^F)$  has constant scalar curvature  $S^F$  and

$$\bar{S} = \frac{S^F}{f^2} - 2l\frac{f''}{f} - l(l-1)\frac{(f')^2}{f^2} + l^2(\lambda_1 + \lambda_2)\frac{f'}{f} + l[\lambda_1^2 + \lambda_2^2 - (l+1)\lambda_1\lambda_2]. \quad (32)$$

*Proof.* Considering that  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2g_F$  and  $P = \frac{\partial}{\partial t}$ , then by Proposition 3.7 we get the equation (32). With the fact that  $S^F$  is function defined on  $F$ , and  $f$  is function defined on  $I$ , then using variables separation we complete the proof of this Proposition.  $\square$

**Proposition 3.18.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2g_F$ ,  $P \in \Gamma(TF)$ . If  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$ , then

- (1) If  $\lambda_1 + \lambda_2 \neq 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{\pi}\lambda_1\lambda_2 = 0$ , and  $\text{div}_F P$  is a constant, then  $S^F$  is a constant;
- (2) If  $\lambda_1 = -\lambda_2 \neq 0$  and  $g_F(P, P)$  is a constant, then  $S^F$  is a constant;



(3) If  $\lambda_1 + \lambda_2 \neq 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n}\lambda_1\lambda_2 \neq 0$ , and  $\text{div}_F P$ ,  $g_F(P, P)$  are constants, then  $S^F$  is a constant.

*Proof.* Considering that  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P \in \Gamma(TF)$ , then by Proposition 3.8, we can get

$$\bar{S} = \frac{S^F}{f^2} - 2l \frac{f''}{f} - l(l-1) \frac{(f')^2}{f^2} + [l\bar{n}\lambda_1\lambda_2 - l(\lambda_1^2 + \lambda_2^2)] f^2 g_F(P, P) + l(\lambda_1 + \lambda_2) \text{div}_F P. \quad (33)$$

Then by variables separation, we have:

(1) If  $\lambda_1 + \lambda_2 \neq 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n}\lambda_1\lambda_2 = 0$ , and  $\text{div}_F P$  is a constant, we can obtain  $\bar{S} = \frac{S^F}{f^2} - 2l \frac{f''}{f} - l(l-1) \frac{(f')^2}{f^2} + l(\lambda_1 + \lambda_2) \text{div}_F P$ . Then  $S^F$  is a constant;

(2) If  $\lambda_1 = -\lambda_2 \neq 0$  and  $g_F(P, P)$  is a constant, which means  $\lambda_1 + \lambda_2 = 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n}\lambda_1\lambda_2 \neq 0$ , and  $g_F(P, P)$  is a constant, we can obtain  $\bar{S} = \frac{S^F}{f^2} - 2l \frac{f''}{f} - l(l-1) \frac{(f')^2}{f^2} + [l\bar{n}\lambda_1\lambda_2 - l(\lambda_1^2 + \lambda_2^2)] f^2 g_F(P, P)$ . then  $S^F$  is a constant;

(3) It is obvious.

(4) If  $\lambda_1 + \lambda_2 = 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n}\lambda_1\lambda_2 = 0$ , then we can get  $\lambda_1 = \lambda_2 = 0$ , which is a contradiction.  $\square$

In (32), we make the change of variable  $f(t) = \sqrt{v(t)}$  and have the following equation:

$$v''(t) + \frac{l-3}{4} \frac{v'(t)^2}{v(t)} - \frac{l}{2} (\lambda_1 + \lambda_2) v'(t) + \left[ (l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l} \right] v(t) - \frac{S^F}{l} = 0. \quad (34)$$

**Remark 8.** When  $\lambda_1 = \lambda_2 = 1$ , equations (32), (33), (34) respectively become (20), (21), (22) in [12], then by Proposition 3.17, we can get Corollary 27 in [12], and by Proposition 3.18.(3), we can get Corollary 28 in [12].

**Theorem 3.19.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2 g_F$ ,  $P = \frac{\partial}{\partial t}$ , and  $\dim F = l = 3$ . Then  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$  if and only if  $(F, \nabla^F)$  has constant scalar curvature  $S^F$  and

(1)  $\bar{S} < \frac{27}{16}(\lambda_1 + \lambda_2)^2 + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$  and  $\bar{S} \neq 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$v(t) = c_1 e^{\left(\left(\frac{3}{2}(\lambda_1 + \lambda_2) + \sqrt{\frac{9}{4}(\lambda_1 + \lambda_2)^2 - \frac{4}{3}\bar{S} + 4\lambda_1^2 + 4\lambda_2^2 - 16\lambda_1\lambda_2}\right)/2\right)t} + c_2 e^{\left(\left(\frac{3}{2}(\lambda_1 + \lambda_2) - \sqrt{\frac{9}{4}(\lambda_1 + \lambda_2)^2 - \frac{4}{3}\bar{S} + 4\lambda_1^2 + 4\lambda_2^2 - 16\lambda_1\lambda_2}\right)/2\right)t} + \frac{S^F}{12\lambda_1\lambda_2 - 3\lambda_1^2 - 3\lambda_2^2 + \bar{S}};$$

(2)  $\bar{S} = \frac{27}{16}(\lambda_1 + \lambda_2)^2 + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$v(t) = c_1 e^{\frac{3}{4}(\lambda_1 + \lambda_2)t} + c_2 t e^{\frac{3}{4}(\lambda_1 + \lambda_2)t} + \frac{S^F}{12\lambda_1\lambda_2 - 3\lambda_1^2 - 3\lambda_2^2 + \bar{S}};$$

(3)  $\bar{S} > \frac{27}{16}(\lambda_1 + \lambda_2)^2 + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$\begin{aligned}
v(t) &= c_1 e^{\frac{3}{4}(\lambda_1 + \lambda_2)t} \cos\left(\left(\sqrt{\frac{4}{3}\bar{S} - 4\lambda_1^2 - 4\lambda_2^2 + 16\lambda_1\lambda_2 - \frac{9}{4}(\lambda_1 + \lambda_2)^2}\right)/2)t\right) \\
&\quad + c_2 e^{\frac{3}{4}(\lambda_1 + \lambda_2)t} \sin\left(\left(\sqrt{\frac{4}{3}\bar{S} - 4\lambda_1^2 - 4\lambda_2^2 + 16\lambda_1\lambda_2 - \frac{9}{4}(\lambda_1 + \lambda_2)^2}\right)/2)t\right) \\
&\quad + \frac{S^F}{12\lambda_1\lambda_2 - 3\lambda_1^2 - 3\lambda_2^2 + \bar{S}}; \\
(4)\bar{S} &= 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2, \text{ and } \lambda_1 + \lambda_2 \neq 0, \quad v(t) = c_1 - \frac{2S^F}{9(\lambda_1 + \lambda_2)} + c_2 e^{\frac{3}{2}(\lambda_1 + \lambda_2)t}; \\
(5)\bar{S} &= 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2, \text{ and } \lambda_1 + \lambda_2 = 0, \quad v(t) = \frac{S^F}{6}t^2 + c_1t + c_2.
\end{aligned}$$

*Proof.* If  $l = 3$ , then we have a simple differential equation as follows:

$$v''(t) - \frac{3}{2}(\lambda_1 + \lambda_2)v'(t) + (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{3})v(t) - \frac{S^F}{3} = 0. \quad (35)$$

If  $\bar{S} \neq 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ , putting  $h(t) = (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{3})v(t) - \frac{S^F}{3}$ , we get  $h''(t) - \frac{3}{2}(\lambda_1 + \lambda_2)h'(t) + (4\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{3})h(t) = 0$ . The above solutions (1) – (3) follow directly from elementary methods for ordinary differential equations.

If  $\bar{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ , and  $\lambda_1 + \lambda_2 \neq 0$ , then  $v''(t) - \frac{3}{2}(\lambda_1 + \lambda_2)v'(t) - \frac{S^F}{3} = 0$ , and we get solution(4).

If  $\bar{S} = 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ , and  $\lambda_1 + \lambda_2 = 0$ , then  $v''(t) - \frac{S^F}{3} = 0$ , and we get solution(5).  $\square$

**Remark 9.** When  $\lambda_1 = \lambda_2 = 1$ , we get Theorem 29 in [12].

**Theorem 3.20.** Let  $M = I \times F$  with the metric tensor  $-dt^2 + f(t)^2g_F$ ,  $P = \frac{\partial}{\partial t}$ , and  $\dim F = l \neq 3$  and  $S^F = 0$ . Then  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$  if and only if

$$\begin{aligned}
(1)\bar{S} &< \frac{l^3}{4(l+1)}(\lambda_1 + \lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2], \\
v(t) &= \left[ c_1 e^{\left(\left(\frac{1}{2}(\lambda_1 + \lambda_2) + \sqrt{\frac{l^2}{4}(\lambda_1 + \lambda_2)^2 - (l+1)[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l}]} \right)/2\right)t} \right. \\
&\quad \left. + c_2 e^{\left(\left(\frac{1}{2}(\lambda_1 + \lambda_2) - \sqrt{\frac{l^2}{4}(\lambda_1 + \lambda_2)^2 - (l+1)[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l}]} \right)/2\right)t} \right]^{\frac{4}{l+1}}; \\
(2)\bar{S} &= \frac{l^3}{4(l+1)}(\lambda_1 + \lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2], \quad v(t) = \left[ c_1 e^{\frac{l}{4}(\lambda_1 + \lambda_2)t} + c_2 t e^{\frac{l}{4}(\lambda_1 + \lambda_2)t} \right]^{\frac{4}{l+1}}; \\
(3)\bar{S} &> \frac{l^3}{4(l+1)}(\lambda_1 + \lambda_2)^2 - l[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2], \\
v(t) &= \left[ c_1 e^{\frac{l}{4}(\lambda_1 + \lambda_2)t} \cos\left(\left(\sqrt{(l+1)[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l}] - \frac{l^2}{4}(\lambda_1 + \lambda_2)^2}\right)/2\right)t \right. \\
&\quad \left. + c_2 e^{\frac{l}{4}(\lambda_1 + \lambda_2)t} \sin\left(\left(\sqrt{(l+1)[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l}] - \frac{l^2}{4}(\lambda_1 + \lambda_2)^2}\right)/2\right)t \right]^{\frac{4}{l+1}}
\end{aligned}$$

*Proof.* In this case, (34) is changed into the simpler form

$$v''(t) + \frac{l-3}{4} \frac{v'(t)^2}{v(t)} - \frac{l}{2}(\lambda_1 + \lambda_2)v'(t) + \left[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\bar{S}}{l}\right]v(t) = 0. \quad (36)$$

Putting  $v(t) = w(t)^{\frac{4}{l+1}}$ , then  $w(t)$  satisfies the equation  $w''(t) - \frac{l}{2}(\lambda_1 + \lambda_2)w'(t) + \frac{l+1}{4}[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}]w(t) = 0$ , by the elementary methods for ordinary differential equations, we prove the above theorem.  $\square$

**Remark 10.** When  $\lambda_1 = \lambda_2 = 1$ , we get Theorem 30 in [12].

When  $\dim F = l \neq 3$ , and  $S^F \neq 0$ , putting  $v(t) = w(t)^{\frac{4}{l+1}}$ , then  $w(t)$  satisfies the following equation :

$$w''(t) - \frac{l}{2}(\lambda_1 + \lambda_2)w'(t) + \frac{l+1}{4}[(l+1)\lambda_1\lambda_2 - \lambda_1^2 - \lambda_2^2 + \frac{\overline{S}}{l}]w(t) - \frac{l+1}{4l}S^F w^{1-\frac{4}{l+1}} = 0. \quad (37)$$

## 4 Multiply Warped Product with a Quarter-symmetric Connection

In this section, firstly we compute curvature of multiply twisted product with a quarter-symmetric connection, secondly we study the special multiply warped product with a quarter-symmetric connection, finally we consider the generalized Kasner space-times with a quarter-symmetric connection.

### 4.1 Connection and Curvature

By Lemma 2.3 and the equation (3), we have the following two Propositions:

**Proposition 4.1.** Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product, If  $X, Y \in \Gamma(TB)$ ,  $U \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$  and  $P \in \Gamma(TB)$ . Then:

- (1)  $\overline{\nabla}_X Y = \overline{\nabla}_X^B Y$ ;
- (2)  $\overline{\nabla}_X U = \frac{Xb_i}{b_i} U$ ;
- (3)  $\overline{\nabla}_U X = \left[ \frac{Xb_i}{b_i} + \lambda_1 \pi(X) \right] U$ ;
- (4)  $\overline{\nabla}_U W = 0$  if  $i \neq j$ ;
- (5)  $\overline{\nabla}_U W = U(\ln b_i)W + W(\ln b_i)U - \frac{g_{F_i}(U, W)}{b_i} \text{grad}_{F_i} b_i - b_i g_{F_i}(U, W) \text{grad}_B b_i + \nabla_U^{F_i} W - \lambda_2 g(U, W)P$  if  $i = j$ .

**Proposition 4.2.** Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product. If  $X, Y \in \Gamma(TB)$ ,  $U \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$  and  $P \in \Gamma(TF_r)$  for a fixed  $r$ , then:

- (1)  $\overline{\nabla}_X Y = \nabla_X^B Y - \lambda_2 g(X, Y)P$ ;
- (2)  $\overline{\nabla}_X U = \frac{Xb_i}{b_i} U + \lambda_1 \pi(U)X$ ;

- (3)  $\bar{\nabla}_U X = \frac{Xb_i}{b_i} U$ ;  
(4)  $\bar{\nabla}_U W = \lambda_1 g(W, P) U$  if  $i \neq j$ ;  
(5)  $\bar{\nabla}_U W = U(\ln b_i) W + W(\ln b_i) U - \frac{g_{F_i}(U, W)}{b_i} \text{grad}_{F_i} b_i - b_i g_{F_i}(U, W) \text{grad}_B b_i + \bar{\nabla}_U^{F_i} W$   
if  $i = j$ .

By Lemmas 2.3, 2.4 and the equation (4), we have the following two Propositions:

**Proposition 4.3.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product. If  $X, Y, Z \in \Gamma(TB)$ ,  $V \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$ ,  $U \in \Gamma(TF_k)$  and  $P \in \Gamma(TB)$ , then:*

- (1)  $\bar{R}(X, Y)Z = \bar{R}^B(X, Y)Z$ ;  
(2)  $\bar{R}(V, X)Y = -\left[\frac{H_B^{b_i}(X, Y)}{f} + \lambda_2 \frac{Pb_i}{b_i} g(X, Y) + \lambda_1 \lambda_2 \pi(P)g(X, Y) + \lambda_1 g(Y, \nabla_X P) - \lambda_1^2 \pi(X)\pi(Y)\right] V$ ;  
(3)  $\bar{R}(X, Y)V = 0$ ;  
(4)  $\bar{R}(V, W)X = VX(\ln b_i)W - WX(\ln b_i)V$  if  $i = j$ ;  
(5)  $\bar{R}(V, W)U = 0$  if  $i = j \neq k$  or  $i \neq j \neq k$ ;  
(6)  $\bar{R}(X, V)W = \bar{R}(V, W)X = \bar{R}(V, X)W = 0$  if  $i \neq j$ ;  
(7)  $\bar{R}(X, V)W = WX(\ln b_i)V - g(V, W)\left[\frac{\nabla_X^B(\text{grad}_B b_i)}{b_i} + \text{grad}_{F_i} \frac{X(\ln b_i)}{b_i^2} + \lambda_1 \frac{Pb_i}{b_i} X + \lambda_2 \nabla_X P + \lambda_1 \lambda_2 \pi(P)X - \lambda_2^2 \pi(X)P\right]$  if  $i = j$ ;  
(8)  $\bar{R}(U, V)W = -g(V, W)U\left[\frac{g_B(\text{grad}_B b_i, \text{grad}_B b_k)}{b_i b_k} + \lambda_1 \frac{Pb_i}{b_i} + \lambda_2 \frac{Pb_k}{b_k} + \lambda_1 \lambda_2 \pi(P)\right]$  if  $i = j \neq k$ ;  
(9)  $\bar{R}(U, V)W = g(U, W)\text{grad}_B(V(\ln b_i)) - g(V, W)\text{grad}_B(U(\ln b_i)) + R^{F_i}(U, V)W - \left[\frac{|\text{grad}_B b_i|_B^2}{b_i^2} + (\lambda_1 + \lambda_2) \frac{Pb_i}{b_i} + \lambda_1 \lambda_2 \pi(P)\right][g(V, W)U - g(U, W)V]$  if  $i = j = k$ .

**Proposition 4.4.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product. If  $X, Y, Z \in \Gamma(TB)$ ,  $V \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$ ,  $U \in \Gamma(TF_k)$  and  $P \in \Gamma(TF_r)$  for a fixed  $r$ , then:*

- (1)  $\bar{R}(X, Y)Z = R^B(X, Y)Z + \lambda_2 \left[g(X, Z) \frac{Yb_r}{b_r} - g(Y, Z) \frac{Xb_r}{b_r}\right] P + \lambda_1 \lambda_2 \pi(P)[g(X, Z)Y - g(Y, Z)X]$ ;  
(2)  $\bar{R}(V, X)Y = -\frac{H_B^{b_i}(X, Y)}{b_i} V - \lambda_1 \lambda_2 \pi(P)g(X, Y)V$  if  $i \neq r$ ;  
(3)  $\bar{R}(V, X)Y = -\frac{H_B^{b_i}(X, Y)}{b_i} - \lambda_1 \pi(V) \frac{Yb_i}{b_i} X - \lambda_2 g(X, Y) \nabla_V P - g(X, Y)[\lambda_1 \lambda_2 \pi(P)V - \lambda_2^2 \pi(V)P]$  if  $i = r$ ;  
(4)  $\bar{R}(X, Y)V = \lambda_1 \pi(V) \left[\frac{Xb_r}{b_r} Y - \frac{Yb_r}{b_r} X\right]$ ;  
(5)  $\bar{R}(V, W)X = -\lambda_1 \delta_i^r \frac{Xb_i}{b_i} \pi(V)W + \lambda_1 \delta_j^r \frac{Xb_j}{b_j} \pi(W)V$  if  $i \neq j$ ;  
(6)  $\bar{R}(V, W)X = VX(\ln b_i)W - WX(\ln b_i)V - \lambda_1 \delta_i^r \frac{Xb_i}{b_i} [\pi(V)W - \pi(W)V]$  if  $i = j$ ;  
(7)  $\bar{R}(V, W)U = 0$  if  $i = j \neq k$  or  $i \neq j \neq k$ ;  
(8)  $\bar{R}(X, V)W = \lambda_1 \frac{Xb_r}{b_r} \pi(W)V$  if  $i \neq j$ ;

$$\begin{aligned}
(9) \bar{R}(X, V)W &= WX(\ln b_i)V - g(V, W)\frac{\nabla_X^B(\text{grad}_B b_i)}{b_i} - \text{grad}_{F_i}(X \ln b_i)g_{F_i}(V, W) \\
&\quad + \lambda_1 \frac{X b_r}{b_r} \pi(W)V - \lambda_1 g(W, \nabla_V P)X - \lambda_2 g(V, W)\frac{X b_r}{b_r} P - \lambda_1 \lambda_2 g(V, W)\pi(P)X \\
&\quad + \lambda_1^2 \pi(W)\pi(V)X \text{ if } i \neq j; \\
(10) \bar{R}(U, V)W &= -g(V, W)\frac{g_B(\text{grad}_B b_i, \text{grad}_B b_k)}{b_i b_k} U - \lambda_1 g(W, \nabla_V P)U - \lambda_2 g(V, W)\nabla_U P - \\
&\quad \lambda_1 \lambda_2 \pi(P)g(V, W)U + \lambda_2^2 g(V, W)\pi(U)P + \lambda_1^2 \pi(W)[\pi(V)U - \pi(U)V] \text{ if } i = j \neq k; \\
(11) \bar{R}(U, V)W &= g(U, W)\text{grad}_B(V \ln b_i) - g(V, W)\text{grad}_B(U \ln b_i) + R^{F_i}(U, V)W \\
&\quad - \frac{|\text{grad}_B b_i|_B^2}{b_i^2} [g(V, W)U - g(U, W)V] + \lambda_1 \lambda_2 \pi(P)[g(U, W)V - g(V, W)U] \\
&\quad \text{if } i = j = k \neq r; \\
(12) \bar{R}(U, V)W &= g(U, W)\text{grad}_B(V \ln b_i) - g(V, W)\text{grad}_B(U \ln b_i) + R^{F_i}(U, V)W \\
&\quad - \frac{|\text{grad}_B f|_B^2}{f^2} [g(V, W)U - g(U, W)V] + \lambda_1 [g(W, \nabla_U P)V - g(W, \nabla_V P)U] \\
&\quad + \lambda_2 [g(U, W)\nabla_V P - g(V, W)\nabla_U P] + \lambda_1 \lambda_2 \pi(P)[g(U, W)V - g(V, W)U] \\
&\quad + \lambda_2^2 [g(V, W)\pi(U) - g(U, W)\pi(V)]P + \lambda_1^2 \pi(W)[\pi(V)U - \pi(U)V] \text{ if } i = j = k = r.
\end{aligned}$$

where  $\delta_i^r$  denotes the Kronecker symbol.

By Propositions 4.3 and 4.4 and the definition of the Ricci curvature tensor, we have the following two Propositions:

**Proposition 4.5.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product,  $\dim M = \bar{n}$ ,  $\dim B = n$ ,  $\dim F_i = l_i$ . If  $X, Y, Z \in \Gamma(TB)$ ,  $V \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$  and  $P \in \Gamma(TB)$ , then:*

$$\begin{aligned}
(1) \bar{Ric}(X, Y) &= \bar{Ric}^B(X, Y) + \sum_{i=1}^m l_i \left[ \frac{H_B^{b_i}(X, Y)}{b_i} + \lambda_2 \frac{P b_i}{b_i} g(X, Y) + \lambda_1 \lambda_2 \pi(P)g(X, Y) \right. \\
&\quad \left. + \lambda_1 g(Y, \nabla_X P) - \lambda_1^2 \pi(X)\pi(Y) \right]; \\
(2) \bar{Ric}(X, V) &= \bar{Ric}(V, X) = (l_i - 1)[VX(\ln b_i)]; \\
(3) \bar{Ric}(V, W) &= 0 \text{ if } i \neq j; \\
(3) \bar{Ric}(V, W) &= \bar{Ric}^{F_i}(V, W) + \left\{ \frac{\Delta_B b_i}{b_i} + (l_i - 1) \frac{|\text{grad}_B b_i|_B^2}{b_i^2} + \sum_{s \neq i} l_s \frac{g_B(\text{grad}_B b_i, \text{grad}_B b_s)}{b_i b_s} \right. \\
&\quad \left. + [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] \pi(P) + \lambda_2 \text{div}_B P + \lambda_2 \sum_{s \neq i} l_s \frac{P b_s}{b_s} + [(\bar{n} - 1)\lambda_1 + (l_i - 1)\lambda_2] \frac{P b_i}{b_i} \right\} g(V, W) \\
&\quad \text{if } i = j.
\end{aligned}$$

where  $\text{div}_B P = \sum_{k=1}^n \varepsilon_k \langle \nabla_{E_k} P, E_k \rangle$ , and  $E_k, 1 \leq k \leq n$  is an orthonormal base of  $B$  with  $\varepsilon_k = g(E_k, E_k)$ .

As a Corollary of Proposition 4.5, we have:

**Corollary 4.6.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product, and  $\dim F_i = l_i > 1$ ,  $P \in \Gamma(TB)$ , then  $(M, \bar{\nabla})$  is mixed Ricci-flat if and only if  $M$  can be*

expressed as a multiply warped product. In particular, if  $(M, \overline{\nabla})$  is Einstein, then  $M$  can be expressed as a multiply warped product.

*Proof.* By Proposition 4.5.(2) and (3), similar to the proof of Theorem 1 in [5], we get this Corollary.  $\square$

**Proposition 4.7.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product,  $\dim M = \overline{n}$ ,  $\dim B = n$ ,  $\dim F_i = l_i$ . If  $X, Y, Z \in \Gamma(TB)$ ,  $V \in \Gamma(TF_i)$ ,  $W \in \Gamma(TF_j)$  and  $P \in \Gamma(TF_r)$  for a fixed  $r$ , then:*

$$\begin{aligned}
(1) \overline{Ric}(X, Y) &= \overline{Ric}^B(X, Y) + \sum_{i=1}^m l_i \frac{H_B^{b_i}(X, Y)}{b_i} + [(\overline{n} - 1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P)g(X, Y) \\
&\quad + \lambda_2 g(X, Y) \operatorname{div}_{F_r} P; \\
(2) \overline{Ric}(X, V) &= (l_i - 1)[VX(\ln b_i)] + [(\overline{n} - 1)\lambda_1 - \lambda_2]\pi(V) \frac{Xb_r}{b_r}; \\
(3) \overline{Ric}(V, X) &= (l_i - 1)[VX(\ln b_i)] + [\lambda_2 - (\overline{n} - 1)\lambda_1]\pi(V) \frac{Xb_r}{b_r}; \\
(4) \overline{Ric}(V, W) &= 0 \text{ if } i \neq j; \\
(4) \overline{Ric}(V, W) &= Ric^{F_i}(V, W) + g(V, W) \left\{ \frac{\Delta_B b_i}{b_i} + (l_i - 1) \frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{s \neq i} l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s} \right. \\
&\quad \left. + [(\overline{n} - 1)\lambda_1\lambda_2 - \lambda_2^2]\pi(P) \right\} + [(\overline{n} - 1)\lambda_1 - \lambda_2]g(W, \nabla_V P) + [\lambda_2^2 + (1 - \overline{n})\lambda_1^2]\pi(V)\pi(W) \\
&\quad + \lambda_2 g(V, W) \operatorname{div}_{F_r} P \text{ if } i = j.
\end{aligned}$$

As a Corollary of Proposition 4.7, we have:

**Corollary 4.8.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product, and  $\dim F_i = l_i > 1$ ,  $P \in \Gamma(TF_r)$ , then  $(M, \overline{\nabla})$  is mixed Ricci-flat if and only if one of the following two conditions is satisfied:*

- (1)  $\lambda_2 = (\overline{n} - 1)\lambda_1$ , and  $M$  can be expressed as a multiply warped product;
- (2)  $\lambda_2 \neq (\overline{n} - 1)\lambda_1$ ,  $M$  can be expressed as a multiply warped product and  $b_r$  is only dependent on  $F_r$ ;

*In particular, if  $(M, \overline{\nabla})$  is Einstein, then  $M$  can be expressed as a multiply warped product.*

*Proof.* By Proposition 4.7.(2) and (3), we have that  $(M, \overline{\nabla})$  is mixed Ricci-flat if and only if  $VX(\ln b_i) = 0$ ,  $[(\overline{n} - 1)\lambda_1 - \lambda_2]\pi(V) \frac{Xb_r}{b_r} = 0$ . Similar to the proof of Corollary 4.6, we get that

$\lambda_2 = (\overline{n} - 1)\lambda_1$ , and  $M$  can be expressed as a multiply warped product.

$\lambda_2 \neq (\overline{n} - 1)\lambda_1$ ,  $M$  can be expressed as a multiply warped product. When  $i \neq r$ ,  $\pi(V) = 0$ . When  $i = r$ , by  $\pi(V) \frac{Xb_r}{b_r} = 0$ , then  $b_r$  depends only on  $F_r$ .  $\square$

By Proposition 4.5 and the definition of the scalar curvature, we have the following:

**Proposition 4.9.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product,  $\dim M = \bar{n}$ ,  $\dim B = n$ ,  $\dim F_i = l_i$ . If  $P \in \Gamma(TB)$ , then the scalar curvature  $\bar{S}$  has the following expression:*

$$\begin{aligned} \bar{S} = \bar{S}^B &+ 2 \sum_{i=1}^m l_i \frac{\Delta_B b_i}{b_i} + \sum_{i=1}^m \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^m l_i (l_i - 1) \frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s} \\ &+ \sum_{i=1}^m l_i [(\bar{n} - 1)\lambda_1 + (n + l_i - 1)\lambda_2] \frac{P b_i}{b_i} + \lambda_2 \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{P b_s}{b_s} + \sum_{i=1}^m l_i [(\bar{n} + n - 1)\lambda_1 \lambda_2 \\ &- (\lambda_1^2 + \lambda_2^2)] \pi(P) + (\lambda_1 + \lambda_2) \sum_{i=1}^m l_i div_B P. \end{aligned}$$

By Proposition 4.7 and the definition of the scalar curvature, we have the following:

**Proposition 4.10.** *Let  $M = B \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply twisted product,  $\dim M = \bar{n}$ ,  $\dim B = n$ ,  $\dim F_i = l_i$ . If  $P \in \Gamma(TF_r)$ , then the scalar curvature  $\bar{S}$  has the following expression:*

$$\begin{aligned} \bar{S} = \bar{S}^B &+ 2 \sum_{i=1}^m l_i \frac{\Delta_B b_i}{b_i} + \sum_{i=1}^m \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^m l_i (l_i - 1) \frac{|grad_B b_i|_B^2}{b_i^2} + \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{g_B(grad_B b_i, grad_B b_s)}{b_i b_s} \\ &+ [\bar{n}(\bar{n} - 1)\lambda_1 \lambda_2 + (1 - \bar{n})(\lambda_1^2 + \lambda_2^2)] \pi(P) + (\bar{n} - 1)(\lambda_1 + \lambda_2) div_{F_r} P. \end{aligned}$$

**Remark 11.** (1) *It is easy to see that Propositions 3.1 – 3.8 are Corollaries of Propositions 4.1, 4.2, 4.3, 4.4, 4.5, 4.7, 4.9, 4.10, respectively.*

(2) *When  $\lambda_1 = \lambda_2 = 1$ , we get Propositions 1, 2, 4, 5, 7, 9, 12, 13 in [12], by Propositions 4.1, 4.2, 4.3, 4.4, 4.5, 4.7, 4.9, 4.10, respectively.*

## 4.2 Special Multiply Warped Product with a Quarter-symmetric Connection

Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product with the metric tensor  $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$  and  $I$  is an open interval in  $\mathbb{R}$ , and  $b_i \in C^\infty(I)$ ,  $\dim M = \bar{n}$ ,  $\dim I = 1$ ,  $\dim F_i = l_i$ .

Similar to the proof method of Theorem 3.11, we have:

**Theorem 4.11.** *Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product with the metric tensor  $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ ,  $P = \frac{\partial}{\partial t}$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if the following conditions are satisfied*

- (1)  $(F_i, \nabla^{F_i})$  is Einstein with the Einstein constant  $\alpha_i$ ,  $i \in \{1, \dots, m\}$ ;
- (2)  $\sum_{i=1}^m l_i \left( \lambda_2 \frac{b'_i}{b_i} - \frac{b''_i}{b_i} + \lambda_1^2 - \lambda_1 \lambda_2 \right) = \alpha$ ;

$$(3) \quad \alpha_i - b_i b_i'' + (1 - l_i)(b_i')^2 + (\lambda_2 b_i^2 - b_i b_i') \sum_{s \neq i} l_s \frac{b_s'}{b_s} + [\lambda_2^2 + (1 - \bar{n})\lambda_1 \lambda_2] b_i^2 + [(\bar{n} - 1)\lambda_1 + (l_i - 1)\lambda_2] b_i b_i' = \alpha b_i^2.$$

**Theorem 4.12.** *Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product with the metric tensor  $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ ,  $P \in \Gamma(TF_r)$  with  $g_{F_r}(P, P) = 1$  and  $\bar{n} > 2$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if the following conditions are satisfied for any  $i \in \{1, \dots, m\}$ :*

- (1)  $(F_i, \nabla^{F_i})(i \neq r)$  is Einstein with the Einstein constant  $\alpha_i, i \in \{1, \dots, m\}$ ;
- (2)  $b_r$  is a constant and  $\sum_{i=1}^m l_i \frac{b_i''}{b_i} = \mu_0$ ;  $\text{div}_{F_r} P = \mu_1$ ;  $\mu_0 - \lambda_2 \mu_1 + \alpha = [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_r^2$ , where  $\mu_0, \mu_1$  are constants;
- (3)  $\text{Ric}^{F_r}(V, W) + \bar{\alpha} g_{F_r}(V, W) = [(\bar{n} - 1)\lambda_1^2 - \lambda_2^2] \pi(V) \pi(W) - [(\bar{n} - 1)\lambda_1 - \lambda_2] g(W, \nabla_V P)$ , for  $V, W \in \Gamma(TF_r)$ ;
- (4)  $\alpha_i - b_i b_i'' + [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_i^2 b_r^2 - b_i b_i' \sum_{s \neq i} l_s \frac{b_s'}{b_s} - (l_i - 1)(b_i')^2 = (\alpha - \lambda_2 \mu_1) b_i^2$ .

*Proof.* By Proposition 4.7(2) and  $g_{F_r}(P, P) = 1$ , we have that  $b_r$  is a constant. By Proposition 4.7(1), we have

$$\overline{\text{Ric}}\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}\right) = \sum_{i=1}^m l_i \frac{b_i''}{b_i} + [\lambda_2^2 + (1 - \bar{n})\lambda_1 \lambda_2] b_r^2 - \lambda_2 \text{div}_{F_r} P = -\alpha.$$

By variables separation, we have

$$\sum_{i=1}^m l_i \frac{b_i''}{b_i} = \mu_0; \quad \text{div}_{F_r} P = \mu_1; \quad \mu_0 - \lambda_2 \mu_1 + \alpha = [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_r^2,$$

then we get (2). By Proposition 4.7(3), we have

$$\begin{aligned} \overline{\text{Ric}}(V, W) &= \text{Ric}^{F_i}(V, W) + b_i^2 g_{F_i}(V, W) \left\{ -\frac{b_i''}{b_i} + (l_i - 1) \frac{-(b_i')^2}{b_i^2} + \sum_{s \neq i} l_s \frac{-b_i' b_s'}{b_i b_s} \right. \\ &\quad \left. + [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] \pi(P) \right\} + [(\bar{n} - 1)\lambda_1 - \lambda_2] g(W, \nabla_V P) \\ &\quad + [\lambda_2^2 + (1 - \bar{n})\lambda_1^2] \pi(V) \pi(W) + \lambda_2 g(V, W) \text{div}_{F_r} P. \end{aligned}$$

When  $i \neq r$ , then  $\nabla_V P = \pi(V) = 0$ , so

$$\begin{aligned} \overline{\text{Ric}}(V, W) &= \text{Ric}^{F_i}(V, W) + b_i^2 g_{F_i}(V, W) \left\{ -\frac{b_i''}{b_i} + (l_i - 1) \frac{-(b_i')^2}{b_i^2} + \sum_{s \neq i} l_s \frac{-b_i' b_s'}{b_i b_s} \right. \\ &\quad \left. + [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_r^2 \right\} + \lambda_2 \mu_1 b_i^2 g_{F_i}(V, W) = \alpha b_i^2 g_{F_i}(V, W). \end{aligned}$$



By variables separation, we have that  $(F_i, \nabla^{F_i})(i \neq r)$  is Einstein with the Einstein constant  $\alpha_i$  and

$$\alpha_i - b_i b_i'' + [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_i^2 b_r^2 - b_i b_i' \sum_{s \neq i} l_s \frac{b_s'}{b_s} - (l_i - 1)(b_i')^2 = (\alpha - \lambda_2 \mu_1) b_i^2.$$

Then we get (1) and (4).

When  $i = r$  and  $b_r$  is a constant, then

$$\begin{aligned} Ric^{F_r}(V, W) + b_r^2 \{ [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_r^2 + \lambda_2 \mu_1 - \alpha \} g_{F_r}(V, W) \\ = [(\bar{n} - 1)\lambda_1^2 - \lambda_2^2] \pi(V) \pi(W) - [(\bar{n} - 1)\lambda_1 - \lambda_2] g(W, \nabla_V P), \end{aligned}$$

let  $\bar{\alpha} = b_r^2 \{ [(\bar{n} - 1)\lambda_1 \lambda_2 - \lambda_2^2] b_r^2 + \lambda_2 \mu_1 - \alpha \}$ , we get (3).  $\square$

**Theorem 4.13.** *Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product and  $P = \frac{\partial}{\partial t}$ . If  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$ , then each  $(F_i, \nabla^{F_i})$  has constant scalar curvature  $S^{F_i}$ .*

*Proof.* By Proposition 4.9, we have

$$\begin{aligned} \bar{S} = & -2 \sum_{i=1}^m l_i \frac{b_i''}{b_i} + \sum_{i=1}^m \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^m l_i (l_i - 1) \frac{-(b_i')^2}{b_i^2} + \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{-b_i' b_s'}{b_i b_s} \\ & + \sum_{i=1}^m l_i [(\bar{n} - 1)\lambda_1 + l_i \lambda_2] \frac{b_i'}{b_i} + \lambda_2 \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{b_s'}{b_s} \\ & - \sum_{i=1}^m l_i [\bar{n} \lambda_1 \lambda_2 - (\lambda_1^2 + \lambda_2^2)]. \end{aligned} \quad (38)$$

Note that each  $S^{F_i}$  is function defined on  $F_i$ , using variables separation we complete this proof.  $\square$

**Theorem 4.14.** *Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product and  $P \in \Gamma(TF_r)$ . If  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$ , then*

- (1) *each  $(F_i, \nabla^{F_i})(i \neq r)$  has constant scalar curvature  $S^{F_i}$ ;*
- (2) *If  $\lambda_1 + \lambda_2 \neq 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n} \lambda_1 \lambda_2 = 0$ , and  $\text{div}_{F_r} P$  is a constant, then  $S^{F_r}$  is a constant;*
- (3) *If  $\lambda_1 = -\lambda_2 \neq 0$  and  $g_{F_r}(P, P)$  is a constant, then  $S^{F_r}$  is a constant;*
- (4) *If  $\lambda_1 + \lambda_2 \neq 0$  and  $\lambda_1^2 + \lambda_2^2 - \bar{n} \lambda_1 \lambda_2 \neq 0$ , and  $\text{div}_{F_r} P, g_{F_r}(P, P)$  are constants, then  $S^{F_r}$  is a constant.*

*Proof.* By Proposition 4.10, we have

$$\begin{aligned} \overline{S} = & -2 \sum_{i=1}^m l_i \frac{b_i''}{b_i} + \sum_{i=1}^m \frac{S^{F_i}}{b_i^2} + \sum_{i=1}^m l_i (l_i - 1) \frac{-(b_i')^2}{b_i^2} + \sum_{i=1}^m \sum_{s \neq i} l_i l_s \frac{-b_i' b_s'}{b_i b_s} \\ & + [\overline{n}(\overline{n} - 1) \lambda_1 \lambda_2 + (1 - \overline{n})(\lambda_1^2 + \lambda_2^2)] b_r^2 g_{F_r}(P, P) + (\overline{n} - 1)(\lambda_1 + \lambda_2) \operatorname{div}_{F_r} P \end{aligned} \quad (39)$$

then similar to the proof of Proposition 3.18, we complete this Proposition.  $\square$

**Remark 12.** When  $\lambda_1 = \lambda_2 = 1$ , we get Theorems 15, 16, Propositions 18, 19 in [12] by Theorems 4.11 – 4.14, respectively.

### 4.3 Generalized Kasner Space-times with a Quarter-symmetric Connection

In this section, we consider the Einstein and scalar curvature of generalized Kasner space-times with a quarter-symmetric connection. We recall the definition of generalized Kasner space-times in [5].

**Definition 4.15.** A generalized Kasner space-time  $(M, g)$  is a Lorentzian multiply warped product of the form  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  with the metric tensor  $g = -dt^2 \oplus \phi^{2p_1} g_{F_1} \oplus \phi^{2p_2} g_{F_2} \cdots \oplus \phi^{2p_m} g_{F_m}$ , where  $\phi : I \rightarrow (0, \infty)$  is smooth and  $p_i \in \mathbb{R}$ , for any  $i \in \{1, \dots, m\}$  and also  $I = (t_1, t_2)$ .

We introduce the following parameters  $\zeta = \sum_{i=1}^m l_i p_i$  and  $\eta = \sum_{i=1}^m l_i p_i^2$  for generalized Kasner space-times. By Theorem 4.11 and direct computations, we get the following:

**Proposition 4.16.** Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  be a generalized Kasner space-time and  $P = \frac{\partial}{\partial t}$ . Then  $(M, \overline{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if the following conditions are satisfied for any  $i \in \{1, \dots, m\}$ :

- (1)  $(F_i, \nabla^{F_i})$  is Einstein with the Einstein constant  $\alpha_i, i \in \{1, \dots, m\}$ ;
- (2)  $\zeta \left( \lambda_2 \frac{\phi'}{\phi} - \frac{\phi''}{\phi} \right) - (\eta - \zeta) \frac{(\phi')^2}{\phi^2} + (\lambda_1^2 - \lambda_1 \lambda_2)(\overline{n} - 1) = \alpha$ ;
- (3)  $\frac{\alpha_i}{\phi^{2p_i}} - p_i \frac{\phi''}{\phi} - (\zeta - 1) p_i \frac{(\phi')^2}{\phi^2} + \{ \lambda_2 \zeta + [(\overline{n} - 1) \lambda_1 - \lambda_2] p_i \} \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + (\overline{n} - 1) \lambda_1 \lambda_2$ .

By the equation (38), we obtain the following:

**Proposition 4.17.** Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  be a generalized Kasner space-time and  $P = \frac{\partial}{\partial t}$ . Then  $(M, \overline{\nabla})$  has constant scalar curvature  $\overline{S}$  if and only if

each  $(F_i, \nabla^{F_i})$  has constant scalar curvature  $S^{F_i}$  and

$$\overline{S} = \sum_{i=1}^m \frac{S^{F_i}}{\phi^{2p_i}} - 2\zeta \frac{\phi''}{\phi} - (\eta + \zeta^2 - 2\zeta) \frac{(\phi')^2}{\phi^2} + (\lambda_1 + \lambda_2)\zeta(\overline{n} - 1) \frac{\phi'}{\phi} + (\overline{n} - 1)(\lambda_1^2 + \lambda_2^2 - \overline{n}\lambda_1\lambda_2). \quad (40)$$

Next, we first give a classification of four-dimensional generalized Kasner space-times with a quarter-symmetric connection and then consider Ricci tensors and scalar curvatures of them.

**Definition 4.18.** Let  $M = I \times_{b_1} F_1 \times_{b_2} F_2 \cdots \times_{b_m} F_m$  be a multiply warped product with the metric tensor  $g = -dt^2 \oplus b_1^2 g_{F_1} \oplus b_2^2 g_{F_2} \cdots \oplus b_m^2 g_{F_m}$ . Then:

- (1)  $(M, g)$  is said to be of type (I) if  $m = 1$  and  $\dim F = 3$ ;
- (2)  $(M, g)$  is said to be of type (II) if  $m = 2$  and  $\dim F_1 = 1$  and  $\dim F_2 = 2$ ;
- (3)  $(M, g)$  is said to be of type (III) if  $m = 3$  and  $\dim F_1 = 1$ ,  $\dim F_2 = 1$ ,  $\dim F_3 = 1$ .

#### 4.3.1 Classification of Einstein Type (I) generalized Kasner space-times with a quarter-symmetric connection

By Theorem 3.16, we have given a classification of Einstein Type (I) generalized Kasner space-times with a quarter-symmetric connection.

#### 4.3.2 Type (I) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Theorem 3.19, we have given a classification of Type (I) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature.

#### 4.3.3 Classification of Einstein Type (II) generalized Kasner space-times with a quarter-symmetric connection

Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2$  be an Einstein Type (II) generalized Kasner space-times and  $P = \frac{\partial}{\partial t}$ . Then  $\alpha_1 = 0$  because of  $\dim F_1 = 1$ .  $\zeta = p_1 + 2p_2$ ,  $\eta = p_1^2 + 2p_2^2$ . By Proposition 4.16, we have

$$\zeta \left( \lambda_2 \frac{\phi'}{\phi} - \frac{\phi''}{\phi} \right) - (\eta - \zeta) \frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 - \lambda_1\lambda_2) = \alpha. \quad (41a)$$

$$-p_1 \frac{\phi''}{\phi} - (\zeta - 1)p_1 \frac{(\phi')^2}{\phi^2} + [\lambda_2 \zeta + (3\lambda_1 - \lambda_2)p_1] \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2. \quad (41b)$$

$$\frac{\alpha_2}{\phi^{2p_2}} - p_2 \frac{\phi''}{\phi} - (\zeta - 1)p_2 \frac{(\phi')^2}{\phi^2} + [\lambda_2 \zeta + (3\lambda_1 - \lambda_2)p_2] \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2. \quad (41c)$$

where  $\alpha_2$  is a constant. Consider the following two cases:

Case (i) ( $\zeta = 0$ ). In this case,  $p_2 = -\frac{1}{2}p_1$ ,  $\eta = \frac{3}{2}p_1^2$ . Then by equations (41a) – (41c), we have:

$$-\eta \frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 - \lambda_1 \lambda_2) = \alpha. \quad (42a)$$

$$p_1 \left[ -\frac{\phi''}{\phi} + \frac{(\phi')^2}{\phi^2} + (3\lambda_1 - \lambda_2) \frac{\phi'}{\phi} \right] = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2. \quad (42b)$$

$$\frac{\alpha_2}{\phi^{-p_1}} - \frac{1}{2}p_1 \left[ -\frac{\phi''}{\phi} + \frac{(\phi')^2}{\phi^2} + (3\lambda_1 - \lambda_2) \frac{\phi'}{\phi} \right] = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2. \quad (42c)$$

$\eta = 0$ . We have  $p_i = 0$ , by the equation (42a), we get  $\alpha = 3\lambda_1^2 - 3\lambda_1 \lambda_2$ ; by the equation (42b), we get  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2$ ; then we have  $\lambda_2^2 = 3\lambda_1^2$  and by the equation (42c), we get  $\alpha_2 = 0$ . So we have  $\lambda_2^2 = 3\lambda_1^2$ ,  $p_i = 0$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1 \lambda_2 = \lambda_2^2 - 3\lambda_1 \lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ .

$\eta \neq 0$ . We have  $p_i \neq 0$ .

1)  $\alpha_2 = 0$ . By equations (42b), (42c), we get  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2$  and

$$-\frac{\phi''}{\phi} + \frac{(\phi')^2}{\phi^2} + (3\lambda_1 - \lambda_2) \frac{\phi'}{\phi} = 0; \quad (43a)$$

$$\frac{(\phi')^2}{\phi^2} = \frac{3\lambda_1^2 - \lambda_2^2}{\eta}. \quad (43b)$$

1'.  $3\lambda_1^2 - \lambda_2^2 < 0$ , we have no solution;

2'.  $3\lambda_1^2 - \lambda_2^2 = 0$ , we have  $\phi = c$ , then by the equation (42b), we get  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2$ .

So we have  $\lambda_2^2 = 3\lambda_1^2$ ,  $p_1 \neq 0$ ,  $p_2 \neq 0$ ,  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\phi = c$ .

3'.  $3\lambda_1^2 - \lambda_2^2 > 0$ , we have  $\phi = c_0 e^{\pm \sqrt{\frac{3\lambda_1^2 - \lambda_2^2}{\eta}} t}$ , by the equation (43a), we get  $\lambda_2 = 3\lambda_1$ , considering that  $3\lambda_1^2 - \lambda_2^2 > 0$ , we have  $\lambda_1^2 < 0$ , which is a contradiction.

2)  $\alpha_2 \neq 0$ . By equations (42b), (42c), we get  $\frac{\alpha_2}{\phi^{-p_1}} = \frac{3}{2}(\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2)$ , so  $\phi = c$ ; then by the equation (42b), we get  $\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 = 0$ , then  $\alpha_2 = 0$ , this is a contradiction.

Case (ii) ( $\zeta \neq 0$ ). Then  $\eta \neq 0$ . Putting  $\phi = \psi^{\frac{\zeta}{\eta}}$ , then  $\psi'' - \lambda_2 \psi' + (\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2) \frac{\eta}{\zeta^2} \psi = 0$ . Hence:

$$(1) \alpha < \frac{\lambda_2^2 \zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1 \lambda_2, \quad \psi = c_1 e^{\frac{\lambda_2 + \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2) \frac{\eta}{\zeta^2}}}{2} t} + c_2 e^{\frac{\lambda_2 - \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2) \frac{\eta}{\zeta^2}}}{2} t},$$

$$(2) \alpha = \frac{\lambda_2^2 \zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1 \lambda_2, \quad \psi = c_1 e^{\frac{\lambda_2}{2} t} + c_2 t e^{\frac{\lambda_2}{2} t};$$

$$(3) \alpha > \frac{\lambda_2^2 \zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1 \lambda_2,$$

$$\psi = c_1 e^{\frac{\lambda_2}{2}t} \cos\left(\frac{\sqrt{4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}-\lambda_2^2}}{2}t\right) + c_2 e^{\frac{\lambda_2}{2}t} \sin\left(\frac{\sqrt{4(\alpha+3\lambda_1\lambda_2-3\lambda_1^2)\frac{\eta}{\zeta^2}-\lambda_2^2}}{2}t\right).$$

We make equations (41a) – (41c) into

$$\frac{\zeta^2}{\eta} \frac{\lambda_2 \psi' - \psi''}{\psi} = \alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2; \quad (44a)$$

$$-\frac{p_1}{\zeta} \frac{(\phi^\zeta)''}{\phi^\zeta} + \frac{\lambda_2 \zeta + (3\lambda_1 - \lambda_2)p_1}{\zeta} \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2; \quad (44b)$$

$$\frac{\alpha_2}{\phi^{2p_2}} - \frac{p_2}{\zeta} \frac{(\phi^\zeta)''}{\phi^\zeta} + \frac{\lambda_2 \zeta + (3\lambda_1 - \lambda_2)p_2}{\zeta} \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2. \quad (44c)$$

When  $p_1 = p_2$ , type (II) spaces turn into type (I) spaces, so we assume  $p_1 \neq p_2$ . By (44b)  $\times p_2 - (44c) \times p_1$ , we get

$$\psi' = \frac{p_1 \alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} \psi^{1-\frac{2p_2\zeta}{\eta}} + \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} \psi. \quad (45)$$

$\alpha < \frac{\lambda_2^2 \zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1 \lambda_2$ ,  $\psi = c_1 e^{at} + c_2 e^{bt}$ , where  $a = \frac{\lambda_2 + \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}$ ,  $b = \frac{\lambda_2 - \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2)\frac{\eta}{\zeta^2}}}{2}$ . By the equation (45), we get

$$ac_1 e^{at} + bc_2 e^{bt} = \frac{p_1 \alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} (c_1 e^{at} + c_2 e^{bt})^{1-\frac{2p_2\zeta}{\eta}} + \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} (c_1 e^{at} + c_2 e^{bt}). \quad (46)$$

1)  $c_1 = 0$ . We have

$$\left[ b - \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} \right] c_2 e^{bt} = \frac{p_1 \alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} (c_2 e^{bt})^{1-\frac{2p_2\zeta}{\eta}} \quad (47)$$

1'.  $b \neq 0$ ,  $p_1 \alpha_2 \neq 0$ . By the equation (47), we get  $p_2 = 0$ , so  $\zeta = p_1$ ,  $\eta = p_1^2$  and  $\frac{\zeta^2}{\eta} = 1$ . Then  $b = \frac{\lambda_2 - \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2)}}{2}$ , so

$$b^2 - \lambda_2 b = 3\lambda_1^2 - 3\lambda_1 \lambda_2 - \alpha. \quad (48)$$

On the other hand,  $\phi^\zeta = \psi = c_2 e^{bt}$ , so by equations (44b), (44c), we get

$$-b^2 + 3\lambda_1 b = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2; \quad (49a)$$

$$\alpha_2 + \lambda_2 b = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2; \quad (49b)$$

(48) + (49a), we get  $(3\lambda_1 - \lambda_2)b = 3\lambda_1^2 - \lambda_2^2$ , if  $\lambda_2 = 3\lambda_1$ , then  $0 = -6\lambda_1^2$ , this is a contradiction with  $\lambda_1 \neq 0$ , so  $\lambda_2 \neq 3\lambda_1$ ,  $b = \frac{3\lambda_1^2 - \lambda_2^2}{3\lambda_1 - \lambda_2}$ . Then by the equation (48), we have  $\alpha = \frac{18\lambda_1^4 - 36\lambda_1^3\lambda_2 + 24\lambda_1^2\lambda_2^2 - 6\lambda_1\lambda_2^3}{(3\lambda_1 - \lambda_2)^2} = \frac{6\lambda_1(\lambda_1 - \lambda_2)(3\lambda_1^2 - 3\lambda_1\lambda_2 + \lambda_2^2)}{(3\lambda_1 - \lambda_2)^2}$ , by the equation (49b), we get  $\alpha_2 = \frac{18\lambda_1^4 - 18\lambda_1^3\lambda_2 + 6\lambda_1\lambda_2^3 - 2\lambda_2^4}{(3\lambda_1 - \lambda_2)^2} = \frac{2(3\lambda_1^2 - \lambda_2^2)(3\lambda_1^2 - 3\lambda_1\lambda_2 + \lambda_2^2)}{(3\lambda_1 - \lambda_2)^2}$ , since  $3\lambda_1^2 - 3\lambda_1\lambda_2 + \lambda_2^2 \neq 0$ , and  $\alpha_2 \neq 0$ , we have  $3\lambda_1^2 \neq \lambda_2^2$ . Considering that  $b < \frac{\lambda_2}{2}$ , we get  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ . So we obtain  $3\lambda_1^2 \neq \lambda_2^2$ ,  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\alpha = \frac{18\lambda_1^4 - 36\lambda_1^3\lambda_2 + 24\lambda_1^2\lambda_2^2 - 6\lambda_1\lambda_2^3}{(3\lambda_1 - \lambda_2)^2}$ ,  $\alpha_1 = 0$ ,  $\alpha_2 = \frac{18\lambda_1^4 - 18\lambda_1^3\lambda_2 + 6\lambda_1\lambda_2^3 - 2\lambda_2^4}{(3\lambda_1 - \lambda_2)^2}$ ,  $\phi = c_0 e^{\frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)p_1}t}$ .

2'.  $b \neq 0$ ,  $p_1\alpha_2 = 0$ .

1°.  $p_1 = 0$ . Then  $\zeta = 2p_2$ ,  $\eta = 2p_2^2$ ,  $\frac{\eta}{\zeta^2} = \frac{1}{2}$ . By the equation (47), we get  $b = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{2\lambda_2}$ , on the other hand,  $b = \frac{\lambda_2 - \sqrt{\lambda_2^2 - 2(\alpha + 3\lambda_1\lambda_2 - 3\lambda_1^2)}}{2}$ , so we get

$$\alpha^2 + (6\lambda_1\lambda_2 - 2\lambda_2^2)\alpha + 3\lambda_1^2\lambda_2^2 - 6\lambda_1^3\lambda_2 + 3\lambda_1^4 = 0. \quad (50)$$

when  $\lambda_1 = \lambda_2$ , we get  $\alpha^2 + 4\lambda_1^2\alpha = 0$ , then  $\alpha = 0$  or  $\alpha = -4\lambda_1^2$ . If  $\alpha = 0$ , we have  $b = \lambda_1 = \lambda_2 < 0$ , by the equation (44c), we get  $\alpha_2 = 0$ ;  $\phi = \psi^{\frac{\zeta}{\eta}} = c_0 e^{\frac{\lambda_1}{p_2}t}$ . If  $\alpha = -4\lambda_1^2$ , then  $\lambda_1 = \lambda_2 > 0$ ,  $b = -\lambda_1$ , then by the equation (44c), we get  $\lambda_1 = 0$ , which is a contradiction.

When  $\lambda_1 \neq \lambda_2$ ,  $\Delta = 8\lambda_2^2(3\lambda_1^2 - \lambda_2^2)$ , if  $\lambda_2^2 > 3\lambda_1^2$ , we have no solution; if  $\lambda_2^2 = 3\lambda_1^2$ , we have  $b = 0$ , which is a contradiction; if  $\lambda_2^2 < 3\lambda_1^2$ , we have  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 + \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$  or  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 - \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$ . By the equation (44c), we have  $\alpha_2 = 0$  and

$$-2b^2 + (3\lambda_1 + \lambda_2)b = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \quad (51)$$

When  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 + \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$ , we have  $b = \frac{\sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}}{2\lambda_2}$ , by the equation (51), we have  $\lambda_1 = \lambda_2$ , which is a contradiction. When  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 - \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$ , we have the same contradiction.

So we get  $\lambda_1 = \lambda_2 < 0$ ,  $p_1 = 0$ ,  $p_2 \neq 0$ ,  $\alpha = \alpha_1 = \alpha_2 = 0$ ,  $\phi = c_0 e^{\frac{\lambda_1}{p_2}t}$ .

2°.  $\alpha_2 = 0$ . Since  $p_1 = 0$  we have discussed, so we assume that  $p_1 \neq 0$ . By the equation (47), we have  $b = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}$ ,  $\phi\zeta = c e^{b\frac{\zeta^2}{\eta}t}$ , then by the equation (44b), we have  $\alpha(\alpha - \lambda_2^2 + 3\lambda_1\lambda_2) = 0$ . So when  $\lambda_2 = 3\lambda_1$ , we have  $\alpha = 0$ ; when  $\lambda_2 \neq 3\lambda_1$ , we have  $\alpha = 0$  or  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$ .

When  $\lambda_2 = 3\lambda_1$ ,  $\alpha = 0$ , then  $b = 0$ , which is a contradiction;

when  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$ , then  $b = 0$ , which is a contradiction;

when  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = 0$ , then  $b = (3\lambda_1 - \lambda_2)\frac{\eta}{\zeta^2}$ , by the equation (44a), we get  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ , then  $b = \frac{3\lambda_1^2 - \lambda_2^2}{3\lambda_1 - \lambda_2}$ . Since  $b < \frac{\lambda_2}{2}$ , we get  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ , by  $\alpha <$

$\frac{\lambda_2^2 \zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1 \lambda_2$ , we get  $\lambda_2^2 < 3\lambda_1^2$ .  $\phi = \psi^{\frac{\zeta}{\eta}} = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta} t}$ .

So we have  $\lambda_2^2 < 3\lambda_1^2$ ,  $\frac{6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ ,  $p_1 \neq 0$ ,  $p_2 \neq -\frac{1}{2}p_1$ ,  $\alpha = \alpha_1 = \alpha_2 = 0$ ,

$$\phi = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta} t}.$$

3'.  $b = 0$ . Then  $\psi = c_2$ , by the equation (44a), we have  $\alpha = 3\lambda_1^2 - 3\lambda_1 \lambda_2$ ; by the equation (44b), we have  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2$ ; so  $\lambda_2^2 = 3\lambda_1^2$ , by the equation (44c), we have  $\alpha_2 = 0$ .

So we have  $\lambda_2^2 = 3\lambda_1^2$ ,  $\zeta \neq 0$ ,  $\eta \neq 0$ ,  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2 = 3\lambda_1^2 - 3\lambda_1 \lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ ,

$$\phi = c.$$

2)  $c_2 = 0$ . We have

$$\left[ a - \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} \right] c_1 e^{at} = \frac{p_1 \alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} (c_1 e^{at})^{1 - \frac{2p_2 \zeta}{\eta}} \quad (52)$$

1'.  $p_1 \alpha_2 \neq 0$ . By the equation (52), we get  $p_2 = 0$ , so  $\zeta = p_1$ ,  $\eta = p_1^2$  and  $\frac{\zeta^2}{\eta} = 1$ . Then  $a = \frac{\lambda_2 + \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2)}}{2}$ , using the equation (52) again, we get  $a - \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{\lambda_2} = \frac{p_1 \alpha_2}{-\lambda_2 p_1} = -\frac{\alpha_2}{\lambda_2}$ , so  $\alpha_2 = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 - \lambda_2 a$ . By the equation (44b), we get  $(3\lambda_1 - \lambda_2)^2 \alpha = 18\lambda_1^4 - 6\lambda_1 \lambda_2^3 + 24\lambda_1^2 \lambda_2^2 - 36\lambda_1^3 \lambda_2$ , if  $3\lambda_1 = \lambda_2$ , then  $0 = -36\lambda_1^4$ , which is a contradiction,

so  $3\lambda_1 \neq \lambda_2$  and  $\alpha = \frac{18\lambda_1^4 - 6\lambda_1 \lambda_2^3 + 24\lambda_1^2 \lambda_2^2 - 36\lambda_1^3 \lambda_2}{(3\lambda_1 - \lambda_2)^2}$ , so  $a = \frac{\lambda_2 + \sqrt{\frac{(6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2)^2}{(3\lambda_1 - \lambda_2)^2}}}{2}$ .

If  $\frac{6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} > 0$ , then  $a = \frac{3\lambda_1^2 - \lambda_2^2}{3\lambda_1 - \lambda_2}$ ,  $\alpha_2 = \frac{18\lambda_1^4 - 2\lambda_2^4 + 6\lambda_1 \lambda_2^3 - 18\lambda_1^3 \lambda_2}{(3\lambda_1 - \lambda_2)^2} = \frac{(3\lambda_1^2 - \lambda_2^2)(3\lambda_1^2 - 3\lambda_1 \lambda_2 + \lambda_2^2)}{(3\lambda_1 - \lambda_2)^2}$ , since  $3\lambda_1^2 - 3\lambda_1 \lambda_2 + \lambda_2^2 = 3(\lambda_1 - \frac{1}{2}\lambda_2)^2 + \frac{1}{4}\lambda_2^2 \neq 0$ , and  $\alpha_2 \neq 0$ , we have  $3\lambda_1^2 \neq \lambda_2^2$ , so  $a \neq 0$ .

If  $\frac{6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ , then  $a = \frac{3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2}$ ,  $\alpha_2 = \frac{18\lambda_1^4 - \lambda_2^4 + 6\lambda_1 \lambda_2^3 - 15\lambda_1^2 \lambda_2^2}{(3\lambda_1 - \lambda_2)^2} \neq 0$ , so  $18\lambda_1^4 - \lambda_2^4 + 6\lambda_1 \lambda_2^3 - 15\lambda_1^2 \lambda_2^2 \neq 0$ .

Hence, we have  $\lambda_2 \neq 3\lambda_1$ ,  $\lambda_2^2 \neq 3\lambda_1^2$ ,  $\frac{6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} > 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\alpha_1 = 0$ ,

$$\alpha = \frac{18\lambda_1^4 - 6\lambda_1 \lambda_2^3 + 24\lambda_1^2 \lambda_2^2 - 36\lambda_1^3 \lambda_2}{(3\lambda_1 - \lambda_2)^2}, \alpha_2 = \frac{18\lambda_1^4 - 2\lambda_2^4 + 6\lambda_1 \lambda_2^3 - 18\lambda_1^3 \lambda_2}{(3\lambda_1 - \lambda_2)^2}, \phi = ce^{\frac{3\lambda_1 - \lambda_2}{(3\lambda_1 - \lambda_2)p_1} t}.$$

or  $\lambda_2 \neq 3\lambda_1$ ,  $18\lambda_1^4 - \lambda_2^4 + 6\lambda_1 \lambda_2^3 - 15\lambda_1^2 \lambda_2^2 \neq 0$ ,  $\frac{6\lambda_1^2 - 3\lambda_1 \lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,

$$\alpha_1 = 0, \alpha = \frac{18\lambda_1^4 - 6\lambda_1 \lambda_2^3 + 24\lambda_1^2 \lambda_2^2 - 36\lambda_1^3 \lambda_2}{(3\lambda_1 - \lambda_2)^2}, \alpha_2 = \frac{18\lambda_1^4 - \lambda_2^4 + 6\lambda_1 \lambda_2^3 - 15\lambda_1^2 \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}, \phi = ce^{\frac{3\lambda_1 \lambda_2 - 3\lambda_1^2}{(3\lambda_1 - \lambda_2)p_1} t}.$$

Especially when  $\lambda_1 = \lambda_2 < 0$ , we have  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\alpha = \alpha_1 = 0$ ,  $\alpha_2 = 2\lambda_1^2$ ,  $\phi = c$ .

2'.  $p_1 \alpha_2 = 0$ . 1°.  $p_1 = 0$ . Then  $\zeta = 2p_2$ ,  $\eta = 2p_2^2$ ,  $\frac{\eta}{\zeta^2} = \frac{1}{2}$ . By the equation (52), we get

$$a = \frac{\alpha - \lambda_2^2 + 3\lambda_1 \lambda_2}{2\lambda_2}, \text{ on the other hand, } a = \frac{\lambda_2 + \sqrt{\lambda_2^2 - 2(\alpha + 3\lambda_1 \lambda_2 - 3\lambda_1^2)}}{2}, \text{ so we get}$$

$$\alpha^2 + (6\lambda_1 \lambda_2 - 2\lambda_2^2)\alpha + 3\lambda_1^2 \lambda_2^2 - 6\lambda_1^3 \lambda_2 + 3\lambda_1^4 = 0. \quad (53)$$

when  $\lambda_1 = \lambda_2$ , we get  $\alpha^2 + 4\lambda_1^2 \alpha = 0$ , then  $\alpha = 0$  or  $\alpha = -4\lambda_1^2$ . If  $\alpha = 0$ , we have

$a = \lambda_1 = \lambda_2 > 0$ , by the equation (44c), we get  $\alpha_2 = 0$ ,  $\phi = \psi^{\frac{\zeta}{\eta}} = c_0 e^{\frac{\lambda_1}{p_2} t}$ . If  $\alpha = -4\lambda_1^2$ , then  $\lambda_1 = \lambda_2 < 0$ ,  $a = -\lambda_1$ , then by the equation (44c), we get  $\lambda_1 = 0$ , which is a contradiction.

When  $\lambda_1 \neq \lambda_2$ ,  $\Delta = 8\lambda_2^2(3\lambda_1^2 - \lambda_2^2)$ , if  $\lambda_2^2 > 3\lambda_1^2$ , we have no solution; if  $\lambda_2^2 = 3\lambda_1^2$ , we have  $a = 0$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\phi = c$ ; if  $\lambda_2^2 < 3\lambda_1^2$ , we have  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 + \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$  or  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 - \sqrt{2\lambda_2^2(3\lambda_1^2 - \lambda_2^2)}$ . By the equation (44c), we have  $\alpha_2 = 0$  and

$$-2a^2 + (3\lambda_1 + \lambda_2)a = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \quad (54)$$

then we have  $\lambda_1 = \lambda_2 > 0$  or  $\lambda_1 = \lambda_2 < 0$ , but when  $\lambda_1 = \lambda_2 < 0$ , it is not satisfies the equation (44c), so we get  $\lambda_1 = \lambda_2 > 0$ ,  $p_1 = 0$ ,  $p_2 \neq 0$ ,  $\alpha = \alpha_1 = \alpha_2 = 0$ ,  $\phi = ce^{\frac{\lambda_1}{p_2} t}$ .

2°.  $\alpha_2 = 0$ . Since  $p_1 = 0$  we have discussed, so we assume that  $p_1 \neq 0$ . By the equation (52), we have  $a = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}$ ,  $\phi^\zeta = ce^{a\frac{\zeta^2}{\eta} t}$ , then by the equation (44b), we have  $\alpha(\alpha - \lambda_2^2 + 3\lambda_1\lambda_2) = 0$ . So when  $\lambda_2 = 3\lambda_1$ , we have  $\alpha = 0$ ; when  $\lambda_2 \neq 3\lambda_1$ , we have  $\alpha = 0$  or  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$ .

When  $\lambda_2 = 3\lambda_1$ ,  $\alpha = 0$ , then  $a = 0$ ,  $\eta = 0$ , which is a contradiction.

When  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2 \neq 0$ , then  $a = 0$  and  $(\lambda_2^2 - 3\lambda_1^2)\eta = 0$ .

If  $\lambda_2^2 \neq 3\lambda_1^2$ , then  $\eta = 0$ , which is a contradiction;

If  $\lambda_2^2 = 3\lambda_1^2$ , which satisfies  $\lambda_2 \neq 3\lambda_1$ , then  $\phi^\zeta = c$ , which satisfies equations (44a)–(44c). so we have  $\lambda_2^2 = 3\lambda_1^2$ ,  $p_1 \neq 0$ ,  $p_2 \neq -\frac{1}{2}p_1$ .  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\phi = c$ .

When  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = 0$ , then  $a = (3\lambda_1 - \lambda_2) \frac{\eta}{\zeta^2}$ , by the equation (44a), we get  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2} > 0$  and  $\lambda_2^2 < 3\lambda_1^2$ , then  $a = \frac{3\lambda_1^2 - \lambda_2^2}{3\lambda_1 - \lambda_2}$ . Since  $a > \frac{\lambda_2}{2}$ , we get  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} > 0$ , considering that  $\frac{p_1^2 + 2p_2^2}{(p_1 + 2p_2)^2} = \frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ , we have  $(3\lambda_1^2 - 3\lambda_1\lambda_2 + \lambda_2^2)p_1^2 - (6\lambda_1^2 - 2\lambda_2^2)p_1p_2 = (3\lambda_1^2 - 6\lambda_1\lambda_2 + 3\lambda_2^2)p_2^2$ , no matter  $\lambda_1 = \lambda_2$  or  $\lambda_1 \neq \lambda_2$ , we can get  $p_2 \neq 0$ .

So we get  $\lambda_2 \neq 3\lambda_1$ ,  $\lambda_2^2 < 3\lambda_1^2$ ,  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} > 0$ ,  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\alpha = \alpha_1 = \alpha_2 = 0$ ,

$$\phi = ce^{\frac{3\lambda_1^2 - \lambda_2^2}{3\lambda_1 - \lambda_2} \frac{\zeta}{\eta} t}.$$

3)  $c_1 \neq 0$ ,  $c_2 \neq 0$ ,  $b \neq 0$ .

1'.  $p_2 \neq 0$ . Then  $e^{at}$ ,  $e^{bt}$  and  $(c_1 e^{at} + c_2 e^{bt})^{1-2p_2} \frac{\zeta}{\eta}$  are linearly independent, by the equation (46), we have  $\left[a - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}\right] c_1 = 0$ ,  $\left[b - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}\right] c_2 = 0$ .

Considering that  $c_1 \neq 0$ ,  $c_2 \neq 0$ , we have  $a = b = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}$ , which is a contradiction.

2'.  $p_2 = 0$ . Then by the equation (46), we have



$$a - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} - \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} = 0, \quad b - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} - \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} = 0.$$

Then  $a = b$ , which is a contradiction.

4)  $c_1 \neq 0, c_2 \neq 0, b = 0$ . Then  $a = \lambda_2 \neq 0$  and

$$ac_1e^{at} = \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} (c_1e^{at} + c_2)^{1-2p_2\frac{\zeta}{\eta}} + \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} (c_1e^{at} + c_2). \quad (55)$$

1'.  $1 - 2p_2\frac{\zeta}{\eta} \neq 0$ .

If  $p_2 \neq 0$ , then  $e^{at}$  and  $(c_1e^{at} + c_2)^{1-2p_2\frac{\zeta}{\eta}}$  are linearly independent, by the equation (55), we have  $a = 0$ , this is a contradiction with  $a = \lambda_2 \neq 0$ .

If  $p_2 = 0$ , using the same method we can get  $a = 0$ , this is a contradiction with  $a = \lambda_2 \neq 0$ .

2'.  $1 - 2p_2\frac{\zeta}{\eta} = 0$ .

Then  $p_2 \neq 0, \eta = 2p_2\zeta$  and the equation (55) becomes

$$\left( \lambda_2 - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} \right) c_1e^{at} - \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} - \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2} = 0. \quad (56)$$

Then

$$\lambda_2 = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2} \frac{\eta}{\zeta^2}. \quad (57)$$

Since  $b = 0$  which means  $\lambda_2 = \sqrt{\lambda_2^2 - 4(\alpha + 3\lambda_1\lambda_2 - 3\lambda_1^2)} \frac{\eta}{\zeta^2}$ , we get  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2$ , then using the equation (57), we have  $\lambda_2 = \frac{3\lambda_1^2 - \lambda_2^2}{\lambda_2} \frac{\eta}{\zeta^2}$  and  $\lambda_2^2 \neq 3\lambda_1^2$ , considering that  $\eta = 2p_2\zeta$ , we get  $p_1 = \frac{6\lambda_1^2 - 4\lambda_2^2}{\lambda_2^2} p_2$ .

If  $3\lambda_1^2 = 2\lambda_2^2$ , then  $p_1 = 0$ , by  $\eta = 2p_2\zeta$  we have  $p_2 = 0$ , which is a contradiction.

If  $3\lambda_1^2 \neq 2\lambda_2^2$ , then  $p_1 \neq 0$ , and by  $\eta = 2p_2\zeta$  we have  $18\lambda_1^4 - 30\lambda_1^2\lambda_2^2 + 11\lambda_2^4 = 0$ , then  $\lambda_1^2 = \frac{5 \pm \sqrt{3}}{6} \lambda_2^2$ , which satisfies  $\lambda_2^2 \neq 3\lambda_1^2$  and  $3\lambda_1^2 \neq 2\lambda_2^2$ . Then  $\alpha_2 = \frac{(p_2 - p_1)(\lambda_2^2 - 3\lambda_1\lambda_2)}{p_1}$ ,  $p_1 = (1 \pm \sqrt{3})p_2 \neq 0$ ,  $\zeta = (3 \pm \sqrt{3})p_2$ ,  $\eta = (6 \pm 2\sqrt{3})p_2^2$ ,  $\phi = ce^{\frac{\lambda_2}{2p_1}t}$ .

So we get  $\lambda_1^2 = \frac{5 \pm \sqrt{3}}{6} \lambda_2^2, p_1 = (1 \pm \sqrt{3})p_2 \neq 0, \alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2, \alpha_1 = 0,$   
 $\alpha_2 = \frac{(p_2 - p_1)(\lambda_2^2 - 3\lambda_1\lambda_2)}{p_1}, \phi = ce^{\frac{\lambda_2}{2p_1}t}.$   
 $\alpha = \frac{\lambda_2^2\zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1\lambda_2, \psi = c_1e^{\frac{\lambda_2}{2}t} + c_2te^{\frac{\lambda_2}{2}t}.$  By the equation (45), we have

$$\left[ \frac{\lambda_2}{2} c_1 + c_2 - a_0 c_1 + \left( \frac{\lambda_2}{2} c_2 - a_0 c_2 \right) t \right] e^{\frac{\lambda_2}{2}t} = \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)} \frac{\eta}{\zeta^2} (c_1 + c_2 t)^{1-2p_2\frac{\zeta}{\eta}} (e^{\frac{\lambda_2}{2}t})^{1-2p_2\frac{\zeta}{\eta}}, \quad (58)$$

where  $a_0 = \frac{\lambda_2}{4} + \frac{3\lambda_1^2 - \lambda_2^2}{\lambda_2^2} \frac{\eta}{\zeta^2}$ .

1)  $c_2 \neq 0$ . Then by the equation (58), we have  $p_2 = 0$ , by the equation (44c), we

get  $\alpha_2 + \lambda_2 \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2$ , then  $\phi^\zeta = c_0 e^{\frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2 - \alpha_2}{\lambda_2}t}$ ; on the other hand,  $\phi^\zeta = \psi^{\frac{\zeta^2}{\eta}} = \left(c_1 e^{\frac{\lambda_2}{2}t} + c_2 t e^{\frac{\lambda_2}{2}t}\right)^{\frac{\zeta^2}{\eta}}$ , then  $c_0 e^{\frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2 - \alpha_2}{\lambda_2}t} = \left(c_1 e^{\frac{\lambda_2}{2}t} + c_2 t e^{\frac{\lambda_2}{2}t}\right)^{\frac{\zeta^2}{\eta}}$ , this is a contradiction with  $c_2 \neq 0$ .

2)  $c_2 = 0$ . Then the equation (58) becomes

$$\left(\frac{\lambda_2}{2}c_1 - a_0c_1\right)e^{\frac{\lambda_2}{2}t} = \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)}\frac{\eta}{\zeta^2}c_1^{1-2p_2\frac{\zeta}{\eta}}\left(e^{\frac{\lambda_2}{2}t}\right)^{1-2p_2\frac{\zeta}{\eta}}. \quad (59)$$

1'.  $a_0 = \frac{\lambda_2}{2}$ . By the equation (59), we get  $p_1\alpha_2 = 0$ . Considering that  $a_0 = \frac{\lambda_2}{4} + \frac{3\lambda_1^2 - \lambda_2^2}{\lambda_2^2}\frac{\eta}{\zeta^2}$ , we have  $\lambda_2^2 \neq 3\lambda_1^2$  and  $\frac{\lambda_2^2}{4}\frac{\zeta^2}{\eta} = 3\lambda_1^2 - \lambda_2^2$ , then  $\alpha = 6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2$ .

If  $p_1 = 0$ , then  $\zeta = 2p_2$ ,  $\eta = 2p_2^2$ ,  $\frac{\eta}{\zeta^2} = \frac{1}{2}$ . By  $\frac{\lambda_2^2}{4}\frac{\zeta^2}{\eta} = 3\lambda_1^2 - \lambda_2^2$ , we have  $\lambda_2^2 = 2\lambda_1^2$  and  $\alpha = 2\lambda_2^2 - 3\lambda_1\lambda_2$ . By the equation (44b), we get  $\phi^\zeta = c_0 e^{\lambda_2 t}$ , then by the equation (44c), we get  $\lambda_2 = \frac{3}{2}\lambda_1$ , this is a contradiction with  $\lambda_2^2 = 2\lambda_1^2$ .

If  $p_1 \neq 0$ ,  $\alpha_2 = 0$ , then  $\psi = c_1 e^{\frac{\lambda_2}{2}t}$ ,  $\phi^\zeta = c_0 e^{\frac{2(3\lambda_1^2 - \lambda_2^2)}{\lambda_2}t}$ , then by the equation (44b), we have  $\alpha = 6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ , which satisfies the equation (44c).  $\phi = c_0 e^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}$ .

So we get  $p_1 \neq 0$ ,  $\alpha = 6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\phi = c_0 e^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}$ .

2'.  $a_0 \neq \frac{\lambda_2}{2}$ . By the equation (59), we have  $p_2 = 0$ , then  $\zeta = p_1$ ,  $\eta = p_1^2$ ,  $\frac{\zeta^2}{\eta} = 1$ , so  $\alpha = \frac{\lambda_2^2}{4} + 3\lambda_1^2 - 3\lambda_1\lambda_2$ ,  $\phi^\zeta = c_1 e^{\frac{\lambda_2}{2}t}$ . By equation (44b), we have  $6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ , by the equation (44c), we have  $\alpha_2 = 3\lambda_1^2 - \frac{5}{4}\lambda_2^2$ .

So we get  $6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\alpha = \frac{\lambda_2^2}{4} + 3\lambda_1^2 - 3\lambda_1\lambda_2$ ,  $\alpha_2 = 3\lambda_1^2 - \frac{5}{4}\lambda_2^2$ ,  $\phi = c e^{\frac{\lambda_2}{2\zeta}t}$ .

$\alpha > \frac{\lambda_2^2\zeta^2}{4\eta} + 3\lambda_1^2 - 3\lambda_1\lambda_2$ ,  $\psi = c_1 e^{\frac{\lambda_2}{2}t}\cos(at) + c_2 e^{\frac{\lambda_2}{2}t}\sin(at)$ ,  $a = \frac{\sqrt{4(\alpha + 3\lambda_1\lambda_2 - 3\lambda_1^2)\frac{\eta}{\zeta^2} - \lambda_2^2}}{2}$ .

By the equation (45), we have

$$\begin{aligned} & \left(\frac{\lambda_2}{2}c_1 + ac_2\right)\cos(at) + \left(-ac_1 + \frac{\lambda_2}{2}c_2\right)\sin(at) \\ &= \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)}\frac{\eta}{\zeta^2}(c_1\cos(at) + c_2\sin(at))^{1-2p_2\frac{\zeta}{\eta}}e^{-\lambda_2 p_2 \frac{\zeta}{\eta}t} \\ &+ \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}(c_1\cos(at) + c_2\sin(at)). \end{aligned} \quad (60)$$

1)  $p_2 \neq 0$ . Then by the equation (60), we get  $p_1\alpha_2 = 0$  and

$$\frac{\lambda_2}{2}c_1 + ac_2 = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_1; \quad (61a)$$

$$-ac_1 + \frac{\lambda_2}{2}c_2 = \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_2. \quad (61b)$$

$(61a) \times c_2 - (61b) \times c_1$ , we get  $c_1^2 + c_2^2 = 0$ , this is a contradiction.

2)  $p_2 = 0$ . Then by the equation (60), we have

$$\frac{\lambda_2}{2}c_1 + ac_2 = \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)}\frac{\eta}{\zeta^2}c_1 + \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_1; \quad (62a)$$

$$-ac_1 + \frac{\lambda_2}{2}c_2 = \frac{p_1\alpha_2}{\lambda_2(p_2 - p_1)}\frac{\eta}{\zeta^2}c_2 + \frac{\alpha - \lambda_2^2 + 3\lambda_1\lambda_2}{\lambda_2}\frac{\eta}{\zeta^2}c_2. \quad (62b)$$

$(62a) \times c_2 - (62b) \times c_1$ , we get  $c_1^2 + c_2^2 = 0$ , this is a contradiction.

So we have no solution in .

According to above discussions, we get the following Theorem:

**Theorem 4.19.** *Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  be a generalized Kasner space-time,  $\dim F_1 = 1$ ,  $\dim F_2 = 2$  and  $P = \frac{\partial}{\partial t}$ . Then  $(M, \bar{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if one of the following conditions is satisfied:*

- (1)  $\lambda_2^2 = 3\lambda_1^2$ ,  $p_1 = p_2 = 0$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ ;
- (2)  $\lambda_2^2 = 3\lambda_1^2$ ,  $p_1^2 + p_2^2 \neq 0$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\phi = c$ ;
- (3)  $\lambda_2^2 \neq 3\lambda_1^2$ ,  $\lambda_2 \neq 3\lambda_1$ ,  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} \neq 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\alpha = \frac{18\lambda_1^4 - 6\lambda_1\lambda_2^3 + 24\lambda_1^2\lambda_2^2 - 36\lambda_1^3\lambda_2}{(3\lambda_1 - \lambda_2)^2}$ ,  
 $\alpha_1 = 0, \alpha_2 = \frac{18\lambda_1^4 - 2\lambda_2^4 + 6\lambda_1\lambda_2^3 - 18\lambda_1^3\lambda_2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\phi = ce^{\frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)p_1}t}$ ;
- (4)  $\lambda_2^2 < 3\lambda_1^2$ ,  $\lambda_2 \neq 3\lambda_1$ ,  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} \neq 0$ ,  $p_1 \neq 0$ ,  $p_2 \neq -\frac{1}{2}p_1$ ,  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\alpha =$   
 $\alpha_1 = \alpha_2 = 0, \phi = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta}t}$ ;
- (5)  $\lambda_2 \neq 3\lambda_1$ ,  $\frac{6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2}{3\lambda_1 - \lambda_2} < 0$ ,  $18\lambda_1^4 - \lambda_2^4 + 6\lambda_1\lambda_2^3 - 15\lambda_1^2\lambda_2^2 \neq 0$ ,  $p_1 \neq 0, p_2 = 0, \alpha_1 =$   
 $0, \alpha = \frac{18\lambda_1^4 - 6\lambda_1\lambda_2^3 + 24\lambda_1^2\lambda_2^2 - 36\lambda_1^3\lambda_2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\alpha_2 = \frac{18\lambda_1^4 - \lambda_2^4 + 6\lambda_1\lambda_2^3 - 15\lambda_1^2\lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\phi = ce^{\frac{3\lambda_1\lambda_2 - 3\lambda_1^2}{(3\lambda_1 - \lambda_2)p_1}t}$ ;
- (6)  $\lambda_1 = \lambda_2$ ,  $p_1 = 0$ ,  $p_2 \neq 0$ ,  $\alpha = \alpha_1 = \alpha_2 = 0$ ,  $\phi = c_0e^{\frac{\lambda_1}{p_2}t}$ ;
- (7)  $\lambda_1^2 = \frac{5 \pm \sqrt{3}}{6}\lambda_2^2$ ,  $p_1 = (1 \pm \sqrt{3})p_2 \neq 0$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2$ ,  $\alpha_1 = 0$ ,  $\alpha_2 = \frac{(p_2 - p_1)(\lambda_2^2 - 3\lambda_1\lambda_2)}{p_1}$ ,  
 $\phi = ce^{\frac{\lambda_2}{2p_1}t}$ ;
- (8)  $6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ ,  $p_1 \neq 0$ ,  $\alpha = 6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ ,  $\alpha_1 = \alpha_2 = 0$ ,  $\phi = ce^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}$ ;
- (9)  $6\lambda_1^2 - 3\lambda_1\lambda_2 - \lambda_2^2 = 0$ ,  $p_1 \neq 0$ ,  $p_2 = 0$ ,  $\frac{\zeta^2}{\eta} = 1$ ,  $\alpha = \frac{\lambda_2^2}{4} + 3\lambda_1^2 - 3\lambda_1\lambda_2$ ,  $\alpha_2 =$   
 $3\lambda_1^2 - \frac{5}{4}\lambda_2^2, \phi = ce^{\frac{\lambda_2}{2}\frac{\zeta}{\eta}t}$ .

#### 4.3.4 Type (II) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Proposition 4.17, then  $(F_2, \nabla^{F_2})$  has constant scalar curvature  $S^{F_2}$  and

$$\bar{S} = \frac{S^{F_2}}{\phi^{2p_2}} - 2\zeta\frac{\phi''}{\phi} - (\eta + \zeta^2 - 2\zeta)\frac{(\phi')^2}{\phi^2} + 3(\lambda_1 + \lambda_2)\zeta\frac{\phi'}{\phi} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2). \quad (63)$$

$\zeta = 0$ . (1)  $\eta = 0$ . then  $p_1 = p_2 = 0$  and  $\bar{S} = S^{F_2} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ ;  
(2)  $\eta \neq 0$ . then  $\bar{S} = \frac{S^{F_2}}{\phi^{2p_2}} - \eta \frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , which means

$$\eta \frac{(\phi')^2}{\phi^2} = \frac{S^{F_2}}{\phi^{2p_2}} - [\bar{S} - 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)]. \quad (64)$$

$\zeta \neq 0$ . Putting  $\phi = \psi^{\frac{2\zeta}{\eta+\zeta^2}}$ , we get

$$-\frac{4\zeta^2}{\eta+\zeta^2}\psi'' + \frac{6(\lambda_1+\lambda_2)\zeta^2}{\eta+\zeta^2}\psi' + (3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2 - \bar{S})\psi + S^{F_2}\psi^{1-\frac{4p_2\zeta}{\eta+\zeta^2}} = 0. \quad (65)$$

#### 4.3.5 Classification of Einstein Type (III) generalized Kasner space-times with a quarter-symmetric connection

Considering  $\dim F_1 = \dim F_2 = \dim F_3 = 1$ , by Remark 4, we get  $\alpha_i = 0$  and by Proposition 4.16, we have

$$\zeta \left( \lambda_2 \frac{\phi'}{\phi} - \frac{\phi''}{\phi} \right) - (\eta - \zeta) \frac{(\phi')^2}{\phi^2} + 3(\lambda_1^2 - \lambda_1\lambda_2) = \alpha; \quad (66a)$$

$$-p_1 \left[ \frac{\phi''}{\phi} + (\zeta - 1) \frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1) \frac{\phi'}{\phi} \right] + \lambda_2 \zeta \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \quad (66b)$$

$$-p_2 \left[ \frac{\phi''}{\phi} + (\zeta - 1) \frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1) \frac{\phi'}{\phi} \right] + \lambda_2 \zeta \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \quad (66c)$$

$$-p_3 \left[ \frac{\phi''}{\phi} + (\zeta - 1) \frac{(\phi')^2}{\phi^2} + (\lambda_2 - 3\lambda_1) \frac{\phi'}{\phi} \right] + \lambda_2 \zeta \frac{\phi'}{\phi} = \alpha - \lambda_2^2 + 3\lambda_1\lambda_2. \quad (66d)$$

$\zeta = \eta = 0$ . By the equation (66a), we have  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2$ , and by the equation (66b), we have  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$ , then we get  $\lambda_2^2 = 3\lambda_1^2$ .

So we obtain  $\lambda_2^2 = 3\lambda_1^2$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_i = 0$ ,  $\zeta = \eta = 0$ .

$\zeta = 0$ ,  $\eta \neq 0$ . (66b) + (66c) + (66d), we get  $\alpha = \lambda_2^2 - 3\lambda_1\lambda_2$ ; by the equation (66a), we have  $\frac{(\phi')^2}{\phi^2} = \frac{3\lambda_1^2 - \lambda_2^2}{\eta}$ .

1)  $3\lambda_1^2 - \lambda_2^2 < 0$ , we have no solution.

2)  $3\lambda_1^2 - \lambda_2^2 = 0$ , then  $\phi = c$ , which satisfies the equation (66a).

3)  $3\lambda_1^2 - \lambda_2^2 > 0$ , then  $\phi = c_0 e^{\pm \sqrt{\frac{3\lambda_1^2 - \lambda_2^2}{\eta}} t}$ , since  $\eta \neq 0$ , so at least one  $p_i \neq 0$ , we assume  $p_1 \neq 0$ , by the equation (66b), we get  $\lambda_2 = 3\lambda_1$ , but by  $3\lambda_1^2 - \lambda_2^2 > 0$ , we get  $\lambda_1^2 < 0$ , which is a contradiction.

So we have  $\lambda_2^2 = 3\lambda_1^2$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_i = 0$ ,  $\zeta = 0$ ,  $\eta \neq 0$ ,  $\phi = c$  in case .

$\zeta \neq 0$ , then  $\eta \neq 0$ . If  $p_1 = p_2 = p_3$ , we get type (I), if  $p_1 = p_2$  or  $p_2 = p_3$  or  $p_1 = p_3$ , we get type (II), so  $p_1 \neq p_2 \neq p_3$ . Let  $\phi = \psi^{\frac{\zeta}{\eta}}$ , then equations (66a) – (66d) becomes

$$\frac{\zeta^2}{\eta} \frac{\lambda_2 \psi' - \psi''}{\psi} = \alpha + 3\lambda_1 \lambda_2 - 3\lambda_2^2 \quad (67a)$$

$$\frac{p_1}{\zeta} \left[ -\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2) \frac{(\phi^\zeta)'}{\phi^\zeta} \right] + \lambda_2 \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 \quad (67b)$$

$$\frac{p_2}{\zeta} \left[ -\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2) \frac{(\phi^\zeta)'}{\phi^\zeta} \right] + \lambda_2 \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 \quad (67c)$$

$$\frac{p_3}{\zeta} \left[ -\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2) \frac{(\phi^\zeta)'}{\phi^\zeta} \right] + \lambda_2 \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 \quad (67d)$$

(67b)  $\times p_2$  – (67c)  $\times p_1$  and considering that  $p_1 \neq p_2$ , we get

$$\lambda_2 \frac{(\phi^\zeta)'}{\phi^\zeta} = \alpha - \lambda_2^2 + 3\lambda_1 \lambda_2 \quad (68a)$$

$$\frac{(\phi^\zeta)'}{\phi^\zeta} = \frac{\alpha}{\lambda_2} - \lambda_2 + 3\lambda_1 \quad (68b)$$

by equations (67b) and (68a), we get  $-\frac{(\phi^\zeta)''}{\phi^\zeta} + (3\lambda_1 - \lambda_2) \frac{(\phi^\zeta)'}{\phi^\zeta} = 0$ , then by the equation (68b), we have

$$\frac{(\phi^\zeta)''}{\phi^\zeta} = \left( \frac{3\lambda_1}{\lambda_2} - 1 \right) \alpha + 9\lambda_1^2 - 6\lambda_1 \lambda_2 + \lambda_2^2 \quad (69)$$

on the other hand, using the equation (68b), we get

$$\phi^\zeta = c_0 e^{\left( \frac{\alpha}{\lambda_2} - \lambda_2 + 3\lambda_1 \right) t} \quad (70)$$

by equations (69) and (70), we obtain  $\alpha^2 + (3\lambda_1 \lambda_2 - \lambda_2^2) \alpha = 0$ , so when  $\lambda_2 = 3\lambda_1$ , we have  $\alpha = 0$ ; when  $\lambda_2 \neq 3\lambda_1$ , we have  $\alpha = 0$  or  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2 \neq 0$ .

1)  $\lambda_2 = 3\lambda_1$ . Then  $\alpha = 0$ , by the equation (70), we get  $\phi^\zeta$  is a constant, then  $\psi$  is a constant, so by the equation (67a), we have  $\lambda_1^2 = 0$ , which is a contradiction with  $\lambda_1 \neq 0$ .

2)  $\lambda_2 \neq 3\lambda_1$ . 1'.  $\alpha = \lambda_2^2 - 3\lambda_1 \lambda_2 \neq 0$ . By the equation (70), we get  $\phi^\zeta$  is a constant, then  $\psi$  is a constant, and  $\phi = c$  is a constant. by the equation (67a), we have  $\alpha = 3\lambda_1^2 - 3\lambda_1 \lambda_2$  and  $\lambda_2^2 = 3\lambda_1^2$ .

So we get  $\lambda_2^2 = 3\lambda_1^2$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1 \lambda_2 = \lambda_2^2 - 3\lambda_1 \lambda_2$ ,  $\alpha_i = 0$ ,  $\zeta \neq 0$ ,  $\eta \neq 0$ ,  $\phi = c$ .

2'.  $\alpha = 0$ . By the equation (67a), we have

$$\psi'' - \lambda_2 \psi' + (3\lambda_1 \lambda_2 - 3\lambda_1^2) \frac{\eta}{\zeta^2} \psi = 0. \quad (71)$$

by the equation (70), we get  $\phi^\zeta = c_0 e^{(3\lambda_1 - \lambda_2)t}$ , then  $\phi = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta}t}$ ,  $\psi = c_1 e^{(3\lambda_1 - \lambda_2)\frac{\eta}{\zeta^2}t}$ ,  
by the equation (71), we get  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ .  
So we have  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = 0$ ,  $\alpha_i = 0$ ,  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\phi = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta}t}$ .

According to above discussions, we get the following Theorem:

**Theorem 4.20.** *Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  be a generalized Kasner space-time for  $p_i \neq p_j$  for  $i, j \in \{1, 2, 3\}$  and  $\dim F_1 = \dim F_2 = \dim F_3 = 1$ , and  $P = \frac{\partial}{\partial t}$ . Then  $(M, \overline{\nabla})$  is Einstein with the Einstein constant  $\alpha$  if and only if one of the following conditions is satisfied:*

- (1)  $\lambda_2^2 = 3\lambda_1^2$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_i = 0$ ,  $\zeta = \eta = 0$ ;
- (2)  $\lambda_2^2 = 3\lambda_1^2$ ,  $\alpha = 3\lambda_1^2 - 3\lambda_1\lambda_2 = \lambda_2^2 - 3\lambda_1\lambda_2$ ,  $\alpha_i = 0$ ,  $\eta \neq 0$ ,  $\phi = c$ ;
- (3)  $\lambda_2 \neq 3\lambda_1$ ,  $\alpha = 0$ ,  $\alpha_i = 0$ ,  $\frac{\eta}{\zeta^2} = \frac{3\lambda_1^2 - \lambda_2^2}{(3\lambda_1 - \lambda_2)^2}$ ,  $\phi = ce^{\frac{3\lambda_1 - \lambda_2}{\zeta}t}$ .

#### 4.3.6 Type (III) generalized Kasner space-times with a quarter-symmetric connection with constant scalar curvature

By Proposition 4.17, we get

$$\overline{S} = -2\zeta \frac{\phi''}{\phi} - (\eta + \zeta^2 - 2\zeta) \frac{(\phi')^2}{\phi^2} + 3(\lambda_1 + \lambda_2)\zeta \frac{\phi'}{\phi} + 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2). \quad (72)$$

$\zeta = \eta = 0$ . Then  $p_1 = p_2 = p_3 = 0$ , and  $\overline{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ .

$\zeta = 0$ ,  $\eta \neq 0$ . Then  $[(\ln \phi)']^2 = \frac{3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2) - \overline{S}}{\eta}$ , so we have:

1)  $\overline{S} > 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , we have no solution.

2)  $\overline{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , then  $\phi = c$ .

3)  $\overline{S} < 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , then  $\phi = c_0 e^{\pm \sqrt{\frac{3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2) - \overline{S}}{\eta}}t}$ .

$\zeta \neq 0$ , then  $\eta \neq 0$ . Putting  $\phi = \psi^{\frac{2\zeta}{\eta + \zeta^2}}$ , then

$$-\frac{4\zeta^2}{\eta + \zeta^2} \psi'' + \frac{6(\lambda_1 + \lambda_2)\zeta^2}{\eta + \zeta^2} \psi' + (3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2 - \overline{S})\psi = 0. \quad (73)$$

So we get

1)  $\overline{S} < \frac{9\zeta^2(\lambda_1 + \lambda_2)^2}{4(\eta + \zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$\begin{aligned} \psi &= c_1 e^{\frac{\frac{3}{2}(\lambda_1 + \lambda_2) + \sqrt{\frac{9(\lambda_1 + \lambda_2)^2}{4} - \frac{(\overline{S} - 3\lambda_1^2 - 3\lambda_2^2 + 12\lambda_1\lambda_2)(\eta + \zeta^2)}{\zeta^2}}}{2}t} \\ &+ c_2 e^{\frac{\frac{3}{2}(\lambda_1 + \lambda_2) - \sqrt{\frac{9(\lambda_1 + \lambda_2)^2}{4} - \frac{(\overline{S} - 3\lambda_1^2 - 3\lambda_2^2 + 12\lambda_1\lambda_2)(\eta + \zeta^2)}{\zeta^2}}}{2}t}; \end{aligned}$$

$$\begin{aligned}
2) \quad \bar{S} &= \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2, \quad \psi = c_1 e^{\frac{3(\lambda_1+\lambda_2)}{4}t} + c_2 t e^{\frac{3(\lambda_1+\lambda_2)}{4}t}, \\
3) \quad \bar{S} &> \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2,
\end{aligned}$$

$$\begin{aligned}
\psi &= c_1 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} \cos \frac{\sqrt{\frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2} - \frac{9(\lambda_1+\lambda_2)^2}{4}}}{2} t \\
&+ c_2 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} \sin \frac{\sqrt{\frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2} - \frac{9(\lambda_1+\lambda_2)^2}{4}}}{2} t
\end{aligned}$$

According to above discussions, we get the following Theorem:

**Theorem 4.21.** *Let  $M = I \times_{\phi^{p_1}} F_1 \times_{\phi^{p_2}} F_2 \cdots \times_{\phi^{p_m}} F_m$  be a generalized Kasner space-time and  $\dim F_1 = \dim F_2 = \dim F_3 = 1$ , and  $P = \frac{\partial}{\partial t}$ . Then  $(M, \bar{\nabla})$  has constant scalar curvature  $\bar{S}$  if and only if one of the following conditions is satisfied:*

- (1)  $\zeta = \eta = 0$ ,  $\bar{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ .
- (2)  $\zeta = 0$ ,  $\eta \neq 0$ , when  $\bar{S} > 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , we have no solution; when  $\bar{S} = 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , then  $\phi = c$ ; when  $\bar{S} < 3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2)$ , then  $\phi = c_0 e^{\pm \sqrt{\frac{3(\lambda_1^2 + \lambda_2^2 - 4\lambda_1\lambda_2) - \bar{S}}{\eta}} t}$ .
- (3)  $\zeta \neq 0$ ,
  - (a)  $\bar{S} < \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$\begin{aligned}
\psi &= c_1 e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2) + \sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4} - \frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}} t \\
&+ c_2 e^{\frac{\frac{3}{2}(\lambda_1+\lambda_2) - \sqrt{\frac{9(\lambda_1+\lambda_2)^2}{4} - \frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2}}}{2}} t,
\end{aligned}$$

- (b)  $\bar{S} = \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,  $\psi = c_1 e^{\frac{3(\lambda_1+\lambda_2)}{4}t} + c_2 t e^{\frac{3(\lambda_1+\lambda_2)}{4}t}$ ;
- (c)  $\bar{S} > \frac{9\zeta^2(\lambda_1+\lambda_2)^2}{4(\eta+\zeta^2)} + 3\lambda_1^2 + 3\lambda_2^2 - 12\lambda_1\lambda_2$ ,

$$\begin{aligned}
\psi &= c_1 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} \cos \frac{\sqrt{\frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2} - \frac{9(\lambda_1+\lambda_2)^2}{4}}}{2} t \\
&+ c_2 e^{\frac{3}{4}(\lambda_1+\lambda_2)t} \sin \frac{\sqrt{\frac{(\bar{S}-3\lambda_1^2-3\lambda_2^2+12\lambda_1\lambda_2)(\eta+\zeta^2)}{\zeta^2} - \frac{9(\lambda_1+\lambda_2)^2}{4}}}{2} t
\end{aligned}$$

**Remark 13.** *When  $\lambda_1 = \lambda_2 = 1$ , we get Propositions 32, 33 Theorems 35, 37, 36, in [12] by Propositions 4.16, 4.17, Theorems 4.19 – 4.21, respectively.*

**Acknowledgement.** This work was supported by NSFC No.11271062 and NCET-13-0721. We would like to thank the referee for his(her) careful reading and helpful comments.

## References

- [1] R. Bishop, B. O'Neill, Manifolds of negative curvature, Trans. Am. Math. Soc.145(1969) 1-49.
- [2] F. Dobarro, E. Dozo, Scalar curvature and warped products of Riemannian manifolds, Trans. Am. Math. Soc. 303(1987) 161-168.
- [3] P. Ehrlich, Y. Jung, S. Kim, Constant scalar curvatures on warped product manifolds, Tsukuba J. Math. 20(1996) No.1 239-265.
- [4] L. Alías, A. Romero, M. Sánchez, Spacelike hypersurfaces of constant mean curvature and Clabi-Bernstein type problems, Tohoku Math. J. 49(1997) 337-345.
- [5] M. Fernández-López, E. García-Río, D. Kupeli, B. Ünal, A curvature condition for a twisted product to be a warped product, Manu. math. 106(2001), 213-217.
- [6] H. A. Hayden, Subspace of a space with torsion, Proc. Lond. Math. Soc. 34(1932) 27-50.
- [7] K. Yano, On semi-symmetric metric connection, Rev. Roumaine Math. Pures Appl. 15(1970) 1579-1586.
- [8] Golab, S., On semi-symmetric and quarter-symmetric linear connections, Tensor N.S., 29(1975), 249-254.
- [9] F. Dobarro, B. Ünal, Curvature of multiply warped products, J. Geom. Phys. 55(2005) 75-106.
- [10] S. Sular and C. Özgür, Warped products with a semi-symmetric metric connection, Taiwanese J. Math. 15 (2011), no. 4, 1701C1719.
- [11] S.Sular and C.Özgür, Warped products with a semi-symmetric non-metric connection, Arab. J. Sci. Eng. 36 (2011), no. 3, 461C473.



- [12] Y. Wang, Multiply Warped products with a semi-symmetric metric connection, Abstract and Applied Analysis. Vol 2014, Article ID 742371, 12pages.
- [13] Y. Wang, Curvature of multiply warped products with an affine connection, Bull. Korean Math. Soc. 50 (2013), No. 5, pp. 1567C1586
- [14] M. Tripathi, A new connection in a Riemannian manifold, arXiv:1207.0199