

The definability of \mathbb{E} in self-iterable mice

Farmer Schlutzenberg¹

Abstract

Let M be a fine structural mouse and let $F \in M$ be such that $M \models "F$ is a total extender" and $(M \restriction \text{lh}(F), F)$ is a premouse. We show that it follows that $F \in \mathbb{E}^M$, where \mathbb{E}^M is the extender sequence of M . We also prove generalizations of this fact.

Let M be a premouse with no largest cardinal and let Σ be a sufficient iteration strategy for M . We prove that if M knows enough of $\Sigma \restriction M$ then \mathbb{E}^M is definable over the universe $[M]$ of M , so if also $[M] \models \text{ZFC}$ then $[M] \models "V = \text{HOD}"$. We show that this result applies in particular to $M = M_{\text{nt}} \restriction \lambda$, where M_{nt} is the least non-tame mouse and λ is any limit cardinal of M_{nt} .

We also show that there is no iterable bicephalus (N, E, F) for which E is type 2 and F is type 1 or 3. As a corollary, we deduce a uniqueness property for maximal $L[\mathbb{E}]$ constructions computed in iterable background universes.

Keywords: inner model, mouse, extender, sequence, self-iterable, bicephalus
2010 MSC: 03E45, 03E55

1. Introduction

Kunen [1] showed that if $V = L[U]$ where U is a normal measure, then U is the unique normal measure. Mitchell [4, 5] constructed inner models with sequences of measurables and proved analogous results regarding these. Steel [18, §8] and Schimmerling/Steel [9, §2] also proved related results, asserting roughly that if P is an iterate of K and E is an extender which is sufficiently certified and $(P \restriction \alpha, E)$ is a premouse, then E is on the extender sequence of P .

In this paper we will consider variants of these results pertaining to fine-structural mice M below a superstrong cardinal. Let $F \in M$ be such that $M \models "F$ is a total (and possibly wellfounded) extender". We are interested in questions such as:

1. Is F in the extender sequence \mathbb{E}_+^M of M ?
2. Is F the extender of an iteration map on M ?²³

Email address: farmer.schlutzenberg@gmail.com (Farmer Schlutzenberg)

¹Thanks to John Steel for discussions on the topic of this paper.

²John Steel suggested this question, under the added assumption that $V = [M]$.

³Where the iteration map is via a fine structural tree on M , as in [3].

We will show in Theorems 3.4–3.9, the central results of the paper, that under further reasonable (but significant) assumptions, the answer (to at least one of the above questions) is in the affirmative, and thus, \mathbb{E}^M is in some sense maximal with regard to extenders. The statements of these theorems are analogous to the results of Steel and Schimmerling/Steel mentioned above. The simplest case is the following theorem. (See §1.1 for a review of terminology and notation. Throughout the paper, we use the definition of *premouse* from [19, §2]. These models do not have extenders of superstrong type on their extender sequence. For the central results, this is the only anti-large cardinal hypothesis we require. For premice with superstrongs things are somewhat different, as discussed in 3.10.)

Theorem (3.4). *Let N be a $(0, \omega_1 + 1)$ -iterable premouse. Let $E \in N$ be such that $(N \parallel \text{lh}(E), E)$ is a premouse and E is total over N . Then $E \in \mathbb{E}^N$.*

Steel first proved the following version of 3.4; see [14] for the proof.

Theorem (Steel). *Let N be a $(0, \omega_1 + 1)$ -iterable premouse with no largest cardinal. Let $E \in N$ be such that $N \models “E$ is a total wellfounded extender such that $\nu = \nu(E)$ is regular and $N \parallel \nu = \text{Ult}(N, E) \parallel \nu”$. Then $E \in \mathbb{E}^N$.*

Note that Steel’s result requires slightly less coherence of E with respect to \mathbb{E}^N (through $\nu(E)$), than does 3.4 (through $\text{lh}(E)$). The “no largest cardinal” is more than enough for Steel’s result, but some such closure was used. On the other hand, in 3.4, $\nu(E)$ need not be a cardinal in N , and we do not explicitly demand that $\text{Ult}(N, E)$ be fully wellfounded; it is only required that $\text{Ult}(N, E)$ be wellfounded through $\text{lh}(E) + 1$ (this is part of the requirement that $(N \parallel \text{lh}(E), E)$ be a premouse).

Theorems 3.5–3.9 are modifications of 3.4 in which E fits on the sequence of an iterate P of N (that is, $(P \parallel \text{lh}(E), E)$ is a premouse). In 3.9, we also relax the requirement that $E \in N$, demanding instead that E be appropriately definable over N (and that the iteration tree leading from N to P is likewise definable). In order to state and prove this result precisely we must develop a coding of iteration trees over premice, which takes some work.

Because these results give criteria for an extender E to be in \mathbb{E}_+^N , or to be in \mathbb{E}_+^P for an iterate P of N , one can try to use them to show that a mouse can recognize its own extender sequence, and we achieve this in certain situations. Given a premouse M , we write $[M]$ for the universe of M . Recall that M_n is the canonical proper class inner model with n Woodin cardinals. Steel showed that for $n \leq \omega$, $K^{M_n} = M_n$,⁴ and therefore \mathbb{E}^{M_n} is definable over $[M_n]$ and $[M_n] \models “V = \text{HOD}”$. It seems to be unknown precisely how far Steel’s result

⁴For $n = 1$, Steel argued as follows. In M_1 , let μ be measurable and δ be Woodin. Let K_μ be Steel’s core model of height μ , as computed in $V_\delta^{M_1}$ (see [18]). Steel showed that $K_\mu = M_1 \restriction \mu$. This yields a definition of \mathbb{E}^{M_1} in $[M_1]$. For $n > 1$, he defines $K \restriction \delta_i$ inductively on i , where δ_i is Woodin and $\delta_{-1} = 0$, defining $K \restriction \delta_i = K(K \restriction \delta_{i-1})$, proceeding much as for M_1 in the interval (δ_{i-1}, δ_i) . In M_n , $K \restriction \delta_i = M_n \restriction \delta_i$ is above- δ_{i-1} , δ_i -iterable.

generalizes. We will show anyway in 4.13 that mice with a measurable limit of Woodins typically have a significant failure of self-iterability, one which seems to present a difficulty in generalizing Steel’s result to this level. Nonetheless, using results in §3, we will prove Theorem 4.9, a consequence of which is the following (we write M_{nt} for the minimal non-tame mouse; see §4):

Theorem. *Suppose M_{nt} exists and is $(\omega, \omega_1, \omega_1 + 1)$ -iterable. Let λ be a limit cardinal of M_{nt} and $N = M_{\text{nt}} \upharpoonright \lambda$. Then \mathbb{E}^N is definable over $\downarrow N$.*

Note that M_{nt} is well beyond a measurable limit of Woodins. Analogous methods work for many tame mice. The method depends on Theorems 4.3 and 4.7, a corollary of which is:

Theorem. *Let Z be a premouse satisfying $\text{ZFC} + “I \text{ am } (\omega, \text{OR}, \text{OR})\text{-iterable}”$. Then \mathbb{E}^Z is definable over $\downarrow Z$ and $\downarrow Z \models “V = \text{HOD}”$.*

The self-iterability hypothesis of this theorem fails if Z has a Woodin cardinal, but Theorems 4.3 and 4.7 themselves are versions suited to premice with Woodin cardinals, and which give the same conclusion. The method breaks down with non-tame mice, because of a more serious lack of self-iterability (see 4.12).

Finally, in §5, we adapt Mitchell and Steel’s “Uniqueness of the next extender” result [3, 9.2] to bicephali B of the form $(\downarrow B, E, F)$, with E type 2 and F type 1 or 3. This has positive implications regarding the uniqueness of maximal $L[\mathbb{E}]$ -constructions. (Although the question is natural, uniqueness in this particular case was not required for the arguments in [3]; and such bicephali were not considered there.)

The paper proceeds as follows. In §1.1 we review notation and terminology. In §2 we review Dodd fine structure for extenders (due to Steel) and prove various related facts. In §3 we state and prove the central results 3.4–3.9, making use of most of the work in §2. However, the proof of 3.4, for example, only depends on the results in §2 in the case that E is type 2, so one can basically skip §2 if one is only interested in the case that E is type 1 or 3, and refer to parts as needed. In §4 we consider, for premice N , the definability of \mathbb{E}^N over $\downarrow N$. This section uses only Theorems 3.6 and 3.8 from §3, and in particular, not Theorem 3.9, nor the material on coding of iteration trees in §3. Finally, §5 is on bicephali. This section can be read independently of §§3,4, and depends only a little on §2.

The pieces of §§2–4 due to the author are taken primarily from the author’s dissertation [14], with various refinements having been incorporated. There was a significant error in a draft of this paper, which is addressed in 4.10 and 4.13.

1.1. Conventions and Notation

General: For $\nu \in \text{OR}$, \mathcal{H}_ν denotes the set of sets hereditarily of size less than ν . For an extensional H , $\text{trcoll}(H)$ denotes the transitive collapse of H . Given any set X , $\text{trclos}(X)$ denotes the transitive closure of X . We use the lexicographic order on $[\text{OR}]^{<\omega}$: $a <_{\text{lex}} b$ iff $a \neq b$ and $\max(a \Delta b) \in b$. We

sometimes identify elements of $[\text{OR}]^{<\omega}$ with strictly descending sequences of ordinals. Let $a \in [\text{OR}]^{<\omega}$ with $a = \{a_0, \dots, a_{k-1}\}$ where $a_i > a_{i+1}$ for all $i + 1 < k$. We write $a \upharpoonright j$ for $\{a_0, \dots, a_{j-1}\}$. Let either σ, τ be sequences, or $\sigma, \tau \in [\text{OR}]^{<\omega}$. We write $\sigma \leq \tau$ iff there is k such that $\sigma = \tau \upharpoonright k$, and write $\sigma \triangleleft \tau$ iff $\sigma \leq \tau$ and $\sigma \neq \tau$.

Premice: Premice are as in [19], except that we officially consider a *premouse* to be an amenable structure $P = (\mathcal{J}_\alpha^\mathbb{E}, \mathbb{E}, \tilde{F})$, where \tilde{F} is the amenable coding of the active extender F of P , as described in [19, 2.9–2.10]. (Note that this coding is independent from squashing, and in this paragraph we consider no squashed premice.) We may blur the distinction between F and \tilde{F} ; whenever we write $(\mathcal{J}_\alpha^\mathbb{E}, \mathbb{E}, F)$ we literally mean P as above. We write $[P]$ for the universe $\mathcal{J}_\alpha^\mathbb{E}$ of P , $F^P = F(P)$ for F , $\mathbb{E}^P = \mathbb{E}(P)$ for the extender sequence \mathbb{E} of P , excluding F^P , and $\mathbb{E}_+^P = \mathbb{E}_+(P)$ is $\mathbb{E}^P \frown F^P$. If P has a largest cardinal it is denoted by $\text{lgcd}(P)$. We write $Q \leq P$ iff Q is an initial segment of P , that is, either $Q = P$ or $Q = (\mathcal{J}_\alpha^{\mathbb{E}^P}, \mathbb{E}^P \upharpoonright \alpha, \mathbb{E}_\alpha^P)$ for some limit ordinal $\alpha < \text{OR}^P$. We write $Q \triangleleft P$ iff $Q \leq P$ and $Q \neq P$. Let $\alpha \leq \text{OR}^P$ be a limit. Then $P|_\alpha$ denotes the $Q \leq P$ such that $\text{OR}^Q = \alpha$; and $P||_\alpha$ denotes $([Q], \mathbb{E}^Q, \emptyset)$ where $Q = P|_\alpha$. The notation $\nu(P) = \nu(F^P)$ is discussed below. *ISC* abbreviates “initial segment condition”. Such notation and terminology is used likewise a little more generally, for example with respect to segmented-premise (defined in 2.9).

Extenders: Our use of the term *extender* allows long, non-total measure spaces, and does not require full wellfoundedness of corresponding ultrapowers (clarified below).⁵ We will only consider those extenders which can be considered as being over (the universe of a) premouse. By restricting our attention to these we simplify some small considerations. Let us clarify. Let M be a premouse. Much as in [19, Definition 2.1 and following paragraph], we say that E is an *extender over* $[M]$ iff there is a structure N (N need not be transitive) and $j : [M] \rightarrow N$ which is Σ_0 -elementary (in the language of set theory) such that $\kappa = \text{cr}(j)$ exists and $\kappa + 1 \subseteq \text{wfp}(N)$, and there is $\delta \in (\kappa, \text{OR}^M]$ and $S \subseteq \text{wfp}(N)$ such that

$$E = \{(x, a) \mid x \in M \ \& \ \exists \xi < \delta [x \subseteq [\xi]^{|a|}] \ \& \ a \in [S]^{<\omega} \cap j(x)\}.$$

We write $\kappa_E = \text{cr}(E)$ for the critical point of E . The *space* of E , denoted

⁵The motivation for this is as follows. Most of the time we will deal with extenders E over some structure M , where E is likely not over V (it does not measure enough sets). We want to be able to use the term *extender* to refer to such E , instead of *partial (pre-)extender*. But for partial extenders, it is difficult to give a useful general notion of *completeness* or *wellfoundedness*. Thus, we prefer not to make any such demand in general (in our definition of *extender*). Although this usage of *extender* diverges from the conventional formal one (being V -total and countably complete), it also seems to be pretty common in informal usage, and is convenient for our present purposes. We also need to deal with long extenders in general.

$\text{space}(E)$, is δ . The *measure domain* of E is

$$\text{meas}(E) = \bigcup_{(\xi, n) \in \delta \times \omega} \mathcal{P}([\xi]^n) \cap M.$$

We say that E is *total* iff $\text{meas}(E) = \bigcup_{(\xi, n) \in \delta \times \omega} \mathcal{P}([\xi]^n)$. The *support* of E is $\text{spt}(E) = S$. We say that E is *short* iff $\text{space}(E) = \kappa_E + 1$. (One could require that $S \subseteq \bigcup_{\xi < \delta} j(\xi)$ and there be at most one $\xi < \delta$ such that $S \subseteq j(\xi)$, as only this part of E is relevant.) If E is short and $S = \lambda \in \text{OR}$, we write $E_{j, \lambda} = E$, and write $E_j = E_{j, j(\kappa_E)}$.

An *amenable transitive structure* is a structure $R = ([R], P_0, \dots, P_{k-1})$ with transitive universe $[R]$ and finitely many predicates P_i , such that each P_i is amenable to $[R]$. An amenable transitive structure $R = ([R], \mathbb{E}, \dots)$ is *pm-based*⁶ iff $([R], \mathbb{E}, \emptyset)$ is a (passive) premouse. Given F and a pm-based R , we say that F is an *extender over R* iff F is an extender over $[R]$. A *pm-based extender* is an extender over (the universe of) some premouse (equivalently, over some pm-based structure). So pm-based extenders can be non-short, and when short, we use the phrase *extender over* (a premouse) where [19] uses *pre-extender over*. Finally, *in this paper all extenders we consider are pm-based extenders*, so at this point we adopt the:

Convention 1.1. *Extender abbreviates pm-based extender.*

Thus, we always explicitly state assumptions on the totality of extenders, and on the wellfoundedness of ultrapowers by extenders.

Let R be a premouse and E an extender over R . Let M be a pm-based structure. Suppose that $\mathcal{H}_{\text{space}(E)}^M = \mathcal{H}_{\text{space}(E)}^R$. Then E is an extender over M . In fact, $N = \text{Ult}_0(M, E)$ is formed as usual (including the predicates of N) and likewise the associated ultrapower embedding $j = i_E^M$, and $j : M \rightarrow N$ is cofinal and Σ_0 -elementary with respect to the language of M ; this is routine by Loś' Theorem. (But N might be illfounded.) We abbreviate i_E^M by i_E if M is understood. Ultrapowers are generally by default at degree 0 (as above), unless context dictates otherwise. If M is a premouse, then $i_E^{M, k}$ denotes the degree k ultrapower embedding $M \rightarrow \text{Ult}_k(M, E)$, if defined. Sometimes we might abbreviate this with i_E^M or i_E . Given $a \in [\text{spt}(E)]^{<\omega}$ and an $\mathbf{r}\Sigma_k^M$ function f , $[a, f]_E^{M, k}$ denotes the object represented by the pair (a, f) in $\text{Ult}_k(M, E)$; we may also write $[a, f]_E^M$ if we wish to suppress k . The notation $\text{Ult}_k(E, F)$, for F an extender, and the notation \circ_k , are introduced in 2.26.

Let E be an extender over M . Note that if $\text{spt}(E) \in \text{OR}$ then for each $\alpha \in \text{spt}(E)$ we have $\alpha = [\{\alpha\}, f]_E^M$ where $f = \text{id}$ has sufficiently large domain. For $\beta \leq \alpha \in \text{spt}(E)$, recall that β is a *generator* of E iff $\beta \neq [a, f]_E^M$ for all f and $a \in [\beta]^{<\omega}$. The *natural length* $\nu(E) = \nu_E$ of E is the maximum of $(\kappa_E^+)^M$ and the strict sup of generators of E . For $X \subseteq \text{spt}(E)$, $E \restriction X$ is the sub-extender of E with support X and domain the least $\delta \leq \text{space}(E)$ such that

⁶ *pm* abbreviates *premouse*.

either $\delta = \text{space}(E)$ or $i_E(\delta) \geq \sup(X)$. (Thus, $E \restriction \text{spt}(E)$ is equivalent to E in terms of the ultrapowers it produces, but might have smaller domain.) Given an active premouse or related structure P , $\text{lh}(F^P)$ denotes the length of F^P , that is, OR^P . We sometimes write $\text{lh}(F)$ to denote $\text{lh}(F^P)$, when P is as above and $F = F^P \restriction (\tau \cup t)$ and F^P is generated by $\tau \cup t$ where $\tau \in \text{OR}$ and $t \in [\text{OR}]^{<\omega}$ (see 2.1).

Fine structure: For definability over premice, we basically follow [19], using the $\text{r}\Sigma_n$ hierarchy and n^{th} core $\mathfrak{C}_n(P)$ basically as there. As in [19], $\mathfrak{C}_0(P)$ denotes the squash P^{sq} of P if P is type 3, and otherwise just denotes P (as mentioned earlier, P is by definition amenable). If P is not type 3, we also define $P^{\text{sq}} = P$, so in all cases, $\mathfrak{C}_0(P) = P^{\text{sq}}$. Also in general, $\mathfrak{C}_0(P)^{\text{unsq}}$ denotes P .

Let $n < \omega$, let P be an n -sound premouse with $\omega < \rho_n^P$, and let $X \subseteq \mathfrak{C}_0(P)$. Let H be (i) the set of points in $\mathfrak{C}_0(P)$ definable over $\mathfrak{C}_0(P)$ with a generalized $\text{r}\Sigma_{n+1}$ term from parameters in $X \cup \{u_n^P\}$ (cf. [3, 2.8.1]). By [3, Proof of Lemma 2.10], H coincides with (ii) the set of points $y \in \mathfrak{C}_0(P)$ such that for some $\text{r}\Sigma_{n+1}$ formula φ and $\vec{x} \in (X \cup \{u_n^P\})^{<\omega}$, y is the unique $y' \in \mathfrak{C}_0(P)$ such that $\mathfrak{C}_0(P) \models \varphi(\vec{x}, y')$ (this uses the n -soundness of P and that \vec{x} can include u_n^P). We write $\text{Hull}_{n+1}^P(X)$ for the structure

$$(H, H \cap \mathbb{E}^{\mathfrak{C}_0(P)}, H \cap F^{\mathfrak{C}_0(P)}),$$

and write $\text{cHull}_{n+1}^P(X)$ for its transitive collapse. We may occasionally identify a type 3 premouse with its squash, so in particular, if P is type 3, then we might write $\text{cHull}_{n+1}^P(X)$ where we really mean $\text{cHull}_{n+1}^P(X)^{\text{unsq}}$. We define $\text{Hull}_\omega^P(X)$ and $\text{cHull}_\omega^P(X)$ similarly. For $\delta \in [\rho_{n+1}^P, \text{OR}^P]$, the δ -core of P is $\text{cHull}_{n+1}^P(\delta \cup \{p_{n+1}^P\})$. Also, $\text{Th}_{n+1}^P(X)$ and $\text{pTh}_{n+1}^P(X)$ respectively denote the generalized and pure $\text{r}\Sigma_{k+1}$ theories of $\mathfrak{C}_0(P)$ in parameters in $X \cup \{u_n^P\}$. (These theories include only generalized or pure $\text{r}\Sigma_{k+1}$ formulas, not negations thereof.) We will make use of the stratification of the pure theories given in [3, Proof of Lemma 2.10]. Because H above is determined by the corresponding pure theory, we also have a corresponding stratification of H ; see 2.23. Given $p \in [\rho_0^P]^{<\omega}$, an $(n+1)$ -solidity witness for (P, p) (or just for p) is a theory

$$\text{cHull}_{n+1}^P(\alpha \cup (p \setminus (\alpha + 1))),$$

where $\alpha \in p$. A generalized $(n+1)$ -solidity witness for (P, p) is analogous, but defined as in [20]. A (generalized) $(n+1)$ -solidity witness for P is that for (P, p_{n+1}^P) . And w_{n+1}^P denotes the set of all $(n+1)$ -solidity witnesses for P .

We take *weak k -embedding* $\pi : M \rightarrow N$ to be defined as in [13] (this definition is due to Steve Jackson). That is, the definition is as usual, except that we add the demand that there be a cofinal set $X \subseteq \rho_k^M$ such that π is $\text{r}\Sigma_{k+1}$ -elementary on $\text{Hull}_{k+1}^M(X)$. (This ensures that the proof of the Shift Lemma goes through as expected. We do not know whether one can prove the Shift Lemma for weak k -embeddings as defined in [3], when $1 \leq k < \omega$.)

Iteration trees: Structures such as premice, phalanxes of premice, bi-cephali, pseudo-premice, etc, we call *premouse-related*. All iteration trees (see

[19]) we consider in this paper will be fine-structural, in that they are much as in [3, §5], and based on premouse-related structures. We will not specify exactly what we mean by the general term *iteration tree*, but it suffices to consider k -maximal trees (see below) and stacks thereof. Let \mathcal{T} be an iteration tree. We write $M_{\alpha+1}^{*\mathcal{T}}$ for the model to which $E_\alpha^\mathcal{T}$ applies after any drop in model, and $i_{\alpha+1,\beta}^{*\mathcal{T}} : M_{\alpha+1}^{*\mathcal{T}} \rightarrow M_\beta^\mathcal{T}$ for the canonical embedding, if it exists. We write $\kappa_\alpha^\mathcal{T} = \text{cr}(E_\alpha^\mathcal{T})$, $\nu_\alpha^\mathcal{T} = \nu(E_\alpha^\mathcal{T})$ and $\text{lh}_\alpha^\mathcal{T} = \text{lh}(E_\alpha^\mathcal{T})$. If $\text{lh}(\mathcal{T}) = \theta + 1$, then $M_\infty^\mathcal{T} = M_\theta^\mathcal{T}$, $b^\mathcal{T} = [0, \theta]_\mathcal{T}$, and if there is no drop along $b^\mathcal{T}$ then $i^\mathcal{T} = i_{0,\theta}^\mathcal{T}$. We say \mathcal{T} is *above* ρ iff $\rho \leq \text{cr}(E_\alpha^\mathcal{T})$ for each $\alpha + 1 < \text{lh}(\mathcal{T})$.

For $k \leq \omega$, the notion k -maximal iteration tree \mathcal{T} (on a k -sound premouse) is defined in [19, Definition 3.4], and equivalently in [3, Definition 6.1.2]. The main points are that for all $\alpha + 1 < \text{lh}(\mathcal{T})$, with $\kappa = \kappa_\alpha^\mathcal{T}$, we have: (i) $\text{lh}(E_\beta^\mathcal{T}) < \text{lh}(E_\alpha^\mathcal{T})$ for all $\beta < \alpha$; (ii) $\text{pred}^\mathcal{T}(\alpha + 1)$ is the least β such that $\kappa < \nu_\beta^\mathcal{T}$; (iii) $M_{\alpha+1}^{*\mathcal{T}}$ is the largest $N \trianglelefteq M_\beta^\mathcal{T}$ such that $E_\alpha^\mathcal{T}$ measures $\mathcal{P}(\kappa) \cap N$; and (iv) $\deg^\mathcal{T}(\alpha + 1)$ is the largest $n \leq \omega$ such that $\kappa_\alpha^\mathcal{T} < \rho_n(M_{\alpha+1}^{*\mathcal{T}})$ and either $[0, \alpha + 1]_\mathcal{T}$ drops or $n \leq k$. We will also extend, in an obvious manner, the term k -maximal iteration tree to trees on premouse-related structures. Any non-obvious details in relation to such will hopefully be clear in context. We often abbreviate k -maximal iteration tree by (k) -maximal (tree). Whenever we use the term *maximal* with regard to an iteration tree, it is in the sense of k -maximal (for some, or for the relevant, k). The iteration trees we consider will all be either maximal, or stacks thereof (but most will be maximal).

Let $k \leq \omega$, let M be a k -sound premouse, and let $\theta \leq \text{OR}$. The notions (k, θ) -iteration strategy for M and (k, θ) -iterability of M are defined in [19, Definition 3.9]. (In particular, such a strategy Σ yields \mathcal{T} -cofinal wellfounded branches exactly for k -maximal trees \mathcal{T} which are according to Σ and have limit length $< \theta$.) For (k, α, θ) -iteration strategy and (k, α, θ) -iterable see [19, Definition 4.4].⁸ Given $\rho \in \text{OR}$ and Σ , we say that Σ is an $((k, \theta)$ -, etc) iteration strategy for M above ρ iff Σ works as an $((k, \theta)$ -, etc) iteration strategy with respect to trees above ρ . We extend this language in the obvious manner to *iteration strategies* and *iterability* for premouse-related structures.

If \mathcal{T} is k -maximal then $\Phi(\mathcal{T})$ denotes the phalanx associated to \mathcal{T} (see [18]). Let M, N be premice and $m, n \leq \omega$ such that M is m -sound and N is n -sound. Let $\delta \leq \text{OR}^M$ and $\lambda \leq \text{OR}^N$ with $\delta \leq \lambda$, $M||\delta = N||\delta$ and δ is a cardinal of N . We write $\mathfrak{P} = ((M, m, \delta), (N, n, \lambda))$ for the phalanx on which maximal trees \mathcal{T} are formed with the usual conditions augmented by the following: (i) $M_{-1}^\mathcal{T} = M$ and $\deg^\mathcal{T}(-1) = m$ and $M_0^\mathcal{T} = N$ and $\deg^\mathcal{T}(0) = n$, (ii) $E_0^\mathcal{T}$ is the first extender of \mathcal{T} and $E_0^\mathcal{T} \in \mathbb{E}_+^N$ with $\lambda \leq \text{lh}(E_0^\mathcal{T})$, (iii) if $\text{cr}(E_\alpha^\mathcal{T}) < \delta$ then $\text{pred}^\mathcal{T}(\alpha + 1) = -1$. For the phalanx notation $\Phi(\mathcal{T}, \iota, E)$ see 2.37.

⁷In [3] this requirement is included in the definition of *iteration tree*, whereas in [19] it is a requirement in the game $\mathcal{G}_k(M, \theta)$. The book [18] seems to weaken this requirement in its use of k -maximal.

⁸The author claims that all assumptions of (k, α, θ) -iterability in the paper can actually be weakened and replaced with $(k, \alpha, \theta)^*$ -iterability (see [16, p. 1202]).

2. Dodd structure

The proof of Theorems 3.4 and 3.5–3.9 will involve the analysis of a comparison. In some cases we will need to deal with the possibility that some generators of extenders are moved by an iteration map resulting from the comparison. We will need to analyze how such generators are moved. This section gives us the tools for this analysis. Parts of this section are also needed in §5. Somewhat restricted versions⁹ of the above-mentioned results basically do not rely on the current section, except that they refer to Definition 2.37 and use a simple case of Lemma 2.38.

We begin by surveying some definitions and notation. These definitions are taken from, or are slight variants of definitions from, [3] and [7]. See §1.1 for background notation, etc.

Definition 2.1 (Generators; Dodd condensation). Let E be an extender¹⁰ over a premouse N . We say that E is *standard* iff $\text{spt}(E) \in \text{OR}$ and $\alpha = [\text{id}, \{\alpha\}]_E^N$ for each $\alpha < \text{spt}(E)$.¹¹ Suppose E is standard and let $\gamma = \text{spt}(E)$ and $\kappa = \kappa_E$.

Let $X \subseteq \gamma$ and $\alpha < \gamma$. Then α is *generated by X* (with respect to E) iff there are $f \in N$ and $b \in [X]^{<\omega}$ such that $\alpha = [b, f]_E^N$. We say that X *generates E* iff every element of γ is generated by X . We write $E \approx E \restriction X$ iff X generates E . We say that E is *finitely generated* iff there is some finite X generating E . For $t \in [\gamma]^{<\omega}$ and $\alpha < \gamma$ we say that α is a *t -generator* iff α is not generated by $\alpha \cup t$. We say E is *weakly amenable (to N)*¹² iff

$$\mathcal{P}(\kappa) \cap N = \mathcal{P}(\kappa) \cap \text{Ult}(N, E).$$

If E is weakly amenable then $(\kappa^+)^E$ denotes $(\kappa^+)^N = (\kappa^+)^{\text{Ult}(N, E)}$.

Suppose E is weakly amenable. We define the *Dodd parameter* $t_E = t(E)$ of E , and *Dodd projectum* $\tau_E = \tau(E)$ of E . Let (k, t_0, \dots, t_{k-1}) be lex-largest in $[\text{OR}]^{<\omega}$ (ordered as in §1.1) such that for each $i < k$,

$$t_i \text{ is the largest } \{t_0, \dots, t_{i-1}\}\text{-generator } \alpha \text{ of } E \text{ such that } \alpha \geq (\kappa^+)^E.$$

Here k exists as $t_{i+1} < t_i$ for $i+1 < k$ (maybe $k = 0$). Then $t_E = \{t_0, \dots, t_{k-1}\}$ and τ_E is the sup of $(\kappa^+)^E$ and all t_E -generators of E .¹³ We say E is *Dodd-sound* iff both

⁹In particular, when the extender E in 3.4 and 3.5–3.9 is type 1 or 3.

¹⁰Our use of the term *extender* is specified in Convention 1.1.

¹¹Note that this is independent of N , and if we consider a degree k ultrapower ($k \leq \omega$) instead of degree 0, we also get the same result; likewise for other notions defined in 2.1.

¹²This is equivalent to the requirement that for every $a \in [\gamma]^{<\omega}$ and $\xi < \text{space}(E)$ and $f \in N$ such that $f : \kappa \rightarrow \mathcal{P}([\xi]^{|\alpha|})$, we have $\{\gamma \mid f(\gamma) \in E_a\} \in N$. Thus, it agrees with the terminology of [20, §2.3] regarding extenders on structures. However, it differs from the condition described in [3, Definition 1.0.8] – we might, for example, have $\text{OR}^N < \text{spt}(E) \in \text{OR}$.

¹³Note that $\tau_E > (\kappa^+)^N$ iff τ_E is a limit of t_E -generators of E (by choice of k). Possibly $k = 0$ and $\tau_E = \nu(E)$; this happens for example if E is the active extender of a type 1 or 3 premouse.

- $E \restriction (t_i \cup t_E \restriction i) \in \text{Ult}(N, E)$ for all $i < k$, and
- if $\tau_E > (\kappa^+)^N$ then $E \restriction (\alpha \cup t_E) \in \text{Ult}(N, E)$ for all $\alpha < \tau_E$.

Now let F be any extender (possibly non-standard) over N . Let E be the (equivalent) standard extender, derived from i_F , such that

$$\text{spt}(E) = \sup_{\alpha \in \text{spt}(F)} ([\{\alpha\}, \text{id}]_F^N + 1)$$

and $\text{space}(E)$ is as small as possible (i.e. $E = E \restriction \text{spt}(E)$). We might apply the preceding terminology to F in two ways, depending on context: (a) applying it literally to E (for example, “ X generates α ” would apply to a set X iff $X \subseteq \text{spt}(E)$, even if $\text{spt}(E) \not\subseteq \text{spt}(F)$); (b) dealing directly with $X \subseteq \text{spt}(F)$, as with the notation “ $F \restriction X$ ” introduced in §1.1.¹⁴ The intended meaning will hopefully be clear in context.

Let $\sigma \in \text{OR}$ and $s, t \in [\text{OR} \setminus \sigma]^{<\omega}$ with $\text{lh}(s) = \text{lh}(t)$. Let E, F be extenders with $\text{meas}(E) = \text{meas}(F)$ and $\sigma, s, t \subseteq \text{spt}(E) \cap \text{spt}(F)$. We write $E \equiv_{\sigma, s, t} F$ iff $E \restriction (\sigma \cup s)$ and $F \restriction (\sigma \cup t)$ are isomorphic, via the mapping of support which is the identity below σ and sends s to t . That is,

$$(x, a \cup s) \in E \iff (x, a \cup t) \in F$$

for all $x \in \text{meas}(E)$ and $a \in [\sigma]^{<\omega}$. +

The first fact below is straightforward to prove; see [7, 3.1].

Fact 2.2. *Let E be a weakly amenable extender with $\text{spt}(E) \in \text{OR}$ and $\kappa = \kappa_E$. Then τ_E is the least $\tau \geq (\kappa^+)^E$ such that there is $t \in [\text{OR}]^{<\omega}$ with $\tau \cup t$ generating E . And t_E is the least $t \in [\text{OR}]^{<\omega}$ witnessing this property of τ_E . In particular, $E \approx E \restriction \tau_E \cup t_E$.*

If $\tau_E > (\kappa^+)^E$ then E is not finitely generated. If $\tau_E > (\kappa^+)^E$ and $\tau_E \in \text{wfp}(\text{Ult}(N, E))$ then τ_E is a cardinal of $\text{Ult}(N, E)$. If $\tau_E = (\kappa^+)^E$ then E is finitely generated, indeed, generated by $t_E \cup \{\kappa\}$.

Fact 2.3 (Steel, [7, 3.2], [12, 4.1]). *Let N be an active, 1-sound, $(0, \omega_1, \omega_1 + 1)$ -iterable premouse. Then F^N is Dodd-sound.¹⁵*

Remark 2.4. Let M be a Dodd-sound active premouse, $F = F^M$, $\mu = \kappa_F$, $t = t^M$, $\tau = \tau^M$, and suppose that $(\mu^+)^M < \tau$. Let $q \in [\text{OR}^M]^{<\omega}$ and $\gamma < \tau$. Then $F \restriction (q \cup \gamma)$ is in M . For F is generated by $t \cup \tau$, so we may pick $\xi \in [\gamma, \tau)$ such that q is generated by $t \cup \xi$. By the second clause of Dodd-soundness, $F \restriction (t \cup \xi) \in M$, but $F \restriction (q \cup \gamma)$ is easily computed from $F \restriction (t \cup \xi)$.

¹⁴That is, let $\pi : N \rightarrow T$ be Σ_0 -elementary, such that F is derived from π . Let $U = \text{Ult}_0(N, F)$ and $k : U \rightarrow T$ the factor map. So $\text{rg}(k)$ is isomorphic to U , but maybe $\alpha < k(\alpha) \in \text{spt}(F)$. In case (b) we deal literally with $X \subseteq \text{spt}(F)$, so might have $k(\alpha) \in X$, whereas in case (a) we deal with the collapsed version of F , and sets $X \subseteq \text{spt}(E)$, so might have $\alpha \in X$ (and might have $\alpha \subseteq X$).

¹⁵We remark that although the proof superficially uses AC, the fact nonetheless follows from ZF alone. This is because one can first pass to an inner model of choice containing the premouse and closed under the iteration strategy. Likewise for solidity, condensation etc.

Remark 2.5. For active premice N such that $\text{Ult}(N|(\kappa^+)^N, F)$ is wellfounded, where $F = F^N$ and $\kappa = \kappa_F$, the definition of Dodd-soundness is optimal, in that no larger fragments of F can possibly appear in $\text{Ult}(N, F)$ (but this seems to use the assumption that N is below superstrong). If $\tau_F > (\kappa^+)^F$ the optimality is clear. Suppose $\tau_F = (\kappa^+)^F$. It is well-known that $\{\kappa\}$ generates $(\kappa^+)^F$. So if $t_F = \emptyset$ then $F \approx F \restriction \{\kappa\}$, so F is just a normal measure. So suppose $t_F \neq \emptyset$. Then we claim that

$$F \approx F \restriction t_F \notin \text{Ult}(N, F).$$

For $F \approx F \restriction t_F \cup \{\kappa\}$, so it suffices to see that κ is generated by $\{\gamma\}$ for every $\gamma \in [\kappa, \lambda)$ where $\lambda = i_F(\kappa)$. So fix $\gamma \in [\kappa, \lambda)$. By [19, Claim 2 of proof of 8.27] (the proof of which uses the non-superstrong assumption), we may fix $f \in N \cap {}^\kappa\kappa$ such that $\gamma \leq i_F(f)(\kappa) < \lambda$. Define $g : \kappa \rightarrow \kappa$ by letting $g(\alpha)$ be the least β such that $\beta = \alpha$ or $f(\beta) \geq \alpha$. Then $g \in N$ and $i_F(g)(\gamma) = \kappa$.

We now introduce a variant of the Dodd parameter and projectum, more analogous to the standard parameter and projectum.

Definition 2.6. Let \mathcal{D} be the class of pairs $(s, \sigma) \in [\text{OR}]^{<\omega} \times \text{OR}$ such that if $s \neq \emptyset$ then $\sigma \leq \min(s)$. We extend $<_{\text{lex}}$ to wellorder \mathcal{D} . Fix $(s_j, \sigma_j) \in \mathcal{D}$ for $j = 1, 2$. Let s_j^* be the strictly decreasing sequence with range s_j . Let $s'_j = s_j^* \hat{\ } \langle \sigma_j \rangle$, a monotone decreasing sequence. Let $(s_1, \sigma_1) <_{\text{lex}} (s_2, \sigma_2)$ iff either $s'_1 \sqsubseteq s'_2$ or there is $i \leq \min(\text{lh}(s_1^*), \text{lh}(s_2^*))$ such that $s_1^* \restriction i = s_2^* \restriction i$ and $s'_1(i) < s'_2(i)$. \dashv

Definition 2.7. Let M be a pm-based structure and $\pi : M \rightarrow N$ be Σ_0 -elementary (N need not be wellfounded). We say π is *cardinal preserving* iff

$$M \models \text{“}\alpha \text{ is an cardinal”} \iff N \models \text{“}\pi(\alpha) \text{ is a cardinal”}$$

for all $\alpha \in \text{OR}^M$. We say π is *Dodd-appropriate* iff π is cardinal preserving, $\mu = \text{cr}(\pi)$ exists and is inaccessible in M , $M|(\mu^+)^M = N|(\mu^+)^N$ and there is $\lambda \leq \pi(\mu)$ such that $\lambda \in \text{wfp}(N)$ and $E_{\pi, \lambda} \notin N$.

Assume π is Dodd-appropriate. Let $\mu = \text{cr}(\pi)$. The *Dodd-fragment parameter, projectum* (s_π, σ_π) of π is the $<_{\text{lex}}$ -least $(s, \sigma) \in \mathcal{D}$ with

$$\sigma \geq (\mu^+)^M \text{ and } E_\pi \restriction (\sigma \cup s) \notin N.$$

If F is a weakly amenable extender over M , $N = \text{Ult}(M, F)$ and $\pi = i_F$ is Dodd-appropriate, then (s_F, σ_F) denotes (s_π, σ_π) . For an active premouse R , (s_R, σ_R) denotes (s_{F^R}, σ_{F^R}) (note that i_{F^R} is Dodd-appropriate). \dashv

Remark 2.8. Let $\pi : M \rightarrow N$ be Dodd-appropriate with N wellfounded. Let $\mu = \text{cr}(\pi)$ and $\pi(\mu) \subseteq S \subseteq \text{OR}^N$ and F be the (possibly long) extender derived from π with support S ; so $E_\pi \subseteq F$. Let $U = \text{Ult}(M, F)$ and $k : U \rightarrow N$ be the factor map. Then $\text{cr}(k) > i_F(\mu) = \pi(\mu)$ and

$$U|_{i_F(\mu)} = N|_{\pi(\mu)} \text{ and } U|(\mu^+)^U = N|(\mu^+)^N = M|(\mu^+)^M, \quad (1)$$

and it easily follows that i_F is Dodd-appropriate and

$$(s_F, \sigma_F) = (s_\pi, \sigma_\pi) \leq_{\text{lex}} (\emptyset, \pi(\mu)).$$

Moreover, $(s_F, \sigma_F) \leq_{\text{lex}} (t_F, \tau_F)$: Suppose not. Let $G = F \upharpoonright (t_F \cup \tau_F)$. Then

$$(t_F, \tau_F) <_{\text{lex}} (s_F, \sigma_F) \leq_{\text{lex}} (\emptyset, \pi(\mu)). \quad (2)$$

Therefore G is short, and because $t_F \cup \tau_F$ generates F , therefore $G \notin U$. But $G \in U$ by line (2) and the definition of (s_F, σ_F) , contradiction. It easily follows that F is Dodd sound iff $(s_F, \sigma_F) = (t_F, \tau_F)$.

There is a characterization of s_π, σ_π analogous to the definition of the Dodd parameter and projectum. Let $s = \langle s_0, \dots, s_{l-1} \rangle \in \text{OR}^{<\omega}$ be the longest possible sequence such that for each $i < l$,

$$s_i \text{ is the largest } \alpha \geq (\mu^+)^M \text{ such that } E_\pi \upharpoonright ((s \upharpoonright i) \cup \alpha) \in N.$$

Note $s_{i+1} < s_i$. Then $s_\pi = s$ and σ_π is the sup of $(\mu^+)^M$ and all α such that $E_\pi \upharpoonright (s_\pi \cup \alpha) \in S$. We omit the proof.

In §5 we apply some of the results of §2 to pm-based structures which are not premice. For example, we will deal with structures of the form $\text{Ult}(P, H)$, where P is type 3 and $\text{cr}(H) = \nu_P$ (computing the ultrapower without squashing P). See [3, §9]. We now set up terminology in relation to this.

Definition 2.9. We say that M is a *segmented-premouse* (*seg-pm*) iff either (a) M is a passive premouse, or (b) there are N, F, \tilde{F} such that:

1. $M = (N, \tilde{F})$,
2. N is a passive premouse,
3. N has a largest cardinal δ ,
4. F is a short extender over N of length OR^N ,
5. $\nu(F) \leq \delta$,
6. $\text{OR}^N = (\delta^+)^{\text{Ult}(N, F)}$,
7. $N = \text{Ult}(N, F) \upharpoonright \text{OR}^N$, and
8. \tilde{F} is the amenable coding of F as in [19, 2.9–2.10], except that δ is used in place of $\nu(F)$; that is, \tilde{F} is the set of all tuples (γ, ξ, a, x) such that
 - $\kappa_F < \xi < (\kappa_F^+)^N$ and $\delta < \gamma < \text{OR}^N$, and
 - $F' \in N \upharpoonright \gamma$ and $(a, x) \in F'$ where $F' = F \cap (N \upharpoonright \xi \times [\delta]^{<\omega})$.

A segmented-premouse which is not a premouse is *proper*. ⊣

Remark 2.10. Premice are segmented-premise. However, active seg-premise P can fail the ISC, or can have $\nu(F^P) < \text{lgcd}(P)$, or both. Proper seg-premise can arise, for example, from taking ultrapowers of type 3 premise at the unsquashed level. There are two main kinds of examples of this. Let Q be a type 3 premouse and $\nu = \nu(F^Q) = \text{lgcd}(Q)$. Let E be a short extender weakly amenable to Q and $P = \text{Ult}(Q, E)$ (formed at the unsquashed level); suppose P is wellfounded. Let $\mu = \text{cr}(E)$. The first example arises in the case that

$$\mu = \text{cof}^Q(\nu) < \nu.$$

For then as discussed in [3, §9] (or see 2.11), i_E^Q is discontinuous at ν at

$$\nu(F^P) = \sup i_E^Q \nu < i_E^Q(\nu) = \text{lgcd}(P).$$

But in this case P satisfies the ISC. The second kind of example results if $\nu = \mu$ (so ν is inaccessible in Q). In this case P fails the ISC, as (by 2.11)

$$\nu < \nu(F^P) = \nu(E),$$

and $F^Q \restriction \nu = F^P \restriction \nu$ but $F^Q \restriction \nu \notin P$.

Regarding fine structure for proper seg-premise P , we only deal with degree 0. That is, we only define $\rho_0 = \text{OR}^P$ and 0-soundness (declared trivially true), and ultrapowers of P are degree 0, formed without squashing. Moreover, whenever we form $U = \text{Ult}_0(P, E)$ for an active seg-premouse P , without squashing P , we define $\widetilde{F^U}$ as usual, that is,

$$\widetilde{F^U} = \bigcup_{x \in P} i_E^P(\widetilde{F^P} \cap x).$$

Lemma 2.11. *Let P be an active seg-pm, $F = F^P$, and H a (possibly long) extender over P , with $W = \text{Ult}_0(P, H)$ a seg-pm.¹⁶ Let $A_F \subseteq \text{lh}(F)$ generate F and $A_H \subseteq \text{spt}(H)$ generate H . Let $j = i_H^P$. For an extender E , given $t \in [\text{spt}(E)]^{<\omega}$, write $G_{E,t}$ for the set of t -generators of E , and $G_E = G_{E, \emptyset}$.*

Then:

- (i) $A_H \cup j^* A_F$ generates F^W . Therefore $G_H \cup j^* G_F$ generates F^W and if $\text{space}(H) \leq \text{cr}(F) + 1$ then $j^* A_F$ generates F^W .
- (ii) $j^* G_{F,t} = G_{F^W, j(t)} \cap \text{rg}(j)$ for each $t \in [\text{OR}^P]^{<\omega}$.
- (iii) If $\text{cr}(F) < \text{cr}(H)$ then $G_H \subseteq G_{F^W}$.

So if $\text{cr}(F) < \text{cr}(H)$ then $G_H \cup j^ G_F$ is a generating set of generators of F^W ; and if $\text{space}(H) \leq \text{cr}(F) + 1$ then $j^* G_F$ is a generating set of generators.¹⁷*

¹⁶Seg-premousehood requires just that W is wellfounded and i_H^P is continuous at $(\kappa_F^+)^P$.

¹⁷However, F^W can also have generators outside of $G_H \cup j^* G_F$. For example, suppose that P is a type 2 premouse, $\text{cr}(F) < \xi = \text{cr}(H) < \nu(F)$, W is wellfounded, and H is short, with G_H bounded in $j(\xi)$. So W is a premouse and $j(\xi)$ is a cardinal of W , so (by the ISC) is a limit of generators of F^W , whereas ξ is not a limit point of $G_H \cup j^* G_F$.

Proof. Part (i): The first sentence is proven like [3, 9.1]: Let $\alpha < \nu(F^W)$. Let

$$\psi : \text{Ult}_0(P, F) \rightarrow \text{Ult}_0(W, F^W)$$

be the map induced by j . Let $f \in P$ and $b \in [A_H]^{<\omega}$ with $\alpha = j(f)(b)$. Let $g \in P$ and $c \in [A_F]^{<\omega}$ with $f = i_F^P(g)(c)$. Then α is generated by $b \cup j(c)$ with respect to F^W . For $\psi \upharpoonright P = j$ and $\psi \circ i_F^P = i_{F^W}^W \circ j$, so

$$\alpha = j(f)(b) = \psi(f)(b) = \psi(i_F^P(g)(c))(b) = i_{F^W}^W(j(g))(j(c))(b).$$

The second sentence follows easily; note that if $\text{space}(H) \leq \text{cr}(F) + 1$ then $G_H \subseteq j(\text{cr}(F)) = \text{cr}(F^W)$.

Part (ii): Because being a t -generator is an $\text{r}\Pi_1$ property of t .

Part (iii): Suppose $\kappa = \text{cr}(F) < \text{cr}(H)$ and $\alpha \in G_H$ but $\alpha \notin G_{F^W}$. Fix $b \in [\alpha]^{<\omega}$ and $f : [\kappa]^{|\mathbf{b}|} \rightarrow \kappa$ with $f \in W$, such that

$$i_{F^W}^W(f)(b) = \alpha.$$

Then $f \in P$; in fact $(\kappa^+)^P = (\kappa^+)^W < \text{cr}(H)$ because W is a premouse.¹⁸ Let $\mu \in P$ be such that $j(\mu) > \alpha$. Let

$$g = i_{F^P}^P(f) \cap ([\mu]^{|\mathbf{b}|} \times \mu).$$

Then $g \in P$ and

$$j(g) = i_{F^W}^W(f) \cap ([j(\mu)]^{|\mathbf{b}|} \times j(\mu)),$$

so $j(g)(b) = \alpha$, contradicting the fact that $\alpha \in G_H$. \square

Lemma 2.12 below is essentially [2, 2.1.4]. It can be deduced from 2.11, using that $\tau_F \cup t_F$ generates F , and, by $\text{r}\Pi_1$ -elementarity, that j sends total fragments of F to total fragments of F^W .

Lemma 2.12. *Let P, F, H, W, j be as in 2.11, with $\text{space}(H) \leq \tau_F$. Assume that F is Dodd sound. Then F^W is Dodd sound. Moreover, $t_{F^W} = j(t_F)$ and $\tau_{F^W} = \sup(j^{\text{``}}\tau_F)$.*

Lemma 2.13. *Let P, F, H, W, j be as in 2.11. Assume $\sigma_F \leq \text{cr}(H)$ and*

$$\mathcal{P}(\sigma_F) \cap P = \mathcal{P}(\sigma_F) \cap W.$$

Then F^W is not Dodd sound. Moreover, $s_{F^W} = j(s_F)$ and $\sigma_{F^W} = \sigma_F$.

Proof. Let $\sigma = \sigma_F$ and $s = s_F$ and $s' = j(s)$. We have $(s_{F^W}, \sigma_{F^W}) \geq_{\text{lex}} (s', \sigma)$ as j maps fragments of F to fragments of F^W . As $\sigma \leq \text{cr}(H)$,

$$F \equiv_{\sigma, s, s'} F^W$$

¹⁸Not just a protomouse; see [2].

(cf. 2.1). So $F \upharpoonright (\sigma \cup s)$ and $F^W \upharpoonright (\sigma \cup s')$ are coded by identical sets in $\mathcal{P}(\sigma)$, and $\mathcal{P}(\sigma) \cap P = \mathcal{P}(\sigma) \cap W$, so

$$F^W \upharpoonright (\sigma \cup s') \notin W.$$

Therefore $(s_{F^W}, \sigma_{F^W}) = (s', \sigma)$. And as in the proof of 2.11(iii), $\text{cr}(H)$ is an s' -generator of F^W , so F^W is not Dodd sound. \square

The proof of 2.15 to follow is similar to that of [7, 4.4]. Moreover, 2.17 improves the conclusion of [7, 4.4] to $0 = 1$. Recall from [3] that the premouse language uses the constant symbols $\dot{\mu}, \dot{\nu}, \dot{\gamma}$, interpreted by $\mu^P = \text{cr}(F^P)$, $\nu^P = \nu(F^P)$ and $\gamma^P =$ the largest witness to the ISC for P , when P is type 2.

Definition 2.14. Let P be a premouse, $F = F^P$, $\mu = \mu^P$, $X \subseteq \text{OR}^P$. Then $\Lambda^P(X)$ denotes the set of all $x \in P$ such that $x = [a, f]_{F^P}^P$ for some $f \in P$ and $a \in [X]^{<\omega}$ (note we may take $f \in P[(\mu^+)^P]$). \dashv

Lemma 2.15. Let P be a type 2 premouse, $F = F^P$, $\mu = \mu^P$, $X \subseteq \text{OR}^P$ and $\Lambda = \Lambda^P(X)$. Suppose $\nu^P, \gamma^P \in \Lambda$. Then

$$\Lambda = \{x \in P \mid x \in \text{Hull}_1^P(X \cup (\mu^+)^P)\}.$$

Proof. Clearly

$$\mu \cup X \subseteq \Lambda \subseteq H =_{\text{def}} \text{Hull}_1^P(X \cup (\mu^+)^P).$$

Let us show $H \subseteq \Lambda$. We have $\mu \in \Lambda$ because $\nu^P \in \Lambda$ and by the argument in 2.5. So $(\mu^+)^P \cup X \subseteq \Lambda$, so it suffices to see that $\Lambda \preccurlyeq_{\text{r}\Sigma_1} P$, and for this we define a premouse \bar{P} and an $\text{r}\Sigma_1$ -elementary $\pi : \bar{P} \rightarrow P$ such that $\text{rg}(\pi) = \Lambda$. Recall here that $\text{r}\Sigma_1$ -elementarity is with respect to the language of premice, incorporating $\dot{\mu}, \dot{\nu}, \dot{\gamma}$.

Let $F^* = F \upharpoonright (\Lambda \cap \text{OR})$, $Q = P[(\mu^+)^P]$ and $\bar{R} = \text{Ult}(Q, F^*)$ and $R = \text{Ult}(Q, F)$. Let $\varrho : \bar{R} \rightarrow R$ be the natural factor map. Let $\varrho(\bar{\nu}) = \nu^P$ and $\xi = (\bar{\nu}^+)^{\bar{R}}$ and $\pi = \varrho \upharpoonright (\bar{R} \restriction \xi)$. So $(\mu^+)^P < \text{cr}(\pi)$ and $\text{rg}(\pi) = \Lambda$ and $\varrho(\xi) = \text{OR}^P$. Let \bar{F} be the (μ, ξ) -extender derived from $i_{F^*}^Q$; so $(X, a) \in \bar{F}$ iff $(X, \pi(a)) \in F^*$. Let $\bar{P} = (\bar{R} \restriction \xi, \bar{F})$, where \bar{F} codes \bar{F} as in [19, 2.9–2.10], but replacing the triple $(\alpha, E_\alpha, \nu(E_\alpha))$ of [19] with $(\xi, \bar{F}, \bar{\nu})$. So $\pi : \bar{P} \rightarrow P$. Let $\gamma^{\bar{P}} = \pi^{-1}(\gamma^P)$, $\nu^{\bar{P}} = \bar{\nu}$, $\mu^{\bar{P}} = \mu$. It suffices to see that π is $\text{r}\Sigma_1$ -elementary (so \bar{P} is a premouse).

By construction, $\pi(\dot{\gamma}^{\bar{P}}, \dot{\nu}^{\bar{P}}, \dot{\mu}^{\bar{P}}) = (\gamma^P, \nu^P, \mu^P)$.¹⁹ Now $\varrho \circ i_{\bar{F}}^Q = i_F^Q$. So for $A \in \mathcal{P}([\mu]^{<\omega}) \cap P$ and $\gamma < \xi$, we have

$$\pi(i_{\bar{F}}^Q(A) \cap [\gamma]^{<\omega}) = i_F^Q(A) \cap [\pi(\gamma)]^{<\omega}.$$

So $\pi \restriction \bar{F} \subseteq \widetilde{F^P}$. Since $\pi \upharpoonright (\mu^+)^P = \text{id}$, [19, 2.9] shows that π is cofinal in P . It follows that π is $\text{r}\Sigma_1$ -elementary. \square

¹⁹This is the main reason for incorporating the assumption that $\gamma^P, \nu^P \in \Lambda$. There are standard counterexamples when these assumptions fail; see the proof that iterable 1-sound mice are Dodd-sound.

Part (ii) of the following lemma is proved in almost the same way; part (i) is easy:

Lemma 2.16. Let P, F, μ be as in 2.15, $\xi \in \text{OR}^P$ and $q \in [\text{OR}^P]^{<\omega}$, with $(\mu^+)^P \leq \xi$. Then (i) if $\text{cHull}_1^P(q \cup \xi) \in P$ then $F \restriction (q \cup \xi) \in P$, and (ii) if $F \restriction (q \cup \xi) \in P$ and $\gamma^P, \nu^P \in \Lambda^P(q \cup \xi)$ then $\text{cHull}_1^P(q \cup \xi) \in P$.

Corollary 2.17. Let P be a type 2, Dodd sound premouse, and $\mu = \mu^P$. Then $\tau_P = \rho_1^P \cup (\mu^+)^P$.

Proof. Let $\gamma = \rho_1^P \cup (\mu^+)^P$. Easily $\tau_P \geq \gamma$, so suppose $\tau_P > \gamma$. Let $X = q \cup \gamma$, where $q \in [\text{OR}^P]^{<\omega}$ is such that $p_1^P, \nu^P, \gamma^P \in \Lambda^P(q)$. By 2.4, $F^P \restriction X \in P$, so by 2.16, $\text{cHull}_1^P(X) \in P$, so $\text{Th}_1^P(\rho_1^P \cup \{p_1^P\}) \in P$, contradiction. \square

Lemma 2.18. Let $k \geq 1$ and let P be a k -sound, type 2, Dodd sound premouse. Let H be a short extender, weakly amenable to P , with $\text{cr}(H) < \rho_k^P$ and $R = \text{Ult}_k(P, H)$ wellfounded. Let $j = i_H^{P,k}$.

Then R is Dodd sound. Moreover, $t_R = j(t_P)$.

If $\tau_P = ((\mu^P)^+)^P$ or $k > 1$ then $\tau_R = j(\tau_P)$.

If $k = 1$ then $\tau_R = \sup j^{\text{“}}\tau_P$.

Proof. ²⁰ If $k > 1$, elementarity considerations give the lemma. So suppose $k = 1$. Let $F = F^P$, $\mu = \mu^P$, $t = t_F$ and $\tau = \tau_F$. Now j maps fragments of F to fragments of F^R by elementarity.

Suppose $\tau = (\mu^+)^P$. Then F is generated by $t \cup \{\mu\}$, an $\text{r}\Pi_2$ condition, preserved by j , so R is Dodd sound with $t_R = j(t_P)$ and $\tau_R = (j(\mu)^+)^R = j(\tau_P)$. But j is continuous at $(\mu^+)^P$, as $(\mu^+)^P$ is $\text{r}\Sigma_1^P$ -regular, as $\mu < \rho_1^P$.

Now suppose $\tau > (\mu^+)^P$. It suffices to see that $j(t) \cup \sup j^{\text{“}}\tau$ generates F^R . Suppose not. Let $\kappa = \text{cr}(H)$. Let a, f be such that $\gamma = [a, f]_H^{P,1}$ is a $j(t)$ -generator for F^R with $\gamma \geq \sup j^{\text{“}}\tau$, where $f : [\kappa]^n \rightarrow P$ is given by a $\text{r}\Sigma_1^P(\{q\})$ term for some $q \in P$. Since the statement “ α is a $j(t)$ -generator” is $\text{r}\Pi_1$, $\text{rg}(f)$ includes a τ -cofinal set of t -generators. We have

$$\text{rg}(f) \subseteq J = \text{Hull}_1^P(\kappa \cup \{q\}).$$

Let $\lambda < \tau$ be large enough that

$$\mu, q, \nu^P, \gamma^P \in \Lambda = \Lambda_F^P(t \cup \lambda).$$

Then Λ has no t -generators above λ . But by 2.15, $J \subseteq \Lambda$, contradiction. \square

For the iteration trees \mathcal{U} we will encounter, $E_\alpha^\mathcal{U}$ will always be weakly amenable (over $M_{\alpha+1}^{*\mathcal{U}}$), but ostensibly may not be close to $M_{\alpha+1}^{*\mathcal{U}}$. We now show that weak amenability gives a little more fine structural preservation than was established in [3]. The argument is related to that for [3, 6.2(Claim 5)] (though there, the extenders were always close to the relevant models), and also related to some of the preceding lemmas.

²⁰If R is $(0, \omega_1, \omega_1 + 1)$ -iterable, then the Dodd soundness of R follows from 2.3. But we don't want to assume this much iterability here.

Definition 2.19. Let N be a k -sound premouse with $\omega < \rho_k^N$. The $(k+1)$ -solidity parameter, projectum of N , denoted $(z_{k+1}^N, \zeta_{k+1}^N)$, is the $<_{\text{lex}}$ -least pair $(z, \zeta) \in \mathcal{D}$ such that $\zeta \geq \omega$ and $\text{Th}_{k+1}^N(\zeta \cup \{z\}) \notin N$. (See 2.6.) \dashv

Remark 2.20. The (rather trivial) requirement that $\zeta \geq \omega$ is just made to simplify the comparison between ζ_{k+1} and ρ_{k+1} . By [3, 2.10], one can equivalently replace “Th” in 2.19 with “pTh”. The author does not know an example, but it seems that ζ_{k+1}^N might fail to be a cardinal of N . However, the following facts follow easily from the definition; let $(z, \zeta) = (z_{k+1}^N, \zeta_{k+1}^N)$ and $(p, \rho) = (p_{k+1}^N, \rho_{k+1}^N)$:

- There is a characterization of (z, ζ) like that of the Dodd-solidity ordinals in 2.8.
- $\rho \leq \zeta \leq \rho_k^N$ and $(z, \zeta) \leq_{\text{lex}} (p, \rho)$.
- N is $(k+1)$ -solid iff $p = z$ iff $\rho = \zeta$.
- N is $(k+1)$ -sound iff $N = \text{Hull}_{k+1}^N(\zeta \cup z)$.

(The last item follows from the others: Suppose $N = \text{Hull}_{k+1}^N(\zeta \cup z)$ but $\rho < \zeta$. Note $p = z \cup q$ for some $q \subseteq [\rho, \zeta]$ with $q \neq \emptyset$. But then $(p, \rho) <_{\text{lex}} (z, \zeta)$, a contradiction.)

Lemma 2.21. Let N be a k -sound premouse. Let E be a short extender weakly amenable to N with $\text{cr}(E) < \rho_k^N$. Let $U = \text{Ult}_k(N, E)$ and $j = i_E^{N, k}$. Suppose U is wellfounded. Then $z_{k+1}^U = j(z_{k+1}^N)$ and $\zeta_{k+1}^U = \sup j \zeta_{k+1}^N$.

Proof. Let $(z, \zeta) = (z_{k+1}^N, \zeta_{k+1}^N)$ and $z' = j(z)$ and $\zeta' = \sup j \zeta$.

First observe that $(z', \zeta') \leq_{\text{lex}} (z_{k+1}^U, \zeta_{k+1}^U)$: By the proof of [3, 4.6], if $(b, \alpha) \in \mathcal{D}$ and $\text{Th}_{k+1}^N(\alpha \cup \{b\}) \in N$ then

$$\text{Th}_{k+1}^U(j(\alpha) \cup \{j(b)\}) \in U.$$

It remains to see that

$$t' =_{\text{def}} \text{pTh}_{k+1}^U(\zeta' \cup \{z'\}) \notin U. \quad (3)$$

So suppose $t' \in U$. We show $t =_{\text{def}} \text{pTh}_{k+1}^N(\zeta \cup \{z\}) \in N$, a contradiction.

Let $\kappa = \text{cr}(E)$. If $\kappa \geq \zeta$ then $t \approx t'$ by $\text{r}\Sigma_{k+1}$ elementarity (exchange z with z'). So by weak amenability, $t \in N$. (The details in 2.13 are analogous.)

So $\kappa < \zeta$. Also, $\zeta < \rho_k^N$. For otherwise $\zeta = \rho_k^N$. But $\text{pTh}_{k+1}^N(\rho_k^N) \notin N$ (since pTh_{k+1}^N has access to parameters p_k^N, u_{k-1}^N), so $z = \emptyset$, and $\rho_k^U = \zeta'$, so

$$t' = \text{pTh}_{k+1}^U(\rho_k^U) \notin U,$$

contradiction.

Since $\zeta < \rho_k^N$ there is $f \in N$ and $b \in [\nu(E)]^{<\omega}$ such that $f : [\kappa]^{<\omega} \rightarrow N$ and $t' = [b, f]_E^{N, k}$. (We get $f \in N$ even if $\rho_k^N = (\zeta^+)^N$, because in this case there is no $\text{r}\Sigma_k^N$ singularization of ρ_k^N .) We will compute t from f , giving $t \in N$. Now

pTh_{k+1} is a set of $\text{r}\Sigma_{k+1}$ formulas. We may assume that for each $x \in [\kappa]^{<\omega}$, $f(x)$ is a set of $\text{r}\Sigma_{k+1}$ formulas in parameters in $\zeta \cup \{z, u_k^N\}$.

For $\alpha < \zeta$, let

$$t_\alpha^N = \text{pTh}_{k+1}^N(\alpha \cup \{z\})$$

and let $f_\alpha : [\kappa]^{|b|} \rightarrow N$ be defined

$$f_\alpha(x) = f(x) \upharpoonright (\alpha \cup \{z, u_k^N\}),$$

where $T \upharpoonright X$ is the restriction of a theory T to parameters in X . For $\alpha < \zeta'$, let

$$t_\alpha^U = \text{pTh}_{k+1}^U(\alpha \cup \{z'\}).$$

Case 1. For each $\gamma < \zeta$, $j(t_\gamma^N) = t_{j(\gamma)}^U$.

We leave this case to the reader; it is simpler than the next one.

Case 2. For some $\gamma < \zeta$, $j(t_\gamma^N) \neq t_{j(\gamma)}^U$.

Fix such a γ . Let $<^*$ be the prewellorder on $\text{pTh}_{k+1}^N(N)$ defined in [3, Proof of 2.10]. It follows that t_γ is cofinal in $<^*$, and that $\text{cof}^N(<^* \upharpoonright t_\gamma) = \kappa$. Let $h \in N$ be such that $h : \kappa \rightarrow t_\gamma$ and h is cofinal, increasing and continuous with respect to $<^*$. For $\alpha \in [\gamma, \zeta)$ let $g_\alpha : \kappa \rightarrow N$ be defined

$$g_\alpha(\beta) = t_\alpha^N \upharpoonright^{<^*} h(\beta)$$

where $T \upharpoonright^{<^*} \varphi$ is the restriction of a theory T (of the appropriate kind) to formulas ψ such that $\psi <^* \varphi$. Then $g_\alpha \in N$ (maybe $\langle g_\alpha \rangle_\alpha \notin N$) and

$$[g_\alpha, \{\kappa\}]_E^{N,k} = t_{j(\alpha)}^U.$$

We may assume that $\kappa = \min(b)$. Let $g'_\alpha : [\kappa]^{|b|} \rightarrow N$ be defined

$$g'_\alpha(x) = g_\alpha(\min(x)).$$

So there is $X_\alpha \in E_b$ such that $g'_\alpha \upharpoonright X_\alpha = f_\alpha \upharpoonright X_\alpha$.

Define the relation $R \subseteq \zeta \times ([\kappa]^{|b|})^2$ by

$$R(\alpha, x_1, x_2) \iff [\max(x_1) < \min(x_2) \text{ and } f_\alpha(x_1) \subseteq f_\alpha(x_2)].$$

So $R \in N$. Define the relation $\psi \subseteq \zeta \times [\kappa]^{|b|}$ by

$$\psi(\alpha, x_1) \iff R(\alpha, x_1, x_2) \text{ holds for } E_b\text{-almost all } x_2.$$

Fix $\alpha \in [\gamma, \zeta)$. Note that if $x_1, x_2 \in X_\alpha$ and $\max(x_1) < \min(x_2)$, then $R(\alpha, x_1, x_2)$. So if $x_1 \in X_\alpha$ then $\psi(\alpha, x_1)$. On the other hand, if $\psi(\alpha, x_1)$ then there is $x_2 \in X_\alpha$ such that $R(\alpha, x_1, x_2)$, and so $f_\alpha(x_1) \subseteq t_\alpha$. So let $S_\alpha = \{x_1 \mid \psi(\alpha, x_1)\}$. Then $S_\alpha \in N$ as $R \in N$ and by weak amenability, and

$$t_\alpha = \bigcup_{x_1 \in S_\alpha} f_\alpha(x_1).$$

Suppose $\text{cof}^N(\zeta) \leq \kappa$. Fix a cofinal function $g : \kappa \rightarrow \zeta$ with $g \in N$ and $\text{rg}(g) \subseteq [\gamma, \zeta)$. Again by weak amenability and as $R \in N$,

$$S' =_{\text{def}} \{(\beta, x) \mid \psi(g(\beta), x)\} \in N.$$

But then

$$t = \bigcup_{(\beta, x) \in S'} f_{g(\beta)}(x) \in N,$$

as required.

Now suppose instead that $\text{cof}^N(\zeta) > \kappa$. Let R_α be the α -section of R . For $\alpha_1, \alpha_2 \in [\gamma, \zeta)$ with $\alpha_1 < \alpha_2$, we have $R_{\alpha_2} \subseteq R_{\alpha_1}$. So $R_{\alpha_1} = R_{\alpha_2}$ for all sufficiently large $\alpha_1, \alpha_2 < \zeta$. Fix such an α_1 . Then $S =_{\text{def}} S_\alpha$ is independent of $\alpha \in [\alpha_1, \zeta)$, and $S \in N$. So let $S' = [\alpha_1, \zeta) \times S$. Then

$$t = \bigcup_{(\alpha, x) \in S'} f_\alpha(x) \in N,$$

completing the proof. $\square(2.21)$

Notation 2.22. Let N be a k -sound seg-pm. Let $\mathcal{F} = \langle F_\alpha \rangle_{\alpha < \lambda}$ be a sequence of weakly amenable short extenders. For $\xi \leq \lambda$, let $\langle N_\alpha \rangle_{\alpha \leq \xi}$ be the degree k iterated ultrapower based on N , using the extenders in \mathcal{F} , if it is defined. That is, $N_0 = N$,

$$N_{\alpha+1} = \text{Ult}_k(N_\alpha, F_\alpha),$$

and we take direct limits at limit α . If N_α is illfounded, or F_α is not an extender over N_α , or $\text{cr}(F_\alpha) \geq \rho_k^{P_\alpha}$, then for all $\beta > \alpha$, N_β is undefined. Say that \mathcal{F} is *k-pre-good for N* iff N_λ is defined (but is maybe illfounded), and *k-good* iff *k-pre-good* and wellfounded. Suppose \mathcal{F} is *k-pre-good for N* . We write

$$\text{Ult}_k(N, \mathcal{F}) = N_\lambda$$

and

$$i_{\mathcal{F}}^{N, k} : N \rightarrow \text{Ult}_k(N, \mathcal{F})$$

for the ultrapower embedding. If N is active, $\text{Ult}_k(F^N, \mathcal{F})$ and $\text{Ult}_k(\text{id}, \langle F^N \rangle \frown \mathcal{F})$ both denote $F(\text{Ult}_k(N, \mathcal{F}))$.

Following the (kind of) calculations used in [3, Proof of Lemma 2.10]:

Definition 2.23 (Minimal Skolem terms). Let φ be an $\text{r}\Sigma_{k+1}$ formula of $n+1$ free variables. The *minimal Skolem term associated to φ* is denoted $\text{m}\tau_\varphi$, and has n variables. Let R be a k -sound premouse with $\omega < \rho_k^R$. We define the partial function

$$\text{m}\tau_\varphi^R : \mathfrak{C}_0(R)^n \rightarrow \mathfrak{C}_0(R),$$

by induction on k , maintaining the following: Let $\Gamma = \text{r}\Sigma_{k+1}^R(\{p_k^R, u_{k-1}^R\})$. The graph of $\text{m}\tau_\varphi^R$ is Γ , and Γ is closed under substitution of minimal Skolem terms;

this is uniform jointly in φ and in k -sound premice R of the same type. Moreover, for $X \subseteq \mathfrak{C}_0(R)$,

$$\text{Hull}_{k+1}^R(X) = \{\text{m}\tau_\varphi^R(\vec{x}, u_k^R) \mid \varphi \text{ is } \text{r}\Sigma_{k+1} \text{ \& } \vec{x} \in X^{<\omega}\}.^{21}$$

If $k = 0$ then $\text{m}\tau_\varphi^R$ is just the usual $\text{r}\Sigma_1$ Skolem function associated to φ .

Suppose $k \geq 1$. Let $u = u_{k-1}^R$ and $p = p_k^R$. Let ψ be an $\text{r}\Sigma_1$ formula and $\varphi(v, w)$ be the $\text{r}\Sigma_{k+1}$ formula

$$\exists t \in T_k[\psi(t, v, w)].$$

Let T be a set of $\text{r}\Sigma_k$ formulas in parameters in $\alpha \cup \{u, p\}$, where $\alpha < \rho_k^N$. Let $\gamma < \alpha$, let $\vec{\beta} = (\beta_0, \beta_1, \beta_2, \beta_3) \in \alpha^4$, let $\vec{r} = (r_0, r_1, r_2, r_3)$ be minimal $\text{r}\Sigma_k$ Skolem terms. Write s_i for the term-in-parameters $r_i(\beta_i, u, p)$. Let $x \in \mathfrak{C}_0(R)$. Say T codes a witness to $\exists w\varphi(x, w)$ at $(\gamma, \vec{\beta}, \vec{r})$ iff

- $x = s_2^R$ (that is, $r_2^R(\beta_2, u, p)$ is defined and $= x$),
- T contains the formula “ $\exists t, q, x', w[t = s_0, q = s_1, x' = s_2, w = s_3, t \text{ is a set of } \text{r}\Sigma_k \text{ formulas in parameters in } \gamma \cup \{q\}, \text{ and } \psi(t, x', w)]$ ”,
- for each $\text{r}\Sigma_k$ formula ϱ and $\vec{\gamma} \in \gamma^{<\omega}$, T contains the formula

$$“\exists t, q[t = s_0, q = s_1 \text{ and } t \text{ contains the formula } “\varrho(\vec{\gamma}, q)”]”$$

iff T contains the formula “ $\exists q[q = s_1 \text{ and } \varrho(\vec{\gamma}, q)]$ ”.

Let $\varphi(v, w)$ be $\text{r}\Sigma_{k+1}$. Then $\varphi'(v, y)$ is the formula “ $\varphi(v, y)$ and letting $(\alpha, \gamma, \vec{\beta}, \vec{r})$ be lex-least such that $\text{Th}_k(\alpha \cup \{p_k\})$ codes a witness to $\exists w\varphi(v, w)$, then $y = r_3(\beta_3, u_{k-1}, p_k)$ ” (the formula is to be interpreted over k -sound premice S , with u_{k-1}, p_{k-1} interpreted, of course, as u_{k-1}^S, p_k^S). Note that:

- φ' is $\text{r}\Sigma_{k+1}(\{u_{k-1}, p_k\})$ (uniformly so),
- $\varphi \mapsto \varphi'$ is recursive (when we choose φ' naturally),
- $\mathfrak{C}_0(R) \models \exists w\varphi(x, w)$ iff $\mathfrak{C}_0(R) \models \exists w\varphi'(x, w)$ iff $\mathfrak{C}_0(R) \models \exists! w\varphi'(x, w)$.

We now define $\text{m}\tau_\varphi^R$, for $\varphi \text{ r}\Sigma_{k+1}$. Let $\vec{x} \in \mathfrak{C}_0(R)^n$. If $\mathfrak{C}_0(R) \models \neg \exists y\varphi(\vec{x}, y)$, then $\text{m}\tau_\varphi^R(\vec{x})$ is undefined. Otherwise

$$\text{m}\tau_\varphi^R(\vec{x}) = \text{the unique } w \in \mathfrak{C}_0(R) \text{ such that } \mathfrak{C}_0(R) \models \varphi'(\vec{x}, w). \quad \dashv$$

Corollary 2.24. *Let N be a k -sound premouse. Let $\mathcal{F} = \langle F_\alpha \rangle_{\alpha < \lambda}$ be a sequence of weakly amenable short extenders, k -good for N . Let $N_\alpha = \text{Ult}_k(N, \mathcal{F} \restriction \alpha)$. (So N_λ is well-defined and wellfounded.) Let $j = i_{\mathcal{F}}^{N, k}$. Then:*

²¹Recall here that $\text{Hull}_{k+1}^R(X)$ includes u_k^R by definition; likewise $\text{Th}_{k+1}^R(X)$ is a theory in parameters in $X \cup \{u_k^R\}$.

1. $z_{k+1}^{N_\lambda} = j(z_{k+1}^N)$ and $\zeta_{k+1}^{N_\lambda} = \sup j \ulcorner \zeta_{k+1}^N \urcorner$.
2. If N, N_λ are both $(k+1)$ -solid then $p_{k+1}^{N_\lambda} = j(p_{k+1}^N)$ and $\rho_{k+1}^{N_\lambda} = \sup j \ulcorner \rho_{k+1}^N \urcorner$.
3. If N is not $(k+1)$ -solid then neither is N_λ .
4. N_λ is $(k+1)$ -sound iff N is $(k+1)$ -sound and $\text{cr}(F_\alpha) < \rho_{k+1}^{N_\alpha}$ for all $\alpha < \lambda$.

Proof. Let $\rho = \rho_{k+1}^N$, $p = p_{k+1}^N$, $\zeta = \zeta_{k+1}^N$, $z = z_{k+1}^N$, $\zeta' = \sup j \ulcorner \zeta \urcorner$ and $z' = j(z)$.

Part 1 is by induction on λ , using 2.21. We just verify that the induction does not fail at a limit stage. We do have the necessary theories in N_λ as usual. So suppose

$$t' = \text{Th}_{k+1}^{N_\lambda}(\zeta' \cup z') \in N_\lambda.$$

Let $\alpha < \lambda$ and $i_{0\alpha} = i_{\mathcal{F}[0, \alpha]}^{N, k}$ and $i_{\alpha\lambda} = i_{\mathcal{F}[\alpha, \lambda]}^{N_\alpha, k}$ and t^* be such that $i_{\alpha\lambda}(t^*) = t'$. Let $z^* = i_{0\alpha}(z)$ and $\zeta^* = \sup i_{0\alpha} \ulcorner \zeta \urcorner$. Then by $\text{r}\Sigma_{k+1}$ elementarity,

$$\text{Th}_{k+1}^{N_\alpha}(\zeta^* \cup z^*) = t^* \in N_\alpha,$$

contradicting the inductive hypothesis.

Part 2 follows from part 1 and 2.20.

Part 3: Suppose that N is not $(k+1)$ -solid. Then by 2.20, $\rho < \zeta$. For a premouse M and $q \in M$, let $\zeta_{k+1}^M(q)$ and $z_{k+1}^M(q)$ be the relativization of the (ζ, z) notions for M at degree $k+1$, to theories in the expanded language with a constant symbol \dot{q} interpreted by q . Clearly the preceding results relativize (for example, $i(z_{k+1}^M(q)) = z_{k+1}^R(i(q))$ for appropriate ultrapower maps $i : M \rightarrow R$, etc). Now $\zeta_{k+1}^N(p) = \rho$ and $z_{k+1}^N(p) = \emptyset$. So

$$\rho_{k+1}^{N_\lambda} \leq \zeta_{k+1}^{N_\lambda}(j(p)) = \sup j \ulcorner \zeta_{k+1}^N(p) \urcorner < \zeta'.$$

So again by 2.20, N_λ is not $(k+1)$ -solid.

Part 4: Suppose that N is not $(k+1)$ -sound. Then by 2.20,

$$N \neq H =_{\text{def}} \text{Hull}_{k+1}^N(\zeta \cup z),$$

and since $\zeta' \leq j(\zeta)$ it suffices to see

$$N_\lambda \neq \text{Hull}_{k+1}^{N_\lambda}(j(\zeta) \cup z').$$

So let $x \in N \setminus H$. We claim that

$$j(x) \notin \text{Hull}_{k+1}^{N_\lambda}(j(\zeta) \cup z'), \quad (4)$$

which suffices. Now clearly $\zeta < \rho_k^N$. Stratify H via the methods of [3, 2.10] (cf. 2.23). So

$$H = \bigcup_{\zeta \leq \alpha < \rho_k^N} H_\alpha,$$

where H_α is the set of all $y \in H$ such that for some $\beta < \zeta$ and minimal $\text{r}\Sigma_{k+1}$ Skolem term r ,

$$t_\alpha = \text{Th}_k^N(\alpha \cup \{p_k^N\})$$

codes a witness to the fact that $y = r(z, \beta, u_k^N)$. (Here if $k = 0$, t_α should be replaced with the α^{th} “fragment” of $\mathfrak{C}_0(N)$ in the usual manner.)

Let $\alpha \in (\zeta, \rho_k^N)$ with

$$x, z, u_k^N \in \text{Hull}_k^N(\alpha \cup \{p_k^N\}),$$

or $x, z, u_k^N \in t_\alpha$ if $k = 0$, and α a limit if $k > 0$. The fact that $x \notin H_\alpha$ is then an $\text{r}\Sigma_k$ assertion about (x, z, u_k^N, t_α) .²² But

$$j(t_\alpha) = \text{Th}_k^{N_\lambda}(j(\alpha) \cup \{p_k^{N_\lambda}\}),$$

and j is $\text{r}\Sigma_k$ -elementary, so $j(x) \notin H_{j(\alpha)}^{N_\lambda}$, where $H_\gamma^{N_\lambda}$ is defined over N_λ analogously. Since $\rho_k^{N_\lambda} = \sup j^{\text{“}}\rho_k^N$, line (4) follows. So by 2.20, N_λ is not $(k+1)$ -sound.

Now suppose instead that N is $(k+1)$ -sound. Let ξ be least such that either

$$\xi = \lambda \text{ or } \text{cr}(F_\xi) \geq \rho' = \sup i_{0\xi}^{\text{“}}\rho.$$

Let $p' = i_{0\xi}(\rho)$. Then by standard arguments, $\rho_{k+1}^{N_\xi} = \rho'$, $p_{k+1}^{N_\xi} = p'$ and N_ξ is $(k+1)$ -sound. (Alternatively, some of those arguments can be avoided by noting that by part 1 and some other arguments,

$$N_\xi = \text{Hull}_{k+1}^{N_\xi}(\zeta_{k+1}^{N_\xi} \cup \{z_{k+1}^{N_\xi}\}),$$

so by 2.20, N_ξ is $(k+1)$ -sound.) So we are done if $\lambda = \xi$, and if $\lambda > \xi$ then note that $\text{cHull}_{k+1}^{N_\lambda}(\zeta' \cup z') = N_\xi \neq N_\lambda$, so N_λ is not sound. \square (2.24)

Remark 2.25. The previous result is optimal in the sense that a degree 0 ultrapower of a 1-sound but non-2-sound structure, can be fully sound. For let N be a 1-sound premouse such that for some $\kappa < \lambda < \text{OR}^N$, $N = \mathcal{J}_\kappa(N|\lambda)$ and $\rho_1^N = \lambda$ and λ is $\text{r}\Sigma_1^N$ -regular, and suppose there is a short extender E weakly amenable to N with $\text{cr}(E) = \kappa$. Let $U_1 = \text{Ult}_1(N, E)$ and $U_0 = \text{Ult}_0(N, E)$, and suppose that each U_i is wellfounded. Then it is easy to see that $U_0 \triangleleft U_1$, and so U_0 is fully sound. We can arrange this situation with N being non-2-sound (and so $\rho_2^N < \lambda$, but we might have either $\kappa < \rho_2^N$ or $\rho_2^N \leq \kappa$).

We now analyze extenders used in iteration trees. The analysis decomposes such extenders into linear compositions of Dodd sound extenders, identified via Dodd-fragment parameters and projecta. The associativity of extenders underlies the analysis.

²²The $\text{r}\Sigma_k$ complexity arises in identifying some terms which output x, z, u_k^N .

Definition 2.26. Let P be a k -sound segmented-premouse, $F = F^P$, and E a short extender over P with $\text{cr}(E) < \rho_k^P$. Then $E \circ_k P$ denotes $\text{Ult}_k(P, E)$; $E \circ_k F$ or $\text{Ult}_k(F, E)$ denotes $F^{\text{Ult}_k(P, E)}$. Write \circ for \circ_0 . In the absence of parentheses, we take association of \circ_k to the right: $E \circ_k F \circ_l Q = E \circ_k (F \circ_l Q)$. \dashv

Lemma 2.27 (Associativity of Extenders). *Let P, Q be seg-premice. Suppose $F = F^P \neq \emptyset$, F is over Q , Q is k -sound and $\kappa_F < \rho_k^Q$. Let E be a short extender over P such that $(\kappa_F^+)^P < \kappa_E$.²³ Let $U = \text{Ult}_k(Q, F)$ and $U_E^P = \text{Ult}_0(P, E)$. Suppose U and U_E^P are wellfounded. Then*

$$(E \circ F) \circ_k Q = E \circ_k (F \circ_k Q),$$

$$i_{E \circ F}^{Q, k} = i_E^{U, k} \circ i_F^{Q, k}.$$

Proof. Let $\lambda = \text{OR}^P$. Then λ is a U -cardinal, $\lambda < \rho_k^U$ and $\mathcal{H}_\lambda^U = [P]$. So E is an extender over \mathcal{H}_λ^U , hence over U . Let $j = i_E^{U, k}$. Note j is continuous at λ , $j(\lambda) = \text{OR}^{U_E^P}$ and $\mathcal{H}_{j(\lambda)}^{\text{Ult}_k(U, E)} = [U_E^P]$ and $j \upharpoonright \mathcal{H}_\lambda^U = i_E^P$. By 2.11,

$$\nu_{E \circ F} = \nu_E \cup \sup i_E^P \nu_F.$$

And letting $\xi = \max(\nu_F, \text{lgcd}(P))$, for $A \in P \cap \mathcal{P}([\kappa_F]^{<\omega})$,

$$i_{E \circ F}(A) \cap [\nu_{E \circ F}]^{<\omega} = i_E^P(i_F(A) \cap [\xi]^{<\omega}) \cap [\nu_{E \circ F}]^{<\omega}. \quad (5)$$

Define an isomorphism $\psi : \text{Ult}_k(Q, E \circ F) \rightarrow \text{Ult}_k(U, E)$ by

$$\psi([\tau_q, i_E^P(a) \cup b]_{E \circ F}^{Q, k}) = [\tau'_{(q', a)}, b]_E^{U, k},$$

for k -terms τ_q defined from parameter $q \in Q$, and $a \in [\nu_F]^{<\omega}$, $b \in [\nu_E]^{<\omega}$, and where $q' = i_F^{Q, k}(q)$ and τ' is defined from τ by converting the appropriate arguments to parameters. Los' Theorem, (5) and the fact that degree k embeddings respect the T_k predicate, show ψ is well-defined and $\text{r}\Sigma_k$ -elementary; surjectivity is clear. Moreover, ψ commutes with the ultrapower embeddings, so $i_{E \circ F}^{Q, k} = i_E^{U, k} \circ i_F^{Q, k}$. \square

Corollary 2.28. *Let P_n, \dots, P_0, Q be segmented-premice. Suppose that for each i , $E_i = F^{P_i} \neq \emptyset$, $\text{cr}(E_{i+1}) > \text{cr}(E_i)$, E_{i+1} is over P_i , E_0 is over Q , and $\text{cr}(E_0) < \rho_k^Q$. Then writing $Q_0 = Q$ and $Q_{i+1} = \text{Ult}_k(Q_i, E_i)$,*

$$\begin{aligned} ((\dots (E_n \circ E_{n-1}) \circ \dots) \circ E_0) \circ_k Q &= E_n \circ_k (\dots \circ_k (E_0 \circ_k Q)); \\ i_{((\dots (E_n \circ E_{n-1}) \circ \dots) \circ E_0)}^{Q, k} &= i_{E_n}^{Q_n, k} \circ \dots \circ i_{E_0}^{Q_0, k}. \end{aligned}$$

Definition 2.29 (Dodd core). Let G be an extender such that (σ_G, s_G) are defined. The *Dodd core* of G , denoted $\mathfrak{C}_D(G)$, is the transitive collapse of $G \upharpoonright \sigma_G \cup s_G$.²⁴ We often identify $\mathfrak{C}_D(G)$ with its trivial completion. \dashv

²³We initially had the stronger assumption that $\kappa_F < \kappa_E$ and E is weakly amenable; the referee noticed that it might be enough to assume that $(\kappa_F^+)^P < \kappa_E$, which it is.

²⁴That is, let $E_0 = G \upharpoonright \sigma_G \cup s_G$, let $s' = [s, \text{id}]_{E_0}^M$ where G is over M , and let $j = i_{E_0}^M$. Then $\mathfrak{C}_D(G)$ is the extender derived from j with support $\sigma_G \cup s'$.

Remark 2.30. Let S be a k -sound premouse such that every $E \in \mathbb{E}_+^S$ is Dodd sound. Let \mathcal{W} be a k -maximal tree on S . Let $\alpha < \text{lh}(\mathcal{W})$ and $G \in \mathbb{E}_+(M_\alpha^\mathcal{W})$, with G not Dodd sound.

By elementarity, $G = F(M_\alpha^\mathcal{W})$, so α is the unique α' such that $G \in \mathbb{E}_+(M_{\alpha'}^\mathcal{W})$. Write $\alpha_G = \alpha$. Lemmas 2.12–2.18 show $\deg^\mathcal{T}(\alpha_G) = 0$ and $\mathfrak{C}_D(G) = F(M_{\beta+1}^{\ast\mathcal{W}})$, where β is the least β' such that (i) $\beta' + 1 \leq_\mathcal{W} \alpha_G$ and \mathcal{W} does not drop in model in $(\beta' + 1, \alpha_G]_\mathcal{W}$, and (ii) $F(M_{\beta'+1}^\mathcal{W})$ is not Dodd sound; condition (ii) can equivalently be replaced with (ii') $\text{cr}(i_{\beta'+1, \alpha_G}^{\ast\mathcal{W}}) \geq \tau_F$ where $F = F(M_{\beta'+1}^{\ast\mathcal{W}})$.

Also, $\gamma = \text{pred}^\mathcal{W}(\beta+1)$ is the unique ordinal γ' such that $\mathfrak{C}_D(G) \in \mathbb{E}_+(M_{\gamma'}^\mathcal{W})$. For \mathcal{W} is normal and $\text{lh}(E_\gamma^\mathcal{W}) \leq \text{OR}(M_{\beta+1}^{\ast\mathcal{W}}) = \text{lh}(\mathfrak{C}_D(G))$.²⁵

For Dodd sound $G \in \mathbb{E}_+(M_\alpha^\mathcal{W})$, α_G denotes the least β with $G \in \mathbb{E}_+(M_\beta^\mathcal{W})$.

Definition 2.31 (Dodd ancestry). Let \mathcal{W}, α be as in 2.30 and $G \in \mathbb{E}_+(M_\alpha^\mathcal{W})$. We define the (Dodd) ancestry of G in \mathcal{W} , denoted $\text{Da}^\mathcal{W}(G)$, and the ordering $<_{\text{Da}}^\mathcal{W}$, recursively on α_G . If G is Dodd-sound let $\text{Da}^\mathcal{W}(G) = \emptyset$. Suppose G is not Dodd-sound. Let γ be such that $\mathfrak{C}_D(G) \in \mathbb{E}_+(M_\gamma^\mathcal{W})$ (by 2.30, γ is unique). Define $\text{Da}^\mathcal{W}(G)$ to be the sequence $d = \langle d_\beta \rangle_{\beta \in D}$ with domain

$$D = \{\beta \mid \beta + 1 \in (\gamma, \alpha_G]_\mathcal{W}\},$$

such that $d_\beta = \text{Da}^\mathcal{W}(E_\beta^\mathcal{W})$.

Now recursively define the relation $<_{\text{Da}}^\mathcal{W}$ by:

$$\beta <_{\text{Da}}^\mathcal{W} \gamma \iff \exists \beta' \in \text{dom}(\text{Da}^\mathcal{W}(E_\gamma^\mathcal{W}))[\beta = \beta' \text{ or } \beta <_{\text{Da}}^\mathcal{W} \beta'].$$

We also write $E_\beta^\mathcal{W} <_{\text{Da}}^\mathcal{W} E_\gamma^\mathcal{W}$ to mean $\beta <_{\text{Da}}^\mathcal{W} \gamma$. ⊣

Note that in 2.31, G is Dodd-sound iff $\text{Da}^\mathcal{W}(G) = \emptyset$.

Figure 1 presents a typical Dodd ancestry. An extender E is represented by the symbol \lfloor , with $\text{cr}(E)$ and $\text{lh}(E)$ corresponding to the lower and upper bounds of the symbol respectively. $E <_{\text{Da}} F$ iff E is pictured to the left of F , within its vertical bounds. So Dodd sound extenders have no extenders to their left. In the figure, $G <_{\text{Da}} H <_{\text{Da}} J$ and $G <_{\text{Da}} J$, but $G, H \not<_{\text{Da}} I$.

Lemma 2.32. Let \mathcal{W}, G be as in 2.31. Then:

- (a) For all $\beta <_{\text{Da}}^\mathcal{W} \gamma$ we have $\text{cr}(E_\gamma^\mathcal{W}) < \text{cr}(E_\beta^\mathcal{W}) < \text{lh}(E_\beta^\mathcal{W}) < \text{lh}(E_\gamma^\mathcal{W})$.
- (b) Let $\gamma_1, \gamma_2 \in \text{dom}(\text{Da}^\mathcal{W}(G))$, with $\gamma_1 < \gamma_2$. For all $\gamma'_1 \leq_{\text{Da}}^\mathcal{W} \gamma_1$ and $\gamma'_2 \leq_{\text{Da}}^\mathcal{W} \gamma_2$, we have $\gamma'_1 < \gamma'_2$.
- (c) Let $\gamma \in \text{dom}(\text{Da}^\mathcal{W}(G))$ and let $\lambda = \text{pred}^\mathcal{W}(\gamma + 1)$. Suppose that $E_\gamma^\mathcal{W} = F(M_\gamma^\mathcal{W})$. Then \mathcal{W} drops in model at some $\beta \in (0, \gamma]_\mathcal{W}$ such that $\beta > \lambda$.

²⁵Literally here, we mean $\text{OR}(M_{\beta+1}^{\ast\mathcal{W}}) = \text{lh}(G')$ where G' is the trivial completion of $\mathfrak{C}_D(G)$.

(vi) if $m > 0$ then $\text{cr}(G) < \nu_\zeta^{\mathcal{W}\alpha}$.

Proof. We prove the lemma by induction on α_G for $\alpha_G + 1 < \text{lh}(\mathcal{W})$, with a subinduction on $\alpha \in [1, \lambda]$. So assume it holds for $H = E_{\alpha_H}^{\mathcal{W}}$ for each $\alpha_H < \alpha_G$.

If G is Dodd sound then $\lambda = 1$, and we use $\zeta = \alpha_G$ and $m = 0$.

Suppose G is not Dodd sound. So $\lambda > 1$. We use 2.30 in the following.

Suppose $\alpha = 1$. We have $G_1 = \mathfrak{C}_D(G)$. Set $m = 0$ and ζ with $G_1 \in \mathbb{E}_+(M_\zeta^{\mathcal{W}})$ and $\beta + 1$ such that $\zeta = \text{pred}^{\mathcal{W}}(\beta + 1) \leq_{\mathcal{W}} \alpha_G$. Then (v) holds because

$$\text{cr}(G) = \text{cr}(G_1) < \tau_{G_1} \leq \text{cr}(i_{\beta+1, \alpha_G}^{*\mathcal{W}}) < \nu_\zeta^{\mathcal{W}}.$$

Suppose $\alpha = \beta + 1 > 1$. So $G_{\beta+1} = \text{Ult}_0(G_\beta, F_\beta)$. Let β' be such that $F_\beta = \mathfrak{C}_D(E_{\beta'}^{\mathcal{W}})$. Let $\gamma' \leq_{\text{Da}}^{\mathcal{W}} \alpha_G$ be such that $\beta' \in \text{dom}(\text{Da}^{\mathcal{W}}(E_{\gamma'}^{\mathcal{W}}))$. Let $\varepsilon' = \text{pred}^{\mathcal{W}}(\beta' + 1)$. Let ζ be such that $F_\beta \in \mathbb{E}_+(M_\zeta^{\mathcal{W}})$. Then define $m \in [1, \omega)$ and $\mathcal{X} = \mathcal{W}_{\beta+1}$ by setting $E_\zeta^{\mathcal{X}} = F_\beta$, and $E_{\zeta+i}^{\mathcal{X}} = F(M_{\zeta+i}^{\mathcal{X}})$ for $1 \leq i < m$, until we reach $M_{\zeta+m}^{\mathcal{X}}$ with active extender F with $\text{cr}(F) = \text{cr}(G)$. Each $E_{\zeta+i}^{\mathcal{X}}$ is a sub-extender of some $G' \leq_{\text{Da}}^{\mathcal{W}} G$, and applies to the same premouse in \mathcal{X} as does G' in \mathcal{W} . So $\text{cr}(E_{\zeta+i+1}^{\mathcal{X}}) < \text{cr}(E_{\zeta+i}^{\mathcal{X}})$.

For instance, $\text{pred}^{\mathcal{X}}(\zeta + 1) = \varepsilon'$. For by 2.30 and normality of \mathcal{W} ,

$$\text{cr}(F_\beta) = \text{cr}(E_{\beta'}^{\mathcal{W}}) < \min(\nu_\zeta^{\mathcal{W}}, \nu_{\varepsilon'}^{\mathcal{W}}),$$

and for each $\delta < \zeta$, we have $\text{lh}(E_\delta^{\mathcal{W}}) \leq \nu_\zeta^{\mathcal{W}} < \text{lh}(F_\beta)$. In particular, $\zeta \geq \varepsilon'$. But $\mathcal{W} \restriction \zeta + 1 = \mathcal{X} \restriction \zeta + 1$, so $\text{pred}^{\mathcal{X}}(\zeta + 1) = \varepsilon'$. If $\gamma' = \alpha_G$ then $m = 1$. Otherwise $m > 1$, and $E_{\zeta+1}^{\mathcal{X}} = F(M_{\zeta+1}^{\mathcal{X}})$ is a subextender of $F(M_{\beta'+1}^{\mathcal{W}})$, and each apply to the same premouse, etc.

We claim that $\mathcal{W}_{\beta+1} = \mathcal{X}$ is as desired. Certainly \mathcal{X} is normal, and its models are wellfounded, since they embed into models of \mathcal{W} . Now let $H_0 = F_\beta$ and let $H_{i+1} = F(M_{\zeta+i+1}^{\mathcal{X}})$ for $i < m$. Then using 2.28, $F(M_{\zeta+m}^{\mathcal{X}})$ is

$$(\dots((H_0 \circ H_1) \circ H_2) \circ \dots \circ H_m) = H_0 \circ H_1 \circ \dots \circ H_m. \quad (6)$$

Let α_D be such that $\mathfrak{C}_D(G) \in \mathbb{E}_+(M_{\alpha_D}^{\mathcal{W}})$. Let $\iota = \text{pred}^{\mathcal{X}}(\zeta + m)$. So

$$\iota = \max([\alpha_D, \alpha_G]_{\mathcal{W}} \cap \beta' + 1).$$

Let $X = \{\delta' \mid \delta' + 1 \in (\alpha_D, \iota]_{\mathcal{W}}\}$. Then

$$H_m = \text{Ult}_0(\mathfrak{C}_D(G), \langle E_{\delta'}^{\mathcal{W}} \rangle_{\delta' \in X}). \quad (7)$$

By induction, for $\delta' \in X$, $E_{\delta'}^{\mathcal{W}} = \text{Ult}_0(F_\delta, \mathcal{F} \restriction (\delta, \varepsilon))$, where $\mathcal{F} \restriction (\delta, \varepsilon)$ is the Dodd decomposition of $E_{\delta'}^{\mathcal{W}}$ (so $F_\delta = \mathfrak{C}_D(E_{\delta'}^{\mathcal{W}})$). But then

$$\text{Ult}_0(M_{\delta'+1}^{*\mathcal{W}}, E_{\delta'}^{\mathcal{W}}) = \text{Ult}_0(M_{\delta'+1}^{*\mathcal{W}}, \mathcal{F} \restriction (\delta, \varepsilon)),$$

and the ultrapower maps agree. This fact is a straightforward extension of 2.27; we omit the details. Therefore $H_m = \text{Ult}_0(\mathfrak{C}_D(G), \mathcal{F} \restriction [1, \xi])$, where $\mathcal{F} \restriction [1, \xi)$

is the concatenation of the Dodd decompositions of $E_{\delta'}^{\mathcal{W}}$ for $\delta' \in X$, and the ultrapower map is that corresponding to (7). We get a similar representation for each H_i , thus partitioning $\mathcal{F} \restriction [1, \beta]$ into $m + 1$ intervals. Finally, using (6) and the extension of 2.27, then $F(M_{\zeta+m}^{\mathcal{X}})$ is

$$\text{Ult}_0(\mathfrak{C}_D(G), \mathcal{F} \restriction [1, \beta + 1)) = G_{\beta+1}.$$

We leave the remaining details and limit case to the reader. \square (2.35)

Lemma 2.36. *Adopt the hypotheses and notation of 2.35. Let P be an m -sound seg-pm, with G over P and $\text{cr}(G) < \rho_m^P$. Suppose that $\text{Ult}_m(P, G)$ is wellfounded. Then for $\alpha \leq \lambda$, we have $\text{Ult}_m(P, G_\alpha) = \text{Ult}_m(P, \mathcal{F} \restriction \alpha)$ and $i_{G_\alpha}^{P,m} = i_{\mathcal{F} \restriction \alpha}^{P,m}$, and the degree m core sequence of $i_G^{P,m}$ is*

$$\left\langle Q_\alpha = \text{Ult}_m(P, \mathcal{F} \restriction \alpha), i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Q_\alpha, m} \right\rangle_{\alpha \leq \lambda}.$$

In particular, $i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Q_\alpha, m}$ is not superstrong.

Suppose that P is $(m + 1)$ -solid. Then $i_{\mathcal{F} \restriction \alpha}^{P,m}$ and $i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Q_\alpha, m}$ are m -embeddings, preserve p_{m+1} , and are cofinal at ρ_{m+1} .

Proof Sketch. For the (inductively established) characterization of the core sequence of $i_G^{P,m}$, given Q_α and factor embedding $j_\alpha = i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Q_\alpha, m}$ as above, note that

$$\text{Ult}_m(P, G) = \text{Ult}_m(Q_\alpha, \mathcal{F} \restriction [\alpha, \lambda]),$$

and show that the natural factor embedding

$$k : \text{Ult}_m(Q_\alpha, F_\alpha) \rightarrow \text{Ult}_m(P, G)$$

maps the maximal fragments of F_α in $\text{Ult}_m(Q_\alpha, F_\alpha)$ (corresponding to $(s_{F_\alpha}, \sigma_{F_\alpha})$) to those of the extender derived from $i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Q_\alpha, m}$. This follows the argument for 2.13, and that $\sigma_{F_\alpha} < \text{cr}(F_\beta)$ when $\alpha < \beta$. For the second paragraph, use 2.24 and commutativity. \square

Definition 2.37. Let \mathcal{W} be a k -maximal tree on a k -sound premouse N , of length $\zeta + 1$. Let $P = M_\zeta^{\mathcal{W}}$ and $\gamma \leq \text{OR}^P$, with $\text{lh}(E_\alpha^{\mathcal{W}}) < \gamma$ for every $\alpha + 1 < \text{lh}(\mathcal{W})$. Let E be an extender such that $(P \restriction \gamma, E)$ is a premouse.

The *potential k -maximal tree* $\mathcal{W} \hat{\ } \langle E \rangle$ is the “putative iteration tree” \mathcal{W}^+ on N of length $\zeta + 2$, extending \mathcal{W} , with $E_\zeta^{\mathcal{W}^+} = E$, and with $\text{pred}^{\mathcal{W}^+}(\zeta + 1)$, etc, determined by the rules for k -maximality. We say the tuple $(k, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ is *potential for N* .

As usual $\Phi(\mathcal{W})$ denotes the phalanx associated to \mathcal{W} (see [18]). Let (k, \mathcal{W}, \dots) be potential for N . Let $U = M_{\zeta+1}^{\mathcal{W}^+}$ and $d = \text{deg}^{\mathcal{W}^+}(\zeta + 1)$. Let $\iota = \text{lgcd}(P \restriction \text{lh}(E))$. We write $\Phi(\mathcal{W}, \iota, E)$ for the phalanx

$$\langle (\Phi(\mathcal{W}), < \iota), (U, d), \text{lh}(E) \rangle.$$

Normal trees \mathcal{U} on $\Phi(\mathcal{W}, \iota, E)$, indexed with ordinals $\geq \zeta + 1$ (the first model of \mathcal{U} is $M_{\zeta+1}^{\mathcal{U}}$), must satisfy the usual conditions for k -maximality, except/including that (i) $M_{\zeta+1}^{\mathcal{U}} = U$, (ii) $\text{lh}(E) < \text{lh}(E_{\zeta+1}^{\mathcal{U}})$, (iii) if $\text{cr}(E_{\alpha}^{\mathcal{U}}) < \iota$ then $\text{pred}^{\mathcal{U}}(\alpha + 1) = \delta \leq \zeta$ and $M_{\alpha+1}^{*\mathcal{U}} \trianglelefteq M_{\delta}^{\mathcal{W}}$ and $\deg^{\mathcal{U}}(\alpha + 1)$ are as usual (like for $\Phi(\mathcal{W})$), and (iv) if $\text{cr}(E_{\alpha}^{\mathcal{U}}) = \iota$ then $\text{pred}^{\mathcal{U}}(\alpha + 1) = \zeta + 1$ and $M_{\alpha+1}^{*\mathcal{U}} = U$ and $\deg^{\mathcal{U}}(\zeta + 1) = d$. \dashv

We now establish a phalanx iterability criterion guaranteeing that an extender is on the sequence of a premouse. The lemma should be compared with [18, 8.6] (to which it is very similar); there, the exchange ordinal is ν_E instead of ι . The lower exchange ordinal leads to our need to appeal to the Dodd-structure analysis in the proof. (And in our application of the lemma later, our iterability proof only seems to give iterability with respect to ι .)

Lemma 2.38. Let M be an ω -sound premouse projecting to ω . Suppose that every $E \in \mathbb{E}_+^M$ is Dodd sound. Let $\mathfrak{c} < \text{OR}^M$ and let Σ be an above- \mathfrak{c} , $(\omega, \omega_1 + 1)$ -strategy for M . Let

$$(\omega, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$$

be potential for M , with \mathcal{W} via Σ . Let $\iota = \text{lgcd}(P || \text{lh}(E))$. Suppose that $\mathfrak{c} < \text{lh}(E)$ and \mathfrak{c} is a cutpoint of $P || \text{lh}(E)$.²⁶

Then $E \in \mathbb{E}_+^P$ iff the phalanx $\mathfrak{P} = \Phi(\mathcal{W}, \iota, E)$ is normally $(\omega_1 + 1)$ -iterable.

Proof. We have $\zeta + 1 = \text{lh}(\mathcal{W})$ and $P = M_{\zeta}^{\mathcal{W}}$ and $(P || \text{lh}(E), E)$ is a premouse. Let $\delta = \text{pred}^{\mathcal{W}^+}(\zeta + 1)$ and $Q = M_{\zeta+1}^{*\mathcal{W}^+} \trianglelefteq M_{\delta}^{\mathcal{W}}$ and $m = \deg^{\mathcal{W}^+}(\zeta + 1)$ and

$$i_E = i_E^{Q, m} : Q \rightarrow \text{Ult}_m(Q, E) = M_{\zeta+1}^{\mathcal{W}^+}.$$

The proof that \mathfrak{P} is iterable assuming $E \in \mathbb{E}_+^P$ we mostly leave to the reader since we won't use this fact. The point is that $\Phi(\mathcal{W}^+) = \Phi(\mathcal{W}, \nu(E), E)$ is iterable since M is iterable above \mathfrak{c} , and there is a direct correspondence between normal trees \mathcal{U} on \mathfrak{P} and normal trees \mathcal{V} on $\Phi(\mathcal{W}^+)$. (For example when E is type 2: Let $\beta + 1 < \text{lh}(\mathcal{U})$ with $\text{pred}^{\mathcal{U}}(\beta + 1) = \zeta + 1$ and $\text{cr}(E_{\beta}^{\mathcal{U}}) = \iota$. Then $M_{\beta+1}^{*\mathcal{U}} = M_{\zeta+1}^{\mathcal{W}^+}$ and $\deg^{\mathcal{U}}(\beta + 1) = m$, whereas $\text{pred}^{\mathcal{V}}(\beta + 1) = \zeta$ and $M_{\beta+1}^{*\mathcal{V}} = P || \text{lh}(E)$ and $\deg^{\mathcal{V}}(\beta + 1) = 0$. It follows that

$$M_{\beta+1}^{\mathcal{U}} = \text{Ult}_m(Q, F(M_{\beta+1}^{\mathcal{V}})),$$

and in particular, $M_{\beta+1}^{\mathcal{U}}$ is wellfounded.)

So assume \mathfrak{P} is iterable. In comparison of \mathfrak{P} vs $\Phi(\mathcal{W})$, the resulting trees are above \mathfrak{c} (but we allow $\text{cr}(E) < \mathfrak{c}$). For if $P || \text{lh}(E)$ is active then $\text{cr}(F^{P || \text{lh}(E)}) \geq \mathfrak{c}$ because \mathfrak{c} is a cutpoint of $P || \text{lh}(E)$. So suppose there is an active premouse N such that $P || \text{lh}(E) \triangleleft N$ and $\text{lh}(E)$ is a cardinal of N and $\text{cr}(F^N) < \mathfrak{c}$. Then

²⁶Note we don't assume $\mathfrak{c} \leq \text{cr}(E)$. But if $P || \text{lh}(E)$ is active then $\mathfrak{c} \leq \text{cr}(F^{P || \text{lh}(E)})$.

$\text{cr}(F^N) = \iota < \mathfrak{c}$, by the ISC and since \mathfrak{c} is a cutpoint of $P||\text{lh}(E)$. So ι is inaccessible in $P||\text{lh}(E)$, so E is type 2 or 3, so

$$\text{cr}(E) \text{ is } < \iota\text{-strong in } P||\text{lh}(E), \text{ as witnessed by } \mathbb{E}^{P||\text{lh}(E)}.$$

The ultrapower map $j : P||\text{lh}(E) \rightarrow \text{Ult}(P||\text{lh}(E), F^N)$ preserves this statement. But then by the ISC, $P||\text{lh}(E)$ does not have a cutpoint above ι , contradiction.

So we get a successful ω -maximal comparison $(\mathcal{X}', \mathcal{Y}')$, with \mathcal{Y}' such that $\mathcal{W} \smallfrown \mathcal{Y}'$ is via Σ . Write

$$\mathcal{X} = \mathcal{W}^+ \smallfrown \mathcal{X}' \text{ and } \mathcal{Y} = \mathcal{W} \smallfrown \mathcal{Y}',$$

so \mathcal{Y} is an ω -maximal tree on M with $\mathcal{Y} \restriction (\zeta + 1) = \mathcal{W}$, and $\mathcal{X} \restriction (\zeta + 2) = \mathcal{W}^+$ and \mathcal{X} is ω -maximal except that $E_\zeta^\mathcal{X} = E$ so maybe $E \notin \mathbb{E}_+(M_\zeta^\mathcal{X})$, and maybe the exchange ordinal $\iota < \nu_E$. So we can't apply the Closeness Lemma [3, 6.1.5] to \mathcal{X} . But for all $\alpha + 1 < \text{lh}(\mathcal{X})$, $E_\alpha^\mathcal{X}$ is weakly amenable over $M_{\alpha+1}^{*\mathcal{X}}$. Let $Z = M_\infty^\mathcal{X}$. Clearly $Z \leq M_\infty^\mathcal{Y}$. By 2.24, Z is unsound, so $Z = M_\infty^\mathcal{Y}$. Let $\alpha + 1 \in b^\mathcal{X}$ be least such that $(\alpha + 1, \infty]_\mathcal{X}$ does not drop in model or degree, and let $n = \deg^\mathcal{X}(\alpha + 1)$. Since $Z = M_\infty^\mathcal{Y}$, therefore Z is $n + 1$ -solid, so by 2.24,

$$p_{n+1}^Z = i_{\alpha+1, \infty}^{*\mathcal{X}}(p_{n+1}(M_{\alpha+1}^{*\mathcal{X}}))$$

and $\rho_{n+1}^Z = \rho_{n+1}(M_{\alpha+1}^{*\mathcal{X}})$. We have $\mathcal{W} = \mathcal{Y} \restriction (\zeta + 1)$. Standard arguments now show that $b^\mathcal{X}$ is above U and $b^\mathcal{X}$ does not drop in model or degree above U (that is, $\zeta + 1 \in b^\mathcal{X}$ and $(\zeta + 1, \infty]_\mathcal{X}$ does not drop in model or degree), so

$$j^\mathcal{X} =_{\text{def}} i_{\zeta+1, \infty}^{*\mathcal{X}} : U \rightarrow Z.$$

Similar arguments show that $\mathcal{Y} \neq \mathcal{W}$, and letting $\beta + 1 = \min(b^\mathcal{Y} \setminus (\zeta + 1))$, that $\delta = \text{pred}^\mathcal{Y}(\beta + 1)$, $M_{\beta+1}^{*\mathcal{Y}} = Q = M_{\zeta+1}^{*\mathcal{W}^+}$ and $\deg^\mathcal{Y}(\beta + 1) = m$ and $(\beta + 1, \infty]_\mathcal{Y}$ does not drop in model or degree, so

$$j^\mathcal{Y} = i_{\beta+1, \infty}^{*\mathcal{Y}} : Q \rightarrow Z.$$

We have $j^\mathcal{X} \circ i_E = j^\mathcal{Y}$, since these maps preserve p_{m+1} and do not move the generators that generate Q . If $\nu_E \leq \text{cr}(j^\mathcal{X})$, standard arguments now show $E \in \mathbb{E}_+^P$. So assume E is type 2, with largest generator γ , and

$$\text{cr}(j^\mathcal{X}) = \iota \leq \gamma < (\iota^+)^U = \text{lh}(E).$$

Let $G = E_\beta^\mathcal{Y}$. So $G \restriction \nu_G \subseteq E_{j^\mathcal{Y}}$.

Claim 1. G is the only extender used on $b^\mathcal{Y}$, and $\nu_G = j^\mathcal{X}(\gamma + 1)$.

Proof. Let $\sigma \in [\zeta + 1, \infty]_\mathcal{X}$ be least such that

$$\sigma = \infty \text{ or } \text{cr}(i_{\sigma, \infty}^\mathcal{X}) > i_{\zeta+1, \sigma}^\mathcal{X}(\gamma).$$

Let $(\iota_1, \gamma_1) = i_{\zeta+1, \sigma}^{\mathcal{X}}(\iota, \gamma)$. Now $E \restriction \iota \in U$ and

$$E_{j^{\mathcal{Y}}} \restriction \iota_1 = j^{\mathcal{X}}(E \restriction \iota) \in Z,$$

since $\text{cr}(E) < \text{cr}(j^{\mathcal{X}})$ and $j^{\mathcal{X}} \circ i_E = j^{\mathcal{Y}}$. It follows that $\iota_1 < \nu_G$ and $G \restriction \iota_1 \in Z$. Since $p_{m+1}^Z = j^{\mathcal{X}} \circ i_E(p_{m+1}^Q)$, Z has the $(m+1)$ -hull property at $\gamma_1 + 1$, i.e.,

$$\text{Hull}_{m+1}^Z((\gamma_1 + 1) \cup \{p_{m+1}^Z\}) \cap \mathcal{P}(\gamma_1 + 1) = Z \cap \mathcal{P}(\gamma_1 + 1) \quad (8)$$

(cf. [18, Example 4.3 and following Remark]). But $\gamma_1 < ((\iota_1)^+)^Z$ and γ_1 is a generator of G . For if $f : [\mu]^{<\omega} \rightarrow \mu$ with $f \in Q$, then

$$\gamma \notin i_E(f) "[\gamma]^{<\omega} \implies \gamma_1 \notin j^{\mathcal{X}}(i_E(f)) "[\gamma_1]^{<\omega} \implies \gamma_1 \notin i_G^{Q,m}(f) "[\gamma_1]^{<\omega}.$$

Therefore $\gamma_1 + 1 = \nu_G$. It follows that

$$M_\sigma^{\mathcal{X}} = \text{Hull}_{m+1}^Z((\gamma_1 + 1) \cup \{p_{m+1}^Z\}) = M_{\beta+1}^{\mathcal{Y}},$$

but then in fact $M_\sigma^{\mathcal{X}} = Z = M_{\beta+1}^{\mathcal{Y}}$, proving the claim. \square (Claim 1)

By Claim 1, E is a subextender of G . We will refine this observation, using the Dodd structure analysis. Now 2.35, 2.36 apply to \mathcal{Y}, G . Let $\mathcal{F} = \langle F_\alpha \rangle_{\alpha < \lambda}$ be the Dodd decomposition of G . Let

$$G_\alpha = \text{Ult}_0(\text{id}, \mathcal{F} \restriction \alpha).$$

Let $\langle Z_\alpha, j_\alpha \rangle_{\alpha \leq \lambda}$ be the degree m core sequence of $i_G^{Q,m} = j^{\mathcal{Y}}$. For $\alpha \leq \beta$ let $j_{\alpha\beta} = j_\beta^{-1} \circ j_\alpha$. By 2.36,

$$Z_\alpha = \text{Ult}_m(Q, \mathcal{F} \restriction \alpha),$$

$$j_{\alpha\beta} = i_{\mathcal{F} \restriction [\alpha, \beta]}^{Z_\alpha, m} \text{ and } j_\alpha = i_{\mathcal{F} \restriction [\alpha, \lambda]}^{Z_\alpha, m},$$

and $j_{\alpha\beta}, j_\alpha$ are m -embeddings which preserve p_{m+1} .

Claim 2. *There is $\varepsilon \leq \lambda$ such that $Z_\varepsilon = U$ and $G_\varepsilon = E$.*

Proof. We will inductively define m -embeddings $i_\alpha : Z_\alpha \rightarrow U$ such that (see Figure 2 for a partial summary):

- $j_\alpha = j^{\mathcal{X}} \circ i_\alpha$ (so i_α preserves p_{m+1}),
- $i_\beta \circ j_{\alpha\beta} = i_\alpha$ for $\alpha \leq \beta$,
- if $\alpha > 0$ then $\iota, \gamma \in \text{rg}(i_\alpha)$,
- if $Z_\alpha \neq U$ or $i_\alpha \neq \text{id}$ then $\text{cr}(i_\alpha) < \iota$ (so if $\alpha > 0$ then $i_\alpha(\text{cr}(i_\alpha)) \leq \iota$).

Case 1. $\alpha = 0$.

We have $Z_0 = Q$ and $j_0 = j^{\mathcal{Y}}$. Set $i_0 = i_E$.




Figure 2: Commuting maps for $\alpha \leq \beta$

Case 2. $\alpha = 1$.

Observe first that $E \restriction \gamma \in U$. This follows the ISC if $E \restriction \gamma$ is not type Z, so suppose otherwise.²⁷ Then ι is the largest generator of $E \restriction \gamma$, and

$$E \equiv_{\iota, \{\iota\}, \{\iota_1\}} G.$$

But $G \restriction \gamma_1 \in Z$, by [12], so $G \restriction \iota \cup \{\iota_1\} \in Z$. Since $\mathcal{P}(\iota) \cap Z = \mathcal{P}(\iota) \cap U$, then $E \restriction \iota + 1 \in U$, so $E \restriction \gamma \in U$ as required.

Let $s = s_E$ and $\sigma = \sigma_E$. Then $\gamma = \max(s)$ since $\nu_E = \gamma + 1$ and $E \restriction \gamma \in U$. Also, $\sigma \leq \iota$ since $E \approx E \restriction \iota \cup \{\gamma\}$. So as above, $G \restriction \sigma \cup j^X(s) \notin Z$, but for each X such that $E \restriction X \in U$, we have $G \restriction j^X(X) = j^X(E \restriction X) \in Z$. Therefore $s_G = j^X(s)$ and $\sigma_G = \sigma$. It follows that

$$Z_1 = \text{cHull}_{m+1}^U(\sigma \cup \{s\} \cup \{p_{m+1}^U\}).$$

Let $i_1 : Z_1 \rightarrow U$ be the uncollapse. Then $\gamma = \max(s) \in \text{rg}(i_1)$, so $\iota \in \text{rg}(i_1)$.

Now suppose $\iota \leq \text{cr}(i_1)$. Since $\gamma \in \text{rg}(i_1)$, then $\gamma < \text{cr}(i_1)$. Since i_1 is an m -embedding preserving p_{m+1} , therefore $U \subseteq \text{rg}(i_1)$, so $Z_1 = U$ and $i_1 = \text{id}$.

Case 3. $\alpha = \beta + 1 > 1$.

Suppose $Z_\beta \neq U$ and $i_\beta : Z_\beta \rightarrow U$ with

$$\kappa = \text{cr}(i_\beta) = \text{cr}(j_\beta) = \text{cr}(F_\beta) < \iota.$$

Now F_β is over Z_β , F_β is on the extender sequence of a premouse (by 2.30) and $Z_\beta \cap \mathcal{P}(\kappa) = Z \cap \mathcal{P}(\kappa) = U \cap \mathcal{P}(\kappa)$. Therefore κ is inaccessible in Z_β and in U . Since $\iota \in \text{rg}(i_\beta)$ we have $\kappa' = i_\beta(\kappa) \leq \iota$.

Now $E_{i_\beta} \notin U$, for otherwise $j^X(\kappa') = j_\beta(\kappa)$ and $E_{j_\beta} = j^X(E_{i_\beta}) \in Z$, contradicting 2.36. So i_β is Dodd-appropriate. Then $j^X(s_{i_\beta}) = s_{j_\beta}$ and $\sigma_{i_\beta} = \sigma_{j_\beta}$ because $\sigma \leq \kappa' \leq \iota$ and $j^X \circ i_\beta = j_\beta$. (In fact, therefore $s_{i_\beta} = s_{j_\beta}$ as $s_{i_\beta} \subseteq \iota$.) Now proceed as in Case 2.

Case 4. α is a limit.

This case follows from the commutativity of the maps before stage α .

²⁷We can't quote [12] here since we don't know that $(P \parallel \text{lh}(E), E)$ is iterable.

This defines all i_α . Now since $Z_\lambda = Z$ and $j_\lambda = \text{id}$, there is $\varepsilon \leq \lambda$ such that $Z_\varepsilon = U$ and $i_\varepsilon = \text{id}$, so $i_{G_\varepsilon}^{Q,m} = i_E$, so $G_\varepsilon = E$. \square (Claim 2)

Fix ε as in Claim 2. Let \mathcal{W}_ε be as in 2.35. Then $E = G_\varepsilon \in \mathbb{E}_+(M_\infty^{\mathcal{W}_\varepsilon})$. But \mathcal{W} and \mathcal{W}_ε are both normal trees via Σ , using only extenders F with $\text{lh}(F) < \text{lh}(E)$, and $M_\infty^{\mathcal{W}} \parallel \text{lh}(E) = M_\infty^{\mathcal{W}_\varepsilon} \parallel \text{lh}(E)$. Therefore $\mathcal{W} = \mathcal{W}_\varepsilon$, and $E \in \mathbb{E}_+^P$, as required. \square (2.38)

3. Extender maximality

Let N be a $(k+1)$ -sound mouse with $\omega < \rho_{k+1}^N$. Our proofs of Theorems 3.4-3.9 require the formation of $(k+1)$ -sound, $\text{r}\Sigma_{k+1}$ -elementary, proper hulls of N , containing a given parameter. The following lemma helps with this.

Lemma 3.1. Let N, n be such that N is an $(n+1)$ -sound premouse and either

- (i) N is $(n, \omega_1, \omega_1 + 1)$ -iterable, or
- (ii) N is $(n+3)$ -sound and $(n+3, \omega_1 + 1)$ -iterable.

Let $\theta \in [\omega, \rho_{n+1}^N)$ be an N -cardinal and $x \in \mathfrak{C}_0(N)$. Then $\exists q \in \mathfrak{C}_0(N)$ such that letting

$$H = \text{Hull}_{n+1}^N(\theta \cup \{q\})$$

and $M = \text{trcoll}(H)$ and $\pi : M \rightarrow N$ be the uncollapse, then $x \in H$, $M \triangleleft N$, $\rho_{n+1}^M = \theta$, $q = \pi(p_{n+1}^M)$, and $p_{n+1}^N = q \setminus \alpha$ for some α .²⁸

Proof. We may assume N is countable, and if (ii) holds, then by replacing N with $\text{Hull}_{n+4}^N(\emptyset)$, we may assume $\rho_{n+4}^N = \omega$ and N is $(n+4)$ -sound. For the assertion in the second paragraph of the lemma $\text{r}\Pi_{n+4}$, given $\rho_{n+3} > \omega$. This uses that $\{p_{n+1}^N\}$ is an $\text{r}\Pi_{n+3}^N$ singleton. Also, $\{u_n^N\}$ is an $\text{r}\Pi_{n+2}^N$ singleton. We prove both of these facts by induction on n . Suppose that $\{u_n^N\}$ is $\text{r}\Pi_{n+2}^N$; we show that $\{p_{n+1}^N\}$ is $\text{r}\Pi_{n+3}^N$. By the induction hypothesis, we may use $u = u_n^N$ as a parameter. But p_{n+1}^N is the unique $p \in [\text{OR}(\mathfrak{C}_0(N))]^{<\omega}$ such that

- (a) p is $(n+1)$ -solid for N , and
- (b) $N = \text{Hull}_{n+1}^N(\rho_{n+1}^N \cup \{p\})$

(note that in both (a) and (b), u is an implicit parameter). Condition (a) is $\text{r}\Sigma_{n+2}^N(\{u\})$, because we only need to assert the existence of generalized solidity witnesses. And (b) holds iff for every $x \in \mathfrak{C}_0(N)$ there is $\alpha \in \text{OR}^N$ such that $x \in \text{Hull}_{n+1}^N(\{p, \alpha\})$ and there is a generalized $(n+1)$ -witness for $p \cup (\alpha + 1)$ (apply the latter condition with $x = p_{n+1}^N$ and note that $\alpha < \rho_{n+1}^N$); this is $\text{r}\Pi_{n+3}^N(\{u\})$. It follows easily that $\{u_{n+1}^N\}$ is also $\text{r}\Pi_{n+3}^N$.

²⁸Given N, n, x, θ, q as in 3.1, we say that q witnesses 3.1 with respect to (N, n, x, θ) .

For notational simplicity, we assume $n = 1$, but the proof easily generalizes. Let w_0 be the set of 2-solidity witnesses for p_2^N . Let $p \in (\rho_2^N)^{<\omega}$ be $<_{\text{lex}}$ -least such that

$$w_0, x \in \text{Hull}_2^N(\theta \cup \{p, p_2^N\}).$$

If $p = \emptyset$ then $q = \emptyset$ witnesses the lemma (by degree 2 condensation²⁹), so assume $p \neq \emptyset$. Let $\gamma = \max(p)$. Let

$$H_0 = \text{cHull}_2^N((\gamma + 1) \cup \{p_2^N\}).$$

We have $H_0 \in N$ since $\gamma < \rho_2^H$. Let $\pi : H_0 \rightarrow N$ be the uncollapse. Let $\gamma_1 = \text{cr}(\pi) = (\gamma^+)^{H_0}$. Let $\delta = \text{card}^N(\gamma) = \text{card}^N(\gamma_1) = \rho_2^{H_0}$. Let $R \leq N$ be least such that $\gamma_1 \leq \text{OR}^R$ and R projects to δ . For a theory t and parameters a, b , write $t_{a/b}$ for the theory resulting by replacing a with b .

Claim. *Let $t = \text{pTh}_2^N((\gamma + 1) \cup \{p_2^N\})$. Then $t_{p_2^N/\dot{p}}$ is $\text{r}\Sigma_2^R$, where \dot{p} is some constant symbol.*

Proof. Suppose (i) holds. Let \mathfrak{P} be the phalanx $((N, 1, \delta), (H_0, 1), \gamma_1)$ (see 1.1). Compare \mathfrak{P} with N , forming normal/1-maximal trees \mathcal{T} on \mathfrak{P} and \mathcal{U} on N . The details, including iterability and the analysis of the comparison, mostly follow the proof of solidity in [3, §8] and [19], using weak Dodd-Jensen as in [19].

We get that $b^\mathcal{T}$ is above H_0 and non-dropping (with $\deg^\mathcal{T}(b^\mathcal{T}) = 1$), and $M_\infty^\mathcal{T} \leq M_\infty^\mathcal{U}$. If $M_\infty^\mathcal{T} \triangleleft M_\infty^\mathcal{U}$ then $M_\infty^\mathcal{T}$ is sound, and it follows³⁰ that

$$M_\infty^\mathcal{T} = H_0 = R \triangleleft N,$$

which suffices. Otherwise let $Q = M_\infty^\mathcal{T} = M_\infty^\mathcal{U}$. Note that \mathcal{U} is non-trivial. In fact, $b^\mathcal{U}$ drops in model, since $\rho_2^Q \leq \delta < \rho_2^N$, and by 2.24.³¹ Moreover, letting $\alpha + 1 = \min(b^\mathcal{U} \setminus \{0\})$, $b^\mathcal{U}$ drops in model at $\alpha + 1$, but not in model or degree beyond there. So letting $R^* = M_{\alpha+1}^{*\mathcal{U}} \triangleleft N$, we have $\delta = \rho_2^{R^*} \leq \text{cr}(i_{R^*, Q}^\mathcal{U})$, so $R^* = R$; also $\deg^\mathcal{U}(b^\mathcal{U}) = 1$. So all $\text{r}\Sigma_2^Q$ subsets of δ are $\text{r}\Sigma_2^R$, which suffices.

Now suppose (ii) holds. Let $\mathfrak{P} = ((N, 3, \delta), (H_0, 1), \gamma_1)$. Compare \mathfrak{P} with N , again producing \mathcal{T} on \mathfrak{P} and \mathcal{U} on N , with \mathcal{U} being 3-maximal. Argue as in the previous case, but using the fact that N is $(n + 4)$ -sound and $\rho_{n+4}^N = \omega$, with standard fine structure, in place of weak Dodd-Jensen. \square (Claim)

Now $H_0 = \text{cHull}_2^N(\gamma_1 \cup \{p_2^N\})$ (recall $\gamma_1 = (\gamma^+)^{H_0}$), so H_0 is a 2-solidity witness for $p_2^N \cup \{\gamma_1\}$. Let $q_1 = p_2^N \cup \{\gamma_1\}$ and $w_1 = w \cup \{H_0\}$. Note that $R \cup \{R\} \subseteq \text{Hull}_1^N(\delta \cup \{\gamma_1\})$ by choice of R , so by the claim and choice of γ, γ_1 ,

$$x, w_1 \in \text{Hull}_2^N(\delta \cup \{q_1\}).$$

²⁹If (i) fails then we don't have the usual iterability assumptions for degree 2 condensation, but one can easily modify the proof thereof using the fine structural assumptions in (ii).

³⁰Here the traditional argument involves showing that $E_\alpha^\mathcal{T}$ is close to $M_{\alpha+1}^{*\mathcal{T}}$, which takes some extra work because we are iterating a phalanx. An alternative is to use 2.24 to deduce that if \mathcal{T} is non-trivial then $M_\infty^\mathcal{T}$ is not sound.

³¹In fact, $\rho_2^Q = \delta$, again either by closeness of extenders or by 2.24.

Let $p' \in [\text{OR}]^{<\omega}$ be $<_{\text{lex}}$ -least with

$$x, w_1 \in \text{Hull}_2^N(\theta \cup \{p', q_1\}).$$

So $p' \subseteq \delta$. If $p' = \emptyset$ we set $q = q_1$. Otherwise, repeat the preceding argument with q_1, w_1, p' in place of $q_0 = p_2^N, w_0, p$, and so on, producing

$$q = p_2^N \cup \{\gamma_1, \gamma_2, \dots, \gamma_k\},$$

with $\gamma_1 > \gamma_2 > \dots > \gamma_k$ (and k as large as possible). Let $M = \text{cHull}_2^N(\theta \cup \{q\})$ and $\pi : M \rightarrow N$ be the uncollapse. We have $\rho_\omega^M = \rho_2^M = \theta$ is a cardinal of N , M is sound and $M \in N$. Degree 2 condensation (arguing as in Footnote 29 in case (ii)) gives $M \triangleleft N$. $\square(3.1)$

Definition 3.2. Let N be an $(n+1)$ -sound premouse. We say that N is *n-reasonable* if every $E \in \mathbb{E}_+^N$ is Dodd sound and N satisfies the conclusion of 3.1 with respect to n . We say that N is *reasonable* iff N is ω -sound and n -reasonable for all n . \dashv

Note that n -reasonableness is first order, and a consequence of $(n+1)$ -soundness and $(n, \omega_1, \omega_1 + 1)$ -iterability. And by condition (ii) of 3.1, it is also a consequence of ω -soundness and $(\omega, \omega_1 + 1)$ -iterability, and therefore:

Corollary 3.3. Let N be a $(0, \omega_1 + 1)$ -iterable premouse. Then every proper segment of N is reasonable.

We now state the most central theorems of the paper. They share the same basic theme, but differ in certain details. The first was stated in the introduction:

Theorem 3.4. Let N be a $(0, \omega_1 + 1)$ -iterable premouse. Let $E \in N$ be such that $(N \upharpoonright \text{lh}(E), E)$ is a premouse and E is total over N . Then $E \in \mathbb{E}^N$.

Theorem 3.5. Let N be a $(0, \omega_1 + 1)$ -iterable premouse. Suppose there are cofinally many $\gamma < \text{OR}^N$ such that $N \upharpoonright \gamma$ is admissible. Suppose that if N has a largest cardinal κ then $N \models \text{"cof}(\kappa) \text{ is not measurable"}$.

Let $(0, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ be potential for N , with $E \in N$ and $\zeta < \omega$. Suppose that $b^{\mathcal{W}^+}$ does not drop in model. Then $E \in \mathbb{E}_+^P$.

Note that in 3.5 it is possible that $\text{lh}(E) = \text{OR}^P$, given that $[0, \zeta]_{\mathcal{V}}$ drops. In this case, the conclusion is that $E = F^P$.

Theorem 3.6. Let N be a premouse and $\omega < \eta < \text{OR}^N$ with η an N -cardinal and $N \models \text{"}\eta^{++} \text{ exists"}$. Let $R \in N$ be a reasonable premouse with $[R] = (\mathcal{H}_{\eta^+})^N$.³² Let $\mathfrak{c} \in \text{OR}^N$ and $\Sigma \in N$ ³³ with

$$N \models \text{"}\Sigma \text{ is an above-}\mathfrak{c}, (\omega, \eta^+)\text{-strategy for } R\text{"}.$$

Let $(\omega, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ be potential for R and such that:

³² $[M]$ denotes the universe of M .

³³Note that $\Sigma \subseteq N \upharpoonright (\eta^{++})^N$, so it makes sense to have $\Sigma \in N$.

- $\mathcal{W}, E \in N$ and \mathcal{W} is above η , of length $< (\eta^+)^N$ and via Σ ,
- $\mathfrak{c} < \text{lh}(E)$ and \mathfrak{c} is a cutpoint of $P \restriction \text{lh}(E)$,³⁴
- $\text{cr}(E) < \eta < \text{lh}(E)$ (therefore $b^{\mathcal{W}^+}$ does not drop).³⁵

Then $E \in \mathbb{E}_+^P$.

In both 3.6 above and 3.7 below, \mathcal{W} is above \mathfrak{c} because it is via Σ , but we do not assume $\mathfrak{c} \leq \text{cr}(E)$ explicitly (but $\mathfrak{c} \leq \text{cr}(E)$ follows from the conclusion that $E \in \mathbb{E}_+^P$). Note that in 3.6, although Σ is an above- \mathfrak{c} strategy and possibly $\mathfrak{c} < \eta$, we assume also that \mathcal{W} is above η , and since R has largest cardinal η , this is a serious restriction on \mathcal{W} . It ensures that \mathcal{W} is equivalent to a tree \mathcal{V} on some $R' \triangleleft R$, and $\mathcal{V} \in R$. However, 3.7 makes no such assumption.

Theorem 3.7. *Let N be a premouse and $\eta \geq \omega$ regular in N with $N \models “\eta^+ \text{ exists}”$. Let $R \in N$ be a reasonable premouse with $\lfloor R \rfloor = \mathcal{H}_\eta^N$. Let $\mathfrak{c} < \eta$ be a cutpoint of R . Suppose that if τ is the largest cardinal of R and $\mathfrak{c} \leq \text{cof}^N(\tau)$ then $\text{cof}^N(\tau)$ is not the critical point of any R -total $F \in \mathbb{E}^R$.*

Let $\Sigma \in N$ be such that $N \models “\Sigma \text{ is an above-}\mathfrak{c}, (\omega, \eta)\text{-strategy for } R”$. Let $(\omega, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ be potential for R with $\mathcal{W}, E \in N$ and \mathcal{W} via Σ , $\text{lh}(\mathcal{W}) < \eta$ and $\mathfrak{c} < \text{OR}^P$, and $b^{\mathcal{W}^+}$ non-dropping. Then $E \in \mathbb{E}_+^P$.

In the following two variants, the tree $\mathcal{W} \in N$, and \mathcal{W} is via Σ , but Σ is external, not assumed to be in N . The first is actually a corollary of the second, but it contains the main point and its statement is simpler, so we give it first:

Theorem 3.8. *Let N be a premouse and Σ a $(0, \theta^+ + 1)$ -strategy for N , where $\theta \geq \omega$. Let $\eta < \text{OR}^N$ be an N -cardinal. Let $(0, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ be potential for $N \restriction \eta$, such that $\mathcal{W}, E \in N$, and \mathcal{W} follows the strategy for $N \restriction \eta$ induced by Σ , $\text{lh}(\mathcal{W}) < \theta^+$, and $b^{\mathcal{W}^+}$ does not drop. Then $E \in \mathbb{E}_+^P$.*

Theorem 3.9 below is finer than the previous results. Hypothesis (d) of 3.9, which asserts that \mathcal{W} and E are suitably definable over N , will be made precise in 3.26.

Theorem 3.9. *Let $k < \omega$. Let N be a $(k+1)$ -sound k -reasonable premouse and $\mathfrak{c} \in \text{OR}$ be a cutpoint of N . Let $\theta \in \text{OR}$ and Σ be an above- $\mathfrak{c}, (k, \theta^+ + 1)$ -strategy for N .³⁶ Let $(k, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$ be potential for N , such that:*

- (a) \mathcal{W} is via Σ with $\text{lh}(\mathcal{W}) < \min(\theta^+, \rho_0^N)$,³⁷
- (b) $\mathfrak{c} < \text{lh}(E)$,
- (c) $b^{\mathcal{W}^+}$ does not drop, and $\text{cr}(E) < \rho_{k+1}^{\mathcal{W}}(M_v^{\mathcal{W}})$ where $v = \text{pred}^{\mathcal{W}^+}(\zeta + 1)$,
- (d) \mathcal{W} and E are $\mathfrak{r}\Sigma_{k+1}^N$ in the codes.³⁸

³⁴The hypothesis “ $\mathfrak{c} < \text{lh}(E)$ ” is redundant if \mathcal{W} is non-trivial, as \mathcal{W} is above \mathfrak{c} .

³⁵The hypothesis “ $\eta < \text{lh}(E)$ ” is redundant if \mathcal{W} is non-trivial, as \mathcal{W} is above η .

³⁶Actually if $\theta \geq \omega_1$ then a (k, θ^+) -strategy suffices.

³⁷ $\text{lh}(\mathcal{W}) < \rho_0^N$ will actually follow from the assumption that \mathcal{W} is $\mathfrak{r}\Sigma_{k+1}^N$ in the codes.

³⁸Cf. 3.26

Then $E \in \mathbb{E}_+^P$.

Note that in 3.9, we might have $\text{lh}(E) = \text{OR}^P$ even with $[0, \zeta]_{\mathcal{W}}$ non-dropping. By 2.24, the assumptions of 3.9 imply $\rho_{k+1}(M_v^{\mathcal{W}}) = \sup i_{0,v}^{\mathcal{W}} \rho_{k+1}^N$.

Remark 3.10. The literal generalization of these results to “premise” M with enough extenders of superstrong type on \mathbb{E}_+^M fails. For suppose E is a total type 2 extender on \mathbb{E}^M , $U = \text{Ult}(M, E)$, $\kappa = \text{lgcd}(M|\text{lh}(E))$, $F \in \mathbb{E}_+^U$, $\kappa = \text{cr}(F)$, F is of superstrong type and U -total. Then $\text{Ult}(M|\text{lh}(E), F)$ and $U|\text{lh}(F)$ are distinct active premice, with the same reduct. If M is iterable and $M \models \text{ZF}^-$, then the other hypotheses of 3.9 also hold.

This doesn’t appear to be a strong failure of 3.9, however, since both extenders are on the extender sequence of normal iterates of M . (Also, in Jensen’s λ -indexing, these extenders are not indexed at the same point.)

In the proof of 3.6 we will need to consider ostensibly illfounded ultrapowers, and fine structural embeddings between them. Toward this we define generalizations of the usual fine structural notions for such possibly illfounded “premise”, at least to the extent that these notions are well defined. The definitions and calculations are almost the usual ones; however when $U = \text{Ult}_m(M, E)$ is illfounded there is a subtlety in the definition of ρ_m^U , T_m^U and $\text{r}\Sigma_{m+1}^U$, as in this case it seems possible that the natural candidate for ρ_m^U is a proper segment of OR^U which is however not a cut “in” U , i.e. there is no $\rho \in \text{OR}^U$ such that

$$\rho_m^U = \{\alpha \in U \mid U \models \alpha \in \rho\}.$$

So we give a detailed description of things in order to handle this issue. ³⁹

Definition 3.11. Let $N = ([N], \mathbb{E}, F)$ be a structure in the premouse language, possibly illfounded. Suppose that $N \models$ “I am a premouse” and N has wellfounded ω ; so N is correct about what formulas are. We define the fine structure of N as far as possible. Write

$$\text{OR}^N = \{\alpha \in [N] \mid N \models \alpha \in \text{OR}\}.$$

Suppose first that $N \models$ “I am not type 3”. Define $\rho_0^N = \text{OR}^N$. We say that N is 0-sound and 0-feasible. Define $\text{r}\Sigma_1^N$ as usual, and for $X \subseteq N$, define $\text{Hull}_1^N(X)$ as usual: the set of all $y \in N$ such that for some Σ_1 formula φ and $\vec{x} \in X^{<\omega}$, y is the unique $y' \in N$ such that $N \models \varphi(\vec{x}, y')$.

Define ρ_1^N as the set of all $\alpha \in \text{OR}^N$ such that either $\alpha < \omega$ or for all $p \in N$, we have $\text{Th}_1^N(\alpha \cup \{p\}) \in N$. Note that ρ_1^N is an \in^N -initial segment of OR^N , but we might have $\rho_1^N \subsetneq \text{OR}^N$ but ρ_1^N not “in” N . Given $p \in [\text{OR}^N]^{<\omega}$, we

³⁹In an earlier version of this paper, the author had not noticed the issue with ρ_m^U and simply claimed that there was no problem in adapting the fine structure to illfounded structures, and omitted any discussion thereof. A question from the anonymous referee regarding the topic lead to the author’s noticing the issue.

define the 1-*solidity* of p for N as usual. We say that N is 1-*sound* iff there is $p \in [\text{OR}^N \setminus \rho_1^N]^{<\omega}$ such that p is 1-solid for N and $N = \text{Hull}_1^N(\rho_1^N \cup \{p\})$.

Suppose N is 1-sound. Define p_1^N to be the unique p witnessing this, and define u_1^N as usual. (Uniqueness: Suppose $p <_{\text{lex}}^N q$ are both such. By the 1-solidity of q , there is a “limit ordinal” $\alpha \in \text{OR}^N$ such that “ $\rho_1^N \subseteq \alpha$ ” and $\text{Th}_1^N(\alpha \cup \{p\}) \in N$. But $N = \text{Hull}_1^N(\alpha \cup \{p\})$, and the usual diagonalization argument then constructs a new subset of $\alpha^{<\omega}$, a contradiction.) Define T_1^N as usual; that is, the set of all $t = \text{Th}_1^N(\alpha \cup \{q\})$ for some $\alpha \in \rho_1^N$ and $q \in N$. So $T_1^N \subseteq N$. Define $\text{r}\Sigma_2^N$ and $\text{Hull}_2^N(X)$ from T_1^N as usual. (So $u_1^N \in \text{Hull}_2(X)$ by definition.)

We say that N is 1-*feasible* iff N is 1-sound and either $\rho_1^N = \text{OR}^N$ or ρ_1^N is “in” N . If N is n -feasible where $n \geq 1$ and $\rho_n^N > \omega$, define ρ_{n+1}^N , $(n+1)$ -solidity and $(n+1)$ -soundness as above. If N is $(n+1)$ -sound define $p_{n+1}^N, u_{n+1}^N, T_{n+1}^N, \text{r}\Sigma_{n+2}^N, \text{Hull}_{n+2}^N$ (note these definitions do not require $(n+1)$ -feasibility) and also $(n+1)$ -feasibility as above.

If N is instead type 3 then these things are adapted in the obvious manner to $\mathfrak{C}_0(N) = N^{\text{sq}}$. (Here N^{sq} is well-defined as $\nu(F^N)$ is well-defined by the premouse axioms.)

Let M, N be as above and $\pi: \mathfrak{C}_0(M) \rightarrow \mathfrak{C}_0(N)$. We say that π is a *virtual (weak/near) m -embedding* given the usual conditions on π , using the parameters etc defined as above. For example, for virtual weak m :⁴⁰

- M, N are m -sound,
- for $k \leq m$ we have $\pi(p_k^M) = p_k^N$,
- for $k < m$ we have either
 - $\rho_k^M = \text{OR}(\mathfrak{C}_0(M))$ and $\rho_k^N = \text{OR}(\mathfrak{C}_0(N))$, or
 - $\rho_k^M < \text{OR}(\mathfrak{C}_0(M))$ and $\pi(\rho_k^M) = \rho_k^N$,
- π is $\text{r}\Sigma_m$ -elementary,
- there is a cofinal $X \subseteq \rho_m^M$ such that π is $\text{r}\Sigma_{m+1}$ -elementary on $\text{Hull}_{m+1}^M(X)$.

Because M, N are m -sound (hence $(m-1)$ -feasible if $m > 0$) we have defined all the fine structural notions needed to make sense of these requirements.

We also extend the definition of *reasonable* to possibly illfounded structures N as above, in the obvious fashion. \dashv

Definition 3.12. Let P, Q be active premice and $\sigma: \mathfrak{C}_0(P) \rightarrow \mathfrak{C}_0(Q)$ a weak 0-embedding. Then $\psi_\sigma: \text{Ult}_0(P, F^P) \rightarrow \text{Ult}_0(Q, F^Q)$ denotes the map induced by σ through the Shift Lemma. (Note $\sigma \subseteq \psi_\sigma$.) \dashv

Lemma 3.13. Let $m \leq n \leq \omega$. Let M, N be premice, M is m -sound, N is n -sound, $\pi: M \rightarrow N$ a weak m -embedding. Let P, Q be active premice,

⁴⁰Cf. the definition of weak m -embedding in §1.1.

$\sigma : P \rightarrow Q$ a weak 0-embedding. Let $E = F^P$, $F = F^Q$, $\kappa = \kappa_E$, $\mu = \kappa_F$. Suppose

$$M \models (\kappa^+)^M = P \models (\kappa^+)^P \text{ and } N \models (\mu^+)^N = Q \models (\mu^+)^Q$$

and $\pi \restriction (\kappa^+)^M \subseteq \sigma$. Suppose $\kappa < \rho_m^M$ and $\mu < \rho_n^M$. Let

$$U = \text{Ult}_m(M, E) \text{ and } W = \text{Ult}_n(N, F)$$

(U, W might be illfounded). Let $i = i_E^{M,m}$ and $j = i_F^{N,n}$ and $\psi : U \rightarrow W$ be the Shift Lemma map.

Then U is m -sound, W is n -sound, i a virtual m -embedding, j a virtual n -embedding, ψ a virtual weak m -embedding, $\psi \circ i = j \circ \pi$,

$$\psi_\sigma \restriction (\text{OR}^P + 1) \subseteq \psi.$$

Moreover, if M is κ -sound and $\rho_{m+1}^M \leq \kappa$ then

$$U = \text{Hull}_{m+1}^U(\nu(E) \cup \{i(p_{m+1}^M)\}).$$

Proof Sketch. For simplicity we assume $m = 2$ and M, N are not type 3. Note that (U could be illfounded and) ρ_2^U might not be “in” U .

To verify that U is 2-sound and i is a virtual 2-embedding, one proves:

1. $\text{r}\Sigma_0$ - and $\text{r}\Sigma_1$ -Los theorem holds. Hence i is $\text{r}\Sigma_1$ -elementary.
2. Let $b \in [\nu(E)]^{<\omega}$ and

$$f, g : [\kappa]^{|b|} \rightarrow M$$

be $\text{r}\Sigma_2^M$ with $\text{rg}(g) \subseteq \rho_1^M$. Define

$$h : [\kappa]^{|b|} \rightarrow M,$$

$$h(u) = \text{Th}_1^M(g(u) \cup \{f(u)\}).$$

Note that h is $\text{r}\Sigma_2^M$. Let $\tilde{f} = [f, b]_E^{M,2}$ and likewise \tilde{g}, \tilde{h} . Then

$$U \models “\tilde{h} = \text{Th}_1(\tilde{g} \cup \{\tilde{f}\})”.$$

3. If $\rho_1^M = \text{OR}^M$ then $\rho_1^U = \text{OR}^U$, and if $\rho_1^M < \text{OR}^M$ then $i(\rho_1^M) \leq \rho_1^U$.
4. $i(p_1^M)$ is 1-solid for U , as witnessed by $i(w_1^M)$ (w_1^M is the set of 1-solidity witnesses for M).
5. $U = \text{Hull}_1^U(i(\rho_1^M) \cup \{i(p_1^M)\})$.
6. U is 1-sound and 1-feasible with $p_1^U = i(p_1^M)$ and $u_1^U = i(u_1^M)$ and either $\rho_1^M = \text{OR}^M$ and $\rho_1^U = \text{OR}^U$, or $\rho_1^M < \text{OR}^M$ and $\rho_1^U = i(\rho_1^M)$ (use diagonalization to see $\rho_1^U \leq i(\rho_1^M)$).
7. $\text{r}\Sigma_2$ -Los theorem holds. Hence i is $\text{r}\Sigma_2$ -elementary.

8. Let $f : [\kappa]^{|b|} \rightarrow M$ be $\text{r}\Sigma_2^M(\{q\})$ and $\kappa \leq \alpha < \rho_2^M$. Let $t = \text{Th}_2^M(\alpha \cup \{q\})$. Define $h : [\kappa]^{|b|} \rightarrow M$ by $h(u) =$ the $\text{r}\Sigma_2$ -theory in parameters $\alpha \cup \{f(u)\}$ determined by t and substitution. Then h is $\text{r}\Sigma_2^M(\{(q, t)\})$ and

$$U \models "h = \text{Th}_2(i(\alpha) \cup \{\tilde{f}\})".$$

9. $\sup i \rho_2^M \leq \rho_2^U$.
10. $i(p_2^M)$ is 2-solid for U , as witnessed by $i(w_2^M)$.
11. Let $b \in [\nu(E)]^{<\omega}$ and φ be $\text{r}\Sigma_2$ and $q \in M$ be such that $\varphi(q, \cdot, \cdot)$ defines a function $f : [\kappa]^{|b|} \rightarrow M$ over M . Let $\tilde{f} = [f, b]_E^{M,2}$. Then

$$U \models "\tilde{f} \text{ is the unique } y \text{ such that } \varphi(i(q), b, y)".$$

12. $U = \text{Hull}_2^U(\text{rg}(i) \cup \nu(E)) = \text{Hull}_2^U(i \rho_2^M \cup \{i(p_2^M)\} \cup \nu(E))$.
13. U is 2-sound with $p_2^U = i(p_2^M)$ and $i(u_2^M) = u_2^U$ and $\rho_2^U = \sup i \rho_2^M$ (but maybe U is not 2-feasible).
14. i is $\text{r}\Sigma_3$ -elementary; this follows from properties 8, 12 and $\text{r}\Sigma_1$ -elementarity, just like when U is wellfounded.

Similarly, $j : N \rightarrow W$ is also a virtual n -embedding, and the $\text{r}\Sigma_n^N$ -Los theorem holds, etc. The usual calculations using all these facts (including $\text{r}\Sigma_2$ -Los theorem for both ultrapowers), now give that ψ is a virtual weak 2-embedding, commutativity holds etc.

Finally suppose that M is κ -sound and $\rho_3^M \leq \kappa$. So $M = \text{Hull}_3^M(\kappa \cup \{p_3^M\})$. By the $\text{r}\Sigma_3^M$ -elementarity of i ,

$$\text{rg}(i) = \text{Hull}_3^U(\kappa \cup \{i(p_3^M)\}).$$

But $U = \text{Hull}_2^U(\text{rg}(i) \cup \nu(E))$ (as above), so $U = \text{Hull}_3^U(\nu(E) \cup \{i(p_3^M)\})$. \square

Lemma 3.14. Let $\pi : \mathfrak{C}_0(M) \rightarrow \mathfrak{C}_0(N)$ be a virtual weak k -embedding. Let φ be $\text{r}\Sigma_{k+1}$ and $x \in \mathfrak{C}_0(M)$. If $\mathfrak{C}_0(M) \models \varphi(x)$ then $\mathfrak{C}_0(N) \models \varphi(\pi(x))$.

Proof. If $k = 0$ this is immediate. If $k > 0$ it holds because $\pi "T_k^M \subseteq T_k^N$, which holds just as for a weak k -embedding.⁴¹ \square

Proof of 3.6. We argue overall as in the proof of the ISC in [3, §10]. First, for motivation, make the following simplifying assumptions: that $R = N|(\eta^+)^N$, \mathcal{W} is trivial, so E coheres \mathbb{E}^N , and that we can compare $\text{Ult}(N, E)$ with N , producing trees \mathcal{U}, \mathcal{V} such that $M_\infty^{\mathcal{U}} = M_\infty^{\mathcal{V}}$, $b^{\mathcal{U}}$ and $b^{\mathcal{V}}$ do not drop, $i^{\mathcal{U}} \circ i_E = i^{\mathcal{V}}$, and $\nu_E \leq \text{cr}(i^{\mathcal{U}})$. Then standard arguments using the ISC show that E is used in \mathcal{V} , and $E \in \mathbb{E}^N$, as required.

⁴¹That is, with *weak k -embedding* defined as in §1.1.

Now drop the above simplifying assumptions. We don't even know, for instance, that $\text{Ult}(N, E)$ is wellfounded. So we will instead arrange a situation similar to that in the preceding paragraph, but with R replaced by a certain hull M , which will be $k+1$ -sound for some k , with $\rho_{k+1}^M = \omega$. Denoting the resulting collapses of E , \mathcal{W} , etc, by \bar{E} , $\bar{\mathcal{W}}$, etc, and letting $\iota = \text{lged}(P \upharpoonright \text{lh}(E))$, we will show that the phalanx $\mathfrak{P} = \Phi(\bar{\mathcal{W}}, \bar{\iota}, \bar{E})$ (see 2.37) is sufficiently iterable in N to allow comparison with M . If $\bar{\iota} = \nu(\bar{E})$ (that is, E is type 1 or 3), then the proof is completed as in the previous paragraph. In general we will rely on the Dodd structure analysis from §2, appealing to 2.38.

We now commence with the details. Because \mathcal{W} is above η , it is equivalent to a tree \mathcal{V} on some $R' \triangleleft R$ such that $\rho_{\omega}^{R'} = \eta$. Note then that $\mathcal{V}, E \in R$. Let $k < \omega$ be large enough to reflect the facts about $x = (\mathcal{V}, E, \mathfrak{c})$ into the model M defined below. Now by assumption R is reasonable. So we can fix $q \in R^{<\omega}$ witnessing 3.1 with respect to $(R, k, x, \theta = \omega)$. Let

$$M = \text{cHull}_{k+1}^R(\{q\})$$

and $\pi : M \rightarrow R$ be the uncollapse. So M is sound, reasonable, $\rho_{k+1}^M = \omega$, $p_{k+1}^M = \pi^{-1}(q)$, and π is a near k -embedding. Since $R \models \text{ZFC}^-$ and $\pi, \Sigma \in N$ and $\omega_1^N < (\eta^+)^N$,

$$N \models \text{"}M \text{ is above-}\bar{\mathfrak{c}}, (\omega, \omega_1 + 1)\text{-iterable"}$$

Let $\bar{\mathcal{V}} = \pi^{-1}(\mathcal{V})$, etc. Let $\bar{\mathcal{W}}$ be the same tree except with $M_0^{\bar{\mathcal{W}}} = M$ (we have $M_0^{\bar{\mathcal{V}}} \triangleleft M$). So $(\omega, \bar{\mathcal{W}}, \bar{\zeta}, \bar{P}, \bar{E}, \bar{\mathcal{W}}^+)$ is potential for M and $\bar{\mathfrak{c}}$ is a cutpoint of $\bar{P} \upharpoonright \text{lh}(\bar{E})$. Let

$$\bar{U} = \text{Ult}_k(M, \bar{E}) = M_{\infty}^{\bar{\mathcal{W}}^+}.$$

We will show that \bar{U} is wellfounded, and define the phalanx $\mathfrak{P} = \Phi(\bar{\mathcal{W}}, \bar{\iota}, \bar{E})$. (See 2.37. Note that \mathfrak{P} has last model \bar{U} , at degree k .)

Claim. \bar{U} is wellfounded and $N \models \text{"}\mathfrak{P} \text{ is } (\omega_1 + 1)\text{-iterable"}$.

Proof. We first find $M' \triangleleft P \upharpoonright \text{lh}(E)$ and a virtual weak k -embedding $\psi' : \bar{U} \rightarrow M'$ such that $\psi' \in N$ and $\pi \upharpoonright (\max(\bar{\iota}, \bar{\mathfrak{c}}) + 1) \subseteq \psi'$. It follows that \bar{U} is wellfounded, and so ψ' is in fact a weak k -embedding. Working in N , we will then lift normal trees \mathcal{T} on \mathfrak{P} to above- \mathfrak{c} , normal trees \mathcal{U} on $\Phi(\mathcal{W})$. Since \mathcal{W} is via $\Sigma \in N$, this suffices. We will use π, ψ' , which are in N , as initial copying maps, using π to lift the models of $\Phi(\bar{\mathcal{W}})$ to (some) models of $\Phi(\mathcal{W})$ and ψ' to lift \bar{U} to M' .

Let $U = \text{Ult}_0(R, E) = \text{Ult}_{\omega}(R, E)$. (We don't assume U is wellfounded.) Let $j = i_E^R$, $\bar{j} = i_{\bar{E}}^{M, k}$ and $\psi : \bar{U} \rightarrow U$ be the Shift Lemma map. By 3.13 and in the sense of 3.11, \bar{U} is k -sound, U is fully sound, ψ is a virtual weak k -embedding, $\psi \circ \bar{j} = j \circ \pi$ and $\pi \upharpoonright (\text{lh}(\bar{E}) + 1) \subseteq \psi$.

By elementarity, U is reasonable (in the sense of 3.11). Let $x' = (j(q), \nu(E), \mathfrak{c})$ and $q' \in U^{<\omega}$ "witness 3.1" with respect to (U, k, x', ι) (in the obvious sense). Let

$$M' = \text{cHull}_{k+1}^U(\iota \cup q')$$

and let $\pi' : M' \rightarrow U$ be the uncollapse, so π' is a virtual near k -embedding. Then $M' \triangleleft U \upharpoonright \text{lh}(E)$ since $\text{lh}(E)$ is a cardinal in U , so $M' \triangleleft P \upharpoonright \text{lh}(E)$ and M' is wellfounded. Moreover,

$$(\nu(E) + 1) \cup \{\mathfrak{c}, j(q)\} \subseteq \text{rg}(\pi').$$

We claim that $\text{rg}(\psi) \subseteq \text{rg}(\pi')$. First let us verify that $\text{rg}(\psi \circ \bar{j}) \subseteq \text{rg}(\pi')$. Let $y \in M$. Let φ be $\text{r}\Sigma_{k+1}$ with $y = \text{m}\tau_\varphi^M(\bar{q})$ (see 2.23). Then by commutativity and elementarity of j ,

$$\psi(\bar{j}(y)) = j(\pi(y)) = \text{m}\tau_\varphi^U(j(q)) \in \text{rg}(\pi'),$$

as desired. But

$$\bar{U} = \text{Hull}_k^{\bar{U}}(\text{rg}(\bar{j}) \cup \nu(\bar{E}))$$

(see the proof of 3.13) and ψ is $\text{r}\Sigma_k$ -elementary and $\psi \upharpoonright \nu(\bar{E}) \subseteq \nu(E) \subseteq \text{rg}(\pi')$, which suffices.

It follows that $\psi' : \bar{U} \rightarrow M'$ is a virtual weak k -embedding where

$$\psi'(x) = (\pi')^{-1}(\psi(x)).$$

But then \bar{U} is wellfounded, so ψ' is in fact a weak k -embedding.

So we have M', ψ' as desired. We now lift trees \mathcal{T} on \mathfrak{P} to \mathcal{U} on $\Phi(\mathcal{W})$, via π, ψ' . The details are mostly standard; we just make a couple of key points.

Note $\pi \upharpoonright (\max(\bar{\iota}, \bar{\mathfrak{c}}) + 1) \subseteq \psi'$ and

$$\psi'(\bar{\iota}, \bar{\mathfrak{c}}) = \iota, \mathfrak{c} < \psi'(\text{lh}(\bar{E})) < \text{lh}(E) = \pi(\text{lh}(\bar{E})).$$

For the models $M_\alpha^{\bar{\mathcal{W}}}$ we use copy map

$$\pi \upharpoonright M_\alpha^{\bar{\mathcal{W}}} : M_\alpha^{\bar{\mathcal{W}}} \rightarrow M_{\pi(\alpha)}^{\mathcal{W}},$$

and for $M_0^{\mathcal{T}} = \bar{U}$ we use

$$\pi_0 = \psi' : \bar{U} \rightarrow M' \triangleleft M_\zeta^{\mathcal{W}} = M_0^{\mathcal{U}}.$$

For $\alpha > 0$ we then proceed to define $\pi_\alpha : M_\alpha^{\mathcal{T}} \rightarrow M_\alpha^{\mathcal{U}}$ inductively as usual. Now $\text{lh}(\bar{E}) < \text{lh}(E_0^{\mathcal{T}})$, so

$$\iota, \mathfrak{c} < \psi'(\text{lh}(\bar{E})) < \text{lh}(E_0^{\mathcal{U}}) \leq \text{OR}^{M'} < \text{lh}(E),$$

and $\psi' \upharpoonright (\text{lh}(\bar{E}) + 1) \subseteq \pi_\alpha$ for all $\alpha \geq 0$. It follows that if $\text{cr}(E_\alpha^{\mathcal{T}}) < \bar{\iota}$ then $\text{root}^{\mathcal{U}}(\alpha + 1) = \pi(\text{root}^{\mathcal{T}}(\alpha + 1))$ and \mathcal{T}, \mathcal{U} agree about drops and degree at $\alpha + 1$, and if there is a drop in model then $\pi(M_{\alpha+1}^{\mathcal{T}}) = M_{\alpha+1}^{\mathcal{U}}$. If $\text{cr}(E_\alpha^{\mathcal{T}}) = \bar{\iota}$ then $M_{\alpha+1}^{\mathcal{T}} = \bar{U}$ and $M_{\alpha+1}^{\mathcal{U}} = M'$ and $\deg^{\mathcal{T}}(\alpha + 1) = \deg^{\mathcal{U}}(\alpha + 1) = k$.⁴²

Because $\bar{\mathfrak{c}} < \text{lh}(\bar{E})$ is a cutpoint of $\bar{P} \upharpoonright \text{lh}(\bar{E})$, we get that $\bar{\mathfrak{c}}$ is a cutpoint of $M_\alpha^{\mathcal{T}}$ for all α just as at the start of the proof of 2.38. It follows that \mathcal{U} is above \mathfrak{c} , so we can use Σ for forming \mathcal{U} . \square

⁴²If we had used $\bar{\iota} + 1$ in place of $\bar{\iota}$ as an exchange ordinal in defining \mathfrak{P} , and $\bar{\iota} = \text{cr}(E_\alpha^{\mathcal{T}})$, then $P' = M_{\alpha+1}^{\mathcal{T}} \triangleleft \bar{P}$ and $M_{\alpha+1}^{\mathcal{U}} = M'$. We have not constructed appropriate embeddings for lifting such P' to M' , so the copying process would break down.

Now since M is above- \bar{c} , $(\omega, \omega_1 + 1)$ -iterable in N , the claim above combined with Lemma 2.38 implies $\bar{E} \in \mathbb{E}_+^P$, so $E \in \mathbb{E}_+^P$, completing the proof.⁴³ \square

The proofs of the variants are similar, proceeding by defining an iterable phalanx \mathfrak{P} and appealing to Lemma 2.38. So we just give a sketch.

Proof of 3.4. We may assume N is passive, and note that every extender in \mathbb{E}^N is Dodd sound and every proper segment of N is reasonable. Let $\gamma < \text{OR}^N$ be such that E is definable over $N|\gamma$ (so $\text{lh}(E) \leq \gamma$). We have $\text{cr}(E) < \rho_\omega^{N|\gamma}$ as E is N -total. Using 3.1 let $k < \omega$ and $q \in [\gamma]^{<\omega}$ be such that $M = \text{cHull}_{k+1}^{N|\gamma}(\{q\})$ is sound (and $\rho_{k+1}^M = \omega$) and the facts about E reflect to an extender \bar{E} defined similarly over M . We have $\text{cr}(\bar{E}) < \rho_k^M$. Let $\bar{U} = \text{Ult}_k(M, \bar{E})$ and $\bar{c} = \text{lgcd}(M || \text{lh}(\bar{E}))$ and define the phalanx

$$\mathfrak{P} = ((M, k, \bar{c}), (\bar{U}, k), \text{lh}(\bar{E})).$$

It suffices to see that \mathfrak{P} is $(\omega_1 + 1)$ -iterable, as then we can use 2.38. But we have the Shift Lemma map $\psi : \bar{U} \rightarrow i_{E|}^{N,0}(N|\gamma)$, and we can find $M' \triangleleft N || \text{lh}(E)$ and a suitable weak k -embedding $\psi' : \bar{U} \rightarrow M'$ as in the preceding proof. \square

Proof of 3.5. We may assume that N is countable with largest cardinal κ . Since $E \in N$, $\text{lh}(E) < \text{OR}^N$, so we may also assume N is passive. Let λ be such that $\kappa < \lambda < \text{OR}^N$ and $E \in N|\lambda$ and $\rho_\omega^{N|\lambda} = \kappa$ and $\text{cof}^N(\kappa) = \text{cof}^{N|\lambda}(\kappa)$. Let γ be least $> \lambda$ such that $N|\gamma$ is admissible. We have \mathcal{W} on N and $\text{lh}(\mathcal{W}) < \omega$. We claim that there is a 0-maximal tree \mathcal{V} on $N|\gamma$ using the same extenders (with the same indices) as does \mathcal{W} , and moreover, \mathcal{W}, \mathcal{V} have the same tree, drop and degree structure, and letting $l = \{\kappa, \text{lh}(E_0^\mathcal{W}), \dots, \text{lh}(E_{\zeta-1}^\mathcal{W})\}$, then for $m < \text{lh}(\mathcal{W})$,

- $M_m^\mathcal{V}$ is a $\Delta_1^{N|\gamma}(l)$ -definable transitive class of $N|\gamma$ and:
- if $[0, m]_\mathcal{W}$ does not drop then $M_m^\mathcal{V} = i_{0m}^\mathcal{W}(N|\gamma)$ has height γ , and
- if $[0, m]_\mathcal{W}$ drops then $M_m^\mathcal{V} = M_m^\mathcal{W} \in N|\gamma$.

For suppose $[0, m]_\mathcal{W}$ does not drop. Then $i_{0m}^\mathcal{W}$ is continuous at κ and $i_{0m}^\mathcal{W}(N|\kappa) = \text{Ult}_0(N|\kappa, F)$ where F is the branch extender, since $N \models \text{“cof}(\kappa) \text{ is not measurable”}$. Therefore if $\rho_\omega^{N|\alpha} = \kappa$ then $i_{0m}^\mathcal{W}(N|\alpha)$ is determined by $i_{0m}^\mathcal{W}(t)$ where $t \subseteq \kappa$ is the appropriate theory, and $i_{0m}^\mathcal{W}(t) = \bigcup_{\beta < \kappa} i_{0m}^\mathcal{W}(t \restriction \beta)$. This gives a $\Delta_1^{N|\gamma}(l)$ description of the function $N|\alpha \mapsto i_{0m}^\mathcal{W}(N|\alpha)$ for such $\alpha < \gamma$. Therefore $M_m^\mathcal{V}$ is $\Delta_1^{N|\gamma}(l)$. Using the admissibility of $N|\gamma$, it follows that $\text{OR}(M_m^\mathcal{V}) = \gamma$ and $M_m^\mathcal{V}$ is admissible. So $M_m^\mathcal{V} = i_{0m}^\mathcal{V}(N|\gamma)$ by the minimality of γ .

Now considerations as above show that if the conclusion of the theorem fails, the failure is a first-order sentence satisfied by $N|\gamma$ (taking l, E to be minimal

⁴³In 2.38, $b^{\mathcal{W}^+}$ is allowed to drop, but we needed $b^{\mathcal{W}^+}$ to be non-dropping in order to prove that \mathfrak{P} is iterable.

counterexamples with respect to $N|\gamma$ and with properties as above, then they also yield a counter-example with respect to N). So we may assume that l, E are minimal. Fix $0 < k < \omega$ such that the failure and all properties established reflect into $M = \text{cHull}_{k+1}^{N|\gamma}(\emptyset)$. Let $\pi : M \rightarrow N|\gamma$ be the uncollapse. Write $\pi(\bar{E}) = E$ etc.

Now $\bar{\mathcal{V}}$ is a 0-maximal tree on M , but is in fact equivalent to a k -maximal tree. For $\rho_1^M = \rho_k^M = \bar{\kappa}$, so it suffices to see that the relevant $\text{r}\Sigma_k^M$ functions with range $\subseteq \bar{\kappa}$ are bounded on measure one sets. So let $\mu < \bar{\kappa}$ and $F \in \mathbb{E}^M$ be M -total with $\text{cr}(F) = \mu$. Let $b \in [\nu(F)]^{<\omega}$. Let $f : [\mu]^{|b|} \rightarrow \bar{\kappa}$ be an $\text{r}\Sigma_k^M$ function. We want $A \in E_b$ such that $f \restriction A$ is bounded in $\bar{\kappa}$. If κ is singular in N then $\bar{\kappa}$ is singular in M with cofinality $\neq \mu$, which easily gives such an A . Suppose κ is regular in N . Let $\pi(f)$ be the $\text{r}\Sigma_k^{N|\gamma}$ -function defined over $N|\gamma$ as f is over M . Because π is a near k -embedding and $\pi(\bar{\kappa}) = \kappa = \rho_k^{N|\gamma}$, we have $\pi(f) : [\pi(\mu)]^{|b|} \rightarrow \kappa$, but $\pi(f) \in N$, so $\pi(f)$ is bounded in κ . But the fact that there is a bound on $\pi(f)$ is an $\text{r}\Sigma_{k+1}$ assertion about the relevant parameters, and therefore f is bounded in $\bar{\kappa}$, so $A = \mu^{|b|}$ suffices.

Now let $\bar{\mathcal{V}}^+ = \bar{\mathcal{V}} \hat{\ } \langle E \rangle$ as a k -maximal potential tree. Let $\bar{U} = M_\infty^{\bar{\mathcal{V}}^+}$. As in the proof of 3.6 we obtain a suitable lifting map $\psi' : \bar{U} \rightarrow M'$ with $M' \triangleleft M_\infty^{\mathcal{V}} \parallel \text{lh}(E)$. (Let $\mathcal{W}^+ = \mathcal{W} \hat{\ } \langle E \rangle$ and $\mathcal{V}^+ = \mathcal{V} \hat{\ } \langle E \rangle$, both 0-maximal. As above, $M_\infty^{\mathcal{V}^+} = i_{0\infty}^{\mathcal{W}^+}(N|\gamma)$ and $i_{0\infty}^{\mathcal{V}^+} = i_{0\infty}^{\mathcal{W}^+} \restriction N|\gamma$. So $i_{0\infty}^{\mathcal{V}^+}$ is fully elementary and we can define M' as a hull of $M_\infty^{\mathcal{V}^+}$.) Define the phalanx $\mathfrak{P} = \Phi(\bar{\mathcal{V}}, \bar{\iota}, \bar{U})$, where $\bar{\iota} = \text{lgcd}(\bar{P} \parallel \text{lh}(E))$, and show that \mathfrak{P} is iterable in V , contradicting 2.38. \square

Proof of 3.7. Because η is regular in N and by our assumptions about $\text{cof}^R(\tau)$ (when R has a largest cardinal τ), there is $\gamma < \eta$ and an ω -maximal tree \mathcal{V} on $N|\gamma$ equivalent to \mathcal{W} . (Choose γ so that: For $\alpha < \text{lh}(\mathcal{W})$, if $[0, \alpha]_{\mathcal{W}}$ drops let $Q_\alpha = M_\alpha^{\mathcal{W}}$ and otherwise let $Q_\alpha = i_{0\alpha}^{\mathcal{W}}(N|\gamma)$; then $M_\alpha^{\mathcal{V}} = Q_\alpha$ and the factor map $\sigma_\alpha : M_\alpha^{\mathcal{V}} \rightarrow Q_\alpha$ is just the identity.) Because $\text{lh}(\mathcal{W}) < \eta$, we have $\mathcal{V}, E \in R$. We set $M = \text{cHull}_{k+1}^R(\{q\})$ with an appropriate q, k , and $\pi : M \rightarrow R$ the uncollapse with $\mathcal{V}, E \in \text{rg}(\pi)$. The rest is much like before. \square

Proof of 3.8. This is basically like for 3.6, but the iterability is established in V , not in N . Let $\gamma < \text{OR}^N$ with $\mathcal{W}, E \in \mathcal{J}(N|\gamma)$. Let $x \in N|\gamma$ with \mathcal{W}, E definable over $N|\gamma$ from x . Let k be sufficiently large and let $q \in N|\gamma$ witness 3.1 with respect to $(N|\gamma, k, x, \theta = \omega)$. Let

$$M = \text{cHull}_{k+1}^{N|\gamma}(\{q\})$$

and $\pi : M \rightarrow N|\gamma$ be the uncollapse. Let $\pi(\bar{x}) = x$ etc. Let $\bar{\mathcal{W}}, \bar{E}$, defined over M from \bar{x} , be the preimages of \mathcal{W}, E . Let $\bar{\mathcal{X}}$ be the k -maximal tree on $N|\gamma$ determined by \mathcal{W} ; so $\bar{\mathcal{X}}$ is via the strategy induced by Σ . Similarly, let $\bar{\mathcal{X}}^+$ be the k -maximal tree on M determined by $\bar{\mathcal{W}}$. So $\Phi(\bar{\mathcal{X}})$ is $(\theta^+ + 1)$ -iterable (in V), since π lifts its models appropriately to models of $\Phi(\mathcal{X})$. And $(k, \bar{\mathcal{X}}, \bar{\zeta}, \bar{P}^*, \bar{E}, \bar{\mathcal{X}}^+)$ is potential for M (with the obvious definitions for $\bar{P}^*, \bar{\mathcal{X}}^+$). Let $\bar{U} = M_\infty^{\bar{\mathcal{X}}^+}$. Then \bar{U} is wellfounded and the phalanx $\mathfrak{P} = \Phi(\bar{\mathcal{X}}, \bar{\iota}, \bar{E})$ is $(\theta^+ + 1)$ -iterable, where

$\bar{\iota} = \text{lgcd}(\bar{P}^* || \text{lh}(\bar{E}))$. For we can find $M' \triangleleft P | \text{lh}(E)$ and an appropriate lifting map $\psi : \bar{U} \rightarrow M'$, essentially as in the proof of 3.6. So by 2.38, $\bar{E} \in \mathbb{E}_+^P$, so $E \in \mathbb{E}_+^P$, as desired. \square

None of the remainder of §3 is needed in §§4,5.

We next proceed toward the proof of 3.9. But first we need to make its statement precise, in terms of the manner in which \mathcal{W} and E are definable over N . This involves coding iteration trees on N , definably over N .

Remark 3.15. Squashing of premice complicates our tree coding. If N is type 3, we will only have direct representations for elements of N^{sq} . But a tree \mathcal{T} on N can have $\nu(F^N) < \text{lh}(E_0^{\mathcal{T}}) < \text{OR}(N)$, so $E = E_0^{\mathcal{T}} \notin N^{\text{sq}}$. One option is to represent such an E with a pair (a, f) such that $i_{F^N}(f)(a) = E$. But instead, we will adjust the rules of normal iteration trees, to rule out such \mathcal{T} .

Definition 3.16. Given an iteration tree \mathcal{T} and $\alpha + 1 < \text{lh}(\mathcal{T})$, α is \mathcal{T} -exceptional iff $M_\alpha^{\mathcal{T}}$ is type 3 and $\nu(M_\alpha^{\mathcal{T}}) < \text{lh}(E_\alpha^{\mathcal{T}}) < \text{OR}(M_\alpha^{\mathcal{T}})$. A tree \mathcal{T}' on a k -sound premouse N is *pre-adjusted k -maximal* iff there is a k -maximal tree \mathcal{T} on N such that \mathcal{T}' has index set

$$\text{lh}(\mathcal{T}) \cup \{(\alpha, 0) \mid \alpha + 1 < \text{lh}(\mathcal{T}) \text{ \& } \alpha \text{ is } \mathcal{T}\text{-exceptional}\},$$

ordered with $\alpha < (\alpha, 0) < \beta$ for $\alpha < \beta$, and if α is \mathcal{T} -exceptional, then $E_\alpha^{\mathcal{T}'} = F(M_\alpha^{\mathcal{T}})$ and $E_{(\alpha, 0)}^{\mathcal{T}'} = E_\alpha^{\mathcal{T}}$, and otherwise $E_\alpha^{\mathcal{T}'} = E_\alpha^{\mathcal{T}}$, and $<_{\mathcal{T}'}$, $D^{\mathcal{T}'}$, $\deg^{\mathcal{T}'}$ are determined as for k -maximal trees.

Given $N, \mathcal{T}, \mathcal{T}'$ as above, $\text{adj}(\mathcal{T})$ denotes the unique tree \mathcal{U} on N with $\mathcal{U} \cong \mathcal{T}'$ and the index set of \mathcal{U} an ordinal. Such trees \mathcal{U} are *adjusted k -maximal*.

Given $\mathcal{W}, P, \gamma, E, \mathcal{W}^+$ as in 2.37, we define the corresponding *adjusted potential k -maximal tree* $\mathcal{U} = \text{adj}(\mathcal{W}^+)$ similarly. In particular, if P is type 3 and $\nu(F^P) < \gamma < \text{OR}(P)$, then for some α , $\text{lh}(\mathcal{U}) = \alpha + 3$, $E_\alpha^{\mathcal{U}} = F^P$ and $E_{\alpha+1}^{\mathcal{U}} = E$. \dashv

Remark 3.17. We make some remarks on the preceding definition; see [13] for more details. We use notation as in 3.16. For \mathcal{T}' -indices $x < y$ with $E_y^{\mathcal{T}'}$ defined, we have $\nu_x^{\mathcal{T}'} \leq \nu_y^{\mathcal{T}'}$, and if $\nu_x^{\mathcal{T}'} = \nu_y^{\mathcal{T}'}$ then $x = \alpha$ and $y = (\alpha, 0)$ for some \mathcal{T} -exceptional α . The tree structure of $\mathcal{T}' \upharpoonright \text{OR}$ is essentially that of \mathcal{T} ; but for example if $\text{pred}^{\mathcal{T}}(\beta + 1) = \alpha$ and α is \mathcal{T} -exceptional and $\nu(M_\alpha^{\mathcal{T}}) \leq \text{cr}(E_\beta^{\mathcal{T}})$ then $\text{pred}^{\mathcal{T}'}(\beta + 1) = (\alpha, 0)$. (Thus, we also might have $\gamma < \text{lh}(\mathcal{T})$ such that $\gamma >_{\mathcal{T}} \beta + 1 >_{\mathcal{T}} \alpha$ but $\gamma \not>_{\mathcal{T}'} \alpha$.) In this case, \mathcal{T} drops in model at $\beta + 1$ (but with $M_{\beta+1}^{*\mathcal{T}} \notin (M_\alpha^{\mathcal{T}})^{\text{sq}}$), and $M_{\beta+1}^{*\mathcal{T}'} = M_{\beta+1}^{*\mathcal{T}}$. So for each $\alpha < \text{lh}(\mathcal{T})$, $M_\alpha^{\mathcal{T}'} = M_\alpha^{\mathcal{T}}$, and if $\alpha <_{\mathcal{T}} \beta$ and $(\alpha, \beta]_{\mathcal{T}}$ does not drop in model then $\alpha <_{\mathcal{T}'} \beta$ and $(\alpha, \beta]_{\mathcal{T}'}$ does not drop in model and $i_{\alpha\beta}^{\mathcal{T}'} = i_{\alpha\beta}^{\mathcal{T}}$. Moreover, for \mathcal{T} -exceptional α , whenever \mathcal{T}' forms an ultrapower of some $P \trianglelefteq M_{(\alpha, 0)}^{\mathcal{T}'}$, we have $P \triangleleft M_{(\alpha, 0)}^{\mathcal{T}'} | \text{OR}(M_\alpha^{\mathcal{T}})$.

The map $\mathcal{T} \mapsto \text{adj}(\mathcal{T})$ is well-defined and 1-1, and, for e.g., $(k, \omega_1, \omega_1 + 1)$ -iterability is equivalent to the corresponding iterability for adjusted k -maximal trees. By coding only adjusted k -maximal trees, we avoid the problem of 3.15.

Given an iteration tree \mathcal{T} on a premouse N , and $\alpha + 1 < \text{lh}(\mathcal{T})$, let $\lambda_\alpha^\mathcal{T} = \nu(E_\alpha^\mathcal{T})$ if $M_\alpha^\mathcal{T}$ is type 3 and $E_\alpha^\mathcal{T} = F(M_\alpha^\mathcal{T})$, and let $\lambda_\alpha^\mathcal{T} = \text{lh}(E_\alpha^\mathcal{T})$ otherwise. So $\nu(E_\alpha^\mathcal{T}) \leq \lambda_\alpha^\mathcal{T}$. If \mathcal{T} is adjusted k -maximal, and $\alpha < \beta < \text{lh}(\mathcal{T})$, then $\lambda_\alpha^\mathcal{T}$ is a cardinal of $M_\beta^\mathcal{T}$, $M_\alpha^\mathcal{T} \restriction \lambda_\alpha^\mathcal{T} = M_\beta^\mathcal{T} \restriction \lambda_\beta^\mathcal{T}$, and if $\beta + 1 < \text{lh}(\mathcal{T})$ then $\lambda_\alpha^\mathcal{T} \leq \nu(E_\beta^\mathcal{T})$.

The precise meaning of the following lemma is described in the definition to follow:

Lemma 3.18 (Coding of finite trees). Let N be a k -sound premouse and \mathcal{T} be a finite, adjusted k -maximal tree on N . In the codes, the models and embeddings of \mathcal{T} are $\mathbf{r}\Delta_{k+1}^N$, and the $\mathbf{r}\Sigma_{\deg \mathcal{T}(n)+1}^N$ satisfaction relation for $M_n^\mathcal{T}$ is $\mathbf{r}\Sigma_{k+1}^N$.⁴⁴

Definition 3.19. Let N, k, \mathcal{T} be as above. We define $\mathbf{r}\Delta_{k+1}^N$ relations

$$\mathbf{M}_n(x), \approx_n(x, y), e_n(x, y), \mathbb{E}_n(x), F_n(x), i_{nm}(x, y)$$

(e.g. $\mathbf{M}_n(x)$ is in two variables n, x) such that properties (1)-(8) below hold. (The intended meaning is as follows: $\mathbf{M}_n \subseteq \mathfrak{C}_0(N)$ is a class of codes for elements of $\mathfrak{C}_0(M_n^\mathcal{T})$, and $\approx_n, e_n, \mathbb{E}_n, F_n$ represent equality, membership, the internal extender sequence and active extender predicates of $\mathfrak{C}_0(M_n^\mathcal{T})$ with respect to the coding, and i_{nm} codes $i_{nm}^\mathcal{T}$.) We will have:

- (1) $\mathbf{M}_n \subseteq \mathfrak{C}_0(N)$, and \approx_n is an equivalence relation on \mathbf{M}_n .
- (2) $e_n \subseteq \mathbf{M}_n \times \mathbf{M}_n$ and $\mathbb{E}_n, F_n \subseteq \mathbf{M}_n$, and e_n, \mathbb{E}_n, F_n each respect \approx_n .
- (3) $M_n^\mathcal{T} = (|M_n^\mathcal{T}|, \in, \mathbb{E}(M_n^\mathcal{T}), F(M_n^\mathcal{T})) \cong (\mathbf{M}_n, e_n, \mathbb{E}_n, F_n) / \approx_n$.
- (4) $i_{nm} \neq \emptyset$ just when $i_{nm}^\mathcal{T}$ is defined, and in this case, $i_{nm} \subseteq \mathbf{M}_n \times \mathbf{M}_m$ and respects (\approx_n, \approx_m) , and $i_{nm}^\mathcal{T} \cong i_{nm} / (\approx_n, \approx_m)$.

We start with $(\mathbf{M}_0, \approx_0, e_0, \mathbb{E}_0, F_0) = (\mathfrak{C}_0(N), =, \in, \mathbb{E}^{\mathfrak{C}_0(N)}, F^{\mathfrak{C}_0(N)})$ and $i_{00} = \text{id} : \mathbf{M}_0 \rightarrow \mathbf{M}_0$. Suppose we have defined \mathbf{M}_n , etc. We make some definitions. Let

$$\varsigma_n : \mathbf{M}_n \rightarrow \mathfrak{C}_0(M_n^\mathcal{T})$$

be the natural surjection, that is, $\varsigma_n(x)$ is the image of $[x]_{\approx_n}$ under Mostowski collapse. If $\varsigma_n(x) = x'$, we say (x, n) is a *code* for (x', n) (or just x is a code for x' if n is understood).

Let $\mathbf{t} = \mathbf{t}^\mathcal{T} = (\prec_\mathcal{T}, \mathscr{D}^\mathcal{T}, \deg^\mathcal{T}, d^\mathcal{T})$ where⁴⁵

$$d^\mathcal{T} = \{(\alpha, \beta) \in \text{lh}(\mathcal{T})^2 \mid \alpha <_\mathcal{T} \beta \text{ and } (\alpha, \beta]_\mathcal{T} \cap D^\mathcal{T} \neq \emptyset\}.$$

⁴⁴The $\mathbf{r}\Delta_{k+1}^N$, $\mathbf{r}\Sigma_{\deg \mathcal{T}(n)+1}^N$ and $\mathbf{r}\Sigma_{k+1}^N$ referenced here are all pure, not generalized. Likewise elsewhere in this paper, unless explicitly stated otherwise.

⁴⁵ $d^\mathcal{T}$ is presently redundant, but included for consistency with later notation for transfinite trees.

Let $\text{lh}(\mathbf{t}) = \text{lh}(\mathcal{T})$. Let $\rho_{k-1}^{-N} = \rho_{k-1}^N$ if $\rho_{k-1}^N < \rho_0^N$, and $\rho_{k-1}^{-N} = \emptyset$ otherwise. We say p is *good for* (\mathcal{T}, m) ⁴⁶ iff $m < \text{lh}(\mathcal{T})$ and

$$p = (p_k^N, u_{k-1}^N, \rho_{k-1}^{-N}, \mathbf{t}, \vec{x})$$

for some $\vec{x} = (x_0, \dots, x_{\text{lh}(\mathcal{T})-2}) \in N^{\text{lh}(\mathcal{T})-1}$ and for each $n < m$, we have (i) $M_n(x_n)$, (ii) if $E_n^T = F(M_n^T)$ then $\varsigma_n(x_n) = \emptyset$, and (iii) if $E_n^T \neq F(M_n^T)$ then $\varsigma_n(x_n) = \text{lh}(E_n^T)$. We say p is *good for* \mathcal{T} iff p is good for $(\mathcal{T}, \text{lh}(\mathcal{T}) - 1)$. Note that “good for (\mathcal{T}, m) ” makes reference to M_n , etc, only for $n < m$. We require p to be good for (\mathcal{T}, m) in order to define M_m , \approx_m , etc. It will become clear that we can find p which is good for \mathcal{T} .

We continue with the properties of the coding, giving upper bounds for its complexity, and the complexity of the satisfaction relation for models of \mathcal{T} . In order to assist in the calculations, we also define functions which yield standard parameters and translation procedures associated to \mathcal{T} , in the codes: **Crit** _{n} is a code for $\text{cr}(E_n^T)$; **Star** _{$n+1$} is a code for $\text{OR}(M_{n+1}^{*T})$; given $m \neq n$, **Shift** translates codes for ordinals within the support of $E_{\min(m,n)}^T$ between M_m and M_n ; **Def** converts a given definition over $\mathfrak{C}_0(M_{n+1}^T)$ for a set $A \subseteq \text{cr}(E_n^T)$ to a definition for A over $\mathfrak{C}_0(M_{n+1}^{*T})$ (both definitions are of the relevant complexity, from parameters); and **MeasDef** converts a given finite set a of E_n^T -generators into a $\Sigma_1^{M_n^{*T}}$ -definition of $(E_n^T)_a$. The latter two functions are just effective implementations of the proofs of [3, 4.5] and the Closeness Lemma [3, 6.1.5], which the reader should probably have in mind:

- (5) The definitions of M_n, \approx_n , etc, are $\text{r}\Delta_{k+1}^N(\{p\})$ in any parameter p good for (\mathcal{T}, n) ; the definitions used only depend on k and the type of N .
- (6) *Satisfaction*: Let **Sat** _{n} (φ, x) iff $n < \text{lh}(\mathcal{T})$, φ is an $\text{r}\Sigma_{\deg^T(n)+1}$ formula, $x = (x_0, \dots, x_{\ell-1})$ for some $\ell < \omega$, $M_n(x_i)$ holds for each $i < \ell$, and

$$\mathfrak{C}_0(M_n^T) \models \varphi(\varsigma_n(x_0), \dots, \varsigma_n(x_{\ell-1})).$$

Then **Sat** is $\text{r}\Sigma_{k+1}^N(\{p\})$ for good p , uniformly.

The set of triples (n, φ, x) such that

$$n < \text{lh}(\mathcal{T}), \mathcal{D}_{\deg}^T \cap (0, n]_{\mathcal{T}} \neq \emptyset \text{ and } \text{Sat}_n(\varphi, x)$$

is $\text{r}\Delta_{k+1}^N(\{p\})$ for good p , uniformly.

- (7) *Parameters*: There are functions **Crit**, **Star**, uniformly $\text{r}\Sigma_{k+1}^N(\{p\})$ for good p , such that:

- (a) $\text{dom}(\text{Crit}) = \text{lh}(\mathcal{T}) - 1$. We have $M_n(\text{Crit}_n)$ and $\varsigma_n(\text{Crit}_n) = \text{cr}(E_n^T)$.

⁴⁶We have encoded in good p more information than strictly necessary, so as to skip some calculations.

(b) $\text{dom}(\text{Star}) = \text{lh}(\mathcal{T}) \setminus \{0\}$. If $n+1 \notin \mathcal{D}^\mathcal{T}$ then $\text{Star}_{n+1} = \emptyset$; otherwise

$$\mathbf{M}_{\text{pred}^\mathcal{T}(n+1)}(\text{Star}_{n+1}) \text{ holds and } \varsigma_{\text{pred}^\mathcal{T}(n+1)}(\text{Star}_{n+1}) = \text{OR}(M_{n+1}^{\ast\mathcal{T}}).$$

(8) *Translations*: There are functions **Shift**, **Def**, **MeasDef**, uniformly $\text{r}\Sigma_{k+1}^N(\{p\})$ for good p , such that:

(a) *Shifting generators*: $\text{dom}(\text{Shift})$ is the set of tuples (m, n, x) such that

(i) $m, n < \text{lh}(\mathcal{T})$ and $m \neq n$ and $\mathbf{M}_m(x)$, and

(ii) $\varsigma_m(x) \in \lambda_{\min(m,n)}^\mathcal{T}$.

If $\text{Shift}_{mn}(x) = y$ then $\mathbf{M}_n(y)$ and $\varsigma_n(y) = \varsigma_m(x)$.

(b) *Definitions of sets*; cf. [3, 4.5]: We have $\text{dom}(\text{Def}) =$

$$\{(n+1, x, \varphi) \mid n+1 < \text{lh}(\mathcal{T}) \ \& \ \mathbf{M}_{n+1}(x) \ \& \ \varphi \text{ is } \text{r}\Sigma_{\text{deg}^\mathcal{T}(n+1)+1}^\mathcal{T}\}.$$

Let $m = \text{pred}^\mathcal{T}(n+1)$. Then $\text{Def}_{n+1}(x, \varphi) = (x', \varphi')$ where:

(i) $\mathbf{M}_m(x')$ with $\varsigma_m(x') \in \mathfrak{C}_0(M_{n+1}^{\ast\mathcal{T}})$,

(ii) φ' is an $\text{r}\Sigma_{\text{deg}^\mathcal{T}(n+1)+1}^\mathcal{T}$ formula, and

(iii) for all $\alpha < \text{cr}(E_n^\mathcal{T})$, we have

$$\mathfrak{C}_0(M_{n+1}^\mathcal{T}) \models \varphi(\varsigma_{n+1}(x), \alpha) \iff \mathfrak{C}_0(M_{n+1}^{\ast\mathcal{T}}) \models \varphi'(\varsigma_m(x'), \alpha).$$

(c) *Definitions of measures*; cf. [3, 6.1.5]: We have $\text{dom}(\text{MeasDef}) =$

$$\{(n, a) \mid n+1 < \text{lh}(\mathcal{T}) \ \& \ \mathbf{M}_n(a) \ \& \ \varsigma_n(a) \in [\lambda_n^\mathcal{T}]^{<\omega}\},$$

and $\text{MeasDef}_n(a) = (x, \varphi)$ is such that, letting $k = \text{pred}^\mathcal{T}(n+1)$ and $C = \mathfrak{C}_0(M_{n+1}^{\ast\mathcal{T}})$, we have:

(i) $\mathbf{M}_k(x)$ and $\varsigma_k(x) \in C$ and φ is $\text{r}\Sigma_1$, and

(ii) $(E_n^\mathcal{T})_{\varsigma_n(a)} = \{A \in C \mid C \models \varphi(A, \varsigma_k(x))\}.$

This completes the list of properties. Now suppose p is good for (\mathcal{T}, n) , and we have defined the following things:

- $(\mathbf{M}_j, \approx_j, e_j, \mathbb{E}_j, F_j)$, i_{jk} and Sat_j for $j, k \leq n$,
- Crit_j , MeasDef_j , Star_{j+1} , Def_{j+1} for $j < n$, and
- Shift_{jk} for $j, k \leq n$ with $j \neq k$.

Suppose $n+1 < \text{lh}(\mathcal{T})$, and p is actually good for $(\mathcal{T}, n+1)$. We want to define \mathbf{M}_{n+1} etc. We have $\mathbf{M}_n(x_n)$ and x_n determines $E_n^\mathcal{T}$. Let Crit_n be the natural code in \mathbf{M}_n , determined by x_n , for $\text{cr}(E_n^\mathcal{T})$. Let $k = \text{pred}^\mathcal{T}(n+1)$; note that p determines k and determines whether $k \in \mathcal{D}^\mathcal{T}$. If $k = n$ let $c = \text{Crit}_n$; otherwise let

$$c = \text{Shift}_{nk}(\text{Crit}_n).$$

So $\mathbf{M}_k(c)$ and $\varsigma_k(c) = \text{cr}(E_n^\mathcal{T})$ in general. If $k \in \mathcal{D}^\mathcal{T}$, let Star_{n+1} be the natural code in \mathbf{M}_k for $\text{OR}(M_{n+1}^{\ast\mathcal{T}})$, determined by c and x_k . Using these codes, take $\mathbf{M}_{n+1}(x)$ to be the natural formula asserting “ $x = ((q, t), b)$ where:

- $M_k(q)$ and $\varsigma_k(q) \in \mathfrak{C}_0(M_{n+1}^{*\mathcal{T}})$,
- t is a generalized $\deg^{\mathcal{T}}(n+1)$ term,
- $M_n(b)$ and $\varsigma_n(b) \in [\lambda_n^{\mathcal{T}}]^{<\omega}$.

Here we use Sat_k to assert “ $\varsigma_k(q) \in \mathfrak{C}_0(M_{n+1}^{*\mathcal{T}})$ ”, etc. Now to define \approx_{n+1} , Sat_{n+1} , etc, we use Los’ Theorem. For this we need to translate statements about measure one sets of $E_n^{\mathcal{T}}$ to statements over $M_k^{\mathcal{T}}$. For this, the main arguments follow the proofs of [3, 4.5] and [3, 6.1.5], observing that everything is sufficiently effective, and in particular, that we get $\text{r}\Sigma_{k+1}^N(\{p\})$ functions. We will use degree $k+1$ min-terms to uniformize $\text{r}\Sigma_{k+1}^N(\{p\})$ relations and obtain our functions. We will just sketch the definition of MeasDef_n , assuming $k < n$; the case $k = n$ is much easier. Let $\kappa = \text{cr}(E_n^{\mathcal{T}})$, so $\kappa < \nu(E_k^{\mathcal{T}})$.

Assume $E_n^{\mathcal{T}} \neq F(M_n^{\mathcal{T}})$, so $\lambda_n^{\mathcal{T}} = \text{lh}(E_n^{\mathcal{T}})$. Given $a \in M_n$ with $\varsigma_n(a) \in [\lambda_n^{\mathcal{T}}]^{<\omega}$, we can identify via $\text{r}\Delta_1$ -satisfaction for $M_n^{\mathcal{T}}$ (and our good parameter p), those codes $c \in M_n$ such that $\gamma = \varsigma_n(c)$ is the position of $(E_n^{\mathcal{T}})_{\varsigma_n(a)}$ in the canonical wellorder of $M_n^{\mathcal{T}}$ (so $\gamma \in (\kappa^{++})^{M_n^{\mathcal{T}}}$).

Using Shift , such codes c are translated to M_{k+1} -codes c' for γ , since if $k+1 < n$ then $(\kappa^{++})^{M_n^{\mathcal{T}}} \leq \lambda_{k+1}^{\mathcal{T}}$. Write $c' = ((q, t), b)$. Let $\mu = \text{cr}(E_k^{\mathcal{T}})$ and

$$\ell = \text{pred}^{\mathcal{T}}(k+1).$$

Now $\gamma < (\kappa^{++})^{M_n^{\mathcal{T}}} \leq (\kappa^{++})^{M_{k+1}^{\mathcal{T}}}$, so there is $f : [\mu]^{|\varsigma_k(b)|} \rightarrow \mu$ such that $f \in \mathfrak{C}_0(M_{k+1}^{*\mathcal{T}})$ and

$$i_{k+1}^{*\mathcal{T}}(f)(\varsigma_k(b)) = \gamma = t^{M_{k+1}^{\mathcal{T}}}(i_{k+1}^{*\mathcal{T}}(\varsigma_\ell(q)), \varsigma_k(b)).$$

We can identify the M_ℓ -codes d for such functions f sufficiently effectively; the main point here is that we can refer to Sat_{k+1} (or just \approx_{k+1}) to check equality (we also have the M_k -code cr_k for μ , so can check that $f : [\mu]^{|\varsigma_k(b)|} \rightarrow \mu$). Such M_ℓ -codes d can then be converted to M_k -codes d' using Shift .

The set of tuples (a, b, d') as above is $\text{r}\Sigma_{k+1}^N(\{p\})$, so there is an $\text{r}\Sigma_{k+1}^N(\{p\})$ function selecting some such b, d' as a function of a . Let $b^* = \varsigma_k(b)$ and $f = \varsigma_k(d')$. Note that

$$(E_n^{\mathcal{T}})_{\varsigma_n(a)} \text{ is } \text{r}\Sigma_1^{M_k^{\mathcal{T}}|\text{lh}(E_k^{\mathcal{T}})}(\{b^*, f\}),$$

and uniformly so. Now $M_k^{\mathcal{T}}|\text{lh}(E_k^{\mathcal{T}}) \trianglelefteq M_{n+1}^{*\mathcal{T}}$ and p incorporates the code x_k for $\text{lh}(E_k^{\mathcal{T}})$, so this almost suffices. However, it might be that $M_{n+1}^{*\mathcal{T}}$ is active type 3 and $M_k^{\mathcal{T}}|\text{lh}(E_k^{\mathcal{T}}) \triangleleft M_{n+1}^{*\mathcal{T}}$ but $M_k^{\mathcal{T}}|\text{lh}(E_k^{\mathcal{T}}) \not\trianglelefteq \mathfrak{C}_0(M_{n+1}^{*\mathcal{T}})$, in which case the relevant parameters are not available directly to $\mathfrak{C}_0(M_{n+1}^{*\mathcal{T}})$. But because \mathcal{T} is adjusted k -maximal, note that $M_{n+1}^{*\mathcal{T}} \triangleleft M_k^{\mathcal{T}}$, and then it is easy to convert our (parameters for) definitions of measures over $M_k^{\mathcal{T}}|\text{lh}(E_k^{\mathcal{T}})$ into (parameters for) definitions for those measures over $\mathfrak{C}_0(M_{n+1}^{*\mathcal{T}})$, using the method above. This completes the definition of $\text{MeasDef}_n(a)$ assuming $E_n^{\mathcal{T}} \neq F(M_n^{\mathcal{T}})$.

If instead $E_n^{\mathcal{T}} = F(M_n^{\mathcal{T}})$, use the proof of [3, 6.1.5], effectivized through the function Def , combined with the preceding argument. We leave the remaining details to the reader.

Using standard calculations with Los Theorem, it is now straightforward to define \approx_{n+1} , etc, and \mathbf{Sat}_{n+1} , in terms of \mathbf{Sat}_k and $\mathbf{MeasDef}_n$, and see the relevant properties. The definition of \mathbf{Sat}_{n+1} also easily yields \mathbf{Def}_{n+1} . For $\mathbf{Shift}_{n,n+1}$, we convert an \mathbf{M}_n -code c for an ordinal $< \lambda_n^{\mathcal{T}}$ to the \mathbf{M}_{n+1} -code $((\emptyset, t_{\text{id}}), c)$, where \emptyset is the natural \mathbf{M}_k -code for the empty set and t_{id} the term for the identity map. For $\mathbf{Shift}_{n+1,n}$, an \mathbf{M}_{n+1} -code $((q, t), b)$ for an ordinal $\alpha < \lambda_n^{\mathcal{T}}$ can be converted to some \mathbf{M}_n -code for α , using min-term uniformization to make the selection. \dashv

To summarize, we have:

Lemma 3.20. Let N, k, \mathcal{T} be as above. Then there is p which is good for \mathcal{T} . For any p which is good for \mathcal{T} , the coded satisfaction relation \mathbf{Sat} , defined relative to p , is $\mathbf{r}\Sigma_{k+1}(\{p\})$.

We next extend the coding to infinite trees \mathcal{T} . The plan is as follows. A code for an object $z \in \mathfrak{C}_0(M_\alpha^{\mathcal{T}})$ will be of the form (α, x) , and x will essentially consist of some finite support for z , in terms of functions and generators coming from finitely many earlier models $M_\beta^{\mathcal{T}}$. We demand that the full tree structure \mathbf{t} of \mathcal{T} , and a function \mathbf{lh} specifying the sequence of extenders used in the tree are given in advance, with both suitably definable over N from some parameter. The specification of the extenders must naturally be given through the coding, analogous to the finite tree case; that is, $(\alpha, \mathbf{lh}(\alpha))$ will be a code for $\mathbf{lh}(E_\alpha^{\mathcal{T}})$ (or for \emptyset if $E_\alpha^{\mathcal{T}} = F(M_\alpha^{\mathcal{T}})$). (So \mathbf{lh} is the analog of the parameter \vec{x} used in the finite case.) We now proceed to the details.

Definition 3.21. Let N be a k -sound premouse. Let $1 \leq \lambda \leq \rho_0^N$ and

$$\mathbf{t} = (<_{\mathbf{t}}, D^{\mathbf{t}}, \deg^{\mathbf{t}}, d^{\mathbf{t}})$$

where $<_{\mathbf{t}}$ is an iteration tree order, $D^{\mathbf{t}}$ a drop structure and $\deg^{\mathbf{t}}$ a degree structure, each on λ , and $d^{\mathbf{t}}$ as before.

Suppose $\lambda < \omega$. For $\vec{x} \in N^{\lambda-1}$ let

$$p = p(\mathbf{t}, \vec{x}) = (p_k^N, u_{k-1}^N, \rho_{k-1}^{-N}, \mathbf{t}, \vec{x});$$

We say p is *coherent* (for N) iff there is an adjusted k -maximal tree \mathcal{T} on N such that p is good for \mathcal{T} .

Now remove the restriction that $\lambda < \omega$, but suppose that \mathbf{t} is $\mathbf{r}\Delta_{k+1}^N$. Fix an $\mathbf{r}\Sigma_{k+1}^N$ function \mathbf{lh} with domain λ . The intention is that \mathbf{lh} is a sequence of codes for extender lengths, for a tree \mathcal{T} with structure \mathbf{t} . The elements $z \in \mathfrak{C}_0(M_\alpha^{\mathcal{T}})$ are coded with pairs (α, x) , where x describes a finite support for z in terms of codes for earlier models in \mathcal{T} . We define $\mathcal{T} \upharpoonright (\alpha + 1)$, and the collection C_α of all x such that (α, x) is a code, and the interpretation $\varsigma(\alpha, x) \in \mathfrak{C}_0(M_\alpha^{\mathcal{T}})$, by recursion on α .

First, $C_0 = \mathfrak{C}_0(N)$, and $\varsigma(0, x) = x$.

Now let $\eta \geq 1$, and suppose:

- we have defined the adjusted k -maximal tree $\mathcal{T} \upharpoonright \eta$ on N , with structure $\mathfrak{t} \upharpoonright \eta$,
- for all $\beta < \eta$, we have defined C_β and $\varsigma(\beta, x)$ for all $x \in C_\beta$, and

$$\mathfrak{C}_0(M_\beta^T) = \{\varsigma(\beta, x) \mid x \in C_\beta\}.$$

Then we say that $(\mathfrak{t}, \mathfrak{h})$ is η -coherent. (Note that 1-coherence is trivial.)

Suppose η is a limit and $\eta < \lambda$. We set $[0, \eta]_{\mathcal{T}} = [0, \eta]_{\mathfrak{t}}$. We set C_η to be the set of all pairs $(\beta + 1, x)$ such that $\beta + 1 <_{\mathfrak{t}} \eta$ and $(\beta + 1, \eta)_{\mathfrak{t}}$ does not drop in model and $x \in C_{\beta+1}$. We define

$$\varsigma(\eta, (\beta + 1, x)) = i_{\beta+1, \eta}^{\mathcal{T}}(\varsigma(\beta + 1, x)).$$

Assuming that $M_\eta^{\mathcal{T}}$ is wellfounded, note that $(\eta + 1)$ -coherence follows.

Suppose $\eta = \alpha + 1 < \lambda$ and that $\mathfrak{h}(\alpha) \in C_\alpha$. Let $\gamma = \varsigma(\alpha, \mathfrak{h}(\alpha))$. Suppose either:

- $\gamma = 0$ and $E = F(M_\alpha^T) \neq \emptyset$, or
- $\gamma < \rho_0(M_\alpha^T)$ and $E = F(M_\alpha^T | \gamma) \neq \emptyset$.

Then we set $E_\alpha^{\mathcal{T}} = E$; suppose this determines an adjusted k -maximal tree $\mathcal{T} \upharpoonright (\alpha + 2)$ (including wellfoundedness). Let $\beta = \text{pred}^{\mathfrak{t}}(\alpha + 1)$ and $M^* = M_{\alpha+1}^{*\mathcal{T}}$. Then $C_{\alpha+1}$ is the set of pairs of the form

$$x = ((\beta, q, u), (\alpha, a))$$

such that $q \in C_\beta$ and

$$q' = \varsigma(\beta, q) \in \mathfrak{C}_0(M^*),$$

u is an $\text{r}\Sigma_n$ -Skolem term, where $n = \deg^{\mathcal{T}}(\alpha + 1)$, $a \in C_\alpha$ and

$$a' = \varsigma(\alpha, a) \in [\lambda_\alpha^{\mathcal{T}}]^{<\omega}.$$

For these objects, we define

$$\varsigma(\alpha + 1, x) = [f_{q', u}, a']_{E_\alpha^{\mathcal{T}}}^{M^*, n}.$$

Note then that $(\eta + 1)$ -coherence follows.

A *code* (relative to $(\mathfrak{t}, \mathfrak{h})$) is a pair (α, x) such that $(\mathfrak{t}, \mathfrak{h})$ is $(\alpha + 1)$ -coherent and $x \in C_\alpha$. (So $\varsigma(\alpha, x)$ is defined for codes (α, x) .) We say that $(\mathfrak{t}, \mathfrak{h})$ is *coherent* iff it is λ -coherent. \dashv

Remark 3.22. For the definability of the coding, and the associated satisfaction relation, etc, we will use the fact that adjusted k -maximal trees \mathcal{T} are natural direct limits of finite such trees $\bar{\mathcal{T}}$. In [15, §2.3], finite supports for k -maximal iteration trees were discussed. We assume the reader is familiar with such methods. Adjusted trees were not considered there, but it is straightforward to adapt those methods to adjusted trees. In particular, we have that for each

adjusted k -maximal tree \mathcal{T} of length λ and for each finite set $J \subseteq \lambda \times V$ with $J_\alpha \subseteq \mathfrak{C}_0(M_\alpha^\mathcal{T})$, there is a *finite support* $S \subseteq \lambda \times V$ (defined much as in [15]) with $J \subseteq S$. Moreover, for any such S , letting $\tilde{\mathcal{T}} = \tilde{\mathcal{T}}_S$ be the corresponding finite hull of \mathcal{T} (finite meaning that $\text{lh}(\tilde{\mathcal{T}})$ is finite), the natural “copy map”

$$\pi_{\bar{\alpha}} : M_{\bar{\alpha}}^{\tilde{\mathcal{T}}} \rightarrow M_{\alpha}^{\mathcal{T}}$$

is a near $\deg^{\mathcal{T}}(\alpha)$ -embedding (here $\alpha \in I_S$ where I_S is the projection of S on the left coordinate, and $\deg^{\tilde{\mathcal{T}}}(\bar{\alpha}) = \deg^{\mathcal{T}}(\alpha)$).⁴⁷ The iteration maps commute in the obvious manner with the maps $\pi_{\bar{\alpha}}$.

If \mathcal{T} is our coded tree, then given $\tilde{\mathcal{T}}, \bar{\alpha}, \alpha$ as above and $\bar{z} \in \mathfrak{C}_0(M_{\bar{\alpha}}^{\tilde{\mathcal{T}}})$ and $z = \pi_{\bar{\alpha}}(\bar{z})$, we will convert the code (α, x) for z into a code c for \bar{z} ; here c is a code in the sense of our earlier coding for finite trees (3.19). We will write $c = \text{tc}(\alpha, x)$ (tc for *transitive collapse*); we remark that c depends on the choice of $\tilde{\mathcal{T}}$ though (which below is determined through a *support* S). This code conversion will be appropriately effective, and since $\pi_{\bar{\alpha}}$ is a near $\deg^{\mathcal{T}}(\alpha)$ -embedding, we can define the $\text{r}\Sigma_{\deg^{\mathcal{T}}(\alpha)+1}$ satisfaction relation for $M_{\alpha}^{\mathcal{T}}$ in terms of that for such models $M_{\bar{\alpha}}^{\tilde{\mathcal{T}}}$. The details, to follow, are basically an effective version of material from [15, §2.3].

Definition 3.23. Let $N, k, \mathfrak{t}, \mathfrak{lh}$ be as in 3.21 (we don’t assume coherence), with N iterable for finite k -maximal trees. Let $S \subseteq \lambda \times \mathfrak{C}_0(N)$ and let I_S be the projection of S on its left coordinate. We say S is a *support* with respect to $(\mathfrak{t}, \mathfrak{lh})$ iff:

1. S is finite and $0 \in I_S$.
2. If $(\alpha + 1, x) \in S$ then $(\alpha, \mathfrak{lh}(\alpha)) \in S$ and $x = ((\beta, q, u), (\alpha, a))$ where:
 - $\beta = \text{pred}^{\mathfrak{t}}(\alpha + 1)$,
 - u is an $\text{r}\Sigma_{\deg^{\mathfrak{t}}(\alpha+1)}$ Skolem term, and
 - $(\alpha, a), (\beta, q) \in S$ and $\beta + 1 \in I_S$.
3. If $(\alpha, x) \in S$ and α is a limit then

$$\max(\alpha \cap I_S) = \gamma + 1 <_{\mathfrak{t}} \alpha$$

and $x = (\beta + 1, y) \in S$ where $\beta + 1 <_{\mathfrak{t}} \alpha$ and $(\beta + 1, \alpha)_{\mathfrak{t}}$ does not drop in model or degree. Moreover, if $\alpha + 1 \in I_S$ and $x = \mathfrak{lh}(\alpha)$ then $\beta + 1 < \gamma + 1$.⁴⁸

If $(\alpha, x) \in S$, say that (α, x) is a *potential code*. Now we want to translate from the codes for infinite trees discussed in 3.21 and our earlier coding for finite

⁴⁷However, as in [15], those $\alpha \in I_S$ such that $\alpha < \max(I_S)$ but $\alpha + 1 \notin I_S$, do not correspond to any $\bar{\alpha}$; this is also discussed further below.

⁴⁸The last requirement helps ensure that when S is a support, then \mathfrak{t}_S is an adjusted k -maximal tree; see [15] for further explanation.

trees in 3.19. Toward this, we compute the *transitive collapse* $\text{tc}(\alpha, x)$ of (α, x) , with respect to a given support S with $(\alpha, x) \in S$ (though (α, x) might not actually be a code in the sense of 3.21). Let

$$E_S = \{\alpha \in I_S \mid \alpha + 1 \in I_S\},$$

$$P_S = I_S \setminus (E_S \cup \{\max(I_S)\}).$$

Let $\mathbf{t}_S = \mathbf{t} \upharpoonright I_S$. We consider \mathbf{t}_S as a padded tree structure, indexed with ordinals in I_S , with padding occurring at precisely the ordinals in P_S , and extenders used at the ordinals in E_S .⁴⁹ Let $\text{tc}(\mathbf{t}_S)$ denote the non-padded tree structure which is otherwise isomorphic to \mathbf{t}_S . We enlarge S to a support which includes canonical potential codes for the empty set with respect to each $\alpha \in I_S$, and which is closed under (coded) images under iteration maps. So, recursively on $\alpha \in I_S$, we define e_α , such that (α, e_α) codes the empty set: Set $e_0 = \emptyset$. Given $\alpha + 1 \in I_S$ and $\beta = \text{pred}^t(\alpha + 1)$ (so $\beta \in I_S$ also), let

$$e_{\alpha+1} = ((\beta, e_\beta, c), (\alpha, e_\alpha)),$$

where $c = c(v)$ is the Skolem term for the constant function with value v . Given a limit $\alpha \in I_S$ and $\gamma + 1 = \max(I_S \cap \alpha)$, let

$$e_\alpha = (\gamma + 1, e_{\gamma+1}).$$

Now for $\gamma, \delta \in I_S$ such that $\gamma \leq_t \delta$ and $(\gamma, \delta]_t \cap D^t = \emptyset$, and given a pair of the form $d = (\gamma, x)$, we define a pair $d^{\gamma\delta}$ of the form (δ, y) . Set $d^{\gamma\gamma} = d$. Suppose $\gamma < \delta$. Suppose first that $I_S \cap (\gamma, \delta)_t = \emptyset$. If $\delta = \alpha + 1$ then note that $\gamma = \text{pred}^t(\alpha + 1)$, and we set

$$d^{\gamma, \alpha+1} = (\alpha + 1, ((\gamma, x, c), (\alpha, e_\alpha)))$$

with c as before. If δ is a limit (so $\gamma = \max(I_S \cap \delta)$ and γ is a successor) then we set

$$d^{\gamma\delta} = (\delta, (\gamma, d)).$$

Now if instead $I_S \cap (\gamma, \delta)_t \neq \emptyset$ then letting $\beta = \max(I_S \cap (\gamma, \delta)_t)$, we set

$$d^{\gamma\delta} = (d^{\gamma\beta})^{\beta\delta}.$$

Now let $S' = S \cup \{(\alpha, e_\alpha) \mid \alpha \in I_S\}$ and S'' the closure of S' under $d \mapsto d^{\gamma\delta}$ (for $\gamma, \delta \in I_S$). Note S', S'' are (finite) supports with $I_{S''} = I_{S'} = I_S$.

For $(\alpha, x) \in S''$ we will define $\text{tc}(\alpha, x)$, recursively on α . Here if (\mathbf{t}, \mathbf{h}) is coherent, determining tree \mathcal{T} , and (α, x) is a code, then \mathbf{t}_S is a finite length adjusted k -maximal tree $\tilde{\mathcal{T}}$, a sub-tree of \mathcal{T} , and with \mathbf{M}_n the coding of $M_n^{\tilde{\mathcal{T}}}$ defined as in 3.19, then we will have $\text{tc}(\alpha, x) \in \mathbf{M}_n$. The definition is as follows:

1. $\text{tc}(0, x) = x$,

⁴⁹ P for *Padding* and E for *Extenders*.

2. $\text{tc}(\alpha + 1, ((\beta, q, u), (\alpha, a))) = ((\text{tc}(\beta, q), u), \text{tc}(\alpha, a)),$
3. for limit α , $\text{tc}(\alpha, d) = \text{tc}(\gamma + 1, d^{\beta+1, \gamma+1})$ where $d = (\beta + 1, y)$ and $\gamma + 1 = \max(I_S \cap \alpha)$.

Let $p(\mathfrak{t}, S) = p(\text{tc}(\mathfrak{t}_S), \vec{x})$ where

$$\vec{x} = \langle x_i \rangle_{i < \text{card}(E_S)} = \langle \text{tc}(\alpha, \mathfrak{lh}(\alpha)) \rangle_{\alpha \in E_S}.$$

We say $(\mathfrak{t}, \mathfrak{lh})$ is *pre-coherent* (for N) iff:

- for every finite $J \subseteq \lambda$ there is a support S such that $J \subseteq I_S$, and
- $p(\mathfrak{t}, S)$ is coherent for every support S .

By essentially the arguments in [15], if $(\mathfrak{t}, \mathfrak{lh})$ is coherent, then it is pre-coherent. Note that if S_1, S_2 are supports with $I_{S_1} \subseteq I_{S_2}$ then $S_1 \cup S_2$ is a support. Therefore if $(\mathfrak{t}, \mathfrak{lh})$ is pre-coherent then the collection of all supports is directed under \subseteq .

Suppose $(\mathfrak{t}, \mathfrak{lh})$ is pre-coherent. Then we can make sense of “ $M_\alpha^\mathcal{T}$ ” for each $\alpha < \lambda$, even if this structure is illfounded, as the direct limit of models $M_{\bar{\alpha}}^\mathcal{T}$, where \mathcal{T} is a finite tree determined by a support S and $\bar{\alpha}$ the corresponding collapse of α . We write \mathbf{M}_α^* , etc, for the codeset for elements of $\mathfrak{C}_0(M_\alpha^\mathcal{T})$, etc. So, we define $\text{r}\Delta_{k+1}^N(N)$ relations $\mathbf{M}_\alpha^*(x)$, etc, and $\text{r}\Sigma_{k+1}^N(N)$ relations $\mathbf{Sat}_\alpha^*(\varphi, x)$, etc (we define analogues of all relations defined in 3.19) as follows. For coherent p , let \mathbf{M}^p , etc, denote the relations \mathbf{M} , etc, defined as in 3.19 from p , and let \mathcal{T}^p be the corresponding tree on N . For $x \in N$, set $x \in \mathbf{M}_\alpha^* \iff$

$$\exists S \left[S \text{ is a support for } (\mathfrak{t}, \mathfrak{lh}) \ \& \ (\alpha, x) \in S \ \& \ \mathbf{M}_{\text{ot}(\alpha \cap E_S)}^{p(\mathfrak{t}, S)}(\text{tc}(\alpha, x)) \right].$$

The remaining relations (\mathbf{Sat}_α^* , etc) are defined similarly. \dashv

Remark 3.24. Let $N, k, \mathfrak{t}, \mathfrak{lh}$ be as in 3.21 with $(\mathfrak{t}, \mathfrak{lh})$ pre-coherent. Then there is a unique adjusted putative k -maximal tree \mathcal{T} on N , such that:

- $\text{lh}(\mathcal{T}) \leq \lambda$ and either $\text{lh}(\mathcal{T}) = \lambda$ or \mathcal{T} has a last, illfounded model,
- $\mathfrak{t} \restriction \text{lh}(\mathcal{T}) = \mathfrak{t}^\mathcal{T}$,
- for each $\alpha < \text{lh}(\mathcal{T})$, we have

$$M_\alpha^\mathcal{T} \cong (\mathbf{M}_\alpha^*, e_\alpha^*, \mathbb{E}_\alpha^*, F_\alpha^*) / \approx_\alpha^*,$$

and letting $\varsigma_\alpha^* : \mathbf{M}_\alpha^* \rightarrow M_\alpha^\mathcal{T}$ be the natural surjection, if $E_\alpha^\mathcal{T} = F(M_\alpha^\mathcal{T})$ then $\varsigma_\alpha^*(\mathfrak{lh}(\alpha)) = \emptyset$, and otherwise $\varsigma_\alpha^*(\mathfrak{lh}(\alpha)) = \text{lh}(E_\alpha^\mathcal{T})$.

Moreover, \mathbf{Sat}_α^* defines $\text{r}\Sigma_{\text{deg}\mathcal{T}(\alpha)+1}$ satisfaction for $M_\alpha^\mathcal{T}$ (in the codes), etc, as in 3.19. Here the natural surjection ς_α^* results from the considerations in 3.22, and the fact that \mathbf{Sat}_α^* is correct also follows from 3.22.

We continue to assume that $(\mathfrak{t}, \mathfrak{lh})$ is pre-coherent. Note then that $(\mathfrak{t}, \mathfrak{lh})$ is coherent iff the tree \mathcal{T} above has wellfounded models (hence length λ). And if $(\mathfrak{t}, \mathfrak{lh})$ is coherent then $C_\alpha = \{x \mid M_\alpha^*(x)\}$ and $\varsigma(\alpha, x) = \varsigma_\alpha^*(x)$ for all $x \in C_\alpha$ (where C_α and ς are as in 3.21).

The set of all coherent p (in the sense for codes for finite trees) is a Boolean combination of $\mathbf{r}\Sigma_{k+1}^N(\{q\})$ sets, where $q = (p_k^N, u_{k-1}^N, \rho_{k-1}^{-N})$, uniformly in N (k -sound, of the same type). For example, with such complexity, one can assert that for each n , $\mathbf{M}_n(x_n)$ and x_n codes either \emptyset or the index of an extender on $\mathbb{E}(\mathfrak{C}_0(M_n^T))$, and that drops occur exactly where \mathfrak{t} specifies. (Note that asserting that an extender does not cause a drop in model can require an $\mathbf{r}\Pi_{k+1}^N(\{q\})$ assertion.)

Let X be the set of tuples $(x, \varphi_0, \varphi_1, \varphi_2) \in \mathfrak{C}_0(N)$ such that each φ_i is an $\mathbf{r}\Sigma_{k+1}^N$ formula and for some $\lambda \leq \rho_0^N$:

- $\varphi_0(x, \cdot)$ defines over $\mathfrak{C}_0(N)$ an iteration tree structure \mathfrak{t} on λ , and $\varphi_1(x, \cdot)$ defines its complement,
- $\varphi_2(x, \cdot, \cdot)$ defines over $\mathfrak{C}_0(N)$ a function \mathfrak{lh} with domain λ , and
- $(\mathfrak{t}, \mathfrak{lh})$ is pre-coherent.

Note that X is Π_1 -over- $\mathbf{r}\Sigma_{k+1}^N(\{q\})$, for

$$q = (\varphi_0, \dots, \varphi_3, x, p_k^N, u_{k-1}^N, \rho_{k-1}^{-N}),$$

uniformly in N . Therefore this property (when true) is passed downward under near k -embeddings.

The following definition completes the description of our coding.

Definition 3.25. Let \mathcal{T} be a k -maximal tree on k -sound N . Then

$$(x, \varphi_{\mathfrak{lh}}, \varphi_{\mathfrak{t}}, \varphi_{\neg\mathfrak{t}}) \in \mathfrak{C}_0(N)$$

is *transfinite-good* for \mathcal{T} iff there is $(\mathfrak{t}, \mathfrak{lh})$ which is coherent and yields \mathcal{T} as in 3.24 and $\mathfrak{lh}, \mathfrak{t}, \neg\mathfrak{t}$ are $\mathbf{r}\Sigma_{k+1}^N(\{x\})$, via the formulas $\varphi_{\mathfrak{lh}}, \varphi_{\mathfrak{t}}, \varphi_{\neg\mathfrak{t}}$. \dashv

Remark 3.26. We now clarify hypothesis 3.9(d) and $\mathcal{X}^+ = \text{adj}(\mathcal{W}^+)$. Let $\zeta' + 2 = \text{lh}(\mathcal{X}^+)$. So $E_{\zeta'}^{\mathcal{X}^+} = E$ and either $[\text{lh}(E) = \text{OR}^P$ and $P = M_{\zeta'}^{\mathcal{X}^+}]$ or

$$\text{lh}(E) < \text{OR}^P \text{ and } P|\text{lh}(E) = \mathfrak{C}_0(M_{\zeta'}^{\mathcal{X}^+})|\text{lh}(E).$$

Let $\mathcal{X} = \mathcal{X}^+ \upharpoonright (\zeta' + 1)$. Then 3.9(d) first asserts that if $\text{lh}(\mathcal{X}) \geq \omega$ then there is p which is transfinite-good for \mathcal{X} . If $\text{lh}(\mathcal{X}) \geq \omega$, let $M_{\zeta'}^*$ be defined from p via 3.25 and 3.21. If $\text{lh}(\mathcal{W}) < \omega$, let p be good for \mathcal{X} and let $M_{\zeta'}$ be defined from p as in 3.19. Let $E^* \subseteq M_{\zeta'}^*$ (or $E^* \subseteq M_{\zeta'}$) be the set of codes for elements of \bar{E} (the $P|\text{lh}(E)$ -amenable predicate for E). Then 3.9(d) secondly asserts that E^* is $\mathbf{r}\Sigma_{k+1}^N$.

Proof of 3.9. We follow the notation used in the preceding discussion of coding. Let $\mathcal{X}^+, \mathcal{X}, \zeta', p$ be as in 3.26. Let $e \in N$ be such that (\dagger) E^* is $\Sigma_{k+1}^N(\{e\})$. Using k -reasonableness let $q_0 \in \text{OR}(N)^{<\omega}$ witness 3.1 with respect to (N, k, x, ω) where

$$x = (\mathfrak{c}, p, \zeta', e, p_{k+1}^N).$$

Let $M = \text{cHull}_{k+1}^N(\{q_0\})$, let $\pi : M \rightarrow N$ be the uncollapse, let $\pi(\bar{p}) = p$ and let $\bar{\mathcal{X}}$ be defined over M from \bar{p} as \mathcal{X} is defined over N from p ; “bars” denote preimages under π in general.

Claim 1. *We have:*

- (a) *The hypotheses of 3.9 and properties mentioned in 3.26 and (\dagger) regarding $N, \mathcal{W}, \mathcal{X}, p$, etc, excluding “ $\text{cr}(E) < \rho_{k+1}(M_v^{\mathcal{W}})$ ”, hold regarding $M, \bar{\mathcal{W}}, \bar{\mathcal{X}}, \bar{p}$, etc. Moreover, $\text{cr}(\bar{E}) < \rho_k(M_v^{\bar{\mathcal{W}}})$.*
- (b) *$\bar{E} \in \mathbb{E}_+(M_\infty^{\bar{\mathcal{X}}})$ iff $E \in \mathbb{E}_+(M_\infty^{\mathcal{X}})$.*
- (c) *$\bar{\mathcal{X}}$ is a hull of \mathcal{X} (see [6]), via hull maps $\pi \upharpoonright \text{lh}(\bar{\mathcal{X}}) : \text{lh}(\bar{\mathcal{X}}) \rightarrow \text{lh}(\mathcal{X})$ and $\pi_\alpha : M_\alpha^{\bar{\mathcal{X}}} \rightarrow M_{\pi(\alpha)}^{\mathcal{X}}$, where $\pi_\alpha \circ \bar{\varsigma}_\alpha = \varsigma_{\pi(\alpha)} \circ \pi \upharpoonright \bar{M}_\alpha$.*
- (d) *π_α is a near $\deg^{\bar{\mathcal{X}}}(\alpha)$ -embedding.*

Proof sketch. The premousehood of $(\bar{P} \parallel \text{lh}(\bar{E}), \bar{E})$ uses that the premouse axioms are Q-formulas, E^* is $\text{r}\Sigma_{k+1}^N(\{e\})$, $M_{\zeta'}^{\mathcal{X}}$ is $\text{r}\Delta_{k+1}^N(\{p, \zeta'\})$, and π is $\text{r}\Sigma_{k+1}$ elementary. The rest of (a) and (b) are similar. Parts (c) and (d) are proved by induction through $\text{lh}(\bar{\mathcal{X}})$; cf. [15, §2.3]. Here part (d) involves an adaptation of the argument in [8], or alternatively uses the uniformity of the definition of Sat^* in 3.23.⁵⁰ \square

Let $\iota = \text{lgcd}(P \parallel \text{lh}(E))$. Let \mathfrak{P} be the phalanx $\Phi(\bar{\mathcal{X}}, \bar{\iota}, \bar{E})$ and \mathfrak{P}' the phalanx $\Phi(\bar{\mathcal{W}}, \bar{\iota}, \bar{E})$. Note that $\mathfrak{P}, \mathfrak{P}'$ have the same last model $\bar{U} = M_\infty^{\bar{\mathcal{X}^+}} = M_\infty^{\bar{\mathcal{W}^+}}$. By 2.38, the following claim completes the proof.

Claim 2. *\bar{U} is wellfounded and \mathfrak{P}' is $(\omega_1 + 1)$ -iterable.*

Proof. The $(\omega_1 + 1)$ -iterability of \mathfrak{P}' follows easily from that of \mathfrak{P} , and we prove the latter. We lift trees on \mathfrak{P} to trees on $\Phi(\mathcal{X})$, using the maps π_α from Claim 1(c), and a weak k -embedding $\psi : \bar{U} \rightarrow M'$ with some $M' \triangleleft P \parallel \text{lh}(E)$. We will get M' and ψ through a variant of the earlier arguments. Let $v' = \text{pred}^{\mathcal{X}}(\zeta' + 1)$. We have

$$\text{cr}(E) < \rho_{k+1}(M_{v'}^{\mathcal{X}}) = \sup i_{0v'}^{\mathcal{X}} \text{ “} \rho_{k+1}^N \text{”}.$$

Let $\mu < \rho_{k+1}^N$ be a cardinal of N with $\text{cr}(E) \leq i_{0v'}^{\mathcal{X}}(\mu)$. Again using k -reasonableness let $q_1 \in \text{OR}(N)^{<\omega}$ witness 3.1 with respect to (N, k, q_0, μ) . Let

$$K_\mu = \text{cHull}_{k+1}^N(\mu \cup \{q_1\}).$$

⁵⁰The proof of uniformity itself used the argument of [8].

So $K_\mu \triangleleft N \parallel \rho_{k+1}^N$. We have the natural near k -embeddings

$$\sigma : M \rightarrow K_\mu \text{ and } \tau : K_\mu \rightarrow N,$$

with $\pi = \tau \circ \sigma$. Let $\tau(\hat{q}_0) = q_0$. Now

$$N \models \forall \varphi \left[\varphi \text{ is } \mathbf{r}\Sigma_{k+1} \Rightarrow \forall z \in \mu^{<\omega} \left[\varphi(q_0, z, u_k) \Rightarrow K_\mu \models \varphi(\hat{q}_0, z, u_k^{K_\mu}) \right] \right],$$

and this is an $\mathbf{r}\Pi_{k+1}$ assertion $\psi(q_0, \hat{q}_0, \mu, K_\mu, u_k)$.⁵¹ Let $U = M_\infty^{\mathcal{X}^+}$. We have virtual k -embeddings

$$j = i^{\mathcal{X}^+} : N \rightarrow U \text{ and } \bar{j} = i^{\bar{\mathcal{X}}^+} : M \rightarrow \bar{U}.$$

So $j(K_\mu) \triangleleft U$ and $U \models \psi(j(z))$ where $z = (q_0, \hat{q}_0, \mu, K_\mu, u_k)$.

Let $\varrho : \bar{U} \rightarrow U$ be given by the Shift Lemma. So ϱ is a virtual weak k -embedding.⁵² Define a (putative) virtual weak k -embedding $\varrho' : \bar{U} \rightarrow j(K_\mu)$ as follows. First set

$$\varrho' \circ \bar{j} = j \circ \sigma.$$

Now $\bar{U} = \text{Hull}_k^{\bar{U}}(\text{rg}(\bar{j}) \cup \nu(\bar{E}))$ and

$$\text{rg}(\varrho) = \text{Hull}_k^U(\text{rg}(\varrho \circ \bar{j}) \cup \varrho^* \nu(\bar{E})).$$

We set

$$\text{rg}(\varrho') = \text{Hull}_k^{j(K_\mu)}(\text{rg}(\varrho' \circ \bar{j}) \cup \varrho^* \nu(\bar{E})).$$

Note that the two hulls are isomorphic (to \bar{U}), and ϱ' is a virtual weak k -embedding, because $U \models \psi(j(z))$ and $\varrho^* \nu(\bar{E}) \subseteq j(\mu)$.

Now $U \models \text{"}j(K_\mu) \text{ is reasonable"}$. So let $q_2 \in j(K_\mu)^{<\omega}$ witness this for $(j(K_\mu), k, r, \iota)$ where $\iota = \text{lgcd}(P \parallel \text{lh}(E))$ and $r = (j(\hat{q}_0), \nu(E), \mathbf{c})$. Let

$$M' = \text{cHull}_{k+1}^{j(K_\mu)}(\iota \cup \{q_2\})$$

and $\varsigma : M' \rightarrow j(K_\mu)$ be the uncollapse. Note $M' \triangleleft P \parallel \text{lh}(E)$ (hence is wellfounded) and $\text{rg}(\varrho') \subseteq \text{rg}(\varsigma)$. Define $\psi : \bar{U} \rightarrow M'$ by $\psi = \varsigma^{-1} \circ \varrho'$. Then ψ is a weak k -embedding, and the rest is like before. \square (Claim 2)(3.9)

⁵¹It's not clear that this would hold if we allowed φ to be *generalized* $\mathbf{r}\Sigma_{k+1}$.

⁵²It's not clear yet that ϱ is a virtual near k -embedding, because the proof of [8] depends on strong closeness with respect to $(M_\epsilon^{\mathcal{X}}, \bar{E}), (M_\epsilon^{\mathcal{X}}, E)$, but here, the measures of E, \bar{E} are ostensibly not definable at all over $M_\epsilon^{\mathcal{X}}, M_\epsilon^{\bar{\mathcal{X}}}$.

4. Definability of \mathbb{E}

Given a mouse M and $x \in M$, we can ask whether \mathbb{E}^M is definable over $\lfloor M \rfloor$ from x . If so,⁵³

$$\lfloor M \rfloor \models "V = \text{HOD}_x".$$

Steel showed that for $n \leq \omega$, $K^{M_n} = M_n$, so \mathbb{E}^{M_n} is $\lfloor M_n \rfloor$ -definable (from $x = \emptyset$, i.e. without parameters); see [10] for $n < \omega$. In this context, K is fully iterable “in intervals between Woodin cardinals”. As we show in 4.13, this degree of self-iterability fails for mice with a measurable limit of Woodins. The author does not know if one can prove a generalization of Steel’s result at this level.

We next prove, in 4.3 and 4.7, that \mathbb{E}^M is nonetheless $\lfloor M \rfloor$ -definable for various mice M with measurable limits of Woodins, and also give a new proof for M_n .⁵⁴ Both theorems require a certain degree of self-iterability; 4.7 requires less, but requires also that the internal strategy agrees with a fuller external strategy for the mouse. We then prove in 4.9 that 4.7 applies in particular to $M_{\text{nt}}|\delta$, where M_{nt} is the least non-tame mouse (see 4.4) and δ is its largest (Woodin) cardinal. This approach to proving “ $V = \text{HOD}$ ” (in a mouse) is essentially limited to tame mice (cf. 4.12), as the degree of self-iterability required typically fails beyond there.

We remark that, using a quite different approach, Woodin has proved the following weaker version of 4.3. Let M be a tame mouse satisfying PS, and having no Woodin cardinals. Suppose $M \models$ “Every proper segment of me is fully iterable”. Then $\lfloor M \rfloor \models "V = \text{HOD}"$. However, Woodin’s proof does not seem to show that \mathbb{E}^M is definable over $\lfloor M \rfloor$. Also, Nam Trang and Martin Zeman have recently found a proof of self-iterability sufficient to apply 4.3, very different from what we do here, and which works in a different context.

Lemma 4.1. *Let N be a premouse. Let $\gamma < \text{OR}^N$ be an N -cardinal. Suppose $N \models \text{“}\gamma^{++} \text{ exists”}$. Let $R, S \in N$ be reasonable premice such that*

$$\lfloor R \rfloor = \lfloor S \rfloor = (\mathcal{H}_{\gamma^+})^N.$$

Let $\mathfrak{c}, \mathfrak{d} < (\gamma^+)^N$ be cutpoints of R, S respectively. Suppose $R|\mu = S|\mu$ where $\mu = \max(\mathfrak{c}, \mathfrak{d})$, and $N \models \text{“}R(S) \text{ is above-}\mathfrak{c} \text{ (above-}\mathfrak{d}), (\omega, \gamma^+ + 1)\text{-iterable”}$. Then $R = S$.

Proof. We work in N . Suppose $R \neq S$, and let η be least such that

$$R|(\eta^+)^N \neq S|(\eta^+)^N.$$

Let $R' = R|(\eta^+)^N$ and $S' = S|(\eta^+)^N$. Let Σ be an above- \mathfrak{c} , $(\omega, \eta^+ + 1)$ -strategy for R' , and Γ likewise but above- \mathfrak{d} for S' . Consider the (possibly

⁵³One might want to assume that $\lfloor M \rfloor \models \text{ZFC}$, in order to know that HOD_x is defined as usual.

⁵⁴Of course, the fact that $K^{M_n} = M_n$ gives more information than just the fact that \mathbb{E}^{M_n} is definable over $\lfloor M_n \rfloor$; and significantly, the “ K ” in Steel’s result has a corresponding generic absoluteness which does not seem to be present in our results here.

partial) comparison $(\mathcal{T}, \mathcal{U})$ of R' with S' , formed using Σ, Γ . (If the comparison reaches length $\eta^+ + 1$, then we stop.) We take $(\mathcal{T}, \mathcal{U})$ to be padded as usual.

We claim $(\mathcal{T}, \mathcal{U})$ is above η . For $\eta < \text{lh}(E)$ and $\max(\mathfrak{c}, \mathfrak{d}) < \text{lh}(E)$ for all extenders E used in $(\mathcal{T}, \mathcal{U})$. Suppose α is such that $(\mathcal{T}, \mathcal{U}) \restriction \alpha + 1$ is above η but $\text{cr}(E_\alpha^\mathcal{U}) < \eta$. Then by 3.6, $E_\alpha^\mathcal{U} \in \mathbb{E}_+(M_\alpha^\mathcal{T})$, so $M_\alpha^\mathcal{T} \restriction \text{lh}(E_\alpha^\mathcal{U}) = M_\alpha^\mathcal{U} \restriction \text{lh}(E_\alpha^\mathcal{U})$, contradiction.

Now suppose \mathcal{T} is non-trivial. Then there is $Q \triangleleft R$ such that $\rho_\omega^Q = \eta$ and \mathcal{T} can be considered a tree on Q . This is because \mathcal{T} is above η . Moreover, $b^\mathcal{T}$ drops. The same things hold for \mathcal{U} . Therefore the comparison is successful (and has length $< (\eta^+)^N$), and (still assuming \mathcal{T} is non-trivial), \mathcal{U} is trivial, and $S \triangleleft M_\infty^\mathcal{T}$. But then $E_0^\mathcal{T} \in [R] = [S] \in M_\infty^\mathcal{T}$, contradiction. \square

Lemma 4.2. *Let N be a premouse and let $\eta < \text{OR}^N$ be an N -cardinal such that $N \models \text{"}\eta^{++} \text{ exists"}$. Suppose $N \restriction (\eta^+)^N$ is reasonable, \mathfrak{c} is a cutpoint of $N \restriction (\eta^+)^N$ and*

$$N \models \text{"}N \restriction \eta^+ \text{ is above-}\mathfrak{c}, (\omega, \eta^+ + 1)\text{-iterable"}$$

Let $\mathcal{H} = (\mathcal{H}_{\eta^{+3}})^N$ (if $(\eta^{+3})^N = \text{OR}^N$ then set $\mathcal{H} = [N]$). Then

- (a) $\{N \restriction (\eta^+)^N\}$ is a $\Sigma_2^\mathcal{H}(\{N \restriction \mathfrak{c}\})$ singleton, uniformly in N , η and \mathfrak{c} .
- (b) If $\mathfrak{c} \leq \eta$ then $\{N \restriction (\eta^+)^N\}$ is a $\Sigma_2^\mathcal{H}(\{N \restriction \eta\})$ singleton, uniformly in N , η .

Proof. Part (a): By 4.1, $\mathcal{H} \models \text{"}N \restriction \eta^+ \text{ is the unique reasonable premouse } R \text{ such that (i) } [R] = \mathcal{H}_{\eta^+}, \text{ (ii) } N \restriction \mathfrak{c} \triangleleft R, \text{ (iii) } \mathfrak{c} \text{ is a cutpoint of } R, \text{ and (iv) } R \text{ is above-}\mathfrak{c}, (\omega, \eta^+ + 1)\text{-iterable.}"$

Part (b): If $\mathfrak{c} \leq \eta$ then by 4.1, $\mathcal{H} \models \text{"}N \restriction \eta^+ \text{ is the unique reasonable premouse } R \text{ such that } N \restriction \eta \triangleleft R \text{ and for some } \mathfrak{c} \leq \eta, \text{ items (i), (iii) and (iv) above hold.}"$ \square

We can now deduce the first main theorem of this section. In the theorem, if $N \not\models \text{ZFC}$ then we take HOD^N to be the union of all transitive sets coded by a set of ordinals X such that for some $\gamma < \text{OR}^N$, X is definable from ordinal parameters over \mathcal{H}_γ^N .

Theorem 4.3. *Let N be a premouse satisfying PS, all of whose proper segments are reasonable. Suppose that for each N -cardinal $\eta < \text{OR}^N$ there is $\mathfrak{c} \leq \eta$ such that \mathfrak{c} is a cutpoint of N and*

$$N \models \text{"}N \restriction \eta^+ \text{ is above-}\mathfrak{c}, (\omega, \eta^+ + 1)\text{-iterable"}$$

Then \mathbb{E}^N is definable over $[N]$ and $[N] \models \text{"}V = \text{HOD"}$. In fact, $\{N \restriction (\eta^+)^N\}$ is $\Sigma_2^\mathcal{H}$ where $\mathcal{H} = (\mathcal{H}_{\eta^{+3}})^N$, uniformly in η .

Proof. Using 4.2(b), we can define $N \restriction \eta$, for N -cardinals η , by induction on η , uniformly in η . (Given $N \restriction \eta$, we determine $N \restriction (\eta^+)^N$ by an application of 4.2(b).) Moreover, the induction leading to and producing $N \restriction (\eta^+)^N$ is in fact Σ_2 -definable over $(\mathcal{H}_{\eta^{+3}})^N$ (from no parameters). \square

We can now give an alternate proof of Steel's result that $\lfloor M_n \rfloor \models "V = \text{HOD}"$ for $n \leq \omega$: Theorem 4.3 applies to $N = M_n$, using \mathfrak{c} as the supremum of the Woodin cardinals δ of N such that $\delta \leq \eta$. (See [17] for the proof of self-iterability in the $n < \omega$ case, and [19, §7] for the $n = \omega$ case.) Now recall:

Definition 4.4. A premouse M is *non-meek* iff there is $E \in \mathbb{E}_+^M$ such that $M \restriction \text{cr}(E) \models$ "There is a proper class of Woodins". A premouse M is *non-tame* iff there is $E \in \mathbb{E}_+^M$ and δ such that $\text{cr}(E) \leq \delta \leq \nu(E)$ and $M \restriction \text{lh}(E) \models$ " δ is Woodin". *Meek/tame* means not non-meek/non-tame. We write M_{nt} for the least non-tame, sound, $(\omega, \omega_1 + 1)$ -iterable mouse, assuming such exists. \dashv

We show below that if a non-tame mouse exists then M_{nt} is well-defined; that is, there is a unique minimal sound non-tame mouse M which projects to ω . We will show that $\lfloor M_{\text{nt}} \rfloor$ can define $\mathbb{E}^{M_{\text{nt}} \restriction \lambda}$, but we need a variant of the results above to achieve this, as in this case the self-iterability hypothesis of 4.3 fails.

Definition 4.5. Let N be a premouse and Σ a (possibly partial) iteration strategy for N . We say that Σ is *extender-maximal* iff for every

$$(k, \mathcal{W}, \zeta, P, E, \mathcal{W}^+)$$

potential for N , with \mathcal{W} via Σ , $\text{pred}^{\mathcal{W}^+}(\zeta + 1) = 0$, and $b^{\mathcal{W}^+}$ does not drop in model or degree, then $E \in \mathbb{E}_+(M_\zeta^{\mathcal{W}})$. \dashv

Lemma 4.6. Let R, S be passive sound premice with $\lfloor R \rfloor = \lfloor S \rfloor$, with largest cardinal η , and $R \restriction \eta = S \restriction \eta$. Suppose there are extender-maximal, above- η , $(\omega, \eta^+ + 1)$ -iteration strategies for R, S . Then $R = S$.

Proof. This is like 4.1. Let Σ_R, Σ_S be strategies witnessing the assumption. Compare (R, S) using (Σ_R, Σ_S) , producing trees $(\mathcal{T}, \mathcal{U})$. It suffices to see that $(\mathcal{T}, \mathcal{U})$ is above η . This follows from extender-maximality. \square

Analogously to Theorem 4.3, we now conclude:

Theorem 4.7. Let N be a premouse satisfying PS, and Σ a $(0, \theta^+ + 1)$ -strategy for N , where $\theta = \text{card}(N)$. For an N -cardinal η , let Σ_η be the partial strategy induced by Σ for trees which are in N , on $N \restriction (\eta^+)^N$, above η , and of length $\leq (\eta^+)^N$. Suppose that $\Sigma_\eta \in N$ for each such η . Then $N \models$ " Σ_η is extender-maximal" and \mathbb{E}^N is definable over $\lfloor N \rfloor$, so $\lfloor N \rfloor \models "V = \text{HOD}"$.

Proof. Because N is iterable, all its proper segments are reasonable. Note that because $\Sigma_\eta \in N$, N is closed under Σ_η and $N \models$ " Σ_η is an above- η , $(\omega, \eta^+ + 1)$ -strategy for $N \restriction \eta^+$ ". The extender-maximality of Σ_η follows from 3.8.

Now work in $\lfloor N \rfloor$; we define \mathbb{E}^N . We identify $N \restriction \eta$ inductively on N -cardinals η . Suppose we have $N \restriction \eta$. Then by 4.6 and the properties of Σ_η , note that $N \restriction \eta^+$ is the unique premouse R such that $\lfloor R \rfloor = \mathcal{H}_{\eta^+}$ and $R \restriction \eta = N \restriction \eta$ and there is an extender-maximal, above- η , $(\omega, \eta^+ + 1)$ -iteration strategy for R . \square

We will next show that whenever λ is a limit cardinal of M_{nt} , then the hypotheses of 4.7 apply to $M_{\text{nt}}|\lambda$, with Σ the $(0, \omega_1 + 1)$ -strategy for $M_{\text{nt}}|\lambda$ induced by the unique $(\omega, \omega_1 + 1)$ -strategy for M_{nt} . The general method of proof is standard, but some non-standard details are involved. The method also works for many mice simpler than M_{nt} .

Lemma 4.8. (a) *Let N be a non-tame premouse with no non-tame proper segment. Let $X \subseteq \mathfrak{C}_0(N)$. Then $\text{cHull}_1^N(X)$ is non-tame. Therefore if N is $(0, \omega_1 + 1)$ -iterable then M_{nt} is well-defined,*

$$M_{\text{nt}} = \text{cHull}_1^N(\emptyset),$$

$\rho_1(M_{\text{nt}}) = \omega$ and $p_1(M_{\text{nt}}) = \emptyset$.

(b) *Let M be a type 3 premouse with $\nu(M)$ regular in M . Let $\nu(M) = [a, f]_{F^M}^M$ with $a, f \in M$. Let \mathcal{T} be a 0-maximal tree on M such that $i^{\mathcal{T}}$ exists. Let $R = M_{\infty}^{\mathcal{T}}$. Then*

$$\nu(R) = [i^{\mathcal{T}}(a), i^{\mathcal{T}}(f)]_{F^R}^R.$$

Therefore if M is non-tame then so is R .

Proof. (a) N is type 3 and $\nu(F^N) = [\{\kappa\}, f]_{F^N}^N$ where $\kappa = \text{cr}(F^N)$ and

$$f : \kappa \rightarrow \kappa$$

is such that $f(\alpha)$ is the least Woodin δ of N such that $\delta > \alpha$. Now $\kappa, f \in \text{Hull}_1^N(X)$. Let $H = \text{cHull}_1^N(X)$ and let $\bar{\kappa}, \bar{f} \in H$ be the collapses of κ, f . One can show

$$\nu(F^H) = [\{\bar{\kappa}\}, \bar{f}]_{F^H}^H.$$

Therefore H is non-tame.

(b) Let

$$\psi : \text{Ult}(M^{\text{sq}}, F^M) \rightarrow \text{Ult}(R^{\text{sq}}, F^R)$$

be induced by $i^{\mathcal{T}}$; i.e.,

$$\psi([a, f]_{F^M}^{M^{\text{sq}}}) = [i^{\mathcal{T}}(a), i^{\mathcal{T}}(f)]_{F^R}^{R^{\text{sq}}}.$$

We need to see that ψ is continuous at $\nu(F^M)$. So let $\gamma < \psi(\nu(F^M))$. Let $b, g \in R^{\text{sq}}$ be such that $\gamma = [b, g]_{F^R}^{R^{\text{sq}}}$. Because \mathcal{T} is only 0-maximal, there is $f \in M^{\text{sq}}$ and some a such that $(b, g) = i^{\mathcal{T}}(f)(a)$. Therefore there are $h \in M^{\text{sq}}$ and $\alpha < \nu(F^R)$ such that

$$\gamma \in i_{F^R} \circ i^{\mathcal{T}}(h) \text{ `` } [\alpha]^{<\omega}.$$

There is $\beta < \nu(F^M)$ such that $\alpha < i^{\mathcal{T}}(\beta)$. It follows that

$$\gamma \in \psi(i_{F^M}(h) \text{ `` } [\beta]^{<\omega}).$$

We may assume

$$i_{F^M}(h) \text{ `` } [\beta]^{<\omega} \subseteq \nu(F^M).$$

But $\nu(F^M)$ is regular in $\text{Ult}(M, F^M)$, so $i_{F^M} \text{ `` } [\beta]^{<\omega}$ is bounded in $\nu(F^M)$. It follows that $\gamma < \nu(F^R)$, as required. \square

Theorem 4.9. Assume M_{nt} exists (so is $(\omega, \omega_1 + 1)$ -iterable). Let Σ be the unique $(\omega, \omega_1 + 1)$ -strategy for M_{nt} . Let $\eta < \text{OR}^{M_{\text{nt}}}$ be an M_{nt} -cardinal and \mathfrak{c} be the sup of all $\delta \leq \eta$ such that $\delta = 0$ or δ is Woodin in M_{nt} . (So \mathfrak{c} is a cutpoint of $M_{\text{nt}} \parallel \text{OR}(M_{\text{nt}})$, but maybe not a strong cutpoint.) Suppose $\mathfrak{c} < \eta$. Let $\lambda > \eta$ be a limit cardinal of M_{nt} .

If $\eta = (\mathfrak{c}^+)^{M_{\text{nt}}}$, let $\xi = \mathfrak{c}$; if $\eta > (\mathfrak{c}^+)^{M_{\text{nt}}}$, let $\xi = (\mathfrak{c}^+)^{M_{\text{nt}}}$. Let Γ be the partial strategy induced by Σ for 0-maximal above- ξ trees on $M_{\text{nt}} \restriction \eta$ which are in $M_{\text{nt}} \restriction \lambda$.

Then Γ is a class of $M_{\text{nt}} \restriction \lambda$.

Therefore $M_{\text{nt}} \restriction \lambda \models \text{“}\Gamma \text{ is an extender-maximal, above-}\xi, (0, \lambda)\text{-strategy for } M_{\text{nt}} \restriction \eta\text{”}$, and $\mathbb{E}^{M_{\text{nt}} \restriction \lambda}$ is definable over $[M_{\text{nt}} \restriction \lambda]$.

Remark 4.10. A draft of this paper claimed that $M_{\text{nt}} \restriction \lambda \models \text{“}Z \restriction \eta \text{ is above-}\mathfrak{c}, \eta\text{-iterable”}$. But this is false for some $\mathfrak{c}, \eta, \lambda$, as we will show in 4.13. The earlier putative proof of this claim ignored the fact that if \mathfrak{c} is measurable in M_{nt} , hence a limit of Woodins, and E is an M_{nt} -total extender with $\text{cr}(E) = \mathfrak{c}$, then $\text{Ult}(M_{\text{nt}} \restriction \eta, E)$ has Woodins above $\text{lh}(E)$, which means that correct \mathbb{Q} -structures for trees using such an E can be non-tame.

Proof. The last sentence follows from 3.8 (which gives the extender-maximality) and 4.7. So we just need to see that Γ is a class of $M_{\text{nt}} \restriction \lambda$. We may easily assume that η is a successor cardinal of M_{nt} and $\lambda \leq \kappa = \text{cr}(F^{M_{\text{nt}}})$. We will consider first the case that $\lambda = \kappa$, and then deduce the general case from this one. Let $N = M_{\text{nt}} \restriction \eta$ and $Z = M_{\text{nt}} \restriction \kappa$.

Note that Γ is everywhere \mathbb{Q} -structure guided (as η is a successor cardinal and \mathfrak{c} is a cutpoint; \mathfrak{c} might be a measurable limit of Woodins, but if $(\mathfrak{c}^+)^{M_{\text{nt}}} < \eta$ then the trees we consider are above $\xi = (\mathfrak{c}^+)^{M_{\text{nt}}}$).

Let $\mathcal{W} \in Z$ be a limit length, above- ξ tree on N , via Γ . Work in Z . Let $\mathbb{C} = \langle N_\alpha \rangle_{\alpha \leq \Omega}$ be the maximal fully backgrounded $L[\mathbb{E}]$ -construction above $N_0 = M(\mathcal{W})$, with all background extenders E^* such that $E^* \in \mathbb{E}^Z$ and $\text{cr}(E^*) > \delta(\mathcal{W})$, and where Ω is least such that either $\Omega = \kappa$ or the $\delta(\mathcal{W})$ -core Q of N_Ω is a \mathbb{Q} -structure for $M(\mathcal{W})$. (Here each N_α is a premouse with $M(\mathcal{W}) \trianglelefteq N_\alpha$ and $\delta(\mathcal{W})$ a cutpoint, not just an $M(\mathcal{W})$ -premouse.)

If $\Omega < \kappa$ then in Z we can use Q to determine b as usual, completing the proof in this case (that $\lambda = \kappa$). So assume $\Omega = \kappa$ for a contradiction. Let $H = F^{M_{\text{nt}}}$, let $R = i_H(N_\kappa)$, let $\delta = \nu(H)$, let $G = H \restriction R \times [\delta]^{<\omega}$, let

$$U = \text{Ult}(R \restriction (\kappa^+)^R, G),$$

let $S = U \restriction (\delta^+)^U = R \restriction (\delta^+)^U$, let G' be the trivial completion of G , and $Q = (S, G')$. Given an iterate M' of M_{nt} and iteration map $j : M_{\text{nt}} \rightarrow M'$ with $\delta(\mathcal{W}) < \text{cr}(j)$, let $Q(M')$ denote “ $j(Q)$ ”, that is, the structure defined over M' from $M(\mathcal{W})$ in the same manner as we have just defined Q over M_{nt} from $M(\mathcal{W})$.

Claim. Q is a non-tame premouse and is $(0, \omega_1 + 1)$ -iterable above $\delta(\mathcal{W})$.

Assume the claim. If $M(\mathcal{W}) \neq Q$ then $M(\mathcal{W}) \triangleleft Q$ and $\text{OR}^{M(\mathcal{W})}$ is Woodin in Q . So comparing Q with the tame $\delta(\mathcal{W})$ -sound \mathbb{Q} -structure extending $M(\mathcal{W})$ leads to contradiction, using 4.8(b). So the following completes the proof in this case:

Proof of Claim. In this proof we write $\delta(P)$ to denote $\text{lgcd}(P)$, for premiss P .

Non-tameness and the ISC for Q is via the usual proof that Woodinness is absorbed by $L[\mathbb{E}]$ -constructions. For iterability, we define a 0-maximal strategy for Q , lifting trees \mathcal{T} on Q to 0-maximal trees \mathcal{T}' on M_{nt} via our strategy for M_{nt} , using a variant of the lift/resurrect procedure of [3, §12]. We assume the reader is familiar with that procedure, and just explain the differences. The differences arise because Q is not built by a standard $L[\mathbb{E}]$ -construction: S is built by $i_F(\mathbb{C})$ in $\text{Ult}(M_{\text{nt}}, F)$, but F^Q is not available to $\text{Ult}(M_{\text{nt}}, F)$.

We will decompose \mathcal{T} into the form $\mathcal{U} \smallfrown \mathcal{V}$, where $(\mathcal{U}, \mathcal{V})$ is a stack of two normal trees, and decompose \mathcal{T}' into the form $\mathcal{U}' \smallfrown \langle F \rangle \smallfrown \mathcal{V}'$, also a stack of two normal trees $(\mathcal{U}' \smallfrown \langle F \rangle, \mathcal{V}')$. We lift \mathcal{U} to \mathcal{U}' via almost the usual procedure. The tree \mathcal{V} begins at the least θ such that the appropriate ancestor for E_θ^T is not directly available to $\mathfrak{C}_0(M_\theta^{T'})$. We set $F = F(M_\theta^{T'})$, and then lift \mathcal{V} to \mathcal{V}' via the usual procedure.

The tree \mathcal{U}' results from lifting/resurrection of \mathcal{U} to M_{nt} , except that for each $\alpha < \text{lh}(\mathcal{U})$, if $[0, \alpha]_{\mathcal{U}}$ does not drop then the lift of $M_\alpha^{\mathcal{U}}$ is $Q(M_\alpha^{\mathcal{U}'})$ (the “image” of Q), whereas if $[0, \alpha]_{\mathcal{U}}$ drops then the lift of $M_\alpha^{\mathcal{U}}$ is $\mathfrak{C}_{n_\alpha}(N_{\gamma_\alpha}^{\mathbb{C}_\alpha})$, where

$$\mathbb{C}_\alpha = i_{F(M_\alpha^{\mathcal{U}'})}^{M_\alpha^{\mathcal{U}'}}(i_{0, \alpha}^{\mathcal{U}'}(\mathbb{C})),$$

$\gamma_\alpha < \text{lh}(\mathbb{C}_\alpha)$, and $n_\alpha = \deg^T(\alpha)$. So the lift is in $\text{Ult}(M_\alpha^{\mathcal{U}'}, F(M_\alpha^{\mathcal{U}'}))$, but may not be in $M_\alpha^{\mathcal{U}'}$. Likewise for partial resurrections. Resurrection, if necessary, is computed in $\text{Ult}(M_\alpha^{\mathcal{U}'}, F(M_\alpha^{\mathcal{U}'}))$.

We choose \mathcal{U} as long as possible such that for all $\alpha + 1 < \text{lh}(\mathcal{U})$,

- (i) if $[0, \alpha]_{\mathcal{U}}$ does not drop then either $E_\alpha^{\mathcal{U}} \in \mathfrak{C}_0(M_\alpha^{\mathcal{U}})$ or $E_\alpha^{\mathcal{U}} = F(M_\alpha^{\mathcal{U}})$,⁵⁵ and
- (ii) letting $E^* = E_\alpha^{\mathcal{U}'}$, the ancestor of the lift of $E_\alpha^{\mathcal{U}}$, either $E^* \in \mathfrak{C}_0(M_\alpha^{\mathcal{U}'})$ or $E^* = F(M_\alpha^{\mathcal{U}'})$ (the latter occurs just when $[0, \alpha]_{\mathcal{U}}$ does not drop and $E_\alpha^{\mathcal{U}} = F(M_\alpha^{\mathcal{U}})$).

Let $\beta = \text{pred}^{\mathcal{U}}(\alpha + 1)$ and $k = \deg^{\mathcal{U}}(\alpha + 1)$. Let

$$\sigma : M_{\alpha+1}^{*\mathcal{U}} \rightarrow \mathfrak{C}_k(N_\eta^{\mathbb{C}_\beta})$$

be the partial resurrection map determined by $\text{cr}(E_\alpha^{\mathcal{U}})$. If $\mathfrak{C}_k(N_\eta^{\mathbb{C}_\alpha}) \in \mathfrak{C}_0(M_\alpha^{\mathcal{U}'})$ or $\mathfrak{C}_k(N_\eta^{\mathbb{C}_\alpha}) = \mathfrak{C}_0(Q(M_\alpha^{\mathcal{U}'}))$ then define $\pi_{\alpha+1}$ as usual. Suppose otherwise. Then

$$\text{Ult}_0(\text{Ult}_0(M_\beta^{\mathcal{U}'}, F(M_\beta^{\mathcal{U}'})), E_\alpha^{\mathcal{U}'}) = \text{Ult}_0(M_{\alpha+1}^{\mathcal{U}'}, F(M_{\alpha+1}^{\mathcal{U}'}))$$

⁵⁵ As Q is type 3, when $[0, \alpha]_{\mathcal{U}}$ does not drop, we can have $\nu(M_\alpha^{\mathcal{U}}) < \text{lh}(E_\alpha^{\mathcal{U}}) < \text{OR}(M_\alpha^{\mathcal{U}})$, in which case (i) fails.

and the ultrapower maps commute. Let

$$j : \text{Ult}_0(M_\beta^{\mathcal{U}'}, F(M_\beta^{\mathcal{U}'})) \rightarrow \text{Ult}_0(M_{\alpha+1}^{\mathcal{U}'}, F(M_{\alpha+1}^{\mathcal{U}'}))$$

be the resulting ultrapower map (via $E_\alpha^{\mathcal{U}'}$). Then we define $\pi_{\alpha+1}$ in the usual way, except that j replaces $i_{\beta, \alpha+1}^{\mathcal{U}'}$.

Note that by our choice of \mathcal{U} , the first drop in model along any \mathcal{U} -branch is to some $M_{\alpha+1}^{*\mathcal{U}} \triangleleft \mathfrak{C}_0(M_\beta^{\mathcal{U}})$, where $\beta = \text{pred}^{\mathcal{U}}(\alpha + 1)$. So $M_{\alpha+1}^{*\mathcal{U}} \in \text{dom}(\pi_\beta)$. (But the resurrection of $\pi_\beta(M_{\alpha+1}^{*\mathcal{U}})$ may not be in $M_\beta^{\mathcal{U}'}$.)

If $\mathcal{V} = \emptyset$ then $F, \mathcal{V}' = \emptyset$.

Suppose $\mathcal{V} \neq \emptyset$. Then $\text{lh}(\mathcal{U})$ is a successor $\theta + 1$, and either (i) or (ii) above fails for $\alpha = \theta$. We set $E_\theta^{\mathcal{T}'} = F = F(M_\theta^{\mathcal{U}'})$. Let $\lambda = i_F(\text{cr}(F))$. We will define $\mathcal{T}' = \mathcal{U}' \hat{\wedge} \langle F \rangle \hat{\wedge} \mathcal{V}'$, indexed with

$$(0, 1, \dots, \theta, \theta^*, \theta + 1, \dots).$$

Let $M = M_\theta^{\mathcal{U}'}$. Then $M_{\theta^*}^{\mathcal{T}'}|_\lambda = \text{Ult}(M, F)|_\lambda$ and $i_{0, \theta^*}^{\mathcal{T}'}(\mathbb{C}) = C_\theta$. Suppose (i) fails for $\alpha = \theta$. Let

$$\psi : \text{Ult}(M_\theta^{\mathcal{U}}, F(M_\theta^{\mathcal{U}})) \rightarrow Q(\text{Ult}(M, F))$$

be induced by π_θ , through the Shift Lemma, and let $E^* = \psi(E_\theta^{\mathcal{U}})$. By 4.8(b), $\psi(\delta(M_\theta^{\mathcal{U}})) = \delta(M_\theta^{\mathcal{U}'})$, so $\delta(M_\theta^{\mathcal{U}'}) < \text{lh}(E^*)$. If instead (i) holds, use $\psi = \pi_\theta$ to lift $E_\theta^{\mathcal{U}}$ to an extender E^* . In either case let $E_{\theta^*}^{\mathcal{T}'}$ be the ancestor of E^* , according to \mathbb{C}_θ . Note that $\text{lh}(F)$ is a cutpoint of $M_{\theta^*}^{\mathcal{U}'}$, and $\text{lh}(F) < \text{cr}(E_{\theta^*}^{\mathcal{T}'})$.

Let σ be the map resurrecting E^* to $E_{\theta^*}^{\mathcal{T}'}$. Let $\gamma \in \text{OR}$ be least such that $\sigma(\psi(\gamma)) \geq \delta(M_\theta^{\mathcal{U}'})$. Then γ is a limit cardinal cutpoint of $M_\theta^{\mathcal{T}}|_{\text{lh}(E_\theta^{\mathcal{T}})}$. For the limit cardinality is clear; and if $G \in \mathbb{E}_+(M_\theta^{\mathcal{T}}|_{\text{lh}(E_\theta^{\mathcal{T}})})$ overlaps γ then the ancestor G' of $\sigma(\psi(G))$ is such that $\text{cr}(G') < \delta(M_\theta^{\mathcal{U}'}) < \text{lh}(G')$, contradicting the tameness of $\text{Ult}(M, F)|_\lambda$. Also, $\gamma < \text{lh}(E_\theta^{\mathcal{T}})$.

Let \mathcal{V} be the remainder of \mathcal{T} , so $\mathcal{T} = \mathcal{U} \hat{\wedge} \mathcal{V}$. Then by the previous paragraph, \mathcal{V} is on $M_\theta^{\mathcal{U}}$ and is above γ . We lift \mathcal{V} to a normal tree \mathcal{V}' on $M_{\theta^*}^{\mathcal{T}'}$, above $\delta(M_\theta^{\mathcal{T}'})$. For this we use the standard process; we need not lift to any $Q(M_\alpha^{\mathcal{V}'})$. This is because $[0, \beta]_{\mathcal{T}}$ drops in model for every β such that $\theta <_{\mathcal{T}} \beta$. For suppose $[0, \theta]_{\mathcal{T}}$ does not drop and let $\alpha + 1 < \text{lh}(\mathcal{T})$ with $\text{pred}^{\mathcal{T}}(\alpha + 1) = \theta$; then $\text{lh}(E_\theta^{\mathcal{T}})$ has cardinality $\leq \gamma$ in $M_\theta^{\mathcal{T}}$, and therefore $\alpha + 1 \in D^{\mathcal{T}}$. If (i) fails this is immediate, and (i) holds it is because $\gamma < \text{lh}(E_\theta^{\mathcal{T}}) < M_\theta^{\mathcal{T}}$ and $\psi(\gamma) < \delta(M_\theta^{\mathcal{T}'})$, but $\sigma(\psi(\gamma)) \geq \delta(M_\theta^{\mathcal{T}'})$, and σ results from resurrection of projecting structures (so if $(\gamma^+)^{M_\theta^{\mathcal{T}}} < \text{lh}(E_\theta^{\mathcal{T}})$, then $\sigma(\psi(\gamma)) = \psi(\gamma)$, contradiction). We leave the remaining details to the reader. \square (Claim)

Now consider the case that $\lambda \leq \kappa$ is a limit cardinal of M_{nt} . As before, we have $\mathcal{W} \in Z = M_{\text{nt}}|_\lambda$ and want to compute $\Gamma(\mathcal{W})$. We simply look for some R with $N \triangleleft R \triangleleft M_{\text{nt}}|_\lambda$ and $\mathcal{W} \in R \models \text{ZFC}$ and $R \models$ “My $L[\mathbb{E}]$ -construction above $M(\mathcal{W})$ reaches a Q-structure Q for $M(\mathcal{W})$ as above”. We use this Q to compute b . Since $M_{\text{nt}}|_\kappa$ has these properties and $\lambda \leq \kappa$ is a limit cardinal

of M_{nt} , by condensation there is also some such $R \triangleleft M|\lambda$, and any such R computes the correct Q-structure, so we are done. This completes the proof of the theorem. \square

Remark 4.11. The methods above can be adapted to many other canonical tame mice, for example, the least proper class mouse with an inaccessible limit of Woodin cardinals, etc.

It is known that under sufficient large cardinal assumptions and for sufficiently complex reals x , $\text{HOD}^{L[x]} \cap \mathbb{R} = M_1 \cap \mathbb{R}$. So if x is a sound mouse projecting to ω which extends $M_1^\#$, then $L[x]$ is a mouse satisfying “ $\mathbb{R} \not\subseteq \text{HOD}$ ”.

Steel asked: (1) Suppose M is a mouse satisfying ZFC. Does $[M]$ satisfy “ $V = \text{HOD}_X$ for some $X \subseteq \omega_1^M$ ”? Naturally we can also consider the variant of (1) in which we require X to be a real.

Some related questions are: (2) To what extent can the methods used to prove 4.9 be adapted to more arbitrary (non-“canonical”) tame mice? (3) Do or can non-tame mice know enough of their own iteration strategies in order that one can answer (1) using the methods of this paper?

It turns out that non-tame mice very typically do *not* know enough of their own iteration strategy to suffice for (3). This is demonstrated in 4.12 below, which is a simple variant of an observation possibly due to Steel; see [11, 1.1]. However, the author has since answered (1) positively; the proof uses methods quite different to those of this paper. Moreover, if M is tame then we may take X to be a real, but it is not known to the author whether the same holds for non-tame M . These results are to appear. The author has also shown that various canonical non-tame mice, for example M_{adr} , satisfy “ $V = \text{HOD}$ ”.

Proposition 4.12. *Let N be a countable premouse satisfying ZFC and Σ an $(\omega, \omega_1 + 1)$ -strategy for N . Let $P \triangleleft N$ be active and suppose τ, δ, η are such that*

$$\text{cr}(F^P) < \tau < \delta < \text{OR}^P \leq \eta < (\tau^+)^N,$$

τ is a cardinal of N , $P \models$ “ δ is Woodin”, and η is a cutpoint of N . Let Σ' be the strategy induced by Σ , for 0-maximal trees \mathcal{T} on P such that $\mathcal{T} \in N$, \mathcal{T} is above τ , and of length $\leq (\tau^+)^N$.⁵⁶ Then $\Sigma' \notin N$.

Proof. For simplicity we assume that $\delta < \nu_E$. The argument in this case can be adapted to the case that $\delta = \nu_E$, using part of the argument for 4.8.

Suppose $\Sigma' \in N$. Let κ be the least measurable of P such that $\kappa > \tau$. Let \mathbb{B} be the δ -generator extender algebra of P at δ , using extenders E with $\text{cr}(E) > \kappa$. Working in N , let \mathcal{T} on P be given by first iterating κ out past η and then making $N|\eta$ generic for $i^{\mathcal{T}}(\mathbb{B})$ over $M_\infty^{\mathcal{T}}$. Because $\Sigma' \in N$, this succeeds with $\mathcal{T} \in N$ of length $< (\tau^+)^N$. Let $E = F(M_\infty^{\mathcal{T}})$, so $\text{cr}(E) = \text{cr}(F^P) < \tau$, E is N -total and $\text{Ult}(N, E)$ is wellfounded, because Σ' is induced by Σ . Moreover,

⁵⁶That is, $\Sigma'(\mathcal{T})$ is defined for such \mathcal{T} of length $\leq (\tau^+)^N$.

$N|\eta$ is generic over $\text{Ult}(N, E)$. Let $Q \trianglelefteq N$ be least such that $\rho_\omega^Q = \tau$ and $i^\mathcal{T}(\delta) \leq \text{OR}^Q$. Then

$$Q \notin \text{Ult}(N, E)[N|\eta]$$

since δ is a cardinal in $\text{Ult}(N, E)[N|\eta]$. But $Q \in \text{Ult}(N, E)[N|\eta]$ as in the proof of [11, 1.1],⁵⁷ contradiction. \square (4.12)

We finally adapt the example above to show that the above- $(\mathfrak{c}^+)^{M_{\text{nt}}}$ iterability of $M_{\text{nt}}|\eta$ in M_{nt} established in 4.9 cannot in general be improved to above- \mathfrak{c} iterability, and that the same holds for typical non-meek mice. The example also indicates that if non-meek mice satisfy “ $V = K$ ” (whatever this means) then there must be new difficulties involved in the proof, as the core model K involved would be less iterable than that in the case of M_n :

Proposition 4.13. *Let N be a countable premouse satisfying ZFC and Σ a $(0, \omega_1 + 1)$ -strategy for N . Let \mathfrak{c} be a limit of Woodin cardinals of N and suppose there is $E \in \mathbb{E}^N$ such that*

$$\mathfrak{c} = \text{cr}(E) < (\mathfrak{c}^+)^N < \text{lh}(E).$$

Suppose there is a cutpoint \mathfrak{d} of N such that $\text{lh}(E) \leq \mathfrak{d} < \text{OR}^N$. Then

$$N \models \text{“}N|\text{lh}(E) \text{ is not above-}\mathfrak{c}, (0, \mathfrak{d}^+ + 1)\text{-iterable”}.$$

Proof. Suppose otherwise. Let $E \in \mathbb{E}^N$ be least witnessing the assumption. Let $\bar{U} = \text{Ult}(N|(\mathfrak{c}^+)^N, E)$. Then $i_E^N(\mathfrak{c})$ is a limit of Woodins of U . Let δ be the least Woodin of \bar{U} such that $\delta > \text{lh}(E) = (\mathfrak{c}^{++})^{\bar{U}}$. Form an above- $\text{lh}(E)$ tree \mathcal{U} on \bar{U} , first linearly iterating past \mathfrak{d} , and then iterating to make $N|\mathfrak{d}$ generic for the above- $\text{lh}(E)$ extender algebra of $M_\infty^\mathcal{T}$ at $\delta^* = i^\mathcal{U}(\delta)$. By the iterability assumption, this succeeds, with $\text{lh}(\mathcal{U}) < (\mathfrak{d}^+)^N$. Let $\mathcal{T} = E \restriction \mathcal{U}$. Now \mathcal{T} can be considered a tree \mathcal{T}' on N , with last model W (\mathcal{T}' has wellfounded models; otherwise pull a counterexample to wellfoundedness down below $N|(\mathfrak{c}^+)^N$ for a contradiction). Let $P \triangleleft N$ be least such that $\delta^* \leq \text{OR}^P$ and $\rho_\omega^P < \delta^*$. Let $P^* = i^{\mathcal{T}'}(P) \in W$.

Now δ^* is regular in $W[N|\mathfrak{d}]$, so $P \notin W[N|\mathfrak{d}]$. By absoluteness as in the proof of 4.12, it follows that in $W[N|\mathfrak{d}][G]$, where $G \subseteq \text{Col}(\omega, \delta^*)$ is $W[N|\mathfrak{d}]$ -generic, there are P_1, P_2, π_1, π_2 such that P_i is a sound premouse, $P_1 \not\trianglelefteq P_2 \not\trianglelefteq P_1$, $N|\mathfrak{d} \trianglelefteq P_i$, \mathfrak{d} is a cutpoint of P_i , $\rho_\omega^{P_i} < \delta^*$, and $\pi_i : P_i \rightarrow P^*$ is elementary. The existence of such objects embedded into P^* is therefore forced over W by $\text{Col}(\omega, \delta^*)$. This pulls back elementarily to N, P and some Woodin $\delta' < \mathfrak{c}$. But this contradicts the iterability of N in V . \square

⁵⁷Sketch: In the generic extension after collapsing η , we can search for a Q of the right form, extending $N|\eta$, and embedded into $i_E(Q)$. But then by uniqueness we get Q in the ground model.

5. Mixed bicephali

In this section we generalize Mitchell and Steel’s “Uniqueness of the next extender” result from [3, §9]:

Theorem 5.1. *Let R be an $(\omega_1 + 1)$ -iterable countable transitive model of ZFC. Work in R . Let $\mathbb{C} = \langle N_\alpha \rangle_{\alpha \leq \lambda}$ be a fully backgrounded $L[\mathbb{E}]$ -construction. Suppose there are fully backgrounded extenders $E, F \neq \emptyset$ such that (N_λ, E) and (N_λ, F) are premice. Then $E = F$.⁵⁸*

The assumption that $R \models \text{ZFC}$ is made for simplicity; one can as usual get by with less than this. In the case that E, F are either both type 2, or neither is type 2, this already follows from the bicephalus arguments in [3]. The remaining case, that say E is type 2 and F is not, was not considered in [3] (this case was not necessary for the results there). However, the uniqueness question in this case is of course also natural, and we verify it here.

In what follows we use $\delta(P) = \delta^P$ to denote $\text{lged}(P)$.

Definition 5.2. A *mixed bicephalus* is a structure (N, E, F) , where (N, E) is a type 2 premouse and (N, F) is a type 1 or type 3 premouse.

A *mis-bicephalus* is a structure (N, E, F) , where (N, E) is a type 2 premouse and (N, F) is a segmented-premouse.⁵⁹

Given a mixed or mis-bicephalus $B = (N, E, F)$, $F_0^B = E$ and $F_1^B = F$. \dashv

Remarks similar to those in 2.10, regarding fine structure and ultrapowers, also apply to mis-bicephali. Ultrapowers of such are always constructed at the unsquashed level. This works like for type 2 bicephali in [3, §9]. The reader should now recall the two examples of proper seg-pms in 2.10; in the comparison argument to follow we need to deal with these and more general examples.

In a normal iteration tree \mathcal{T} , the exchange ordinal $\nu_\alpha^{\mathcal{T}}$ associated with extender $E_\alpha^{\mathcal{T}}$ is usually the strict sup of generators $\nu(E_\alpha^{\mathcal{T}})$. For our trees on mixed bicephali, it is convenient to tweak this in special cases; the tweak makes the proof of iterability of bicephali slightly smoother.

Definition 5.3. Let N be a mixed bicephalus. The *maximal iteration game* on N is defined as usual, except that for $\alpha + 1 < \text{lh}(\mathcal{T})$: Set $\nu_\alpha^{\mathcal{T}} = \nu(E_\alpha^{\mathcal{T}})$, unless $[0, \alpha]_{\mathcal{T}}$ does not drop and $E_\alpha^{\mathcal{T}} = F_1(M_\alpha^{\mathcal{T}})$, in which case set $\nu_\alpha^{\mathcal{T}} = \delta(M_\alpha^{\mathcal{T}})$.⁶⁰ Then $\text{pred}^{\mathcal{T}}(\alpha + 1)$ is the least β such that $\text{cr}(E_\alpha^{\mathcal{T}}) < \nu_\beta^{\mathcal{T}}$. We say N is α -iterable iff player II has a winning strategy in the maximal game on N of length α . \dashv

⁵⁸We leave the precise interpretation of “fully backgrounded” to the reader; it should be enough to guarantee the $(0, \omega_1 + 1)$ -iterability of B in the proof.

⁵⁹“mis-” abbreviates “mixed segmented-”. *Segmented-premouse* is defined in 2.9.

⁶⁰Steve Jackson noticed that we could have instead set $\nu_\alpha^{\mathcal{T}} = \nu(E_\alpha^{\mathcal{T}})$ in all cases. However, doing so would slightly complicate the standard proof of iterability of the bicephalus (when the bicephalus is constructed by fully backgrounded $L[\mathbb{E}]$ -construction in an iterable background universe).

Lemma 5.4. *There is no $(\omega_1 + 1)$ -iterable mixed bicephalus.*

From now on we write *bicephalus* for *mis-bicephalus*.

Proof. Suppose P is otherwise. We may assume P is countable. As in [3, §9], we compare P with itself, producing padded 0-maximal trees \mathcal{T}, \mathcal{U} . For $\alpha + 1 < \text{lh}(\mathcal{T})$, say $E_\alpha^\mathcal{T}$ is \mathcal{T} -special iff $[0, \alpha]_\mathcal{T}$ does not drop and $E_\alpha^\mathcal{T} = F_1(M_\alpha^\mathcal{T})$. We may and do require that if $E_\alpha^\mathcal{T}$ is \mathcal{T} -special then $E_\alpha^\mathcal{U} \neq \emptyset$ (we have

$$F_0(M_\alpha^\mathcal{T}) \neq F_1(M_\alpha^\mathcal{U}),$$

because $F_0^P \neq F_1^P$). Likewise for \mathcal{U} .

We claim that the comparison fails. This is as in [3, §9], but we discuss a detail not discussed there. Suppose the comparison succeeds, with

$$Q = M_\infty^\mathcal{T} \trianglelefteq R = M_\infty^\mathcal{U}.$$

Then $b^\mathcal{T}$ drops (otherwise $F_0^Q \neq F_1^Q$ and the comparison can continue). In [3], it is argued that therefore Q is unsound, so $Q = R$, so $b^\mathcal{U}$ drops, and standard fine structure leads to contradiction. The appeal to fine structure assumes that enough extenders used in \mathcal{T}, \mathcal{U} are close to the models to which they apply. But [3, 6.1.5] was not proven for bicephali. We deal with this by proving:

(*1) If $[0, \delta + 1]_\mathcal{T}$ drops then $E_\delta^\mathcal{T}$ is close to $(M_{\delta+1}^*)^\mathcal{T}$; likewise for \mathcal{U} .⁶¹

Suppose that $E_\alpha^\mathcal{T}$ is \mathcal{T} -special. Let $\kappa = \text{cr}(E_\alpha^\mathcal{T})$. If $\gamma \in [0, \alpha]_\mathcal{T}$ is least such that $\gamma = \alpha$ or $\text{cr}(i_{\gamma\alpha}^\mathcal{T}) > \text{cr}(F_1(M_\gamma^\mathcal{T}))$, then $\kappa = \text{cr}(F_1(M_\gamma^\mathcal{T}))$. So for $\delta < \gamma$, we have $\nu_\delta^\mathcal{T} < \kappa$, and if $\gamma < \alpha$, then $\kappa < \text{cr}(i_{\gamma\alpha}^\mathcal{T}) < \nu_\gamma^\mathcal{T}$. Therefore $\text{pred}^\mathcal{T}(\alpha + 1) = \gamma$ and $\text{cr}(E_\alpha^\mathcal{T}) = \text{cr}(F_1(M_\gamma^\mathcal{T}))$.

Applying this to all \mathcal{T} -special extenders, we get:

(*2) If $E_\alpha^\mathcal{T}$ is \mathcal{T} -special, γ is as above and $\delta + 1 \in (\gamma, \alpha]_\mathcal{T}$, then $E_\delta^\mathcal{T}$ is not \mathcal{T} -special.

By these and similar considerations regarding those α such that $[0, \alpha]_\mathcal{T}$ does not drop and $E_\alpha^\mathcal{T} = F_0(M_\alpha^\mathcal{T})$, we have:

If $[0, \delta + 1]_\mathcal{T}$ drops then either $\text{lh}(E_\delta^\mathcal{T}) < \text{OR}(M_\delta^\mathcal{T})$ or $[0, \delta]_\mathcal{T}$ drops.

By this and the argument of [3, 6.1.5], (*1) follows.

We examine the generators of \mathcal{T} -special extenders. Given a short extender E weakly amenable to a premouse M , let $\varrho(E)$ denote the least $\varrho \geq (\kappa_E^+)^M$ such that $E \restriction \varrho \notin \text{Ult}_0(M, E)$ and $E \restriction \varrho$ is not type Z. (Note that $\varrho(E)$ is independent of the choice of M .)

⁶¹Establishing “closeness” in general would require consideration of $\Sigma_1^{(M, F_1, F_2)}$ in the language using both F_1, F_2 , which we prefer to avoid. We could alternatively make do with semi-closeness, using 2.24, but we also need the consequences of the proof of (*1) later.

If M is a premouse and $E \in \mathbb{E}_+^M$, then $\varrho(E) = \nu(E)$. Suppose $[0, \alpha]_{\mathcal{T}}$ is non-dropping. Then $\varrho(F_1(M_\alpha^{\mathcal{T}})) = \sup i_{0\alpha}^{\mathcal{T}} \delta(P)$. Moreover,

$$\varrho(F_1(M_\alpha^{\mathcal{T}})) < \nu(F_1(M_\alpha^{\mathcal{T}}))$$

iff there is $\beta <_{\mathcal{T}} \alpha$ such that

$$i_{0\alpha}^{\mathcal{T}} \delta(P) = i_{0\beta}^{\mathcal{T}} \delta(P) \subseteq \text{cr}(i_{\beta\alpha}^{\mathcal{T}}). \quad (9)$$

In this case, we have that

$$\nu(F_1(M_\alpha^{\mathcal{T}})) = \sup_{\gamma < \alpha} (\nu'(E_\gamma^{\mathcal{T}})), \quad (10)$$

where $\nu'(E) = \nu(E)$ if E is type 2 or type 3, and $\nu'(E) = \text{cr}(E) + 1$ if E is type 1. These statements can be proven inductively along the branch $[0, \alpha]_{\mathcal{T}}$, using 2.11 and (*2) for the successor case, and $\text{r}\Pi_1$ elementarity in the limit case.

It follows that if $[0, \alpha]_{\mathcal{T}}$ is non-dropping, then $\nu(F_1(M_\alpha^{\mathcal{T}})) \leq \delta(M_\alpha^{\mathcal{T}})$, and therefore $M_\alpha^{\mathcal{T}}$ is a bicephalus, and player II does not win through 5.3(a).

So the comparison fails, reaching length $\omega_1 + 1$. We take a hull and get $\pi : H \rightarrow V_\theta$ elementary with H countable. Let $\kappa = \text{cr}(\pi)$. Then $M_\kappa^{\mathcal{T}} \in H$ and $\pi \upharpoonright M_\kappa^{\mathcal{T}} = i_{\kappa\omega_1}^{\mathcal{T}}$, and likewise with \mathcal{U} . Also

$$M_\kappa^{\mathcal{T}}|(\kappa^+)^{M_\kappa^{\mathcal{T}}} = M_\kappa^{\mathcal{U}}|(\kappa^+)^{M_\kappa^{\mathcal{U}}},$$

and the (κ, ω_1) -extenders derived from $i_{\kappa, \omega_1}^{\mathcal{T}}$ and $i_{\kappa, \omega_1}^{\mathcal{U}}$ are identical. Denote this extender G . We have $M_{\omega_1}^{\mathcal{T}}|_{\omega_1} = M_{\omega_1}^{\mathcal{U}}|_{\omega_1}$, so $M_{\omega_1}^{\mathcal{U}}$ agrees with $M_{\omega_1}^{\mathcal{T}}$ about V_{ω_1} .

Let $\alpha + 1 = \min(\kappa, \omega_1]_{\mathcal{T}}$ and $\beta + 1 = \min(\kappa, \omega_1]_{\mathcal{U}}$. So $E_\alpha^{\mathcal{T}}$, G and $E_\beta^{\mathcal{U}}$ measure the same sets and are compatible through $\min(\nu_\alpha^{\mathcal{T}}, \nu_\beta^{\mathcal{U}})$. Now

$$\varrho(G) = \varrho(E_\alpha^{\mathcal{T}}) \leq \nu(E_\alpha^{\mathcal{T}}) \leq \nu_\alpha^{\mathcal{T}}.$$

Likewise for \mathcal{U}, β , so $\varrho(E_\alpha^{\mathcal{T}}) = \varrho(G) = \varrho(E_\beta^{\mathcal{U}})$.

Case 1. $\nu(E_\alpha^{\mathcal{T}}) = \varrho(E_\alpha^{\mathcal{T}}) = \varrho(G) = \varrho(E_\beta^{\mathcal{U}}) = \nu(E_\beta^{\mathcal{U}})$.

If $\alpha = \beta$, then also $\text{lh}(E_\alpha^{\mathcal{T}}) = \text{lh}(E_\beta^{\mathcal{U}})$, so the compatibility implies $E_\alpha^{\mathcal{T}} = E_\beta^{\mathcal{U}} = E_\alpha^{\mathcal{U}}$, contradiction. So assume $\alpha < \beta$. Then $\nu(E_\beta^{\mathcal{U}}) < \text{lh}(E_\alpha^{\mathcal{T}}) < \text{lh}(E_\beta^{\mathcal{U}})$, and $\text{lh}(E_\alpha^{\mathcal{T}})$ is a cardinal in $M_\beta^{\mathcal{U}}|_{\text{lh}(E_\beta^{\mathcal{U}})}$, so $E_\beta^{\mathcal{U}}$ is indexed too late for a premouse, so is \mathcal{U} -special. So $M_\beta^{\mathcal{U}}$ has form $(N, F_0, E_\beta^{\mathcal{U}})$, and by our rules on using \mathcal{U} -special extenders, $E_\beta^{\mathcal{T}} \neq \emptyset$. So $M_\beta^{\mathcal{T}}|_{\text{OR}^N}$ is active.

Now OR^N is a cardinal of $\text{Ult}_0(M_\beta^{\mathcal{U}}, E_\beta^{\mathcal{U}})$. But $E_\alpha^{\mathcal{T}} \upharpoonright \nu(E_\alpha^{\mathcal{T}}) = E_\beta^{\mathcal{U}} \upharpoonright \nu(E_\beta^{\mathcal{U}})$ and $M_\alpha^{\mathcal{T}}|_\kappa = M_\beta^{\mathcal{U}}|_\kappa$. So $\text{Ult}_0(M_\alpha^{\mathcal{T}}, E_\alpha^{\mathcal{T}})$ and $M_{\alpha+1}^{\mathcal{T}}$ agree with $\text{Ult}_0(M_\beta^{\mathcal{U}}, E_\beta^{\mathcal{U}})$ past $\text{OR}^N + 1$, and agree with $M_\beta^{\mathcal{U}}$ strictly below OR^N . But then for all $\gamma \in (\alpha, \beta)$, $E_\gamma^{\mathcal{T}} = \emptyset$. So $M_\beta^{\mathcal{T}} = M_{\alpha+1}^{\mathcal{T}}$, so $M_\beta^{\mathcal{T}}|_{\text{OR}^N}$ is passive, contradiction.

Case 2. $\varrho(E_\alpha^{\mathcal{T}}) < \nu(E_\alpha^{\mathcal{T}})$.

By the case hypothesis, $E_\alpha^{\mathcal{T}}$ is \mathcal{T} -special. We digress for a moment.

Definition 5.5. Let E be a short extender, weakly amenable over N . For $\varrho \leq \nu(E)$, we say $E \restriction \varrho$ is a *natural segment of E* iff either (i) $\varrho = 0$, or (ii) $\varrho \geq (\kappa_E^+)^N$ and $\nu(E \restriction \varrho) = \varrho$ and

$$\mathcal{P}(\varrho) \cap \text{Ult}_0(N, E) = \mathcal{P}(\varrho) \cap \text{Ult}_0(N, E \restriction \varrho).$$

Let the *segmentation* of E be

$$S = \{\varrho \mid E \restriction \varrho \text{ is a natural segment of } E\}.$$

For $\sigma, \varrho \in S$ with $\sigma \leq \varrho$ let

$$j_{\sigma\varrho} : \text{Ult}_0(N, E \restriction \sigma) \rightarrow \text{Ult}_0(N, E \restriction \varrho)$$

be the factor embedding, and if $\sigma < \varrho$, let $\kappa_\sigma = \text{cr}(j_{\sigma\varrho})$. If $\varrho = \min(S \setminus (\sigma + 1))$, let G_σ be the (κ_σ, ϱ) extender derived from $j_{\sigma\varrho}$. We call the extenders G_σ the *natural factors of E* . \dashv

If N is a premouse and $E = F^N$, then clearly $S = \{0, \nu(E)\}$. If $E = E_\alpha^\mathcal{T}$, then $\varrho(E) = \min(S \setminus \{0\})$. Let $\beta \in [0, \alpha']_\mathcal{T}$ be least as in line (9) (with α' replacing α). Then $S \setminus \{0\}$ consists of those ordinals ϱ of either the form $\varrho = \varrho(E)$, or the form $\varrho = \nu(E_\delta^\mathcal{T})$ for some δ such that $\delta + 1 \in (\beta, \alpha']_\mathcal{T}$. Moreover,

$$G_0 = E_{\alpha'}^\mathcal{T} \restriction \varrho(E_{\alpha'}^\mathcal{T})$$

and if $\sigma \in S \setminus \{0\}$ and there is $\varrho \in S$ such that $\sigma < \varrho$ then

$$G_\sigma = E_\delta^\mathcal{T} \restriction \nu(E_\delta^\mathcal{T}),$$

where δ is least such that $\delta + 1 \leq_\mathcal{T} \alpha'$ and $\text{cr}(E_\delta^\mathcal{T}) \geq \sigma$. Further, $E_\delta^\mathcal{T}$ is not \mathcal{T} -special. These remarks follow from the calculations in [18, 4.3 and following Remark] using fact (*2) and 2.27, much as in the proof of 2.35.

We now continue with Case 2. Let $S = \langle \varrho_\xi \rangle_{\xi < \omega_1}$ be the segmentation of G and let $\langle G_\sigma \rangle_{\sigma \in S}$ be the sequence of natural factors of G . This sequence is just the concatenation of the sequences of natural factors of the extenders $E_\delta^\mathcal{T}$ for $\delta + 1 \in (\kappa, \omega_1]_\mathcal{T}$. Likewise for the natural factors of the extenders $E_\delta^\mathcal{U}$ for $\delta + 1 \in (\kappa, \omega_1]_\mathcal{U}$.

We have $\varrho_0 = 0$ and $\varrho_1 = \varrho(E_\alpha^\mathcal{T}) < \nu(E_\alpha^\mathcal{T})$, and $E_\alpha^\mathcal{T}$ is \mathcal{T} -special. So

$$G_{\varrho_0} = E_\alpha^\mathcal{T} \restriction \varrho_1 = E_\beta^\mathcal{U} \restriction \varrho_1$$

and there is γ_0 such that $E_{\gamma_0}^\mathcal{T}$ is non- \mathcal{T} -special and $\varrho_2 = \nu(E_{\gamma_0}^\mathcal{T})$ and

$$G_{\varrho_1} = E_{\gamma_0}^\mathcal{T} \restriction \varrho_2.$$

Now if $\varrho_1 < \nu(E_\beta^\mathcal{U})$ then there is γ' such that $E_{\gamma'}^\mathcal{U}$ is non- \mathcal{U} -special and $\varrho_2 = \nu(E_{\gamma'}^\mathcal{U})$ and

$$G_{\varrho_1} = E_{\gamma'}^\mathcal{U} \restriction \varrho_2,$$

but then $(M_{\gamma_0}^{\mathcal{T}} \parallel \text{lh}(E_{\gamma_0}^{\mathcal{T}}), E_{\gamma_0}^{\mathcal{T}})$ is a premouse, and likewise for $E_{\gamma'}^{\mathcal{U}}$, so $E_{\gamma_0}^{\mathcal{T}} = E_{\gamma'}^{\mathcal{U}}$, contradiction. So $\varrho_1 = \nu(E_{\beta}^{\mathcal{U}})$.

Now let $\alpha_0 = \alpha$ and $\beta_0 = \beta$ and

$$\beta_1 + 1 = \min((\beta_0 + 1, \omega_1]_{\mathcal{U}}).$$

So $\varrho_2 = \varrho(E_{\beta_1}^{\mathcal{U}})$ and

$$G_{\varrho_1} = E_{\beta_1}^{\mathcal{U}} \restriction \varrho_2.$$

Now like before, $\varrho_2 < \nu(E_{\beta_1}^{\mathcal{U}})$ (here if $\text{lh}(E_{\gamma_0}^{\mathcal{T}}) < \text{lh}(E_{\beta_1}^{\mathcal{U}})$, use the argument from Case 1). So there is a non- \mathcal{U} -special $E_{\delta_1}^{\mathcal{U}}$ such that $\varrho_3 = \nu(E_{\delta_1}^{\mathcal{U}})$ and

$$G_{\varrho_2} = E_{\delta_1}^{\mathcal{U}} \restriction \varrho_3.$$

But then as before,

$$G_{\varrho_2} = E_{\alpha_2}^{\mathcal{T}} \restriction \varrho(E_{\alpha_2}^{\mathcal{T}})$$

for some \mathcal{T} -special $E_{\alpha_2}^{\mathcal{T}}$ with $\varrho(E_{\alpha_2}^{\mathcal{T}}) < \nu(E_{\alpha_2}^{\mathcal{T}})$. So by (*2), $E_{\alpha_0}^{\mathcal{T}}$ has only two natural factors, G_{ϱ_0} and G_{ϱ_1} .

This pattern continues through ω stages, producing a sequence of overlapping \mathcal{T} - and \mathcal{U} -special extenders $E_{\alpha_{2i}}^{\mathcal{T}}$ and $E_{\beta_{2i+1}}^{\mathcal{U}}$, for $i < \omega$, each having exactly two natural factors. The natural factors of $E_{\alpha_{2i}}^{\mathcal{T}}$ are $G_{\varrho_{2i}}$ and $G_{\varrho_{2i+1}}$; the natural factors of $E_{\beta_{2i+1}}^{\mathcal{U}}$ are $G_{\varrho_{2i+1}}$ and $G_{\varrho_{2i+2}}$.

For $i < \omega$ let $\kappa_i = \text{cr}(G_{\varrho_i})$ and $F_{2i} = E_{\alpha_{2i}}^{\mathcal{T}}$. So $\kappa_{2i} = \text{cr}(F_{2i})$. The next claim shows that $M_{\omega_1}^{\mathcal{T}}$ is illfounded, a contradiction.

Claim. *For each $i < \omega$ we have $i_{F_{2i}}(\kappa_{2i}) > \kappa_{2i+2}$.*

Proof. We will take $i = 0$ for simplicity, but the general case is similar. We have

$$E_{\gamma_0}^{\mathcal{T}} \restriction \nu(E_{\gamma_0}^{\mathcal{T}}) = G_{\varrho_1} = E_{\beta_1}^{\mathcal{U}} \restriction \varrho(E_{\beta_1}^{\mathcal{U}}),$$

and $\varrho_2 = \varrho(E_{\beta_1}^{\mathcal{U}}) < \nu(E_{\beta_1}^{\mathcal{U}})$. Let $\zeta <_{\mathcal{U}} \beta_1$ be least such that $\varrho_2 \leq \text{cr}(i_{\zeta\beta_1}^{\mathcal{U}})$. Then $\gamma_0 \geq \zeta$ because otherwise there is some extender used along $[0, \zeta]_{\mathcal{U}}$ which sends its critical point $> \varrho_2$, but then $\varrho(E_{\beta_1}^{\mathcal{U}}) > \varrho_2$, a contradiction. And $\varrho_2 = \nu(F_1(M_{\zeta}^{\mathcal{U}}))$ and

$$F_1(M_{\zeta}^{\mathcal{U}}) \restriction \varrho_2 = G_{\varrho_1}.$$

So $E_{\gamma_0}^{\mathcal{T}}$ and $F_1(M_{\zeta}^{\mathcal{U}})$ are essentially the same extender. Therefore

$$M_{\zeta}^{\mathcal{U}} \parallel \text{OR}(M_{\zeta}^{\mathcal{U}}) = \text{Ult}(M_{\gamma_0}^{\mathcal{T}}, E_{\gamma_0}^{\mathcal{T}}) \parallel \text{OR}(M_{\zeta}^{\mathcal{U}}),$$

which implies that $E_{\xi}^{\mathcal{U}} = \emptyset$ for all $\xi \in [\zeta, \gamma_0)$, so $M_{\gamma_0}^{\mathcal{U}} = M_{\zeta}^{\mathcal{U}}$, and implies that either

$$\text{lh}(F_1(M_{\zeta}^{\mathcal{U}})) = \text{lh}(E_{\gamma_0}^{\mathcal{T}}) \quad \& \quad F_1(M_{\zeta}^{\mathcal{U}}) = E_{\gamma_0}^{\mathcal{T}} \quad \& \quad E_{\gamma_0}^{\mathcal{U}} = F_0(M_{\zeta}^{\mathcal{U}})$$

or

$$\text{lh}(F_1(M_{\zeta}^{\mathcal{U}})) > \text{lh}(E_{\gamma_0}^{\mathcal{T}}) \quad \& \quad E_{\gamma_0}^{\mathcal{U}} = \emptyset = E_{\gamma_0+1}^{\mathcal{T}} \quad \& \quad E_{\gamma_0+1}^{\mathcal{U}} = F_0(M_{\zeta}^{\mathcal{U}}).$$

Note that, in any case, letting $F = F_1(M_\zeta^\mathcal{U})$, we have

$$i_{G_{e_1}}(\kappa_1) = i_F(\kappa_1) > \delta(M_\zeta^\mathcal{U}) \geq \kappa_2.$$

The fact that $i_F(\kappa_1) > \delta(M_\zeta^\mathcal{U})$ is because F is not of superstrong type, which is because F_1^P is not of superstrong type (this is easily preserved under iteration). The last inequality is because $\kappa_2 = \text{cr}(E_{\delta_1}^\mathcal{U})$ and $M_{\delta_1+1}^{*\mathcal{U}} = M_\zeta^\mathcal{U}$.

But $i_{F_0}(\kappa_0) > i_{\gamma_0+1}^*(\kappa_1) = i_F(\kappa_1)$. This completes the proof of the claim (in the case that $i = 0$). \square

This completes Case 2, and by symmetry, the proof. \square

Using the lemma, one can establish “Uniqueness of the next extender”:

Theorem 5.1, Proof Sketch. Suppose not. By [3, §9, §12], $B = (N_\lambda, E, F)$ is a mixed bicephalus. But B is $(0, \omega_1 + 1)$ -iterable, by the proof in [3, §12]. But the iterability of B contradicts Lemma 5.4.

(At the request of the referee, we provide a sketch of the iterability proof. It is almost the same as the case for a bicephalus with two active type 2 extenders. Given a tree \mathcal{T} on B and its lift \mathcal{U} on R , and given $\alpha < \text{lh}(\mathcal{T})$ such that $[0, \alpha]_\mathcal{T}$ does not drop, write

$$\begin{aligned} M_\alpha^\mathcal{T} &= M_\alpha = (N_\alpha, F_{0\alpha}, F_{1\alpha}), \\ B_\alpha &= i_{0\alpha}^\mathcal{U}(B) = (A_\alpha, E_{0\alpha}, E_{1\alpha}). \end{aligned}$$

We will have then a lifting map

$$\pi_\alpha : M_\alpha \rightarrow B_\alpha,$$

with

$$\pi_\alpha \circ i_{0\alpha}^\mathcal{T} = i_{0\alpha}^\mathcal{U} \upharpoonright B,$$

and in particular, $\pi_\alpha(\delta(M_\alpha)) = \delta(B_\alpha)$. Moreover,

$$\pi_\alpha : (N_\alpha, F_{i\alpha}) \rightarrow (A_\alpha, E_{i\alpha})$$

is a weak 0-embedding for both $i = 0$ and $i = 1$ (that is, Σ_0 -elementary, and Σ_1 -elementary on an \in -cofinal set). So π_α lifts extenders from \mathbb{E}^{N_α} to \mathbb{E}^{B_α} , and lifts $F_{i\alpha}$ to $E_{i\alpha}$. Moreover, in the case that $E_\alpha^\mathcal{T} = F_{1\alpha}$, because we use the exchange ordinal $\nu_\alpha^\mathcal{T} = \delta(M_\alpha^\mathcal{T})$ and since $i_{0\alpha}^\mathcal{U}$ is fully elementary, we have

$$\pi_\alpha(\nu_\alpha^\mathcal{T}) = \delta(B_\alpha) = \nu(E_{1\alpha}),$$

so in this case we have the usual sort of correspondence of exchange ordinals $\nu_\alpha^\mathcal{T}$ and $\nu_\alpha^\mathcal{U}$. \square

References

- [1] Kenneth Kunen. Some applications of iterated ultrapowers in set theory. *Annals of Mathematical Logic*, 1(2):179–227, 1970.
- [2] William Mitchell, Ernest Schimmerling, and John R. Steel. The covering lemma up to a Woodin cardinal. *Annals of Pure and Applied Logic*, 84(2):219–255, 1997.
- [3] William Mitchell and John R. Steel. *Fine structure and iteration trees*. Number 3 in Lectures Notes in Logic. Springer-Verlag, 1994.
- [4] William J. Mitchell. Sets constructed from sequences of ultrafilters. *Journal of Symbolic Logic*, 39(1):57–66, 1974.
- [5] William J. Mitchell. Sets constructed from sequences of measures: revisited. *Journal of Symbolic Logic*, 48(3):600–609, 1983.
- [6] Grigor Sargsyan. *Hod Mice and the Mouse Set Conjecture*. Memoirs of the American Mathematical Society, 2015.
- [7] Ernest Schimmerling. Combinatorial principles in the core model for one Woodin cardinal. *Annals of Pure and Applied Logic*, 74(2):153–201, 1995.
- [8] Ernest Schimmerling and John R. Steel. Fine structure for tame inner models. *Journal of Symbolic Logic*, 61(2):621–639, 1996.
- [9] Ernest Schimmerling and John R. Steel. The maximality of the core model. *Transactions of the American Mathematical Society*, 351(8):3119–3141, 1999.
- [10] Ralf Schindler. Core models in the presence of Woodin cardinals. *Journal of Symbolic Logic*, 71(4):1145–1154, 2006.
- [11] Ralf Schindler and John Steel. The self-iterability of $L[E]$. *Journal of Symbolic Logic*, 74(3):751–779, 2009.
- [12] Ralf Schindler, John R. Steel, and Martin Zeman. Deconstructing inner model theory. *Journal of Symbolic Logic*, 67(2):721–736, 2002.
- [13] Farmer Schlutzenberg. Reconstructing copying and condensation. To appear, available at <https://sites.google.com/site/schlutzenberg/home-1/research/papers-and-preprints>.
- [14] Farmer Schlutzenberg. *Measures in mice*. PhD thesis, University of California, Berkeley, 2007. Arxiv: 1301.4702.
- [15] Farmer Schlutzenberg. Homogeneously Suslin sets in tame mice. *Journal of Symbolic Logic*, 77(4):1122–1146, 2012.
- [16] J. R. Steel. Core models with more woodin cardinals. *The Journal of Symbolic Logic*, 67(3):1197–1226, 2002.

- [17] John R. Steel. Projectively well-ordered inner models. *Annals of Pure and Applied Logic*, 74(1):77–104, 1995.
- [18] John R. Steel. *The core model iterability problem*. Number 8 in Lecture Notes in Logic. Springer-Verlag, 1996.
- [19] John R. Steel. An outline of inner model theory. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of set theory*, volume 3, chapter 19. Springer, first edition, 2010.
- [20] Martin Zeman. *Inner Models and Large Cardinals*. Walter de Gruyter, 2002.