

Thermalization in a periodically driven fully-connected quantum Ising ferromagnet

ANGELO RUSSOMANNO^{1,2,3}, ROSARIO FAZIO^{5,6} and GIUSEPPE E. SANTORO^{2,3,4}

¹ *Department of Physics, Bar-Ilan University (RA), Ramat Gan 52900, Israel*

² *SISSA, Via Bonomea 265, I-34136 Trieste, Italy*

³ *CNR-IOM Democritos National Simulation Center, Via Bonomea 265, I-34136 Trieste, Italy*

⁴ *International Centre for Theoretical Physics (ICTP), P.O.Box 586, I-34014 Trieste, Italy*

⁵ *Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56127 Pisa, Italy*

⁶ *Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, 117543 Singapore*

PACS 75.10.Pq – Spin chain models

PACS 05.30.Rt – Quantum phase transitions

PACS 03.65.-w – Quantum mechanics

Abstract – By means of a Floquet analysis, we study the quantum dynamics of a fully connected Lipkin-Ising ferromagnet in a periodically driven transverse field showing that thermalization in the steady state is intimately connected to properties of the $N \rightarrow \infty$ classical Hamiltonian dynamics. When the dynamics is ergodic, the Floquet spectrum obeys a Wigner-Dyson statistics and the system satisfies the eigenstate thermalization hypothesis (ETH): Independently of the initial state, local observables relax to the $T = \infty$ thermal value, and Floquet states are delocalized in the Hilbert space. On the contrary, if the classical dynamics is regular no thermalization occurs. We further discuss the relationship between ergodicity and dynamical phase transitions, and the relevance of our results to other fully-connected periodically driven models (like the Bose-Hubbard), and possibilities of experimental realization in the case of two coupled BEC.

Introduction. Recent experimental advances in ultracold atomic systems [1–4] and femtosecond resolved spectroscopies [5] have made the study of out-of-equilibrium closed many-body quantum systems no longer a purely academic question. The key problem in this context are the properties of the final/steady state after the system has undergone a time-dependent perturbation [6, 7]. Depending on the nature of the perturbation, particular aspects acquire a prominent role. For a gentle (quasi-adiabatic) driving, the distance of the evolved final state from the instantaneous ground state carries precious information on the crossing of quantum critical points [8] and on the accuracy of quantum adiabatic computation [9] and quantum annealing protocols [10–12]. In the opposite case of a sudden quench, the focus is on the (possible) thermal properties of the steady state. Thermalization is expected in “classically ergodic” systems, where the Hamiltonian behaves as a random matrix [13], its eigenstates obey the *eigenstate thermalization hypothesis* (ETH) [6, 14–16], and relaxation to the micro-canonical ensemble, with vanishing fluctuations in the thermodynamic limit, follows.

Randomness of the eigenstates implies a strong connection between thermalization and delocalization in Hilbert space [6, 17–21].

Our goal is understanding the properties of the long-time dynamics of a many-body quantum system undergoing a periodic driving. The interest in periodically driven systems — a long-standing topic in quantum chaos of small quantum systems [22] — has risen again vigorously only quite recently, with a focus on many-body dynamics [23–31] and its properties of stability and ergodicity [32]. In a recent work [23] it was proposed that a periodically driven closed quantum system might display, in the thermodynamic limit, a tendency towards a fully coherent “periodic steady state” — a kind of diagonal ensemble [6, 16] for periodically driven systems —, where destructive interference effects average to zero for long times the transient fluctuations induced by off-diagonal Floquet matrix elements. This effect has been explicitly demonstrated on a periodically driven quantum Ising chain, as a direct consequence of the smooth continuous nature of the Floquet quasi-energy spectrum. A

quantum Ising chain, being integrable, does not show thermalization: The energy per site results from a GGE average [25, 33–35] and stays always well below the infinite temperature ($T = \infty$) value [23]. On the contrary, one might conjecture [23, 35] that when a classically ergodic system is periodically driven, the “steady state” would show thermalization to the $T = \infty$ ensemble. This is indeed shown in recent works [27, 36] on two nonintegrable periodically kicked spin chain models, consistently with the Floquet states obeying ETH at $T = \infty$ [28]. The same phenomena are observed in Ref. [26] where the case of interacting hard-core bosons is considered and the Floquet states are shown to obey ETH being random superpositions of unperturbed eigenstates. Analogous conclusions for a Bose-Hubbard chain are reported in Ref. [37]. These results call for a more detailed scrutiny of the relation between integrability and thermalization in periodically driven quantum many-body systems.

In the present letter we address this relation by studying a periodically driven fully-connected quantum Ising ferromagnet: a very clear prototype model whose long-time dynamics can be analyzed reliably up to the thermodynamic limit. The very rich phenomenology we are going to describe should occur also in a driven two-mode Bose-Hubbard model, whose Hamiltonian is equivalent to a fully-connected spin system [31] and is experimentally feasible by modulating the inter-well barrier height in a double-well BEC realization [38]. Systems of this kind can be mapped, for large N , onto large- S quantum spins with effective Planck’s constant $\hbar/N$. Hence, for $N = \infty$ their dynamics is that of a one-dimensional classical *non-linear* Hamiltonian system $\mathcal{H}(Q, P, t)$ [39]. At variance with the perfectly regular classical dynamics observed after a quantum quench [39, 40], we will show how rich is the periodic driving case: we can see classically regular motion, chaos and even full ergodicity by choosing appropriately the parameters of the driving. We will solve numerically the Schrödinger equation at finite “large” N . In both classically ergodic and regular cases, the intensive observables relax to a steady periodic regime: stroboscopic time fluctuations vanish in the large- N limit. However, the classically ergodic cases are very different from the regular ones: while the latter show a sensitivity to the initial state and never thermalize, the former effectively thermalize towards a $T = \infty$ ensemble. This is a consequence, as we will show, of the Floquet states obeying ETH at $T = \infty$ and being delocalized in the Hilbert space.

Model. Here we focus on a fully-connected quantum Ising ferromagnet with a periodically driven transverse field, the smooth-driving counterpart of the kicked-top of Ref. [41]. Consider N spin-1/2, $\hat{\mathbf{S}}_{i=1\dots N}$, and the total spin operator $\hat{\mathbf{S}} = \sum_i \hat{\mathbf{S}}_i$. The fully connected p -spin transverse field quantum Ising ferromagnet [42] is written as $\hat{H}_p(t) = -(NJ/2) \hat{m}_z^p - N\Gamma(t) \hat{m}_x$, where J is the longitudinal coupling, $\Gamma(t)$ is a (time-dependent) transverse field, and $\hat{m}_{x/z} = 2\hat{S}^{x/z}/N$ are rescaled magnetization

operators. $\hat{H}_p(t)$ commutes with $\hat{\mathbf{S}}^2$, and the equilibrium ground state is a state of *maximum spin*, $S = S_{max} = N/2$, belonging to the $(N+1)$ -dimensional multiplet of spin eigenstates $|S = N/2, M\rangle$. Since $[\hat{H}_p(t), \hat{\mathbf{S}}^2] = 0$, the Schrödinger dynamics starting from an initial state $|\psi_0\rangle$ with $S = N/2$ will always remain in that sector. If $m = 2M/N$ are the eigenvalues of $\hat{m}_z$, the multiplet of interest has $m = -1 + 2j/N$ with $j = 0, \dots, N$; we denote it as $|S = N/2, M\rangle_z \rightarrow |m\rangle$. The case $p = 2$ corresponds to the Ising-anisotropic version [43] of the so-called Lipkin model [44] (see also Refs. [45, 46]):

$$\hat{H}_{p=2}(t) = -\frac{2J}{N} \sum_{i,j} \hat{S}_i^z \hat{S}_j^z - 2\Gamma(t) \sum_i \hat{S}_i^x. \quad (1)$$

When Γ is constant, $\hat{H}_{p=2}$ has a quantum critical point (QCP) at $\Gamma_c/J = 1$ separating a large- Γ quantum paramagnet from a low- Γ ferromagnet; for $p > 2$ the transition is first order [47]. The non-equilibrium quantum dynamics of these models has so far been discussed in the cases of quantum annealing [10–12] across the QCP [42, 47–49], and of a sudden quench of $\Gamma(t)$ [39, 40], in the context of dynamical phase transitions. Here we will consider its non-equilibrium coherent dynamics under a periodic transverse field, more specifically $\Gamma(t) = \Gamma_0 + A \sin(\omega_0 t)$. The properties of a fully-connected spin chain undergoing a smooth periodic driving have been discussed with the rotating wave approximation in the context of non-equilibrium phase transitions in Ref. [30] and from the perspective of many-body coherent destruction of tunneling in the limit of high driving frequency in Ref. [29]; here we take a different point of view and focus on the regularity/ergodicity properties of the quantum many body system.

To discuss its exact quantum dynamics, we have to expand the state $|\psi(t)\rangle$ on the $(N+1)$ -dimensional basis $|m\rangle$ as $|\psi(t)\rangle = \sum_m \psi_m(t) |m\rangle$, the Schrödinger equation reads:

$$i\hbar \frac{\partial}{\partial t} \psi_m = -\frac{N}{2} J m^p \psi_m - \frac{N}{2} \Gamma(t) \sum_{\alpha=\pm 1} h_m^\alpha \psi_{m+\alpha \frac{2}{N}}, \quad (2)$$

with $h_m^\pm = \sqrt{1 - m^2 + 2(1 \mp m)/N}$. The system becomes increasingly classical for $N \rightarrow \infty$: indeed the commutator $[\hat{m}_x, \hat{m}_y] = i(2/N) \hat{m}_z$ vanishes in that limit. A careful semi-classical analysis [39, 42] reveals that the expectation values of the magnetization are effectively described, for $N = \infty$, by a one-dimensional classical Hamiltonian of the form

$$\mathcal{H}_p(Q, P, t) = -\frac{J}{2} Q^p - \Gamma(t) \sqrt{1 - Q^2} \cos(2P), \quad (3)$$

with the identification $\langle \psi(t) | \hat{m}_z | \psi(t) \rangle \rightarrow Q(t)$, and $\langle \psi(t) | \hat{m}_x | \psi(t) \rangle \rightarrow \sqrt{1 - Q^2(t)} \cos(2P(t))$. After a sudden quench of Γ , energy conservation gives an integrable $\mathcal{H}_p(Q, P)$; under a periodic $\Gamma(t)$, on the contrary, classical chaos in the (Q, P) phase space can emerge [50]. In

the rest of the paper we will focus on the case $p = 2$, whose theoretical and experimental importance relies also in the fact that Eq. (1) describes the exact dynamics [31] and Eq. (3) with $p = 2$ the mean field dynamics [51] of two coupled-trapped Bose-Einstein Condensates under a time-periodic modulation.

Floquet analysis. The natural framework to study the time evolution of a periodically driven quantum system is the Floquet theory [52, 53]. It states that there exists a basis of solutions of the Schrödinger equation which are periodic “up to a phase factor” $e^{-i\mu_\alpha t} |\phi_\alpha(t)\rangle$, where $|\phi_\alpha(t)\rangle = |\phi_\alpha(t + \tau)\rangle$, $\tau = 2\pi/\omega_0$ being the period. The Floquet quasi-energies μ_α and modes $|\phi_\alpha(t)\rangle$ are obtained by diagonalizing the evolution operator over one period $\hat{U}(\tau) |\phi_\alpha(0)\rangle = e^{-i\mu_\alpha \tau} |\phi_\alpha(0)\rangle$, with $\mu_\alpha \in [-\omega_0/2, \omega_0/2]$. If we consider the stroboscopic dynamics at times $t_n = n\tau$, since $\hat{U}(n\tau) = \hat{U}^n(\tau)$, the Schrödinger evolution is completely determined by $|\phi_\alpha(0)\rangle$ and μ_α : the state can be written as $|\psi(n\tau)\rangle = \sum_\alpha e^{-in\mu_\alpha \tau} R_\alpha |\phi_\alpha(0)\rangle$, where $R_\alpha = \langle \phi_\alpha(0) | \psi_0 \rangle$. We focus on the $p = 2$ case, considering intensive observables like the energy-per-site

$$e_{\psi_0}(n\tau) = \frac{1}{N} \langle \psi(n\tau) | \hat{H}_{p=2}(0) | \psi(n\tau) \rangle. \quad (4)$$

Provided the Floquet spectrum is non-degenerate (which we have verified numerically), we can easily evaluate the finite- N stroboscopic infinite-time average of e_{ψ_0} and the corresponding squared fluctuations $\delta e_{\psi_0}^2 = \bar{e}_{\psi_0}^2 - \bar{e}_{\psi_0}^2$:

$$\bar{e}_{\psi_0} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} e_{\psi_0}(k\tau) = \sum_\alpha |R_\alpha|^2 e_{\alpha\alpha} \quad (5)$$

$$\delta e_{\psi_0}^2 = \sum_{\alpha \neq \beta} |R_\alpha|^2 |R_\beta|^2 |e_{\alpha\beta}|^2, \quad (6)$$

where $e_{\alpha\beta} = \langle \phi_\alpha(0) | \hat{H}_{p=2}(0) | \phi_\beta(0) \rangle / N$. We can think of $\bar{e}_{\psi_0}$ as a “Floquet diagonal ensemble average” [6, 16, 35].

Results. The phenomenology of the driven model is quite rich, depending on Γ_0 , A and ω_0 . It will be helpful to use the equilibrium phase diagram as a guide (although the dynamics has no strict relation to the different equilibrium phases). When $\Gamma_0 < \Gamma_c = J$, a classical hyperbolic point at $(Q, P) = (0, 0)$ [39] makes the system prone to chaos even with a small A [50]: the stroboscopic Poincaré sections in Fig. 1(a) show an instance of ergodic phase space with fully developed chaos. At the quantum level [13], the corresponding distribution of Floquet quasi-energy spacings $P(S)$ — with $S_\alpha = \rho(\mu_\alpha)(\mu_{\alpha+1} - \mu_\alpha)$, where $\rho(\omega) = \langle \sum_\alpha \delta(\omega - \mu_\alpha) \rangle_\Delta$ is the density of quasi-energies smoothed over a “mesoscopic” scale Δ [13, 54] — is well described by an orthogonal Wigner-Dyson distribution [13] $P_{\text{WD}}(S) = \frac{\pi}{2} S \exp(-\frac{\pi}{4} S^2)$. When $\Gamma_0 > \Gamma_c$, inside the equilibrium paramagnetic phase, on the contrary, the classical motion tends to be more regular, and a larger A is needed for a substantial chaotic component in phase space: for $\Gamma_0 = 3J$, an $A/J = 0.5$ still shows very regular classical motion, Fig. 1(b), and a Poisson level statistics




Fig. 1: (Top) Schematic equilibrium phase diagram with the two representative driving cases illustrated. (a) and (b) Poincaré sections of the classical dynamics at stroboscopic times $t_n = n\tau$ in the (Q, P) phase space. The parameters are $\Gamma_0/J = 0.25$, $A/J = 0.45$ for (a), and $\Gamma_0/J = 3$, $A/J = 0.5$ for (b), with $\omega_0/J = 2$ and $\Gamma(t) = \Gamma_0 + A \sin(\omega_0 t)$. Chaos is fully developed in (a); the motion is regular in (b).

$P_P(S) = e^{-S}$ [55]. In the last case, the Poisson statistics gives rise, for very large N , to a Floquet spectrum with a non-extensive number of quasi-degeneracies; nevertheless relaxation to the Floquet-diagonal ensemble and Eqs. (5) are still valid, as shown in Refs. [7, 56] for the analogous case of a quantum quench with degeneracies in the energy spectrum. Results for the regular and ergodic cases are very similar to those found in the kicked-top problem [41]. Here we will focus on these two paradigmatic cases; see Ref. [35] for details about the intermediate situations.

We now turn to the evolution of the observables. If $e_{\psi_0}(n\tau)$ relaxes to a periodic steady regime, its asymptotic stroboscopic value must equal Eq. (5), and time fluctuations, Eq. (6), have to vanish. In Fig. 2(a) we show $e_{\psi_0}(n\tau)$ in the ergodic case: it clearly relaxes to $\bar{e}_{\psi_0}$. Two important points are in order:

- i) δe_{ψ_0} vanishes as $\sim N^{-1/2}$ (Fig. 2(b)), showing indeed relaxation in the thermodynamic limit;
- ii) $\bar{e}_{\psi_0}$ is independent of the initial state $|\psi_0\rangle$, up to differences of order $N^{-1/2}$, and equal to the $T = \infty$ thermal average $e_{T=\infty} = \frac{1}{N(N+1)} \text{Tr}_{S_{\text{max}}} [\hat{H}(0)]$ (the trace is restricted to the $S_{\text{max}} = N/2$ subspace).

This is true for every $|\psi_0\rangle$: as Fig. 2(c) shows, the Floquet diagonal terms $e_{\alpha\alpha}$ entering in Eq. (5)) are all equal




Fig. 2: (a) Stroboscopic evolution of energy $e_{\psi_0}(n\tau)$ in the ergodic case $\Gamma_0/J = 0.25$, $A/J = 0.45$, $\omega_0/J = 2$, for two different initial states ψ_0 , the symmetry-broken ground state (GS) and an eigenstate $|m\rangle$ of $2\hat{S}_z/N$, simulations are for $N = 800$. Here we can see that $e_{\psi_0}(n\tau)$ converges to $e_{T=\infty}$, in panel b) that the fluctuations δe_{ψ_0} (panel b) decay as $N^{-1/2}$. (c) $e_{\alpha\alpha} = H_{\alpha\alpha}/N$ (the diagonal Floquet matrix elements) vs the Floquet quasi-energy μ_α for $N = 800$: $e_{\alpha\alpha}$ is almost constantly equal to $e_{T=\infty}$ value, in agreement with $T = \infty$ -ETH.

to $e_{T=\infty}$, up to fluctuations of order $N^{-1/2}$, and one can see almost by inspection that property *ii*) follows whatever is the initial state, thanks to the normalization $\sum_\alpha |R_\alpha|^2 = 1$. Indeed, when the dynamics is ergodic, all the Floquet states are equivalent: they are superpositions of energy eigenstates with random phases and each one is equivalent to the $T = \infty$ thermal ensemble: they behave as eigenstates of a random matrix [13–15, 54, 57, 58] and obey the Eigenstate Thermalization Hypothesis at $T = \infty$. Concerning fluctuations, we have verified numerically (see Ref. [35] and the Supplementary Material) that the Floquet off-diagonal terms $|e_{\alpha\beta}|$ in Eq. (6) scale like $N^{-1/2}$ and, consequently, so does δe_{ψ_0} , whatever is the initial state (property *i*). There is also an analytical argument leading to this scaling. It relies on the fact that the Floquet states obey $T = \infty$ -ETH and are indeed uniform superpositions with random phases of the eigenstates of the Hamiltonian

$$|\phi_\alpha\rangle = \frac{1}{\sqrt{N+1}} \sum_{n=1}^{N+1} e^{-i\theta_n^\alpha} |n\rangle, \quad (7)$$

where θ_n^α are independent random variables uniformly distributed in $[0, 2\pi]$. Using this formula, the fact that $|n\rangle$ are eigenstates of $\hat{H}(0)$ and the central limit theorem, it is easy to show that the distributions of the real and the imaginary part of $e_{\alpha\beta}$ have a variance scaling like $\sim 1/N$ and indeed $\delta e_{\psi_0} \sim N^{-1/2}$. The detailed derivation is reported in the Supplementary Material. Thermalization to $T = \infty$ by means of ETH applies whenever the system is classically ergodic and is valid for the energy as well as for all the other intensive observables.

The physics in the classically regular case is very differ-




Fig. 3: Similar to Fig. 2, but for the classically regular case $\Gamma_0/J = 3$, $A/J = 0.5$, $\omega_0/J = 2$. Here the initial states are two different eigenstates $|m\rangle$ of $2\hat{S}_z/N$ and the ground state. (In panel a) we omit the results for the GS which show very small fluctuations around $e = -3$). Notice that $\bar{e}_{\psi_0}$ depends on $|\psi_0\rangle$, while the (very small) fluctuations decay with N with a scaling exponent dependent on $|\psi_0\rangle$ (panel b), and $e_{\alpha\alpha}$ fluctuates wildly with μ_α (panel c).

ent: Fig. 3(a) shows examples of $e_{\psi_0}(n\tau)$: We see relaxation to the Floquet diagonal ensemble (δe_{ψ_0} is practically invisible and scales to 0), but the asymptotic value $\bar{e}_{\psi_0}$ strongly depends on $|\psi_0\rangle$. The Floquet states behave here very differently: the diagonal terms $e_{\alpha\alpha}$ strongly depend on α (see Fig. 3(c)) and there is no ETH. We show numerically in the Supplementary material that also in this case the off-diagonal terms $|e_{\alpha\beta}|$ scale on average to zero like $N^{-1/2}$. We also find that these terms show larger fluctuations than in the ergodic case; this is the reason why δe_{ψ_0} scales, in the regular case, in a less smooth way with a scaling exponent depending on $|\psi_0\rangle$ (see the bottom-left panel of Fig. 3, where for the eigenstates $|m\rangle$ of $2\hat{S}_z/N$ we have $\delta e_m \sim N^{-1/2}$ and for the ground state $\delta e_{GS} \sim N^{-1}$).

Consistently with this picture, the order parameter stroboscopic time average $\bar{m}_{z\psi_0}$ vanishes if ergodicity is at play, even if $\Gamma(t)$ is always within the equilibrium ferromagnetic phase. We see this in Fig. 4, where we show $m_{z\psi_0}(n\tau)$ for two different $|\psi_0\rangle$. Although different in the details, this finding is in line with the “dynamical transition” found upon quenching from the ferromagnetic phase [39, 40]. Indeed, by taking $|\psi_0\rangle$ as the broken-symmetry ferromagnetic [GS] at Γ_0 and considering the dependence of $\bar{m}_{zGS}$ on the driving field amplitude A , we observe a transition at a critical value A_c (see the inset) independent of the number of particles N . As done in Ref. [39] for the case of quantum quench, we can give a classical phase space interpretation of this fact. In the classical $N \rightarrow \infty$ limit the ground state is a point in the phase space; if Γ_0 is in the broken symmetry phase this point has coordinates $P = 0$ and $Q = \pm\sqrt{1 - \Gamma_0^2}$. For $A < A_c$ this point falls in regular region of the phase space (bot-




Fig. 4: (Upper panel) Stroboscopic evolution of $m_{z,\psi_0}(n\tau)$ starting from $|\psi_0\rangle = |m = -0.75\rangle$ and $|GS\rangle$ in the ergodic case, showing a quick convergence towards $\bar{m}_{z,\psi_0} = 0$. Inset: $\bar{m}_{z,GS}$ vs A for $\Gamma_0/J = 0.25$ and $\omega_0/J = 2$, showing a dynamical phase transition (at $A_c \sim 0.07$) similar to that found in Ref. [39]. (Lower panels) Classical phase space interpretation of the transition: for $A < A_c$ (left panel, $A = 0.03$) the phase space point representative of the ground state is in a regular broken-symmetry region of the phase space; for $A > A_c$ (right panel, $A = 0.45$) it is in a chaotic region symmetric around $Q = 0$.

tom left panel of Fig. 4) and its subsequent dynamics is trapped in a torus [50] which is not symmetric around $(m_z)_{N \rightarrow \infty} = Q = 0$. Instead, when $A > A_c$, the ground state phase space point falls in a chaotic region symmetric around $Q = 0$ (bottom right panel) which is ergodically explored by the subsequent dynamics, and so the time average of $(m_z)_{N \rightarrow \infty} = Q$ vanishes. As a matter of fact, by changing the parameters A and ω_0 of the driving, we would find a whole critical line separating regions of phase space where the symmetry is broken from regions where the symmetry is dynamically restored.

It is interesting to explore the connection between thermalization and delocalization of states in the Hilbert space [18, 20, 21, 27]. An indicator for delocalization is the *inverse participation ratio* (IPR) [59] of a state on a given basis. If we consider the basis $\{|E_n\rangle\}$ of the eigenstates of $\hat{H}(0)$, we can study the localization of a given Floquet state $|\phi_\alpha(0)\rangle$ by calculating $\mathcal{I}_E(\phi_\alpha) = \sum_n |\langle E_n | \phi_\alpha(0) \rangle|^4$. For the ergodic case, we find a clear delocalization: upon averaging over α , we have $\langle \mathcal{I}_E(\phi_\alpha) \rangle_\alpha \sim 1/N$, with fluctuations that scale to 0 for $N \rightarrow \infty$, consistently with the Floquet states obeying $T = \infty$ -ETH (see [35]). In the classically regular cases, on the contrary, $\langle \mathcal{I}_E(\phi_\alpha) \rangle_\alpha$ is fi-

nite and almost independent of N , again with fluctuations scaling to 0, marking localization of the Floquet states. We can gauge delocalization in the Hilbert space [60] also by means of the probability amplitude to remain in $|\psi_0\rangle$, $G_{\psi_0}(n\tau) = \langle \psi_0 | \psi(n\tau) \rangle = \langle \psi_0 | \hat{U}^n(\tau) | \psi_0 \rangle$ whose square $\mathcal{F}_{\psi_0}(t) = |G_{\psi_0}(t)|^2 = |\langle \psi_0 | \psi(t) \rangle|^2$ can be seen as a dynamical fidelity [61]. Interestingly, one can show that

$$\bar{\mathcal{F}}_{\psi_0} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} |G_{\psi_0}(k\tau)|^2 = \sum_{\alpha} |R_{\alpha}|^4 \equiv \mathcal{I}_F(\psi_0),$$

which is the IPR of the initial state $|\psi_0\rangle$ in the Floquet basis. In the ergodic cases, we find that $\bar{\mathcal{F}}_{\psi_0} \sim N^{-1}$: *any* $|\psi_0\rangle$ appears as *extended* in the Floquet basis, the possibility of Anderson — here “dynamical” — localization is excluded, in full agreement with results on the kicked-top [41]. Classically regular cases show a diversity of possibilities: some states are definitely localized in the Floquet basis, while other states show anomalous scaling of the IPR, $\mathcal{I}_F(\psi_0) \sim N^{-\lambda}$ with $0 < \lambda < 1$, but further work is necessary to precisely understand the physics behind this.

Conclusions and perspectives. We have shown how, in a periodically driven fully connected spin model, classical ergodicity translates, at the quantum level, into the system heating up to $T = \infty$, with Floquet states obeying $T = \infty$ -ETH and being delocalized in the Hilbert space. On the contrary, if the classical dynamics is regular no thermalization occurs. We expect that a similar behaviour can be seen in other fully-connected models like Bose-Hubbard or Dicke models [39]. We can propose an experimental set-up to verify our predictions: modulating the inter-well barrier height in a double-well BEC realization [38] a driven two-mode Bose Hubbard Hamiltonian equivalent to our model [31] can be realized. Due to ergodicity and the exponential separation of trajectories, quantum effects can be easily observed, becoming evident after a few periods even if N is large [35] and we are well inside the semi-classical regime. In this work we have considered the signatures of quantum many body ergodicity manifesting in the behaviour of *local* observables, which have a clear classical limit. The next step is to study the behaviour of genuinely quantum *non-local* objects like the entanglement entropy whose analysis in connection to quantum phase transitions in static fully-connected spin chains has been carried out in Refs. [62, 63].

We acknowledge useful discussions with G. Biroli, E.G. Dalla Torre, M. Fabrizio, A. Polkovnikov, A. Silva and V. Smelyanskiy. Research was supported by the Coleman-Soref foundation, by MIUR, through PRIN-20087NX9Y7 and RPIN-2010LLKJBX, by SNSF, through SINERGIA Project CRSII2 136287 1, by the EU-Japan Project LEM-SUPER, and by the EU FP7 under grant agreements n. 280555, n. 600645 (IP-SIQS), n. 618074 (STREP-TERMIQ).

REFERENCES

- [1] GREINER M., MANDEL O., HANSCH T. W. and BLOCH I., *Nature (London)*, **419** (2002) 51.
- [2] KINOSHITA T., WENGER T. and WEISS D. S., *Nature (London)*, **440** (2006) 900.
- [3] JAKSCH D. and ZOLLER P., *Annals of Physics*, **315** (2005) 52.
- [4] TROTZKY S., CHEN Y.-A., FLESCH A., CULLOCH I. P. M., SCHOLLWÖCK U., EISERT J. and BLOCH I., *Nature Physics*, **8** (2012) 325.
- [5] DANTUS M. and GROSS P., *Ultrafast Spectroscopy in Encyclopaedia of Applied Physics* (Wiley) 2004.
- [6] POLKOVNIKOV A., SENGUPTA K., SILVA A. and VENGALATTORE M., *Rev. Mod. Phys.*, **83** (2011) 863.
- [7] REIMANN P., *Phys. Rev. Lett.*, **101** (2008) 190403.
- [8] ZUREK W. H., DORNER U. and ZOLLER P., *Phys. Rev. Lett.*, **95** (2005) 105701.
- [9] FARHI E., GOLDSTONE J., GUTMANN S., LAPAN J., LUNDGREN A. and PREDA D., *Science*, **292** (2001) 472.
- [10] KADOWAKI T. and NISHIMORI H., *Phys. Rev. E*, **58** (1998) 5355.
- [11] SANTORO G. E., MARTOŇÁK R., TOSATTI E. and CAR R., *Science*, **295** (2002) 2427.
- [12] SANTORO G. E. and TOSATTI E., *J. Phys. A: Math. Gen.*, **39** (2006) R393.
- [13] HAAKE F., *Quantum Signatures of Chaos* (2nd ed) (Springer) 2001.
- [14] DEUTSCH J. M., *Phys. Rev. A*, **43** (1991) 2046.
- [15] SREDNICKI M., *Phys. Rev. E*, **50** (1994) 888.
- [16] RIGOL M., DUNJKO V. and OLSHANII M., *Nature*, **452** (2008) 854.
- [17] BASKO D. M., ALEINER I. L. and ALTSHULER B., *Annals of Physics*, **321** (2006) 1126.
- [18] SANTOS L. F. and RIGOL M., *Phys. Rev. E*, (2010) 031130.
- [19] PAL A. and HUSE D. A., *Phys. Rev. B*, **82** (2010) 174411.
- [20] CANOVI E., ROSSINI D., FAZIO R., SANTORO G. E. and SILVA A., *Phys. Rev. B*, **83** (2011) 094431.
- [21] NEUENHAHN C. and MARQUARDT F., *Phys. Rev. E*, **85** (2012) 060101(R).
- [22] CASATI G. and MOLINARI L., *Prog. Theor. Phys. Supp.*, **98** (1989) 287.
- [23] RUSSOMANNO A., SILVA A. and SANTORO G. E., *Phys. Rev. Lett.*, **109** (2012) 257201.
- [24] BASTIDAS V., EMARY C., SCHALLER G. and BRANDES T., *Phys. Rev. A*, **86** (2012) 063627.
- [25] LAZARIDES A., DAS A. and MOESSNER R., *Phys. Rev. Lett.*, **112** (2014) 150401.
- [26] LAZARIDES A., DAS A. and MOESSNER R., *Phys. Rev. E*, **90** (2014) 012110.
- [27] PONTE P., CHANDRAN A., PAPIĆ Z. and ABANIN D. A., *Annals of Physics*, **353** (2015) 196.
- [28] KIM H., IKEDA T. N. and HUSE D. A., *Phys. Rev. E*, **90** (2014) 052105.
- [29] GONG J., MORALES-MOLINA L. and HÄNGGI P., *Phys. Rev. Lett.*, **103** (2009) 133002.
- [30] ENGELHARDT G., BASTIDAS V., EMARY C. and BRANDES T., *Phys. Rev. E*, **87** (2013) 052110.
- [31] KHRIPKOV C., COHEN D. and VARDI A., *Phys. Rev. E*, **87** (2013) 012910.
- [32] CITRO R., DALLA TORRE E. G., D’ALESSIO L., POLKOVNIKOV A., BABADI M., OKA T. and DEMLER E., *arXiv*, (2015) cond-mat/1501.05660.
- [33] JAYNES E. T., *Phys. Rev.*, **108** (1957) 171.
- [34] RIGOL M., DUNJKO V., YUROVSKY V. and OLSHANII M., *Phys. Rev. Lett.*, **98** (2007) 050405.
- [35] RUSSOMANNO A., Ph.D. thesis SISSA, Trieste (2014).
- [36] D’ALESSIO L. and RIGOL M., *Phys. Rev. X*, **4** (2014) 041048.
- [37] LIU D. E., *arXiv*, (2014) cond-mat/1410.2962v1.
- [38] ALBIEZ M., GATI R., FOLLING J., HUNSMANN S., CRISTIANI M. and OBERTHALER M. K., *Phys. Rev. Lett.*, **95** (2005) 010402.
- [39] SCIOLLA B. and BIROLI G., *JSTAT*, (2011) P11003.
- [40] MAZZA G. and FABRIZIO M., *Phys. Rev. B*, **86** (2012) 184303.
- [41] HAAKE F., KUŚ M. and SCHARF R., *Z. Phys. B*, **65** (1986) 381.
- [42] BAPST V. and SEMERJIAN G., *JSTAT*, (2012) P06007.
- [43] BOTET R. and JULLIEN R., *Phys. Rev. B*, **28** (1983) 3955.
- [44] LIPKIN H., MESHKOV N. and GLICK A., *Nuclear Physics*, **62** (1965) 188.
- [45] LATORRE J. I., ORÚS R., RICO E. and VIDAL J., *Phys. Rev. A*, **71** (2005) 064101.
- [46] RIBEIRO P., VIDAL J. and MOSSERI R., *Phys. Rev. Lett.*, **99** (2007) 050402.
- [47] JÖRG T., KRZAKALA F., KURCHAN J., MAGGS A. C. and PUJOS J., *EPL*, **89** (2010) 40004.
- [48] DAS A., SENGUPTA K., SEN D. and CHAKRABARTI B. K., *Phys. Rev. B*, **74** (2006) 144423.
- [49] CANEVA T., FAZIO R. and SANTORO G. E., *Phys. Rev. B*, **78** (2008) 104426.
- [50] BERRY M. V., *Regular and irregular motion in Topics in Nonlinear Mechanics*, edited by JONNA S., Vol. 46 (Am.Inst.Ph.) 1978 pp. 16–120.
- [51] SMERZI A., FANTONI S., GIOVANAZZI S. and SHENOY S. R., *Phys. Rev. Lett.*, **79** (1997) 4950.
- [52] SHIRLEY J. H., *Phys. Rev.*, **138** (1965) B979.
- [53] GRIFONI M. and HÄNGGI P., *Phys. Rep.*, **304** (1998) 229.
- [54] BERRY M. V., *Semiclassical mechanics of regular and irregular motion in Les Houches, Session XXXVI, 1981 — Chaotic Behaviour of Deterministic Systems*, edited by G. IOOS, R. H. G. HELLEMAN R. S., (North-Holland Publishing Company) 1983 pp. 174–271.
- [55] BERRY M. V. and TABOR M., *Proc. Roy. Soc. A*, **349** (1976) 101.
- [56] REIMANN P. and KASTNER M., *New J. Phys.*, **14** (2012) 043020.
- [57] BOHIGAS O., GIANNONI M. J. and SCHMIT C., *Phys. Rev. Lett.*, **52** (1984) 1.
- [58] PERES A., *Phys. Rev. A*, **30** (1984) 504.
- [59] EDWARDS J. T. and THOULESS D. J., *J. Phys. C*, (1972) 807.
- [60] ANDERSON P. W., *Phys. Rev.*, **109** (1958) 1492.
- [61] SHARMA S., RUSSOMANNO A., SANTORO G. E. and DUTTA A., *EPL*, **106** (2014) 67003.
- [62] FILIPPONE M., DUSUEL S. and VIDAL J., *Phys. Rev. A*, **83** (2011) 022327.
- [63] BARTHEL T., DUSUEL S. and VIDAL J., *Phys. Rev. Lett.*, **97** (2006) 220402.

Supplementary Material: Scaling of the fluctuations. — We show in the upper panel Fig. 5 the scaling of the average off-diagonal terms giving rise to the fluctuations (Eq. (6) of the main text)

$$\overline{|e_{\alpha\beta}|} = \frac{1}{N(N+1)} \sum_{\alpha \neq \beta} |e_{\alpha\beta}|,$$

both in the regular and the ergodic case. On the lower panel we show the scaling of the fluctuations of the off-diagonal terms

$$\delta(|e_{\alpha\beta}|) = \sqrt{|e_{\alpha\beta}|^2 - \overline{|e_{\alpha\beta}|}^2}.$$

We see that in the regular case these fluctuations are larger: for that reason, in this case, δe_{ψ_0} scales to zero in a less regular fashion with a scaling exponent dependent on $|\psi_0\rangle$, as we can see in the bottom-right panel of Fig. 3 of the main text.

When the dynamics is ergodic, we can show analytically that $\delta e_{\psi_0} \sim 1/\sqrt{N}$ whatever is the initial state. The argument runs as follows. Because in the ergodic case the Floquet states obey ETH at $T = \infty$ (Eq. (7) of the main text), they can be written as uniform superpositions with random phases of the eigenstates of the Hamiltonian

$$|\phi_\alpha\rangle = \frac{1}{\sqrt{N+1}} \sum_{n=1}^{N+1} e^{-i\theta_n^\alpha} |n\rangle \quad (8)$$

where θ_n^α are independent random variables distributed uniformly in $[0, 2\pi]$. Exploiting that $|n\rangle$ are eigenstates of $\hat{H}(0)$, the Floquet off-diagonal terms $e_{\alpha\beta}(0)$ entering in Eq. (6) of the manuscript can be written as

$$e_{\alpha\beta}(0) = \frac{1}{N+1} \sum_n e^{i(\theta_n^\alpha - \theta_n^\beta)} e_{nn}. \quad (9)$$

where $e_{nn} = \langle n | \hat{H}(0) | n \rangle / N$ do not scale with N . Because the e_{nn} are real, we can separate real and imaginary part of $e_{\alpha\beta}(0)$

$$\begin{aligned} \Re e_{\alpha\beta}(0) &= \frac{1}{N+1} \sum_n \cos(\theta_n^\alpha - \theta_n^\beta) e_{nn} \\ \Im e_{\alpha\beta}(0) &= \frac{1}{N+1} \sum_n \sin(\theta_n^\alpha - \theta_n^\beta) e_{nn}. \end{aligned} \quad (10)$$

Since θ_n^α and θ_n^β are distributed uniformly in $[0, 2\pi]$, then the distributions of $\cos(\theta_n^\alpha - \theta_n^\beta)$ and $\sin(\theta_n^\alpha - \theta_n^\beta)$ are symmetric around zero with a certain *finite* variance smaller than 1. By applying the central limit theorem, we see that both the real and the imaginary part of $e_{\alpha\beta}(0)$ are random variables with vanishing average and variance scaling like $\sim 1/N$. (The situation does not change by taking into account that e_{nn} can have different signs.) From this fact it easily follows that $\langle |e_{\alpha\beta}|^2 \rangle \sim 1/N$, where the average is performed on the random distribution of the θ_n^α .




Fig. 5: (Upper panel) Average of the $|e_{\alpha\beta}|$ vs N in the regular and in the ergodic case. (Lower panel) Fluctuations of the $|e_{\alpha\beta}|$ in the regular and the ergodic case.

Considering the overlaps R_α , whose square also appears in the fluctuations, Eq. (6) of the main text, we find

$$R_\alpha = \langle \psi_0 | \phi_\alpha(0) \rangle = \frac{1}{\sqrt{N+1}} \sum_n \psi_n^* e^{-i\theta_n^\alpha}. \quad (11)$$

where the initial state is expanded in the energy basis as $|\psi_0\rangle = \sum_n \psi_n |n\rangle$. We have to distinguish two cases: *i*) when the initial state is localized in the energy basis and *ii*) when it is extended.

i) In the first case, the number of states $|n\rangle$ such that $\psi_n \neq 0$ does not scale with N : we can schematize this situation by assuming $\psi_{\bar{n}} = 1$ for some $\bar{n}$ and $\psi_n = 0$ for $n \neq \bar{n}$. So $|R_\alpha|^2 = |R_\beta|^2 \approx \frac{1}{N+1}$ and the fluctuations (Eq. (6) of the manuscript) are

$$\begin{aligned} \delta e_{\psi_0}^2 &\simeq \frac{1}{(N+1)^2} \sum_{\alpha \neq \beta} |e_{\alpha\beta}|^2 \\ &\approx \frac{1}{(N+1)^2} \sum_{\alpha \neq \beta} \langle |e_{\alpha\beta}|^2 \rangle \sim \frac{N(N-1)}{N(N+1)^2} \sim \frac{1}{N}. \end{aligned} \quad (12)$$

In the second equality we have exploited that the Floquet states are random and self-averaging to replace

$|e_{\alpha\beta}|^2$ with its average over the probability distribution, $\langle |e_{\alpha\beta}|^2 \rangle$, whose scaling like $\sim 1/N$ we exploit in the third equality. In the fourth equality we have used that the number of terms in the sum is $N(N-1)$.

ii) Because we are only interested in scalings, we can schematize this situation by assuming that, for all n , $\psi_n \approx \frac{1}{\sqrt{N+1}}$. By applying the central limit theorem to the R_α in Eq. (11), we see that they are random variables with vanishing average and variance scaling like $1/N$. Hence, the fluctuations in Eq. (6) of the manuscript would be given by

$$\begin{aligned} \delta e_{\psi_0}^2 &= \sum_{\alpha \neq \beta} |R_\alpha|^2 |R_\beta|^2 |e_{\alpha\beta}|^2 \\ &\leq \sum_{\alpha \neq \beta} \langle |R_\alpha|^2 \rangle \langle |R_\beta|^2 \rangle \langle |e_{\alpha\beta}|^2 \rangle \sim \frac{N(N-1)}{N^3} \sim \frac{1}{N}, \end{aligned} \quad (13)$$

where the inequality comes from the fact that we are neglecting correlations between the R_α and $e_{\alpha\beta}$.