ON NONLOCAL NONLINEAR ELLIPTIC PROBLEM WITH THE FRACTIONAL LAPLACIAN

LI MA

ABSTRACT. In this paper, we study a nonlocal elliptic problem with the fractional Laplacian on \mathbb{R}^n . We show that the problem has infinite positive solutions in $C^{\tau}(\mathbb{R}^n) \cap H^{\alpha}_{loc}(\mathbb{R}^n)$. Moreover each of these solutions tends to some positive constant limit at infinity.

Mathematics Subject Classification (2000): 35J60, 53C21, 58J05 **Keywords**: fractional Laplacian,

1. INTRODUCTION

We prove the following result.

Theorem 1. Assume $0 < \alpha < n$ and p > 1. Let $\omega : R_+ \rightarrow R_+$ be the monotone non-increasing function such that

(1)
$$\int_{1}^{\infty} \frac{\omega(r)}{r} dr = A < \infty \pounds$$

Then there is a positive constant θ such that for smooth functions k(x) and K(x) with $|K(x)| \leq \theta \omega(|x|)(1 + |x|)^{-\tau}$ and $0 \leq k(x) \leq \theta \omega(|x|)(1 + |x|)^{-\tau}$ on \mathbb{R}^n for some $\tau \geq \alpha$, the problem

(2)
$$(-\Delta)^{\alpha/2}u + k(x)u = K(x)u^p, \quad in \ R^n$$

has infinite positive solutions in $C^{\tau}(\mathbb{R}^n) \cap H^{\alpha}_{loc}(\mathbb{R}^n)$. Moreover, each of these solutions tends to some positive constant limit at infinity.

We prove the above result by using the Perron method and similar argument as in Lin's work [3]. The Perron method is based on the maximum principle developed in [1]. Similar argument had been carried out in [4] for nonlinear subelliptic equations on R^n .

We denote by *C* the uniform constants, which may vary in different inequalities or formulae.

Denote by $B_1(0)$ the unit ball in \mathbb{R}^n .

We present the potential analysis as in [2] in section 2 and we prove the main result in section 3.

Date: May 26th, 2014.

^{*} The research is partially supported by the National Natural Science Foundation of China (No. 11271111) and SRFDP 20090002110019.

2. Preliminary

Let $\omega : \mathbb{R}^n \to \mathbb{R}$ be a radially symmetric monotone non-increasing function. Let $f : \mathbb{R}^n \to \mathbb{R}$ be a locally Holder continuous function with the decay growth as below:

(3)
$$|f(x)| \le C\omega(|x|)|x|^{-\tau}$$

where C > 0 and $\tau > 2\alpha$ are uniform constants.

Lemma 2. Define

$$w(x) = c_n \int_{\mathbb{R}^n} \frac{f(y)}{|x - y|^{n - \alpha}} dy,$$

where $c_n = [n(n-2)|B_1(0)|]^{-1}$. Then w(x) is well-defined and near ∞ we have

(4)
$$|w(x)| \leq \begin{cases} C|x|^{\alpha-n}\omega(|x|) & \text{if } \tau > n\\ C\omega(|x|)\log|x| & \text{if } \tau = n\\ C\omega(|x|)|x|^{\alpha-\tau} & \text{if } \alpha \le \tau \end{cases}$$

where $c_n = 1/n(n-2)|B_1(0)|$.

Proof. Clearly we have

$$|w(x)| \le C \int_{\mathbb{R}^n} \frac{\omega(|x|)}{|x-y|^{n-\alpha}(1+|y|^{\tau})} dy.$$

We now divide the the whole space R^n into three parts:

$$D_1 = \{ y \in \mathbb{R}^n; |y - x| \le |x|/2 \},\$$
$$D_2 = \{ y \in \mathbb{R}^n; |x|/2 \le |y - x| \le 2|x| \},\$$

and

$$D_3 = \{ y \in R^n; |y - x| \ge 2|x| \}.$$

Set, for j = 1, 2, 3,

$$I_{j} = \int_{D_{j}} \frac{\omega(|x|)}{|x - y|^{n - \alpha} (1 + |y|^{\tau}} dy.$$

Then

$$|w(x)| \leq C(I_1 + I_2 + I_3).$$

One can show as in [2] that w(x) has the upper bounds as in (4).

We now assume that $\omega(x)$ satisfies (1).

Lemma 3. Assume $f \ge 0$ on \mathbb{R}^n and $f(x) \ge C\omega(|x|)|x|^{-\tau}$ for some $\tau \ge \alpha$ and C > 0. The Riesz potential w(x) defined in Lemma 1 has the following lower bounds at ∞ :

(5)
$$w(x) \ge \begin{cases} C|x|^{\alpha-n}\omega(|x|) & \text{if } \tau > n\\ C\omega(|x|)\log|x| & \text{if } \tau = n\\ C\omega(|x|)|x|^{\alpha-\tau} & \text{if } \alpha \le \tau n \end{cases}$$

Proof. As in the proof above, we have

$$w(x) = \left(\int_{D_1} + \int_{D_2} + \int_{D_3}\right) \frac{c_n f(y)}{|x - y|^{n - \alpha}} dy \ge \int_{D_2} \frac{c_n f(y)}{|x - y|^{n - \alpha}} dy := J.$$

One can show as in [2] that J has the lower bounds as in (5).

3. Proof of Theorem 1

The argument presented below is similar to LIn's work [3]. We just do it briefly.

Take some constants $\theta_1 > 0$ and 0 < a < 1. We first define the function $U_a(x)$ on \mathbb{R}^n by solving the nonlocal equation

$$(-\Delta)^{\alpha/2}u = -\frac{C\omega(x)}{(1+|x|)^{\tau}}, \quad on \ R^n$$

for $C \in (0, \theta_1)$ with $0 < U_a(x) \le a$ and $U_a(x) \to a$ at infinity. We can verify that U_a is the lower solution to (2).

Then we define the function $U^{a}(x)$ on \mathbb{R}^{n} by solving the nonlocal equation

$$(-\Delta)^{\alpha/2}u = \frac{C\omega(x)}{(1+|x|)^{\tau}}, \quad on \ R^n$$

for the same C > 0 with $a \le U^a(x) < 1$ and $U^a(x) \to a$ at infinity. We can verify that U^a is the upper solution to (2).

Then we can use the Perron method [1][3] to get the desired solution u(x) to (2) such that $U_a(x) \le u(x) \le U^a(x)$ on \mathbb{R}^n .

This then completes the proof of Theorem 1.

References

- L.Caffarelli, L. Silvestre, regularity Theory for fully nonlinear integro-Differential Equations Comm. Pure Appl. Math. Vol. LXXII 0597-0638(2009.
- [2] Y.Li, W.M.Ni, On conformally scalar curvature equations in Rⁿ, Duke Math. J., 57(1988)895-924
- [3] F.H.Lin, On the elliptic equation $D_i[a_{ij}(x)D_jU k(x)U + K(x)U^p = 0,$ PAMS,95(2)(1985)219-226
- [4] Li MA, Dezhong Chen, Yang Yang, Some results for subelliptic equations, Acta. Math.Sinica. English series, 22(2006)1695-1704.

MA: DEPARTMENT OF MATHEMATICS, HENAN NORMAL UNIVERSITY, XINXIANG, 453007, CHINA *E-mail address*: lma@tsinghua.edu.cn