A note on modules over Quantum Laurent Polynomials

Ashish Gupta

December 17, 2021

Abstract

A twisted group algebra F * A of a finitely generated free abelian group A over a field F is in general noncommutative, but A may contain nontrivial subgroups C so that the subalgebra F * C is commutative. In this paper, we show that an F * A-module M which is finitely generated as an F * C-module for any commutative subalgebra F * C must be artinian. We also show that M must be torsion-free as F * C-module and the Gelfand–Kirillov dimension of M must equal the rank of C. We then apply these results to modules over finitely generated nilpotent groups of class 2.

1 Introduction

A quantum Laurent polynomial algebra $P(\mathfrak{q}, F)$ over a field F is defined as the associative F-algebra generated over F by the variables u_1, \dots, u_n and their inverses, satisfying the relations,

 $u_i u_j = q_{ij} u_j u_i,$

where $q_{ij} \in F$ are nonzero scalars and \mathfrak{q} is the $n \times n$ matrix (q_{ij}) . The q_{ij} satisfy

$$q_{ii} = 1 = q_{ij}q_{ji}, \qquad i, j = 1, \cdots, n.$$

These algebras have been called by various names. They have been called the *multiplicative analouges* of the Weyl algebra, the McConnell–Pettit algebras, the quantum Laurent polynomial algebras and the quantum tori. They arise as localizations of group algebras (see [Br]) and also play an important role in noncommutative geometry (see [M]).

The quantum Laurent polynomial algebras are precisely the twisted group algebras (see [P1, Chapter 1]) F * A of a finitely generated free abelian group A over a field F. In this paper we shall mainly use the notation F * A for a quantum Laurent polynomial algebra. Sometimes the following more explicit notation shall be needed:

$$F_{\mathfrak{q}}[u_1^{\pm 1},\cdots,u_n^{\pm 1}]$$

where $\mathbf{q} = (q_{ij})$.

Subsequently the situation with an arbitrary number of generators $u_1^{\pm 1}, \dots, u_n^{\pm 1}$ was studied in [MP]. In [MP], it was noted that the Krull and the global dimensions of F * A must coincide. An interesting and important criterion was given for these dimensions in [MP, Corollary 3.8] in terms of certain localizations (see [MP,]).

In [AG]. Modules over quantum Laurent polynomials have been considered in [A1], [?], [?], [BG] and [MP]. In this note we employ the methods of [AG] and [MP] and the work in [BG] in order to prove the following theorem for modules over F * A:

Theorem A. Let M be a nonzero finitely generated F * A-module, where F * A has center F. Let C < A be a subgroup which contains a subgroup C_0 of finite index such that $F * C_0$ is commutative. If M is finitely generated as an F * C-module then,

- (i) $\mathcal{GK}(M) = \operatorname{rk}(C)$,
- (ii) M is F * C-torsion-free,
- (iii) M is artinian,
- (iv) M is cyclic.

In this theorem $\mathcal{GK}(M)$ denotes the Gelfand–Kirillov dimension of M. It follows from this result that each finitely generated F * A-module with GK dimension one must be artinian (see Corollary 2.4).

Artinian modules arise in other situations also. For example it was shown in [Ar] that if the multiparameters q_{ij} , where $1 \leq i < j \leq n$, are independent in F^* , then each finitely generated periodic F * A-module is artinian and cyclic.

In Proposition 3.7 we give an application of this theorem to finitely generated modules over finitely generated and torsion-free nilpotent groups H of class 2. If k is field we denote by kH the group algebra of H over k. A kH-module M is said to be *reduced* if the annihilator of M in $k\zeta H$ is a prime ideal P and M is $k\zeta H/P$ -torsion-free. Every finitely generated kH-module is filtered by a finite series each section of which is reduced with some prime ideal of $k\zeta H$. The following theorem which is a special case of is deduced

Theorem B. Let H be a finitely generated torsion-free nilpotent group of class 2. A finitely generated kH-module Q which is reduced with annihilator P, and is finitely generated over kL for some abelian subgroup $L \leq H$ is torsion-free as kL/P.kL-module. Moreover if ζH is cyclic then Q is also artinian.

2 Background on F * A

The twisted group algebra F * A of a finitely generated free abelian group A over a field F has as an F-basis a copy $\overline{A} := \{\overline{a}, a \in A\}$ of A. The multiplication of basis elements $\overline{a}_1, \overline{a}_2$ is defined by

$$\bar{a}_1\bar{a}_2 = \lambda(a_1, a_2)\overline{a_1a_2},$$

where $\lambda : A \times A \to F^*$ is a function. The associativity condition requires that the map λ be a 2-cocycle (see [P1, Chapter 1]). The scalars $\kappa \in F$ commute with the basis elements. Thus $\kappa \bar{a} = \bar{a}\kappa$ for $\kappa \in F$ and $a \in A$. Thus any $\alpha \in F * A$ can be uniquely expressed in the form $\alpha = \sum_{a \in A} \lambda_a \bar{a}$, where $\lambda_a = 0$ for all but finitely many $a \in A$. The subset of elements $a \in A$ such that $\lambda_a \neq 0$ is known as the support of α in A. For a subgroup B of A, the subset of elements $\beta \in F * A$ such that the support of β lies in B, has the structure of a twisted group algebra of B over F. This is denoted as F * B. As already mentioned, F * A has as an F-basis a copy $\{\bar{a}, a \in A\}$ of A. By

a diagonal change in basis we may also take the subset $\{\kappa_a \bar{a}, a \in A, \kappa_a \in F^*\}$ as an *F*-basis for F * A. As explained in [P1, Chapter 1], with the help of a diagonal change in basis it can be shown that F * Z is commutative whenever Z is an infinite cyclic subgroup of A. By [Br, Theorem A], the maximal rank of a subgroup B of A such that F * B is commutative is equal to the Krull and the global dimension of F * A. Examples of F * A for which the commutativity of F * B implies that B is an infinite cyclic subgroup of A can be found in [MP].

It is known (e.g. [P2, Lemma 37.8]) that for each subgroup $B \leq A$, the monoid of nonzero elements of F * B is an Ore subset of F * A. Thus we may localize F * A at $X := F * B \setminus \{0\}$. This localization $(F * A)X^{-1}$ is itself a crossed product (see [P2, Chapter 1]) D * A/B of the group A/B over the division ring D of quotients of F * B. In this connection we shall adopt a useful notation that was introduced in [MP]: let $\{x_1, \dots, x_n\}$ be a basis for A such that $\{x_1, \dots, x_k\}$ is basis for B, where $k \leq n - 1$. Then $(F * A)X^{-1}$ is denoted as:

$$F(x_1,\cdots,x_k)[x_{k+1},\cdots,x_n],$$

where the localised generators are enclosed in roundparenthesess. As shown in [MP], the unit group \mathcal{U} of F * A is the set

$$\mathcal{U} = \{\kappa \bar{a}, \kappa \in F^*, a \in A\}$$

The derived subgroup \mathcal{U}' of \mathcal{U} is a subgroup of F^* . Thus \mathcal{U} is nilpotent of class at most 2. For subsets X and Y of A, we denote by \overline{X} the subset $\{\overline{x}, x \in X\}$ of \mathcal{U} , and by $[\overline{X}, \overline{Y}]$ the subgroup of \mathcal{U}' generated by the commutators $[\overline{x}, \overline{y}]$, where $x \in X$ and $y \in Y$. Since \mathcal{U}' is central in \mathcal{U} , hence by [?, Chapter 5],

$$\begin{split} [\bar{x}_1 \bar{x}_2, \bar{y}] &= [\bar{x}_1, \bar{y}] [\bar{x}_2, \bar{y}] \\ [\bar{x}, \bar{y}_1 \bar{y}_2] &= [\bar{x}, \bar{y}_1] [\bar{x}, \bar{y}_2] \end{split}$$

A dimension for studying modules over a crossed product D * A of a free finitely generated abelian group A over a division ring D, denoted as dim(M), was introduced in [BG] and was shown to coincide with the Gelfand–Kirillov dimension ([KL]).

Definition 2.1 (Definition 2.1 of [BG]). Let M be a D * A-module. The dimension dim(M) of M is defined to be the greatest integer r, $0 \le r \le rk(A)$, so that for some subgroup B in A with rank r, M is not D * B-torsion.

Proposition 2.2 (Lemma 2.2(1) of [BG]). Let

$$0 \to M_1 \to M \to M_2 \to 0,$$

be an exact sequence of D * A-modules. Then,

 $\dim(M) = \sup(\dim(M_1), \dim(M_2)).$

3 The proofs of Theorems A and B

The following lemma shall be key tool in the proof of Theorem A.

Lemma 3.1. Suppose that F * A has a finitely generated module M and A has a subgroup C with A/C torsion free, $\operatorname{rk}(C) = \mathcal{GK}(M)$, and F * C commutative. Suppose moreover that M is not F * C-torsion. Then C has a virtual complement E in A such that F * E is commutative. In fact given \mathbb{Z} -bases $\{x_1, \dots, x_r\}$ and $\{x_1, \dots, x_r, x_{r+1}, \dots, x_n\}$ for C and A respectively, there exist monomials $\mu_j, j = r+1, \dots, n$, in F * C, and an integer s > 0 such that the monomials $\mu_j \bar{x_j}^s$ commute in F * A.

Proof. Let $\bar{x}_i \bar{x}_j = q_{ij} \bar{x}_j \bar{x}_i$, where, $i, j = 1, \dots, n$ and $q_{ij} \in F^*$. We set $S = F * C \setminus \{0\}$ and denote the quotient field $(F * C)S^{-1}$ by F_S . Then $(F * A)S^{-1}$ is a crossed product

$$R = F(x_1, \cdots, x_r)[x_{r+1}, \cdots, x_n].$$

The corresponding module of fractions MS^{-1} is nonzero as M is not S-torsion by the hypothesis. Furthermore, by the hypothesis, $\mathcal{GK}(M) = \operatorname{rk}(C)$ and so in view of [BG, Lemma 2.3], MS^{-1} is finite dimensional as F_S -space. It is shown in [AG, Section 3] that if R has a module that is one dimensional over F_S then there exist monomials $\mu_i \in F * C$ such that the monomials $\mu_i \bar{x}_i$, where $r+1 \leq i \leq n$, commute mutually. Since the scalars in F are central in F * A, we may assume that $\mu_i \in \overline{C}$. If c_i is the element of C corresponding to μ_i , we may in this case take

$$E = \langle c_{r+1}x_{r+1}, \cdots, c_nx_n \rangle$$

the s-fold exterior power $M' := \wedge^s (MS_{F_S}^{-1})$, where, $s = \dim_{F_S} (MS^{-1})$ is a one dimensional module over the crossed product R' obtained from R by raising the cocycle of R to its s-th power. More explicitly R' is the crossed product

$$R' = R = F(x_1, \cdots, x_r)[x_{r+1}, \cdots, x_n]$$

but with $[\bar{x}_i, \bar{x}_j] = q'_{ij}$, where

$$q'_{ij} = q_{ij}, \quad 1 \le i \le r; \qquad q'_{ij} = q^s_{ij}, \quad r < i, j \le n.$$
 (1)

Then there are monomials $\mu_j \in F * C$ such that the monomials $\mu_j \bar{x}_j$, where $j = r+1, \dots, n$ commute with respect to the cocycle defined in equation 1. It easily follows from this that the monomials $\{\mu_j \bar{x}_j^s\}_{j=r+1}^n$ commute in F * A.

Remark 3.2. The above lemma remains valid if we drop the assumption that A/C is torsion-free. In this case A has a subgroup A' of finite index such that A'/C is torsion-free. Moreover in view of [BG, Lemma 2.7], M is finitely generated and has GK dimension equal to rk(C) as F * A'-module.

We are now ready to prove,

Theorem A. Let M be a nonzero finitely generated F * A-module where F * A has center F. Let C < A be a subgroup having a subgroup C_0 of finite index such that $F * C_0$ is commutative. If M is finitely generated as an F * C-module then,

- (i) $\mathcal{GK}(M) = \operatorname{rk}(C),$
- (ii) M is F * C-torsion-free,
- (iii) M is artinian,
- (iv) M is cyclic.

Proof. (i) We shall denote the module M regarded as F * C-module as M_C . By hypothesis M_C is finitely generated and so M_{C_0} is also finitely generated where M_{C_0} denotes M_C viewed as $F * C_0$ -module. By [BG, Lemma 2.7],

$$\mathcal{GK}(M) = \mathcal{GK}(M_C) = \mathcal{GK}(M_{C_0})$$

. If $\mathcal{GK}(M) < \operatorname{rk}(C)$ we may pick a subgroup $E_0 < C_0$ with $\operatorname{rk}(E_0) < \operatorname{rk}(C_0)$ such that M_{C_0} is not $F * E_0$ -torsion and $\mathcal{GK}(M) = \operatorname{rk}(E_0)$ (section ??). Moreover $F * E_0$ is commutative since $F * C_0$ is commutative by the hypothesis. By Lemmma 3.1 and Remark 3.2, E_0 has a (virtual) complement E_1 in A such that $F * E_1$ is commutative. Since $\operatorname{rk}(E_1) + \operatorname{rk}(C_0)$ exceeds $\operatorname{rk}(A)$, Hence $E_1 \cap C_0 > \langle 1 \rangle$. Moreover as E_1E_0 has finite index in A, hence E_1C_0 has finite index in A. But $\overline{E_1 \cap C_0}$ is central in $F * E_1C_0$ and hence F * A has center larger than F. This is contrary to the hypothesis in the theorem.

(ii) Suppose that the F * C-torsion submodule T of M is nonzero. We recall that T is an F * A-submodule of M. Applying part (i) of the theorem just established to T we obtain $\mathcal{GK}(T) = \operatorname{rk}(C)$. We shall denote the F * C-module structure on T by T_C . We note that T_C is finitely generated. Moreover

$$\mathcal{GK}(T_C) = \operatorname{rk}(T) = \operatorname{rk}(C)$$

by [BG, Lemma 2.7]. It then follows from [BG, Proposition 2.6], that there exists $t \in T$ so that $\operatorname{ann}_{F*C}(t) = 0$. But by the definition of T this is $\operatorname{ann}_{F*C}(t) \neq 0$. Hence T = 0 and M is F*C-torsion-free.

(iii) We first note that each nonzero subfactor of M has the same GK dimension as M. Let N and L be submodules of M such that Q := N/L is nonzero. As Q finitely generated over F * C it is F * C-torsion free by part (ii) of the theorem shown above. It follows from Definition 2.1 and Proposition 2.2 that

$$\mathcal{GK}(Q) = \operatorname{rk}(C) = \mathcal{GK}(M).$$

$$M = M_0 > M_1 > M_2 > \cdots,$$
(2)

Now let

be a strictly descending sequence of submodules of M. By the above $\mathcal{GK}(M_i/M_{i+1}) = \mathcal{GK}(M)$ for all $i \geq 0$. By [MP, Lemma 5.6] and [MP, 5.9] the sequence (2) must become constant after a finite number of steps. Hence M is artinian.

(iv) Since F * A has center F, it is simple by [MP, Proposition 1.3]. It follows from [Ba, Corollary 1.5] that M is cyclic.

Corollary 3.3. Each finitely generated module with Gelfand-Kirillov dimension less than or equal to one over an F * A with center exactly F is artinian and cyclic.

Proof. We recall that an F * A-module with GK dimension zero is finite dimensional (e.g. [KL]) over the base field F and so is artinian. By [W, Lemma 2.9], an F * A-module with GK dimension one is finitely generated over some F * C with C infinite cyclic. Since F * C must be commutative, the result follows from Theorem A.

Proposition 3.4. A finitely generated F * A-module M satisfying

$$\mathcal{GK}(N) + \dim(F * A) = \operatorname{rk}(A)$$

has finite length.

Proof. By [?, Theorem A], dim(F * A) is equal to the maximal rank of a subgroup B of A such that F * B is commutative. Hence in view of [?]B, the minimum possible GK dimension of a nonzero finitely generated F * A-module is $\operatorname{rk}(A) - \dim(F * A) = \mathcal{GK}(M)$. Thus in any (strictly) descending sequence $M = M_0 > M_1 > \cdots >$ of submodules of M, $\mathcal{GK}(M_i/M_{i+1}) = \mathcal{GK}(M)$ for each $i \ge 0$. It then p follows from [MP, Lemma 5.6] and [MP, 5.9] that this sequence must halt. Hence M has finite length.

We now investigate the situation in which a submodule of a finitely generated F * A-module is finitely generated over a commutative subalgebra F * C for C < A.

Definition 3.5. A nonzero F * A-module N is called critical when N/L has strictly smaller GK dimension than N for each nonzero proper submodule L of N.

3.1 Applications to nilpotent groups

We shall now consider some applications to finitely generated torsion-free nilpotent groups of class 2. Throughout this section H stands for such a group and k denotes a field. We denote the group algebra of H over k by kH and by ζH the center of H.

In [G], the following definition is given.

Definition 3.6. A finitely generated kH-module M is said to be reduced if there is a prime ideal P of $k\zeta H$ such that M is annihilated by P and is $k\zeta H/P$ -torsion-free.

It is remarked in [G] that every finitely generated kH-module is filtered by a finite series in which each section is reduced. We are now ready to prove

Proposition 3.7. Let H be a finitely generated torsion-free nilpotent group of class 2. Let M be a finitely generated kH-module reduced with annihilator P in $k\zeta H$. If M is finitely generated as a kL-module for some abelian subgroup L < H then,

- (i) M is torsion-free as kL/P.kL-module,
- (ii) for any descending sequence

$$M_1 \ge M_2 \ge \cdots \ge$$

of submodules of M, there is an integer s such that M_k/M_{k+1} is $k\zeta H/P$ -torsion for all $k \geq s$,

Proof. We note that there is no harm in assuming that $L > \zeta H$. As in [G], let K denote the field of fractions of $k\zeta H/P$ and set

$$\widehat{M} = M \otimes_{k \in H/P} K.$$

Then \widehat{M} is a finitely generated module for $kH/P.kH \otimes_{k\zeta H/P} K$ and the latter is a twisted group algebra K * A, where $A = H/\zeta H$. Moreover the center of K * A is exactly K. By the hypothesis in the theorem, M is finitely generated over kL and so over kL/P.kL. Thus \widehat{M} is finitely generated over $kL/P.kL \otimes_{k\zeta H/P} K$ which is the subalgebra K * C for $C = L/\zeta H$. Moreover since L is abelian, therefore, K * C is commutative. By Theorem A, \widehat{M} is K * C-torsion-free. In other words \widehat{M} is torsion-free as $kL/P.kL \otimes_{k\zeta H/P} K$ -module. As M is reduced it is by definition $k\zeta H/P$ -torsion free. Hence the natural kH/P.kH-linear map $\mu : M \to \widehat{M}$ via $y \mapsto y \otimes 1$ is a monomorphism. Therefore M must be kL/P.kL-torsion free. This shows part(i). As a consequence we obtain

Theorem B. Let H be a finitely generated torsion-free nilpotent group of class 2. For a finitely generated kH-module N which is reduced with annihilator P and is finitely generated over kL for some abelian subgroup $L \leq H$, N is torsion-free as kL/P.kL-module. Moreover if ζH is cyclic then N has finite length.

References

- [A1] Artamonov, V. A., 'Projective modules over quantum algebras of polynomials', Sb. Math. 82 (1995), 261–269.
- [A3] Artamonov, V. A. 'General quantum polynomials: irreducible modules and Morita equivalence', *Izv. Math.* 63 (1999), 3–36.
- [AG] Aljadeff E., Ginosar Y., On the global dimension of multiplicative Weyl algebra, Arch. Math. 62 (1994), 401 - 407.
- [Ar] Artamonov, V. A., General quantum polynomials: irreducible modules and Morita equivalence, Izv. Math. 63 (1999), 3–36.
- [Ba] Bavula, V. V., Module structure of the tensor product of simple algebras of Krull dimension 1, An. Stiint. Univ. Ovidius Constanta Ser. Mat. 4 (1996), 7–21.
- [Br] Brookes, C.J.B., Crossed Products and finitely presented groups, J. Group Theory 3 (2000), 433–444.
- [BG] Brookes C.J.B., Groves J.R.J., Modules over crossed products of a division ring with an abelian group I, J. Algebra 229 (2000), 25–54.
- [G] Groves J.R.J., Modules over infinite nilpotent groups, J. Aust. Math. Soc. 71 (2001) 211–221.
- [KL] Krause G.R., Lenagan T. H., Growth of algebras and Gelfand-Kirillov dimension, Res. Notes Math., Pitman, 1985.
- [M] Mannin Yu., Topics in noncommutative geometry, Princeton Univ. Press, 1991.
- [MP] McConnell, J. C., Pettit, J.J., 'Crossed products and multiplicative analouges of Weyl algebras', J. Lond. Math. Soc. 38 (1988), 47–55.
- [P1] Passman D.S, The algebraic structure of group rings Wiley, New York, 1977.
- [P2] Passman D.S, Infinite crossed products, Academic Press, London, 1989.
- [R] R]J. E. Roseblade, Prime Ideals in Group Rings of Polycyclic Groups, Proc. London Math. Soc., May 1978; s3-36: 385 - 447.
- [W] Wadsley, S.J., Homogeneity and Rigidity of the Brookes-Groves invariant for the noncommutative torus J. Algebra 297 (2006), 417-437.