## DIAMETER OF THE COMMUTING GRAPH OF $\mathbb{R}^{n \times n}$

YAROSLAV SHITOV

ABSTRACT. The vertices of commuting graph of  $\mathbb{R}^{n \times n}$  are non-scalar matrices; the edges are defined as pairs (u, v) satisfying uv = vu. One can see [1] that, for  $n \geq 3$ , the diameter of this graph is at least four; we give a short proof that it is exactly four. A longer proof of this result has been known [2, 3].

If the eigenvalues of a are simple, then  $\mathbb{R}^n$  is a direct sum of invariant subspaces each of which has dimension either 1 or 2; in this case, a commutes with a nontrivial idempotent. The distance between a pair of idempotents is at most 2 by Theorem 11 of [1], so that, for any matrices  $a_0$  and  $a_4$ , there are sequences  $a_0(t), \ldots, a_4(t)$  of non-scalar matrices such that  $a_0(t)$  and  $a_4(t)$  converge to  $a_0$  and  $a_4$ , respectively, and  $a_{j-1}(t)$  commutes with  $a_j(t)$ . Multiplying  $a_i(t)$  by a constant and adding a scalar matrix, we can get a singular matrix  $\alpha_i(t)$  with unit norm. Since the unit sphere is compact, every  $\alpha_i(t)$  is dense in some  $\alpha_i$ , which is non-scalar and real up to adding a scalar matrix. Passing to the limit, we see that  $\alpha_{j-1}$  and  $\alpha_j$  commute.

## References

- S. Akbari, A. Mohammadian, H. Radjavi, P. Raja, On the diameters of commuting graphs, Linear Algebra Appl. 418 (2006) 161–176.
- [2] J. M. Grau, A. Oller-Marcen, C. Tasis, On the diameter of the commuting graph of the full matrix ring over the real numbers, preprint (2014) arXiv:1412.0541.
- [3] C. Miguel, A note on a conjecture about commuting graphs, Linear Algebra Appl. 438 (2013) 4750–4756.

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, 20 MYASNITSKAYA ULITSA, MOSCOW 101000, RUSSIA

E-mail address: yaroslav-shitov@yandex.ru

<sup>2000</sup> Mathematics Subject Classification. 05C50. Key words and phrases. Commuting graph.