

# Data-Discriminants of Likelihood Equations <sup>\*</sup>

Jose Israel Rodriguez <sup>†</sup>

Xiaoxian Tang <sup>‡</sup>

January 27, 2022

## Abstract

Maximum likelihood estimation (MLE) is a fundamental computational problem in statistics. The problem is to maximize the likelihood function with respect to given data on a statistical model. An algebraic approach to this problem is to solve a very structured parameterized polynomial system called likelihood equations. For general choices of data, the number of complex solutions to the likelihood equations is finite and called the ML-degree of the model.

The only solutions to the likelihood equations that are statistically meaningful are the real/positive solutions. However, the number of real/positive solutions is not characterized by the ML-degree. We use discriminants to classify data according to the number of real/positive solutions of the likelihood equations. We call these discriminants data-discriminants (DD). We develop a probabilistic algorithm for computing DDs. Experimental results show that, for the benchmarks we have tried, the probabilistic algorithm is more efficient than the standard elimination algorithm. Based on the computational results, we discuss the real root classification problem for the 3 by 3 symmetric matrix model.

## 1 Introduction

We begin the introduction with an illustrative example. Suppose we have a weighted four-sided die such that the *probability*  $p_i$  of observing side  $i$  ( $i = 0, 1, 2, 3$ ) of the die satisfies the constraint  $p_0 + 2p_1 + 3p_2 - 4p_3 = 0$ . We toss the die 1000 times and record a 4-dimensional *data* vector  $(u_0, u_1, u_2, u_3)$ , where  $u_i$  is the number of times we observe the side  $i$ . We want to determine the probability distribution  $(p_0, p_1, p_2, p_3) \in \mathbb{R}_{>0}^4$  that best explains the data subject to the constraint. One approach is by *maximum likelihood estimation* (MLE):

---

<sup>\*</sup>This research paper is partly supported by 2014 NIMS Thematic Program on Applied Algebraic Geometry.

<sup>†</sup>ACMS Department, University of Notre Dame, Notre Dame, IN 46556; jo.ro@nd.edu. This material is based upon work supported by the National Science Foundation under Award No. DMS-1402545, as well as the hospitality of the Simons Institute.

<sup>‡</sup>CAMP, National Institute for Mathematical Sciences, Jeonmin-Dong, Yuseong-gu, Daejeon, Republic of Korea 305-811; tangxiaoxian@nims.re.kr

**Maximize the likelihood function  $p_0^{u_0} p_1^{u_1} p_2^{u_2} p_3^{u_3}$  subjected to**

$$p_0 + 2p_1 + 3p_2 - 4p_3 = 0, p_0 + p_1 + p_2 + p_3 = 1,$$

$$p_0 > 0, p_1 > 0, p_2 > 0, \text{ and } p_3 > 0.$$

For some statistical models, the MLE problem can be solved by well known *hill climbing* algorithms such as the EM-algorithm. However, the hill climbing method can fail if there is more than one local maximum. Fortunately, it is known that the MLE problem can be solved by solving the system of *likelihood equations* [15, 2]:

$$F_0 = p_0\lambda_1 + p_0\lambda_2 - u_0$$

$$F_1 = p_1\lambda_1 + 2p_1\lambda_2 - u_1$$

$$F_2 = p_2\lambda_1 + 3p_2\lambda_2 - u_2$$

$$F_3 = p_3\lambda_1 - 4p_3\lambda_2 - u_3$$

$$F_4 = p_0 + 2p_1 + 3p_2 - 4p_3$$

$$F_5 = p_0 + p_1 + p_2 + p_3 - 1$$

where  $\lambda_1$  and  $\lambda_2$  are newly introduced indeterminates (Lagrange multipliers) for formulating the likelihood equations. More specifically, for given  $(u_0, u_1, u_2, u_3)$ , if  $(p_0, p_1, p_2, p_3)$  is a critical point of the likelihood function, then there exist complex numbers  $\lambda_1$  and  $\lambda_2$  such that  $(p_0, p_1, p_2, p_3, \lambda_1, \lambda_2)$  is a solution of the polynomial system. For randomly chosen data  $u_i$ , the likelihood equations have 3 complex solutions. However, only solutions with positive coordinates  $p_i$  are statistically meaningful. A solution with all positive  $p_i$  coordinates is said to be a positive solution. So an important problem is *real root classification* (RRC):

**For which  $u_i$ , the polynomial system has 1, 2 and 3 real/positive solutions?**

According to the theory of computational (real) algebraic geometry [26, 20], the number of (real/positive) solutions only changes when the data  $u_i$  goes across some “special” values (see Theorem 2). The set of “special”  $u_i$  is a (*projective*) *variety* (see Lemma 4 in [20]) in (3 dimensional complex projective space) 4-dimensional complex space. The number of real/positive solutions is uniform over each open connected component determined by the variety. In other words, the “special”  $u_i$  plays the similar role as the *discriminant* for univariate polynomials. The first step of RRC is calculating the “special”  $u_i$ , leading to the discriminant problem:

**How to effectively compute the “special”  $u_i$ ?**

Geometrically, the “special”  $u_i$  is a projection of a variety. So in principle, it can be computed by *elimination* ( see Chapter 3, page 115–128 in [6]). For instance, by the command `eliminate` in Macaulay2 [10], we compute that the “special”  $u_i$  in the illustrative example form a hypersurface defined by a homogenous polynomial in  $(u_0, u_1, u_2, u_3)$  (see Example 1). However, for most MLE problems, due to the large size of likelihood equations, the elimination computation is too expensive. In this paper, we discuss the “discriminant” problem for the likelihood equations. The contributions of the paper are listed as follows.

- For likelihood equations, we show that the “special”  $u_i$  form a projective variety. We call the homogenous polynomial that generates the codimension 1 component of the projective variety the *data-discriminant*. This name distinguishes it from the *weight-discriminant* for the likelihood equations (which

replaces the condition  $p_0 + \cdots + p_n = 1$  with the condition  $h_0 p_0 + \cdots + h_n p_n = 1$  with parameters  $h_0, \dots, h_n$ ).

- For algebraic statistical models, we develop a probabilistic algorithm to compute data-discriminants. We implement the algorithm in Macaulay2. Experimental results show that the probabilistic algorithm is more efficient than the standard elimination algorithm.
- We discuss the real root classification for the  $3 \times 3$  symmetric matrix model, which inspire future work.

We remark that our work can be viewed as following the numerous efforts in applying computational algebraic geometry to tackle MLE and critical points problems [15, 2, 1, 16, 25, 12, 8, 13, 18, 14, 21].

The paper is organized as follows. The formal definition of the data-discriminant is introduced in Section 2. The standard elimination algorithm and the probabilistic algorithm are presented in Section 3. Experimental results comparing the two algorithms are shown in Section 4. The real root classification of the  $3 \times 3$  symmetric matrix model and conclusion are given in Section 5.

## 2 Definition

In this section, we discuss how to define “data-discriminant”. We assume the readers are familiar with elimination theory (see Chapter 3 in [6]).

**Notation 1.** Let  $\mathbb{P}$  denote the projective closure of the complex numbers  $\mathbb{C}$ . For homogeneous polynomials  $g_1, \dots, g_s$  in  $\mathbb{Q}[p_0, \dots, p_n]$ ,  $\mathcal{V}(g_1, \dots, g_s)$  denotes the projective variety in  $\mathbb{P}^n$  defined by  $g_1, \dots, g_s$ . Let  $\Delta_n$  denote the  $n$ -dimensional probability simplex  $\{(p_0, \dots, p_n) \in \mathbb{R}^{n+1} | p_0 > 0, \dots, p_n > 0, p_0 + \cdots + p_n = 1\}$ .

**Definition 1.** [15](Algebraic Statistical Model and Model Invariant) The set  $X$  is said to be an algebraic statistical model if  $X = \mathcal{V}(g_1, \dots, g_s) \cap \Delta_n$  where  $g_1, \dots, g_s$  define an irreducible generically reduced projective variety. Each  $g_k$  ( $1 \leq k \leq s$ ) is said to be a model invariant of  $X$ .

For a given algebraic statistical model, there are several different ways to formulate the likelihood equations [15]. In this section, we introduce the Lagrange likelihood equations and define the data-discriminant for this formulation. One can similarly define data-discriminants for other formulations of the likelihood equations.

**Notation 2.** For any  $f_1, \dots, f_m$  in the polynomial ring  $\mathbb{Q}[x_1, \dots, x_k]$ ,  $\mathcal{V}_a(f_1, \dots, f_m)$  denotes the affine variety in  $\mathbb{C}^k$  defined by  $f_1, \dots, f_m$  and  $\langle f_1, \dots, f_m \rangle$  denotes the ideal generated by  $f_1, \dots, f_m$ . For an ideal  $I$  in  $\mathbb{Q}[x_1, \dots, x_k]$ ,  $\mathcal{V}_a(I)$  denotes the affine variety defined by  $I$ .

**Definition 2.** [13](Lagrange Likelihood Equations and Correspondence) Given an algebraic statistical model  $X$ . The system of polynomial equations below is said to be the Lagrange likelihood equations of  $X$ :

$$\begin{aligned} F_0 &= p_0(\lambda_1 + \frac{\partial g_1}{\partial p_0} \lambda_2 + \cdots + \frac{\partial g_s}{\partial p_0} \lambda_{s+1}) - u_0 = 0 \\ &\dots \\ F_n &= p_n(\lambda_1 + \frac{\partial g_1}{\partial p_n} \lambda_2 + \cdots + \frac{\partial g_s}{\partial p_n} \lambda_{s+1}) - u_n = 0 \end{aligned}$$

$$\begin{aligned}
F_{n+1} &= g_1(p_0, \dots, p_n) = 0 \\
&\dots \\
F_{n+s} &= g_s(p_0, \dots, p_n) = 0 \\
F_{n+s+1} &= p_0 + \dots + p_n - 1 = 0
\end{aligned}$$

where  $g_1, \dots, g_s$  are the model invariants of  $X$  and  $u_0, \dots, u_n, p_0, \dots, p_n, \lambda_1, \dots, \lambda_{s+1}$  are indeterminates (also denoted by  $\mathbf{u}, \mathbf{p}, \Lambda$ ). More specifically,

- $p_0, \dots, p_n, \lambda_1, \dots, \lambda_{s+1}$  are unknowns,
- $u_0, \dots, u_n$  are parameters.

$\mathcal{V}_a(F_0, \dots, F_{n+s+1})$ , namely the set

$$\{(\mathbf{u}, \mathbf{p}, \Lambda) \in \mathbb{C}^{n+1} \times \mathbb{C}^{n+1} \times \mathbb{C}^{s+1} \mid F_0 = 0, \dots, F_{n+s+1} = 0\},$$

is said to be the Lagrange likelihood correspondence of  $X$  and denoted by  $\mathcal{L}_X$ .

**Notation 3.** Let  $\pi$  denote the canonical projection from the ambient space of the Lagrange likelihood correspondence to the  $\mathbb{C}^{n+1}$  associated to the  $\mathbf{u}$  indeterminants  $\pi: \mathbb{C}^{n+1} \times \mathbb{C}^{n+s+2} \rightarrow \mathbb{C}^{n+1}$ .

Given an algebraic statistical model  $X$  and a data vector  $\mathbf{u} \in \mathbb{R}_{>0}^n$ , the maximum likelihood estimation (MLE) problem is to **maximize the likelihood function**  $p_0^{u_0} \dots p_n^{u_n}$  **subject to**  $X$ . The MLE problem can be solved by computing  $\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X$ . More specifically, if  $\mathbf{p}$  is a regular point of  $\mathcal{V}(g_1, \dots, g_s)$ , then  $\mathbf{p}$  is a critical point of the likelihood function if and only if there exist  $\Lambda \in \mathbb{C}^{s+1}$  such that  $(\mathbf{u}, \mathbf{p}, \Lambda) \in \mathcal{L}_X$ . Theorem 1 states that for a general data vector  $\mathbf{u}$ ,  $\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X$  is a finite set and the cardinality of  $\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X$  is constant over a dense Zariski open set, which inspires the definition of ML-degree. For details, see [15].

**Theorem 1.** [15] For an algebraic statistical model  $X$ , there exist an affine variety  $V \subset \mathbb{C}^{n+1}$  and a non-negative integer  $N$  such that for any  $\mathbf{u} \in \mathbb{C}^{n+1} \setminus V$ ,

$$\#\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X = N.$$

**Definition 3.** [15](ML-Degree) For an algebraic statistical model  $X$ , the non-negative integer  $N$  stated in Theorem 1 is said to be the ML-degree of  $X$ .

**Notation 4.** For any  $S$  in  $\mathbb{C}^{n+1}$ ,  $\mathcal{I}(S)$  denotes the ideal

$$\{D \in \mathbb{Q}[\mathbf{u}] \mid D(a_0, \dots, a_n) = 0, \forall (a_0, \dots, a_n) \in S\}.$$

$\overline{S}$  denotes the affine closure of  $S$  in  $\mathbb{C}^{n+1}$ , namely  $\mathcal{V}_a(\mathcal{I}(S))$ .

**Definition 4.** For an algebraic statistical model  $X$ , suppose  $F_0, \dots, F_{n+s+1}$  are defined as in Definition 2. Let  $J$  denote

$$\det \begin{bmatrix} \frac{\partial F_0}{\partial p_0} & \dots & \frac{\partial F_0}{\partial p_n} & \frac{\partial F_0}{\partial \lambda_1} & \dots & \frac{\partial F_0}{\partial \lambda_{s+1}} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial F_{n+s+1}}{\partial p_0} & \dots & \frac{\partial F_{n+s+1}}{\partial p_n} & \frac{\partial F_{n+s+1}}{\partial \lambda_1} & \dots & \frac{\partial F_{n+s+1}}{\partial \lambda_{s+1}} \end{bmatrix}.$$

Then, we have the following:

- $\mathcal{L}_{X\infty}$  denotes the set of non-properness of  $\pi$ , i.e., the set of the  $u \in \overline{\pi(\mathcal{L}_X)}$  such that there does not exist a compact neighborhood  $U$  of  $u$  where  $\pi^{-1}(U) \cap \mathcal{L}_X$  is compact;
- $\mathcal{L}_{XJ}$  denotes  $\overline{\pi(\mathcal{L}_X \cap \mathcal{V}_a(J))}$ ;
- $\mathcal{L}_{Xp}$  denotes  $\overline{\pi(\mathcal{L}_X \cap \mathcal{V}_a(\prod_{k=0}^n p_k))}$ .

The geometric meaning of  $\mathcal{L}_{Xp}$  and  $\mathcal{L}_{XJ}$  are as follows. The first,  $\mathcal{L}_{Xp}$ , is the projection of the intersection of the Lagrange likelihood correspondence with the coordinate hyperplanes. The second,  $\mathcal{L}_{XJ}$ , is the projection of the intersection of the Lagrange likelihood correspondence with the hypersurface defined by  $J$ . Geometrically,  $\mathcal{L}_{XJ}$  is the closure of the union of the projection of the singular locus of  $\mathcal{L}_X$  and the set of critical values of the restriction of  $\pi$  to the regular locus of  $\mathcal{L}_X$  (see Definition 2 in [20]).

The Lagrange likelihood equations define an affine variety. As we continuously deform the parameters  $u_i$ , coordinates of a solution can tend to infinity. Geometrically,  $\mathcal{L}_{X\infty}$  is the set of the data  $\mathbf{u}$  such that the Lagrange likelihood equations have some solution  $(\mathbf{p}, \Lambda)$  at infinity; this is the closure of the set of “non-properness” as defined in the page 1, [19] and page 3, [23]. It is known that the set of non-properness of  $\pi$  is closed and can be computed by Gröbner bases (see Lemma 2 and Theorem 2 in [20]).

The ML-degree encaptures geometry of the likelihood equations over the complex numbers. However, statistically meaningful solutions occur over real numbers. Below, Theorem 2 states that  $\mathcal{L}_{X\infty}$ ,  $\mathcal{L}_{XJ}$  and  $\mathcal{L}_{Xp}$  define open connected components such that the number of real/positive solutions is uniform over each open connected component. Theorem 2 is a corollary of *Ehresmann’s theorem* for which there exists semi-algebraic statements since 1992 [5].

**Theorem 2.** For any algebraic statistical model  $X$ ,

- if  $\mathcal{C}_1, \dots, \mathcal{C}_t$  are the open connected components of

$$\mathbb{R}^{n+1} \setminus (\mathcal{L}_{X\infty} \cup \mathcal{L}_{XJ}),$$

then for each  $k$  ( $1 \leq k \leq t$ ), for any  $\mathbf{u} \in \mathcal{C}_k$ ,

$$\#\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X \cap \mathbb{R}^{n+s+2}$$

is a constant;

- if  $\mathcal{C}_1, \dots, \mathcal{C}_t$  are the open connected components of

$$\mathbb{R}^{n+1} \setminus (\mathcal{L}_{X\infty} \cup \mathcal{L}_{XJ} \cup \mathcal{L}_{Xp}),$$

then for each  $k$  ( $1 \leq k \leq t$ ), for any  $\mathbf{u} \in \mathcal{C}_k$ ,

$$\#\pi^{-1}(\mathbf{u}) \cap \mathcal{L}_X \cap (\mathbb{R}_{>0}^{n+1} \times \mathbb{R}^{s+1})$$

is a constant.

Before we give the definition of data-discriminant, we study the structures of  $\mathcal{L}_{X\infty}$ ,  $\mathcal{L}_{XJ}$ , and  $\mathcal{L}_{Xp}$  below.

- Proposition 1 shows that the structure of the likelihood equations forces  $\mathcal{L}_{Xp}$  to be contained in the union of coordinate hyperplanes defined by  $\prod_{k=0}^n u_k$ .
- Proposition 2 shows that the structure of the likelihood equations forces  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$  to be a projective variety.
- Similarly as the proof of Proposition 2, we can also show that the structure of the likelihood equations forces  $\mathcal{L}_{X\infty} \setminus \{\mathbf{0}\}$  to be a projective variety.

**Proposition 1.** *For any algebraic statistical model  $X$ ,*

$$\mathcal{L}_{Xp} \subset \mathcal{V}_a(\prod_{k=0}^n u_k).$$

**Proof.** By Definition 2, for any  $k$  ( $0 \leq k \leq n$ ),

$$u_k = p_k(\lambda_1 + \frac{\partial g_1}{\partial p_1} \lambda_2 + \cdots + \frac{\partial g_s}{\partial p_1} \lambda_{s+1}) - F_k.$$

Hence,

$$u_k \in \langle F_k, p_k \rangle \cap \mathbb{Q}[u_k] \subset \langle F_0, \dots, F_{n+s+1}, p_k \rangle \cap \mathbb{C}[\mathbf{u}]$$

So

$$\mathcal{V}_a(\langle F_0, \dots, F_{n+s+1}, p_k \rangle \cap \mathbb{C}[\mathbf{u}]) \subset \mathcal{V}_a(u_k)$$

By the Closure Theorem [6],

$$\mathcal{V}_a(\langle F_0, \dots, F_{n+s+1}, p_k \rangle \cap \mathbb{C}[\mathbf{u}]) = \overline{\pi(\mathcal{L}_X \cap \mathcal{V}_a(p_k))}$$

Therefore,

$$\begin{aligned} \mathcal{L}_{Xp} &= \overline{\pi(\mathcal{L}_X \cap \mathcal{V}_a(\prod_{k=0}^n p_k))} \\ &= \overline{\pi(\mathcal{L}_X \cap \bigcup_{k=0}^n \mathcal{V}_a(p_k))} \\ &= \bigcup_{k=0}^n \overline{\pi(\mathcal{L}_X \cap \mathcal{V}_a(p_k))} \\ &\subset \bigcup_{k=0}^n \mathcal{V}_a(u_k) \\ &= \mathcal{V}_a(\prod_{k=0}^n u_k). \square \end{aligned}$$

**Remark 1.** Generally,  $\mathcal{L}_{Xp} \neq \mathcal{V}_a(\prod_{k=0}^n u_k)$ . For example, suppose the algebraic statistical model is  $\mathcal{V}_a(p_0 - p_1) \cap \Delta_1$ . Then  $\mathcal{L}_{Xp} = \emptyset \neq \mathcal{V}_a(u_0 u_1)$ .

**Notation 5.**  $\mathcal{D}_{Xp}$  denotes the product  $\prod_{k=0}^n u_k$ .

**Proposition 2.** *For an algebraic statistical model  $X$ , we have  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$  is a projective variety in  $\mathbb{P}^n$ , where  $\mathbf{0}$  is the zero vector  $(0, \dots, 0)$  in  $\mathbb{C}^{n+1}$ .*

**Proof.** By the formulation of the Lagrange likelihood equations, we can prove that  $\mathcal{I}(\pi(\mathcal{L}_X \cap \mathcal{V}_a(J)))$  is a homogeneous ideal by the two basic facts below, which can be proved by Definition 2 and basic algebraic geometry arguments.

**C1.** For every  $\mathbf{u}$  in  $\pi(\mathcal{L}_X \cap \mathcal{V}_a(J))$ , each scalar multiple  $\alpha\mathbf{u}$  is also in  $\pi(\mathcal{L}_X \cap \mathcal{V}_a(J))$ .

**C2.** For any  $S \subset \mathbb{C}^{n+1}$ , if for any  $\mathbf{u} \in S$  and for any scalar  $\alpha \in \mathbb{C}$ ,  $\alpha\mathbf{u} \in S$ , then  $\mathcal{I}(S)$  is a homogeneous ideal in  $\mathbb{Q}[\mathbf{u}]$ .

That means the ideal  $\mathcal{I}(\pi(\mathcal{L}_X \cap \mathcal{V}_a(J)))$  is generated by finitely many homogeneous polynomials  $D_1, \dots, D_m$ . Therefore,  $\mathcal{L}_{XJ} = \mathcal{V}_a(\mathcal{I}(\pi(\mathcal{L}_X \cap \mathcal{V}_a(J)))) = \mathcal{V}_a(D_1, \dots, D_m)$ . So  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\} = \mathcal{V}(D_1, \dots, D_m) \subset \mathbb{P}^n$ .  $\square$

**Notation 6.** For an algebraic statistical model  $X$ , we define the notation  $\mathcal{D}_{XJ}$  according to the codimension of  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$  in  $\mathbb{P}^n$ .

- If the codimension is 1, then assume  $\mathcal{V}(D_1), \dots, \mathcal{V}(D_K)$  are the codimension 1 irreducible components in the minimal irreducible decomposition of  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$  in  $\mathbb{P}^n$  and  $\langle D_1 \rangle, \dots, \langle D_K \rangle$  are radical.  $\mathcal{D}_{XJ}$  denotes the homogeneous polynomial  $\prod_{j=1}^K D_j$ .

- If the codimension is greater than 1, then our convention is to take  $\mathcal{D}_{XJ} = 1$ .

Similarly, we use the notation  $\mathcal{D}_{X\infty}$  to denote the projective variety  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$ . Now we define the “data-discriminant” of Lagrange likelihood equations.

**Definition 5. (Data-Discriminant)** For a given algebraic statistics model  $X$ , the homogeneous polynomial  $\mathcal{D}_{X\infty} \cdot \mathcal{D}_{XJ} \cdot \mathcal{D}_{Xp}$  is said to be the data-discriminant (DD) of Lagrange likelihood equations of  $X$  and denoted by  $\mathcal{D}_X$ .

**Remark 2.** Note that DD can be viewed as a generalization of the “discriminant” for univariate polynomials. So it is interesting to compare DD with border polynomial (BP) [26] and discriminant variety (DV) [20]. DV and BP are defined for general parametric polynomial systems. DD is defined for the likelihood equations but can be generalized to any square and generic zero-dimensional system. Generally, for any square and generic zero-dimensional system,  $\mathcal{V}_a(\text{DD}) \subset \text{DV} \subset \mathcal{V}_a(\text{BP})$ . Note that due to the special structure of likelihood equations, DD is a homogenous polynomial despite being an affine system of equations. However, generally, DV is not a projective variety and BP is not homogenous.

**Example 1 (Linear Model).** The algebraic statistic model for the four sided die story in Section 1 is given by

$$X = \mathcal{V}(p_0 + 2p_1 + 3p_2 - 4p_3) \cap \Delta_3.$$

The Lagrange likelihood equations are the  $F_0 = 0, \dots, F_5 = 0$  shown in Section 1. The Lagrange likelihood correspondence is  $\mathcal{L}_X = \mathcal{V}_a(F_0, \dots, F_5) \subset \mathbb{C}^{10}$ . If we choose generic  $(u_0, u_1, u_2, u_3) \in \mathbb{C}^4$ ,  $\pi^{-1}(u_0, u_1, u_2, u_3) \cap \mathcal{L}_X = 3$ , namely the ML-degree is 3. The data-discriminant is the product of  $\mathcal{D}_{X\infty}$ ,  $\mathcal{D}_{Xp}$  and  $\mathcal{D}_{XJ}$ , where

$$\mathcal{D}_{X\infty} = u_0 + u_1 + u_2 + u_3, \mathcal{D}_{Xp} = u_0 u_1 u_2 u_3, \text{ and}$$

$$\begin{aligned} \mathcal{D}_{XJ} = & 441u_0^4 + 4998u_0^3u_1 + 20041u_0^2u_1^2 + 33320u_0u_1^3 + 19600u_1^4 - 756u_0^3u_2 + 20034u_0^2u_1u_2 + 83370u_0u_1^2u_2 + 79800u_1^3u_2 - 5346u_0^2u_2^2 + \\ & 55890u_0u_1u_2^2 + 119025u_1^2u_2^2 + 4860u_0u_2^3 + 76950u_1u_2^3 + 18225u_2^4 - 1596u_0^3u_3 - 11116u_0^2u_1u_3 - 17808u_0u_1^2u_3 + 4480u_1^3u_3 + 7452u_0^2u_2u_3 - 7752u_0u_1u_2u_3 + \end{aligned}$$

$$49680u_1^2u_2u_3 - 17172u_0u_2^2u_3 + 71460u_1u_2^2u_3 + 27540u_2^3u_3 + 2116u_0^2u_3^2 + 6624u_0u_1u_3^2 - 4224u_1^2u_3^2 - 9528u_0u_2u_3^2 + 15264u_1u_2u_3^2 + 14724u_2^2u_3^2 - 1216u_0u_3^3 - 512u_1u_3^3 + 3264u_2u_3^3 + 256u_3^4.$$

By applying the well known partial cylindrical algebraic decomposition (PCAD) [4] method to the data-discriminant above, we get that for any  $(u_0, u_1, u_2, u_3) \in \mathbb{R}_{>0}^4$ ,

- if  $\mathcal{D}_{XJ}(u_0, u_1, u_2, u_3) > 0$ , then the system of likelihood equations has 3 distinct real solutions and 1 of them is positive;
- if  $\mathcal{D}_{XJ}(u_0, u_1, u_2, u_3) < 0$ , then the system of likelihood equations has exactly 1 real solution and it is positive.

The answer above can be verified by the `RealRootClassification` [26, 3] command in Maple 17. In this example, the  $\mathcal{D}_{X\infty}$  does not effect the number of real/positive solutions since it is always positive when each  $u_i$  is positive. However, generally,  $\mathcal{D}_{X\infty}$  plays an important role in real root classification. Also remark that the real root classification is equivalent to the positive root classification for this example but it is not true generally (see Example 6).

### 3 Algorithm

In this section, we discuss how to compute  $\mathcal{D}_X$ . We assume that  $X$  is a given statistical model,  $F_0, \dots, F_{n+s+1}$  are defined as in Definition 2, and  $J$  is defined as in Definition 4. We rename  $F_0, \dots, F_{n+s+1}$  as  $F_0, \dots, F_m$ . Subsection 3.1 presents the standard elimination algorithm for reference and Subsection 3.2 presents our main algorithm (Algorithm 2).

#### 3.1 Standard Elimination Algorithm

Considering the data-discriminant as a projection drives a natural algorithm to compute it. This is the standard elimination algorithm in symbolic computation:

- we compute the  $\mathcal{L}_{XJ}$  by *elimination* and then get  $\mathcal{D}_{XJ}$  by the *radical equidimensional decomposition* (see Definition 3 in [20]). The algorithm is formally described in the Algorithm 1;

---

#### Algorithm 1: DX-J

---

**input** :  $F_0, \dots, F_m, J$   
**output**:  $\mathcal{D}_{XJ}$   
1  $\mathcal{G}_u \leftarrow$  the generator polynomial set of the elimination ideal  $\langle F_0, \dots, F_m, J \rangle \cap \mathbb{Q}[\mathbf{u}]$   
2  $\mathcal{D}_{XJ} \leftarrow$  the codimension 1 component of the equidimensional radical decomposition of  $\langle \mathcal{G}_u \rangle$   
3 **return**  $\mathcal{D}_{XJ}$

---

- we compute  $\mathcal{L}_{X\infty}$  by the Algorithm `PROPERNESSDEFECTS` presented in [20] and then get  $\mathcal{D}_{X\infty}$  by the radical equidimensional decomposition. We omit the formal description of the algorithm.

The previous algorithms in this subsection can not be used to compute DDs of algebraic statistical models in a reasonable time, see Tables 1–2 in Section 4. This motivates the exploration of a more practical method found in the next subsection.



### 3.2 Probabilistic Algorithm

First, we prepare the lemmas, then we present the main algorithm (Algorithm 2).

- Lemma 1 is used to linearly transform parameter space.
- Corollary 1 and Lemma 2 are used to compute the totally degree of  $\mathcal{D}_{XJ}$ .
- Corollary 2 is used in the sampling for interpolation.

**Lemma 1.** *For any  $G \in \mathbb{Q}[\mathbf{u}]$ , there exists an affine variety  $V$  in  $\mathbb{C}^n$  such that for any  $(a_1, \dots, a_n) \in \mathbb{C}^n \setminus V$ , the total degree of  $G$  equals the degree of  $B(t_0, t_1, \dots, t_n)$  w.r.t. to  $t_0$ , where*

$$B(t_0, t_1, \dots, t_n) = G(t_0, a_1 t_0 + t_1, \dots, a_n t_0 + t_n)$$

**Proof.** Suppose the total degree of  $G$  is  $d$  and  $G_d$  is the homogeneous component of  $G$  with total degree  $d$ . For any  $(1, a_1, \dots, a_n) \in \mathbb{C}^{n+1} \setminus \mathcal{V}_a(G_d)$ , let  $B(t_0, t_1, \dots, t_n) = G(t_0, a_1 t_0 + t_1, \dots, a_n t_0 + t_n)$ . It is easily seen that the degree of  $B$  w.r.t.  $t_0$  equals  $d$ .  $\square$

**Corollary 1.** *For any  $G \in \mathbb{Q}[\mathbf{u}]$ , there exists an affine variety  $V$  in  $\mathbb{C}^{2n+2}$  such that for any*

$$(a_0, b_0, \dots, a_n, b_n) \in \mathbb{C}^{2n+2} \setminus V,$$

*the total degree of  $G$  equals the degree of  $B(t)$  where*

$$B(t) = G(a_0 t + b_0, \dots, a_n t + b_n).$$

**Lemma 2.** *There exists an affine variety  $V$  in  $\mathbb{C}^{2n+2}$  such that for any  $(a_0, b_0, \dots, a_n, b_n) \in \mathbb{C}^{2n+2} \setminus V$ , if*

$$\langle A(t) \rangle = \langle F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J \rangle \cap \mathbb{Q}[t]$$

*where  $F_i(t)$  is the polynomial by replacing  $u_i$  with  $a_i t + b_i$  in  $F_i$  ( $i = 0, \dots, n$ ) and*

$$B(t) = \mathcal{D}_{XJ}(a_0 t + b_0, \dots, a_n t + b_n),$$

*then  $\langle B(t) \rangle = \sqrt{\langle A(t) \rangle}$ .*

**Proof.** By the definition of  $\mathcal{D}_{XJ}$  (Notation 6), there exists an affine variety  $V_1$  such that for any  $(a_0, b_0, \dots, a_n, b_n) \in \mathbb{C}^{2n+2} \setminus V_1$ ,  $\langle B(t) \rangle$  is radical. Thus, we only need to show that there exists an affine variety  $V_2$  in  $\mathbb{C}^{2n+2}$  such that for any  $(a_0, b_0, \dots, a_n, b_n) \in \mathbb{C}^{2n+2} \setminus V_2$ ,  $\mathcal{V}_a(\langle B(t) \rangle) = \mathcal{V}_a(\langle A(t) \rangle)$ .

Suppose  $\pi_t$  is the canonical projection:  $\mathbb{C} \times \mathbb{C}^{m+1} \rightarrow \mathbb{C}$ . For any

$$t^* \in \pi_t(\mathcal{V}_a(F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J)),$$

let  $u_i^* = a_i t^* + b_i$  (for  $i = 0, \dots, n$ ), then  $(u_0^*, \dots, u_n^*) \in \pi(\mathcal{L}_X \cap \mathcal{V}_a(J))$ . Hence  $\mathcal{D}_{XJ}(u_0^*, \dots, u_n^*) = 0$  and so  $B(t^*) = 0$ . Thus

$$B(t) \in \mathcal{I}(\pi_t(\mathcal{V}_a(F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J))).$$

Therefore,

$$\begin{aligned}\mathcal{V}_a(A(t)) &= \mathcal{V}_a(\mathcal{I}(\pi_t(\mathcal{V}_a(F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J))) \\ &\subset \mathcal{V}_a(B(t)).\end{aligned}$$

For any  $t^* \in \mathcal{V}_a(\langle B(t) \rangle)$ , let  $u_i^* = a_i t^* + b_i$  for  $i = 0, \dots, n$ , then  $(u_0^*, \dots, u_n^*) \in \mathcal{V}_a(\mathcal{D}_{XJ}) \subset \mathcal{L}_{XJ}$ . By the Extension Theorem [6], there exists an affine variety  $V_2 \subset \mathbb{C}^{2n+2}$  such that if  $(a_0, b_0, \dots, a_n, b_n) \notin V_2$ , then  $(u_0^*, \dots, u_n^*) \in \pi(\mathcal{L}_X \cap \mathcal{V}_a(J))$ , thus

$$t^* \in \pi_t(\mathcal{V}_a(F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J)) \subset \mathcal{V}_a(A(t)). \square$$

**Corollary 2.** *There exists an affine variety  $V$  in  $\mathbb{C}^n$  such that for any  $(a_1, \dots, a_n) \in \mathbb{C}^n \setminus V$ , if*

$$\langle A(u_0) \rangle = \langle F_0, F_1^* \dots, F_n^*, F_{n+1}, \dots, F_m, J \rangle \cap \mathbb{Q}[u_0]$$

where  $F_i^*$  is the polynomial by replacing  $u_i$  with  $a_i$  in  $F_i$  ( $i = 1, \dots, n$ ) and

$$B(u_0) = \mathcal{D}_{XJ}(u_0, a_1, \dots, a_n),$$

then  $\langle B(u_0) \rangle = \sqrt{\langle A(u_0) \rangle}$ .

---

### Algorithm 2: (Main Algorithm) InterpolationDX-J

---

```

input :  $F_0, \dots, F_m, J$ 
output:  $\mathcal{D}_{XJ}$ 
1  $a_1, \dots, a_n \leftarrow \text{LinearOperator}(F_0, \dots, F_m, J)$ 
2 for  $i$  from 1 to  $n$  do
3    $F'_i \leftarrow$  replace  $u_i$  in  $F_i$  with  $a_i u_0 + u_i$ 
4  $\text{NewSys} \leftarrow F_0, F'_1, \dots, F'_n, F_{n+1}, \dots, F_m, J$ 
5  $d, d_0, \dots, d_n \leftarrow \text{Degree}(\text{NewSys})$ 
6 for  $j$  from 1 to  $d$  do
7    $\left[ \begin{array}{l} \text{Rename all the monomials of the set} \\ \text{as } U_{j,1}, \dots, U_{j,N_j} \end{array} \right. \quad \{u_1^{\alpha_1} \dots u_n^{\alpha_n} | \alpha_1 + \dots + \alpha_n = j, 0 \leq \alpha_i \leq d_i\}$ 
8  $N \leftarrow \max(N_1, \dots, N_d)$ 
9 for  $k$  from 1 to  $N$  do
10   $b_{k,1}, \dots, b_{k,n} \leftarrow$  random integers
11   $A(u_0) \leftarrow \text{Intersect}(\text{NewSys}, b_{k,1}, \dots, b_{k,n})$ 
12   $C_{d,k}^*, \dots, C_{1,k}^* \leftarrow$  the coefficients of  $A(u_0)$  w.r.t  $u_0^0, \dots, u_0^{d-1}$ 
13 for  $j$  from 1 to  $d$  do
14   $\mathcal{M}_j \leftarrow N_j \times N_j$  matrix whose  $(k, r)$ -entry is  $U_{j,r}(b_{k,1}, \dots, b_{k,n})$ 
15   $C_j \leftarrow (U_{j,1}, \dots, U_{j,N_j}) \mathcal{M}_j^{-1} (C_{j,1}^*, \dots, C_{j,N_j}^*)^T$ 
16  $\mathcal{D}_{XJ} \leftarrow$  replace  $u_1, \dots, u_n$  in  $u_0^d + \sum_{i=0}^{d-1} C_{d-i} u_0^i$  with  $u_1 - a_1 \cdot u_0, \dots, u_n - a_n \cdot u_0$ 
17 Return  $\mathcal{D}_{XJ}$ 

```

---

We show an example to explain the basic idea of the probabilistic algorithm and how the lemmas work in the algorithm.

**Example 2 (Toy Example for Interpolation Idea).** Suppose the radical of the elimination ideal  $\langle F, J \rangle \cap \mathbb{Q}[\mathbf{u}]$  is generated by  $D(u_0, u_1, u_2)$ , where  $F = u_0p^2 + u_1p + u_2$  and  $J = 2u_0p + u_1$ . We already know that  $D$  is homogenous and equals  $u_1^2 - 4u_0u_2$ . Rather than by the standard elimination algorithm, we compute  $D$  by the steps below.

- First, we substitute  $u_0 = t + 11, u_1 = 3t + 2$  and  $u_2 = 5t + 6$  into  $F$  and  $J$  (the integers 1, 11, 3, 2, 5 and 6 are randomly chosen). We compute the radical of the elimination ideal  $\langle F(t, p), J(t, p) \rangle \cap \mathbb{Q}[t]$  and get  $\langle 11t^2 + 232t + 260 \rangle$ . By Lemma 2,  $D(t + 11, 3t + 2, 5t + 6) = 11t^2 + 232t + 260$ . By Corollary 1, the total degree of  $D$  is 2 (it geometrically means the random line  $u_0 = t + 11, u_1 = 3t + 2, u_2 = 5t + 6$  intersect our desired hypersurface at 2 points in the parameter space and it is exactly the definition of the degree of hypersurface). Similarly, we compute the degree of  $D$  w.r.t  $u_0, u_1$  and  $u_2$  and get 1, 2 and 1, respectively. So all the possible monomials in  $D$  are  $u_1^2, u_0u_1, u_1u_2$  and  $u_0u_2$ .

- Assume  $D = u_1^2 + (C_1u_0 + C_2u_2)u_1 + C_3u_0u_2$ . We first substitute  $u_0 = 13$  and  $u_2 = 4$  into  $F$  and  $J$ . We compute the radical of the elimination ideal  $\langle F(u_1, p), J(u_1, p) \rangle \cap \mathbb{Q}[u_1]$  and get  $\langle u_1^2 - 208 \rangle$ . By Corollary 2,  $D(13, u_1, 4)$  equals  $u_1^2 - 208$ . Hence,  $13C_1 + 4C_2 = 0$  and  $52C_3 = -208$ . Therefore,  $C_3 = -4$ . We need one more evaluation to solve  $C_1$  and  $C_2$ . So we substitute  $u_0 = 7$  and  $u_2 = 3$  into  $F$  and  $J$ . Similarly, we get  $7C_1 + 3C_2 = 0$  and thus  $C_1 = C_2 = 0$ . Therefore,  $D = u_1^2 - 4u_0u_2$  (the integers 13, 4, 7, 3 are randomly chosen).

This example is “nice”. Because the degree of  $D$  w.r.t  $u_1$  equals the total degree of  $D$ . In general case, if there is no  $u_i$  such that the degree of  $D$  w.r.t  $u_i$  equals the total degree, then we should apply the linear transformation to change the parameter coordinates before interpolation. Lemma 1 guarantees the linear transformation makes sense.

---

### Algorithm 3: Intersect

---

**input :**  $F_0, \dots, F_m, J$  and integers  $b_1, \dots, b_n$   
**output:**  $\mathcal{D}_{XJ}(u_0, b_1, \dots, b_n)$

- 1 **for**  $i$  **from** 1 **to**  $n$  **do**
- 2    $F_i^* \leftarrow$  replace  $u_i$  in  $F_i$  with  $b_i$
- 3  $A(u_0) \leftarrow$  the generator of the elimination ideal  $\langle F_0, F_1^*, \dots, F_n^*, F_{n+1}, \dots, F_m, J \rangle \cap \mathbb{Q}[u_0]$
- 4  $A(u_0) \leftarrow$  the monic generator of  $\sqrt{\langle A(u_0) \rangle}$
- 5 **return**  $A(u_0)$

---

Now we are prepared to introduce the probabilistic algorithm for computing the  $\mathcal{D}_{XJ}$ . We explain the main algorithm (Algorithm 2) and all the sub-algorithms (Algorithms 4–6) below.

**Algorithm 5 (Degree).** The probabilistic algorithm terminates correctly by Corollary 1 and Lemma 2.

**Algorithm 4 (LinearOperator).** The probabilistic algorithm terminates correctly by Lemma 1.

**Algorithm 3 (Intersect).** The probabilistic algorithm terminates correctly by Corollary 2.

**Algorithm 2 (InterpolationDX-J).**

**Lines 1–5.** We compute the total degree of  $\mathcal{D}_{XJ}$  and the degrees of  $\mathcal{D}_{XJ}$  w.r.t  $u_0, \dots, u_d$ :  $d, d_0, \dots, d_n$

---

**Algorithm 4: LinearOperator**


---

```

input :  $F_0, \dots, F_m, J$ 
output:  $a_1, \dots, a_n$  such that the total degree of  $\mathcal{D}_{XJ}$  equals the degree of  $\mathcal{D}_{XJ}(u_0, a_1 \cdot u_0 + u_1, \dots, a_n \cdot u_0 + u_n)$  w.r.t  $u_0$ 
1  $d, d_0, \dots, d_n \leftarrow \text{Degree}(F_0, \dots, F_m, J)$ 
2 if  $d = d_0$  then
3   return  $0, \dots, 0$ 
4 else
5   repeat
6     for  $i$  from 1 to  $n$  do
7        $a_i \leftarrow$  a random integer
8        $F'_i \leftarrow$  replace  $u_i$  in  $F_i$  with  $a_i \cdot u_0 + u_i$ 
9        $\text{NewSys} \leftarrow F_0, F'_1, \dots, F'_n, F_{n+1}, \dots, F_m, J$ 
10       $d, d_0, \dots, d_n \leftarrow \text{Degree}(\text{NewSys})$ 
11   until  $d = d_0$ 
12 return  $a_1, \dots, a_n$ 

```

---

by Algorithm 5. Algorithm 4 guarantees that  $d_0 = d$  by applying a proper linear transformation  $u_1 = a_1 \cdot u_0 + u_1, \dots, u_n = a_n \cdot u_0 + u_n$ .

**Lines 6–7.** Suppose  $\mathcal{D}_{XJ} = u_0^d + C_1 u_0^{d-1} + \dots + C_{d-1} u_0 + C_d$  where  $C_1, \dots, C_d \in \mathbb{Q}[u_1, \dots, u_n]$  and the total degree of  $C_j$  is  $j$ . For  $j = 1, \dots, n$ , we estimate all the possible monomials of  $C_j$  by computing the set

$$\{u_1^{\alpha_1} \dots u_n^{\alpha_n} \mid \alpha_1 + \dots + \alpha_n = j, 0 \leq \alpha_i \leq d_i\}$$

Assume the cardinality of the set is  $N_j$  and rename these monomials as  $U_{j,1}, \dots, U_{j,N_j}$ . Then we assume

$$C_j = c_{j,1} U_{j,1} + \dots + c_{j,N_j} U_{j,N_j}$$

where  $c_{j,1}, \dots, c_{j,N_j} \in \mathbb{Q}$ . The rest of the algorithm is to compute  $c_{j,1}, \dots, c_{j,N_j}$ .

**Lines 8–12.** For each  $j$ , for  $k = 1, \dots, N_j$ , for a random integer vector  $\mathbf{b}_k = (b_{k,1}, \dots, b_{k,n})$ , we compute  $\mathcal{D}_{XJ}(u_0, \mathbf{b}_k)$  by Algorithm 3. That means to compute the function value  $C_j(\mathbf{b}_k)$  without knowing  $\mathcal{D}_{XJ}$ .

**Lines 13–15.** For each  $j$ , we solve a square linear equation system for the unknowns  $c_{j,1}, \dots, c_{j,N_j}$ :

$$c_{j,1} U_{j,1}(\mathbf{b}_k) + \dots + c_{j,N_j} U_{j,N_j}(\mathbf{b}_k) = C_j(\mathbf{b}_k),$$

$$(k = 1, \dots, N_j)$$

It is known that we can choose nice  $\mathbf{b}_k$  probabilistically such that the coefficient matrix of the linear equation system is non-singular.

**Lines 16.** We apply the inverse linear transformation in the parameter space to get the  $\mathcal{D}_{XJ}$  for the original  $F_0, \dots, F_m$ .

We can also apply the interpolation idea to Algorithm PROPERNESSDEFECTS [20] and get a probabilistic algorithm to compute the  $\mathcal{D}_{X\infty}$ . We omit the formal description of the algorithm.

**Remark 3.** According to the Notation 6, when the codimension of  $\mathcal{L}_{XJ} \setminus \{\mathbf{0}\}$  ( $\mathcal{L}_{X\infty} \setminus \{\mathbf{0}\}$ ) is greater than 1, we define  $\mathcal{D}_{XJ}$  ( $\mathcal{D}_{X\infty}$ ) is 1. Therefore, it is no more true that the number of real/positive solutions still remains

---

**Algorithm 5: Degree**


---

```

input :  $F_0, \dots, F_m, J$ 
output:  $d, d_0, \dots, d_n$ , where  $d$  is the total degree of  $\mathcal{D}_{XJ}$  and  $d_i$  is the degree of  $\mathcal{D}_{XJ}$  w.r.t each  $u_i$  ( $i = 0, \dots, n$ )
1 for  $i$  from 0 to  $n$  do
2    $F_0^*, \dots, F_n^* \leftarrow$  replace  $u_0, \dots, u_{i-1}, u_{i+1}, \dots, u_n$  in  $F_0, \dots, F_n$  with random integers
3    $A(u_i) \leftarrow$  the generator of the elimination ideal  $\langle F_0^*, \dots, F_n^*, F_{n+1}, \dots, F_m, J \rangle \cap \mathbb{Q}[u_i]$ 
4    $A(u_i) \leftarrow$  the generator of  $\sqrt{\langle A(u_i) \rangle}$ 
5    $d_i \leftarrow$  degree of  $A(u_i)$ 
6    $a_i, b_i \leftarrow$  random integers
7  $F_0(t), \dots, F_n(t) \leftarrow$  replace  $u_0, \dots, u_n$  with  $a_0 \cdot t + b_0, \dots, a_n \cdot t + b_n$  in  $F_0, \dots, F_n$ 
8  $A(t) \leftarrow$  the generator of the elimination ideal  $\langle F_0(t), \dots, F_n(t), F_{n+1}, \dots, F_m, J \rangle \cap \mathbb{Q}[t]$ 
9  $A(t) \leftarrow$  the generator of  $\sqrt{\langle A(t) \rangle}$ 
10  $d \leftarrow$  degree of  $A(t)$ 
11 return  $d, d_0, \dots, d_n$ 

```

---

constant over the region determined by the data-discriminant. That means if the output of the Algorithm 2 is 1, we should use the standard method (elimination or computing Gröbner base).

## 4 Experimental Timings

We have implemented the probabilistic algorithm in Ma-caulay2. We have also implemented the standard algorithm in Macaulay2 to do comparisons (Tables 1 and 2). Some of the necessary implementation details are shown below.

- In the Algorithm 1. Line 1, Algorithm 3. Line 3 and Algorithm 5. Lines 3 and 8, we use the Macaulay2 command `eliminate` to compute the elimination ideals.

- The probabilistic algorithm is implemented in two different ways. The first implementation is to interpolate at once, which is exactly the same as the Algorithm 2. The second implementation is to interpolate step by step. For example, suppose the  $\mathcal{D}_{XJ}$  is a polynomial in  $u_0, u_1, u_2$  and  $u_3$ , we first compute  $\mathcal{D}_{XJ}(u_0, u_1, u_2^*, u_3^*)$  by interpolation for some chosen integers  $u_2^*$  and  $u_3^*$ . And then we compute  $\mathcal{D}_{XJ}(u_0, u_1, u_2, u_3^*)$  by interpolation. At this time, it is easy to recover  $\mathcal{D}_{XJ}$  since  $\mathcal{D}_{XJ}$  is homogeneous. The algorithm is naive to describe so we omit the formal description.

We run Algorithms 1 and 2 for many examples to set benchmarks by a 3.2 GHz Inter Core i5 processor (8GB total memory) under OS X 10.9.3. There are two kinds of benchmarks, the random models and literature models.

- We generate 2 groups of “random models”. The first group of random models are generated as follows. We first generate a random homogenous polynomial in 3 variables  $p_0, p_1$  and  $p_2$  with total degree 2. Suppose this homogenous polynomial is a model variant. We repeat the process for 10 rounds and get 10 random models. We call this group of 10 models 2deg-models. Similarly, we generate the group of 3deg-models. The Table 1 provides the timings of Algorithm 1 and Algorithm 2 (with two different implementations) for 2deg-models and 3deg-models.

- The literature models are the examples presented in the literatures [15, 7, 13]. Table 2 provides the timings of Algorithm 1 and Algorithm 2 (with two different implementations) for the literature models.

Table 1: Timings of Computing  $\mathcal{D}_{XJ}$  for Random Models (s: seconds; h: hours; S1: Strategy 1; S2: Strategy 2)

2deg-models			3deg-models		
Algorithm 1	Algorithm 2		Algorithm 1	Algorithm 2	
	S1	S2		S1	S2
4.9s	0.8s	0.6s	>2h	800.4s	901.2s
3.0s	0.7s	0.6s	>2h	777.3s	871.5s
5.0s	0.8s	0.6s	>2h	1428.9s	1499.5s
5.4s	0.8s	0.7s	>2h	1118.9s	1192.9s
6.3s	0.8s	0.7s	>2h	448.9s	489.8s
3.9s	0.7s	0.6s	>2h	1279.6s	1346.1s
2.0s	0.7s	0.5s	>2h	1286.5s	1409.0s
1.7s	0.7s	0.5s	>2h	1605.9s	1620.9s
3.8s	0.8s	0.6s	>2h	1099.4s	1242.6s
5.8s	0.8s	0.7s	>2h	1229.0s	1288.7s

For Examples 3–5 in the Table 2, the model invariants for these models are list below. Example 6 is given in Section 5.1.

**Example 3 (Random Censoring (Example 2.2.2 in [7])).**

$$2p_0p_1p_2 + p_1^2p_2 + p_1p_2^2 - p_0^2p_{12} + p_1p_2p_{12}$$

**Example 4 ( $3 \times 3$  Zero-Diagonal Matrix [13]).**

$$\det \begin{bmatrix} 0 & p_{12} & p_{13} \\ p_{21} & 0 & p_{23} \\ p_{31} & p_{32} & 0 \end{bmatrix}$$

**Example 5 (Grassmannian of 2-planes in  $\mathbb{C}^4$  [15, 13]).**

$$p_{12}p_{34} - p_{13}p_{24} + p_{14}p_{23}$$

In the Tables 1–2, the columns “Algorithm 1” give the timings of Algorithm 1. The columns “Algorithm 2” give the timings of Algorithm 2, where “S1” and “S2” means the first and second implementations, respectively. The red data means the computation has not finished and received no output. It is seen from the tables that

- for all the benchmarks we have tried, the Algorithm 2 is more efficient than Algorithm 1;
- for the random models and Example 3, the two implementations of Algorithm 2 have almost the same efficiency;
- for Examples 4–6, the second implementation (interpolation step by step) of Algorithm 2 is more efficient than the first implementation (interpolation at once). In fact, it takes the same time for the two implementations to get sample points. But it takes more time for the first implementation to compute the inverse of  $\mathcal{M}_j$  in Algorithm 2. Line 13, which is a large size matrix with rational entries.
- for Example 6, with the standard elimination algorithm, our computer runs out of memory after 12 days.

Note that for each benchmark, the output of Algorithm 2 is the same as Algorithm 1 when both algorithms terminate.

Table 2: Timings of Computing  $\mathcal{D}_{XJ}$  for Literature Models (s: seconds; h: hours; d: days; S1: Strategy 1; S2: Strategy 2)

Models	Algorithm 1	Algorithm 2	
		S1	S2
Example 3	11.1s	5.3s	6.4s
Example 4	36446.4s	360.2s	56.3s
Example 5	>16h	>16h	2768.2s
Example 6	>12d	>30d	30d

## 5 Conclusions and last example

In order to classify the data according to the number of real/positive solutions of likelihood equations, we study the data-discriminant and develop a probabilistic algorithm to compute it. Experiments show that the probabilistic algorithm is more practical than the standard elimination algorithm. This is our first application of real root classification method on the MLE/likelihood equations problem. Our future work aims to

- improve Algorithm 2 (note that Algorithm 2 is applying evaluation/interpolation technique to the standard method. It is not the first time that such an approach is investigated. In [9, 24], Newton–Hensel lifting has been applied to compute (parametric) geometric resolutions. It is hopeful that Algorithm 2 will be more powerful if we apply the Newton-Hensel lifting techniques to balance the time consuming of the evaluation and lifting steps);
- study the data-discriminants of different formulations of likelihood equations for the same algebraic statistical model
- develop algorithms for computing real root classification for likelihood equations.

More broadly, the ideas in Subsection 3.2 and Algorithm 2 can be applied to compute discriminants when the Newton polytope is known.

### 5.1 $3 \times 3$ symmetric matrix model

We end the paper with the discussion of real root classification on the  $3 \times 3$  symmetric matrix model.

Consider the following story with dice. A gambler has a coin, and two pairs of three-sided dice. The coin and the dice are all unfair. However, the two dice in the same pair have the same weight. He plays the same game 1000 rounds. In each round, he first tosses the coin. If the coin lands on side 1, he tosses the first pair of dice. If the coin lands on side 2, he tosses the second pair of dice. After the 1000 rounds, he records a  $3 \times 3$  data matrix  $[\bar{u}_{ij}]$  ( $i, j = 1, 2, 3$ ) where  $\bar{u}_{ij}$  is the the number of times for him to get the sides  $i$  and  $j$  with respect to the two dice. By the matrix  $[\bar{u}_{ij}]$ , he is trying to estimate the probability  $\bar{p}_{ij}$  of getting the sides  $i$  and  $j$  with respect to the two dice.

It is easy to check that the matrix

$$\begin{bmatrix} \bar{p}_{11} & \bar{p}_{12} & \bar{p}_{13} \\ \bar{p}_{21} & \bar{p}_{22} & \bar{p}_{23} \\ \bar{p}_{31} & \bar{p}_{32} & \bar{p}_{33} \end{bmatrix}$$

is symmetric and has at most rank 2. Let

$$p_{ij} = \begin{cases} \bar{p}_{ij} & i = j \\ \frac{1}{2}\bar{p}_{ij} & i < j \end{cases}, \quad u_{ij} = \begin{cases} \bar{u}_{ij} & i = j \\ \bar{u}_{ij} + \bar{u}_{ji} & i < j \end{cases}.$$

We have an algebraic statistical model below.

**Example 6 ( $3 \times 3$  Symmetric Matrix Model).** The algebraic statistical model for the dice story is given by

$$X = \mathcal{V}(g) \cap \Delta_5,$$

where

$$g = \det \begin{bmatrix} 2p_{11} & p_{12} & p_{13} \\ p_{12} & 2p_{22} & p_{23} \\ p_{13} & p_{23} & 2p_{33} \end{bmatrix},$$

$$\Delta_5 = \{(p_{11}, \dots, p_{33}) \in \mathbb{R}_{>0}^6 \mid p_{11} + p_{12} + p_{13} + p_{22} + p_{23} + p_{33} = 1\}.$$

The gambler's problem is equivalent to **maximizing the likelihood function**  $\frac{\Pi p_{ij}^{u_{ij}}}{(\Sigma p_{ij})^{\Sigma u_{ij}}}$  ( $i \leq j$ ) **subjected to**  $\mathcal{V}(g) \cap \Delta_5$ . According to the Definition 2, we present the Langrange likelihood equations below.

$$\begin{aligned} F_0 &= p_{11}\lambda_1 + (8p_{22}p_{33} - 2p_{23}^2)p_{11}\lambda_2 - u_{11} = 0 \\ F_1 &= p_{12}\lambda_1 + (2p_{13}p_{23} - 4p_{12}p_{33})p_{12}\lambda_2 - u_{12} = 0 \\ F_2 &= p_{13}\lambda_1 + (2p_{12}p_{23} - 4p_{13}p_{22})p_{13}\lambda_2 - u_{13} = 0 \\ F_3 &= p_{22}\lambda_1 + (8p_{11}p_{33} - 2p_{13}^2)p_{22}\lambda_2 - u_{22} = 0 \\ F_4 &= p_{23}\lambda_1 + (2p_{12}p_{13} - 4p_{11}p_{23})p_{23}\lambda_2 - u_{23} = 0 \\ F_5 &= p_{33}\lambda_1 + (8p_{11}p_{22} - 2p_{12}^2)p_{33}\lambda_2 - u_{33} = 0 \\ F_6 &= g(p_{11}, p_{12}, p_{13}, p_{22}, p_{23}, p_{33}) = 0 \\ F_7 &= p_{11} + p_{12} + p_{13} + p_{22} + p_{23} + p_{33} - 1 = 0 \end{aligned}$$

where  $p_{11}, p_{12}, p_{13}, p_{22}, p_{23}, p_{33}, \lambda_1$  and  $\lambda_2$  are unknowns and  $u_{11}, u_{12}, u_{13}, u_{22}, u_{23}$  and  $u_{33}$  are parameters.

We have 8 equations in 8 unknowns with 6 parameters and the ML-degree is 6 [15]. By the Algorithm 2, we have computed  $\mathcal{D}_{XJ}$ , which has 1307 terms with total degree 12. By a similar computation, we get  $\mathcal{D}_{X\infty}$ <sup>1</sup> whose last factor is exactly  $g(u_{11}, \dots, u_{33})$  and all the other factors are positive when each  $u_i$  is positive.

For the data-discriminant  $\mathcal{D}_X$  we have computed above, we have also computed<sup>2</sup> at least one rational point (sample point) from each open connected component of  $\mathcal{D}_X \neq 0$  using RAGLIB[22, 17, 11]. With these sample points we can solve the real root classification problem on the open cells. By testing all 236 sample points, we see that if  $g(u_{11}, \dots, u_{33}) \neq 0$ , then<sup>3</sup>

<sup>1</sup>See  $\mathcal{D}_{XJ}$  and  $\mathcal{D}_{X\infty}$  on the second author's website:  
sites.google.com/site/rootclassification/publications/DD

<sup>2</sup>The sample points were first successfully computed by one of the anonymous referees.

<sup>3</sup>This proves the real version of the RRC conjecture in the previous version of this manuscript.



– if  $\mathcal{D}_{XJ}(u_{11}, \dots, u_{33}) > 0$ , then the system has 6 distinct real solutions and there can be 6 positive solution or 2 positive solutions;

– if  $\mathcal{D}_{XJ}(u_{11}, \dots, u_{33}) < 0$ , then the system has 2 distinct real (positive) solutions.

With 2 of these sample points, we see that the sign of  $\mathcal{D}_X$  is not enough to classify the positive solutions. For example, for the sample point  $(u_{11} = 1, u_{12} = 1, u_{13} = \frac{280264116870825}{295147905179352825856}, u_{22} = 1, u_{23} = \frac{34089009205592922038535}{141080698675730650759168}, u_{33} = \frac{32898355113670387769001}{141080698675730650759168})$ , the system has 6 distinct positive solutions. While for the sample point  $(u_{11} = 1, u_{12} = 1, u_{13} = 199008, u_{22} = 30, u_{23} = 2022, u_{33} = 1)$ , the system has also 6 real solutions but only 2 positive solutions<sup>4</sup>.

## 6 Acknowledgments

We thank Professors Bernd Sturmfels, Hoon Hong, Jonathan Hauenstein and Frank Sottile for their valuable advice. We also thank Professors Mohab Safey El Din and Jean-Charles Faugere for their software advice on RAGlib and FGB respectively. We also especially thank the anonymous referees for their insightful suggestions to greatly improve the paper.

## References

- [1] M.-L. G. Buot, S. Hoşten, and D. Richards. Counting and locating the solutions of polynomial systems of maximum likelihood equations, ii: The behrens-fisher problem. *Statistica Sinica*, 17:1343–1354, 2007.
- [2] F. Catanese, S. Hoşten, A. Khetan, and B. Sturmfels. The maximum likelihood degree. *Amer. J. Math.*, 128(3):671–697, 2006.
- [3] C. Chen, J. H. Davenport, J. P. May, M. M. Maza, B. Xia, and R. Xiao. Triangular decomposition of semi-algebraic systems. In *Proceedings of ISSAC’10*, pages 187–194. ACM New York, 2010.
- [4] G. E. Collins and H. Hong. *Partial Cylindrical Algebraic Decomposition for Quantifier Elimination*. Springer, 1998.
- [5] M. Coste and M. Shiota. Nash triviality in families of nash manifolds. *Inventiones Mathematicae*, 108(1):349–368, 1992.
- [6] D. A. Cox, J. Little, and D. O’Shea. *Ideals, varieties, and algorithms: an introduction to computational algebraic geometry and commutative algebra*. Springer, 2007.
- [7] M. Drton, B. Sturmfels, and S. Sullivant. *Lectures on algebraic statistics*. Springer, 2009.
- [8] J.-C. Faugère, M. Safey El Din, and P.-J. Spaenlehauer. Gröbner bases and critical points: the unmixed case. In *Proceedings of ISSAC’12*, pages 162–169. ACM New York, 2012.
- [9] M. Giusti, G. Lecerf, and B. Salvy. A Gröbner free alternative for polynomial system solving. *Journal of Complexity*, 17:154–211, 2001.
- [10] D. R. Grayson and M. E. Stillman. *Macaulay2, a software system for research in algebraic geometry*.

---

<sup>4</sup>This disproves the positive version of the RRC conjecture in the previous version of this manuscript.

- [11] A. Greuet and M. Safey El Din. Probabilistic algorithm for the global optimization of a polynomial over a real algebraic set. *SIAM Journal on Optimization*, 24(3):1313–1343, 2014.
- [12] E. Gross, M. Drton, and S. Petrović. Maximum likelihood degree of variance component models. *Electronic Journal of Statistics*, 6:993–1016, 2012.
- [13] E. Gross and J. I. Rodriguez. Maximum likelihood geometry in the presence of data zeros. In *Proceedings of ISSAC’14*, pages 232–239. ACM New York, 2014.
- [14] J. Hauenstein, J. I. Rodriguez, and B. Sturmfels. Maximum likelihood for matrices with rank constraints. *Journal of Algebraic Statistics*, 5:18–38, 2014.
- [15] S. Hoşten, A. Khetan, and B. Sturmfels. Solving the likelihood equations. *Foundations of Computational Mathematics*, 5(4):389–407, 2005.
- [16] S. Hoşten and S. Sullivant. The algebraic complexity of maximum likelihood estimation for bivariate missing data. In *Algebraic and geometric methods in statistics*, pages 123–133. Cambridge University Press, 2009.
- [17] H. Hong and M. Safey El Din. Variant quantifier elimination. *Journal of Symbolic Computation*, 47(7):883–901, 2012.
- [18] J. Huh and B. Sturmfels. *Likelihood geometry*, pages 63–117. Springer International Publishing, 2014.
- [19] Z. Jelonek. Testing sets for properness of polynomial mappings. *Mathematische Annalen*, 315:1–35, 1999.
- [20] D. Lazard and F. Rouillier. Solving parametric polynomial systems. *Journal of Symbolic Computation*, 42(6):636–667, 2005.
- [21] J. I. Rodriguez. Maximum likelihood for dual varieties. In *Proceedings of the 2014 Symposium on Symbolic-Numeric Computation, SNC ’14*, pages 43–49, New York, NY, USA, 2014. ACM.
- [22] M. Safey El Din and E. Schost. Polar varieties and computation of one point in each connected component of a smooth real algebraic set. In *Proceedings of ISSAC’03*, pages 224–231. ACM Press, 2003.
- [23] M. Safey El Din and E. Schost. Properness defects of projections and computation of in each connected component of a real algebraic set. *Discrete and Computational Geometry*, 32(3):417–430, 2004.
- [24] E. Schost. Computing parametric geometric resolutions. *Applicable Algebra in Engineering, Communication and Computing*, 13(5):349–393, 2003.
- [25] C. Uhler. Geometry of maximum likelihood estimation in gaussian graphical models. *Annals of Statistics*, 40(1):238–261, 2012.
- [26] L. Yang, X. Hou, and B. Xia. A complete algorithm for automated discovering of a class of inequality-type theorems. *Science in China Series F Information Sciences*, 44(1):33–49, 2001.