

THE ACYCLICITY OF THE FROBENIUS FUNCTOR FOR MODULES OF FINITE FLAT DIMENSION

THOMAS MARLEY AND MARCUS WEBB

ABSTRACT. Let R be a commutative Noetherian local ring of prime characteristic p and $f : R \rightarrow R$ the Frobenius ring homomorphism. For $e \geq 1$ let $R^{(e)}$ denote the ring R viewed as an R -module via f^e . Results of Peskine, Szpiro, and Herzog state that for finitely generated modules M , M has finite projective dimension if and only if $\mathrm{Tor}_i^R(R^{(e)}, M) = 0$ for all $i > 0$ and all (equivalently, infinitely many) $e \geq 1$. We prove this statement holds for arbitrary modules using the theory of flat covers and minimal flat resolutions.

1. INTRODUCTION

Let R be a commutative Noetherian ring of prime characteristic p and $f : R \rightarrow R$ the Frobenius map. For $e \geq 1$ let $R^{(e)}$ be the ring R considered as an R -module via f^e ; i.e., for $r \in R, s \in R^{(e)}$, $r \cdot s := r^{p^e} s$. A classic result of Kunz [8] states that R is regular if and only if R is reduced and $R^{(e)}$ is flat as an R -module for all (equivalently, some) $e \geq 1$. Subsequently, Peskine and Szpiro [9, Théorème 1.7] proved that if \mathbf{P} is a finite projective resolution of a finitely generated R -module M then for all $e \geq 1$, $R^{(e)} \otimes_R \mathbf{P}$ is a projective resolution of $R^{(e)} \otimes_R M$; that is, finitely generated modules of finite projective dimension are acyclic objects with respect to the Frobenius functors $R^{(e)} \otimes_R -$. A year later Herzog [6, Satz 3.1] showed the converse holds in the case the Krull dimension of R is finite; namely, if M is a finitely generated R -module and \mathbf{P} is a projective resolution of M such that $R^{(e)} \otimes_R \mathbf{P}$ is acyclic for infinitely many integers e , then M has finite projective dimension. An interesting question is whether these results hold for arbitrary R -modules, not just finitely generated ones. In Corollary 4.5(a) and Theorem 5.2, we give an affirmative answer to this question:

Theorem 1.1. *Let R be a Noetherian ring of prime characteristic and M an R -module. Then the following hold:*

- (a) *If $\mathrm{fd}_R M < \infty$ then for all $i, e > 0$, $\mathrm{Tor}_i^R(R^{(e)}, M) = 0$ and $\mathrm{fd}_{R^{(e)}} R^{(e)} \otimes_R M = \mathrm{fd}_R M$.*
- (b) *If R has finite Krull dimension and $\mathrm{Tor}_i^R(R^{(e)}, M) = 0$ for all $i > 0$ and infinitely many integers e , then $\mathrm{pd}_R M < \infty$.*

Here $\mathrm{fd}_R M$ and $\mathrm{pd}_R M$ denote the flat and projective dimensions of M , respectively. Note that part (a) implies that if $\mathrm{pd}_R M < \infty$ then $\mathrm{pd}_{R^{(e)}} R^{(e)} \otimes_R M \leq \mathrm{pd}_R M$ for all e (although we do not know of any examples where the inequality is strict). We also prove an analogue of Theorem 1.1 for injective dimension in the case the Frobenius map is finite (cf. Corollaries 4.5(b) and 5.3).

Date: December 29, 2014.

2010 Mathematics Subject Classification. Primary 13D07; Secondary 13A35 .

Key words and phrases. flat dimension, Frobenius map, Bass number, flat cover.

A special case of part (a) of Theorem 1.1 is well-known and is what inspired this investigation: Suppose $\underline{x} = x_1, \dots, x_n$ is a regular sequence on R and $\mathbf{C} = \mathbf{C}(\underline{x}; R)$ is the Čech complex on R with respect to \underline{x} . Then \mathbf{C} is a finite flat resolution of the local cohomology module $H_{(\underline{x})}^n(R)$ and $R^{(e)} \otimes_R \mathbf{C} \cong \mathbf{C}(\underline{x}; R^{(e)}) \cong \mathbf{C}(\underline{x}^{p^e}; R)$ is acyclic since $x_1^{p^e}, \dots, x_n^{p^e}$ is an R -sequence.

The proofs of Peskine-Szpiro and Herzog both reduce to the case R is local and utilize minimal projective (free) resolutions. The minimality condition for such projective resolutions can be expressed by saying that all the differentials are zero when tensored with the residue field. It can be easily shown that projective resolutions of this type do not necessarily exist for arbitrary modules. However, flat resolutions with this kind of minimality condition do exist for a large class of modules (i.e., cotorsion modules), which we show is sufficient to prove Theorem 1.1. The theory of flat covers, cotorsion modules, and minimal flat resolutions, as developed by Enochs and Xu in [4] and [5], are essential ingredients in all our arguments. In Section 2, we summarize some basic properties of these notions as well as prove some auxiliary results which are used in later sections. Minimal flat resolutions have many properties analagous to those of minimal injective resolutions. In particular, the flat modules appearing in a minimal flat resolution of a module M are uniquely determined (up to isomorphism) by invariants which we call the *Enochs-Xu* numbers of M . The Enochs-Xu numbers of a module are in some sense the dual of the Bass numbers of a module. In Section 3, we prove a vanishing result for Enochs-Xu numbers which will be crucial in our proof of part (a) of Theorem 1.1.

Sections 4 and 5 are devoted to the proofs of parts (a) and (b), respectively, of Theorem 1.1. For both parts it is sufficient to consider the case when R is a local ring, in which case every module of finite flat dimension has finite projective dimension by the Jensen-Raynaud-Gruson theorem ([7, Proposition 6] and [10, Seconde partie, Théorème 3.2.6]). Our strategy is to apply the methods of Peskine-Szpiro and Herzog to minimal flat resolutions. There are several difficulties which arise: the finitistic flat dimension may exceed the depth of the ring (cf. [2, Corollary 5.3]); minimal flat resolutions do not in general localize; and the modules appearing in minimal flat resolutions are not generally finitely generated. We are able to overcome these difficulties by, in part, proving that the depths of the modules in degrees exceeding depth R in finite minimal flat resolutions are infinite, and showing that, in the case the Frobenius map is finite, the acyclicity of Frobenius for minimal flat resolutions of cotorsion modules commutes with the colocalization functor $\mathrm{Hom}_R(R_p, -)$. And while the flat modules in question are not typically finitely generated, we are able to reduce to the case where they are completions of free R -modules.

Acknowledgments: The authors would like to thank Edgar Enochs for his help in our understanding of flat covers. The authors also express their gratitude to everyone in the UNL algebra seminar, in particular Lucho Avramov, Peder Thompson, Mark Walker, and Wenliang Zhang, for their useful feedback on this project.

2. FLAT COVERS AND COTORSION MODULES

In this section we collect the results we will regarding flat covers, cotorsion modules, and minimal flat resolutions. We refer the reader to [13] for background on this material.

Definition 2.1. Let M be an R -module. An R -homomorphism $\varphi : F \rightarrow M$ is called a *flat cover* of M if the following hold:

- (a) F is flat;
- (b) for every map $\psi : G \rightarrow M$ with G flat, there exists a homomorphism $g : G \rightarrow F$ such that $\psi = \varphi g$; and
- (c) if $h : F \rightarrow F$ satisfies $\varphi = \varphi h$ then h is an isomorphism.

By abuse of language, we sometimes refer to the module F as the ‘flat cover’ of M , rather than the homomorphism φ . In [3], it is proved that flat covers exist for all modules over all rings. However, in this work we will only need the existence of flat covers for modules over commutative Noetherian rings of finite Krull dimension, which was proved in [12]. It is easily seen that flat covers are surjective (since every module is a homomorphic image of a flat module), that a flat cover of a module is unique up to isomorphism, and that the flat cover of a flat module is an isomorphism.

Definition 2.2. An R -module M is called *cotorsion* if $\text{Ext}_R^1(F, M) = 0$ for every flat R -module F .

It is easily seen that if M is cotorsion then $\text{Ext}_R^i(F, M) = 0$ for all $i \geq 1$ and all flat R -modules F . By [4, Lemma 2.2 and Corollary], the kernel of a flat cover is cotorsion and a flat cover of a cotorsion module is cotorsion. For an R -module M we let $C_R(M)$ denote the cotorsion envelope of M , which exists by [13, Theorem 3.4.6]. The cotorsion envelope of a flat module is flat by [13, Theorem 3.4.2]. If F is flat and cotorsion then $F \cong \prod_{p \in \text{Spec } R} T(p)$ where each $T(p)$ is the completion with respect to the pR_p -adic topology of a free R_p -module $G(p)$; furthermore, the ranks of the free R_p -modules $G(p)$ are uniquely determined by F ([4, Theorem]). For each $p \in \text{Spec } R$, let $\pi(p, F)$ denote the rank (possibly infinite) of $G(p)$.

Given an R -module M , a *minimal flat resolution* of M is a complex

$$\mathbf{F} : \quad \cdots \rightarrow F_i \xrightarrow{\partial_i} F_{i-1} \rightarrow \cdots \rightarrow F_0 \rightarrow 0$$

such that $H_i(\mathbf{F}) = 0$ for $i > 0$, $H_0(\mathbf{F}) \cong M$, and for each i , the natural map $F_i \rightarrow \text{coker } \partial_{i+1}$ is a flat cover. It is clear that every R -module has a minimal flat resolution, and that any two minimal flat resolutions of M are (chain) isomorphic. Since the flat cover of a flat module is an isomorphism, it follows that if $\text{fd}_R M = n < \infty$ then the length of any minimal flat resolution of M is n . Note that, by the remarks in the preceding paragraph, if \mathbf{F} is a minimal flat resolution of M then F_i is cotorsion for all $i \geq 1$. For each $i \geq 1$ and $p \in \text{Spec } R$ we set $\pi_i(p, M) := \pi(p, F_i)$. For $i = 0$ and $p \in \text{Spec } R$, we set $\pi_0(p, M) := \pi(p, C_R(F_0))$. Note that if M is cotorsion, so is F_0 and thus $\pi_0(p, M) = \pi(p, F_0)$ for all $p \in \text{Spec } R$. We call the invariants $\pi_i(p, M)$ the *Enochs -Xu numbers* of M . Note that $\text{fd}_R M \leq n$ if and only if $\pi_i(p, M) = 0$ for all $p \in \text{Spec } R$ and $i > n$.

We now state a theorem of Enochs and Xu:

Theorem 2.3. ([5, Theorem 2.1 and 2.2]) *Let R be a Noetherian ring. Then for any R -module M we have:*

- (a) $\pi_i(p, M) = \pi_i(p, C_R(M))$ for all $i \geq 0$ and $p \in \text{Spec } R$.
- (b) If M is cotorsion then $\pi_i(p, M) = \dim_{k(p)} \text{Tor}_i^{R_p}(k(p), \text{Hom}_R(R_p, M))$ for all $i \geq 0$ and $p \in \text{Spec } R$.

In particular, note that part (a) of this theorem implies that $\text{fd}_R M = \text{fd}_R C_R(M)$.

We now establish some additional results on cotorsion modules which are needed in sections 3 and 4. We thank Edgar Enochs for showing us a proof of part (b) of the following lemma, which is implicit in [5]:

Lemma 2.4. *Let R be a Noetherian ring and S a flat R -algebra.*

- (a) *If C is a cotorsion R -module and T a flat S -module then $\text{Hom}_R(T, C)$ is a cotorsion S -module.*
- (b) *If F is a flat and cotorsion R -module and $p \in \text{Spec } R$ then $\text{Hom}_R(R_p, F)$ is a flat and cotorsion R_p -module.*

Proof. To prove (a), let A be a flat S -module. As S is flat over R , any flat S -module is also flat as an R -module. Thus, $\text{Ext}_R^i(P \otimes_S A, C) = 0$ for all $i > 0$ and all projective S -modules P . Thus, we have a Grothendieck spectral sequence $\text{Ext}_R^p(\text{Tor}_q^S(T, A), C) \Rightarrow \text{Ext}_S^{p+q}(A, \text{Hom}_R(T, C))$ (cf. [11, Theorem 11.54]). The spectral sequence collapses, giving $\text{Ext}_S^i(A, \text{Hom}_R(T, C)) \cong \text{Ext}_R^i(T \otimes_S A, C) = 0$ for $i > 0$, as $T \otimes_S A$ is a flat S -module and therefore flat as an R -module as well. Hence, $\text{Hom}_R(T, C)$ is a cotorsion S -module.

For (b), as F is flat and cotorsion we have $F \cong \prod_{q \in \text{Spec } R} T(q)$ where each $T(q)$ is the completion of a free R_q -module. We'll show that $\text{Hom}_R(R_p, F) \cong \prod_{q \subseteq p} T(q)$, which is flat and cotorsion by [4, Theorem]. For each $q \in \text{Spec } R$ let $\rho_q : F \rightarrow T(q)$ be the projection map. Let $\varphi \in \text{Hom}_R(R_p, F)$. Then, as R_p is divisible by each element in $R \setminus p$, the same is true for the image $(\rho_q \varphi)(R_p)$ for all q . Suppose $q \not\subseteq p$. Let $r \in q \setminus p$. Then for all $n \geq 1$ we have $(\rho_q \varphi)(R_p) \subseteq r^n (\rho_q \varphi)(R_p) \subseteq q^n T(q)$. As $T(q)$ is separated in the qR_q -adic topology, we conclude that $(\rho_q \varphi)(R_p) = 0$. Thus, $\text{Hom}_R(R_p, F) \cong \text{Hom}_R(R_p, G)$, where $G = \prod_{q \subseteq p} T(q)$. But since each $T(q)$ for $q \subseteq p$ is an R_p -module, G is an R_p -module. Thus, $\text{Hom}_R(R_p, F) \cong G$. \square

Lemma 2.5. *Let R be a Noetherian ring of finite Krull dimension and M an R -module. Suppose M has a (left) cotorsion resolution \mathbf{C} ; i.e., there exists an exact sequence*

$$\cdots \rightarrow C_i \xrightarrow{\varphi_i} C_{i-1} \rightarrow \cdots \rightarrow C_0 \rightarrow M \rightarrow 0$$

where each C_i is cotorsion. Then

- (a) *M is cotorsion.*
- (b) *For any flat R -module T , $\text{Hom}_R(T, \mathbf{C})$ is a cotorsion resolution of $\text{Hom}_R(T, M)$.*

Proof. For each integer $i \geq 0$ let $K_i = \text{coker } \varphi_{i+1}$. Let T be a flat R -module. Since $\text{Ext}_R^i(T, C_j) = 0$ for all $i \geq 1$ and $j \geq 0$, we have that $\text{Ext}_R^1(T, M) \cong \text{Ext}_R^{i+1}(T, K_i)$ for all $i \geq 0$. Let $n = \dim R$. Then $\text{Ext}_R^{n+1}(T, K_n) = 0$ as $\text{pd}_R T \leq n$ by the Jensen-Raynaud-Gruson theorem ([7, Proposition 6] and [10, Seconde partie, Théorème 3.2.6]). Thus, $\text{Ext}_R^1(T, M) = 0$ and M is cotorsion. This proves (a).

For part (b), first note that $\text{Hom}_R(T, C_i)$ is cotorsion for all i by Lemma 2.4(a). Let $K_i = \text{coker } \varphi_{i+1}$ as above. Since each K_i has a cotorsion resolution, K_i is cotorsion by part (a). Applying $\text{Hom}_R(T, -)$ to the exact sequences $0 \rightarrow K_{i+1} \rightarrow C_i \rightarrow K_i \rightarrow 0$, we obtain that

$$0 \rightarrow \text{Hom}_R(T, K_{i+1}) \rightarrow \text{Hom}_R(T, C_i) \rightarrow \text{Hom}_R(T, K_i) \rightarrow 0$$

is exact for all i . Splicing these short exact sequences, we see that $\text{Hom}_R(T, \mathbf{C})$ is a cotorsion resolution of $\text{Hom}_R(T, M)$. \square

We remark that for any R -modules L , M , and N there is a natural R -module homomorphism

$$L \otimes_R \operatorname{Hom}_R(M, N) \xrightarrow{\psi} \operatorname{Hom}_R(M, L \otimes_R N)$$

such that for $\ell \in L$, $f \in \operatorname{Hom}_R(M, N)$ and $m \in M$, $\psi(\ell \otimes f)(m) = \ell \otimes f(m)$.

Lemma 2.6. *Let R be a Noetherian ring of finite Krull dimension. Let M , T , and F be R -modules such that M is finitely generated, T is flat, and F is flat and cotorsion. Then the natural map $\psi : M \otimes_R \operatorname{Hom}_R(T, F) \rightarrow \operatorname{Hom}_R(T, M \otimes_R F)$ is an isomorphism.*

Proof. We first note that the lemma is clear in the case M is a finitely generated free R -module. Let \mathbf{G} be a free resolution of M consisting of finitely generated free R -modules. As F is flat and cotorsion, $\mathbf{G} \otimes_R F$ is a cotorsion resolution of $M \otimes_R F$. By Lemma 2.5(b), we obtain that $\operatorname{Hom}_R(T, \mathbf{G} \otimes_R F)$ is a cotorsion resolution of $\operatorname{Hom}_R(T, M \otimes_R F)$. Now consider the commutative diagram:

$$\begin{array}{ccccccc} G_1 \otimes_R \operatorname{Hom}_R(T, F) & \longrightarrow & G_0 \otimes_R \operatorname{Hom}_R(T, F) & \longrightarrow & M \otimes_R \operatorname{Hom}_R(T, F) & \longrightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & \downarrow & & \\ \operatorname{Hom}_R(T, G_1 \otimes_R F) & \longrightarrow & \operatorname{Hom}_R(T, G_0 \otimes_R F) & \longrightarrow & \operatorname{Hom}_R(T, M \otimes_R F) & \longrightarrow & 0 \end{array}$$

where the rows are exact and the vertical arrows are the natural maps. Since the first two vertical maps are isomorphisms, so is the third by the Five Lemma. \square

We will need the following colocalization result for Tor :

Proposition 2.7. *Let R be a Noetherian ring of finite Krull dimension. Let L be a finitely generated R -module and M a cotorsion R -module. Suppose $\operatorname{Tor}_i^R(L, M) = 0$ for all $i > 0$. Then $\operatorname{Tor}_i^{R_p}(L_p, \operatorname{Hom}_R(R_p, M)) = 0$ for all $p \in \operatorname{Spec} R$ and $i > 0$.*

Proof. Let $p \in \operatorname{Spec} R$. It suffices to prove that $\operatorname{Tor}_i^R(L, \operatorname{Hom}_R(R_p, M)) = 0$ for all $i > 0$. Let \mathbf{F} be a minimal flat resolution of M . As M is cotorsion, so is F_i for all i . As $\operatorname{Tor}_i^R(L, M) = 0$ for all $i > 0$ we have that $L \otimes_R \mathbf{F}$ is acyclic. We remark that $L \otimes_R F_i$ is cotorsion for all i . To see this, let \mathbf{G} be a free resolution of L by finitely generated free R -modules. Then $\mathbf{G} \otimes_R F_i$ is a cotorsion resolution of $L \otimes_R F_i$. Hence, $L \otimes_R F_i$ is cotorsion by Lemma 2.5(a). By Lemma 2.5(b), $\operatorname{Hom}_R(R_p, L \otimes_R \mathbf{F})$ is acyclic. By Lemma 2.6, $L \otimes_R \operatorname{Hom}_R(R_p, \mathbf{F}) \cong \operatorname{Hom}_R(R_p, L \otimes_R \mathbf{F})$ as complexes. Thus, $L \otimes_R \operatorname{Hom}_R(R_p, \mathbf{F})$ is acyclic. However, $\operatorname{Hom}_R(R_p, \mathbf{F})$ is a flat resolution of $\operatorname{Hom}_R(R_p, M)$ by Lemmas 2.4(a) and 2.5(b). Thus, $\operatorname{Tor}_i^R(L, \operatorname{Hom}_R(R_p, M)) \cong H_i(L \otimes_R \operatorname{Hom}_R(R_p, \mathbf{F})) = 0$ for $i > 0$. \square

3. A VANISHING RESULT FOR BASS NUMBERS AND ENOCHS-XU NUMBERS

Recall that for an R -module M and $p \in \operatorname{Spec} R$, the Bass number $\mu_i(p, R)$ is defined to be $\dim_{k(p)} \operatorname{Ext}_{R_p}^i(k(p), M_p)$, where $k(p)$ is the field R_p/pR_p . If \mathbf{I} is a minimal injective resolution of M , then $\mu_i(p, M)$ is the cardinality of the number of copies of $E_R(R/p)$ in any decomposition of I^i into indecomposable modules. It is well-known that if M is finitely generated and $\operatorname{id}_R M < \infty$ then $\mu_i(p, M) = 0$ for all $p \in \operatorname{Spec} R$ and $i > \operatorname{depth} R_p$. The following proposition shows this result still holds even if M is not finitely generated.

Proposition 3.1. *Let R be a Noetherian ring and M an R -module such that $\operatorname{id}_R M < \infty$. Then $\mu_i(p, M) = 0$ for all $p \in \operatorname{Spec} R$ and $i > \operatorname{depth} R_p$.*

Proof. By localizing at p , we can assume R is local with maximal ideal m and that $p = m$. Let $\mu_i(M) := \mu_i(m, M)$, $s := \text{depth } R$, and $k := R/m$. Certainly $\mu_i(M) = 0$ for $i > \text{id}_R M$. Suppose $r > s$ and $\mu_i(M) = 0$ for all $i > r$. It suffices to show that $\mu_r(M) = 0$. Let $x_1, \dots, x_s \in m$ be a regular sequence and $I = (x_1, \dots, x_s)$. Since $\text{depth } R/I = 0$ we have an exact sequence

$$0 \longrightarrow k \longrightarrow R/I \longrightarrow R/J \longrightarrow 0$$

where $J = (I, y)$ for some $y \in m$. Since $\text{pd}_R R/I = s$, $\text{Ext}_R^i(R/I, M) = 0$ for $i > s$. From the long exact sequence on Ext , we have $\text{Ext}_R^i(k, M) \cong \text{Ext}_R^{i+1}(R/J, M)$ for $i > s$. As $r > s$ and $\mu_i(M) = 0$ for all $i > r$, we obtain $\text{Ext}_R^i(R/J, M) = 0$ for $i > r + 1$. Let $n := \text{id}_R M$, \mathbf{I} a minimal injective resolution of M and $\mathbf{L} := \text{Hom}_R(R/J, \mathbf{I})$. Then $H^i(\mathbf{L}) = \text{Ext}_R^i(R/J, M) = 0$ for $i > r + 1$, giving us an exact sequence

$$0 \longrightarrow K \longrightarrow L^{r+1} \xrightarrow{\partial^{r+1}} L^{r+2} \longrightarrow \dots \longrightarrow L^n \longrightarrow 0,$$

where $K = \ker \partial^{r+1}$. Now, for each $i \geq 0$ we have

$$\begin{aligned} L^i &= \text{Hom}_R(R/J, I^i) \\ &\cong \text{Hom}_R(R/J, \oplus E_R(R/p)^{\mu_i(p, M)}) \\ &\cong \bigoplus_{p \in V(J)} E_{R/J}(R/p)^{\mu_i(p, M)}. \end{aligned}$$

Note that for $p \neq m$, $\text{Tor}_j^R(k, E_{R/J}(R/p)) = 0$ for all j since $E_{R/J}(R/p)$ is an R_p -module. Since $\mu_i(M) = 0$ for $i \geq r + 1$, we see that $\text{Tor}_j^R(k, L_i) = 0$ for all j and all $i \geq r + 1$. Hence, from the long exact sequence above, we see that $\text{Tor}_j^R(k, K) = 0$ for all j . Now, we have an exact sequence

$$K \longrightarrow \text{Ext}_R^{r+1}(R/J, M) \longrightarrow 0.$$

Tensoring this sequence with k and noting that $\text{Ext}_R^{r+1}(R/J, M) \cong \text{Ext}_R^r(k, M)$, we conclude that $k \otimes_R \text{Ext}_R^r(k, M) = 0$. Hence, $\mu_r(M) = 0$. \square

Corollary 3.2. *Let R be a Noetherian ring and M an R -module such that $\text{fd}_R M < \infty$. Then $\text{Tor}_i^{R_p}(k(p), M_p) = 0$ for all $p \in \text{Spec } R$ and $i > \text{depth } R_p$.*

Proof. By localizing at p , we may assume R is local and $p = m$. Let $k = R/m$, $E = E_R(k)$, and let $(-)^v = \text{Hom}_R(-, E)$ be the Matlis dual functor. Since the Matlis dual is exact and T^v is injective for any flat R -module T , we see that $\text{id}_R M^v < \infty$. By Proposition 3.1 and [11, Corollary 11.54], we have $\text{Tor}_i^R(k, M)^v \cong \text{Ext}_R^i(k, M^v) = 0$ for $i > \text{depth } R$. Hence, $\text{Tor}_i^R(k, M) = 0$ for $i > \text{depth } R$. \square

We now prove the analogue of Proposition 3.1 for the Enochs-Xu numbers of a module of finite flat dimension:

Proposition 3.3. *Let R be a Noetherian ring and M an R -module such that $\text{fd}_R M < \infty$. Then $\pi_i(p, M) = 0$ for all $p \in \text{Spec } R$ and $i > \text{depth } R_p$.*

Proof. By Theorem 2.3(a), we may assume M is a cotorsion module. Let $p \in \text{Spec } R$. Note that $\text{Hom}_R(R_p, M)$ is a cotorsion R_p -module by Lemma 2.4(a) and $\text{fd}_{R_p} \text{Hom}_R(R_p, M) < \infty$ by Lemmas 2.4(b) and 2.5(b). Using part (b) of Theorem 2.3, we have that $\pi_i(p, M) =$

$\pi_i(pR_p, \text{Hom}_R(R_p, M))$ for all $i \geq 0$. Resetting notation, we can now assume R is a Noetherian local ring and $p = m$, the maximal ideal of R . By Theorem 2.3(b) and Corollary 3.2, we have $\pi_i(m, M) = \dim_{R/m} \text{Tor}_i^R(R/m, M) = 0$ for $i > \text{depth } R$. \square

4. AN ACYCLICITY THEOREM FOR FINITE FLAT RESOLUTIONS

We begin by making a convention regarding depth for (possibly) non-finitely generated modules. Let R be a Noetherian local ring with maximal ideal m and M an arbitrary R -module. For $i \geq 0$, let $H_m^i(M)$ denote the i th local cohomology module with support in m . We define the *depth* of M by

$$\text{depth } M := \inf\{i \in \mathbb{N}_0 \mid H_m^i(M) \neq 0\}.$$

Note that under this definition, $\text{depth } M = \infty$ if and only if $H_m^i(M) = 0$ for all i . We remark that if F is a flat R -module then $H_m^i(F) \cong H_m^i(R) \otimes_R F$; hence, $\text{depth } F \geq \text{depth } R$. We'll need the following result concerning the depths of certain cotorsion flat modules:

Lemma 4.1. *Let $\varphi : R \rightarrow S$ be a homomorphism of Noetherian local rings. Let m and n be the maximal ideals of R and S , respectively, and assume $\varphi(m) \subseteq n$. Let F be a cotorsion flat R -module such that $\pi(m, F) = 0$. Then for every S -module N we have $\text{depth}_S N \otimes_R F = \infty$. In particular, $\text{depth } F = \infty$.*

Proof. We first remark that as F is a flat R -module, $H_n^i(N \otimes_R F) \cong H_n^i(N) \otimes_R F$ for all i . Note that, as $mS \subseteq n$, $\text{Supp}_R H_n^i(N) \subseteq \{m\}$ for all i . Thus, it suffices to prove that given any R -module M with $\text{Supp}_R M \subseteq \{m\}$ then $M \otimes_R F = 0$. As F is flat and cotorsion, we have $F \cong \prod_{p \in \text{Spec } R} T(p)$ where $T(p)$ is the pR_p -adic completion of a free R_p -module of rank $\pi(p, F)$. As $\pi(m, F) = 0$, we can write this decomposition as $F \cong \prod_{p \neq m} T(p)$. Let M be an R -module of finite length. As R is Noetherian, M is finitely presented and hence $M \otimes_R F \cong \prod_{p \neq m} (M \otimes_R T(p)) = 0$, since $M \otimes_R R_p = 0$ for all $p \neq m$. Suppose now that M is an arbitrary R -module such that $\text{Supp}_R M \subseteq \{m\}$. Then M is the direct limit of the direct system $\{M_\alpha\}$ consisting of the finite length R -submodules of M (with morphisms the inclusion maps). Thus

$$M \otimes_R F \cong (\varinjlim M_\alpha) \otimes_R F \cong \varinjlim (M_\alpha \otimes_R F) = 0.$$

\square

Our next result is the acyclicity lemma of Peskine and Szpiro [9], but stated for complexes whose modules are not necessarily finitely generated. The proof is largely *mutatis mutandis*, but we include it for the convenience of the reader.

Proposition 4.2. ([9, Lemme d'acyclicité 1.8]) *Let R be a Noetherian local ring and consider a bounded complex \mathbf{T} of R -modules*

$$\mathbf{T} : \quad 0 \rightarrow T_s \xrightarrow{f_s} T_{s-1} \rightarrow \cdots \xrightarrow{f_0} T_0 \rightarrow 0.$$

Suppose the following two conditions hold for each $i > 0$:

- (1) $\text{depth } T_i \geq i$;
- (2) $\text{depth } H_i(\mathbf{T}) = 0$ or $H_i(\mathbf{T}) = 0$.

Then $H_i(\mathbf{T}) = 0$ for all $i > 0$.

Proof. If $s = 0$ there is nothing to show, so we assume $s > 0$. Since $H_s(\mathbf{T})$ is a submodule of T_s and $\text{depth } T_s \geq s \geq 1$, we see that $\text{depth } H_s(\mathbf{T}) > 0$. Thus, $H_s(\mathbf{T}) = 0$. Suppose for some $j \geq 1$ we have that $H_i(\mathbf{T}) = 0$ for $j < i \leq s$. It suffices to show that $H_j(\mathbf{T}) = 0$. Let $C_j = \text{im } f_{j+1}$. Then the truncated complex

$$0 \longrightarrow T_s \xrightarrow{f_s} T_{s-1} \longrightarrow \cdots \xrightarrow{f_{j+2}} T_{j+1} \longrightarrow C_j \longrightarrow 0$$

is exact. Using the inequalities $\text{depth } T_i \geq i$ and induction, one can show that $\text{depth } C_j \geq j + 1$. Let $K_j = \ker f_j$ and consider the exact sequence

$$0 \longrightarrow C_j \longrightarrow K_j \longrightarrow H_j(\mathbf{T}) \longrightarrow 0.$$

Since $K_j \subseteq T_j$ and $\text{depth } T_j \geq j \geq 1$, we conclude that $\text{depth } K_j \geq 1$. Since $\text{depth } C_j \geq 2$, we see that $\text{depth } H_j(\mathbf{T}) \geq 1$. Thus, $H_j(\mathbf{T}) = 0$. \square

We now prove the main result of this section, which generalizes [9, Corollary 1.10] to modules of finite flat dimension:

Theorem 4.3. *Let $\varphi : R \rightarrow S$ be a homomorphism of Noetherian rings such that for every $q \in \text{Spec } S$ and $p = \varphi^{-1}(q)$ one has that $\text{depth } S_q \geq \text{depth } R_p$. Let M be an R -module of finite flat dimension. Then*

- (a) $\text{Tor}_i^R(S, M) = 0$ for all $i > 0$.
- (b) $\text{fd}_S S \otimes_R M \leq \text{fd}_R M$.
- (c) If $k(p) \otimes_R S \neq 0$ for all $p \in \text{Spec } R$, then $\text{fd}_S S \otimes_R M = \text{fd}_R M$.

Proof. We prove part (a) by way of contradiction. Suppose $\text{Tor}_i^R(S, M) \neq 0$ for some $i \geq 1$. Let $q \in \text{Spec } S$ be a prime minimal with respect to the property that $\text{Tor}_i^R(S, M)_q \neq 0$ for some $i \geq 1$ and let $p := \varphi^{-1}(q)$. By replacing R , S , and M with R_p , S_q , and M_p , we may assume (R, m) and (S, n) are local rings, $\varphi(m) \subseteq n$, $\text{Tor}_i^R(S, M) \neq 0$ for some $i \geq 1$, and $\text{Supp}_S \text{Tor}_i^R(S, M) \subseteq \{n\}$ for all $i \geq 1$. Let \mathbf{F} be a minimal flat resolution of M and $\mathbf{L} := S \otimes_R \mathbf{F}$. Since $H_i(\mathbf{L}) \cong \text{Tor}_i^R(S, M)$ and $\text{Supp}_S \text{Tor}_i^R(S, M) \subseteq \{n\}$ for all $i \geq 1$, we have that $\text{depth } H_i(\mathbf{L}) = 0$ or $H_i(\mathbf{L}) = 0$ for all $i \geq 1$.

We claim that $\text{depth}_S L_i \geq i$ for all i . Since L_i is a flat S -module, $\text{depth}_S L_i \geq \text{depth } S$. Suppose $i > \text{depth } S$. Then F_i is a cotorsion flat R -module and $\pi(m, F_i) = \pi_i(m, M) = 0$ by Proposition 3.3 since $i > \text{depth } R$. Then $\text{depth}_S L_i = \infty$ by Lemma 4.1. This proves the claim. By Proposition 4.2, we obtain that $H_i(\mathbf{L}) = \text{Tor}_i^R(S, M) = 0$ for all $i \geq 1$, which is a contradiction. This proves part (a).

To see part (b), let \mathbf{F} be a flat resolution of M of length $n = \text{fd}_R M$. Then $S \otimes_R \mathbf{F}$ is an S -flat resolution of $S \otimes_R M$, since $\text{Tor}_i^R(S, M) = 0$ for all $i > 0$ by part (a). Hence, $\text{fd}_S S \otimes_R M \leq n$.

To prove part (c) it suffices by part (b) to show that $\text{fd}_S S \otimes_R M \geq \text{fd}_R M$. Let $n = \text{fd}_R M$ and J an ideal maximal with respect to $\text{Tor}_n^R(R/J, M) \neq 0$. Then by [1, Proposition 2.2], $J = p$ is prime and $\text{Tor}_n^R(R/p, M)$ is a (nonzero) $k(p)$ -module, where $k(p) = R_p/pR_p$. Let (\mathbf{F}, ∂) be a flat resolution of M of length n . Then we have an exact sequence

$$0 \longrightarrow \text{Tor}_n^R(R/p, M) \longrightarrow k(p) \otimes_R F_n \xrightarrow{1 \otimes \partial_n} k(p) \otimes_R F_{n-1}.$$

Now, tensoring with S over R (which is the same as tensoring by $S \otimes_R k(p)$ over $k(p)$, which is flat over $k(p)$), we have

$$0 \longrightarrow S \otimes_R \text{Tor}_n^R(R/p, M) \longrightarrow S \otimes_R k(p) \otimes_R F_n \xrightarrow{1 \otimes 1 \otimes \partial_n} S \otimes_R k(p) \otimes_R F_{n-1}$$

is exact. Reassociating, we obtain an exact sequence

$$0 \longrightarrow S \otimes_R \operatorname{Tor}_n^R(R/p, M) \longrightarrow (k(p) \otimes_R S) \otimes_S (S \otimes_R F_n) \xrightarrow{(1 \otimes 1) \otimes (1 \otimes \partial_n)} (k(p) \otimes_R S) \otimes_S (S \otimes_R F_{n-1}).$$

Since $S \otimes_R \mathbf{F}$ is a flat S -resolution of $S \otimes_R M$, this sequence shows that

$$\operatorname{Tor}_n^S(k(p) \otimes_R S, S \otimes_R M) \cong S \otimes_R \operatorname{Tor}_n^R(R/p, M) \cong S \otimes_R k(p)^\ell,$$

which is nonzero, since $l \neq 0$ and $S \otimes_R k(p) \neq 0$. Hence, $\operatorname{fd}_S S \otimes_R M \geq n$. \square

In the case the ring homomorphism $\varphi : R \longrightarrow S$ is finite, we have the following dual result for modules of finite injective dimension:

Corollary 4.4. *Let $\varphi : R \longrightarrow S$ be as in Theorem 4.3 and assume S is a finitely generated R -module. Let M be an R -module of finite injective dimension. Then*

- (a) $\operatorname{Ext}_R^i(S, M) = 0$ for all $i > 0$.
- (b) $\operatorname{id}_S \operatorname{Hom}_R(S, M) \leq \operatorname{id}_R M$.
- (b) If $k(p) \otimes_R S \neq 0$ for all $p \in \operatorname{Spec} R$, then $\operatorname{id}_S \operatorname{Hom}_R(S, M) = \operatorname{id}_R M$.

Proof. For part (a), it suffices to prove the statement locally at every prime ideal (as S is a f.g. R -module). So assume R is local and let $(-)^v$ denote the Matlis dual functor for R . By [11, Theorem 11.57], $\operatorname{Tor}_i^R(A, M^v) \cong \operatorname{Ext}_R^i(A, M)^v$ for all $i \geq 0$ and all finitely generated R -modules A . In particular, we have that $\operatorname{fd}_R M^v = \operatorname{id}_R M < \infty$. Applying part (a) of Theorem 4.3, we have $\operatorname{Ext}_R^i(S, M)^v \cong \operatorname{Tor}_i^R(S, M^v) = 0$ for all $i > 0$. Thus, $\operatorname{Ext}_R^i(S, M) = 0$ for all $i > 0$.

As in Theorem 4.3, part (b) follows readily from part (a).

To prove (c), it suffices to show that $\operatorname{id}_S \operatorname{Hom}_R(S, M) \geq \operatorname{id}_R M$. Let $n = \operatorname{id}_R M$. Then there exists $p \in \operatorname{Spec} R$ such that $\operatorname{Ext}_{R_p}^n(k(p), M_p) \neq 0$. An argument analagous to the one in the proof of part (c) of Theorem 4.3 shows that

$$\operatorname{Ext}_{S_p}^n(k(p) \otimes_R S, \operatorname{Hom}_R(S, M)_p) \cong \operatorname{Hom}_{R_p}(S \otimes_R k(p), \operatorname{Ext}_{R_p}^n(k(p), M_p)),$$

which is nonzero since $k(p) \otimes_R S \neq 0$ and $\operatorname{Ext}_{R_p}^n(k(p), M_p) \neq 0$. Thus, $\operatorname{id}_S \operatorname{Hom}_R(S, M) \geq n$. \square

We now apply our results to the Frobenius map. The following corollary generalizes Théorèmes 1.7 and 4.15 of [9], which were proved for finitely generated modules.

Corollary 4.5. *Let R be a Noetherian ring of prime characteristic, M an R -module, and $e \geq 1$ an integer.*

- (a) If $\operatorname{fd}_R M < \infty$ then $\operatorname{Tor}_i^R(R^{(e)}, M) = 0$ for all $i > 0$ and $\operatorname{fd}_{R^{(e)}} R^{(e)} \otimes_R M = \operatorname{fd}_R M$.
- (b) If the Frobenius map of R is finite and $\operatorname{id}_R M < \infty$ then $\operatorname{Ext}_R^i(R^{(e)}, M) = 0$ for all $i > 0$ and $\operatorname{id}_{R^{(e)}} \operatorname{Hom}_R(R^{(e)}, M) = \operatorname{id}_R M$.

Proof. It suffices to prove both (a) and (b) in the case $e = 1$. If $f : R \longrightarrow R$ is the Frobenius map, then for all $q \in \operatorname{Spec} R$, $f^{-1}(q) = q$ and $k(q) \otimes_R R^{(e)} \neq 0$ for all $e \geq 1$. The conclusions now follow from Theorem 4.3 and Corollary 4.4. \square

5. PROOF OF THE CONVERSE

In this section we prove part (b) of Theorem 1.1. Before doing so, we need to establish some results on completions of free modules. Let (R, m) be a Noetherian local ring and M an R -module. We let \widehat{M} denote the m -adic completion of M ; i.e., $\widehat{M} := \varprojlim M/m^n M$. If M is separated (i.e., $\cap_n m^n M = 0$), then the natural map $M \rightarrow \widehat{M}$ is injective. We note that for any $n \geq 1$, $M/m^n M \cong \widehat{M/m^n M} \cong \widehat{M}/\widehat{m^n M}$. In particular, if M is separated, then $\widehat{m^n M} \cap M = m^n M$. Finally, as m^n is finitely generated, we have that $m^n \widehat{M} = \widehat{m^n M}$.

Lemma 5.1. *Let (R, m) be a Noetherian local ring, F a free R -module, and N a finitely generated R -module. Let $t := t(R)$ be the least integer such that $m^t \cap H_m^0(R) = 0$. (Such a t exists by the Artin-Rees lemma.) Then the following hold:*

- (a) $H_m^0(\widehat{F}) = H_m^0(F)$.
- (b) For all $n \geq t$, $m^n \widehat{F} \cap H_m^0(\widehat{F}) = 0$.
- (c) $\widehat{F} \otimes_R N \cong \widehat{F \otimes_R N}$.

Proof. Let X be a basis for F . Then $F \cong \bigoplus_{\alpha \in X} R_\alpha$, where $R_\alpha = R$ for all α . We write this as $F = \bigoplus_\alpha R$ for short. Note that $m^n F = \bigoplus_\alpha m^n$ for all $n \geq 1$ and $H_m^0(F) = \bigoplus_\alpha H_m^0(R)$. In particular, F is separated. We also observe that the topology on $H_m^0(F)$ induced from the m -adic topology on F coincides with the m -adic topology on $H_m^0(F)$. (One can check this on each component.) Thus, $F/\widehat{H_m^0(F)} \cong \widehat{F}/\widehat{H_m^0(F)}$. Clearly, $\widehat{H_m^0(F)} = H_m^0(F)$ and $H_m^0(F) \subseteq H_m^0(\widehat{F})$. Therefore, to prove (a) it suffices to show that $H_m^0(F/\widehat{H_m^0(F)}) = 0$. If $R = H_m^0(R)$, this is clear. Otherwise, we can replace R by $R/H_m^0(R)$ and assume $H_m^0(R) = 0$. In this case, $H_m^0(F) = 0$ and it suffices to prove that $H_m^0(\widehat{F}) = 0$. Let $x \in m$ be a non-zero-divisor on R . Then $0 \rightarrow F \xrightarrow{\mu_x} F$ is exact, where μ_x is multiplication by x . We claim that $0 \rightarrow \widehat{F} \xrightarrow{\mu_x} \widehat{F}$ is exact. To see this, it suffices to show that the topology induced by $\mu_x^{-1}(m^n F)$ on F coincides with the m -adic topology on F . As x is a non-zero-divisor on R , there exists (by the Artin-Rees lemma) an integer t such that $(m^n :_R x) \subseteq m^{n-t}$ for all $n \geq t$. Then $\mu_x^{-1}(m^n F) = \bigoplus_\alpha (m^n :_R x) \subseteq m^{n-t} F$ for all $n \geq t$. Hence the topologies coincide, and x is a non-zero-divisor on \widehat{F} . Thus, $H_m^0(\widehat{F}) = 0$ and part (a) is proved.

Using part (a) and that $m^n \widehat{F} \cap F = m^n F$, we have for all $n \geq t$

$$\begin{aligned} m^n \widehat{F} \cap H_m^0(\widehat{F}) &= m^n F \cap H_m^0(F) \\ &= \bigoplus_\alpha (m^n \cap H_m^0(R)) \\ &= 0. \end{aligned}$$

Thus, (b) is proved.

We now prove part (c). Let $R^r \xrightarrow{\varphi} R^s \rightarrow N \rightarrow 0$ be a presentation for N . Let $K = \text{im } \varphi$. By the Artin-Rees lemma, there exists an integer $t \geq 1$ such that $m^n R^s \cap K \subseteq m^{n-t} K$ for all $n \geq t$. As F is free, we have $0 \rightarrow F \otimes_R K \rightarrow F \otimes_R R^s \rightarrow F \otimes_R N \rightarrow 0$ is exact. Then

for $n \geq t$,

$$\begin{aligned} m^n(F \otimes_R R^s) \cap (F \otimes_R K) &= (\oplus_\alpha m^n R^s) \cap \oplus_\alpha K \\ &= \oplus_\alpha (m^n R^s \cap K) \\ &\subseteq \oplus_\alpha m^{n-t} K \\ &\subseteq m^{n-t}(F \otimes_R K). \end{aligned}$$

Thus, the induced and m -adic topologies on $F \otimes_R K$ coincide. Therefore,

$$0 \longrightarrow \widehat{F \otimes_R K} \longrightarrow \widehat{F \otimes_R R^s} \longrightarrow \widehat{F \otimes_R N} \longrightarrow 0$$

is exact. Composing with the surjection $\widehat{F \otimes_R R^r} \xrightarrow{1 \otimes \varphi} \widehat{F \otimes_R R^s}$ and noting that $\widehat{A} \otimes_R R^n$ is naturally isomorphic to $\widehat{A^n}$ for any R -module A and any positive integer n , we obtain the commutative diagram:

$$\begin{array}{ccccccc} \widehat{F \otimes_R R^r} & \xrightarrow{1 \otimes \varphi} & \widehat{F \otimes_R R^s} & \longrightarrow & \widehat{F \otimes_R N} & \longrightarrow & 0 \\ \downarrow \cong & & \downarrow \cong & & & & \\ \widehat{F} \otimes_R R^r & \xrightarrow{1 \otimes \varphi} & \widehat{F} \otimes_R R^s & \longrightarrow & \widehat{F} \otimes_R N & \longrightarrow & 0. \end{array}$$

Hence, $\widehat{F \otimes_R N} \cong \widehat{F} \otimes_R N$ by the Five Lemma. \square

Before proving part (b) of Theorem 1.1, we introduce some notation used in the proof. For $e \geq 1$ we let m_e denote the maximal ideal of $R^{(e)}$. For $x \in R$, we let x_e denote the element x considered as an element in $R^{(e)}$. Thus, $xR^{(e)} = x_e^{p^e} R^{(e)}$. Given an R -module N we let $N^{(e)}$ denote the $R^{(e)}$ -module $R^{(e)} \otimes_R N$. If $\psi : L \rightarrow N$ is an R -module homomorphism, $\psi^{(e)}$ denotes the map $1 \otimes_R \psi : L^{(e)} \rightarrow N^{(e)}$.

Theorem 5.2. *Let R be a Noetherian ring of prime characteristic p and M an R -module. Assume R has finite Krull dimension. Suppose $\mathrm{Tor}_i^R(R^{(e)}, M) = 0$ for all $i > 0$ and infinitely many integers e . Then $\mathrm{pd}_R M < \infty$.*

Proof. We first note that it suffices to prove $\mathrm{fd}_R M < \infty$ by the Jensen-Raynaud-Gruson theorem. Let $d := \dim R$. As $\mathrm{fd}_R M < \infty$ if and only if $\mathrm{Tor}_{d+1}^{R_q}(M_q, N_q) = 0$ for every R -module N and $q \in \mathrm{Spec} R$, we can assume R is local. By a standard argument, there exists a faithfully flat local R -algebra (T, n) such that $T^{(e)}$ is f.g. as a T -module for all e (e.g., [8, Section 3]). Note that $\mathrm{fd}_R M = \mathrm{fd}_T(T \otimes_R M)$ and $\mathrm{Tor}_i^T(T^{(e)}, T \otimes_R M) \cong T^{(e)} \otimes_{R^{(e)}} \mathrm{Tor}_i^R(R^{(e)}, M)$ for all i and e . Hence, by replacing R by T , we may assume $R^{(e)}$ is a finitely generated R -module for all e .

Let $\varphi : F \rightarrow M$ be a flat cover and $K = \ker \varphi$. By [4, Lemma 2.2], K is cotorsion. Note that $\mathrm{fd}_R K < \infty$ if and only if $\mathrm{fd}_R M < \infty$ and $\mathrm{Tor}_i^R(R^{(e)}, K) \cong \mathrm{Tor}_{i+1}^R(R^{(e)}, M)$ for all $i \geq 1$ and all e . Hence, we may assume that M is cotorsion.

We assume that $\mathrm{fd}_R M = \infty$ and derive a contradiction. Choose $q \in \mathrm{Spec} R$ minimal with respect to the property that $\mathrm{fd}_{R_q} \mathrm{Hom}_R(R_q, M) = \infty$. By Proposition 2.7, if $\mathrm{Tor}_i^R(R^{(e)}, M) = 0$ for some e and all $i > 0$, then $\mathrm{Tor}_i^{R_q}(R_q^{(e)}, \mathrm{Hom}_R(R_q, M)) = 0$ for all $i > 0$. Note also that $\mathrm{Hom}_R(R_q, M)$ is a cotorsion R_q -module by Lemma 2.4(a) and that $R_q^{(e)}$ is finitely generated as an R_q -module for all $e \geq 1$. Thus, by replacing R with R_q and

M with $\text{Hom}_R(R_q, M)$, we may assume $\text{fd}_R M = \infty$ and $\text{fd}_{R_q} \text{Hom}_R(R_q, M) < \infty$ for all $q \neq m$.

Let \mathbf{F} be a minimal flat resolution of M and let $\varphi_i : F_i \rightarrow F_{i-1}$ denote the differentials of \mathbf{F} . Since M is cotorsion, so is F_i for all i . By the proof of [5, Theorem 2.2], $\varphi_i \otimes_R 1_{R/m} = 0$ for all i ; i.e., $\varphi_i(F_i) \subseteq mF_{i-1}$ for all i . Then $\varphi_i^{(e)}(F_i^{(e)}) \subseteq m_e^{[p^e]} F_{i-1}^{(e)}$ for all i and e .

Let $s := \text{depth } R$ and $\mathbf{x} \in m$ a regular sequence on R of length s . Let $t := t(R/(\mathbf{x}))$ as defined in Lemma 5.1, and let e be an integer such that $m^{[p^e]} \subseteq m^t$ and $\text{Tor}_i^R(R^{(e)}, M) = 0$ for all $i > 0$. Let S denote the $R^{(e)}$ -algebra $R^{(e)}/(\mathbf{x}_e)$ and $n = m_e S$, the maximal ideal of S . Then $H_n^0(S) \neq 0$ and $n^t \cap H_n^0(S) = 0$ by definition of t . Since $\text{pd}_{R^{(e)}} S = s$, we have that $\text{Tor}_i^{R^{(e)}}(S, M^{(e)}) = 0$ for all $i > s$. As $\mathbf{F}^{(e)}$ is an $R^{(e)}$ -flat resolution of $M^{(e)}$, we obtain that $H_i(S \otimes_{R^{(e)}} \mathbf{F}^{(e)}) = 0$ for all $i > s$. For each $i \geq 0$ let $C_i = \text{im } \varphi_{i+1}$. Then for all $i \geq 1$ we have an exact sequence of $R^{(e)}$ -modules

$$0 \rightarrow C_i^{(e)} \rightarrow F_i^{(e)} \rightarrow C_{i-1}^{(e)} \rightarrow 0.$$

From the remarks above, we have that $C_i^{(e)} = \varphi_{i+1}^{(e)}(F_{i+1}^{(e)}) \subseteq m_e^{[p^e]} F_i^{(e)} \subseteq m_e^t F_i^{(e)}$ for all i . Since $H_i(S \otimes_R \mathbf{F}^{(e)}) = 0$ for all $i > s$, we have that

$$0 \rightarrow S \otimes_{R^{(e)}} C_i^{(e)} \rightarrow S \otimes_{R^{(e)}} F_i^{(e)} \rightarrow S \otimes_{R^{(e)}} C_{i-1}^{(e)} \rightarrow 0$$

is exact for all $i > s - 1$. Note that $S \otimes_{R^{(e)}} C_i^{(e)} \subseteq n^t(S \otimes_{R^{(e)}} F_i^{(e)})$ for all $i > s - 1$.

By our assumptions, we have that $\text{fd}_{R_q} \text{Hom}_R(R_q, M) \leq d - 1$ for all $q \neq m$, where $d = \dim R$. By Theorem 2.3(b), this implies that $\pi(q, F_i) = \pi_i(q, M) = \pi_i(qR_q, \text{Hom}_R(R_q, M)) = 0$ for all $i \geq d$ and all $q \neq m$. Consequently, for $i \geq d$, F_i is the completion of a free R -module G_i of rank $r_i := \pi_i(m, M)$. By Lemma 5.1(c), for $i \geq d$ we have that $F_i^{(e)} \cong R^{(e)} \otimes_R \widehat{G_i} \cong \widehat{G_i^{(e)}}$ and $S \otimes_{R^{(e)}} F_i^{(e)} \cong S \otimes_{R^{(e)}} \widehat{G_i^{(e)}}$, which is the n -adic completion of a free S -module of rank r_i . Note that since $\text{fd}_R M = \infty$, $r_i \neq 0$ for all $i \geq d$. For all $i \geq d - 1$, let $B_i = S \otimes_{R^{(e)}} C_i^{(e)}$ and $V_i = S \otimes_{R^{(e)}} F_i^{(e)}$. From above, for $i \geq d$ we have exact sequences of S -modules

$$0 \rightarrow B_i \rightarrow V_i \rightarrow B_{i-1} \rightarrow 0$$

where V_i is the completion of a (nonzero) free S -module and $B_i \subseteq n^t V_i$. In particular, by Lemma 5.1(a), $H_n^0(V_i) \neq 0$ for all $i \geq d$. By Lemma 5.1(b), $H_n^0(B_i) \subseteq n^t V_i \cap H_n^0(V_i) = 0$ for all $i \geq d$. This implies that $H_n^0(V_i) = 0$ for $i \geq d + 1$, a contradiction. \square

In the case the Frobenius map is finite, we obtain the converse to part (b) of Corollary 4.5. This generalizes [6, Satz 5.2], which was proved in the case the module M is finitely generated.

Corollary 5.3. *Let R be a Noetherian ring of prime characteristic. Assume that R has finite Krull dimension and that the Frobenius map is finite. Let M be an R -module and suppose that $\text{Ext}_R^i(R^{(e)}, M) = 0$ for all $i > 0$ and for infinitely many integers e . Then $\text{id}_R M < \infty$.*

Proof. Since $R^{(e)}$ is a finitely generated R -module for all e and $\dim R < \infty$, we may assume R is local. Let $(-)^v$ be the Matlis dual functor. Then, as in the proof of Corollary 4.4(a), $\text{id}_R M = \text{fd}_R M^v$. By [11, Theorem 11.57], $\text{Tor}_i^R(R^{(e)}, M^v) \cong \text{Ext}_R^i(R^{(e)}, M)^v = 0$ for all $i > 0$ and infinitely many e . By Theorem 5.2, $\text{id}_R M = \text{fd}_R M^v < \infty$. \square

REFERENCES

1. A. Auslander and D. Buchsbaum, *Homological Dimension in Noetherian rings II*, Trans. Amer. Math. Soc. **88** (1958), 194–206.
2. H. Bass, *Injective dimension in Noetherian rings*, Trans. Amer. Math. Soc. **102** (1962), 319–327.
3. L. Bican, R. El Bashir, and E. Enochs, *All modules have flat covers*, Bull. London Math. Soc. **33** (2001), no. 4, 385–390.
4. E. Enochs, *Flat covers and flat cotorsion modules*, Proc. Amer. Math. Soc. **92** (1984), 179–184.
5. E. Enochs and J. Xu, *On Invariants dual to the Bass numbers*, Proc. Amer. Math. Soc. **125** (1997), 951–960.
6. J. Herzog, *Ringe der Charakteristik p und Frobeniusfunktoren*, Math. Z. **140** (1974), 67–78.
7. C. Jensen, *On the vanishing of $\varprojlim^{(i)}$* , J. Algebra **15** (1970), 151–166.
8. E. Kunz, *Characterizations of regular local rings for characteristic p* , Amer. J. Math. **91** (1969), 772–784.
9. C. Peskine and L. Szpiro, *Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck*, Inst. Hautes Études Sci. Publ. Math. No. 42 (1973), 47–119.
10. M. Raynaud and L. Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89.
11. J. Rotman, *An Introduction to Homological Algebra*, Pure and Applied Mathematics no. **85**, Academic Press, Inc., New York-London, 1979.
12. J. Xu, *The existence of flat covers over Noetherian rings of finite Krull dimension*, Proc. Amer. Math. Soc. **123** (1995), 27–32.
13. J. Xu, *Flat covers of modules*, Lecture Notes in Mathematics, vol. 1634, Springer-Verlag, Berlin, 1996.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE 68588-0130
E-mail address: tmarley1@unl.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEBRASKA-LINCOLN, LINCOLN, NE 68588-0130
E-mail address: mwebb4@math.unl.edu