

# Bicoloring covers for graphs and hypergraphs

Tapas Kumar Mishra      Sudebkumar Prasant Pal

Department of Computer Science and Engineering  
IIT Kharagpur 721302  
India

## Abstract

Let the *bicoloring cover number*  $\chi^c(G)$  for a hypergraph  $G(V, E)$  be the minimum number of bicolourings of vertices of  $G$  such that every hyperedge  $e \in E$  of  $G$  is properly bicolored in at least one of the  $\chi^c(G)$  bicolourings. We establish a tight bound for  $\chi^c(K_n^k)$ , where  $K_n^k$  is the complete  $k$ -uniform hypergraph on  $n$  vertices. We investigate the relationship between  $\chi^c(G)$ , matchings, hitting sets,  $\alpha(G)$  (independence number) and  $\chi(G)$  (chromatic number). We design a factor  $O(\frac{\log n}{\log \log n - \log \log \log n})$  approximation algorithm for computing a bicoloring cover. We define a new parameter for hypergraphs - "cover independence number  $\gamma(G)$ " and prove that  $\log \frac{|V|}{\gamma(G)}$  and  $\frac{|V|}{2\gamma(G)}$  are lower bounds for  $\chi^c(G)$  and  $\chi(G)$ , respectively. We show that  $\chi^c(G)$  can be approximated by a polynomial time algorithm achieving approximation ratio  $\frac{1}{1-t}$ , if  $\gamma(G) = n^t$ , where  $t < 1$ . We also construct a particular class of hypergraphs  $G(V, E)$  called *cover friendly* hypergraphs where the ratio of  $\alpha(G)$  to  $\gamma(G)$  can be arbitrarily large. We prove that for any  $t \geq 1$ , there exists a  $k$ -uniform hypergraph  $G$  such that the *clique number*  $\omega(G) = k$  and  $\chi^c(G) > t$ . Let  $m(k, x)$  denote the minimum number of hyperedges such that some  $k$ -uniform hypergraph  $G$  with  $m(k, x)$  hyperedges does not have a bicoloring cover of size  $x$ . We show that  $2^{(k-1)x-1} < m(k, x) \leq x \cdot k^2 \cdot 2^{(k+1)x+2}$ . Let the *dependency*  $d(G)$  of  $G$  be the maximum number of hyperedge neighbors of any hyperedge in  $G$ . We propose an algorithm for computing a bicoloring cover of size  $x$  for  $G$  if  $d(G) \leq (\frac{2^{x(k-1)}}{e} - 1)$  using  $nx + kx \frac{m}{d}$  random bits.

Keywords: Hypergraph bicoloring, local lemma, probabilistic method, Kolmogorov complexity, approximation

## 1 Introduction

We define the *bicoloring cover number*  $\chi^c(G)$  for a hypergraph  $G(V, E)$  as the minimum number of bicolourings such that every hyperedge  $e \in E$  of  $G$  is properly bicolored in at least one of the  $\chi^c$  bicolourings. Let  $\mathcal{X}$  be a set of bicolourings  $\{X_1, X_2, \dots, X_t\}$ . Then  $\mathcal{X}$  is a bicoloring cover for  $G$  if for each hyperedge  $e$  of  $G$ , there is an integer  $i \in \{1, 2, \dots, t\}$ , such that  $e$  is non-monochromatic with respect to bicoloring  $X_i$ .

Consider the scenario where  $n$  doctors can each be assigned one of two kinds of *tasks*; either he can see patients or perform operations. All doctors are equivalent and can perform only one of the two tasks in each *group*. There are  $m$  groups made from this set of  $n$  doctors viz.,  $E_1, E_2, \dots, E_m$ , where each group is of size  $k$ . Any doctor can be a member of multiple groups. In order to provide proper treatment, all the  $k$  members of no group should be assigned the same task; each group must have at least one doctor seeing patients and at least one doctor performing operations.

Given  $n$  doctors and  $m$  groups of doctors, is there a possible allocation of *tasks* to doctors such that none of the groups has all doctors allocated the same task? This problem can be viewed as the hypergraph bicoloring problem for the  $k$ -uniform hypergraph  $G(V, E)$ , where  $n = |V|$ ,  $m = |E|$ . Here, the doctors represent vertices, the groups represent  $k$ -uniform hyperedges, and the tasks assigned to doctors represent the two colors for bicoloring vertices. However, there exist hypergraphs that are not bicolorable. For such hypergraphs, it makes sense to use a bicoloring cover with  $\chi^c$  bicolorings. Instead of all  $m$  groups of doctors being deployed simultaneously, we could have a minimum number  $\chi^c$  of deployments, one for each of the bicolorings from a bicoloring cover for  $G(V, E)$ . Note that in any of these bicolorings, the same doctor can serve in multiple groups. Observe that if we have to deploy each of the  $m$  groups of doctors effectively, then we need at least  $\chi^c$  bicolorings, where each bicoloring yields one shift of duty assignments. The minimum number of shifts required for deploying all the  $m$  groups of doctors, is therefore the bicoloring cover number  $\chi^c(G)$ . Throughout the paper,  $G$  denotes a  $k$ -uniform hypergraph with vertex set  $V$  and hyperedge set  $E$ , unless otherwise stated. We use  $V(G)$  and  $V$ , and  $E(G)$  and  $E$  interchangeably. All logarithms are to the base two unless specified otherwise.

## 1.1 Related works

Graph decomposition is a widely studied problem in graph theory. The main idea of the problem is whether a given graph  $G(V, E)$  can be decomposed into some family of smaller graphs i.e., is there a family of graphs  $\mathcal{H} = \{H_1, \dots, H_j\}$  such that (1).  $V(H_i) \subseteq V(G)$  for all  $H_i \in \mathcal{H}$ , (2).  $\cap_{H_i \in \mathcal{H}} E(H_i) = \emptyset$  and (3).  $\cup_{H_i \in \mathcal{H}} E(H_i) = E(G)$ . In other words, the family of graphs  $\mathcal{H}$  covers  $G$ , or partitions the edge set of  $G$ . If such a  $\mathcal{H}$  exists, then splitting  $G$  into  $\{H_1, \dots, H_j\}$  is called a  $\mathcal{H}$ -decomposition of  $G$ . A kind of decomposition studied requires  $\mathcal{H}$  to a single graph (say  $\{H_1\}$ ) and checks if  $G$  can be decomposed into multiple copies of  $H_1$  with the disjoint intersection condition omitted. Such a decomposition is denoted by  $H_1|G$ . The family  $\mathcal{H}$  may consist of paths, cycles, bipartite graphs or matchings. For instance, consider *matching decomposition*, where in an edge-coloring of  $G$ , each color class is a matching. So, coloring edges of  $G$  by  $\chi_e(G)$  colors properly gives the minimum matching decomposition of the graph. Vizing's theorem [13] states that for all simple graphs  $G$ ,  $\chi_e(G) \leq \Delta(G) + 1$ . As a result, there is always a matching decomposition of  $G$  into  $\mathcal{H}$  of size  $|\mathcal{H}| = \Delta(G) + 1$ . A  $tK_2|G$  decomposition is splitting  $G$  into multiple copies of  $t$   $K_2$ 's i.e., matchings of size  $t$ . Bialostocki and Roditty [2] proved that  $3K_2|G$  if and only if  $3||E(G)||$  and  $\Delta(G) \leq \frac{|E(G)|}{3}$ , with a finite number of exceptions. Alon [1] shown that for every  $t > 1$ , if  $|E(G)| \geq \frac{8}{3}t^2 - 2t$ ,  $tK_2|G$  if and only if  $t||E(G)||$  and  $\Delta(G) \leq \frac{|E(G)|}{t}$ . Along similar lines, a significant amount of study has been done and there is vast literature for various kinds of decomposition of graphs (see [4]). In this paper, we aim to combine the concepts of decomposition and coloring graphs and hypergraphs.

## 1.2 Our contribution

We define  $\chi^c(G)$  for a hypergraph  $G(V, E)$  as the minimum number of bicolorings that guarantees every hyperedge  $e \in E$  of  $G$  is properly bicolored in at least one of the  $\chi^c(G)$  bicolorings. In section 2, (i) we derive a tight bound for  $\chi^c(G)$  for the complete  $k$ -uniform hypergraph  $G$ , (ii) establish upper bounds for  $\chi^c(G)$  based on *matchings* and *hitting sets* of the hypergraph, and, (iii) design polynomial time algorithms for computing bicoloring covers. We also relate  $\chi^c(G)$  with independent sets and chromatic numbers and show that  $\chi^c(G) = \lceil \log \chi(G) \rceil$ .

In section 3, we present an inapproximability result about the impossibility of approximating the bicoloring cover of  $n$ -vertex  $k$ -uniform hypergraphs, to within an additive factor of  $(1 - \epsilon) \log n$ , for any fixed  $\epsilon > 0$  in time polynomial in  $n$ . For a  $k$ -uniform hypergraph  $H(V, E)$ , where  $|V| = n$ ,

we show that the bicoloring cover number  $\chi^c(H)$  is  $O(\frac{\log n}{\log \log n - \log \log \log n})$  approximable.

Let  $C = \{C_i \mid C_i \text{ is a bicoloring cover of } \chi^c(G) \text{ bicolorings that cover } G(V, E)\}$ . Let  $|C| = w$ , where  $w \leq 2^{n\chi^c(G)}$ . Each vertex receives a color bit vector due to a bicoloring cover  $C_i$ . Let  $\gamma_i(G)$  be the size of the largest set of vertices that receive the same color bit vector due to bicoloring cover  $C_i$ . Let  $\gamma(G) = \max_{1 \leq i \leq w} \gamma_i(G)$ . We call  $\gamma(G)$  the *cover independence* number of hypergraph  $G$ . In section 4, we show that for any  $k$ -uniform hypergraph  $G$ ,  $\gamma(G) \geq k - 1$ . We relate  $\gamma(G)$  to  $\chi^c(G)$  and  $\chi(G)$  and derive the lower bounds of  $\log \frac{|V|}{\gamma(G)}$  and  $\frac{|V|}{2\gamma(G)}$  for  $\chi^c(G)$  and  $\chi(G)$ , respectively. We also construct a particular class of hypergraphs  $G(V, E)$  called *cover friendly* hypergraphs where the ratio of  $\alpha(G)$  to  $\gamma(G)$  can be made arbitrarily large. More specifically, we construct  $k$ -uniform hypergraphs  $G(V, E)$  where  $\alpha(G) \geq n^{1-t}$ , whereas  $\gamma(G) = n^t$ , for some small fraction  $0 < t < 0.5$ . We show in Corollary 1 that  $\chi^c(G)$  can be approximated for such cover friendly hypergraphs, with an approximation ratio of  $\frac{1}{1-t}$ , by exploiting the special properties of such hypergraphs. However, using Proposition 1, we can only achieve an approximation ratio of at least  $\frac{1}{t}$  for cover friendly hypergraphs. This implies that we achieve an improvement (reduction) in approximation ratio for estimating  $\chi^c(G)$  for cover friendly hypergraphs by a factor of at least  $\frac{1-t}{t}$  by using the properties of  $\gamma(G)$  in Corollary 1; the approximation ratio achieved using Observation 1 in Proposition 1 is much smaller for cover friendly graphs. Furthermore, our constant factor approximation ratio of  $\frac{1}{1-t}$  for approximating  $\chi^c(G)$  for cover friendly hypergraphs is in sharp contrast to our  $O(\frac{\log n}{\log \log n - \log \log \log n})$  factor algorithm for estimating  $\chi^c(G)$  for general hypergraphs as summarized in Theorem 5 of Section 3.

Let  $H(V', E')$  be the largest  $k$ -uniform subhypergraph of a  $k$ -uniform hypergraph  $G(V, E)$ , where  $V' \subseteq V$ ,  $E' \subseteq E$ , and there is a hyperedge for every subset of  $k$  vertices in  $V'$  i.e.,  $E' = \binom{V'}{k}$ . We define  $\omega(G) = |V'|$ . In Section 4.3, we prove that for any  $t \geq 1$ , there exists a  $k$ -uniform hypergraph  $G$  where  $\omega(G) = k$  and  $\chi^c(G) > t$ . Observe that, for  $k = 2$  (usual graphs), this result implies that triangle-free graphs can have arbitrarily large bicoloring cover numbers.

In sections 5 and 6, we correlate  $\chi^c(G)$  to the number  $|E|$  of hyperedges, and the *dependency*  $d(G)$ , using probabilistic analysis, the Moser-Tardos algorithm [11], and an incremental method based on cuts in hypergraphs. We show that if  $|E| \leq 2^{(k-1)x-1}$ , then a bicoloring cover of size  $x$  can be computed in polynomial time. We use  $m(k, x)$  to denote the minimum number of hyperedges such that some  $k$ -uniform hypergraph  $G$  with  $m(k, x)$  hyperedges does not have a bicoloring cover of size  $x$ . We show that  $2^{(k-1)x-1} < m(k, x) \leq x \cdot k^2 \cdot 2^{(k+1)x+2}$ . Let the *dependency*  $d(G)$  of  $G$  be the maximum number of neighboring hyperedges of a hyperedge in  $G$ . We use the Moser-Tardos constructive approach for Lovász local lemma, as in [11], for computing bicoloring covers of size  $x$ , where the dependency  $d(G)$  of the hypergraph is bounded by  $\frac{2^{x(k-1)}}{e} - 1$ .

## 2 Preliminaries

### 2.1 Bicoloring cover number and chromatic number for hypergraphs

We establish the following result relating  $\chi^c(G)$  and  $\chi(G)$  for arbitrary hypergraphs.

**Theorem 1** *Let  $G(V, E)$  be a hypergraph. Let  $\chi^c(G)$  and  $\chi(G)$  be the bicoloring cover number and chromatic number of  $G$ , respectively. Then,  $\chi^c(G) = \lceil \log \chi(G) \rceil$ .*

To show that  $\lceil \log \chi(G) \rceil \leq \chi^c(G)$ , choose a bicoloring cover  $C$  of size  $\chi^c(G)$  for  $G$ . Each vertex  $v$  of  $G$  is assigned a set of  $\chi^c(G)$  colors (bits 0 or 1), by the  $\chi^c(G)$  bicolorings in the bicoloring cover  $C$ . Assign the decimal equivalent of the  $\chi^c(G)$ -bit pattern for  $v$  as the color for  $v$  to get a vertex-coloring  $C'$  for  $G$ . The total number of colors used is at most  $2^{\chi^c(G)}$ . We claim  $C'$  is a proper vertex-coloring for  $G$ , thereby enforcing the inequality  $\chi(G) \leq 2^{\chi^c(G)}$  or  $\lceil \log \chi(G) \rceil \leq \chi^c(G)$ . For the sake of contradiction, assume that some hyperedge  $e \in E(G)$  is monochromatic under  $C'$ . This

means in each of the  $\chi^c(G)$  bicolings, every vertex of  $e$  gets same color. As a result,  $e$  is not covered by the  $\chi^c(G)$  sized cover, which is a contradiction. Consequently, we have the following lemma.

**Lemma 1** For a hypergraph  $G(V, E)$ ,  $\lceil \log \chi(G) \rceil \leq \chi^c(G)$ .

To prove the second inequality, consider a proper coloring  $C$  of the vertices of  $G$  with  $\chi(G)$  colors. Construct the bicoloring cover  $X$  of size  $\lceil \log \chi(G) \rceil$  by assigning the vertices with two colors determined by the 0/1 bits of the color they were assigned under proper coloring  $C$ ; a vertex  $v$  is assigned the  $i$ th bit of the color assigned to it under coloring  $C$  for the 0/1 bicoloring of  $v$  in the  $i$ th bicoloring of the bicoloring cover  $X$ ,  $1 \leq i \leq \lceil \log \chi(G) \rceil$ . Assume for the sake of contradiction that some  $e \in E(G)$  is not covered under bicoloring cover  $X$ . This means every vertex of  $e$  has the same bit vector of length  $\lceil \log \chi(G) \rceil$ , and therefore has the same color under coloring  $C$ , a contradiction. Consequently, we have the following lemma.

**Lemma 2** For a hypergraph  $G(V, E)$ ,  $\chi^c(G) \leq \lceil \log \chi(G) \rceil$ .

Theorem 1 follows from Lemmas 1 and 2. The following lemma is a direct consequence of Theorem 1.

**Lemma 3** Let  $G(V, E)$  be a  $k$ -uniform hypergraph, and  $\{I_1, I_2, \dots, I_u\}$  be a partition of the vertex set  $V$  into independent sets. Then there exists a bicoloring cover for  $G$  of size  $\lceil \log u \rceil$ .

## 2.2 Matchings, hitting sets and bicoloring covers for hypergraphs

Let  $G(V, E)$  be a  $k$ -uniform hypergraph, with  $|V| = n$  and  $E = \{E_1, E_2, \dots, E_m\}$ , where  $E_i \subseteq V$ ,  $1 \leq i \leq m$ . We have the following bounds for  $\chi^c(G)$  based on the sizes of maximal matchings and hitting sets. The first algorithm *MBC* uses a maximal matching for computing a bicoloring cover. The second algorithm *HBC* uses a hitting set.

---

**Algorithm *MBC*:** Computing bicoloring cover using a Maximal matching  $M$

---

**Data:**  $k$ -uniform hypergraph  $G(V, E)$  with  $|V| = n$ , and a maximal matching  $M$  of  $G$

**Result:** Set  $X$  of bicolings of size  $|X| \leq \log_2 |M| + 2$

Color every vertex in the hyperedges of  $M$  with color 1 and rest of the vertices with color 2;  
 $recMBC(M)$ ;

Color the remaining hyperedge of the matching independently using one bicoloring;

---



---

**Function  $recMBC(M)$**

---

**Input:** A set of hyperedges  $M$

**if** ( $|M| > 1$ ) **then**

Split the hyperedges in  $M$  into two sets  $A, B$  of size  $\lfloor \frac{|M|}{2} \rfloor$  and  $\lceil \frac{|M|}{2} \rceil$  respectively;

Color every vertex in  $A$  with color 1 and every vertex in  $B$  with color 2;

$recMBC(A)$ ;

$recMBC(B)$ ;

---

Let  $M$  be a maximal matching of the  $n$ -vertex  $k$ -uniform hypergraph  $G(V, E)$ . We propose an algorithm *MBC* for computing a bicoloring cover of  $G$  using  $M$ . The algorithm *MBC* takes the hypergraph  $G(V, E)$  and a maximal matching  $M$  of  $G$  as inputs and produces a bicoloring cover  $C1$  for  $G$ . Let  $V_M$  denote the set of vertices in the hyperedges in  $M$ . In the first bicoloring, *MBC* colors every vertex of  $V_M$  with color 0, and, all the vertices in  $V \setminus V_M$  with color 1. Due to the maximality of the matching  $M$ , every hyperedge that contains a vertex from  $V \setminus V_M$  shares at least one vertex with some hyperedge in  $M$ . So, every hyperedge  $e \not\subseteq V_M$  is certainly properly

bicolored. The hyperedges which are not properly bicolored in the first bicoloring are subsets of  $V_M$ . Then  $MBC$  calls a recursive function  $recMBC$  with the matching  $M$  as an argument. All the subsequent bicoloredings are performed by  $recMBC$ . The function  $recMBC$  splits the hyperedges in  $M$  into two sets,  $M1$  and  $M2$  with  $|M1| = \lceil \frac{|M|}{2} \rceil$  and  $|M2| = \lfloor \frac{|M|}{2} \rfloor$ .  $MBC$  colors the vertices of every hyperedge in  $M1$  and  $M2$  with colors 0 and 1, respectively. This gives the second bicoloring. Note that every hyperedge of  $G$  that shares at least one vertex each with a hyperedge in  $M1$  and a hyperedge in  $M2$ , is properly bicolored. Now, vertices of hyperedges in  $M1$  and  $M2$  can be colored independently in the subsequent bicoloredings. The function  $recMBC$  is invoked recursively on  $M1$  and  $M2$ , separately. Note that  $recMBC$  terminates when its argument has a single hyperedge; such a hyperedge can be bicolored using a single bicoloring.

We analyze the number of bicoloring generated by the algorithm  $MBC$  as follows. After the first bicoloring, the problem size is  $|M|$  and the problem size gets halved in each subsequent bicoloring step. So, after  $\log |M|$  bicoloredings, the problem reduces to bicoloring of a single hyperedge, which can be done using a single bicoloring. We summarize our result in the following theorem.

**Theorem 2** For any  $k$ -uniform hypergraph  $G(V, E)$ ,  $\chi^c(G) \leq \log |M| + 2$ , where  $M$  is a maximal matching of  $G$ . Algorithm  $MBC$  computes such a bicoloring cover for  $G(V, E)$  in  $O(n \log |M|)$  time.

---

**Algorithm  $HBC$ :** Computing bicoloring cover using a hitting set  $H$

---

**Data:**  $k$ -uniform hypergraph  $G(V, E)$  with  $|V| = n$  and a hitting set  $H$

**Output:** Set  $X$  of bicoloredings of size  $|X| = \log \lceil \frac{|H|}{k-1} \rceil + 1$

Color every vertex in  $H$  with color 1 and the rest of the vertices with color 2;

Let  $G'(H, E')$  be a hypergraph defined on the vertices of  $H$ , and  $E'$  be all the hyperedges that are monochromatic after the first bicoloring;

$KnCover(G')$ ;

---

Let  $H$  be a hitting set of the hypergraph  $G(V, E)$ . We propose an algorithm  $HBC$  for computing a bicoloring cover of  $G$  using  $H$ .  $HBC$  takes the hypergraph  $G(V, E)$  and the hitting set  $H$  as inputs and produces a bicoloring cover  $C1$ . In the first bicoloring,  $HBC$  colors every vertex in  $H$  with color 0 and all the remaining vertices with the color 1. So, the hyperedges which are monochromatic in the first coloring are subsets of  $H$ . Let  $G'(H, E')$  be a hypergraph on the vertices of  $H$ ,  $E'$  be all the hyperedges that are monochromatic after the first bicoloring.  $HBC$  invokes algorithm  $KnCover$  on hypergraph  $G'(H, E')$  to properly bicolor the hyperedges of  $G'$ . By Corollary ??, we know that  $KnCover$  computes a bicoloring cover for  $G'$  consisting of  $\lceil \log \frac{|H|}{k-1} \rceil$  bicoloredings. These  $\lceil \log \frac{|H|}{k-1} \rceil$  bicoloredings combined with the first bicoloring gives the desired bicoloring cover for  $G$ . So, we get the following theorem.

**Theorem 3** For any  $k$ -uniform hypergraph  $G(V, E)$ ,  $\chi^c(G) \leq \lceil \log \frac{|H|}{k-1} \rceil + 1$ , where  $H$  is a hitting set of  $G$ . Algorithm  $HBC$  computes such a bicoloring cover for  $G(V, E)$  in  $O(n \log \frac{|H|}{k-1})$  time.

As the union of vertices of some maximal matching  $M$  gives a hitting set, replacing  $|H|$  by  $|M|k$ , yields the same bound as in Theorem 2. As the effectiveness of the algorithm followed in proof of Theorem 2 depends on the size of the maximal matching, finding the smallest maximal matching is useful.

### 3 Approximating bicoloring covers

Lovász [9] showed that the decision problem of bicolorability of hypergraphs is NP-complete. Feige and Killian [6] showed that if NP does not have efficient randomized algorithms i.e.,  $NP \not\subseteq ZPP$ ,

then there is no polynomial time algorithm for approximating the chromatic number of an  $n$ -vertex graph within a factor of  $n^{1-\epsilon}$ , for any fixed  $\epsilon > 0$ . Using the above result, Krivelevich [8] demonstrated that for any fixed  $k \geq 3$ , it is impossible to approximate the chromatic number of  $k$ -uniform graphs on  $n$  vertices within a factor of  $n^{1-\epsilon}$  for any fixed  $\epsilon > 0$ , in time polynomial in  $n$ . In Section 3.1, we show that it is impossible to approximate the bicoloring cover of  $k$ -uniform hypergraphs on  $n$  vertices within an additive factor of  $(1 - \epsilon) \log n$  for any fixed  $\epsilon > 0$ , in time polynomial in  $n$ . We also design approximation algorithms for computing bicoloring covers in Section 3.2 using the methods developed in [8].

### 3.1 Inapproximability of the computation of $\chi^c(G)$

Let  $G(V, E)$  be a  $k$ -uniform hypergraph, and  $\chi^c(G)$  and  $\chi(G)$  be the bicoloring cover number and the chromatic number of  $G$ , respectively. Assume that  $G$  has a bicoloring cover of size  $x$  i.e.,  $\chi^c(G) \leq x$ . By Theorem 1,  $\chi(G) \leq 2^x$ . Let  $R$  be an algorithm that computes a bicoloring cover of size  $x$  for graph  $G$ . Suppose  $R$  is a  $\alpha$ -additive approximation algorithm i.e., for any input instance  $G$ , the size of the computed bicoloring cover  $x \leq \chi^c(G) + \alpha$ . Then, using  $R$  we can design an approximation algorithm for proper coloring of  $G$  using  $2^x \leq 2^{\chi^c(G) + \alpha} = 2^{\chi^c(G)} 2^\alpha < \chi(G) 2^{\alpha+1}$  colors. However, in [8], it is established that no polynomial time algorithm can approximate  $\chi(G)$  within a factor of  $n^{1-\epsilon}$ , for any fixed  $\epsilon > 0$ . So, setting  $2^{\alpha+1} = n^{1-\epsilon}$ , we get  $\alpha = (1-\epsilon) \log n - 1$ . Therefore, we have the following theorem.

**Theorem 4** *Under the assumption that  $NP \not\subseteq ZPP$ , no polynomial time algorithm can approximate the bicoloring cover number  $\chi^c(G)$  for  $n$ -vertex  $k$ -uniform hypergraph  $G(V, E)$  within an additive approximation factor of  $(1 - \epsilon) \log n - 1$ , for any fixed  $\epsilon > 0$ .*

### 3.2 An approximation algorithm for computing bicoloring covers

Krivelevich and Sudakov [8] have developed an algorithm  $D(G, p)$  that takes a  $n$ -vertex  $k$ -uniform hypergraph  $G(V, E)$ , and a integer  $p \geq \chi(G)$  as inputs, and computes a proper coloring of the hypergraph  $G$ . The algorithm  $D(G, p)$  uses two algorithms  $C_1$  and  $C_2$  that properly color the hypergraph  $G$  using at most  $8n^{1 - \frac{1}{(k-1)(p-1)+1}}$  and  $\frac{2n \log p}{\log n}$  colors, respectively if  $p \geq \chi(G)$ . In order words,  $D(G, p)$  succeeds in computing an approximate proper coloring if  $p \geq \chi(G)$ . Since the actual value of chromatic number is not known a priori,  $D(G, p)$  is executed with all possible integral values of  $p$  in the range 1 through  $|V|$ . So, the approximation ratio for  $\chi(G)$  using  $D(G, p)$  is  $\min\{\frac{8n^{1 - \frac{1}{(k-1)(p-1)+1}}}{p}, \frac{2n \log p}{p}\}$ . Krivelevich and Sudakov use a value of  $p = \frac{1}{(k-1)} \frac{\log n}{\log \log n}$  so that both the terms in the minimization are of the same order, achieving the approximation ratio of  $O(\frac{n(\log \log n)^2}{(\log n)^2})$ .

In order to compute a bicoloring cover where the number of bicolorings is within a good approximation factor with respect to  $\chi^c(G)$ , we use a similar idea and the algorithms of Krivelevich and Sudakov. From Lemma 1, we know that  $\chi(G) \leq 2^{\chi^c(G)}$ . Suppose we invoke  $D(G, p)$ , where  $p = 2^s$  and  $s \geq \chi^c(G)$ . Then, the algorithms  $C_1$  and  $C_2$  properly color the hypergraph  $G$  using at most  $8n^{1 - \frac{1}{(k-1)(2^s-1)+1}}$  and  $\frac{2ns}{\log n}$  colors, respectively. However, we do not know the value of  $\chi^c(G)$  to begin with. As we know that  $\chi^c(G) \leq \lceil \log \frac{n}{k-1} \rceil$  (see Theorem ??), we run  $D(G, 2^s)$  with all possible values of  $s$  in the range 1 through  $\lceil \log \frac{n}{k-1} \rceil$  and choose the minimum value of  $s$  for which  $D(G, 2^s)$  outputs a proper coloring. From this proper coloring, we can compute a bicoloring cover using the reduction stated in the proof of Lemma 2. Let  $C_{12}$  be the algorithm that (i) takes a  $k$ -uniform hypergraph  $G(V, E)$  and an integer  $s$  as inputs, (ii) runs  $D(G, 2^s)$  for different values of  $s$ , and (iii) computes a bicoloring cover from the proper coloring output of  $D(G, 2^s)$ . From Lemma 2, it is clear that  $C_{12}$  produces a bicoloring cover of size  $\min\left(\log(8n^{1 - \frac{1}{(k-1)(2^s-1)+1}}), \log(\frac{2ns}{\log n})\right)$ . So, the approximation ratio for algorithm  $C_{12}$  is at

most  $\min \left( \frac{\log(8n^{1-\frac{1}{(k-1)(2^s-1)+1}})}{s}, \frac{\log(\frac{2ns}{\log n})}{s} \right)$ . We choose the value of  $s$  that makes both the terms of the same order. Setting  $s = \log(\frac{1}{k-1} \frac{\log n}{\log \log n})$ , the first term becomes  $\log(8n^{1-\frac{1}{(k-1)(2^s-1)+1}}) = \log(8nn^{-\frac{\log \log n}{\log n - (k-2)\log \log n}}) \leq \log(8nn^{-\frac{\log \log n}{\log n}}) = \log(8nn^{-\log_n \log n}) = O(\log(\frac{n \log \log n}{\log n}))$ . The second term becomes  $\log(\frac{2ns}{\log n}) = O(\log(\frac{n \log \log n}{\log n}))$ . Therefore,  $C_{12}$  has an approximation ratio of  $O(\frac{\log n + \log \log \log n - \log \log n}{\log \log n - \log \log \log n}) = O(\frac{\log n}{\log \log n - \log \log \log n} - 1)$ . We have the following theorem.

**Theorem 5** *For any  $n$  vertex  $k$ -uniform hypergraph  $G(V, E)$ , the bicoloring cover number  $\chi^c(G)$  is  $O(\frac{\log n}{\log \log n - \log \log \log n})$  approximable.*

## 4 Lower bounds for the bicoloring cover number

In this section we study the relationship between the bicoloring cover number, independent sets and related concepts. Throughout this section, all the approximation ratios are determined with respect to Algorithm *HBC*. In Section 4.1, we develop certain relationships between the bicoloring cover number  $\chi^c(G)$ , the chromatic number  $\chi(G)$ , and introduce a new parameter  $\gamma(G)$ , which we call the *cover independence number*. We demonstrate a better approximation ratio for  $\chi^c(G)$  for hypergraphs where there is large separation between  $\alpha(G)$  (the *independence number*) and  $\gamma(G)$ . In Section 4.2, we demonstrate examples of  $k$ -uniform hypergraphs  $G(V, E)$ , where there is a large separation between  $\gamma(G)$  and  $\alpha(G)$ . In Section 4.3, using a probabilistic argument we demonstrate the existence of hypergraphs with an arbitrarily large gap between  $\omega(G)$  and  $\chi^c(G)$ . This shows that the lower bound for  $\chi^c(G)$  as in Theorem ?? for arbitrary  $k$ -uniform hypergraphs, is not tight.

### 4.1 Independence number, cover independence number and the bicoloring cover number

A set  $I$  of vertices of any hypergraph  $G(V, E)$  is called an *independent set* if there is no hyperedge of  $G$  in  $I$  i.e., for no hyperedge  $e \in E(G)$ ,  $e \subseteq I$ . The maximum size of any independent set is called the *independence number*  $\alpha(G)$ . Note that  $\chi(G) \geq \frac{|V|}{\alpha}$  ([10]). Combined with Lemma 1, we have the following observations.

**Observation 1** *For a  $k$ -uniform hypergraph  $G$ ,  $\chi^c(G) \geq \log \lceil \frac{|V(G)|}{\alpha(G)} \rceil$ .*

**Proposition 1** *For a  $k$ -uniform hypergraph  $G$ ,  $\chi^c(G)$  can be approximated in polynomial time by a ratio factor  $\frac{1}{1-t}$  algorithm if  $\alpha(G) = n^t$ , where  $t < 1$ .*

**Proof.** Algorithm *HBC* computes a bicoloring cover of size  $\lceil \log \frac{|H|}{k-1} \rceil + 1$  in polynomial time, where  $H$  is a hitting set for  $G$  (see Theorem 3). Following Observation 1, we note that the approximation ratio is at most  $\frac{\log |H| - \log(k-1) + 1}{\log |V(G)| - \log \alpha(G)}$ , which is at most  $\frac{1}{1-t}$  if  $\alpha(G) = n^t$  and  $t < 1$ .  $\square$

From Observation 1 we note that the bicoloring cover number  $\chi^c(G)$  is lower bounded by  $\log \frac{|V|}{\alpha(G)}$ . We introduce the notion of *cover independence* in Section 4.1.1, and show in Theorem 7 of Section 4.1.4 that  $\chi^c(G)$  is lower bounded by  $\log \frac{|V|}{\gamma(G)}$ , where  $\gamma(G)$  is the *cover independence number*. Further, in Section 4.2 we construct hypergraphs called *cover friendly* hypergraphs, where the values of  $\alpha(G)$  and  $\gamma(G)$  are widely separated. Observe that Theorem 7 yields a better lower bound for  $\chi^c(G)$  than that given by Observation 1.

#### 4.1.1 The notion of cover independence

There can be multiple sets of bicolorings of size  $\chi^c(G)$  that cover  $G$ . Let  $w$  be the number of distinct (labeled) bicoloring covers of size  $\chi^c(G)$ , where  $w \leq 2^{n\chi^c}$ . Let the set  $C = \{C_1, \dots, C_w\}$  be the set of all the bicoloring covers of size  $\chi^c(G)$  i.e., the set of all the optimal bicoloring covers for  $G$ .

Let  $C_i = \{X_1^i, \dots, X_{\chi^c(G)}^i\}$ ,  $1 \leq i \leq w$ , be a bicoloring cover of size  $\chi^c(G)$ , where  $X_j^i$  denotes the  $j^{th}$  bicoloring of vertices of  $G$  in the  $i^{th}$  bicoloring cover  $C_i$ . The  $\chi^c(G)$  bicolorings in  $C_i$  define a *color bit vector*  $B_v^i$  of  $\chi^c(G)$  bits for each vertex  $v \in V$ , where the  $j^{th}$  bit of  $B_v^i$  corresponds to the color of  $v$  in the  $j^{th}$  bicoloring  $X_j^i$  in  $C_i$ . Consider the partition  $\mathcal{P}_i = \{V_1^i, V_2^i, \dots, V_q^i\}$  of the vertex set  $V$  of  $G(V, E)$  such that vertices  $u$  and  $v$  belong to the same part, say  $V_k^i$ , if and only if  $B_u^i$  is identical to  $B_v^i$ . The partition  $\mathcal{P}_i$  is called a *canonical partition* of  $V$  due to the optimal bicoloring cover  $C_i$ . Note that  $q$  is the number of distinct color bit vectors, determining the number of parts in the above partition. Let  $\gamma_i(G)$  be the size of the largest set of vertices that receive the same color bit vector for the bicoloring cover  $C_i$ . We define

$$\gamma_i(G) = \max_{1 \leq k \leq q} |V_k^i|,$$

We also define

$$\gamma(G) = \max_{1 \leq i \leq w} \gamma_i(G),$$

We call  $\gamma(G)$  the *cover independence number* of the  $k$ -uniform hypergraph  $G(V, E)$ . Any optimal bicoloring cover  $C_i$  of the hypergraph  $G(V, E)$  with  $\gamma_i(G) = \gamma(G)$  is called a *witness* for  $G$ .

#### 4.1.2 Interpreting the parameter $\gamma(G)$ with examples

We know that there is a unique bipartition for a connected bipartite graph  $G(V, E)$ , where each edge of  $G$  has one vertex in each part. This unique bipartition yields a bicoloring that covers all edges of  $G$ . The value of  $\gamma(G)$  for such graphs is the size of the larger part in this bipartition.




Figure 1: (a)  $BC_1$ ,  $BC_2$  and  $BC_3$  denote three distinct bicoloring covers of size 2 for  $K_4$  where  $\gamma(K_4) = 1$ . (b) Bicoloring cover  $C = \{X_1, X_2\}$  for  $C_7$ , where  $X_1 = \text{red}\{1, 3, 5, 7\}, \text{blue}\{2, 4, 6\}$ ,  $X_2 = \text{red}\{1, 3, 5\}, \text{blue}\{2, 4, 6, 7\}$  and  $\gamma(C_7) \geq |\{1, 3, 5\}| = 3$ .

Note that  $\gamma(G)$  for a complete graph is 1. Observe that in a complete graph there is an edge between each pair of vertices. So, a bicoloring cover  $C$  must color vertices in every pair of vertices with different colors in at least one bicoloring  $C_i$  of the cover  $C$ . For instance, consider the bicoloring covers of  $K_4$ . From Figure 1, it is clear that the size of the largest set of vertices colored with same color in both the bicolorings is 1, in all the three bicoloring covers  $BC_1$ ,  $BC_2$  and  $BC_3$  i.e.,  $\gamma_1(K_4) = \gamma_2(K_4) = \gamma_3(K_4) = 1$ . Therefore,  $\gamma(K_4) \geq 1$ . To see that  $\gamma(K_4) < 2$ , observe that if any pair of vertices (say vertices 1 and 2), are colored with the same color in both the bicolorings, then the edge  $\{1, 2\}$  remains uncovered by the set of bicolorings.

For an odd cycle  $C_n$ ,  $V = \{v_1, \dots, v_n\}$ ,  $\gamma(C_n)$  is  $\frac{n-1}{2}$  and  $\chi^c(C_n) = 2$ . Since an odd cycle is not bicolorable,  $\chi^c(C_n) \geq 2$ . To show that  $\chi^c(C_n) = 2$ , we consider a bicoloring  $X_1$ , where every odd vertex is colored 0, every even vertex is colored 1. The only edge that is not properly colored is  $v_1, v_n$ . A second bicoloring  $X_2$ , which is exactly the same as  $X_1$  except that the color



for  $v_n$  is 1 in  $X_2$ . Note that  $X_2$  properly colors  $\{v_1, v_n\}$ . So,  $C = \{X_1, X_2\}$  is a bicoloring cover for  $C_n$ . So,  $\chi^c(C_n) = 2$ . The vertices  $\{v_1, v_3, \dots, v_{n-2}\}$  are colored with 0 in both  $X_1$  and  $X_2$  in  $C$ . Consequently,  $\gamma(C_n) \geq \frac{n-1}{2}$  (see Fig. 1). Also, observe that any subset of vertices from the odd cycle  $C_n$  with greater than  $\frac{n-1}{2}$  vertices must contain two consecutive vertices. Therefore,  $\gamma(C_n) \leq \frac{n-1}{2}$ .

For any bicolorable hypergraph,  $\chi^c = 1$ , set  $C$  consists of all the proper bicolorings of vertices,  $\gamma_i$  is the size of the larger of the two color classes of  $i^{th}$  proper bicoloring.  $\gamma \geq \gamma_i \geq \frac{n}{2}$ . For example, consider the bicoloring of  $H(V, E)$ , where  $V = \{1, 2, 3, 4, 5\}$ , and  $E$  consists of all the 3-uniform hyperedges except  $\{1, 2, 3\}$  and  $\{1, 2, 4\}$  (see Fig. 2). Certainly  $H$  is bicolorable with bicolorings  $X_1$  and  $X_2$ :  $X_1 = \text{red}\{1, 2, 3\}, \text{blue}\{4, 5\}$ ,  $X_2 = \text{red}\{1, 2, 4\}, \text{blue}\{3, 5\}$ .  $\gamma_1 = \gamma_2 = 3$ . Coloring any four vertices with same color in a bicoloring does not cover all the hyperedges. Hence  $\gamma = 3 \geq \frac{5}{2}$ .




Figure 2: Two bicolorings of  $V = \{1, 2, 3, 4, 5\}$ :  $X_1 = \text{red}\{1, 2, 3\}, \text{blue}\{4, 5\}$ ,  $X_2 = \text{red}\{1, 2, 4\}, \text{blue}\{3, 5\}$ .  $C = \{X_1, X_2\}$  is a bicoloring cover of  $K_5^3$ .  $\gamma(K_5^3) \geq |\{1, 2\}| = 2$ .  $C_1 = \{X_1\}$  and  $C_2 = \{X_2\}$  are two distinct bicoloring covers for  $H = K_5^3 \setminus \{\{1, 2, 3\}, \{1, 2, 4\}\}$ .  $\gamma_1 \geq |\{1, 2, 3\}| = 3$ .  $\gamma_2 \geq |\{1, 2, 4\}| = 3$ .  $\gamma(H) \geq \max(\gamma_1, \gamma_2) = 3$ .

#### 4.1.3 A preliminary lower bound for $\gamma(G)$

For arbitrary  $k$ -uniform hypergraph  $G$ ,  $\gamma(G) \geq k - 1$ . Assume for the sake of contradiction that  $\gamma(G) = l$  for some  $1 \leq l \leq k - 2$ . This implies there exists an optimal bicoloring cover  $C_i$  such that  $\gamma_i(G) = l$ , and there does not exist any optimal bicoloring cover  $C_j$  with  $\gamma_j(G) = l + 1$ . We arrive at a contradiction by showing that there exists an optimal bicoloring cover  $C_j$  with  $\gamma_j(G) = l + 1$ . Consider  $C_i$ , a witness for  $G$ , and let  $V'$  be the set of vertices such that  $|V'| = l = \gamma_i(G) = \gamma(G)$ , where all the vertices in  $V'$  receive identical color bit vectors for the bicoloring cover  $C_i$ . Let us move a vertex  $s$  from  $V \setminus V'$  to  $V'$  by assigning the color bit vector of  $V'$  to  $s$ , thereby obtaining another set of bicolorings  $C_j$  of the same size as  $C_i$ . Let  $V'_{new} = V' \cup \{s\}$ . If we can show that  $C_j$  is a bicoloring cover for the hypergraph  $G$ , then it follows that  $\gamma_j(G) = |V'_{new}| = l + 1$  and we are done. Any hyperedge that does not contain any vertex from  $V'_{new}$  is covered by  $C_j$  (using the same bicolorings as in  $C_i$ ). Note that since  $|V'_{new}| \leq k - 1$ , any hyperedge  $e \in E$  that includes vertices from  $V'_{new}$ , must contain at least one vertex  $t \in V \setminus V'_{new}$ . Since the color bit vectors of  $t$  and  $V'_{new}$  are different for  $C_j$ ,  $e$  is properly bicolored by some bicoloring in the set  $C_j$ . Since these exhaustive cases include every hyperedge in  $G$ ,  $C_j$  is a bicoloring cover for  $G$ , and  $\gamma_j(G) = l + 1$ , a contradiction. Therefore, we have the following theorem.

**Theorem 6** For any  $k$ -uniform hypergraph  $G$ ,  $\gamma(G) \geq k - 1$ .

#### 4.1.4 A lower bound for $\chi^c(G)$ and $\chi(G)$ based on $\gamma(G)$

We now study the significance of the cover independence number  $\gamma(G)$  and its relationship with the bicoloring cover number  $\chi^c(G)$  and chromatic number  $\chi(G)$  of a  $k$ -uniform hypergraph  $G(V, E)$ . Observe that  $\gamma(G)$  is the maximum cardinality of a subset  $S_i \subseteq V$ , of vertices of  $G(V, E)$ , where

the color bit vector for vertices in  $S_i$  remains invariant in the  $i$ th optimal bicoloring cover. Consequently,  $\gamma(G)$  can be used to lower bound the number of such subsets of vertices of  $G$ , where each such subset of vertices is represented by its own color bit vector, as shown in the analysis below.

Consider any optimal bicoloring cover  $\mathcal{C}$  of a  $k$ -uniform hypergraph  $G$ . The bicoloring cover  $\mathcal{C}$  splits  $V(G)$  into a canonical partition  $\mathcal{P}$  of at least  $\lceil \frac{|V(G)|}{\gamma(G)} \rceil$  subsets  $V_1, \dots, V_{\lceil \frac{|V(G)|}{\gamma(G)} \rceil}, \dots, V_q$  of  $G$ , where each vertex in any such set  $S$  of size at most  $\gamma(G)$  receives the color bit vector corresponding to the set  $S$  for the bicoloring cover  $\mathcal{C}$ . If for any  $i, j, i < j$ , there is no hyperedge that shares at least one vertex each with  $V_i$  and  $V_j$  in  $G$ , then we merge  $V_j$  into  $V_i$ . We repeat this process for every  $i, j, i < j$ , till there is at least one hyperedge that shares at least one vertex each with  $V_i$  and  $V_j$ . Let this new bicoloring cover be  $\mathcal{C}1$  and let  $\mathcal{P}1 = \{V_1, \dots, V_p\}$  be the new canonical partition of the vertices of  $G$  due to  $\mathcal{C}1$ , where  $p \leq q$  denotes the number of sets in the canonical partition  $\mathcal{P}1$  of  $V$  due to  $\mathcal{C}1$ . For any  $i, j, 1 \leq i < j \leq p$ ,  $V_i$  and  $V_j$  are now assigned distinct color bit vectors. Let  $\mathcal{C}1 = \{X_1, \dots, X_{\chi^c(G)}\}$ , where  $X_i$  denote the  $i^{th}$  bicoloring in  $\mathcal{C}1$ . By the definition of  $\gamma(G)$ ,  $|V_i| \leq \gamma(G), 1 \leq i \leq p$ , so  $p \geq \lceil \frac{|V(G)|}{\gamma(G)} \rceil$ .

The canonical partition  $\mathcal{P}1$  due to  $\mathcal{C}1$  can be naturally mapped to a complete graph  $H$  with (i) the vertex set  $V(H) = \{1, 2, \dots, p\}$ , where  $p \geq \lceil \frac{|V(G)|}{\gamma(G)} \rceil$ , and the vertex  $i$  corresponds to the part  $V_i$  of the canonical partition  $\mathcal{P}1$ , and (ii) the set  $E(H)$  of edges  $\{i, j\}$ , denoting the existence of a hyperedge  $e$  of  $G$  that shares at least one vertex each with the corresponding sets  $V_i$  and  $V_j$  in the canonical partition  $\mathcal{P}1$ .

**Proposition 2**  $H$  is a complete graph.

**Proof.** According to the definition of  $\mathcal{C}1$ , for every  $i, j, 1 \leq i < j \leq p$ , there is at least one hyperedge that shares at least one vertex each with parts  $V_i$  and  $V_j$  of the canonical partition  $\mathcal{P}1$ . So, for every  $i, j, 1 \leq i < j \leq p$ , there is an edge between vertices  $i$  and  $j$  in  $H$ . So, the proposition holds.  $\square$

Since  $H$  is a complete graph,  $\omega(H) = p \geq \lceil \frac{|V(G)|}{\gamma(G)} \rceil$ . Using Corollary ??, we conclude the following lemma.

**Lemma 4**  $\chi^c(H) \geq \log \lceil \frac{|V(G)|}{\gamma(G)} \rceil$ .

**Lemma 5**  $\chi^c(H) \leq \chi^c(G)$ .

**Proof.** We show that the bicoloring cover  $\mathcal{C}1$  for  $G$  can be modified into a bicoloring cover  $\mathcal{C}1'$  for  $H$ . We construct  $\mathcal{C}1'$  in the following manner. For each  $X_l \in \mathcal{C}1$ , we include a bicoloring  $X'_l$  in  $\mathcal{C}1'$ . We assign the color of vertices of  $V_i$  in  $X_l$  to the vertex  $i$  in  $X'_l$ . In this construction,  $\mathcal{C}1' = \{X'_1, \dots, X'_{\chi^c(G)}\}$ , and  $|\mathcal{C}1'| = \chi^c(G)$ . We need to show that  $\mathcal{C}1'$  is a valid bicoloring cover for  $H$ . Let  $e' = \{i, j\} \in E(H)$ . This implies that there exists a hyperedge  $e$  that shares at least one vertex each with  $V_i$  and  $V_j$ . Suppose  $e$  is covered in bicoloring  $X_l$  of  $\mathcal{C}1$ . This implies that  $V_i$  and  $V_j$  are assigned different colors in  $X_l$ . So, by the construction of  $H$ , vertices  $i$  and  $j$  are colored with different colors in  $X'_l$ , thereby covering  $e'$ . So,  $\mathcal{C}1'$  is a valid bicoloring cover for  $H$  and  $\chi^c(H) \leq |\mathcal{C}1'| = \chi^c(G)$ .  $\square$

Using Lemma 4 and Lemma 5, we have the following theorem.

**Theorem 7** For a  $k$ -uniform hypergraph  $G$ ,  $\chi^c(G) \geq \log \lceil \frac{|V(G)|}{\gamma(G)} \rceil$ .

**Corollary 1** For a  $k$ -uniform hypergraph  $G$ ,  $\chi^c(G)$  can be approximated in polynomial time by a ratio factor  $\frac{1}{1-t}$  algorithm if  $\gamma(G) = n^t$ , where  $t < 1$ .

**Proof.** Algorithm  $HBC$  computes a bicoloring cover of size  $\lceil \log \frac{|H|}{k-1} \rceil + 1$  in polynomial time for  $G$ , where  $H$  is a hitting set of  $G$  (see Theorem 3). Following Theorem 7, we observe that the achieved approximation ratio is at most  $\frac{\log |A| - \log(k-1) + 1}{\log |V(G)| - \log \gamma(G)}$ , which is at most  $\frac{1}{1-t}$  if  $\gamma(G) = n^t$  and  $t < 1$ .  $\square$

Further, we establish the following lower bound for  $\chi(G)$  based on Lemma 2 and Theorem 7.




Figure 3: Example of a hypergraph  $G1(V, E)$  with  $\alpha(G1) > \gamma(G1)$ .  $V = \mathcal{V}_1 \cup \mathcal{V}_2$ , where  $\mathcal{V}_1 = \{1, 2, \dots, 8\}$  and  $\mathcal{V}_2 = \{9, 10, 11, 12\}$ .  $V_0, V_1, V_2$  and  $V_3$  denote the parts with color bit vectors  $(0,0)$ ,  $(0,1)$ ,  $(1,0)$  and  $(1,1)$  respectively.

**Theorem 8** For a  $k$ -uniform hypergraph  $G$ ,  $\chi(G) \geq \frac{|V(G)|}{2\gamma(G)}$ .

**Proof.** From Lemma 2,  $\chi^c(G) \leq \lceil \log \chi(G) \rceil \leq \log \chi(G) + 1$ . Therefore,  $2^{\chi^c(G)} \leq 2 \cdot \chi(G)$ . From Theorem 7,  $2 \cdot \chi(G) \geq 2^{\chi^c(G)} \geq \frac{|V(G)|}{\gamma(G)}$  and the theorem follows.  $\square$

The following proposition establishes the fact that  $\alpha(G)$  is at least as large as  $\gamma(G)$ .

**Proposition 3** For an arbitrary  $k$ -uniform hypergraph  $G(V, E)$ ,  $\alpha(G) \geq \gamma(G)$ .

**Proof.** We prove the proposition by contradiction. Assume that  $\gamma(G) > \alpha(G)$ . Let  $\mathcal{C}$  be one of the bicoloring covers of size  $\chi^c(G)$  that produces a subset of vertices  $\mathcal{V}$  of cardinality  $\gamma(G)$ , such that every vertex in the subset receives the same color in each of the  $\chi^c(G)$  bicolorings of  $\mathcal{C}$ . From our assumption,  $\gamma(G) > \alpha(G)$ , so there must be at least one hyperedge  $e \in E$  such that  $e \subseteq \mathcal{V}$ . From the definition of  $\mathcal{V}$ , it is clear that  $e$  remains monochromatic in all of the  $\chi^c(G)$  bicolorings:  $\mathcal{C}$  cannot be a bicoloring cover of size  $\chi^c(G)$ . This concludes the proof of Proposition 3.  $\square$

The gap between  $\alpha(G)$  and  $\gamma(G)$  becomes a question of great importance for comparing the lower bounds of  $\chi^c(G)$  by Proposition 1 and Corollary 1. In the following section, we generate an example where  $\gamma(G)$  is strictly less than  $\alpha(G)$ , and also generalize the example to construct a class of hypergraphs where  $\alpha(G) \gg \gamma(G)$ .

## 4.2 Construction of hypergraphs with a large gap between $\alpha(G)$ and $\gamma(G)$

### 4.2.1 A small hypergraph demonstrating the separation between $\alpha(G)$ and $\gamma(G)$

We need to show that there exists hypergraphs where there is an arbitrary gap between  $\alpha(G)$  and  $\gamma(G)$ . Consider the 3-uniform hypergraph  $G1(V, E)$ , where  $V = \mathcal{V}_1 \cup \mathcal{V}_2$ ,  $\mathcal{V}_1 = \{1, 2, \dots, 8\}$  and  $\mathcal{V}_2 = \{9, 10, 11, 12\}$  (see Figure 3). The set of hyperedges  $E = E_1 \cup E_2$  is as follows:

- $E_1 = \{\{u, v, w\} | 1 \leq u < v < w \leq 8\}$ , and,
- $E_2 = \{\{u, v, w\} | 1 \leq u < v \leq 8, 9 \leq w \leq 12\} \setminus \{\{1, 5, 9\}, \{2, 6, 10\}, \{3, 7, 11\}, \{4, 8, 12\}\}$ .

Observe that  $\mathcal{V}_2$  is an independent set as it contains no hyperedges. Also, observe that  $G1$  is not bicolorable since it contains a  $K_8^3$  as a subhypergraph (due to hyperedges in  $E_1$ ) and from Theorem ??,

$$\chi^c(G1) \geq \lceil \log \frac{8}{3} \rceil = 2. \quad (1)$$

**Lemma 6** The independence number  $\alpha(G1)$  is five for the hypergraph  $G1$ . Moreover, independent sets of size greater than three for  $G1$  are obtained by adding at most one vertex from  $\mathcal{V}_1$  to subsets of  $\mathcal{V}_2$ .

**Proof.** We prove the above lemma by showing that the maximum sized independent sets of  $G1$  are  $\{\{i, 9, 10, 11, 12\} | i \in \mathcal{V}_1\}$ . Observe that any maximum independent set can contain at most two vertices from  $\mathcal{V}_1$ , since any three vertices in  $\mathcal{V}_1$  introduces a hyperedge in  $G1$ . Suppose  $u$  and  $v$  be any two vertices from  $\mathcal{V}_1$  that are in some maximal independent set,  $u < v$ . If  $u \neq v - 4$ , then we cannot add any vertex to that independent set. If  $u = v - 4$ , then we can add only one vertex  $v + 4$  to that independent set. Such independent sets are of size 3. However, restricting only one vertex  $u$  from  $\mathcal{V}_1$  in the independent set, we can add all the vertices of  $\mathcal{V}_2$ , generating the independent set  $\{u, 9, 10, 11, 12\}$ .  $\square$

In what follows we show that  $\chi^c(G1) \leq 2$ , which combined with Inequality 1 gives  $\chi^c(G1) = 2$ . Consider the bicolorings of vertices:

- $X_1 = \{0, 0, 1, 1, 0, 0, 1, 1, 0, 0, 1, 1\}$ .
- $X_2 = \{0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1\}$ , where  $j^{th}$  entry in  $X_i$  denote the color of the vertex  $j$  in  $i^{th}$  bicoloring,  $1 \leq j \leq 12$ .

Each vertex  $j$ , receives a color bit vector  $(b_{j,2}, b_{j,1})$ , where  $b_{j,2}$  and  $b_{j,1}$  denote the color of  $j$  in  $X_2$  and  $X_1$ , respectively. Let  $C1 = \{X_2, X_1\}$  be a set of bicolorings of  $V$ . Split  $V$  into partition  $P = \{V_0, V_1, V_2, V_3\}$  such that vertex  $j$  is added to part  $V_0, V_1, V_2$  or  $V_3$  if  $j$  receives bits  $(0, 0), (0, 1), (1, 0), (1, 1)$ , respectively. So  $V_0 = \{1, 5, 9\}$ ,  $V_1 = \{2, 6, 10\}$ ,  $V_2 = \{3, 7, 11\}$ ,  $V_3 = \{4, 8, 12\}$ . Note that each  $V_i$ ,  $0 \leq i \leq 3$ , is an independent set, and has a distinct color bit vector associated with it due to the bicoloring cover  $C1$ . The construction of  $G1$  guarantees that every hyperedge of  $G1$  consists of vertices from at least two of the parts.

We wish to show that  $\gamma(G1) = 3 < \alpha(G) = 5$ . For this purpose, we first show that  $C1$  is indeed an optimal bicoloring cover and use  $C1$  to show that  $\gamma(G1) \geq 3$ . Consider a hyperedge  $e = \{u, v, w\} \in E$ . Without loss of generality, we assume that  $u$  and  $v$  lie in different parts in  $P$ . Let the color bit vector of  $u$  ( $v$ ) be  $(a_2, a_1)((b_2, b_1))$ . By definition of the parts, either (i)  $a_1 \neq b_1$ , or (ii)  $a_1 = b_1$  and  $a_2 \neq b_2$ . If  $a_1 \neq b_1$ ,  $e$  is covered by  $X_1$ ; otherwise,  $e$  is covered by  $X_2$ . So,  $C1$  is a valid bicoloring cover for  $G1$  and  $\chi^c(G1) \leq |C1| = 2$ . Combined with Inequality 1, we conclude that  $\chi^c(G1) = 2$ . So,  $C1$  is an optimal bicoloring cover and by definition,  $\gamma(G1) \geq \max(\{|V_i| | 0 \leq i \leq 3\}) = 3$ .

We now show that  $\gamma(G1)$  is 3. For the sake of contradiction we assume that (i)  $C2$  is an optimal bicoloring cover of size two for  $G1(V, E)$ , (ii)  $\gamma(G1) > 3$ , and (iii)  $C2$  is a witness for  $G1$ . From Lemma 6, it is clear that any independent set of size five for  $G1$  is obtained only by adding any single vertex from  $\mathcal{V}_1$  to the set  $\mathcal{V}_2$ . From Lemma 6, we also know that an independent set of size four in  $G1$  is either the set  $\mathcal{V}_2$ , or any set with a single vertex from  $\mathcal{V}_1$  and any three vertices from  $\mathcal{V}_2$ . Let any such independent set be called  $V_4$ . Since  $C2$  is an optimal bicoloring cover, the canonical partition  $P2$  generated from  $C2$  consists of at most four mutually disjoint independent sets of  $G1$ , and since  $C2$  is a witness for  $G1$ , it must contain  $V_4$  as one of the parts. We define  $V1' = V \setminus V_4$ . Observe that  $V1'$  contains either 7 or 8 vertices from  $\mathcal{V}_1$  i.e.,  $|V1' \cap \mathcal{V}_1| \geq 7$ . If  $V_4 = \mathcal{V}_2$  or  $|V_4| = 5$ , then  $V1' \subseteq \mathcal{V}_1$ . If  $V_4$  has one vertex from  $\mathcal{V}_1$ , and  $|V_4| \geq 4$ , then  $V1'$  has one vertex from  $\mathcal{V}_2$  and seven vertices from  $\mathcal{V}_1$ . From the construction of  $E$ , it is clear that vertices from  $V1' \cap \mathcal{V}_1$  form a  $K_7^3$  in  $G1$ . Consider the partition of the vertices of  $V1'$  into independent sets of  $G1$ . Any such independent set of  $G1$  can include at most two vertices of  $V1' \cap \mathcal{V}_1$ ; three vertices from  $V1' \cap \mathcal{V}_1$  always form a hyperedge in  $G1$ . Since  $V_4$  is already a part in canonical partition  $P2$ , there can be at most three more parts in  $P2$  as  $\chi^c(G1) = 2$ . So,  $V_4$  has at most 5 vertices and the other at most 3 parts can include at most  $3 \cdot 2 = 6$  vertices of  $V1' \cap \mathcal{V}_1$ . So, at least one vertex  $u$  of  $V$  ( $u \in V1' \cap \mathcal{V}_1$ ), is not included in the partition  $P2$ . Therefore, no such partition  $P2$  can include every vertex of  $V$ . So, either  $C2$  is not an optimal bicoloring cover, or  $C2$  is not a witness for  $G1$ , a contradiction to our assumption. Consequently,  $\gamma(G1) = 3$ . We conclude that  $\alpha(G1) = 5 > 3 = \gamma(G1)$ .

**Lemma 7** For the hypergraph  $G1$ , the bicoloring cover number  $\chi^c(G1)$  is two, and the cover independence number  $\gamma(G1)$  is three.

#### 4.2.2 An asymptotic construction demonstrating the separation between $\alpha(G)$ and $\gamma(G)$

In order to give a general asymptotic construction, we choose a composite  $n$ , where  $n = p \cdot q$ , and  $p$  and  $q$  are integers,  $q > p > 2$ , and  $q$  is of the form  $2^z$ ,  $z \in \mathbb{N}$ . Let  $t = \log_n p$ . So,  $p = n^t$  and  $q = n^{1-t}$ . Observe that even keeping  $p$  fixed at a certain value, we can indefinitely increase the values of  $n$  and  $q$ , achieving ever increasing ratio  $\frac{q}{p} = n^{1-2t}$ . Since  $p < q$ , it follows that  $0 < t < 0.5$ . Consider the  $n$ -vertex  $k$ -uniform hypergraph  $G(V, E)$  where  $k = n^t$ . We design our hypergraph  $G$  in such a way that  $\alpha(G) = p + q - 2 = n^t + n^{1-t} - 2$  and  $\gamma(G) = p = n^t$ . Let  $V = \mathcal{V}_1 \cup \mathcal{V}_2$ , where  $\mathcal{V}_1 = \{1, 2, \dots, n - q\}$ , and  $\mathcal{V}_2 = \{n - q + 1, n - q + 2, \dots, n\}$ .

Let  $E_1 = \{\{u_1, \dots, u_k\} | u_1 < \dots < u_k \text{ and } \{u_1, \dots, u_k\} \subset \mathcal{V}_1\}$ ,  $E_2 = \{\{u_1, \dots, u_k\} | u_k \in \mathcal{V}_2, u_1 < \dots < u_{k-1}, \text{ and } \{u_1, \dots, u_{k-1}\} \subset \mathcal{V}_1\}$ ,  $E_3 = \{u_1, \dots, u_k | u_1 < \dots < u_k \text{ and for a fixed } r, 1 \leq r \leq q, \{u_1, \dots, u_k\} \subseteq V_r\}$ . Let  $E = E_1 \cup E_2 \setminus E_3$ . Note that  $\mathcal{V}_2$  is an independent set; hyperedges of  $G$  are either subsets of  $\mathcal{V}_1$ , or include at most one vertex from  $\mathcal{V}_2$ .

We partition the vertices of  $V = \{1, \dots, n\}$  into  $q = n^{1-t}$  parts  $\{V_0, \dots, V_{q-1}\}$ , such that the vertex  $i$  is placed in  $V_{(i-1) \bmod q}$ . Since  $q$  divides  $n$ ,  $|V_r| = p = n^t$ ,  $0 \leq r \leq q - 1$ . So, we get a grid-like arrangement of vertices similar to that in Figure 3 with  $p = n^t$  rows and  $q = n^{1-t}$  columns. Also, observe that each  $V_r$  is an independent set since removal of  $E_3$  from  $E_1 \cup E_2$  removes all hyperedges  $e$  that lie completely inside a part  $V_r$ ,  $0 \leq r \leq q - 1$ .

**Lemma 8** For the hypergraph  $G$ ,  $\alpha(G) = p + q - 2$ . Moreover, independent sets of size greater than  $p$  for  $G$  are obtained by adding at most  $p - 2$  vertices from  $\mathcal{V}_1$  to subsets of  $\mathcal{V}_2$ .

**Proof.** Observe that any maximum independent set can contain at most  $k - 1 = p - 1$  vertices from  $\mathcal{V}_1$ ; otherwise, it introduces at least one hyperedge  $e \in E_1$ . Let  $u_1, \dots, u_{p-1}$  be any  $p - 1$  vertices from  $\mathcal{V}_1$  that belong to some independent set, say  $S$ . If every  $u_j$ ,  $1 \leq j \leq p - 1$ , of this independent set  $S$  belongs to the same part  $V_r$ ,  $1 \leq r \leq q$ , then we can add at most one more vertex of the same set  $V_r$  to the independent set  $S$ : this gives an independent set of size  $p$ . Otherwise, if we vertices of  $S$  are spread over two parts  $V_r$  and  $V_{r'}$ ,  $0 \leq r < r' \leq q - 1$ , then adding any other vertex would give a hyperedge of  $G$  and not an independent set. However, if we restrict the independent set to include only  $p - 2$  vertices  $u_1, \dots, u_{p-2}$  from  $\mathcal{V}_1$ , then we can add all the vertices of  $\mathcal{V}_2$  to the independent set, thereby generating an independent set  $\{u_1, \dots, u_{p-2}, 1, \dots, q\}$ , with  $p + q - 2$  vertices.  $\square$

A lower bound for the bicoloring cover number of  $G$  is now estimated as follows.  $G$  has a complete  $k$ -uniform subhypergraph on the vertices of  $\mathcal{V}_1$ , due to the hyperedges of  $E$ . So, using Theorem ??

$$\chi^c(G) \geq \lceil \log \left( \frac{n - q}{k - 1} \right) \rceil = \lceil \log \left( \frac{pq - q}{p - 1} \right) \rceil = \lceil \log q \rceil. \quad (2)$$

By construction,  $\{V_0, \dots, V_{q-1}\}$  is a partition of  $V$  into independent sets. So, using Lemma 3,  $G$  has a bicoloring cover of size  $\lceil \log q \rceil$  i.e.,

$$\chi^c(G) \leq \lceil \log q \rceil. \quad (3)$$

From Inequalities 2 and 3, it is clear that the set of bicolorings that partitions  $V$  into  $\{V_0, \dots, V_{q-1}\}$ , is a bicoloring cover of optimal size. By definition,  $\gamma(G) \geq \max(\{|V_r| | 0 \leq r \leq q - 1\}) = p$ .

We further claim that  $\gamma(G) = p = n^t$ . For the sake of contradiction we assume that (i)  $C2$  is an optimal bicoloring cover for  $G$  of size  $\lceil \log q \rceil$ , (ii)  $\gamma(G) > p$ , and (iii)  $C2$  is a witness for  $G1$ . From Lemma 8 we know that any independent set of size strictly greater than  $p$  for  $G$  must have at most  $p - 2$  vertices from  $\mathcal{V}_1$  and a subset of vertices from  $\mathcal{V}_2$ . Let any such independent set be  $V_q$ . Then,  $|V_q| \leq p + q - 2 \leq \alpha(G)$  and  $|V_q \cap \mathcal{V}_1| \leq p - 2$ . Since  $C2$  is an optimal bicoloring cover, the canonical partition  $P2$  consists of at most  $2^{\chi^c(G)} = 2^{\lceil \log q \rceil} = q$  independent parts. Since  $C2$  is a witness for  $G1$ , the canonical partition  $P2$  of  $C2$  must contain  $V_q$  as one of its parts. We define  $V' = V \setminus V_q$ . Since (i)  $|V_q \cap \mathcal{V}_1| \leq p - 2$ , and (ii)  $|\mathcal{V}_1| = n - q$ , observe that  $V'$  contains at least  $n - q - p + 2$  vertices from  $\mathcal{V}_1$  i.e.,  $|V' \cap \mathcal{V}_1| \geq n - q - p + 2$ . From the definition of  $E$ , the vertices of  $V' \cap \mathcal{V}_1$  form a  $K_{n-q-p+2}^p$  in  $G$ . Consider the partition of the vertices of  $V'$  into independent sets in  $G$ . Any such independent set of  $G$  can include at most  $p - 1$  vertices of  $V' \cap \mathcal{V}_1$ ;  $p$  vertices from  $V' \cap \mathcal{V}_1$  always form a hyperedge in  $G$ . Since  $V_q$  is already a part in  $P2$ , there can be at most  $q - 1$  more parts in  $P2$  as  $\chi^c(G1) = \lceil \log q \rceil$ . Now,  $V_q$  has at most  $p + q - 2$  vertices and the other at most  $q - 1$  parts can include at most  $(q - 1)(p - 1) = pq - q - p + 1 = n - q - p + 1$  vertices of  $V' \cap \mathcal{V}_1$ . So, at least one vertex  $u \in V' \cap \mathcal{V}_1 \subset V$  is not included in the partition  $P2$ . Therefore, no such partition  $P2$  can include every vertex of  $V$ . So, either  $C2$  is not an optimal bicoloring cover or it is not a witness for  $G1$ , a contradiction to our assumption. So,  $\gamma(G) = p = n^t$ . Consequently,  $\alpha(G) > q = n^{1-t} > p = n^t = \gamma(G)$ . This concludes the construction of hypergraphs where  $\gamma(G) = n^t$  and  $\alpha(G) > n^{1-t} > \gamma(G)$ .

As discussed above, there exists  $k$ -uniform hypergraphs  $G(V, E)$  where  $\alpha(G) \geq n^{1-t}$ , whereas  $\gamma(G) = n^t$ , for some small fraction  $0 < t < 0.5$ . We call this special class of hypergraphs *cover friendly hypergraphs*. We have the following lemma.

**Lemma 9** *For the cover friendly  $n$ -vertex  $n^t$ -uniform hypergraph  $G$ ,  $\chi^c(G) = \lceil \log n^{1-t} \rceil$ , and  $\gamma(G) = n^t$ , where  $t < 0.5$ .*

Using the results from the asymptotic construction of  $G$  in Section 4.2.2, note that  $G1$  in Section 4.2.1 (see Figure 3), is an instance of  $G$  where (i)  $|V| = n = 12$ , (ii)  $p = 3$ , (iii)  $q = 4$ , (iv)  $t = \log_n p = \log_{12} 3$ , (v)  $\alpha(G1) = p + q - 2 = 5$ , (vi)  $\chi^c(G1) = \lceil \log q \rceil = 2$ , and (vii)  $\gamma(G1) = p = 3$ . We summarize the general construction in the following theorem.

**Theorem 9** *Let  $G(V, E)$  be a  $n$ -vertex  $k$ -uniform hypergraph, where  $n = pq$ ,  $2 < p < q$ , such that  $q$  is of the form  $2^z$ ,  $p, z \in \mathbb{N}$ , and  $k = n^t$ . Let  $t = \log_n p$ . Let  $V = \mathcal{V}_1 \cup \mathcal{V}_2$ , where  $\mathcal{V}_1 = \{1, 2, \dots, n - q\}$ , and  $\mathcal{V}_2 = \{n - q + 1, n - q + 2, \dots, n\}$ . Let  $E = E_1 \cup E_2 \setminus E_3$ , where  $E_1 = \{\{u_1, \dots, u_k\} | u_1 < \dots < u_k \text{ and } \{u_1, \dots, u_k\} \subset \mathcal{V}_1\}$ ,  $E_2 = \{\{u_1, \dots, u_k\} | u_k \in \mathcal{V}_2, u_1 < \dots < u_{k-1}, \text{ and } \{u_1, \dots, u_{k-1}\} \subset \mathcal{V}_1\}$ ,  $E_3 = \{u_1, \dots, u_k | u_1 < \dots < u_k, \text{ and for a fixed } r, 1 \leq r \leq q, \{u_1, \dots, u_k\} \subseteq V_r\}$ . Then, (i)  $\chi^c(G) = \lceil \log n^{1-t} \rceil$ , (ii)  $\alpha(G) = n^{1-t} + n^t - 2$ , and (iii)  $\gamma(G) = n^t$ .*

For cover friendly hypergraphs, using Corollary 1, we get an approximation ratio of  $\frac{1}{1-t}$  for approximating  $\chi^c(G)$ . However, using Proposition 1, we get an approximation ratio of at least  $\frac{1}{t}$ . So, we get an improvement (reduction) in approximation ratio for  $\chi^c(G)$  by a factor of at least  $\frac{1-t}{t}$ , using the properties of  $\gamma(G)$ . Moreover, a constant approximation ratio of  $\frac{1}{1-t}$  for approximating  $\chi^c(G)$  is guaranteed for cover friendly hypergraphs as opposed to approximation ratio of  $O(\frac{\log n}{\log \log n - \log \log \log n})$  for general hypergraphs given by Theorem 5, exploiting the characteristics of  $\gamma(G)$ .

### 4.3 Clique number and the bicoloring cover number

We define *clique number* for hypergraphs as follows. Let  $H(V', E')$  be the largest induced subhypergraph of a  $k$ -uniform hypergraph  $G(V, E)$ , where  $V' \subseteq V$ ,  $E' \subseteq E$ , and  $E' \subseteq 2^{V'}$ , such that every subset of  $k$  vertices from  $V'$  constitutes a hyperedge in  $H$ . We say that  $|E'| = \binom{|V'|}{k}$ . We define the

clique number  $\omega(G)$  for hypergraph  $G(V, E)$  as the cardinality of the set  $V'$ . Note that  $\omega(G) \geq k$  for any  $k$ -uniform hypergraph. All non-empty triangle-free undirected graphs have clique number two. Observe that a non-empty  $k$ -uniform hypergraph  $G(V, E)$  has  $\omega(G) = k$  provided no induced subhypergraph  $G'(S, E')$  of  $G(V, E)$ , defined on a subset  $S \subseteq V$  where  $|S| = k + 1$ , has all the  $k + 1$   $k$ -sized subsets of  $S$  as hyperedges in  $E'$ . Like triangle-free graphs,  $k$ -uniform hypergraphs with  $\omega(G) = k$  can be quite a rich class of hypergraphs. We prove the following theorem.

**Theorem 10** *For any  $t \geq 1$ , there exists a  $k$ -uniform hypergraph  $G(V, E)$  with  $\omega(G) = k$  and  $\chi^c(G) > t$ .*

By Theorem ??, we know that  $\chi^c(G) \geq \lceil \log \left( \frac{\omega(G)}{k-1} \right) \rceil$ . In reality,  $\chi^c(G)$  can be arbitrarily far apart from  $\omega(G)$ . Analyzing in a manner similar to that in the existential proof of the existence of triangle-free graphs with arbitrarily large chromatic numbers (see [10]), we demonstrate the separation between  $\chi^c(G)$  and  $\omega(G)$  as stated in Theorem 10.

A random  $k$ -uniform hypergraph  $G_{n,p}(V, E)$  is a  $k$ -uniform hypergraph on  $n$  labeled vertices  $V = [n] = \{1, \dots, n\}$ , in which every subset  $e \subset V$  of size  $|e| = k$  is chosen to be a hyperedge of  $G$  randomly, and independently with probability  $p$ , where  $p$  may depend on  $n$ . We use  $G(V, E)$  or simply  $G$  to denote such a random hypergraph. To show the gap between  $\omega(G)$  and  $\chi^c(G)$ , we choose a random  $k$ -uniform hypergraph  $G(V, E)$  with the value of  $p$  set to  $n^{-\frac{k}{k+1}}$ . For showing that  $\chi^c(G) > t$  some arbitrary integer  $t$ , it is sufficient to show that  $G$  contains no independent set of size  $\lceil \frac{n}{2^t} \rceil$ : we know from Observation 1,  $\chi^c(G) \geq \log \frac{|V|}{\alpha(G)} > \log \frac{n}{2^t} = t$ .

Let  $C_I(G)$  and  $C_\omega(G)$  denote the number of independent sets of size  $\lceil \frac{kn}{(k+1)2^t} \rceil$  and the number of complete subgraphs of order  $k + 1$ , respectively, in the (random)  $k$ -uniform  $n$ -vertex hypergraph  $G(V, E)$ . For any event  $x$ , let  $\mathcal{E}(x)$  denote the expectation of  $x$ . We show that  $\text{Prob}(C_I(G) \geq 1) + \text{Prob}(C_\omega(G) \geq \frac{n}{k+1}) < 1$ ; this implies there exists some hypergraph  $G(V, E)$  such that  $C_I(G) = 0$ , as well as  $C_\omega(G) < \frac{n}{k+1}$ . Then, we delete at most  $\frac{n}{k+1}$  vertices from  $G(V, E)$  to generate a new hypergraph  $G'$  where  $C_I(G') = 0$ , as well as  $C_\omega(G') = 0$ .

First we show that  $\text{Prob}(C_I(G) \geq 1)$  with probability strictly less than  $\frac{1}{2}$  as follows. Let  $F$  be some set of  $\lceil \frac{kn}{(k+1)2^t} \rceil$  vertices in  $G$ . The probability that  $F$  is an independent set is  $(1-p)^{\binom{\lceil \frac{kn}{(k+1)2^t} \rceil}{k}}$ . The expectation  $\mathcal{E}(C_I(G))$  is the above probability summed up over all the possible subsets of size  $\lceil \frac{kn}{(k+1)2^t} \rceil$  in  $G$ . We use the following three known inequalities in our analysis: (i)  $\binom{n}{k} < 2^n$ ,  $0 < k < n$ , (ii)  $1 - x \leq e^{-x}$ ,  $0 \leq x \leq 1$ , and (iii)  $\binom{n}{k} > \left(\frac{n}{k}\right)^k$ ,  $0 < k < n$ . Therefore,

$$\begin{aligned} \mathcal{E}(C_I) &= \binom{n}{\lceil \frac{kn}{(k+1)2^t} \rceil} (1-p)^{\binom{\lceil \frac{kn}{(k+1)2^t} \rceil}{k}} && \text{(from the definition of expectation)} \\ &\leq \binom{n}{\lceil \frac{kn}{(k+1)2^t} \rceil} (1-p)^{\left(\frac{kn}{(k+1)2^t}\right)^k} \\ &< 2^n e^{-p \left(\frac{kn}{(k+1)2^t}\right)^k} && \text{(using (i), (ii) and (iii))} \\ &= 2^n e^{-p \left(\frac{n}{(k+1)2^t}\right)^k} \end{aligned} \tag{4}$$

For a sufficiently large value of  $n$  that depends on both  $k$  and  $t$ , we can show that  $2^n e^{-p \left(\frac{n}{(k+1)2^t}\right)^k}$  is strictly less than  $\frac{1}{2}$  (see Appendix A for details). Now, using Markov's inequality we know that  $P(C_I(G) \geq 1) \leq \mathcal{E}(C_I(G)) < \frac{1}{2}$  for sufficiently large values of  $n$ .

Next, we need to show that the probability of existence of complete subhypergraphs of size  $k + 1$  is small. Let  $W$  be some subset of  $k + 1$  vertices in  $G$ . Then,  $W$  is a complete subgraph with probability  $p^{k+1}$ . The expectation  $\mathcal{E}(C_\omega(G))$  is given by  $\mathcal{E}(C_\omega) = \binom{n}{k+1} p^{k+1} < \frac{n^{k+1}}{(k+1)!} \cdot n^{-\frac{k}{k+1}(k+1)} = \frac{n}{(k+1)!}$ . Again, using Markov's inequality,  $P(C_\omega(G) \geq \frac{n}{k+1}) < \frac{1}{k!}$ . Since  $P(C_I(G) \geq 1) + P(C_\omega(G) \geq \frac{n}{k+1}) < 1$ .

$\frac{n}{k+1}) < 1$ , there exists some graph  $G$  such that  $C_I(G) = 0$ , as well as  $C_\omega(G) < \frac{n}{k+1}$ . From each of the at most  $(k+1)$ -sized complete subhypergraphs, we can remove one vertex each (and all the hyperedges incident on it), to eliminate that complete subhypergraph of size  $k+1$ . Note that the removal of such vertices and corresponding hyperedges cannot increase the size of any independent set in  $G$ . This transformation results in a subhypergraph  $G'(V', E')$  of  $G$  such that  $G'$  does not contain any  $(k+1)$ -sized complete subhypergraphs, and  $|V'| \geq n - \frac{n}{k+1} = \frac{kn}{k+1}$ . Moreover,  $G'$  does not contain any independent set of size  $\lceil \frac{kn}{(k+1)2^t} \rceil = \lceil \frac{|V'|}{2^t} \rceil$ , and therefore  $\chi^c(G') > t$ . So, this hypergraph  $G'$  has  $\omega(G') = k$  but  $\chi^c(G') > t$ , for any  $t > 1$ , establishing our claim in Theorem 10.

## 5 Bicoloring covers for sparse hypergraphs

A  $k$ -uniform hyperedge is rendered monochromatic with probability  $\frac{2}{2^k} = 2^{-(k-1)}$  in a random bicoloring of its  $k$  vertices. If the number of hyperedges  $|E|$  in a  $k$ -uniform hypergraph is at most  $2^{k-2}$ , then the probability that some hyperedge is rendered monochromatic in a random bicoloring is at most  $\frac{2^{k-2}}{2^{k-1}} < \frac{1}{2}$ . Since the probability that none of the hyperedges is monochromatic is at least  $\frac{1}{2}$ , we have the following algorithm for computing a bicoloring for  $G$ . Randomly and independently color the vertices of  $G$  and check whether all the hyperedges are properly bicolored. If some hyperedge is rendered monochromatic in the random bicoloring then repeat the random bicoloring step. We can easily verify that the expected number of steps of failure is less than two. Extending similar arguments, we develop the following relationship between the number of hyperedges in a  $k$ -uniform hypergraph and the size of its bicoloring cover.

**Theorem 11** *A  $k$ -uniform hypergraph  $G(V, E)$  with  $|E| \leq 2^{(k-1)x-1}$  has a bicoloring cover of size  $x$  that can be computed in expected polynomial time.*

**Proof.** Since all the  $x$  bicolorings are random and independent, the probability that a specific hyperedge becomes monochromatic in each of the  $x$  bicolorings is  $(\frac{1}{2^{k-1}})^x$ . Choosing the number of hyperedges  $|E| \leq 2^{(k-1)x-1}$ , the probability that some hyperedge becomes monochromatic in each of the  $x$  bicolorings is strictly less than  $\frac{1}{2}$ . In other words, the probability that each of the  $|E|$  hyperedges is non-monochromatic in one or more bicolorings is at least  $\frac{1}{2}$ . Consequently, the hypergraph has a cover of size  $x$  and that can be computed by random coloring of vertices in expected two iterations.  $\square$

Since Theorem 11 gives only a sufficiency condition for a  $k$ -uniform hypergraph to have a bicoloring cover of size  $x$ , it is interesting to estimate the smallest integer  $m$  such that there is no bicoloring cover with  $x$  bicolorings for some  $k$ -uniform hypergraph with  $m$  hyperedges. This number  $m$  is a measure of the tightness of the sufficiency condition given by Theorem 11. We define  $m(k, x)$  as the smallest integer such that there exists a  $k$ -uniform hypergraph  $G$  with  $m(k, x)$  hyperedges, which does not have a bicoloring cover of size  $x$ . If the number of hyperedges in the  $k$ -uniform hypergraph is less than  $m(k, x)$ , then it certainly has a bicoloring cover of size  $x$ . In other words, for any hypergraph of size less than  $m(k, x)$ , there exist at least one set of  $x$  bicolorings of vertices that properly bicolores every hyperedge of the hypergraph. However, if the number of hyperedges is greater than or equal  $m(k, x)$ , then we cannot guarantee the existence of a bicoloring cover of size  $x$  for the hypergraph. Alternatively, there exist at least one  $k$ -uniform hypergraph of size  $m(k, x)$  such that no set of  $x$  bicolorings can properly bicolor every hyperedge in the hypergraph. From Theorem 11, it is obvious that  $m(k, x) > 2^{(k-1)x-1}$ .

We note that  $8 < m(2, 3, 2) \leq 84$ . The lower bound is given by the proper substitution in Theorem 11. The upper bound is obtained from a  $K_9^3$  which does not have a bicoloring cover of size 2 (see Theorem ??). Computing the exact values of  $m(k, x)$  for different values of  $x$  by brute-force is difficult. In order to prove that  $m(k, x) = a$ , for some fixed  $x, k$  and  $a$ , one may find out at



least one  $k$ -uniform hypergraph with  $a$  hyperedges that does not have a bicoloring cover of size  $x$ . So, one may check every hypergraph with  $a$  hyperedges for a bicoloring cover of size  $x$ , which is computationally expensive. In order to estimate an upper bound for  $m(k, x)$ , we consider a (i)  $k$ -uniform hypergraph with  $k + k^2$  vertices, (ii) fix a set  $C1$  of  $x$  independent bicolings of the vertex set  $V$ , and (iii) pick  $m$   $k$ -uniform hyperedges uniformly, independently and randomly. Let the probability that a randomly picked hyperedge  $e$  becomes monochromatic in a random bicoloring be at least  $p$ ; below we estimate a lower bound for  $p$  considering random bicolings. Then, the probability that  $e$  becomes monochromatic in each of the  $x$  bicolings is at least  $p^x$ . So, the probability that  $e$  becomes non-monochromatic in at least one of the  $x$  bicolings is at most  $1 - p^x$ . Since, we are choosing  $m$  hyperedges independently, randomly and uniformly, the probability that every hyperedge from the  $m$  chosen hyperedges becomes non-monochromatic in at least one of the  $x$  of bicolings in  $C1$  is at most  $(1 - p^x)^m$ . Since there are  $2^{n^x}$  ways to perform the  $x$  independent bicolings, the probability each of the  $m$  chosen hyperedges becomes non-monochromatic in at least one set of  $x$  colorings is at most  $2^{n^x}(1 - p^x)^m$ . So, if  $f(n, x, p, m) = 2^{n^x}(1 - p^x)^m < 1$ , then there exists at least one set of  $m$  hyperedges that cannot be covered by any set of  $x$  bicolings.

Now we estimate a lower bound for  $p$ , the probability with which any randomly picked hyperedge becomes monochromatic in any bicoloring of the  $k + k^2$  vertices in  $V$ . Any bicoloring colors some vertices with color 1 and rest with color 2. Let the set of color 1 vertices be of size  $a$ . Then, the total number of monochromatic hyperedges is  $\binom{a}{k} + \binom{n-a}{k}$ . This sum is minimized at  $a = \lceil \frac{n}{2} \rceil$ . Therefore, the probability that a particular random hyperedge  $e$  is monochromatic in one bicoloring is at least  $2 \frac{\binom{\frac{n}{2}}{k}}{\binom{n}{k}} = 2 * \frac{\frac{n}{2}(\frac{n}{2}-1)\dots(\frac{n}{2}-k+1)}{n(n-1)\dots(n-k+1)} > \frac{1}{2^{k-1}} \left(\frac{n-2k}{n-k}\right)^k = \frac{1}{2^{k-1}} \left(1 - \frac{1}{k}\right)^k$  (since  $\frac{\frac{n}{2}}{n} > \frac{\frac{n}{2}-1}{n-1} > \dots > \frac{\frac{n}{2}-k+1}{n-k+1} > \frac{\frac{n}{2}-k}{n-k}$ ). Let  $p = \frac{1}{2^{k-1}} \left(1 - \frac{1}{k}\right)^k$ . For  $k \geq 2$ ,  $\left(1 - \frac{1}{k}\right)^k \geq \frac{1}{4}$ . We find that the expression  $f(n, x, p, m)$  is upper bounded by  $2^{n^x} \left(1 - \left(\frac{1}{2^{k-1}}\right)^x\right)^m < 2^{n^x} e^{-\frac{m}{2^{(k-1)x}}}$ . The last expression becomes unity when  $m$  is set to  $2^{(k-1)x} \cdot n \cdot x \ln 2$ . This implies that there exists a hypergraph with  $m$  hyperedges such that at least one of the  $m$  hyperedges remains monochromatic in each set of  $x$  bicolings. Since  $n = k^2 + k$ , we have  $2^{(k-1)x} \cdot n \cdot x \ln 2 < 2^{(k-1)x} \cdot 2k^2 \cdot x \cdot 2 = xk^2 2^{(k-1)x+2}$ . We state our result in the following theorem.

**Theorem 12**  $2^{(k-1)x-1} < m(k, x) \leq x \cdot k^2 \cdot 2^{(k-1)x+2}$ .

## 6 Computing bicoloring covers for hypergraphs with bounded dependency

The *dependency* of a hyperedge  $e$  in a  $k$ -uniform hypergraph  $G(V, E)$ , denoted by  $d(G, e)$  is the number of hyperedges in the set  $E$  with which  $e$  shares at least one vertex. The *dependency of a hypergraph*  $d(G)$  or simply  $d$ , denotes the maximum dependency of any hyperedge in the hypergraph  $G$ . Lovász local lemma [5, 12, 11] ensures the existence of a proper bicoloring for any  $k$ -uniform hypergraph provided the dependency of the hypergraph is upper bounded by  $\frac{2^{k-1}}{e} - 1$ . Furthermore, the constructive version of Lovász local lemma by Moser and Tardos [11] enables the computation of a bicoloring of a  $k$ -uniform hypergraph with dependency at most  $\frac{2^{k-1}}{e} - 1$  by a randomized algorithm. Chandrasekaran et.al. [3] proposed a derandomization for local lemma that computes a bicoloring in polynomial time. In what follows, we use similar techniques for establishing permissible bounds on the dependency of a hypergraph as a function of the size of its desired bicoloring cover, and for computing such bicoloring covers. The Kolmogorov complexity approach for Lovász local lemma leads to a method that can bicolor a hypergraph whose dependency is at most  $2^k/8$  (see [7]).

Let  $P$  be a finite set of mutually independent random variables in a probability space. We consider a finite set  $\mathcal{A}$  of events, where each event  $A \in \mathcal{A}$  is determined by a subset  $S(A) \subseteq P$  of

the variables in  $P$ . We say that an event  $A \in \mathcal{A}$  is violated if an evaluation of variables in  $S(A)$  results in the occurrence of  $A$ . We have the following lemma due to Moser and Tardos [11].

**Lemma 10** [11] *Let  $P$  be a finite set of mutually independent random variables in a probability space. Let  $\mathcal{A}$  be a finite set of events determined by these variables. For any  $A \in \mathcal{A}$ , let  $\Gamma(A)$  denote the set of all the events in  $\mathcal{A}$  that depend on  $A$ . If there exists an assignment of reals  $x : \mathcal{A} \rightarrow (0, 1)$  such that  $\forall A \in \mathcal{A} : \Pr[A] \leq x(A) \prod_{B \in \Gamma(A)} (1 - x(B))$ , then there exists an assignment of values to the variables in  $P$  not violating any of the events in  $\mathcal{A}$ . Moreover the Moser-Tardos Sequential Solver algorithm [11] resamples an event  $A \in \mathcal{A}$  at most an expected  $x(A)/(1 - x(A))$  times before it finds such an evaluation. Thus, the expected total number of resampling steps is at most  $\sum_{A \in \mathcal{A}} \frac{x(A)}{1 - x(A)}$ .*

In particular, if  $\forall A \in \mathcal{A}$ ,  $x(A)$  is set to  $\frac{1}{d+1}$  and  $\Pr[A] \leq p$ , then the premise of Lemma 10 reduces to  $ep(d+1) \leq 1$ , where  $d$  is the maximum dependency  $\max_{A \in \mathcal{A}} |\Gamma(A)|$  of any event  $A$  in  $\mathcal{A}$ . So, from Lemma 10 with suitable substitutions, we get the following corollary.

**Corollary 2** *Let  $P$  be a finite set of mutually independent random variables in a probability space. Let  $\mathcal{A}$  be a finite set of events determined by these variables, where  $m = |\mathcal{A}|$ . For any  $A \in \mathcal{A}$ , let  $\Gamma(A)$  denote the set of all the events in  $\mathcal{A}$  that depend on  $A$ . Let  $d = \max_{A \in \mathcal{A}} |\Gamma(A)|$ . If  $\forall A \in \mathcal{A} : \Pr[A] \leq p$  and  $ep(d+1) \leq 1$ , then an assignment of the variables not violating any of the events in  $\mathcal{A}$  can be computed using expected  $\frac{1}{d}$  resamplings per event and expected  $\frac{m}{d}$  resamplings in total.*

---

**Algorithm MTC:** Randomized algorithm for computing a bicoloring cover

---

**Data:**  $k$ -uniform hypergraph  $G(V, E)$  with  $d \leq \frac{2^{|X|}(k-1)}{e} - 1$

**Result:** Set  $X$  of bicolorings of size  $|X|$

**for**  $v \in V$  **do**

**for**  $i \in \{1, \dots, |X|\}$  **do**

$r_v^i \leftarrow$  a random evaluation of  $v$  in  $i^{th}$  bicoloring of  $X$ ;

**while**  $\exists A_i \in \mathcal{A} : A_i$  happens i.e., every bicoloring in  $X$  renders  $E_i$  monochromatic **do**

    Pick an arbitrary violated event  $A_i \in \mathcal{A}$ ;

**for**  $v \in E_i$  **do**

**for**  $i \in \{1, \dots, |X|\}$  **do**

$r_v^i \leftarrow$  a random evaluation of  $v$  in  $i^{th}$  bicoloring of  $X$ ;

---

In what follows, we use Corollary 2 and an adaptation of the Moser-Tardos algorithm which we call *MTC*, to compute a bicoloring cover  $X$  of  $x$  bicolorings, for a  $k$ -uniform hypergraph  $G(V, E)$ . Let the event  $A_i$  correspond to the hyperedge  $E_i \in E$  becoming monochromatic in each of the  $x$  random and independent bicolorings. The probability  $p(A_i)$  is at most  $\left(\frac{1}{2^{k-1}}\right)^x$ . So, using Corollary 2, the maximum allowable dependency  $d$  of the hypergraph  $G$  is  $\frac{2^{x(k-1)}}{e} - 1$ , so that  $G$  has a bicoloring cover with  $x$  bicolorings. In order to compute a bicoloring cover for  $G$  with  $x$  bicolorings, where  $d(G) \leq \frac{2^{x(k-1)}}{e} - 1$ , the algorithm *MTC*, repeatedly recolors vertices of monochromatic hyperedges, one at a time. It picks up a monochromatic hyperedge and generates  $x$  random bits 0/1 for each vertex of the monochromatic hyperedge, one bit for each of the  $x$  bicolorings. If there are several monochromatic hyperedges then *MTC* picks up any such hyperedge for recoloring all its vertices with colors 0/1, for each of the  $x$  bicolorings. Each such step is called a resampling step, where one hyperedge gets all its  $k$  vertices recolored for each of the  $x$  bicolorings. The correctness of *MTC* follows from the correctness of the Moser-Tardos constructive version of the local lemma; the algorithm terminates after generating a bicoloring cover with  $x$  bicolorings.

Since *MTC* is a randomized algorithm, it consumes random bits in each resampling step. Let  $T$  be the total number of resampling steps performed. The algorithm *MTC* uses  $nx + Tkx$  random bits for computing a bicoloring cover of size  $x$ , given  $d \leq \frac{2^{x(k-1)}}{e} - 1$ . Here,  $nx$  random bits are for

the initial assignment (one bit per vertex per bicoloring),  $kx$  bits each for each of the  $T$  resampling steps (one bit per vertex of the resampled hyperedge per bicoloring). We know from Corollary 2 that the expected number of resampling steps is  $T = \frac{m}{d}$ . So, the expected number of random bits used by the algorithm is  $nx + kx\frac{m}{d}$ . Since  $d \leq \frac{2^{x(k-1)}}{e} - 1$ , we have  $x \geq \frac{1}{k-1} \log(e(d+1))$ . Therefore, the expected number of random bits used by the algorithm is at least  $\frac{1}{k-1} \log(e(d+1))(n + k\frac{m}{d})$ . We summarize these results as the following theorem.

**Theorem 13** *Let  $G(V, E)$  be a  $k$ -uniform hypergraph,  $n = |V|$ ,  $m = |E|$ . Let the dependency of the hypergraph  $d(G)$  be upper bounded by  $\frac{2^{x(k-1)}}{e} - 1$ , for some  $x \in \mathcal{N}$ . Then, there exists a bicoloring cover of size  $x$ , which can be computed by a randomized algorithm using  $\frac{1}{d}$  expected resamplings per hyperedge and  $\frac{m}{d}$  resamplings in total, using expected  $nx + kx\frac{m}{d}$  random bits.*

We note that Algorithm *MTC* can be derandomized in the same manner as done for the case of bicolorings in [3]. As the dependency of the hypergraph grows, the bicoloring cover size guaranteed by the local lemma also increases. However, from Theorem ??, we know that for any  $k$ -uniform graph  $G(V, E)$ ,  $|V| = n$ ,  $\chi_c(G) \leq \left\lceil \log\left(\frac{n}{k-1}\right) \right\rceil$ . So, the application of this algorithm is practical for the case where it guarantees a cover of size of at most  $\lceil \log(\frac{n}{k-1}) \rceil$ . We can find the maximum dependency for which this algorithm is useful by simply replacing  $x$  in the dependency bound as  $d \leq \frac{2^{x(k-1)}}{e} - 1 \leq \frac{1}{e} 2^{\log(\frac{n}{k-1})(k-1)} - 1$ , that is  $d \leq \frac{1}{e} \left(\frac{n}{k-1}\right)^{(k-1)} - 1$ .

## 7 Concluding remarks

Bounds for bicoloring cover numbers established in this paper are supported by algorithms that generate the bicoloring covers of the corresponding sizes. The algorithms and bounds can be generalized for multicolorings, where more than two colors are used. In such natural extensions to multicolorings, the constraint imposed on every hyperedge can be relaxed so that at least  $p \geq 2$  vertices of the hyperedge are distinctly colored in at least one of the multicolorings.

Throughout the paper, we have used independent bicolorings in our probabilistic analysis. Whether the use of mutually dependent bicolorings would lead to discovery of better bounds for bicoloring cover numbers, remains an open question. Computing the exact value or approximating the cover independence number  $\gamma(G)$  remains an open problem.

## References

- [1] N. Alon. A note on the decomposition of graphs into isomorphic matchings. *Acta Mathematica Hungarica*, 42(3-4):221–223, 1983.
- [2] A. Bialostocki and Y. Roditty. 3k2-decomposition of a graph. *Acta Mathematica Academiae Scientiarum Hungarica*, 40(3-4):201–208, 1982.
- [3] Karthekeyan Chandrasekaran, Navin Goyal, and Bernhard Haeupler. Deterministic algorithms for the lovász local lemma. *SIAM Journal on Computing*, 42(6):2132–2155, 2013.
- [4] F.R.K. Chung and R.L. Graham. Recent results in graph decompositions. In *Combinatorics*, pages 103–123. Cambridge University Press, 1981. Cambridge Books Online.
- [5] P. Erdős and L. Lovász. Problems and results on 3-chromatic hypergraphs and some related questions. *Colloq. Math. Soc. János Bolyai*, 10:609–627, 1975.
- [6] Uriel Feige and Joe Kilian. Zero knowledge and the chromatic number. *Journal of Computer and System Sciences*, 57(2):187 – 199, 1998.

- [7] Lance Fortnow. Computational Complexity: A Kolmogorov Complexity Proof of the Lovász Local Lemma. <http://blog.computationalcomplexity.org/2009/06/kolmogorov-complexity-proof-of-lov.html>, 2009.
- [8] Michael Krivelevich and Benny Sudakov. Approximate coloring of uniform hypergraphs. *Journal of Algorithms*, 49(1):2 – 12, 2003. 1998 European Symposium on Algorithms.
- [9] László Lovász. Coverings and colorings of hypergraphs. In *Proc. 4th Southeastern Conference on Combinatorics, Graph Theory, and Computing, Utilitas Mathematica Publishing, Winnipeg*, pages 3–12, 1973.
- [10] Michael Molloy and Bruce Reed. *Graph Colouring and the Probabilistic Method*. Algorithms and Combinatorics, Vol. 23. Springer-Verlag, Berlin, 2002.
- [11] Robin A. Moser and Gábor Tardos. A constructive proof of the general Lovász Local Lemma. *J. ACM*, 57(2):11:1–11:15, February 2010.
- [12] Rajeev Motwani and Prabhakar Raghavan. *Randomized Algorithms*, pages 115–118. Cambridge University Press, New York, NY, USA, 1995.
- [13] Douglas B. West. *Introduction to Graph Theory*, pages 378–388. Prentice Hall, second edition, 2000.

## A Estimation of $n$ for Inequality 4 in Section 4.3

From Section 4.3, we have,  $p = n^{-\frac{k}{k+1}}$ . Using Inequality 4, we need to choose a value of  $n$  that satisfies the inequality  $2^n e^{-p \left( \frac{n}{(k+1)2^t} \right)^k} < \frac{1}{2}$ . We proceed as follows.

$$\begin{aligned}
2^n e^{-p \left( \frac{n}{(k+1)2^t} \right)^k} &< \frac{1}{2} \\
\Leftrightarrow 2^{n+1} &< e^{p \left( \frac{n}{(k+1)2^t} \right)^k} \\
\Leftrightarrow (n+1) \log_e 2 &< n^{-\frac{k}{k+1}} \left( \frac{n}{(k+1)2^t} \right)^k \\
\Leftrightarrow (k+1)^k 2^{tk} \log_e 2 &< \frac{n^{\frac{k^2}{k+1}}}{n+1}
\end{aligned}$$

This inequality can always be satisfied for a sufficiently large value of  $n > ((k+1)^k * 2^{tk+1} \log_e 2)^{(k+1)/(k^2)}$ .