

Some notes on the k -normal elements and k -normal polynomials over finite fields

Mahmood Alizadeh

*Department of Mathematics, College of Science, Ahvaz Branch, Islamic Azad
University, Ahvaz, Iran*

E-mail: alizadeh@iauahvaz.ac.ir

Abstract

Recently, the k -normal element over finite fields is defined and characterized by Huczynska et al.. In this paper, the characterization of k -normal elements, by using to give a generalization of Schwartz's theorem, which allows us to check if an element is a normal element, is obtained. In what follows, in respect of the problem of existence of a primitive 1-normal element in \mathbb{F}_{q^n} over \mathbb{F}_q , for all q and n , had been stated by Huczynska et al., it is shown that, in general, this problem is not satisfied. Finally, a recursive method for constructing 1-normal polynomials of higher degree from a given 1-normal polynomial over \mathbb{F}_{2^m} is given.

Keywords: Finite field, normal basis, k -normal element, k -normal polynomial, primitive element.

1 Introduction

Let \mathbb{F}_q , be the Galois field of order $q = p^m$, where p is a prime and m is a natural number, and \mathbb{F}_q^* be its multiplicative group. A *normal* basis of \mathbb{F}_{q^n} over \mathbb{F}_q is a basis of the form $N = \{\alpha, \alpha^q, \dots, \alpha^{q^{n-1}}\}$, i.e., a basis that consists of the algebraic conjugates of a fixed element $\alpha \in \mathbb{F}_{q^n}^*$. Recall that an element $\alpha \in \mathbb{F}_{q^n}$ is said to generate a normal basis over \mathbb{F}_q if its conjugates form a basis of \mathbb{F}_{q^n} as a vector space over \mathbb{F}_q . For convenience an element α generating a normal basis, is called a *normal* element.

A monic irreducible polynomial $F(x) \in \mathbb{F}_q[x]$ is called *normal polynomial* or *N-polynomial* if its roots form a normal basis or, equivalently, if they are linearly independent over \mathbb{F}_q . The minimal polynomial of an element in a normal basis $\{\alpha, \alpha^q, \dots, \alpha^{q^{n-1}}\}$ is $m(x) = \prod_{i=0}^{n-1} (x - \alpha^{q^i}) \in \mathbb{F}_q[x]$, which is irreducible over \mathbb{F}_q .

Normal bases have many applications, including coding theory, cryptography and computer algebra systems. The elements in a normal basis are exactly the roots of some N -polynomial. Hence an N -polynomial is just another way of describing a normal basis. The construction of normal polynomials over finite fields is a challenging mathematical problem [7].

Recently, the k -normal element over finite fields is defined and characterized by Huczynska, Mullen, Panario and Thomson [6]. The following definition for checking whether an element is a k -normal element, is given by them.

Definition 1.1 *For each $0 \leq k \leq n-1$, the element $\alpha \in \mathbb{F}_{q^n}$ is called a k -normal element if $\deg(\gcd(x^n - 1, \sum_{i=0}^{n-1} \alpha^{q^i} x^{n-1-i})) = k$.*

Using this mention, a normal element of \mathbb{F}_{q^n} over \mathbb{F}_q is a 0-normal element. The existence of primitive normal elements has been obtained in [1] and [2] for sufficiently large q and n . The *Primitive Normal Basis Theorem* has been established for all q and n in [8] and [4]. Moreover, the existence of primitive 1-normal elements is established in [6], for sufficiently large q^n . Also, the existence of primitive 1-normal elements of \mathbb{F}_{q^n} over \mathbb{F}_q , for all q and n is stated in ([6], Problem 6.2), as a problem.

In this paper, in Section 2, some definitions, notes and results which are useful for our study have been stated. Section 3 is devoted to characterization and construction of k -normal elements (especially, 1-normal elements). Also, in this section, in respect of the problem of existence of primitive 1-normal elements, for all q and n , some value of q and n is given, such that the extension finite field \mathbb{F}_{q^n} is not consists of any primitive 1-normal element. Finally, in Section 4, by definition of k -normal polynomials, given in Section 2, a recursive method for construction 1-normal polynomials of higher degree from a given 1-normal polynomial over \mathbb{F}_{2^m} is given.

2 Preliminary notes

We use the definitions, notations and results given by Huczynska [6], Cohen[3], Gao [5] and Kyuregyan [7], where similar problems are considered. We need the following results for our further study.

Recall that $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}(\alpha)$, the trace of α in \mathbb{F}_{q^n} over \mathbb{F}_q , is given by $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}(\alpha) = \sum_{i=0}^{n-1} \alpha^{q^i}$. For convince, $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}$ is denoted by $Tr_{q^n|q}$. To state further results, recall that the Frobenius map

$$\sigma : \eta \rightarrow \eta^q, \quad \eta \in \mathbb{F}_{q^n}$$

is an automorphism of \mathbb{F}_{q^n} that fixes \mathbb{F}_q . In particular, σ is a linear transformation of \mathbb{F}_{q^n} viewed as a vector space of dimension n over \mathbb{F}_q . For any polynomial $g(x) = \sum_{i=0}^{n-1} g_i x^i \in F_q[x]$, define

$$g(\sigma)\eta = \left(\sum_{i=0}^{n-1} g_i \sigma^i \right) \eta = \sum_{i=0}^{n-1} g_i \eta^{q^i},$$

which is also a linear transformation on F_{q^n} .

The *null space* of $g(\sigma)$ is defined to be the set of all elements $\alpha \in \mathbb{F}_{q^n}$ such that $g(\sigma)\alpha = 0$, we also call it the null space of $g(x)$. For any element $\alpha \in \mathbb{F}_{q^n}$, the monic polynomial $g(x) \in \mathbb{F}_q[x]$ of the smallest degree such that $g(\sigma)\alpha = 0$ is called the σ -order of α . This polynomial is denoted by $Ord_{\alpha, \sigma}(x)$. Note that $Ord_{\alpha, \sigma}(x)$ divides any polynomial $h(x)$ annihilating α . In particular, for every $\alpha \in F_{q^n}$, $Ord_{\alpha, \sigma}(x)$ divides $x^n - 1$, the minimal or characteristic polynomial for σ .

Proposition 2.1 [6] *Let $\alpha \in \mathbb{F}_{q^n}$. Then α is a k -normal element over \mathbb{F}_q if and only if $\deg(Ord_{\alpha, \sigma}(x)) = n - k$.*

Proposition 2.2 [3] *A necessary condition for existence a primitive element with zero trace in \mathbb{F}_{q^n} over \mathbb{F}_q is $n \geq 3$, except when $n = 2$ and $q = 3$. Also if $n \geq 3$, such a primitive element exists, except when $n = 3$ and $q = 4$.*

Recall that, an element $\alpha \in \mathbb{F}_{q^n}$ is called a proper element of \mathbb{F}_{q^n} over \mathbb{F}_q if $\alpha \notin \mathbb{F}_{q^v}$ for any proper divisor v of n . So by this mention, an element $\alpha \in \mathbb{F}_{q^n}$ is called a proper k -normal element of \mathbb{F}_{q^n} over \mathbb{F}_q if α is a k -normal element and a proper element of \mathbb{F}_{q^n} over \mathbb{F}_q .

Definition 2.3 *A monic irreducible polynomial $P(x) \in \mathbb{F}_q$ of degree n is called a k -normal polynomial (or N_k -polynomial) over \mathbb{F}_q if its roots be the k -normal elements of \mathbb{F}_{q^n} over \mathbb{F}_q .*

By using the above definition, an N -polynomial is an N_0 -polynomial. Since, the proper k -normal elements of \mathbb{F}_{q^n} over \mathbb{F}_q are the roots of a k -normal polynomial of degree n over \mathbb{F}_q , hence the k -normal polynomials of degree n over \mathbb{F}_q is just another way of describing the proper k -normal elements of \mathbb{F}_{q^n} over \mathbb{F}_q .

The following proposition is useful for constructing N_1 -polynomials over \mathbb{F}_{2^m} .

Proposition 2.4 [7] *Let $\delta \in \mathbb{F}_{2^s}^*$ and $F_1(x) = \sum_{i=0}^n c_i x^i$ be an irreducible polynomial of degree n over \mathbb{F}_{2^m} . Also let $Tr_{2^m|2}(\frac{c_1 \delta}{c_0}) = 1$ and $Tr_{2^m|2}(\frac{c_{n-1}}{\delta}) = 1$. Then the sequence $(F_u(x))_{u \geq 1}$, defined by*

$$F_{u+1}(x) = x^{n2^{u-1}} F_u(x + \delta^2 x^{-1}) \quad u \geq 1.$$

is a sequence of irreducible polynomials of degree $2^u n$ over \mathbb{F}_{2^m} for every $u \geq 1$.

3 Characterization and construction of k -normal elements and existence of primitive 1-normal elements

In this section, the characterization and construction of k -normal elements (specially, of 1-normal elements), and also a characterization of some extension finite fields, consist a primitive 1-normal element is given. In the continue, in respect

of the problem of existence of the primitive 1-normal elements for all q and n ([6], problem 6.2), some values of q and n is introduced, such that the extension finite field \mathbb{F}_{q^n} is not consists of any primitive 1-normal element over \mathbb{F}_q .

Let p denote the characteristic of \mathbb{F}_q and let $n = n_1 p^e = n_1 t$, with $\gcd(p, n_1) = 1$ and suppose that $x^n - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^n - 1 = (\varphi_1(x)\varphi_2(x) \cdots \varphi_r(x))^t, \quad (1)$$

where $\varphi_i(x) \in \mathbb{F}_q[x]$ are the distinct irreducible factors of $x^n - 1$. For each s , $0 \leq s < n$, let there is a $u_s > 0$ such that, $R_{s,1}(x), R_{s,2}(x), \dots, R_{s,u_s}(x)$ be all of the s degree polynomials dividing $x^n - 1$. So, from (1) we can write $R_{s,i}(x) = \prod_{j=1}^r \varphi_j^{t_{ij}}(x)$, for each $1 \leq i \leq u_s$, $0 \leq t_{ij} \leq t$. Let

$$\phi_{s,i}(x) = \frac{x^n - 1}{R_{s,i}(x)}, \quad (2)$$

for $1 \leq i \leq u_s$. Then, there is a useful characterization of the k -normal elements in \mathbb{F}_{q^n} over \mathbb{F}_q as follows.

Lemma 3.1 *An element $\alpha \in \mathbb{F}_{q^n}$ is a k -normal element if and only if, there is a j , $1 \leq j \leq u_k$, such that*

$$\phi_{k,j}(\sigma)\alpha = 0, \quad (3)$$

and also

$$\phi_{s,i}(\sigma)\alpha \neq 0, \quad (4)$$

for each s , $k < s < n$, and $1 \leq i \leq u_s$.

Proof. According to Proposition 2.1, α is a k -normal element of \mathbb{F}_{q^n} over \mathbb{F}_q if and only if $\deg(\text{Ord}(\alpha)) = n - k$. This is true if and only if the Equations (3) and (4) hold. ■

In the case $k = 0$, the Lemma 3.1 is obtained by Schwartz [12](see also [10], Theorem 4.10). Actually, Lemma 3.1 is a generalization of Schwartz's theorem. The definition 1.1 does give us a method to test if an element is a k -normal element. In fact, the Lemma 3.1 dose give us another way to check if an element is k -normal.

The following results are some immediate consequences of Lemma 3.1 .

Theorem 3.2 *Let $n = n_1 p^e$, for some $e \geq 1$ and $a, b \in \mathbb{F}_q^*$. Then the element α is a proper 1-normal element of \mathbb{F}_{q^n} over \mathbb{F}_q if and only if $a + b\alpha$ is a proper 1-normal element of \mathbb{F}_{q^n} over \mathbb{F}_q .*

Proof. Let $n = n_1 p^e = n_1 t$, then by (1), $x^n - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^n - 1 = (x^{n_1} - 1)^t = (\varphi_1(x)\varphi_2(x) \cdots \varphi_r(x))^t.$$

Let

$$R_{1,1}(x) = \varphi_1(x) = x - 1. \quad (5)$$

Set for each s , $0 < s < n$ and $1 \leq i \leq u_s$, except when $s = i = 1$,

$$\begin{aligned}\phi_{s,i}(x) &= \frac{x^n - 1}{R_{s,i}(x)} \\ &= (x - 1)^t M_{s,i}(x) \\ &= (x - 1) M'_{s,i}(x),\end{aligned}$$

where

$$M'_{s,i}(x) = (x - 1)^{t-1} M_{s,i}(x) = \sum_{v=0}^{n-s-1} t'_{iv} x^v.$$

Hence

$$\phi_{s,i}(x) = \sum_{v=0}^{n-s-1} t'_{iv} x^{v+1} - \sum_{v=0}^{n-s-1} t'_{iv} x^v. \quad (6)$$

From (6), we obtain

$$\begin{aligned}\phi_{s,i}(\sigma)(a + b\alpha) &= \sum_{v=0}^{n-s-1} t'_{iv} (a + b\alpha)^{q^{v+1}} - \sum_{v=0}^{n-s-1} t'_{iv} (a + b\alpha)^{q^v}, \\ &= a \sum_{v=0}^{n-s-1} t'_{iv} + b \sum_{v=0}^{n-s-1} t'_{iv} \alpha^{q^{v+1}} - a \sum_{v=0}^{n-s-1} t'_{iv} - b \sum_{v=0}^{n-s-1} t'_{iv} \alpha^{q^v} \\ &= b \left(\sum_{v=0}^{n-s-1} t'_{iv} \alpha^{q^{v+1}} - \sum_{v=0}^{n-s-1} t'_{iv} \alpha^{q^v} \right) = b \phi_{s,i}(\sigma)(\alpha),\end{aligned} \quad (7)$$

for each s , $0 < s < n$ and $1 \leq i \leq u_s$, except when $s = i = 1$. On the other hand, in the case $s = i = 1$, from (5), we have

$$\phi_{1,1}(x) = \frac{x^n - 1}{R_{1,1}(x)} = \frac{x^n - 1}{x - 1} = x^{n-1} + x^{n-2} + \cdots + x + 1 = \sum_{i=0}^{n-1} x^i.$$

So

$$\begin{aligned}\phi_{1,1}(\sigma)(a + b\alpha) &= \sum_{i=0}^{n-1} (a + b\alpha)^{q^i} \\ &= \sum_{i=0}^{n-1} a + b \sum_{i=0}^{n-1} \alpha^{q^i} \\ &= na + b \phi_{1,1}(\sigma)(\alpha) \\ &= b \phi_{1,1}(\sigma)(\alpha).\end{aligned} \quad (8)$$

From (7) and (8), it follows that

$$\phi_{s,i}(\sigma)(a + b\alpha) = b \phi_{s,i}(\sigma)(\alpha)$$

for each s , $0 < s < n$ and $1 \leq i \leq u_s$. By Lemma 3.1, the proof is completed. ■

Theorem 3.3 *Suppose that α is a proper element of \mathbb{F}_{q^2} . Then α is 1-normal over \mathbb{F}_q if and only if $Tr_{q^2|q}(\alpha) = 0$.*

Proof. By (1), $x^2 - 1$ has the following factorization in $\mathbb{F}_q[x]$:

$$x^2 - 1 = (x - 1)(x + 1).$$

So, we have $R_{1,1}(x) = x - 1$ and $R_{1,2}(x) = x + 1$. Hence, $\phi_{1,1}(x) = x + 1$ and $\phi_{1,2}(x) = x - 1$. Note that, by Lemma 3.1, to complete the proof, it is enough that

$$\phi_{1,1}(\sigma)\alpha = \alpha^q + \alpha = Tr_{q^2|q}(\alpha) = 0. \quad (9)$$

This is satisfied by hypothesis. \blacksquare

Theorem 3.4 *Let n be a prime different from p and q a primitive element modulo n . Suppose that α is a proper element of \mathbb{F}_{q^n} . Then α is a 1-normal element of \mathbb{F}_{q^n} over \mathbb{F}_q if and only if $Tr_{q^n|q}(\alpha) = 0$.*

Proof. Note that

$$x^n - 1 = (x - 1)(x^{n-1} + \cdots + x + 1).$$

Since q is a primitive modulo n , $x^{n-1} + \cdots + x + 1$ is an irreducible polynomial over \mathbb{F}_q . Hence, from (1), we have $R_{1,1}(x) = x - 1$ and $R_{n-1,1}(x) = x^{n-1} + \cdots + x + 1$. So, $\phi_{1,1}(x) = x^{n-1} + \cdots + x + 1$ and $\phi_{n-1,1}(x) = x - 1$. Thus, by Lemma 3.1, $\alpha \in \mathbb{F}_{q^n}$ is a 1-normal element of \mathbb{F}_{q^n} over \mathbb{F}_q if and only if

$$\phi_{1,1}(\sigma)\alpha = \sum_{i=0}^{n-1} \sigma^i(\alpha) = \sum_{i=0}^{n-1} \alpha^{q^i} = Tr_{q^n|q}(\alpha) = 0. \quad (10)$$

and

$$\phi_{n-1,1}(\sigma)\alpha = \alpha^q - \alpha \neq 0. \quad (11)$$

The expressions (10) and (11) are satisfied by hypothesis. This completes the proof. \blacksquare

In the case, the 0-normal elements, similar problems for Theorems 3.2 and 3.4 had been established by Kyuregyan [7] and Pei, Wang and Omura [11], respectively.

The following corollary which characterize some of extension finite fields, consist a primitive 1-normal element, is an immediate consequence of Theorems 3.3, 3.4 and Proposition 2.2.

Corollary 3.5 *Let n be a prime different from p and q a primitive element modulo n . Then, A necessary condition for existence a primitive 1-normal element in \mathbb{F}_{q^n} over \mathbb{F}_q is $n \geq 3$, except when $n = 2$ and $q = 3$. Also if $n \geq 3$, such a primitive 1-normal element exists, except when $n = 3$ and $q = 4$.*

In respect of the problem of existence of a primitive 1-normal element in \mathbb{F}_{q^n} over \mathbb{F}_q , for all q and n , the following notation, which is an immediate consequence of Corollary 3.5 is given.

Notation 3.6 By Corollary 3.5, it is shown that, there are some extension finite fields \mathbb{F}_{q^n} , with out any primitive 1-normal element over \mathbb{F}_q . For this, for example, we can refer to the finite fields \mathbb{F}_4 over \mathbb{F}_2 , \mathbb{F}_{4^3} over \mathbb{F}_4 and \mathbb{F}_{q^2} over \mathbb{F}_q , for each odd prime power $q \neq 3$. Consequently, it is shown that, the problem of existence of a primitive 1-normal element in \mathbb{F}_{q^n} over \mathbb{F}_q , for all q and n is not satisfied.

4 Recursive construction N_1 -polynomials

In this section, we establish a theorem which will show how Lemma 3.1 and Proposition 2.4 can be applied to produce infinite sequences of N_1 -polynomials over \mathbb{F}_{2^m} . Nothing that $\phi_{s,i}(\sigma)\alpha = \sum_{i=0}^{n-s} a_i \alpha^{q^i}$ for any polynomial $\phi_{s,i}(x) = \sum_{i=0}^{n-s} a_i x^i$, we can restate Lemma 3.1 as follows.

Proposition 4.1 Let $F(x)$ be an irreducible polynomial of degree n over \mathbb{F}_q and α be a root of it. Let $x^n - 1$ factors as (1) and let $\phi_{s,i}(x)$ be as in (14). Then $F(x)$ is a N_k -polynomial over \mathbb{F}_q if and only if, there is j , $1 \leq j \leq u_k$, such that

$$L_{\phi_{k,j}}(\alpha) = 0,$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0,$$

for each s , $k < s < n$, and $1 \leq i \leq u_s$, where u_s is the number of all s degree polynomials dividing $x^n - 1$ and $L_{\phi_{s,i}}(x)$ is the linearized polynomial defined by

$$L_{\phi_{s,i}}(x) = \sum_{v=0}^{n-s} t_{iv} x^{q^v} \text{ if } \phi_{s,i}(x) = \sum_{v=0}^{n-s} t_{iv} x^v.$$

The following theorem dose give us a recursive method for construction 1-normal polynomials of higher degree from a given 1-normal polynomial over \mathbb{F}_{2^m} . A similar result, in the case 0-normal polynomials has been established by Kyuregyan in [7]. We use of an analogues technique, which used by Kyuregyan, in the proof of theorem

Theorem 4.2 Let $\delta \in \mathbb{F}_{2^s}^*$ and $F_1(x) = \sum_{i=0}^n c_i x^i$ be N_1 -polynomial of degree the even integer n over \mathbb{F}_{2^m} . Also let $\text{Tr}_{2^m|2}(\frac{c_1 \delta}{c_0}) = 1$ and $\text{Tr}_{2^m|2}(\frac{c_{n-1}}{\delta}) = 1$. Then the sequence $(F_u(x))_{u \geq 1}$, defined by

$$F_{u+1}(x) = x^{n2^{u-1}} F_u(x + \delta^2 x^{-1}) \quad u \geq 1. \quad (12)$$

is a sequence of N_1 -polynomials of degree $2^u n$ over \mathbb{F}_{2^m} for every $u \geq 1$.

Proof. Since $F_1(x)$ is an irreducible polynomial over \mathbb{F}_{2^m} , so Proposition 2.4 and the hypothesis imply that $F_2(x)$ is irreducible over \mathbb{F}_{2^m} .

Let $\alpha \in \mathbb{F}_{2^{mn}}$ be a root of $F_1(x)$. Since $F_1(x)$ is an N_1 -polynomial of degree n over \mathbb{F}_{2^m} by the hypothesis, so $\alpha \in \mathbb{F}_{2^{mn}}$ is a 1-normal element over \mathbb{F}_{2^m} .

Let $n = n_1 2^e$, where n_1 is a non-negative integer with $\gcd(n_1, 2) = 1$ and $e \geq 0$. For convenience we denote 2^e by t . Let $x^n + 1$ has the following factorization in $\mathbb{F}_{2^m}[x]$:

$$x^n + 1 = (x^{n_1} + 1)^t = (\varphi_1(x)\varphi_2(x) \cdots \varphi_r(x))^t, \quad (13)$$

where the polynomials $\varphi_i(x) \in \mathbb{F}_q[x]$ are the distinct irreducible factors of $x^{n_1} + 1$. From (13), for each $1 < s < n$, that, there is a $u_s > 0$ such that, $R_{s,1}(x)$, $R_{s,2}(x)$, \dots , $R_{s,u_s}(x)$ are all of the s degree polynomials dividing $x^n + 1$, we can write $R_{s,i}(x) = \prod_{j=1}^r \varphi_j^{t_{ij}}(x)$, for each $1 \leq i \leq u_s$, $0 \leq t_{ij} \leq t$. Let

$$\phi_{s,i}(x) = \frac{x^n + 1}{R_{s,i}(x)} = \sum_{v=0}^{n-s} t_{iv} x^v, \quad (14)$$

for $1 \leq i \leq u_s$. Since $F_1(x)$ is an N_1 -polynomial of degree n over \mathbb{F}_{2^m} , so by Proposition 4.1, there is a j , $1 \leq j \leq u_1$, such that

$$L_{\phi_{1,j}}(\alpha) = 0, \quad (15)$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0, \quad (16)$$

for each $1 < s < n$, which there is a $u_s > 0$ such that, $R_{s,1}(x)$, $R_{s,2}(x)$, \dots , $R_{s,u_s}(x)$ be all of the s degree polynomials dividing $x^n + 1$, and $1 \leq i \leq u_s$.

Now, we proceed by proving that $F_{u+1}(x)$ is a 1-normal polynomial. Obviously, by Proposition 2.4 and the hypothesis, $F_{u+1}(x)$ is irreducible over \mathbb{F}_{2^m} . Let α_{u+1} be a root of $F_{u+1}(x)$. Note that by (13), the polynomial $x^{2^u n} + 1$ has the following factorization in $\mathbb{F}_{2^m}[x]$:

$$x^{2^u n} + 1 = (\varphi_1(x) \cdot \varphi_2(x) \cdots \varphi_r(x))^{2^u t}, \quad (17)$$

where $\varphi_j(x) \in \mathbb{F}_{2^m}[x]$ are distinct irreducible factors of $x^{n_1} + 1$.

For each $1 < s' < 2^u n$, let there is a $u'_{s'} > 0$ such that, $R'_{s',1}(x)$, $R'_{s',2}(x)$, \dots , $R'_{s',u'_{s'}}(x)$ be all of the s' degree polynomials dividing $x^{2^u n} + 1$. So, from (17) we can write $R'_{s',i'}(x) = \prod_{j=1}^r \varphi_j^{t'_{ij}}(x)$, for each $1 \leq i' \leq u'_{s'}$, $0 \leq t'_{ij} \leq 2^u t$. Let

$$H'_{s',i'}(x) = \frac{x^{2^u n} + 1}{R'_{s',i'}(x)} \quad (18)$$

for $1 \leq i' \leq u'_{s'}$. By Proposition 4.1, $F_{u+1}(x)$ is an N_1 -polynomial of degree $2^u n$ over \mathbb{F}_{2^m} if and only if, there is a j , $1 \leq j \leq u'_1$, such that

$$L_{H'_{1,j}}(\alpha_{u+1}) = 0,$$

and also

$$L_{H'_{s',i'}}(\alpha_{u+1}) \neq 0,$$

for each $1 < s' < 2^u n$, which there is a $u'_{s'} > 0$ such that, $R'_{s',1}(x)$, $R'_{s',2}(x)$, \dots , $R'_{s',u'_{s'}}(x)$ be all of the s' degree polynomials dividing $x^{2^u n} + 1$, and $1 \leq i' \leq u'_{s'}$.

Let $H_{s,i}(x) = \frac{x^{2^u n} + 1}{R_{s,i}(x)}$, for each $1 < s < n$, and $1 \leq i \leq u_s$, satisfied in the stated conditions. We note that by (14)

$$\begin{aligned}
H_{s,i}(x) &= \frac{x^{2^u n} + 1}{R_{s,i}(x)} \\
&= \frac{(x^n + 1)(\sum_{j=0}^{2^u-1} x^{jn})}{R_{s,i}(x)} \\
&= (\sum_{j=0}^{2^u-1} x^{jn}) \phi_{s,i}(x) \\
&= \sum_{v=0}^{n-s} t_{iv} (\sum_{j=0}^{2^u-1} x^{jn+v}).
\end{aligned} \tag{19}$$

It follows that

$$L_{H_{s,i}}(\alpha_{u+1}) = \sum_{v=0}^{n-s} t_{iv} (\sum_{j=0}^{2^u-1} \alpha_{u+1}^{2^{jmn} + 2^{mv}}). \tag{20}$$

From (12), if α_{y+1} is a root of $F_{y+1}(x)$, then $\alpha_{y+1} + \delta^2 \alpha_{y+1}^{-1}$ is a root of $F_y(x)$, and therefore, it may assume that

$$\alpha_{y+1} + \delta^2 \alpha_{y+1}^{-1} = \alpha_y \tag{21}$$

for all $1 \leq y \leq u$, where $\alpha_1 = \alpha$.

Now, by (21) and observing that $F_y(x)$ is an irreducible polynomial of degree $2^{y-1}n$ over \mathbb{F}_{2^m} , we obtain

$$\alpha_{y+1}^{2^{2^{y-1}mn}} + \delta^2 \alpha_{y+1}^{-2^{2^{y-1}mn}} = \alpha_y^{2^{2^{y-1}mn}} = \alpha_y. \tag{22}$$

So from (21) and (22) we have

$$(\alpha_{y+1}^{2^{2^{y-1}mn}} + \alpha_{y+1}) + \delta^2 (\alpha_{y+1}^{-2^{2^{y-1}mn}} + \alpha_{y+1}^{-1}) = 0.$$

Or

$$(\alpha_{y+1}^{2^{2^{y-1}mn}} + \alpha_{y+1})(1 + \frac{\delta^2}{\alpha_{y+1}^{2^{2^{y-1}mn}+1}}) = 0. \tag{23}$$

Also observing that $F_{y+1}(x)$ is an irreducible polynomial of degree $2^y n$ over \mathbb{F}_{2^m} , we have $\alpha_{y+1}^{2^{2^{y-1}mn}} + \alpha_{y+1} \neq 0$, and so $\alpha_{y+1}^{2^{2^{y-1}mn}} = \delta^2 \alpha_{y+1}^{-1}$. It follows that

$$\alpha_{y+1}^{2^{2^{y-1}mn}} + \alpha_{y+1} = \delta^2 \alpha_{y+1}^{-1} + \alpha_{y+1} = \alpha_y \tag{24}$$

for each y , $1 \leq y \leq u$. Hence by (24) we obtain

$$\begin{aligned}
\sum_{j=0}^{2^u-1} \alpha_{u+1}^{2^{jmn}} &= \sum_{j=0}^{2^{u-1}-1} (\alpha_{u+1}^{2^{2^{u-1}-1}mn} + \alpha_{u+1})^{2^{jmn}} = \sum_{j=0}^{2^{u-1}-1} \alpha_u^{2^{jmn}} \\
&= \cdots = \sum_{j=0}^{2^{u-z}-1} \alpha_{u-z+1}^{2^{jmn}} = \sum_{j=0}^{2^{u-z}-1} (\alpha_{u-z+1}^{2^{2^{u-z}-1}mn} + \alpha_{u-z+1})^{2^{jmn}} \\
&= \sum_{j=0}^{2^{u-z}-1} \alpha_{u-z}^{2^{jmn}} = \cdots = \sum_{j=0}^1 \alpha_2^{2^{jmn}} = \alpha_1 = \alpha.
\end{aligned}$$

Therefore, $\sum_{j=0}^{2^u-1} \alpha_{u+1}^{2^{jmn}} = \alpha$. By substituting α for $\sum_{j=0}^{2^u-1} \alpha_{u+1}^{2^{jmn}}$ in (20), we get

$$L_{H_{s,i}}(\alpha_{u+1}) = \sum_{v=0}^{n-s} t_{iv} \alpha^{2^{mv}} = L_{\phi_{s,i}}(\alpha) \quad (25)$$

for each s , $1 \leq s < n$, which there is a $u_s > 0$ such that, $R_{s,1}(x)$, $R_{s,2}(x)$, \cdots , $R_{s,u_s}(x)$ are all of the s degree polynomials dividing $x^n + 1$.

So, by (15) and (25), there is a j , $1 < j < u'_1 = u_1$, such that

$$L_{H'_{1,j}}(\alpha_{u+1}) = L_{H_{1,j}}(\alpha_{u+1}) = L_{\phi_{1,j}}(\alpha) = 0$$

Now we clime that, for each $1 < s' < 2^u n$, which there is a $u'_{s'} > 0$, such that, $R'_{s',1}(x)$, $R'_{s',2}(x)$, \cdots , $R'_{s',u'_{s'}}(x)$ be all of the s' degree polynomials dividing $x^{2^u n} + 1$, and $1 \leq i' \leq u'_{s'}$ we have

$$L_{H'_{s',i'}}(\alpha_{u+1}) \neq 0.$$

Setting $t'_{i'j} = t$ for each $t'_{i'j} > t$ in $R'_{s',i'}(x) = \prod_{j=1}^r \varphi_j^{t'_{i'j}}(x)$, we get

$$(R'_{s',i'}(x))|_{t'_{i'j}=t \text{ for } t'_{i'j} > t} = \prod_{j=1, t'_{i'j}=t \text{ for } t'_{i'j} > t}^r \varphi_j^{t'_{i'j}}(x) = \prod_{j=1}^r \varphi_j^{t_{i_0j}}(x) = R_{s_0,i_0}(x),$$

for some $1 < s_0 < n$ and $1 \leq i_0 \leq u_{s_0}$. Hence, $R_{s_0,i_0}(x)$ divide $R'_{s',i'}(x)$ and so $H'_{s',i'}(x)$ divide $H_{s_0,i_0}(x)$. Consequently, $L_{H'_{s',i'}}(\alpha_{u+1})$ divide $L_{H_{s_0,i_0}}(\alpha_{u+1})$. Therefore, by (16) and (25), the clime is true. The proof is completed. ■

References

- [1] L. Carlitz, Primitive roots in a finite field, Transactions of the American Mathematical Society, 73 (1952), 373-382.
- [2] L. Carlitz, Some problems involving primitive roots in a finite field, Proceedings of the National Academy of Sciences of the USA, 38 (1952), 314-318.

- [3] W. S. D. Cohen, Primitive elements and polynomials with arbitrary trace, *Discrete Math.* 83 (1990), 1-2.
- [4] S. D. Cohen and S. Huczynska, The primitive normal basis theorem without a computer, *Journal of the London Mathematical Society*, 67 (2003), 41-56.
- [5] S. Gao, Normal bases over finite fields, Ph.D. Thesis, Waterloo, (1993).
- [6] S. Huczynska, G. L. Mullen, D. Panario and D. Thomson, Existence and properties of k -normal elements over finite fields, *Finite Fields and Their Applications*, 24 (2013), 170-183.
- [7] M. K. Kyuregyan, Iterated construction of irreducible polynomials over finite fields with linearly independent roots, *Finite Fields Appl.* 10 (2004) 323-341.
- [8] H. W. Lenstra and R. Schoof, Primitive normal bases for finite fields, *Mathematics of Computation*, 48 (1987), 217-231.
- [9] R. Lidl and H. Niederreiter, *Finite Fields: Encyclopedia of Mathematics and its Applications*, Vol. 20 (2nd ed.), Cambridge University Press, (Cambridge) 1997.
- [10] A.J. Menezes, I.F.Blake, X.Gao, R.C.Mullin, S.A.Vanstone, T.Yaghoobian, *Applications of finite fields*, Kluwer Academic publishers, Boston, Dordrecht, Lancaster, 1993.
- [11] D. Pei, C.C. Wang and J.K. Omura, Normal bases of finite field $GF(2^m)$, *IEEE Trans. Info. Th.*, 32 (1986), 285-287.
- [12] S. Schwartz, Irreducible polynomials over finite fields with linearly independent roots, *Math. Slovaca*, Vol. 38, 1988, 147-158.