

Detecting tail behavior: mean excess plots with confidence bounds

BIKRAMJIT DAS and SOUVIK GHOSH

Singapore University of Technology and Design

20 Dover Drive, Singapore 138682

E-mail: bikram@sutd.edu.sg

LinkedIn Corporation

2029 Stierlin Court, Mountain View, CA 94043, USA

E-mail: sghosh@linkedin.com

In many practical situations exploratory plots are helpful in understanding tail behavior of sample data. The Mean Excess plot is often applied in practice to understand the right tail behavior of a data set. It is known that if the underlying distribution of a data sample is in the domain of attraction of a Fréchet, Gumbel or Weibull distributions then the ME plot of the data tend to a straight line in an appropriate sense, with positive, zero or negative slopes respectively. In this paper we construct confidence intervals around the ME plots which assist us in ascertaining which particular maximum domain of attraction the data set comes from. We recall weak limit results for the Fréchet domain of attraction, already obtained in [Das and Ghosh \(2013\)](#) and derive weak limits for the Gumbel and Weibull domains in order to construct confidence bounds. We test our methods on both simulated and real data sets.

AMS 2000 subject classifications: Primary 62G32, 62-09, 60G70; secondary 62G10, 62G15, 60F05.

Keywords: extreme values, regular variation, random set, ME plot, asymptotic theory, confidence bounds.

1. Introduction

This article concerns the use of mean excess plots, a popular exploratory tool used to understand the tail behavior of a univariate data set. Given a sample of data points, one of the first things a sensible data analyst does is to compute a summary statistics. Such a summary statistic might involve calculating measures of central tendencies (mean, median, mode) and those of dispersion (standard deviation, range, etc), plotting a histogram, an empirical cumulative distribution function and so on and so forth. A more curious analyst would ask the question, does it even make sense to calculate the sample mean or standard deviation; would they represent their counterparts in the original population? What if the probability distribution of the population from which the data is sampled does not even have a first or second moment. This is a question that would or perhaps should particularly come to the mind of analysts modeling risk or other extreme events. In a world where data is being used to make serious economic, financial or environmental policy decisions, understanding extreme risks, which relate to the tail behavior of data sets have become increasingly important. This can be easily observed in the world of finance and insurance ([Das, Embrechts and Fasen, 2013](#); [Donnelly and Embrechts, 2010](#); [McNeil, Frey and Embrechts, 2005](#)), telecommunications ([Maulik, Resnick and Rootzén, 2002](#)), environmental statistics ([Davison and Smith, 1990](#)) and many more areas.

The mean excess(ME) plot is a graphical tool that is widely used to understand the tail behavior of a sample; see [Embrechts, Klüppelberg and Mikosch \(1997\)](#); [Davison and Smith \(1990\)](#). A Mean Excess plot, if the mean exists, assists in distinguishing light-tailed data sets from heavy-tailed ones. The inference is based on a visual examination of the slope of a fitted line through the ME plot (to be described in the next section) being zero, less than zero or greater than zero. Clearly, a confidence set around the fitted line would make inference in these cases more meaningful; hence this is the aim of the paper.

*Bikramjit Das gratefully acknowledges Support from SRG ESD 047 and MIT-SUTD IDC grant IDG31300110.

1.1. The ME plot

The ME plot, as described in the introduction, is a popular tool in extreme value analysis. It is a simple graphical test to check whether data conform to a generalised Pareto distribution (GPD). The class of GPD arise naturally in extreme value analysis as limit distributions while using the peaks-over-threshold (POT) method (Beirlant et al., 2004; Davison and Smith, 1990). The cumulative distribution function of a GPD is:

$$G_{\xi,\beta}(x) = \begin{cases} 1 - (1 + \xi x/\beta)^{-1/\xi} & \text{if } \xi \neq 0, \\ 1 - \exp(-x/\beta) & \text{if } \xi = 0, \end{cases} \quad (1.1)$$

where $\beta > 0$, and $x \geq 0$ when $\xi \geq 0$ and $0 \leq x \leq -\beta/\xi$ if $\xi < 0$. Parameters ξ and β are referred to as the *shape* and the *scale* parameter respectively. In extreme value analysis, we are interested in the shape parameter ξ which tells us whether the data is heavy-tailed ($\xi > 0$) or light-tailed ($\xi \leq 0$) or even more specifically if the underlying distribution has finite right end-point ($\xi < 0$). The case $\xi > 0$ and $\beta = 1$ corresponds to the classical Pareto law with tail exponent $1/\xi$.

The ME plot is an empirical graphical plot of the *ME function* of a random variable $X \sim F$ which is defined as:

$$M(u) := \mathbb{E}[X - u | X > u], \quad (1.2)$$

provided $\mathbb{E}X_+ < \infty$. The ME function is also known as the *mean residual life function* for non-negative random variables and is extensively used in reliability theory and survival analysis for data modelling since $M(u)$ completely determines F if $\mathbb{E}(X) < \infty$ (Hall and Wellner, 1981). Suppose we have an iid sample $X_1, \dots, X_n \sim F$. A natural estimate of $M(u)$ is the empirical ME function $\hat{M}(u)$ defined as

$$\hat{M}(u) = \frac{\sum_{i=1}^n (X_i - u) I_{[X_i > u]}}{\sum_{i=1}^n I_{[X_i > u]}}, \quad u \geq 0. \quad (1.3)$$

Denoting $X_{(1)} \geq X_{(2)} \geq \dots \geq X_{(n)}$ to be the order statistics from a sample X_1, \dots, X_n , the ME plot is a plot of the points

$$\mathcal{ME}_n := \{(X_{(k)}, \hat{M}(X_{(k)})) : 1 < k \leq n\}.$$

We study the asymptotic properties of \mathcal{ME}_n for different classes of distributions F . It is well-known that for a random variable $X \sim G_{\xi,\beta}$, we have $E(X) < \infty$ if and only if $\xi < 1$ and in this case, the ME function of X is linear in u :

$$M(u) = \frac{\beta}{1 - \xi} + \frac{\xi}{1 - \xi} u, \quad (1.4)$$

where $0 \leq u < \infty$ if $0 \leq \xi < 1$ and $0 \leq u \leq -\beta/\xi$ if $\xi < 0$.

Interestingly, the linearity of the ME function characterises the GPD class (McNeil, Frey and Embrechts, 2005; Embrechts, Klüppelberg and Mikosch, 1997). From the discussions in Ghosh and Resnick (2010) we have learnt that the empirical ME plot \mathcal{ME}_n above a high order statistics $X_{(k)}$ when appropriately normalised converge in probability to a straight line if F is in the maximal domain of attraction of any generalized extreme value distribution with finite mean (Gumbel, Weibull or Fréchet distribution). Distributional limits for \mathcal{ME}_n in a space of closed sets and confidence intervals around \mathcal{ME}_n can also be computed in many cases and such findings have been discussed in case the underlying data is heavy-tailed (Fréchet domain of attraction) in Das and Ghosh (2013).

In Section 1.2 we collect notations and ideas to be used throughout the paper. See Das and Ghosh (2013) for further elaboration of the concepts of convergence of closed sets (random) in this context. In the main part of the paper we start by consolidating a few results which are already known on the distributional property of ME plot, especially in the heavy-tailed case (Das and Ghosh, 2013); this is covered in Section 2.1. The rest of Section 2 deals with limit results for ME plots in the case where the underlying distribution is either in the Gumbel maximum domain of attraction or in a Weibull maximum domain of attraction. The limit theorems proved in Section 2 is used to create confidence bounds around the ME plots in Section 3. In Section 4 we use the tools developed in Section 2 and 3 to test it out both on simulated data as well as real data sets.

1.2. Miscellany

First we recall the idea of maximum domain of attraction of an extreme value distribution. The class of extreme value distributions is parametrized by a shape parameter $\xi \in \mathbb{R}$ and we define the distribution function G_ξ to be

$$G_\xi(x) = \exp(-(1 + \xi x)^{-1/\xi}), \quad 1 + \xi x > 0,$$

for all real ξ and for $\xi = 0$, the right hand side is interpreted as $\exp(-e^{-x})$.

Definition 1.1. *A distribution function F (or the underlying random variable $X \sim F$) is in the maximum domain of attraction of an extreme value distribution G_ξ if there exists sequences $c_n > 0$ and $d_n \in \mathbb{R}$ such that*

$$F^n(c_n x + d_n) \rightarrow G_\xi(x) \quad \text{for all } x \in \mathbb{R}.$$

The distributions for the cases $\xi > 0$, $\xi = 0$ and $\xi < 0$ are respectively called the Fréchet distribution, the Gumbel distribution and the Weibull distribution. As mentioned in the introduction, if $F \in D(G_\xi)$ for some extreme value distribution with $\xi < 1$, implying that F has finite mean, then the ME function of F is linear with an appropriate slope determined by the parameter ξ ; see [Ghosh and Resnick \(2010\)](#).

Throughout this paper we will take $k := k_n$ to be a sequence increasing to infinity such that $n/k_n \rightarrow \infty$ or $k_n/n \rightarrow 0$. For a distribution function $F(x)$ we write $\bar{F}(x) := 1 - F(x)$ for the tail and the quantile function is

$$b(u) := F^{\leftarrow}(1 - \frac{1}{u}) := \inf \left\{ s : F(s) \geq 1 - \frac{1}{u} \right\} = \left(\frac{1}{1 - F} \right)^{\leftarrow}(u).$$

A function $U : (0, \infty) \rightarrow \mathbb{R}_+$ is regularly varying with index $\rho \in \mathbb{R}$, written $U \in RV_\rho$, if

$$\lim_{t \rightarrow \infty} \frac{U(tx)}{U(t)} = x^\rho, \quad x > 0.$$

If $X \sim F$ we will often have the right-hand tail of F to be regularly varying, that is, $\bar{F} \in RV_{-\alpha}$ for $\alpha \geq 0$, and by abuse of notation we might say $X \in RV_{-\alpha}$. Regular variation is discussed in several books such as [Resnick \(2007, 2008\)](#); [Seneta \(1976\)](#); [Geluk and de Haan \(1987\)](#); [de Haan \(1970\)](#); [de Haan and Ferreira \(2006\)](#); [Bingham, Goldie and Teugels \(1987\)](#).

We use $\mathbb{M}_+(0, \infty]$ to denote the space of nonnegative Radon measures μ on $(0, \infty]$ metrized by the vague metric. Point measures are written as a function of their points $\{x_i, i = 1, \dots, n\}$ by $\sum_{i=1}^n \delta_{x_i}$. See, for example, [\(Resnick, 2008, Chapter 3\)](#).

We will use the following notations to denote different classes of functions: For $0 \leq a < b \leq \infty$,

1. $\mathbb{C}[a, b)$: Continuous functions on $[a, b)$.
2. $\mathbb{D}[a, b)$: Right-continuous functions with finite left limits defined on $[a, b)$.
3. $\mathbb{D}_l[a, b)$: Left-continuous functions with finite right limits defined on $[a, b)$.

It is known that $\mathbb{D}[0, 1]$ is complete and separable under a metric $d_0(\cdot)$ which is equivalent to the Skorohod metric $d_S(\cdot)$ ([Billingsley, 1968](#), p.128), but not under the uniform metric $\|\cdot\|$. As we will see, the limit processes that appear in our analysis below are always continuous. We can check that if x is continuous (in fact uniformly continuous) in $[0, 1]$, for $x_n \in \mathbb{D}[0, 1]$, $\|x_n - x\| \rightarrow 0$ is equivalent to $d_S(x_n, x) \rightarrow 0$ and hence equivalent to $d_0(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$ ([Billingsley, 1968](#), p.124). So we use convergence in uniform metric, for our convenience henceforth. For spaces of the form $\mathbb{D}[a, b)$ or $\mathbb{D}_l[a, b)$ we will consider the topology of local uniform convergence. In some cases we will also consider product spaces of functions and then the topology will be the product topology. For example, $\mathbb{D}_l^2[1, \infty)$ will denote the class of 2-dimensional functions on $[1, \infty)$ which are left-continuous with right limit. The classes of functions defined on the sets $[a, b]$ or $(a, b]$ will have the obvious notation. For further details on notions of convergence and topology for convergences of plots see [Das and Ghosh \(2013\)](#).

2. Limit results for the ME Plots

In this section we find distributional limits for ME plots when it exists. We continue the study of ME plots from [Ghosh and Resnick \(2010\)](#) and [Das and Ghosh \(2013\)](#) and give a complete picture of limit results for ME plots. We cite some of the results from the afore-mentioned papers for completeness.

The basic assumption is that we have an iid sample of data points from some unknown distribution F which belongs to the maximum domain of attraction of one of the three extreme value distributions. The assumption of independence in the sample can be relaxed a bit under certain conditions which we do not explore here.

Suppose X_1, \dots, X_n is an i.i.d. sample from a distribution F . We will work under this assumption for the entire section. The properties of the empirical ME function $\hat{M}(u)$ as an estimator of $M(u)$ has been studied by [Yang \(1978\)](#). It was shown there that $\hat{M}(u)$ is uniformly strongly consistent for $M(u)$: for any $0 < b < \infty$

$$P \left[\lim_{n \rightarrow \infty} \sup_{0 \leq u \leq b} |\hat{M}(u) - M(u)| = 0 \right] = 1.$$

A weak (distributional) limit for $\hat{M}(u)$ was also shown in [Yang \(1978\)](#): for any $0 < b < 1$

$$\sqrt{n} \left(\hat{M}(F^{\leftarrow}(t)) - M(F^{\leftarrow}(t)) \right) \Rightarrow U(t),$$

where $U(t)$ is a Gaussian process on $[0, b]$ with covariance function

$$\Gamma(s, t) = \frac{(1-t)\sigma^2(t) - t\theta^2(t)}{(1-s)(1-t)^2} \quad \text{for all } 0 \leq s \leq t \leq b$$

with

$$\sigma^2(t) = \text{var} \left(X I_{[t < F(X) \leq 1]} \right) \quad \text{and} \quad \theta(t) = E \left(X I_{[t < F(X) \leq 1]} \right).$$

Using Lemma 2.4 in [Das and Ghosh \(2013\)](#) it is easy to see that the ME plots also exhibit the same features. Our interest in ME plots is for detecting right tail behavior of data samples (an equivalent case can be easily made for left tail behavior). Hence the linearity we seek in the ME plot will be for high thresholds. Necessarily, the ME plots we will discuss in the various cases will be transformations of the ME plot over an appropriate quantile, i.e., $\{(X_{(i)}, \hat{M}(X_{(i)})) : 1 < i \leq k\}$ for $k := k(n) < n$ where \hat{M} is as defined in (1.3).

2.1. ME plot in the Fréchet case

First we look at the case where the underlying distribution F is heavy-tailed, in the sense that $F \in D(G_\xi)$ with $\xi > 0$ or in other words, $\bar{F} \in RV_{-1/\xi}$. We define the ME plot as:

$$\mathcal{M}_n := \frac{1}{X_{(k)}} \left\{ \left(X_{(i)}, \hat{M}(X_{(i)}) \right) : i = 2, \dots, k \right\}. \quad (2.1)$$

From ([Ghosh and Resnick, 2010](#), Theorem 3.2), we know that for $0 < \xi < 1$, as $n, k, n/k \rightarrow \infty$,

$$\mathcal{M}_n \xrightarrow{P} \mathcal{M} := \left\{ \left(t, \frac{\xi}{1-\xi} t \right) : t > 1 \right\}$$

The distributional behavior of \mathcal{M}_n depends on whether F has finite second moment or not and has been discussed under certain regularity conditions in [Das and Ghosh \(2013\)](#). We note them down below.

Case 1 ($0 < \xi < 1/2$): For any $0 < \epsilon < 1$ as $n, k, n/k \rightarrow \infty$,

$$\mathcal{MN}_n := \left\{ \left(\left(\frac{i}{k} \right)^{-\xi}, \frac{\xi}{1-\xi} \left(\frac{i}{k} \right)^{-\xi} \right) \right\}$$

$$\begin{aligned}
& + \sqrt{k} \left(\frac{X_{(i)}}{X_{(k)}} - \left(\frac{i}{k} \right)^{-\xi}, \frac{\hat{M}(X_{(i)})}{X_{(k)}} - \frac{\xi}{1-\xi} \left(\frac{i}{k} \right)^{-\xi} \right) : i = \lceil \epsilon k \rceil, \dots, k \Big\} \\
\Rightarrow \mathcal{MN} &:= \left\{ \left(t^{-\xi} + \xi t^{-(1+\xi)} B(t), \frac{\xi}{1-\xi} t^{-\xi} + \xi t^{-1} \int_0^t y^{-(1+\xi)} B(y) dy \right), \epsilon \leq t \leq 1 \right\} \quad \text{in } \mathcal{F}, \quad (2.2)
\end{aligned}$$

where $B(t)$ is the standard Brownian bridge on $[0, 1]$. This is the case where F has a finite second moment and hence the distributional limit has a Brownian component.

Case 2 ($1/2 < \xi < 1$): For any $0 < \epsilon < 1$, as $n, k, n/k \rightarrow \infty$,

$$\begin{aligned}
\mathcal{MN}_n &:= \left\{ \left(\left(\frac{i}{k} \right)^{-\xi}, \frac{\xi}{1-\xi} \left(\frac{i}{k} \right)^{-\xi} \right) \right. \\
& \quad \left. + \left(\sqrt{k} \left(\frac{X_{(i)}}{X_{(k)}} - \left(\frac{i}{k} \right)^{-\xi} \right), \frac{kb(n/k)}{b(n)} \left(\frac{\hat{M}(X_{(i)})}{X_{(k)}} - \frac{\xi}{1-\xi} \left(\frac{i}{k} \right)^{-\xi} \right) \right) : i = \lceil \epsilon k \rceil, \dots, k \right\} \\
\Rightarrow \mathcal{MN} &:= \left\{ \left(t^{-\xi} + \xi t^{-(1+\xi)} B(t), \frac{\xi}{1-\xi} t^{-\xi} + t^{-1} S_{1/\xi} \right), \epsilon \leq t \leq 1 \right\} \quad \text{in } \mathcal{F}, \quad (2.3)
\end{aligned}$$

where $S_{1/\xi}$ is a stable random variable with characteristic function

$$E[e^{itS_{1/\xi}}] = \exp \left\{ -\frac{1}{1-\xi} \Gamma \left(2 - \frac{1}{\xi} \right) \cos \frac{\pi}{2\xi} |t|^{1/\xi} \left[1 - i \operatorname{sgn}(t) \tan \frac{\pi}{2\xi} \right] \right\}, \quad (2.4)$$

and is independent of the standard Brownian Bridge $B(t)$ on $[0, 1]$. This is the case where F has a finite mean but does not have a finite second moment, hence we also observe a non-Gaussian stable weak limit. The results in (2.2) and (2.3) are described in Theorems 4.3 and 4.6 in details in [Das and Ghosh \(2013\)](#).

2.2. ME plot in the Gumbel case

The behaviour (in probability) of ME plot when F is in the maximum domain of attraction of a Gumbel distribution has been discussed in [Ghosh and Resnick \(2010\)](#). We state the following result to recall notations to be used: this follows from Theorems 3.3.26 and 3.4.13(b) in [Embrechts, Klüppelberg and Mikosch \(1997\)](#); see ([Ghosh and Resnick, 2010](#), Theorem 3.9) or ([Resnick, 2008](#), Proposition 1.4) for further details.

Proposition 2.1. *The following are equivalent for a distribution function F with right end point $x_F \leq \infty$:*

1. *F is in the maximum domain of attraction of the Gumbel distribution, i.e.,*

$$F^n(c(n)x + d(n)) \rightarrow \exp \{ -e^{-x} \} \quad \text{for all } x \in \mathbb{R}, \quad (2.5)$$

for some sequence $c(n)$ and $d(n)$.

2. *There exists $z < x_F$ such that F has a representation*

$$\bar{F}(x) = \kappa(x) \exp \left\{ - \int_z^x \frac{1}{a(t)} dt \right\}, \quad \text{for all } z < x < x_F, \quad (2.6)$$

where $\kappa(x)$ is a measurable function satisfying $\kappa(x) \rightarrow \kappa > 0$, $x \rightarrow x_F$, and $a(x)$ is a positive, absolutely continuous function with density $a'(x) \rightarrow 0$ as $x \rightarrow x_F$.

We know from ([Resnick, 2008](#), Proposition 1.1) that a choice of the norming sequence $c(n)$ and $d(n)$ in (2.5) is

$$d(n) = F^{\leftarrow}(1 - n^{-1}) \quad \text{and} \quad c(n) = a(d(n)).$$

Theorem 3.3.26 in [Embrechts, Klüppelberg and Mikosch \(1997\)](#) says that a choice of the auxiliary function $a(x)$ in (2.6) is

$$a(x) = \int_x^{x_F} \frac{\bar{F}(t)}{\bar{F}(x)} dt \quad \text{for all } x < x_F,$$

and for this choice, the auxiliary function is the ME function, i.e., $a(x) = M(x)$. Furthermore, we also know that $a'(x) \rightarrow 0$ as $x \rightarrow x_F$ and this implies that $M(u)/u \rightarrow 0$ as $u \rightarrow x_F$. Define the ME plot in this case as

$$\mathcal{M}_n := \frac{1}{X_{\lceil k/e \rceil} - X_{(k)}} \left\{ \left(X_{(i)} - X_{(k)}, \hat{M}(X_{(i)}) \right) : i = 2, \dots, k \right\}. \quad (2.7)$$

From a minor modification of ([Ghosh and Resnick, 2010](#), Theorem 3.10), we know that as $n, k, n/k \rightarrow \infty$, $\mathcal{M}_n \xrightarrow{P} \mathcal{M} := \{(t, 1) : t > 0\}$. Now we will additionally put one more condition in order to get a weak limit for ME plots in the Gumbel case which is stated as follows.

Assumption 2.2. *The distribution function F satisfies the following:*

$$\sqrt{k} \left(\frac{n}{k} \bar{F}(c(n/k)y + d(n/k)) - e^{-y} \right) \rightarrow 0 \quad (2.8)$$

point-wise and in L_1 -norm in $[0, \infty)$ as $n, k, n/k \rightarrow \infty$.

Now we can state the distributional result for ME plots when F is in the maximum domain of attraction of the Gumbel distribution.

Theorem 2.3. *Suppose X_1, \dots, X_n are i.i.d. observations from a distribution F which is in the maximum domain of attraction of the Gumbel distribution and satisfies Assumption 2.2. Then for any $0 < \epsilon < 1$, as $n, k, n/k \rightarrow \infty$,*

$$\begin{aligned} \mathcal{MN}_n &:= \left\{ \left(-\ln \left(\frac{i}{k} \right), 1 \right) + \sqrt{k} \left(\frac{X_{(i)} - X_{(k)}}{X_{\lceil k/e \rceil} - X_{(k)}} + \ln \left(\frac{i}{k} \right), \frac{\hat{M}(X_{(i)})}{X_{\lceil k/e \rceil} - X_{(k)}} - 1 \right) : i = \lceil \epsilon k \rceil, \dots, k \right\} \\ &\Rightarrow \mathcal{MN} := \left\{ \left(-\ln(t) + eB(e^{-1})\ln(t) + \frac{B(t)}{t}, 1 + eB(e^{-1}) + \frac{1}{t} \int_0^t \frac{B(s)}{s} ds \right), \epsilon \leq t \leq 1 \right\} \quad \text{in } \mathcal{F}, \end{aligned}$$

where $B(t)$ is the standard Brownian bridge on $[0, 1]$.

Proof. The proof is along the same lines of the proof of Theorem 4.3 in [Das and Ghosh \(2013\)](#). Denote the tail empirical measures by

$$\nu_n(\cdot) := \frac{1}{k} \sum_{i=1}^n \varepsilon_{\frac{X_i - d(n/k)}{c(n/k)}}(\cdot) \quad \text{and}, \quad (2.9)$$

$$\hat{\nu}_n(\cdot) := \frac{1}{k} \sum_{i=1}^n \varepsilon_{\frac{X_i - X_{(k)}}{c(n/k)}}(\cdot) \quad (2.10)$$

and define for $k := k(n) < n$ and $0 < t \leq 1$:

$$W_n(t) := \sqrt{k}(\nu_n(-\ln t, \infty] - t) = \sqrt{k} \left(\frac{1}{k} \sum_{i=1}^n \varepsilon_{\frac{X_i - d(n/k)}{c(n/k)}}(-\ln t, \infty] - t \right).$$

We prove in Lemma 2.4 that $W_n \Rightarrow W$ in $D_l(0, 1]$, where W is the standard Brownian motion in $[0, 1]$. Applying Vervaat's lemma ([Resnick, 2007](#), Proposition 3.3, p.59) to (2.2) we get

$$\sqrt{k} \left(\exp \left\{ -\frac{X_{\lceil kt \rceil} - d(n/k)}{c(n/k)} \right\} - t, \nu_n(-\ln t, \infty] - t \right) \Rightarrow (-W(t), W(t)) \quad \text{in } \mathbb{D}_l^2(0, 1].$$

Using the Functional Delta-method ([van der Vaart and Wellner, 1996](#), Theorem 3.9.4) we get

$$\sqrt{k} \left(\frac{X_{(\lceil kt \rceil)} - d(n/k)}{c(n/k)} + \ln t, \nu_n(y, \infty] - e^{-y} \right) \Rightarrow \left(\frac{W(t)}{t}, W(e^{-y}) \right) \quad \text{in } \mathbb{D}_l(0, 1] \times \mathbb{D}[1, \infty), \quad (2.11)$$

and it is easy to check that

$$\sqrt{k} \left(\frac{X_{(\lceil kt \rceil)} - X_{(k)}}{c(n/k)} + \ln t, \hat{\nu}_n(y, \infty] - e^{-y} \right) \Rightarrow \left(\frac{B(t)}{t}, B(e^{-y}) \right) \quad \text{in } \mathbb{D}_l(0, 1] \times \mathbb{D}[1, \infty). \quad (2.12)$$

Then following arguments used in the proof of Theorem 4.3 in [Das and Ghosh \(2013\)](#) we get

$$\begin{aligned} & \sqrt{k} \left(\frac{X_{(\lceil kt \rceil)} - d(n/k)}{c(n/k)} + \ln t, \frac{\hat{M}(X_{(\lceil kt \rceil)})}{c(n/k)} - 1 \right) \\ &= \sqrt{k} \left(\frac{X_{(\lceil kt \rceil)} - d(n/k)}{c(n/k)} + \ln t, \frac{k}{(\lceil kt \rceil - 1)c(n/k)} \int_{\frac{X_{(\lceil kt \rceil)} - d(n/k)}{c(n/k)}}^{\infty} \nu_n(y, \infty] dy - 1 \right) \\ &\Rightarrow \left(\frac{B(t)}{t}, \frac{1}{t} \int_{-\ln t}^{\infty} B(e^{-y}) dy \right) = \left(\frac{B(t)}{t}, \frac{1}{t} \int_0^t \frac{B(s)}{s} ds \right). \end{aligned}$$

The proof the theorem is completed by invoking Lemma 2.4 in [Das and Ghosh \(2013\)](#). \square

Lemma 2.4. As $n \rightarrow \infty, k \rightarrow \infty, n/k \rightarrow \infty$,

$$W_n \Rightarrow W$$

in $D_l(0, 1]$ where W is a Brownian motion in $D_l(0, 1]$.

Proof. We check the conditions C1-C4 of ([Rootzén, 2009](#), Theorem 2.1). In this part of the proof whenever we write ‘ \sim ’ between two expressions, it means the asymptotics hold for $n, k, n/k \rightarrow \infty$. Now following the notations used in the aforementioned paper, we set $r_n = \min\{k^{1/4}, (n/k)^{1/2}\}$ and $l_n = 1$. For any $u, v \in \mathbb{R}$ let

$$N_n(u, v) := \sum_{i=1}^{r_n} \varepsilon_{\frac{X_i - d(n/k)}{c(n/k)}}(u, v].$$

Then for any $\theta < x_T = \infty$ (since F is in a Gumbel domain of attraction, it has right end point $x_T = \infty$) with $0 \leq u, v < \theta$ we have,

$$\mathbb{P}[N_n(u, v) \neq 0] \sim r_n \mathbb{P}[u c(n/k) + d(n/k) < X_1 \leq v c(n/k) + d(n/k)]$$

and

$$\begin{aligned} \mathbb{E}[N_n(u, v)^2 | N_n(u, v) \neq 0] &= \frac{\mathbb{E}[N_n(u, v)^2]}{\mathbb{P}[N_n(u, v) \neq 0]} \\ &\sim 1 + r_n \mathbb{P}[u c(n/k) + d(n/k) < X_1 \leq v c(n/k) + d(n/k)] \\ &\leq 1 + \text{const.} r_n k/n \end{aligned}$$

which is bounded by the choice of r_n . Hence condition C1 holds. Condition C2 holds as the random variables X_i are assumed to be independent. Next note that for any $0 \leq u, v < \infty$,

$$\frac{1}{r_n \bar{F}(d(n/k))} \text{Cov} \left(\sum_{i=1}^{r_n} \varepsilon_{\frac{X_i - d(n/k)}{c(n/k)}}(u, \infty], \sum_{i=1}^{r_n} \varepsilon_{\frac{X_i - d(n/k)}{c(n/k)}}(v, \infty] \right)$$

$$\begin{aligned}
&= \frac{1}{\bar{F}(d(n/k))} \text{Cov} \left(\varepsilon_{\frac{x_1 - d(n/k)}{c(n/k)}}(u, \infty], \varepsilon_{\frac{x_1 - d(n/k)}{c(n/k)}}(v, \infty] \right) \\
&\sim \frac{\bar{F}((u \vee v)c(n/k) + d(n/k))}{\bar{F}(d(n/k))} \\
&\rightarrow \exp(-u \vee v).
\end{aligned}$$

Hence C3 holds and obviously C4 holds because of the choice of r_n . Hence, by (Rootzén, 2009, Theorem 2.1)

$$\sqrt{k}(\nu_n(u, \infty] - \mathbb{E}(\nu_n(u, \infty))) \Rightarrow G \quad \text{in } D[0, \infty),$$

where G is a centered Gaussian process in $[0, \infty)$ with covariance function $\exp(-u \vee v)$ and hence a time change $u \mapsto -\ln u$ gives us that $W_n \Rightarrow W$ in $D_l(0, 1]$ where W is a standard Brownian Motion on $[0, 1]$. \square

2.3. ME plot in the Weibull case

If $F \in D(G_{\xi, \beta})$, then we have the following characterizations for the case $\xi < 0$ (Embrechts, Klüppelberg and Mikosch, 1997; Ghosh and Resnick, 2010):

Proposition 2.5. *If $\xi < 0$ then the following are equivalent:*

1. F has a finite right end point x_F and $\bar{F}(x_F - x^{-1}) \in RV_{1/\xi}$.
2. $F^n(x_F + c(n)x) \rightarrow \exp\{-(-x)^{-1/\xi}\}$ for all $x \leq 0$ where $c(t) = x_F - F(1 - \frac{1}{t})$, $t \geq 1$.
3. There exists a measurable function $\beta(u)$ such that

$$\lim_{u \rightarrow x_F} \sup_{u \leq x \leq x_F} |F_u(x) - G_{\xi, \beta(u)}(x)| = 0.$$

Recall from Ghosh and Resnick (2010), the following result on ME plots (there is a typographical error in the statement of the result there):

Proposition 2.6. *If X_1, \dots, X_n are i.i.d. observations with distribution F which has a finite right end point x_F and satisfies $1 - F(x_F - x^{-1}) \in RV_{1/\xi}$ as $x \rightarrow \infty$, then in \mathcal{F} ,*

$$\mathcal{M}_n := \frac{1}{X_{(1)} - X_{(k)}} \left\{ (X_{(i)} - X_{(k)}, \hat{M}(X_{(i)})) : i = 2, \dots, k \right\} \xrightarrow{P} \mathcal{M} := \left\{ \left(t, \frac{\xi}{1 - \xi}(1 - t) \right) : 0 \leq t \leq 1 \right\}. \quad (2.13)$$

In this paper we obtain the weak limit of the ME plot when the null hypothesis that $\bar{F}(x_F - x^{-1}) \in RV_{1/\xi}$ for some $\xi < 0$ holds. In the same spirit as Das and Ghosh (2013) we deal with the tail empirical process. Denote by ν_n :

$$\nu_n(\cdot) := \frac{1}{k} \sum_{i=1}^n \varepsilon_{\frac{x_F - X_i}{c(n/k)}}. \quad (2.14)$$

Following Theorem 4.2 in Resnick (2007), we can show that

$$\nu_n \Rightarrow \nu \quad \text{in } \mathbb{M}_+[0, \infty)$$

where $\nu[0, x] = x^{-1/\xi}$, $x \geq 0$. Now define for $k := k(n) < n$ and $y \geq 0$:

$$W_n(y) := \sqrt{k} \left(\frac{1}{k} \sum_{i=1}^n \varepsilon_{\frac{x_F - X_i}{c(n/k)}}[0, y^{-\xi}] - \frac{n}{k} \bar{F}(x_F - c(n/k)y^{-\xi}) \right) \quad (2.15)$$

$$= \sqrt{k} (\nu_n[0, y^{-\xi}] - \mathbb{E}(\nu_n[0, y^{-\xi}])). \quad (2.16)$$

The next result in the spirit of (Resnick, 2007, Theorem 9.1) and also similar to Lemma 2.4.

Lemma 2.7. As $n \rightarrow \infty, k \rightarrow \infty, n/k \rightarrow \infty$,

$$W_n \Rightarrow W$$

in $D[0, \infty)$ where W is a Brownian motion in $D[0, \infty)$.

The proof follows by going through the steps of the proof of Lemma 2.4 or (Resnick, 2007, Theorem 9.1). Let us also assume the following:

Assumption 2.8. F satisfies the following

$$[1] \quad \sqrt{k} \left(\frac{n}{k} \bar{F}(x_F - c(n/k)y) - y^{-1/\xi} \right) \rightarrow 0 \quad \text{for all } y \geq 0,$$

$$[2] \quad \sqrt{k} \int_0^1 \left| \frac{n}{k} \bar{F}(x_F - c(n/k)y) - y^{-1/\xi} \right| dy \rightarrow 0.$$

as $n, k, n/k \rightarrow \infty$.

Theorem 2.9. Suppose X_1, \dots, X_n are i.i.d. observations from a distribution F which has a finite right end point x_F and satisfies $1 - F(x_F - x^{-1}) \in RV_{1/\xi}, \xi < 0$ as $x \rightarrow \infty$ and Assumption 2.8 holds. Then for any $0 < \epsilon < 1$, as $n, k, n/k \rightarrow \infty$,

$$\begin{aligned} \mathcal{MN}_n &:= \left\{ \left(1 - \left(\frac{i}{k} \right)^{-\xi}, \frac{\xi}{\xi - 1} \left(\frac{i}{k} \right)^{-\xi} \right) \right. \\ &\quad \left. + \sqrt{k} \left(\frac{X_{(i)} - X_{(k)}}{X_{(1)} - X_{(k)}} - \left(1 - \left(\frac{i}{k} \right)^{-\xi} \right), \frac{\hat{M}(X_{(i)})}{X_{(1)} - X_{(k)}} - \frac{\xi}{\xi - 1} \left(\frac{i}{k} \right)^{-\xi} \right) : i = \lceil \epsilon k \rceil, \dots, k \right\} \\ &\Rightarrow \mathcal{MN} := \left\{ \left(1 - t^{-\xi} + \xi t^{-(1+\xi)} B(t), \frac{\xi}{\xi - 1} t^{-\xi} + \xi t^{-1} \int_0^t y^{-(1+\xi)} B(y) dy \right), \epsilon \leq t \leq 1 \right\} \quad \text{in } \mathcal{F}, \end{aligned}$$

where $B(t)$ is the standard Brownian bridge on $[0, 1]$ restricted to $(0, 1]$.

Remark 2.10. This result is similar to the one obtained for $\bar{F} \in RV_{-1/\xi}$ or $\xi > 0$ in Theorem 4.3 of Das and Ghosh (2013); the subtle difference appears in the fact that we no longer need to restrict the range of ξ as is done there with $0 < \xi < 1/2$, since the integral

$$\int_0^t y^{-(1+\xi)} B(y) dy \stackrel{d}{=} \int_0^t y^{-(1+\xi)} W(y) dy - W(1) \int_0^t y^{-\xi} dy$$

exists if and only if $\int_0^t s^{-2\xi} ds < \infty$ which is always true for $\xi < 0$ and in turn implies that the limit \mathcal{MN} exists. The truncation with ϵ with $\epsilon \leq t \leq 1$ is still necessary to guarantee that the limit set \mathcal{MN} does not blow up for t near 0.

Proof. The proof is omitted here as it follows using similar arguments as in the proof of (Das and Ghosh, 2013, Theorem 4.3). The difference occurs in the fact that we use the weak convergence result mentioned in Lemma 2.7 as our basis and apply a proper version of Vervaat's Lemma and 'converging together' arguments on this to obtain the result. \square

3. Creating confidence bounds from the limit results

In Section 2 we obtain weak limits for ME plots for different values of $\xi \in \mathbb{R}$ where the underlying distribution $F \in D(G_\xi)$. Now, depending on varying values of ξ we construct the different confidence bounds following the results. We resort to Monte Carlo simulation for actually computing the limits since most of them require calculating quantiles of suprema of functionals of Brownian bridges over a finite interval or quantiles of stable distributions.

We need to truncate the ME plot near infinity in all the cases since the weak limits we obtain blow up there (it relates to t near 0 in the limit of \mathcal{MN}_n).

3.1. Fréchet case:

This case has already been discussed in Das and Ghosh (2013) and we recall it here for the sake of completeness. Define the truncated versions of \mathcal{M}_n defined in (2.1) and its limit \mathcal{M} respectively for $0 < \epsilon < 1$ as:

$$\mathcal{M}_n^\epsilon := \frac{1}{X_{(k)}} \left\{ (X_{(i)}, \hat{M}(X_{(i)})) : i = \lceil k\epsilon \rceil, \dots, k \right\} \quad \text{and} \quad \mathcal{M}^\epsilon := \left\{ \left(t^{-1}, \frac{\xi}{1-\xi} t^{-1} \right) : \epsilon \leq t \leq 1 \right\}. \quad (3.1)$$

Then $\mathcal{M}_n^\epsilon \xrightarrow{P} \mathcal{M}^\epsilon$.

Case 1 ($0 < \xi < 1/2$): From (2.2), we have that the $(1 - \alpha)100\%$ confidence band for \mathcal{M}^ϵ as

$$\mathcal{CM}_n^\epsilon := \mathcal{M}_n^\epsilon + \left\{ (x, y) : x \in \left(-\frac{c_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{c_{\alpha/2, \epsilon}}{\sqrt{k}} \right), y \in \left(-\frac{d_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{d_{\alpha/2, \epsilon}}{\sqrt{k}} \right) \right\}, \quad (3.2)$$

where

$$c_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \xi t^{-(1+\xi)} B(t), \quad (3.3)$$

$$d_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \xi t^{-1} \int_0^t y^{-(1+\xi)} B(y) dy. \quad (3.4)$$

Since the weak limit of properly scaled and shifted \mathcal{M}_n^ϵ consists of functionals of the same Brownian Bridge in both components, (3.2) provides an asymptotic confidence bound around \mathcal{M}^ϵ with $P(\mathcal{M}^\epsilon \subset \mathcal{CM}_n^\epsilon) \geq (1 - \alpha)$ for large n .

Case 2 ($1/2 < \xi < 1$): From (2.3), we have the $(1 - \alpha)100\%$ confidence band for \mathcal{M}^ϵ as

$$\mathcal{CM}_n^\epsilon = \left\{ \left(\frac{X_{(\lceil kt \rceil)}}{X_{(k)}}, \frac{\hat{M}(X_{(\lceil kt \rceil)})}{X_{(k)}} \right) + \left(-\frac{c_{\alpha_1/2, \epsilon}}{\sqrt{k}}, \frac{c_{\alpha_1/2, \epsilon}}{\sqrt{k}} \right) \times \left(\frac{X_{(1)} d_{1-\alpha_2/2}}{\lceil kt \rceil X_{(k)}}, \frac{X_{(1)} d_{\alpha_2/2}}{\lceil kt \rceil X_{(k)}} \right) : \epsilon \leq t \leq 1 \right\}, \quad (3.5)$$

where

$$d_\alpha = (1 - \alpha)\text{-th quantile of } S_{1/\xi} \text{ defined in (2.4).}$$

Here $0 < \alpha_1, \alpha_2 < 1$ are chosen such that $(1 - \alpha) = (1 - \alpha_1)(1 - \alpha_2)$. Since the random components in the first and second components in the limit of (2.3) are independent this gives us the right confidence interval so that $P(\mathcal{M}^\epsilon \subset \mathcal{CM}_n^\epsilon) \geq 1 - \alpha$. The above quantiles are calculated using Monte Carlo simulation methods. In real data examples ξ is estimated using a Hill estimator, or any reasonable estimator for the tail index of a heavy-tailed distribution.

3.2. Gumbel case

This is the case where $F \in D(G_0)$. Many well-known distribution functions such as exponential, normal, log-normal distributions fall into this class. First we define the truncated versions of \mathcal{M}_n defined in (2.7) and its limit \mathcal{M} respectively for $0 < \epsilon < 1$ as:

$$\mathcal{M}_n^\epsilon := \frac{1}{X_{(\lceil k/\epsilon \rceil)} - X_{(k)}} \left\{ (X_{(i)} - X_{(k)}, \hat{M}(X_{(i)})) : i = \lceil k\epsilon \rceil, \dots, k \right\} \quad \text{and} \quad \mathcal{M}^\epsilon := \{(-\ln(t), 1) : \epsilon \leq t \leq 1\}. \quad (3.6)$$

Then $\mathcal{M}_n^\epsilon \xrightarrow{P} \mathcal{M}^\epsilon$. Using Theorem 2.3, we have that the $(1 - \alpha)100\%$ confidence band for \mathcal{M}^ϵ as

$$\mathcal{CM}_n^\epsilon := \mathcal{M}_n^\epsilon + \left\{ (x, y) : x \in \left(-\frac{c_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{c_{\alpha/2, \epsilon}}{\sqrt{k}} \right), y \in \left(-\frac{d_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{d_{\alpha/2, \epsilon}}{\sqrt{k}} \right) \right\}, \quad (3.7)$$

where

$$c_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \left\{ eB(e^{-1}) \ln(t) + \frac{B(t)}{t} \right\} \quad (3.8)$$

$$d_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \left\{ eB(e^{-1}) + \frac{1}{t} \int_0^t \frac{B(y)}{y} dy \right\}. \quad (3.9)$$

By the same logic, as the earlier cases, (3.7) provides an asymptotic confidence bound around \mathcal{M}^ϵ with $P(\mathcal{M}^\epsilon \subset \mathcal{CM}_n^\epsilon) \geq (1 - \alpha)$ for large n . The quantiles are obtained using Monte Carlo simulation.

3.3. Weibull case

In this case $F \in D(G_\xi)$ with $\xi < 0$. Many distributions, especially with bounded right hand-tail falls into this category, for example Uniform, Beta, etc. Here we define the truncated versions of \mathcal{M}_n as defined in Proposition 2.6 and its limit \mathcal{M} respectively for $0 < \epsilon < 1$ as:

$$\mathcal{M}_n^\epsilon := \frac{1}{X_{(1)} - X_{(k)}} \left\{ (X_{(i)} - X_{(k)}, \hat{M}(X_{(i)})) : i = 2, \dots, k \right\} \quad \text{and} \quad \mathcal{M}^\epsilon := \left\{ \left(t, \frac{\xi}{1 - \xi}(t - 1) \right) : \epsilon \leq t \leq 1 \right\}. \quad (3.10)$$

Then $\mathcal{M}_n^\epsilon \xrightarrow{P} \mathcal{M}^\epsilon$. Using Theorem 2.9, we have that the $(1 - \alpha)100\%$ confidence band for \mathcal{M}^ϵ as

$$\mathcal{CM}_n^\epsilon := \mathcal{M}_n^\epsilon + \left\{ (x, y) : x \in \left(-\frac{c_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{c_{\alpha/2, \epsilon}}{\sqrt{k}} \right), y \in \left(-\frac{d_{\alpha/2, \epsilon}}{\sqrt{k}}, \frac{d_{\alpha/2, \epsilon}}{\sqrt{k}} \right) \right\}, \quad (3.11)$$

where

$$c_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \xi t^{-(1+\xi)} B(t), \quad (3.12)$$

$$d_{\alpha, \epsilon} = (1 - \alpha)\text{-th quantile of } \sup_{\epsilon \leq t \leq 1} \xi t^{-1} \int_0^t y^{-(1+\xi)} B(y) dy. \quad (3.13)$$

The bounds obtained here are very similar to the one in the Fréchet case. And using the same argument, (3.11) provides an asymptotic confidence bound around \mathcal{M}^ϵ with $P(\mathcal{M}^\epsilon \subset \mathcal{CM}_n^\epsilon) \geq (1 - \alpha)$ for large n . Similar to the previous cases, the quantiles are obtained using Monte Carlo simulation.

4. Examples: Simulated and real data

This section is devoted to application of the methodology developed for constructing confidence intervals around ME plots as derived in the Section 3.




Figure 1. ME Plot for 10000 i.i.d. Generalized Pareto random variables with $\xi = -0.5$ and $\beta = 1$.

Given an iid sample $X_1, \dots, X_n \sim F$, we are concerned with detecting if $F \in D(G_\xi)$ and if so whether ξ is positive (the Fréchet case), zero (the Gumbel case) or negative (the Weibull case). The Fréchet case has been discussed in [Das and Ghosh \(2013\)](#) with examples. Hence we concern ourselves with the other two cases for the simulated examples. First we see how our confidence intervals work in simulated examples, and then use them on real data. In all the plots below, the light blue shade creates a 95% confidence interval and the dark blue shade creates 90% confidence interval.




Figure 2. ME Plot for 10000 i.i.d. $Beta(\alpha, \beta)$ random variables with $\alpha = 2, \beta = 2$.

4.1. Simulated examples: Weibull

In this case we have $F \in D(G_\xi)$ with $\xi < 0$. We consider two families of distributions here.

1. Consider $F \sim GPD(\xi, \beta)$ with pdf given by

$$F(x) = 1 - \left(1 + \xi \frac{x}{\beta}\right)^{-1/\xi}, \quad 1 + \frac{\xi x}{\beta} > 0, \quad \beta > 0, \xi < 0.$$

Of course, $F \in D(G_\xi)$ here and F is in the Weibull domain of attraction if $\xi < 0$. In fact the Uniform $(0, 1)$ falls into this class with $\xi = -1$ and $\beta = 1$.

For our simulation example we take $\xi = -0.5, \beta = 1$ and generate 10000 iid samples from the distribution. The two plots in the left of Figure 1 are Pickands and Moment estimate of ξ for increasing values of top order statistics used. They seem reasonable close to -0.5 . For $k = 800, 1000$ and $\delta = 0.2, 0.3$ we create confidence bounds around the ME plot (in black) which clearly covers the dashed red line with slope -0.5 .

2. Next consider $F \sim \text{Beta}(a, b)$ with pdf given by

$$f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1}, \quad 0 < x < 1, \quad a, b > 0.$$

In this case $F \in D(G_{-1/b})$. We take the example $a = 2, b = 2$ where $F \in D(G_{-0.5})$. As observed in the previous example we see that the Pickands and Moment estimates approximate -0.5 well; see 2. We again create confidence bounds with $k = 800, 1000$ and $\delta = 0.2, 0.3$ and observe that the bounds cover the dashed red line with slope -0.5 .

Thus the detection in the Weibull family looks reasonable.

4.2. Simulated examples: Gumbel

Distributions in the Gumbel domain of attraction are harder to detect since a data sample has to form a plot with slope zero in this case, which is statistically unlikely. Hence confidence bounds help to an extent,

although as we will see through the three examples below that, in practice, a plotting technique is helpful to different degrees in different cases.




Figure 3. ME Plot for 10000 i.i.d. Exponential random variables with $\lambda = 1$.




Figure 4. ME Plot for 10000 i.i.d. Standard Normal random variables.

1. The first example is where F follows $Exp(1)$. We generate 10000 iid samples from the distribution and create ME plots with parameters $k = 800, 1000$ and $\delta = 0.2, 0.3$; see Figure 3. The Pickands and Moment estimates are close to zero and the confidence intervals around the ME plot in the four different

cases all cover the line with slope 0 (and intercept 1) as expected. So here the detection techniques works well.

2. The next data set we look at is a sample generated from F which follows $N(0, 1)$. We again generate 10000 iid samples from the distribution and make ME plots with parameters $k = 800, 1000$ and $\delta = 0.2, 0.3$; see Figure 4. In this case the Pickands estimate is close to zero but the Moment estimate though close zero seem to be an underestimate. The confidence intervals around the ME plot in the four different cases all cover the dashed red line with slope 0 (and intercept 1) up to some point and then it doesn't. We can believe that $F \in D(G_0)$ but the case becomes less convincing than the previous example.




Figure 5. ME Plot for 10000 i.i.d. Standard log-normal random variables.

3. Finally we look into F which is a standard Lognormal distribution. It is known the a Lognormal distribution belongs to $D(G_0)$, but on the other hand we know that it has no finite moments (unlike the Normal or Exponential case). So it is on the one-hand sub-exponential or heavy-tailed although belong to a Gumbel domain of attraction.

We simulate 10000 iid samples from a standard Lognormal distribution and create ME plots as in the previous cases. The results are in Figure 5. Both the Pickands estimate and the moment estimate of the extreme value parameter are much higher than the true value, that is zero. The ME plot with confidence intervals around it miss the target red dashed line of slope zero (and intercept 1); a larger choice of δ would make the confidence intervals large enough to cover the line, but clearly our technique doesn't seem to perform so well here. Since the Lognormal distribution has heavy tails we tend to have a positive slope of the ME plot as would happen in case when F is in the Fréchet domain of attraction. Hence overall for detecting a Gumbel domain of attraction family we need to be more careful with this technique.

4.3. Observed data: Ozone concentration at Zurich urban area

It is of interest for environmental scientists to study *ozone* concentration near urban conglomerations, as its presence in the atmosphere implies health risks related to respiratory diseases. Directive 2008/50/EC of

the European Parliament puts the target value of ozone for its member states to be within $120\mu\text{g}/\text{m}^3$. The directive says that as of January 1, 2010 ozone concentrations should not exceed this limit for more than 25 days in a calendar year where the daily calculation is based on maximum of daily 8-hour averages.




Figure 6. ME Plot for Ozone concentration in Zurich urban area.

We study a data set, freely available from www.eea.europa.eu. The data set contains daily maxima of ozone concentration (in $\mu\text{g}/\text{m}^3$) from one station in Zurich, Switzerland (station code CH 0010A, Zurich-Kaserne) located 410 *mts* above sea-level. Data is observed from January 1, 1992 to December 31, 2009. Measurements were unavailable for 22 days, which we impute by the average value of ozone concentration on the same day for other available years.

As seen in the top left plot in Figure 6 the data clearly admits periodicity. Moreover it is likely that the

data is heteroscedastic. So we homoscedasticize the data by dividing the value on each date by the standard deviation of the values on the same day over all the 18 years of data available. Since our techniques work for stationary data sets, we fit an AR (38) process to the data set (AR(38) is chosen by an AIC criterion) and observe (from the ACF; see second plot from the second line in Figure 6) that the residuals (first plot in the second line) look independent. Now we analyze the extremal behavior of the residuals of the model. The Pickands and Moments estimates give a negative value but close to 0 and we can hypothesise that the sample is from a Weibull domain of attraction family. But, since the value of the parameter is close to 0 we also check whether the data is possibly from a Gumbel domain of attraction family. The confidence bounds (90% deep blue and 95% light blue) are created assuming $F \in D(G_0)$ for $k = 329$ and $k = 658$; which are 5% and 10% of the data set and with $\delta = 0.2, 0.3$. Observe that the 90% bounds tend to reject the hypothesis of the underlying $F \in D(G_0)$ and the 95% do not. This is most likely a result of the parameter being close to zero.

On the other hand using the Pickands estimate to estimate the tail index we get $\xi = -0.14$ (for $k = 658$) and $\xi = -0.11$ (for $k = 329$) and the confidence bounds (again 90% deep blue and 95% light blue) for $\delta = 0.2, 0.3$ covers the straight line with the slopes ξ quite well. Hence we are expect that the underlying distribution is in fact in a Weibull domain off attraction with parameter close to $\xi = 0.1$.

4.4. Observed data: Flow-rates at river Aare

The other data we analyze is maximum daily flow-rate at river Aare. River Aare flows through Switzerland and some manufacturing and power plants located near the river are often concerned about flooding on the river. The data we analyse has been collected from the Federal office of the Environment (FOEN), Bern and generously provided to us by Kernkraftwerk Gösgen-Däniken. It pertains to daily maximum flow-rates of Aare at the measurement station Aare-Murgenthal (2063) measured in m^3/sec from 1st January 1974 to 20th October, 2010. See also www.hydrodaten.admin.ch/d/2063.htm.

Note that the data admits to possibilities of measurement error since automated measurement at the specific station started only in 1993. Moreover, the control authorities aim to maintain the flow-rate of Aare at the Aare-Murgenthal (2063) station below $850 m^3/sec$ and would do so by using opening or closing log-gates. This manually hinders the possibility of the data set being tuitionary. We were informed that such manual intervention has been done a couple of times.

To analyse the data, we first note the seasonality pattern in the data set; see top left plot in Figure 7. Hence as in the previous example we fit an AR process and work with the residuals obtained after the model-fitting. Observe that the Pickands and Moment estimates both indicate towards a small but positive value of the extreme value parameter; but does not completely reject the possibility of it being zero. We again create 90% (dark blue) and 95% (light blue) confidence bounds under the Gumbel assumption for $k = 673$ and $k = 1345$ (again 5% and 10%) of the sample size and $\delta = 0.1, 0.2$. The detection technique seems to reject that the underlying distribution $F \in D(G_0)$.

Now we allow a Pickands estimate to chose the extreme value parameter which gives a value of around $\xi = 0.16$ (for $k = 673$ and $k = 1345$) and the confidence bounds seem to support that the data is from a distribution in the Fréchet domain of attraction. Thus we may conclude that flow-rate data at Murgenthal station is perhaps slightly heavy-tailed even if marginally so.




Figure 7. ME Plot for Aare river flow data.

References

- BEIRLANT, J., GOEGBEUR, Y., TEUGELS, J. and SEGERS, J. (2004). *Statistics of Extremes. Wiley Series in Probability and Statistics*. John Wiley & Sons Ltd., Chichester. Theory and applications, with contributions from Daniel De Waal and Chris Ferro. [MR2108013 \(2005j:62002\)](#)
- BILLINGSLEY, P. (1968). *Convergence of Probability Measures*. John Wiley & Sons Inc., New York.
- BINGHAM, N. H., GOLDIE, C. M. and TEUGELS, J. L. (1987). *Regular Variation*. Cambridge University Press.

- DAS, B., EMBRECHTS, P. and FASEN, V. (2013). Four Theorems and a Financial Crisis. *The International Journal of Approximate Reasoning* **54** 701–716.
- DAS, B. and GHOSH, S. (2013). Weak limits of exploratory plots in the analysis of extremes. *Bernoulli* **19** 308–342.
- DAVISON, A. C. and SMITH, R. L. (1990). Models for exceedances over high thresholds (with discussion). *Journal of Royal Statistical Society, Series B* **52** 393–442.
- DE HAAN, L. (1970). *On Regular Variation and Its Application to the Weak Convergence of Sample Extremes*. Mathematisch Centrum Amsterdam.
- DE HAAN, L. and FERREIRA, A. (2006). *Extreme Value Theory: An Introduction*. Springer-Verlag, New York.
- DONNELLY, C. and EMBRECHTS, P. (2010). The devil is in the tails: actuarial mathematics and the subprime mortgage crisis. *ASTIN Bulletin* **40** 1–33.
- EMBRECHTS, P., KLÜPPELBERG, C. and MIKOSCH, T. (1997). *Modelling Extreme Events for Insurance and Finance*. Springer-Verlag, Berlin.
- GELUK, J. L. and DE HAAN, L. (1987). *Regular Variation, Extensions and Tauberian Theorems*. CWI Tract **40**. Stichting Mathematisch Centrum, Centrum voor Wiskunde en Informatica, Amsterdam.
- GHOSH, S. and RESNICK, S. I. (2010). A Discussion on Mean Excess Plots. *Stochastic Processes and their Applications* **120** 1492–1517.
- HALL, W. J. and WELLNER, J. (1981). Mean residual life. In *Statistics and related topics (Ottawa, Ont., 1980)* 169–184. North-Holland, Amsterdam-New York. [MR665274 \(83k:62136\)](#)
- MAULIK, K., RESNICK, S. I. and ROOTZÉN, H. (2002). Asymptotic independence and a network traffic model. *Journal of Applied Probability* **39** 671–699.
- MCNEIL, A. J., FREY, R. and EMBRECHTS, P. (2005). *Quantitative Risk Management: Concepts, Techniques and Tools*. Princeton University Press, Princeton, NJ.
- RESNICK, S. I. (2007). *Heavy Tail Phenomena: Probabilistic and Statistical Modeling*. Springer Series in Operations Research and Financial Engineering. Springer-Verlag, New York.
- RESNICK, S. I. (2008). *Extreme Values, Regular Variation and Point Processes*. Springer Series in Operations Research and Financial Engineering. Springer, New York. Reprint of the 1987 original.
- ROOTZÉN, H. (2009). Weak convergence of the tail empirical process for dependent sequences. *Stochastic Processes and their Applications* **119** 468–490.
- SENETA, E. (1976). *Regularly Varying Functions*. Springer-Verlag, New York. Lecture Notes in Mathematics, 508.
- VAN DER VAART, A. W. and WELLNER, J. A. (1996). *Weak Convergence and Empirical Processes with Applications to Statistics*. Springer-Verlag, New York.
- YANG, G. L. (1978). Estimation of a biometric function. *The Annals of Statistics* **6** 112–116.